Lesson 1.3

Midpoint & Distance Formula

Two formulas — and how to never mix them up

The trick

$$\text{Midpoint} = \text{Mean}$$

The trick


$$\text{Midpoint} = \text{Mean}$$

$$\text{Distance} = \text{Difference}$$


Midpoint: simple case first



Midpoint: simple case first




Midpoint: simple case first




Add and halve: $\dfrac{-3+3}{2} = 0$ ✓

Midpoint: simple case first



Add and halve: $\dfrac{-3+3}{2} = 0$ ✓ Subtract: $\dfrac{3-(-3)}{2} = 3$ ×

Midpoint: simple case first




Add and halve: $\dfrac{-3+3}{2} = 0$ ✓ Subtract: $\dfrac{3-(-3)}{2} = 3$ ×

So the midpoint **adds** — it's the **average** (mean).

Midpoint formula


$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$



Midpoint formula

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

Example: $A(2, 3)$, $B(8, 7)$




Midpoint formula

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

Example: $A(2, 3)$, $B(8, 7)$

x: $\frac{2+8}{2} = 5$




Midpoint formula

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

Example: $A(2, 3)$, $B(8, 7)$

x: $\frac{2+8}{2} = 5$

y: $\frac{3+7}{2} = 5$



Midpoint formula


$$M = \left(\frac{x_1 + x_2}{2}, \ \frac{y_1 + y_2}{2} \right)$$

Example: $A(2, 3)$, $B(8, 7)$


x: $\frac{2+8}{2} = 5$

y: $\frac{3+7}{2} = 5$


$M = (5, 5)$




Distance: simple case first



Distance: simple case first




Distance: simple case first




Count: 3. That's $5 - 2$ — a **difference**.

Distance = Pythagoras on a grid



Find AB.


Distance = Pythagoras on a grid



Find AB.

Across 3, up 4.

Distance = Pythagoras on a grid




Find AB.

Across 3, up 4.

$d = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$

Distance = Pythagoras on a grid



Find AB.

Across 3, up 4.

$d = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

Two common mistakes

Watch out! Square **each** difference before adding:
$\sqrt{3^2 + 4^2} = 5$, not $3 + 4 = 7$.

Two common mistakes

Watch out! Square **each** difference before adding: $\sqrt{3^2 + 4^2} = 5$, not $3 + 4 = 7$.

Watch out! Testing add vs. subtract? Don't use 0 and 2 — both give 1. Use 2 and 4: the middle is $3 = \frac{2+4}{2}$.

Your turn! Pause the video and try it.

$P(-1, 4)$ and $Q(5, -4)$. Find the midpoint of $\overline{PQ}$ and the length PQ.

Your turn! Pause the video and try it.

$P(-1, 4)$ and $Q(5, -4)$. Find the midpoint of $\overline{PQ}$ and the length PQ.

$$M = \left(\frac{-1+5}{2}, \frac{4+(-4)}{2} \right) = (2, 0)$$

Your turn! Pause the video and try it.

$P(-1, 4)$ and $Q(5, -4)$. Find the midpoint of $\overline{PQ}$ and the length PQ.

$$M = \left(\frac{-1+5}{2}, \frac{4+(-4)}{2} \right) = (2, 0)$$

$$PQ = \sqrt{(5-(-1))^2 + (-4-4)^2} = \sqrt{6^2+8^2} = \sqrt{100} = 10$$

Remember

- **Midpoint = Mean**: add the coordinates, divide by 2.
- **Distance = Difference**: subtract, square, add, square root.
- The distance formula is just the Pythagorean Theorem.

Now practice at your level:

fangmath.com → Geometry → Chapter 1