

Probability Theory on Coin Toss Space

Peiliang Guo

January 14, 2017

Problem 1.

(i) We first prove disjoint additivity: if E_1 and E_2 are disjoint, then $\mathbb{P}(E_1) + \mathbb{P}(E_2) = \mathbb{P}(E_1 \cup E_2)$.

$$\mathbb{P}(E_1) + \mathbb{P}(E_2) = \sum_{\omega \in E_1} \mathbb{P}(\omega) + \sum_{\omega \in E_2} \mathbb{P}(\omega) = \sum_{\omega \in (E_1 \cup E_2)} \mathbb{P}(\omega) = \mathbb{P}(E_1 \cup E_2)$$

Since A and A^C are disjoint,

$$\mathbb{P}(A^C) + \mathbb{P}(A) = \mathbb{P}(\Omega) = 1 \Rightarrow \mathbb{P}(A^C) = 1 - \mathbb{P}(A)$$

(ii) Note that disjoint additivity holds for more than two sets, so the equality case holds trivially.

Disjoint additivity also shows if $E_1 \subset E_2$, then $\mathbb{P}(E_1) \leq \mathbb{P}(E_2)$, since

$$E_2 = E_1 \cup (E_2 \setminus E_1) \Rightarrow \mathbb{P}(E_2) = \mathbb{P}(E_1) + \mathbb{P}(E_2 \setminus E_1) \Rightarrow \mathbb{P}(E_1) \leq \mathbb{P}(E_2)$$

Now, let $E_0 = A_0$, and let $E_k = A_k \setminus (\cup_{i=1}^{k-1} E_i)$ for $k = 2, 3, \dots, N$, then E_i are disjoint and $\cup_{i=1}^N E_i = \cup_{i=1}^N A_i$. So,

$$\mathbb{P}\left(\bigcup_{n=1}^N A_n\right) = \mathbb{P}\left(\bigcup_{n=1}^N E_n\right) = \sum_{n=1}^N \mathbb{P}(E_n) \leq \sum_{n=1}^N \mathbb{P}(A_n)$$

Problem 2.

(i) The probabilities follow binomial probabilities with $p = \tilde{p} = \frac{1}{2}$. i.e.

$$\tilde{\mathbb{P}}(S_3 = 3) = \binom{3}{3} \left(\frac{1}{2}\right)^3 = \frac{1}{8}$$

$$\tilde{\mathbb{P}}(S_3 = 2) = \binom{3}{2} \left(\frac{1}{2}\right)^3 = \frac{3}{8}$$

$$\tilde{\mathbb{P}}(S_3 = 1) = \binom{3}{1} \left(\frac{1}{2}\right)^3 = \frac{3}{8}$$

$$\tilde{\mathbb{P}}(S_3 = 0) = \binom{3}{0} \left(\frac{1}{2}\right)^3 = \frac{1}{8}$$

(ii)

$$\tilde{\mathbb{E}}S_1 = \binom{1}{1} \tilde{p}S_1(H) + \tilde{q}S_1(T) = \frac{1}{2} \cdot 8 + \frac{1}{2} \cdot 2 = 5$$

$$\tilde{\mathbb{E}}S_2 = \binom{2}{2} \tilde{p}^2 S_2(HH) + \binom{2}{1} \tilde{p}\tilde{q}S_2(HT) + \binom{2}{0} \tilde{q}^2 S_2(TT) = \frac{1}{4} \cdot 16 + \frac{2}{4} \cdot 4 + \frac{1}{4} \cdot 1 = 6.25$$

$$\tilde{\mathbb{E}}S_3 = \binom{3}{3} \tilde{p}^3 S_3(HHH) + \binom{3}{2} \tilde{p}^2 \tilde{q}S_3(HHT) + \binom{3}{1} \tilde{p}\tilde{q}^2 S_3(HTT) + \binom{3}{0} \tilde{q}^3 S_3(TTT) = 7.8125$$

The rate of growth of mean stock price under $\tilde{\mathbb{P}}$ is $1 + r = 1.25$.

(iii)

$$\mathbb{P}(S_3 = 32) = \binom{3}{3} \left(\frac{2}{3}\right)^3 = \frac{8}{27}$$

$$\mathbb{P}(S_3 = 8) = \binom{3}{2} \left(\frac{2}{3}\right)^2 \left(\frac{1}{3}\right) = \frac{12}{27}$$

$$\mathbb{P}(S_3 = 2) = \binom{3}{1} \left(\frac{2}{3}\right)^1 \left(\frac{1}{3}\right)^2 = \frac{6}{27}$$

$$\mathbb{P}(S_3 = 0.5) = \binom{3}{0} \left(\frac{1}{3}\right)^3 = \frac{1}{27}$$

$$\mathbb{E}S_1 = \frac{2}{3} \cdot 8 + \frac{1}{3} \cdot 2 = 6$$

$$\mathbb{E}S_2 = \frac{4}{9} \cdot 16 + \frac{4}{9} \cdot 4 + \frac{1}{9} \cdot 1 = 9$$

$$\mathbb{E}S_3 = \frac{8}{27} \cdot 32 + \frac{12}{27} \cdot 8 + \frac{6}{27} \cdot 2 + \frac{1}{27} \cdot 0.5 = 13.5$$

The rate of growth of mean stock price under \mathbb{P} is 1.5.

Problem 3.

By martingale property, for any $0 \leq n \leq N - 1$,

$$M_n = \mathbb{E}_n M_{n+1} \Rightarrow \varphi(M_n) = \varphi(\mathbb{E}_n M_{n+1})$$

By conditional Jensen's inequality, since φ is convex,

$$\varphi(M_n) = \varphi(\mathbb{E}_n M_{n+1}) \leq \mathbb{E}_n[\varphi(M_{n+1})]$$

Thus, $\varphi(M_0), \dots, \varphi(M_N)$ is a submartingale.

Problem 4.

(i) For arbitrary $n \geq 0$, we have

$$\begin{aligned}
\mathbb{E}_n[M_{n+1}] &= \mathbb{E}_n[M_n + X_{n+1}] \\
&= \mathbb{E}_n[M_n \cdot 1] + \mathbb{E}_n[X_{n+1}] && \text{by linearity of C.E.} \\
&= M_n \mathbb{E}_n[1] + \mathbb{E}_n[X_{n+1}] && \text{by taking out what is known} \\
&= M_n + \mathbb{E}[X_{n+1}] && \text{by independence} \\
&= M_n + \frac{1}{2} \cdot 1 + \frac{1}{2} \cdot (-1) \\
&= M_n
\end{aligned}$$

(ii) For arbitrary $n \geq 0$, we have

$$\begin{aligned}
\mathbb{E}_n[S_{n+1}] &= \mathbb{E}_n \left[e^{\sigma M_{n+1}} \left(\frac{2}{e^\sigma + e^{-\sigma}} \right)^{n+1} \right] \\
&= \mathbb{E}_n \left[e^{\sigma M_n} \left(\frac{2}{e^\sigma + e^{-\sigma}} \right)^{n+1} e^{\sigma X_{n+1}} \right] \\
&= \left(\frac{2}{e^\sigma + e^{-\sigma}} \right) \mathbb{E} [e^{\sigma X_{n+1}}] \mathbb{E}_n \left[e^{\sigma M_n} \left(\frac{2}{e^\sigma + e^{-\sigma}} \right)^n \right] \\
&= \left(\frac{2}{e^\sigma + e^{-\sigma}} \right) \left(\frac{1}{2} e^\sigma + \frac{1}{2} e^{-\sigma} \right) S_n \\
&= S_n
\end{aligned}$$

Problem 5.

(i)

$$\begin{aligned}
2I_n &= 2 \sum_{j=0}^{n-1} M_j (M_{j+1} - M_j) = 2 \sum_{j=0}^{n-1} \sum_{i=1}^j X_i X_{j+1} = 2 \left(\sum_{j=1}^n \sum_{i=1}^j X_i X_j - \sum_{j=1}^n X_j^2 \right) \\
&= \left(\sum_{j=1}^n X_j \right)^2 + \sum_{j=1}^n X_j^2 - 2 \sum_{j=1}^n X_j^2 = M_n^2 - n
\end{aligned}$$

the last equality holds because $X_j^2 = 1$ in both cases.

(ii)

$$\begin{aligned}
E_n[f(I_{n+1})] &= E_n \left[f \left(\frac{1}{2} M_{n+1}^2 - \frac{n+1}{2} \right) \right] \\
&= E_n \left[f \left(\frac{1}{2} (M_n^2 + 2X_{n+1}M_n + X_{n+1}^2 - n - 1) \right) \right] \\
&= E_n [f(I_n + M_n X_{n+1})]
\end{aligned}$$

Since I_n and M_n are known at time n , we have

$$\begin{aligned} g(I_n) &= E_n[f(I_{n+1})] = E_n[f(I_n + M_n X_{n+1})] \\ &= \frac{1}{2}f(I_n + M_n) + \frac{1}{2}f(I_n - M_n) \\ &= \frac{1}{2}\left(f(I_n + \sqrt{2I_n + n}) + f(I_n - \sqrt{2I_n + n})\right) \end{aligned}$$

Problem 6.

For any $0 \leq n \leq N - 1$,

$$\begin{aligned} E_n[I_{n+1}] &= E_n[I_n + \Delta_n(M_{n+1} - M_n)] \\ &= I_n + \Delta_n E_n[M_{n+1} - M_n] \\ &= I_n \end{aligned}$$

Problem 7.

Let $X_0 = 0$, $X_1 = 1$ with probability $\frac{1}{2}$, and $X_1 = -1$ with probability $\frac{1}{2}$. For $n \geq 2$, $X_n = X_{n-1} + \varepsilon_n$ with probability $\frac{1}{2}$, and $X_n = X_{n-1} - \varepsilon_n$ with probability $\frac{1}{2}$, where $\varepsilon_n = 1$ if $X_{n-2} > X_{n-1}$, and $\varepsilon_n = 2$ if $X_{n-2} < X_{n-1}$.

Problem 8.

(i) For $n = N - 1$, for any $\omega_1 \dots \omega_n$,

$$\begin{aligned} M'_n(\omega_1 \dots \omega_n) &= \frac{1}{1+r} [\tilde{p} M'_N(\omega_1 \dots \omega_n H) + \tilde{q} M'_N(\omega_1 \dots \omega_n T)] \\ &= \frac{1}{1+r} [\tilde{p} M_N(\omega_1 \dots \omega_n H) + \tilde{q} M_N(\omega_1 \dots \omega_n T)] \\ &= M_N(\omega_1 \dots \omega_n) \end{aligned}$$

The argument can be applied recursively until n reaches 0.

(ii) for any $0 \leq n \leq N - 1$, and for every $\omega_1 \dots \omega_n$,

$$\begin{aligned} \tilde{\mathbb{E}}_n \left[\frac{V_{n+1}}{(1+r)^{n+1}} \right] (\omega_1 \dots \omega_n) &= \frac{1}{(1+r)^{n+1}} [\tilde{p} V_{n+1}(\omega_1 \dots \omega_n H) + \tilde{q} V_{n+1}(\omega_1 \dots \omega_n T)] \\ &= \frac{1}{(1+r)^n} \frac{1}{1+r} [\tilde{p} V_{n+1}(\omega_1 \dots \omega_n H) + \tilde{q} V_{n+1}(\omega_1 \dots \omega_n T)] \\ &= \frac{V_n}{(1+r)^n} \end{aligned}$$

(iii) for any $0 \leq n \leq N - 1$, and for every $\omega_1 \dots \omega_n$, (here we suppress $\omega_1 \dots \omega_n$)

$$\begin{aligned} \tilde{\mathbb{E}}_n \left[\frac{V'_{n+1}}{(1+r)^{n+1}} \right] &= \frac{1}{(1+r)^{n+1}} \tilde{\mathbb{E}}_n \left[\tilde{\mathbb{E}}_{n+1} \left[\frac{V'_N}{(1+r)^{N-n-1}} \right] \right] \\ &= \frac{1}{(1+r)^N} \tilde{\mathbb{E}}_n [V'_N] \\ &= \frac{V'_n}{(1+r)^n} \end{aligned}$$

(iv) Conclusion can be drawn from part (i), since the terminal payoff are the same $V_N = V'_N$, and V_0, V_1, \dots, V_N , and V'_0, V'_1, \dots, V'_N are martingales.

Problem 9.

(i) We have

$$\begin{aligned}\tilde{p}_1(H) &= \frac{1 + r_1(H) - d_1(H)}{u_1(H) - d_1(H)} = \frac{1 + \frac{1}{4} - 1}{\frac{3}{2} - 1} = \frac{1}{2}, \tilde{q}_1(H) = 1 - \tilde{p}_1(H) = \frac{1}{2} \\ \tilde{p}_1(T) &= \frac{1 + r_1(T) - d_1(T)}{u_1(T) - d_1(T)} = \frac{1 + \frac{1}{2} - 1}{4 - 1} = \frac{1}{6}, \tilde{q}_1(T) = 1 - \tilde{p}_1(T) = \frac{5}{6} \\ \tilde{p}_0 &= \frac{1 + r_0 - d_0}{u_0 - d_0} = \frac{1 + \frac{1}{4} - \frac{1}{2}}{2 - \frac{1}{2}} = \frac{1}{2}, \tilde{q}_1(T) = 1 - \tilde{p}_1(T) = \frac{1}{2}\end{aligned}$$

Therefore,

$$\begin{aligned}\tilde{\mathbb{P}}(HH) &= \tilde{p}_0 \tilde{p}_1(H) = \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}, \tilde{\mathbb{P}}(HT) = \tilde{p}_0 \tilde{q}_1(H) = \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4} \\ \tilde{\mathbb{P}}(TH) &= \tilde{q}_0 \tilde{p}_1(T) = \frac{1}{2} \cdot \frac{1}{6} = \frac{1}{12}, \tilde{\mathbb{P}}(TT) = \tilde{q}_0 \tilde{q}_1(T) = \frac{1}{2} \cdot \frac{5}{6} = \frac{5}{12}\end{aligned}$$

Note that

$$\begin{aligned}\tilde{\mathbb{E}} \left[\frac{S_2}{(1+r_0)(1+r_1)} \right] &= \frac{1}{4} \cdot \frac{12}{(1+\frac{1}{4})(1+\frac{1}{4})} + \frac{1}{4} \cdot \frac{8}{(1+\frac{1}{4})(1+\frac{1}{4})} + \\ &\quad \frac{1}{12} \cdot \frac{8}{(1+\frac{1}{4})(1+\frac{1}{2})} + \frac{5}{12} \cdot \frac{2}{(1+\frac{1}{4})(1+\frac{1}{2})} \\ &= 4 = S_0\end{aligned}$$

(ii) $V_2 = (S_2 - 7)^+$, so $V_2(HH) = 12 - 7 = 5$, $V_2(HT) = 8 - 7 = 1$, $V_2(TH) = 8 - 7 = 1$, $V_2(TT) = 0$. Therefore,

$$V_1(H) = \tilde{\mathbb{E}}_1 \left[\frac{V_2}{1+r_1} \right] (H) = \frac{1}{2} \cdot \frac{5}{1+\frac{1}{4}} + \frac{1}{2} \cdot \frac{1}{1+\frac{1}{4}} = \frac{12}{5}$$

$$V_1(T) = \tilde{\mathbb{E}}_1 \left[\frac{V_2}{1+r_1} \right] (T) = \frac{1}{6} \cdot \frac{1}{1+\frac{1}{2}} + 0 = \frac{1}{9}$$

$$V_0 = \tilde{\mathbb{E}} \left[\frac{V_2}{(1+r_0)(1+r_1)} \right] = \frac{1}{4} \cdot \frac{5}{(1+\frac{1}{4})(1+\frac{1}{4})} + \frac{1}{4} \cdot \frac{1}{(1+\frac{1}{4})(1+\frac{1}{4})} + \frac{1}{12} \cdot \frac{1}{(1+\frac{1}{4})(1+\frac{1}{2})} + \frac{5}{12} \cdot \frac{1}{(1+\frac{1}{4})(1+\frac{1}{2})} = \frac{226}{225}$$

(iii)

$$\Delta_0 = \frac{V_1(H) - V_1(T)}{S_1(H) - S_1(T)} = \frac{\frac{12}{5} - \frac{1}{9}}{8 - 2} = \frac{103}{270}$$

(iv)

$$\Delta_1(H) = \frac{V_2(HH) - V_2(HT)}{S_2(HH) - S_2(HT)} = \frac{5 - 1}{12 - 8} = 1$$

Problem 10.

(i) for any $0 \leq n \leq N - 1$, we have

$$\begin{aligned}
\tilde{\mathbb{E}}_n \left[\frac{X_{n+1}}{(1+r)^{n+1}} \right] &= \tilde{\mathbb{E}}_n \left[\frac{\Delta_n Y_{n+1} S_n + (1+r)(X_n - \Delta_n S_n)}{(1+r)^{n+1}} \right] \\
&= \frac{1}{(1+r)^n} \left[\Delta_n S_n \frac{\tilde{\mathbb{E}}_n[Y_{n+1}]}{1+r} + X_n - \Delta_n S_n \right] \\
&= \frac{X_n}{(1+r)^n} + \frac{1}{(1+r)^{n+1}} \Delta_n S_n \left[\tilde{\mathbb{E}}_n[Y_{n+1}] - 1 - r \right] \\
&= \frac{X_n}{(1+r)^n} + \frac{1}{(1+r)^{n+1}} \Delta_n S_n \left[\frac{1+r-d}{u-d} \cdot u + \frac{u-1-r}{u-d} \cdot d - 1 - r \right] \\
&= \frac{X_n}{(1+r)^n}
\end{aligned}$$

(ii) By definition, we have $V_n = \frac{1}{1+r} \tilde{\mathbb{E}}_n[V_{n+1}]$, i.e. V_0, V_1, \dots, V_N is a $\tilde{\mathbb{P}}$ martingale. Also, since $X_N = V_N$ for all $\omega_1 \dots \omega_N$, we have $X_n = V_n$ for all n in all $\omega_1 \dots \omega_N$. Therefore, V_n is indeed the price of the derivative that pays V_N at time N .

(iii) Since $A_n \in (0, 1)$,

$$\tilde{\mathbb{E}}_n \left[\frac{S_{n+1}}{1+r} \right] = \tilde{\mathbb{E}}_n \left[\frac{(1 - A_{n+1})Y_{n+1}S_n}{1+r} \right] < S_n \tilde{\mathbb{E}}_n \left[\frac{Y_{n+1}}{1+r} \right] = \frac{S_n}{1+r} \left(\frac{1+r-d}{u-d} \cdot u + \frac{u-1-r}{u-d} \cdot d \right) = S_n$$

Now, suppose A_n is constant $a \in (0, 1)$,

$$\begin{aligned}
\tilde{\mathbb{E}}_n \left[\frac{S_{n+1}}{(1-a)^{n+1}(1+r)^{n+1}} \right] &= \frac{S_n}{(1-a)^n(1+r)^n} \tilde{\mathbb{E}}_n \left[\frac{(1-a)Y_{n+1}}{(1-a)(1+r)} \right] \\
&= \frac{S_n}{(1-a)^n(1+r)^n} \left(\frac{1+r-d}{u-d} \cdot u \frac{u-1-d}{u-d} \cdot d \right) \\
&= \frac{S_n}{(1-a)^n(1+r)^n}
\end{aligned}$$

Problem 11.

(i)

$$F_N + P_N = S_N - K + (K - S_N)^+ = \begin{cases} S_N - K & , S_N \geq K \\ 0 & , S_N < K \end{cases} = (S_N - K)^+ = C_N$$

(ii)

$$F_n + P_n = \tilde{\mathbb{E}}_n \left[\frac{F_N}{(1+r)^{N-n}} \right] + \tilde{\mathbb{E}}_n \left[\frac{P_N}{(1+r)^{N-n}} \right] = \tilde{\mathbb{E}}_n \left[\frac{F_N + P_N}{(1+r)^{N-n}} \right] = \tilde{\mathbb{E}}_n \left[\frac{C_N}{(1+r)^{N-n}} \right] = C_n$$

(iii)

$$F_0 = \tilde{\mathbb{E}}_0 \left[\frac{F_N}{(1+r)^N} \right] = \tilde{\mathbb{E}}_0 \left[\frac{S_N - K}{(1+r)^N} \right] = \tilde{\mathbb{E}}_0 \left[\frac{S_N}{(1+r)^N} \right] - \frac{K}{(1+r)^N} = S_0 - \frac{K}{(1+r)^N}$$

(iv) At time 0: have 1 share of stock and $F_0 - S_0 = -\frac{K}{(1+r)^N}$ in bank account

At time N : $S_N + (1+r)^N \left(-\frac{K}{(1+r)^N}\right) = S_N - K = F_N$

(v)

$$C_0 = F_0 + P_0 = \tilde{\mathbb{E}}_0 \left[\frac{S_N - S_0(1+r)^N}{(1+r)^N} \right] + P_0 = P_0$$

(vi) Put all subscripts (conditional expectations) to n instead of 0

Problem 12.

At time m , the value V_m of the chooser's option is

$$V_m = \max(C_m, P_m) = P_m + (C_m - P_m)^+ = P_m + F_m^+ = P_m + \left(S_m - \frac{K}{(1+r)^{N-m}} \right)^+$$

Therefore, at time 0, the price of the chooser's option is

$$V_0 = P_0 + C'_0$$

where C' is a call option with strike $\frac{K}{(1+r)^{N-m}}$ and maturity m .

Problem 13.

(i) For every function g , we can write

$$\tilde{\mathbb{E}}_n[g(S_{n+1}, Y_{n+1})] = \tilde{p}g(uS_n, Y_n + uS_n) + \tilde{q}g(dS_n, Y_n + dS_n) = h(S_n, Y_n)$$

Therefore, $(S_n, Y_n), n = 0, 1, \dots, N$ is Markov.

(ii)

$$V_N = f\left(\frac{1}{N+1} \sum_0^N S_n\right) \Rightarrow v_N(s, y) = f\left(\frac{1}{N+1}y\right)$$

Now, for $n = 0, 1, \dots, N-1$,

$$v_n(s, y) = \tilde{p}v_{n+1}(us, y + us) + \tilde{q}v_{n+1}(ds, y + ds)$$

Problem 14.

(i) For every function g , and for $n = M, M+1, \dots, N$,

$$\tilde{\mathbb{E}}_n[g(S_{n+1}, Y_{n+1})] = \tilde{p}g(uS_n, Y_n + uS_n) + \tilde{q}g(dS_n, Y_n + dS_n)$$

and for $n = 0, 1, \dots, M-2$,

$$\tilde{\mathbb{E}}_n[g(S_{n+1}, Y_{n+1})] = \tilde{p}g(uS_n, 0) + \tilde{q}g(dS_n, 0)$$

(ii) For $n = M+1, \dots, N-1$,

$$v_n(s, y) = \frac{1}{1+r} [\tilde{p}v_{n+1}(us, y + us) + \tilde{q}v_{n+1}(ds, y + ds)]$$

For $n = M$,

$$v_n(s) = \frac{1}{1+r} [\tilde{p}v_{n+1}(us, us) + \tilde{q}v_{n+1}(ds, ds)]$$

For $n = 0, 1, \dots, M-1$,

$$v_n(s) = \frac{1}{1+r} [\tilde{p}v_{n+1}(us) + \tilde{q}v_{n+1}(ds)]$$