




Arabidopsis method comparison · one million input templates per library



























Each point is one treatment or biological ATAC replicate; controls are not plotted. Quality was identical across three technical repeats.
FRiP uses SE read spans or PE outer-fragment spans / retained spans; peak widths and coverage are in quality-summary.tsv.
Missing correlation is omitted, never replaced by zero. Baseline regions are a fixed comparison set, not truth. Genrich uses its recorded q-value operating point.


Observed component cost · median and range of three technical repeats
















All libraries in each cohort; indexing and common evaluation excluded. Index costs are reported separately.
Fourteen tool CPUs, 80 GiB per experiment; randomized method order; shared filesystem and OS caches. Error bars are observed ranges, not confidence intervals.
RSS is a sampled sum for the tool process group, not whole-host peak memory. These are component costs, not complete workflow timings.



Nested, seeded subsets of two biological replicates on TAIR10. The five-million depth was measured once per method.
Held-out FRiP denominator is retained fragments in the evaluated replicate; Jaccard uses peak base-pair intersection / union.
Increased depth changes both peak discovery and covered bases; these observations do not establish a universal plant QC threshold.

TAIR10 and Col-CEN v1.2 · one million ATAC pairs



Same reads and policy; unique reciprocal assembly mappings only. At least 95% aligned bases in both directions and 95% round-trip span overlap. See the mapping table for all exclusions. Concordance is not biological accuracy.