

Metropolis-Hastings

Metropolis-Hastings

Sometimes Gibbs samples are too correlated

Metropolis-Hastings

Sometimes Gibbs samples are too correlated

Apply rejection sampling to Markov Chains

Metropolis-Hastings

For $k = 1, 2, \dots$

- Sample x' from a **wrong** $Q(x^k \rightarrow x')$

Metropolis-Hastings

For $k = 1, 2, \dots$

- Sample x' from a **wrong** $Q(x^k \rightarrow x')$
- Accept proposal x' with probability $A(x^k \rightarrow x')$

Metropolis-Hastings

For $k = 1, 2, \dots$

- Sample x' from a **wrong** $Q(x^k \rightarrow x')$
- Accept proposal x' with probability $A(x^k \rightarrow x')$
- Otherwise stay at x^k

$$x^{k+1} = x^k$$

Metropolis-Hastings

For $k = 1, 2, \dots$

- Sample x' from a **wrong** $Q(x^k \rightarrow x')$
- Accept proposal x' with probability $A(x^k \rightarrow x')$
- Otherwise stay at x^k

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

Metropolis-Hastings

For $k = 1, 2, \dots$

- Sample x' from a **wrong** $Q(x^k \rightarrow x')$
- Accept proposal x' with probability $A(x^k \rightarrow x')$
- Otherwise stay at x^k

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

$$\begin{aligned} T(x \rightarrow x) &= Q(x \rightarrow x)A(x \rightarrow x) \\ &\quad + \sum_{x' \neq x} Q(x \rightarrow x')(1 - A(x \rightarrow x')) \end{aligned}$$

Metropolis-Hastings

For $k = 1, 2, \dots$

- Sample x' from a **wrong** $Q(x^k \rightarrow x')$
- Accept proposal x' with probability $A(x^k \rightarrow x')$
- Otherwise stay at x^k

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

$$\begin{aligned} T(x \rightarrow x) &= Q(x \rightarrow x)A(x \rightarrow x) \\ &\quad + \sum_{x' \neq x} Q(x \rightarrow x')(1 - A(x \rightarrow x')) \end{aligned}$$

How to choose A: $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

Detailed Balance

$$\pi(x') = \sum_x \pi(x) T(x \rightarrow x')$$

Detailed Balance

If $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

Then $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

Detailed Balance

If $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

Then $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

Proof $\sum_x \pi(x)T(x \rightarrow x') = \sum_x \pi(x')T(x' \rightarrow x)$

Detailed Balance

If $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

Then $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

Proof
$$\begin{aligned} \sum_x \pi(x)T(x \rightarrow x') &= \sum_x \pi(x')T(x' \rightarrow x) \\ &= \pi(x') \sum_x T(x' \rightarrow x) \end{aligned}$$

Detailed Balance

If $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

Then $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

Proof
$$\begin{aligned} \sum_x \pi(x)T(x \rightarrow x') &= \sum_x \pi(x')T(x' \rightarrow x) \\ &= \pi(x') \underbrace{\sum_x T(x' \rightarrow x)}_1 \end{aligned}$$

Detailed Balance

If $\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$

Then $\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$

Proof
$$\begin{aligned} \sum_x \pi(x)T(x \rightarrow x') &= \sum_x \pi(x')T(x' \rightarrow x) \\ &= \pi(x') \sum_x T(x' \rightarrow x) \\ &= \pi(x') \end{aligned}$$

Metropolis-Hastings

For $k = 1, 2, \dots$

- Sample x' from a **wrong** $Q(x^k \rightarrow x')$
- Accept proposal x' with probability $A(x^k \rightarrow x')$
- Otherwise stay at x^k

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

$$T(x' \rightarrow x') = Q(x' \rightarrow x')$$

How to choose A:

$$\pi(x') = \sum_x \pi(x)T(x \rightarrow x')$$

Metropolis-Hastings

For $k = 1, 2, \dots$

- Sample x' from a **wrong** $Q(x^k \rightarrow x')$
- Accept proposal x' with probability $A(x^k \rightarrow x')$
- Otherwise stay at x^k

$$x^{k+1} = x^k$$

$$T(x \rightarrow x') = Q(x \rightarrow x')A(x \rightarrow x') \quad \text{for all } x \neq x'$$

$$T(x' \rightarrow x) = Q(x' \rightarrow x)$$

How to choose A:

$$\pi(x)T(x \rightarrow x') = \pi(x')T(x' \rightarrow x)$$