

Markov Chain Monte Carlo (MCMC)

- MCMC — silver bullet of probabilistic modeling

Markov Chain Monte Carlo (MCMC)

- MCMC — silver bullet of probabilistic modeling
- Learn how exploit specifics of your problem to speed up MCMC

Markov Chain Monte Carlo (MCMC)

- MCMC — silver bullet of probabilistic modeling
- Learn how exploit specifics of your problem to speed up MCMC
- Understand the limitations

Monte Carlo




Monte Carlo



Monte Carlo


Estimate expected values by sampling



Monte Carlo

Estimate expected values by sampling

$$\frac{\pi}{4} = \mathbb{E}[x^2 + y^2 \leq 1]$$




Monte Carlo

Estimate expected values by sampling

$$\frac{\pi}{4} = \mathbb{E}[x^2 + y^2 \leq 1]$$

$$\approx \frac{1}{M} \sum_{s=1}^M [x_s^2 + y_s^2 \leq 1]$$

$$x_s, y_s \sim \mathcal{U}(0, 1)$$



Monte Carlo

Estimate expected values by sampling

$$\mathbb{E}_{p(x)} f(x)$$

Monte Carlo

Estimate expected values by sampling

$$\mathbb{E}_{p(x)} f(x) \approx \frac{1}{M} \sum_{s=1}^M f(x_s)$$

$$x_s \sim p(x)$$

Monte Carlo

Why do we need to estimate expected values?

Monte Carlo

Why do we need to estimate expected values?

- Full Bayesian inference (see Week 1)

$$p(y \mid x, Y_{\text{train}}, X_{\text{train}})$$

Monte Carlo

Why do we need to estimate expected values?

- Full Bayesian inference (see Week 1)

$$\begin{aligned} p(y \mid x, Y_{\text{train}}, X_{\text{train}}) \\ = \int p(y \mid x, w) p(w \mid Y_{\text{train}}, X_{\text{train}}) dw \end{aligned}$$

Monte Carlo

Why do we need to estimate expected values?

- Full Bayesian inference (see Week 1)

$$\begin{aligned} & p(y \mid x, Y_{\text{train}}, X_{\text{train}}) \\ &= \int p(y \mid x, w) p(w \mid Y_{\text{train}}, X_{\text{train}}) dw \\ &= \mathbb{E}_{p(w \mid Y_{\text{train}}, X_{\text{train}})} p(y \mid x, w) \end{aligned}$$

Monte Carlo

Why do we need to estimate expected values?

- Full Bayesian inference (see Week 1)

$$\begin{aligned} p(y \mid x, Y_{\text{train}}, X_{\text{train}}) \\ &= \int p(y \mid x, w) p(w \mid Y_{\text{train}}, X_{\text{train}}) dw \\ &= \mathbb{E}_{p(w \mid Y_{\text{train}}, X_{\text{train}})} p(y \mid x, w) \end{aligned}$$

$$p(w \mid Y_{\text{train}}, X_{\text{train}}) = \frac{p(Y_{\text{train}} \mid X_{\text{train}}, w) p(w)}{Z}$$

Monte Carlo

Why do we need to estimate expected values?

- Full Bayesian inference (see Week 1)
- M-step of EM-algorithm (see Week 2)

Monte Carlo

Why do we need to estimate expected values?

- Full Bayesian inference (see Week 1)
- M-step of EM-algorithm (see Week 2)

$$\max_{\theta} \mathbb{E}_q \log p(X, T \mid \theta)$$