[College logo]

[University name]

[College name]

Health Anxiety, AI Chatbot Use and Cyberchondria Among Adults in Baghdad: An Online Cross-Sectional Survey

A graduation research project submitted in partial fulfilment of the requirements for the degree of [Degree]

Prepared by:

[Student name 1] · [Student name 2] · [Student name 3]

Supervised by:

[Supervisor name and title]

[Academic year]

Acknowledgment

[To be completed by the authors.]

We thank all the participants who gave their time to complete this survey, and [names of supervisor, department and any others to be acknowledged].

Table of Contents

Summary / Abstract.....viii
1. Introduction.....1
1.1 Background.....1
1.2 Health anxiety.....1
1.3 Cyberchondria.....2
1.4 The relationship between health anxiety and cyberchondria.....3
1.5 From search engines to AI chatbots.....4
1.6 Online searching and the doctor–patient relationship.....5
1.7 Rationale.....5
1.8 Aim and objectives.....6
2. Method.....7
2.1 Study design and setting.....7
2.2 Participants and sampling.....8
2.3 Sample size.....8
2.4 Data collection instrument.....9
2.5 Variables and operational definitions.....10
2.6 Statistical analysis.....11
2.7 Ethical considerations.....14
3. Results.....14
3.1 Characteristics of participants.....14
3.2 Patterns of online health information seeking.....16
3.3 Pre-consultation behaviour and the doctor–patient relationship.....17
3.4 Levels of health anxiety and cyberchondria, and psychometric properties.....18
3.5 Association between health anxiety and cyberchondria.....20
3.6 Factors associated with health anxiety and cyberchondria (bivariate analysis).....22
3.7 Independent predictors (multivariable analysis).....25
3.8 Health anxiety, cyberchondria and pre-consultation behaviour.....27
3.9 Pathways: mediation and moderation.....28
3.10 Sensitivity analyses.....30
4. Discussion.....30
4.1 Principal findings.....30
4.2 Health anxiety and cyberchondria.....31
4.3 Do health anxiety and cyberchondria fuel each other?.....32

4.4 The role of AI chatbots.....33
4.5 Pre-consultation behaviour and the doctor–patient relationship.....34
4.6 Medical and health-science students.....35
4.7 Comparison with regional studies.....35
4.8 Strengths and limitations.....36
5. Conclusions and Recommendations.....38
5.1 Conclusions.....38
5.2 Recommendations.....39
6. References.....41
Appendices.....47
Appendix A. Item-by-item correlations between the SHAI-14 and CSS-12.....47
Appendix B. Study questionnaire.....48

List of Abbreviations

AI	Artificial intelligence
AOR	Adjusted odds ratio
AUC	Area under the receiver operating characteristic curve
CFA	Confirmatory factor analysis
CFI	Comparative fit index
CI	Confidence interval
CSS-12	Cyberchondria Severity Scale, 12-item short form
DWLS	Diagonally weighted least squares
ER	Emergency room
FDR	False discovery rate
GP	General practitioner
HA	Health anxiety
HC3	Heteroskedasticity-consistent standard errors, type 3
IQR	Interquartile range
KMO	Kaiser–Meyer–Olkin measure of sampling adequacy
LLM	Large language model
ML	Maximum likelihood
OLS	Ordinary least squares
OR	Odds ratio
RMSEA	Root mean square error of approximation
SD	Standard deviation
SHAI-14	Short Health Anxiety Inventory, 14-item version
STROBE	Strengthening the Reporting of Observational Studies in Epidemiology
TLI	Tucker–Lewis index
VIF	Variance inflation factor

List of Tables

Table 1. Socio-demographic characteristics of the participants (n = 300).....15

Table 2. Online health information seeking, pre-consultation behaviour and relationship with physicians (n = 300).....16

Table 3. Descriptive statistics, reliability and language equivalence of the SHAI-14 and CSS-12 (n = 300).....19

Table 4. Prevalence of elevated health anxiety at different SHAI-14 cut-offs (n = 300).....19

Table 5. Measurement invariance of the CSS-12 and SHAI-14 across Arabic and English responders (multi-group CFA).....19

Table 6. Correlation between health anxiety (SHAI-14) and cyberchondria (CSS-12) and its dimensions (n = 300).....20

Table 7. Health anxiety (SHAI-14) and cyberchondria (CSS-12) scores by participant characteristics (n = 300).....23

Table 8. Variables significantly associated ($p < 0.05$) with elevated health anxiety and with high cyberchondria.....25

Table 9. Hierarchical robust linear regression of cyberchondria (CSS-12 total) – final model (n = 300).....26

Table 10. Multivariable logistic regression of elevated health anxiety (SHAI ≥ 18; 57 events, n = 300).....26

Table 11. Adjusted associations of health anxiety and cyberchondria with pre-consultation behaviours and the doctor–patient relationship.....28

Table 12. Exploratory mediation models decomposing the association between health anxiety (HA) and the dimensions of cyberchondria (standardised coefficients, n = 300).....28

Table 13. Sensitivity analyses of the main findings.....30

List of Figures

Figure 1. Conceptual framework of the study. Health anxiety is hypothesised to be associated with cyberchondria (H1) independently of covariates (H2), partly through excessive searching and post-search distress (H3), and cyberchondria with pre-consultation behaviour and the doctor–patient relationship (H4). The dashed arrow shows the short-lived relief that maintains the cycle.....7

Figure 2. Study flowchart: questionnaire development, data collection, data preparation and stages of statistical analysis.....14

Figure 3. Tools used and motivations for searching for health information online (n = 300).....17

Figure 4. Behavioural responses after searching, disclosure of searches to physicians and perceived effect on the doctor–patient relationship (n = 300).....18

Figure 5. Scatter plot of health anxiety (SHAI-14) against cyberchondria (CSS-12) with the least-squares line and 95% confidence band (n = 300; points slightly jittered).....21

Figure 6. Spearman correlation matrix of health anxiety, cyberchondria and its dimensions, trust in online/AI information, number of tools, number of motives and age group (FDR-adjusted significance).....22

Figure 7. Cyberchondria (CSS-12) by online tool profile.....24

Figure 8. Adjusted odds ratios for elevated health anxiety (SHAI ≥ 18). Orange: $p < 0.05$.....27

Figure 9. Mediation models: (A) excessive searching as a pathway from health anxiety to post-search distress; (B) post-search distress as a pathway from health anxiety to reassurance seeking.....29

Figure 10. Cyberchondria dimensions by health-anxiety status (mean ± 95% CI).. 29

Figure 11. Item-by-item correlations between SHAI-14 and CSS-12 items (Spearman ρ). * significant after FDR correction across the 168 pairs; ** also significant after Holm–Bonferroni correction.....47

Summary / Abstract

Background: Adults increasingly consult search engines and, more recently, artificial-intelligence (AI) chatbots before seeing a doctor. Health anxiety and cyberchondria (excessive, distressing online health searching) are thought to reinforce each other, but evidence from Iraq, and on AI chatbots, is scarce.

Objectives: To measure the association between health anxiety and cyberchondria among adults in Baghdad, and to describe online and AI health-information seeking and pre-consultation behaviour.

Methods: Cross-sectional online survey of 300 adults in Baghdad (July–September 2026) using the Short Health Anxiety Inventory (SHAI-14) and the Cyberchondria Severity Scale (CSS-12). Analyses included measurement-invariance testing across Arabic and English responders, FDR-corrected group comparisons, robust regression and bootstrapped mediation.

Results: Participants were mostly young (62.3% aged 20–29) and 63.3% had a health-related background; the CSS-12 was invariant across Arabic and English responders. AI chatbots were ticked by 78.0% (88.7% including respondents who named a chatbot), against 56.3% for search engines. Health anxiety correlated with cyberchondria ($r = 0.47$, 95% CI 0.38–0.55) and remained its strongest predictor after adjustment ($\beta = 0.48$), most strongly through excessive searching and post-search distress. Elevated health anxiety (SHAI-14 ≥ 18, a screening threshold rather than a diagnosis) was found in 19.0%; only 0.7% reached the stricter low-prevalence cut-off of 29. After searching, 34.7% delayed or skipped care because they felt reassured and 31.7% never told their doctor, citing fear of judgement; higher cyberchondria was associated with more disclosure. The association between AI use and cyberchondria was not robust to how AI use was defined.

Conclusion: In this convenience sample of mostly young, health-oriented adults, health anxiety and cyberchondria were linked, and AI chatbots were a major channel for online health information, without reliable evidence that their use adds to cyberchondria beyond health anxiety. The cross-sectional design cannot establish direction. Clinicians should ask about online and AI searches without judgement.

Keywords: cyberchondria; health anxiety; artificial intelligence; chatbots; online health information seeking; Iraq.

1. Introduction

1.1 Background

Searching the Internet before seeing a doctor has become an ordinary part of being ill. A symptom is now often typed into a search engine, or described to an AI chatbot, long before it is described to a physician. This shift has clear benefits: information is immediate, free and available at any hour, and it can help people judge whether a symptom needs medical attention. It also has a well-documented downside. For a sizeable minority of people, online searching does not settle a worry but amplifies it, a phenomenon that has come to be called *cyberchondria* [1,2].

Iraq offers a useful and under-studied setting in which to examine this. Internet access in the country has expanded rapidly, with roughly four in five Iraqis online by the start of 2025 [3], and in such a highly connected population the Internet and, increasingly, AI chatbots may function as a first point of contact for health concerns. How this behaviour interacts with health anxiety, and what it means for the consultation that follows, has not been examined in Baghdad.

1.2 Health anxiety

Health anxiety refers to a persistent preoccupation with having, or acquiring, a serious illness, maintained by misinterpretation of normal bodily sensations. It lies on a continuum, from transient worry that most people experience to clinically significant illness anxiety disorder and somatic symptom disorder in the DSM-5 [4]. The cognitive-behavioural model proposes that bodily sensations are interpreted as signs of serious disease and that the resulting anxiety is maintained by safety behaviours such as body checking and reassurance seeking: these behaviours relieve anxiety briefly but

prevent the person from learning that the feared outcome does not occur, so the worry returns [5].

Health anxiety is not rare: its lifetime prevalence is estimated at about 6% in the general population and up to 20% among hospital outpatients, and it generates substantial costs through unnecessary medical contacts [6,7]. A meta-analysis of university student samples found that health anxiety scores rose steadily between 1985 and 2017 [8]. In Iraq, almost half of 1591 respondents in an online survey during the COVID-19 restrictions reported health anxiety [9], and a second national online survey of 1578 Iraqi adults found that home quarantine was accompanied by greater health anxiety and increased reassurance-seeking behaviour [10], suggesting that the population is far from immune to illness worry.

1.3 Cyberchondria

The term cyberchondria was popularised by White and Horvitz, whose analysis of large-scale search logs showed that searches for common, benign symptoms frequently escalated towards rare and serious diagnoses, and that this escalation was driven by the amount and wording of medical content viewed and by the user's own predisposition to escalate [1]. Starcevic and Berle later defined cyberchondria as excessive or repeated online searching for health information that is associated with increasing, rather than relieving, distress or anxiety [2]. Crucially, searching for health information is not itself pathological; it becomes cyberchondria when it is excessive, compulsive and distressing, and when it drives further reassurance seeking [11]. A systematic review described cyberchondria as a possible transdiagnostic, compulsive digital syndrome closely linked to health anxiety and obsessive-compulsive symptoms, while noting that the evidence base was almost entirely cross-sectional and based on online student samples [12].

The most widely used instrument is the Cyberchondria Severity Scale (CSS), originally a 33-item measure with five factors: compulsion, distress, excessiveness, reassurance seeking and mistrust of medical professionals [13]. Its 12-item short form (CSS-12) drops the mistrust dimension and retains four three-item subscales that load substantially on a general cyberchondria factor, supporting both a total and subscale scores [14]. An Arabic version of the CSS-12 has shown acceptable structure and validity among Lebanese adults [15].

1.4 The relationship between health anxiety and cyberchondria

Cyberchondria and health anxiety are closely related yet distinct: in a community sample, cyberchondria overlapped strongly with health anxiety but, beyond that overlap, was associated with greater functional impairment and healthcare utilisation [16]. Searching can also raise anxiety directly: in a randomised experiment, participants who searched Google or a medical app for the causes of experimentally provoked bodily symptoms reported more negative affect, more health anxiety and a stronger wish to consult a physician than controls who waited [17]. Two meta-analyses have quantified the association: health anxiety correlates moderately with general online health information seeking ($r = 0.34$) but strongly with cyberchondria ($r = 0.62$) [18], and a later meta-analysis reported a similar pooled correlation ($r = 0.63$), with subscale correlations ranging from 0.24 to 0.66 [19]. The same work frames health-related Internet use in anxious individuals as a safety behaviour maintained by intermittent reinforcement [19]. Intolerance of uncertainty is also strongly associated with cyberchondria, already in adolescence ($r = 0.55$), presumably because search results rarely deliver a single, certain answer [20].

Whether the two conditions "fuel each other" is the central question. The only large longitudinal study, a four-wave Dutch panel of 5322 adults, found initial evidence of a reciprocal relationship in which health-anxious people

search more and searching in turn raises health anxiety; however, this loop appeared among people with non-clinical health anxiety and not in the clinically anxious subsample, in whom searching seemed to maintain rather than worsen anxiety [21]. A more recent three-wave study of pregnant women likewise identified health anxiety as a prospective risk factor for cyberchondria [22]. The COVID-19 pandemic offered a natural test of the cycle: cyberchondria tracked rising concern and safety behaviours in Croatia [23], and was linked with health anxiety among university students in Turkey [24]. A recent scoping review reported cyberchondria prevalence estimates between 30.7% and 55.6% across studies and consistent associations with health anxiety [25].

1.5 From search engines to AI chatbots

Most of this evidence was gathered when "searching online" meant using a search engine. Large language model (LLM) chatbots such as ChatGPT and Gemini change the experience in important ways: they answer in fluent, personalised, conversational language; they can be questioned repeatedly; and their responses can appear more empathetic than those of physicians [26,27]. In a large US survey, about one in five respondents had used ChatGPT for health information; two-thirds of these users reported scepticism about its output and two-thirds turned to their physicians, yet about a third changed medications or requested referrals based on its answers [28]. Health anxiety has also been identified as a stable predictor of symptom-checker app use, with over half of users meeting a hypochondria classification [29]. In a representative Czech sample of 4775 Internet users, health anxiety was associated with willingness to use generative AI for health information only indirectly, through greater trust in AI [30], and among young dermatology outpatients in Turkey 61% had consulted an AI chatbot (mostly ChatGPT) before their visit, without a detectable increase in anxiety [31]. Whether AI

chatbots relieve worry, by offering calm and contextualised answers, or amplify it, by being endlessly available for another round of reassurance, is an open empirical question.

1.6 Online searching and the doctor–patient relationship

What patients do after searching matters for clinical care. Online information can improve the doctor–patient relationship when patients discuss it with their physician, but can strain it when it is withheld or used to challenge the physician [32,33]. Reassurance seeking may translate into additional consultations, requests for investigations or "doctor shopping", while false reassurance may lead to delayed care. Studies in the Arab region have begun to describe these patterns: cyberchondria was common among adults in Jeddah and associated with smartphone addiction [34], and among Saudi primary-care patients cyberchondria was linked with health anxiety [35]. Across 13 Arabic-speaking countries of the Middle East and North Africa, about a third of 4276 migraine patients had delayed consultation or modified treatment because of online information, and higher trust in online information strongly predicted delayed consultation [36]. No comparable study has been reported from Iraq, and none, to our knowledge, has placed AI chatbots alongside search engines in this population.

1.7 Rationale

This study addresses three gaps. First, it provides, to our knowledge, the first estimate of the association between health anxiety and cyberchondria (using the CSS-12) in a Baghdad-based sample. Second, it examines AI chatbots as a distinct, and now dominant, source of online health information, rather than treating "the Internet" as one entity. Third, it links these psychological constructs to concrete pre-consultation behaviours, namely delaying care, self-medication, urgent visits, multiple consultations,

disclosure of searches and trust in physicians, that matter directly to clinicians.

1.8 Aim and objectives

Aim: To study the relationship between health anxiety and cyberchondria and their correlates among adults in Baghdad who search for health information online, with particular attention to AI chatbots and pre-consultation behaviour.

Specific objectives:

1. To describe online and AI-based health-information seeking and pre-consultation behaviour among adults in Baghdad.
2. To estimate levels of health anxiety and cyberchondria and the strength of their association, overall and by cyberchondria dimension, independently of socio-demographic and digital-use factors (H1, H2); correlates of each construct were examined as secondary analyses.
3. To examine statistical pathways consistent with a mutually reinforcing cycle, and factors that might modify the association (H3).
4. To examine how health anxiety and cyberchondria relate to pre-consultation behaviour, disclosure of searches to physicians and trust in physicians (H4).

Hypotheses: (H1) Health anxiety is positively associated with cyberchondria, most strongly with its distress and excessiveness dimensions. (H2) The association is independent of socio-demographic and digital-use factors. (H3) Excessive searching statistically mediates the link between health anxiety and post-search distress. (H4) Higher cyberchondria is associated with care-seeking behaviours (urgent visits, multiple consultations) and with a more strained doctor–patient relationship. H1 and H2 were the primary, confirmatory hypotheses and H3 and H4 secondary; all other analyses,

including the AI-use comparisons, item-level, moderation and sensitivity analyses, were exploratory.

The conceptual framework of the study, linking these hypotheses to the proposed cycle between health anxiety and the facets of cyberchondria, is shown in Figure 1.




Figure 1. *Conceptual framework of the study. Health anxiety is hypothesised to be associated with cyberchondria (H1) independently of covariates (H2), partly through excessive searching and post-search distress (H3), and cyberchondria with pre-consultation behaviour and the doctor–patient relationship (H4). The dashed arrow shows the short-lived relief that maintains the cycle.*

2. Method

2.1 Study design and setting

This was an analytical cross-sectional study reported in line with the STROBE statement for observational studies [37] and the CHERRIES checklist for web-based surveys [38]. Data were collected online in Baghdad, Iraq, between 17 July 2026 and 11 September 2026.

2.2 Participants and sampling

A non-probability (convenience and snowball) sampling approach was used. The survey link was distributed through online channels (social media and messaging groups) whose members were based in Baghdad. Inclusion criteria were: age 18 years or older, ability to read Arabic or English, and provision of informed consent. Every completed submission was retained for analysis. No duplicate response patterns were found. [Authors to state whether Google Forms limited submissions to one per account.] Because the link spread by snowball sampling, the number of people who saw the invitation is unknown, so view and participation rates could not be computed. Three responses had been annotated by the team during data collection [authors to state the reason for each annotation]; they were kept, and their exclusion was tested in a sensitivity analysis. The resulting sample was predominantly young and from health-related fields (Section 3.1).

2.3 Sample size

The study was powered for its primary objective, the correlation between health anxiety and cyberchondria. Using Fisher's z approximation with two-sided $\alpha = 0.05$ and 80% power, a minimum of 194 participants is required to detect a small-to-moderate correlation ($r = 0.20$), and 85 to detect $r = 0.30$ [39]. Meta-analytic estimates of this association are far larger ($r \approx 0.62$) [18,19], so these targets are conservative. For descriptive proportions, 300 participants give a margin of error of ±5.7% at 95% confidence (worst case, $p = 0.5$), compared with ±5% for the 385 participants of a Cochran-based calculation. The logistic model for elevated health anxiety had 57 events for 11 predictors (≈5 events per variable), which is acknowledged as a limitation of that model.

2.4 Data collection instrument

A bilingual (Arabic–English) self-administered questionnaire was built in Google Forms. All items were required, except the free-text description of chronic disease and the name of the AI chatbot used, which were conditional. The questionnaire had five parts:

1. **Consent and socio-demographic data:** confirmation of age ≥18 and consent; age group; gender; highest education; nature of study or work; diagnosed chronic medical condition (and its type).
2. **Digital health information seeking:** tools used (search engines, AI chatbots, dedicated medical platforms, social media; multiple answers); main AI chatbot; motivation for searching (multiple answers); and trust in medical information generated by online tools/AI (5-point scale).
3. **Health anxiety:** the 14-item Short Health Anxiety Inventory (SHAI-14) [40]. Each item offers four statements scored 0–3, giving a total of 0–42; higher scores indicate greater health anxiety. The SHAI has good reliability and validity in clinical and non-clinical samples, including recent validations in Spain, Sweden and Iran [41–43], and an Arabic version has been validated in Saudi Arabia [44].
4. **Cyberchondria:** the CSS-12 [14], with items rated on a 5-point frequency scale (1 = never to 5 = always), giving a total of 12–60 and four subscales (3–15 each): Excessiveness (items 1–3), Compulsion (4–6), Distress (7–9) and Reassurance seeking (10–12).
5. **Pre-consultation behaviour and relationship with physicians:** behavioural responses after searching (multiple answers), whether search or AI results are disclosed to the physician, the perceived effect of digital information seeking on the relationship with and trust in physicians, and the language mainly relied upon while answering.

Each item was presented in English and Arabic. The research team produced the Arabic wording; its equivalence was examined empirically by comparing the psychometric behaviour of the scales between participants who relied mainly on Arabic and those who relied mainly on English (Section 2.6). For one SHAI item (item 2, noticing aches and pains) the four statements were displayed in a different order from the original scale ("as much as most people" before "less than most people"); all other SHAI and CSS items appeared in their original order. Items were scored according to the meaning of the chosen statement, following the original scoring key. The Arabic wording was produced by the research team rather than through formal cross-cultural adaptation [45], and the published Arabic versions of the CSS-12 and SHAI [15,44,46] were not used. As in the original scale, CSS item 3 refers to entering symptoms into a search engine.

2.5 Variables and operational definitions

- **Elevated health anxiety:** SHAI-14 ≥ 18, the optimal cut-off for distinguishing people with health anxiety from healthy controls in a Swedish diagnostic-accuracy study [42]. Because the same study recommends higher cut-offs in psychiatric (22) and low-prevalence (29) settings, the term "elevated" rather than "clinically relevant" is used, it is not a diagnosis, and prevalence at 15, 22 and 29 is also reported
- **High cyberchondria:** CSS-12 total in the upper tertile of the sample (> 33), since the CSS-12 has no validated clinical cut-off.
- **AI-chatbot user:** ticked "AI chatbots" as a tool or named an AI chatbot in the follow-up question. Thirty-two respondents named a chatbot without ticking the tool option; they were classified as users (the follow-up question was optional and specifically addressed to chatbot users). Because this decision affects the results, AI use is reported both ways

(ticked only, and ticked or named), and all analyses involving AI use were repeated with the stricter definition.

- **Health-related background:** medical students, students in other health fields and healthcare workers.
- **Occupation:** free-text "Other" answers were recoded into the closest category (e.g. secondary-school students and school leavers as non-health students; housewives, engineers and business owners as non-health employed/other). Age groups 40–49, 50–59 and 60+ were merged (≥40) for inferential analysis because of small cell sizes.

2.6 Statistical analysis

Analyses were carried out in Python 3 (pandas, SciPy, statsmodels, pingouin, factor_analyzer and semopy). The analysis code is fully scripted and reproducible. All tests were two-sided with $\alpha = 0.05$.

- **Descriptive statistics:** frequencies and percentages for categorical variables; mean ± SD and median (IQR) for scores. Normality was assessed with the Shapiro–Wilk test, skewness and kurtosis.
- **Psychometrics:** internal consistency by Cronbach's α (with 95% CI) and McDonald's ω [47]; corrected item–rest correlations; sampling adequacy (KMO, Bartlett's test); and confirmatory factor analysis by maximum likelihood (ML), with fit judged by CFI, TLI, RMSEA and SRMR against conventional criteria (CFI/TLI ≥ 0.95, RMSEA ≤ 0.06–0.08, SRMR ≤ 0.08), recognising that fixed cut-offs do not transfer exactly to every model [48]. Diagonally weighted least squares (DWLS) estimates on Pearson covariances are reported for comparison only, because their fit indices are not comparable with these cut-offs. Equivalence of the Arabic and English responders was tested by comparing α values with Feldt's test, factor-loading similarity with Tucker's congruence coefficient [49] (an index of approximate structural similarity) and adjusted mean differences.

Formal measurement invariance was then tested by multi-group CFA (ML with a mean structure), fitting configural, metric (equal loadings) and scalar (equal loadings and intercepts) models in sequence; each step was judged by the χ^2 difference test and by a change in CFI of no more than −0.01 and in RMSEA of no more than 0.015 [50]. Differential item functioning (DIF) was examined for every item by ordinal logistic regression of the item on the scale total, language and their interaction, a standard approach for ordinal items [51], with FDR correction across the 26 items.

- **Correlation analysis:** Pearson and Spearman correlations (bootstrapped 95% CIs) and partial correlations adjusted for covariates. Differences between dependent correlations (health anxiety with each CSS subscale) were tested with Williams' T2 test for dependent correlations. Item-level correlations were computed for every SHAI item with the CSS total and every CSS item with the SHAI total.
- **Group comparisons for every variable:** Mann–Whitney U (rank-biserial r as effect size) or Kruskal–Wallis H (ε^2 as effect size) for SHAI, CSS and each CSS subscale, with Welch's t-test/ANOVA as parametric confirmation and pairwise Mann–Whitney post-hoc comparisons with Holm adjustment. Associations of all categorical variables with elevated health anxiety and high cyberchondria were tested with Pearson χ^2 or Fisher's exact test (Monte-Carlo exact p when expected counts were small), with Cramér's V. Pairwise associations among all categorical variables were also screened.
- **Multiple testing:** because many associations were tested, p-values within each family of tests were adjusted with the Benjamini–Hochberg false discovery rate procedure (q-values) [52]. The families were: the 162 group comparisons; the chi-square tests; the pairwise associations among categorical variables; the 40 coefficients of the behaviour models; the

moderation terms; and the 168 item-level correlations. Unadjusted p-values are reported, and associations that do not survive correction are labelled as exploratory.

- **Multivariable models:** hierarchical OLS regression of CSS on socio-demographics (block 1), digital-use variables (block 2) and SHAI (block 3), with HC3 robust standard errors, standardised coefficients and VIF checks; the reverse model (SHAI on CSS); models for each CSS subscale; quantile regression (25th–90th percentiles) and a quadratic term to examine non-linearity. Logistic regression identified predictors of elevated health anxiety (AORs; apparent AUC and AUC corrected for optimism by 500 bootstrap refits) and of each pre-consultation behaviour; ordinal logistic regression was used for the ordered disclosure and relationship outcomes.
- **Mediation and moderation:** as an exploratory decomposition of the association into CSS facets, mediation models with bootstrap confidence intervals (5000 resamples) estimated how much of the link between health anxiety and post-search distress ran through excessive searching, and related pathways, including a reverse-direction model and a serial model [53]. Moderation of the SHAI–CSS association by trust, AI use, health background, gender, chronic disease, language and age was tested with interaction terms and simple slopes.
- **Sensitivity analyses:** exclusion of annotated responses; exclusion of straight-line responders (identical answers to all items of a scale); alternative SHAI cut-offs; bootstrap of the main regression coefficient; and comparison of early versus late responders as a proxy for non-response bias.

The flow of the study, from questionnaire development to the stages of analysis, is summarised in Figure 2.




Figure 2. *Study flowchart: questionnaire development, data collection, data preparation and stages of statistical analysis.*

2.7 Ethical considerations

Participation was voluntary and anonymous. The first page of the form explained the purpose of the study, and electronic informed consent (with confirmation of age ≥18) was required before any question was shown; participants could withdraw at any time by closing the form. No names or contact details were collected, and data were stored securely and used only for research purposes. [Ethical approval: committee name, approval number and date to be inserted.]

3. Results

3.1 Characteristics of participants

A total of 300 adults completed the questionnaire; all gave consent and no item had missing data. The sample was young, with 187 (62.3%) aged 20–

29 and 56 (18.7%) aged 18–19; 169 (56.3%) were female. Most were currently students (189 (63.0%)); medical students were the largest occupational group (132 (44.0%)), and 190 (63.3%) had a health-related background. A diagnosed chronic condition was reported by 65 (21.7%), most commonly migraine, hypertension, asthma, anaemia, psoriasis and diabetes. Most participants relied mainly on Arabic when answering (212 (70.7%)) (Table 1).

Table 1. Socio-demographic characteristics of the participants (n = 300)

Characteristic	n	%
Age group (years)		
18-19	56	18.7
20-29	187	62.3
30-39	21	7.0
40-49	25	8.3
50-59	9	3.0
60+	2	0.7
Gender		
Female	169	56.3
Male	131	43.7
Highest education		
Currently a student	189	63.0
High school or below	18	6.0
Diploma/Vocational	7	2.3
Bachelor	65	21.7
Postgraduate	21	7.0
Study/work		
Medical student	132	44.0
Other health-science student	43	14.3
Healthcare worker	15	5.0
Non-health student	46	15.3
Non-health employee/other	64	21.3
Health-related background (1=yes)		
Yes	190	63.3
No	110	36.7
Diagnosed chronic condition (1=yes)		
Yes	65	21.7
No	235	78.3
Language relied on		
Arabic	212	70.7
English	88	29.3

3.2 Patterns of online health information seeking

AI chatbots were the most frequently used tool for looking up symptoms: 234 (78.0%) ticked them, or 266 (88.7%) when the 32 respondents who named a chatbot without ticking the option are included, ahead of search engines (169 (56.3%)). Notably, 123 (41.0%) used AI chatbots without any search engine, and ChatGPT was the dominant chatbot (178 (59.3%)). AI use fell with age, from 96% at 18–19 years to 72% at 40 years or older, so the 34 non-users were an older group. The leading motives were convenience (80.0%), deciding whether a doctor visit was needed (57.7%) and instant reassurance (51.0%), and trust in online/AI medical information was mostly moderate (Table 2, Figure 3).

Table 2. Online health information seeking, pre-consultation behaviour and relationship with physicians (n = 300)

Variable	n	%
Tools used (multiple)		
AI chatbots*	266	88.7
Search engines	169	56.3
Dedicated medical platforms	55	18.3
Social media	32	10.7
Main AI chatbot		
ChatGPT	178	59.3
Gemini	71	23.7
Claude	12	4.0
Other AI	3	1.0
None reported	36	12.0
Tool profile		
AI + search engine	143	47.7
AI without search engine	123	41.0
No AI (search/other only)	34	11.3
Motivation (multiple)		
Convenience/accessibility	240	80.0
Check if doctor visit is needed	173	57.7
Instant reassurance	153	51.0
Self-diagnose / double-check prescription	69	23.0
Cost/logistic barriers	54	18.0
Trust in online/AI medical data (1-5)		
1 – Do not trust at all	20	6.7
2 – Slightly	83	27.7
3 – Moderately	170	56.7
4 – Very much	25	8.3
5 – Completely	2	0.7
Behaviour after searching (multiple)		

Delay/skip care (reassured)	104	34.7
Ignore symptom	90	30.0
Visit multiple doctors	76	25.3
Urgent visit/ER/fast diagnosis	66	22.0
Self-medication	64	21.3
Delay care (terrified/overwhelmed)	26	8.7
Disclose search to physician		
Always	81	27.0
Sometimes	124	41.3
Never (fear of judgement)	95	31.7
Effect on doctor relationship		
No negative impact	196	65.3
Somewhat complicated	66	22.0
Significantly reduced trust	38	12.7

Tools, motivations and behaviours allowed multiple answers, so percentages do not sum to 100. AI users include 32 respondents who named a chatbot without ticking the AI option.




Figure 3. *Tools used and motivations for searching for health information online (n = 300).*

3.3 Pre-consultation behaviour and the doctor–patient relationship

After searching, the most frequent response was to delay or skip care because the information was reassuring (34.7%), followed by ignoring the symptom (30.0%); others escalated, consulting several doctors (25.3%) or seeking urgent care (22.0%). Only 27.0% always told their physician about their searches, whereas 31.7% never did so, citing fear of being judged, and 34.7% said searching had complicated their relationship with, or reduced their trust in, their doctor (Figure 4).




Figure 4. *Behavioural responses after searching, disclosure of searches to physicians and perceived effect on the doctor–patient relationship (n = 300).*

3.4 Levels of health anxiety and cyberchondria, and psychometric properties

The mean SHAI-14 score was 12.35 ± 5.83 and 57 (19.0%) (95% CI 15.0–23.8%) scored at or above 18, indicating elevated health anxiety; the estimate depended strongly on the threshold, from 32.7% at 15 to 0.7% at 29 (Table 4). The mean CSS-12 score was 28.86 ± 8.36: participants searched a great deal (excessiveness was the highest-scoring dimension, 9.66 ± 2.94) but searching rarely disrupted daily life (compulsion was the lowest, 5.00 ± 2.58). Because scores deviated modestly from normality (Shapiro–Wilk $p < 0.01$), non-parametric bivariate tests and robust regression standard errors were used.

Both scales were reliable (SHAI-14 $\alpha = 0.81$, $\omega = 0.82$; CSS-12 $\alpha = 0.86$, $\omega = 0.87$; Table 3). Under ML, the four-factor structure of the CSS-12 fitted acceptably (CFI = 0.954, RMSEA = 0.070, SRMR = 0.053) and clearly better than a single factor, whereas the one-factor SHAI-14 model fitted poorly (CFI = 0.830), so the SHAI-14 should not be regarded as strictly unidimensional, consistent with the two-factor solutions reported in recent validations [41,43].

Measurement invariance across Arabic and English responders is summarised in Table 5. The CSS-12 reached full scalar invariance, so its scores can be compared directly across the two languages; the SHAI-14 reached metric but not scalar invariance (scalar $\Delta CFI = -0.018$). No item

showed differential item functioning after FDR correction (smallest q = 0.13), and language did not modify the association between health anxiety and cyberchondria (interaction p = 0.292).

Table 3. Descriptive statistics, reliability and language equivalence of the SHAI-14 and CSS-12 (n = 300)

Scale	Mean ± SD	Median (IQR)	Range	α (95% CI)	ω
SHAI-14 total (0–42)	12.35 ± 5.83	12 (8–16)	1–30	0.81 (0.78–0.84)	0.82
CSS-12 total (12–60)	28.86 ± 8.36	29 (23–34)	12–57	0.86 (0.84–0.89)	0.87
Excessiveness (3–15)	9.66 ± 2.94	10 (8–12)	3–15	0.77 (0.72–0.81)	0.77
Compulsion (3–15)	5.00 ± 2.58	4 (3–6)	3–15	0.88 (0.86–0.90)	0.89
Distress (3–15)	7.24 ± 2.74	7 (5–9)	3–15	0.76 (0.71–0.81)	0.77
Reassurance (3–15)	6.96 ± 2.81	7 (5–9)	3–15	0.76 (0.71–0.81)	0.77

Language equivalence: SHAI mean 12.40±5.73 (Arabic) vs 12.23±6.08 (English), adjusted p = 0.52; CSS mean 28.96±8.21 vs 28.62±8.77, adjusted p = 0.89. CFA fit, ML (primary): CSS-12 four factors CFI 0.954, TLI 0.937, RMSEA 0.070, SRMR 0.053; SHAI-14 one factor CFI 0.830, RMSEA 0.079, SRMR 0.064. DWLS on Pearson covariances (comparison only, not comparable with ML cut-offs): CSS-12 CFI 1.000, RMSEA 0.003; SHAI-14 CFI 0.977, RMSEA 0.038. Measurement invariance across languages: Table 5.

Table 4. Prevalence of elevated health anxiety at different SHAI-14 cut-offs (n = 300)

SHAI-14 score	Basis of the cut-off	n	%	95% CI
≥ 15	Lower threshold (sensitivity)	98	32.7	27.6–38.2
≥ 18	**Optimal versus healthy controls (primary definition)**	**57**	**19.0**	**15.0–23.8**
≥ 22	Optimal in a psychiatric setting	22	7.3	4.9–10.9
≥ 29	Optimal in a low-prevalence setting	2	0.7	0.2–2.4

Cut-offs of 18, 22 and 29 from the Swedish diagnostic-accuracy study of the SHAI-14 [42]; ≥ 18 is used throughout this thesis. 95% CI: Wilson score interval. An elevated score is a screening result, not a diagnosis.

Table 5. Measurement invariance of the CSS-12 and SHAI-14 across Arabic and English responders (multi-group CFA)

Model	χ^2 (df)	CFI	RMSEA	$\Delta\chi^2$ (Δdf)	p	ΔCFI	Decision
CSS-12 (four factors)							
Configural	195.1 (96)	0.937	0.083	—	—	—	Baseline
Metric (equal loadings)	200.7 (104)	0.939	0.079	5.52 (8)	0.701	+0.002	Supported
Scalar (+ intercepts)	209.1 (112)	0.938	0.076	8.41 (8)	0.395	+0.000	Supported
SHAI-14 (one factor)							
Configural	322.6 (154)	0.808	0.086	—	—	—	Baseline
Metric (equal loadings)	332.2 (167)	0.812	0.081	9.61 (13)	0.725	+0.004	Supported
Scalar (+ intercepts)	361.2 (180)	0.794	0.082	29.00 (13)	0.007	−0.018	Not supported

Maximum likelihood with a mean structure; Arabic n = 212, English n = 88. A step is supported when CFI does not fall by more than 0.01 and $\Delta\chi^2$ is non-significant [50]. Item-level DIF (ordinal logistic regression): no item significant after FDR correction (smallest q = 0.13).

3.5 Association between health anxiety and cyberchondria

Health anxiety was positively and moderately correlated with cyberchondria (Pearson r = 0.47, 95% CI 0.38–0.55; Spearman ρ = 0.47, bootstrap 95% CI 0.36–0.56; $p < 0.001$), with health anxiety accounting for 22% of the variance in cyberchondria (Figure 5). The association was essentially unchanged after adjustment for age, gender, health background, chronic disease, trust in online information and AI use (partial r = 0.48, 95% CI 0.38–0.56). Health anxiety was correlated with all four cyberchondria dimensions, but not equally: correlations with excessiveness (r = 0.46) and distress (r = 0.45) were similar to each other and significantly stronger than those with compulsion (r = 0.25) and reassurance seeking (r = 0.24) (Williams' t, all $p < 0.001$) (Table 6, Figure 6).

At item level, fear of serious illness and the interpretation of bodily sensations were most closely linked to cyberchondria, whereas simple awareness of bodily sensations and of aches and pains was not ($q > 0.10$); 111 of the 168 SHAI × CSS item pairs remained significant after FDR correction, the strongest being fear of serious illness with panic on reading about a rare or serious condition (ρ = 0.43; Appendix A, Figure 11). Trust in online/AI information was associated only with excessiveness (ρ = 0.19; Figure 6).

Table 6. Correlation between health anxiety (SHAI-14) and cyberchondria (CSS-12) and its dimensions (n = 300)

Cyberchondria measure	Pearson r (95% CI)	Spearman ρ	p	Adjusted β
CSS-12 total	0.47 (0.38–0.55)	0.47	<0.001	0.48
Excessiveness	0.46 (0.37–0.55)	0.45	<0.001	0.44
Compulsion	0.25 (0.14–0.36)	0.32	<0.001	0.29
Distress	0.45 (0.36–0.54)	0.44	<0.001	0.44
Reassurance	0.24 (0.13–0.35)	0.23	<0.001	0.25

Adjusted β: standardised coefficient of SHAI from robust OLS models adjusted for age, gender, occupation, chronic disease, language, tools, trust and motives. Excessiveness and Distress correlations were each significantly stronger than Compulsion and Reassurance (Williams' t, $p < 0.001$); they did not differ from each other ($p = 0.81$).




Figure 5. *Scatter plot of health anxiety (SHAI-14) against cyberchondria (CSS-12) with the least-squares line and 95% confidence band (n = 300; points slightly jittered).*




Figure 6. *Spearman correlation matrix of health anxiety, cyberchondria and its dimensions, trust in online/AI information, number of tools, number of motives and age group (FDR-adjusted significance).*

3.6 Factors associated with health anxiety and cyberchondria (bivariate analysis)

Every variable was tested against SHAI, CSS and each CSS dimension (162 tests; Table 7). Health anxiety was higher in participants with a chronic condition ($p = 0.004$, $q = 0.028$) and, most clearly, in those searching for instant reassurance (13.76 ± 5.65 vs 10.88 ± 5.65; $q = 0.002$). Differences by age, education and occupation were smaller and borderline after correction (age $q = 0.051$) and are treated as exploratory. Gender was unrelated to either score.

Under the inclusive definition, cyberchondria was higher among AI-chatbot users than non-users (29.49 ± 8.21 vs 23.97 ± 8.05; $q = 0.005$), mainly

in excessiveness and reassurance seeking, and was highest in those who combined AI with search engines (Figure 7); the specific chatbot made no difference ($p = 0.726$).

Table 7. Health anxiety (SHAI-14) and cyberchondria (CSS-12) scores by participant characteristics (n = 300)

Variable	n	SHAI mean ± SD	p (q)	CSS mean ± SD	p (q)
Age group			**0.008 (0.051)**		**0.288 (0.505)**
18-19	56	14.0 ± 5.5		28.6 ± 8.1	
20-29	187	12.4 ± 5.8		28.4 ± 8.0	
30-39	21	10.2 ± 6.4		32.5 ± 11.3	
≥40	36	11.0 ± 5.7		29.7 ± 8.2	
Gender			**0.378 (0.589)**		**0.859 (0.918)**
Female	169	12.7 ± 6.0		29.0 ± 8.7	
Male	131	11.9 ± 5.5		28.6 ± 8.0	
Study/work			**0.005 (0.039)**		**0.820 (0.918)**
Medical student	132	13.3 ± 5.8		28.9 ± 8.1	
Other health-science student	43	12.2 ± 5.5		29.2 ± 9.1	
Healthcare worker	15	10.4 ± 5.1		29.5 ± 7.4	
Non-health student	46	13.0 ± 6.0		27.6 ± 7.8	
Non-health employee/other	64	10.5 ± 5.7		29.4 ± 9.1	
Chronic condition			**0.004 (0.028)**		**0.435 (0.623)**
Yes	65	14.2 ± 6.0		29.9 ± 8.9	
No	235	11.9 ± 5.7		28.6 ± 8.2	
Uses AI chatbots			**0.008 (0.051)**		**<0.001 (0.005)**
Yes	266	12.6 ± 5.7		29.5 ± 8.2	
No	34	10.5 ± 6.7		24.0 ± 8.1	
Uses search engines			**0.111 (0.316)**		**0.113 (0.316)**
Yes	169	12.8 ± 5.9		29.4 ± 7.9	
No	131	11.8 ± 5.7		28.2 ± 8.9	
Uses social media			**0.001 (0.016)**		**0.021 (0.093)**
Yes	32	15.8 ± 6.9		32.5 ± 9.1	
No	268	11.9 ± 5.6		28.4 ± 8.2	
Tool profile			**0.006 (0.039)**		**<0.001 (0.005)**
AI + search engine	143	13.1 ± 5.6		30.1 ± 7.6	
AI without search engine	123	12.0 ± 5.8		28.7 ± 8.8	
No AI (search/other only)	34	10.5 ± 6.7		24.0 ± 8.1	
Motive: reassurance			**<0.001 (0.002)**		**0.085 (0.280)**
Yes	153	13.8 ± 5.7		29.6 ± 7.6	

No	147	10.9 ± 5.7		28.1 ± 9.1	
Motive: self-diagnosis			**0.217 (0.435)**		**0.079 (0.266)**
Yes	69	13.1 ± 6.2		30.4 ± 8.1	
No	231	12.1 ± 5.7		28.4 ± 8.4	
Trust in online/AI			**0.052 (0.190)**		**0.110 (0.316)**
1	20	10.1 ± 6.4		24.8 ± 7.8	
2	83	13.2 ± 6.1		28.9 ± 8.3	
3	170	11.9 ± 5.6		28.9 ± 8.4	
4	25	14.7 ± 5.2		31.8 ± 7.1	
5	2	13.0 ± 1.4		25.5 ± 17.7	
Language			**0.857 (0.918)**		**0.654 (0.802)**
Arabic	212	12.4 ± 5.7		29.0 ± 8.2	
English	88	12.2 ± 6.1		28.6 ± 8.8	

p: Mann–Whitney U (two groups) or Kruskal–Wallis H (three or more groups); q: Benjamini–Hochberg false-discovery-rate adjusted p across all 162 group comparisons. Effect sizes, medians (IQR), Welch tests and post-hoc comparisons are provided in the supplementary workbook.




Figure 7. *Cyberchondria (CSS-12) by online tool profile.*

In categorical analysis (Table 8), elevated health anxiety was more common among social-media users and among those seeking instant reassurance, and participants with elevated health anxiety had higher odds of high cyberchondria (52.6% vs 23.0%; OR = 3.71, 95% CI 2.04–6.76). High cyberchondria was most frequent among participants who always disclosed their searches to their physician (q = 0.015).

Table 8. Variables significantly associated ($p < 0.05$) with elevated health anxiety and with high cyberchondria

Variable	n/N (%) with outcome, by group	Cramér's V	p	q (FDR)
Elevated health anxiety (SHAI ≥ 18)				
Motive: reassurance	No: 15/147 (10.2%); Yes: 42/153 (27.5%)	0.22	<0.001	0.004
Uses social media	No: 44/268 (16.4%); Yes: 13/32 (40.6%)	0.19	0.001	0.015
Delay care (terrified)	No: 47/274 (17.2%); Yes: 10/26 (38.5%)	0.15	0.016	0.107
Motive: convenience	No: 17/60 (28.3%); Yes: 40/240 (16.7%)	0.12	0.039	0.180
High cyberchondria (CSS > 33)				
Elevated health anxiety	No: 56/243 (23.0%); Yes: 30/57 (52.6%)	0.26	<0.001	0.001
Disclose to physician	Always: 33/81 (40.7%); Never : 15/95 (15.8%); Sometimes: 38/124 (30.6%)	0.21	0.001	0.015
Ignore symptom	No: 70/210 (33.3%); Yes: 16/90 (17.8%)	0.16	0.006	0.060
Uses AI chatbots	No: 3/34 (8.8%); Yes: 83/266 (31.2%)	0.16	0.007	0.060
Tool profile	AI + search engine: 49/143 (34.3%); AI without search engine: 34/123 (27.6%); No AI (search/other only): 3/34 (8.8%)	0.17	0.012	0.096
Effect on doctor relationship	No negative impact: 46/196 (23.5%); Significantly reduced trust: 14/38 (36.8%); Somewhat complicated: 26/66 (39.4%)	0.16	0.023	0.141
Main AI chatbot	ChatGPT: 61/178 (34.3%); Claude: 4/12 (33.3%); Gemini: 17/71 (23.9%); None reported: 4/36 (11.1%); Other AI: 0/3 (0.0%)	0.18	0.034	0.180
Delay care (terrified)	No: 74/274 (27.0%); Yes: 12/26 (46.2%)	0.12	0.039	0.180

Pearson χ^2 or Fisher's exact test (Monte-Carlo exact p where expected counts were < 5). Associations with $q \geq 0.05$ are exploratory. Non-significant associations are listed in the supplementary workbook.

3.7 Independent predictors (multivariable analysis)

In the hierarchical regression (Table 9), socio-demographic variables explained little of the variance in cyberchondria ($R^2 = 0.030$); digital-use variables raised R^2 to 0.133, and health anxiety added a further 18.4% ($R^2 = 0.317$; $f^2 = 0.27$, a medium effect). Health anxiety was by far the strongest predictor (B = 0.68, 95% CI 0.53–0.84, $\beta = 0.48$, $p < 0.001$). AI-chatbot use was associated with a 5.3-point higher CSS score under the inclusive definition ($p < 0.001$), but not when only respondents who ticked "AI chatbots" were counted as users (B = 1.49, p = 0.314); the 32 reclassified respondents had the highest cyberchondria of any group (mean 31.6), so there is no reliable evidence that AI use is associated with cyberchondria independently of health anxiety. No other variable was robustly associated, and the model diagnostics supported the stability of the health-anxiety effect (Table 13).

In the reverse model, cyberchondria was the strongest predictor of health anxiety (β = 0.45, p < 0.001), together with chronic disease (B = 1.68, p = 0.029) and searching for instant reassurance (p < 0.001); after adjustment, medical students no longer differed from other groups.

Table 9. Hierarchical robust linear regression of cyberchondria (CSS-12 total) – final model (n = 300)

Predictor	B	95% CI	β	p
Constant	12.04	6.39–17.68		<0.001
Age 18–19 (ref. 20–29)	−0.43	−2.77–1.91	−0.02	0.721
Age 30–39	**6.27**	**0.19–12.34**	**0.19**	**0.043**
Age ≥40	3.98	−0.86–8.81	0.16	0.107
Healthcare worker (ref. non-health employed)	1.06	−4.32–6.44	0.03	0.700
Medical student	0.41	−3.67–4.49	0.02	0.845
Non-health student	−0.73	−5.12–3.66	−0.03	0.745
Other health-science student	1.32	−3.61–6.25	0.06	0.599
Female	−0.05	−1.82–1.72	0.00	0.954
Chronic condition	−0.14	−2.42–2.15	−0.01	0.906
Answered mainly in English	0.51	−1.71–2.74	0.03	0.650
Uses AI chatbots	**5.33**	**2.34–8.32**	**0.20**	**<0.001**
Uses search engines	0.95	−1.00–2.90	0.06	0.342
Uses medical platforms	0.17	−2.10–2.44	0.01	0.885
Uses social media	1.46	−1.61–4.54	0.05	0.351
Trust in online/AI (per point)	0.72	−0.50–1.93	0.06	0.247
Motive: instant reassurance	−1.11	−2.91–0.69	−0.07	0.229
Motive: self-diagnosis	0.99	−1.01–3.00	0.05	0.332
Motive: cost/logistics	0.44	−1.50–2.39	0.02	0.656
Motive: need for a doctor?	0.09	−1.71–1.90	0.01	0.919
Health anxiety (SHAI, per point)	**0.68**	**0.53–0.84**	**0.48**	**<0.001**

Block 1 (socio-demographics) $R^2 = 0.030$; Block 2 (+ digital use) $R^2 = 0.133$, $\Delta R^2 = 0.103$, p = 0.0002; Block 3 (+ SHAI) $R^2 = 0.317$, $\Delta R^2 = 0.184$, p < 0.001; adjusted $R^2 = 0.268$; F = 8.03, p < 0.001. HC3 robust standard errors. β: standardised coefficient.

In logistic regression (Table 10, Figure 8), elevated health anxiety was independently associated with higher cyberchondria (AOR = 3.12 per SD, 95% CI 2.09–4.65), with searching for instant reassurance (AOR = 3.83, 95% CI 1.75–8.42) and with younger age (AOR = 0.61 per age-group step). Discrimination was good (AUC 0.83; 0.79 after bootstrap correction for optimism), but with about five events per predictor the odds ratios are likely to be optimistic and are treated as exploratory.

Table 10. Multivariable logistic regression of elevated health anxiety (SHAI ≥ 18; 57 events, n = 300)

Predictor	AOR	95% CI	p
Age group (per step)	**0.61**	**0.38–0.96**	**0.034**
Female	1.23	0.63–2.40	0.551

Health-related background	1.05	0.47–2.35	0.909
Chronic condition	1.51	0.64–3.53	0.345
Answered mainly in English	0.94	0.45–1.97	0.871
Uses AI chatbots	0.61	0.22–1.69	0.346
Uses search engines	1.43	0.69–2.96	0.330
Trust in online/AI (per point)	0.74	0.46–1.21	0.232
Motive: instant reassurance	**3.83**	**1.75–8.42**	**<0.001**
Motive: self-diagnosis	0.93	0.42–2.09	0.867
Cyberchondria (per 1 SD)	**3.12**	**2.09–4.65**	**<0.001**

AUC = 0.83 (apparent; optimism-corrected 0.79); McFadden pseudo-R^2 = 0.22; likelihood-ratio $p < 0.001$; robust (HC1) standard errors.




Figure 8. *Adjusted odds ratios for elevated health anxiety (SHAI ≥ 18). Orange: $p < 0.05$.*

3.8 Health anxiety, cyberchondria and pre-consultation behaviour

When health anxiety and cyberchondria were entered together (Table 11), cyberchondria rather than health anxiety went with behaviour: higher CSS scores were associated with urgent visits (AOR = 1.51), fear-driven delay (AOR = 1.59), less ignoring of symptoms (AOR = 0.65) and a more strained relationship with physicians (ordinal OR = 1.34). Only one behavioural association survived FDR correction across the 40 tests: higher cyberchondria was associated with lower odds of never disclosing searches to the physician (AOR = 0.57, 95% CI 0.43–0.78, q = 0.012). Higher health anxiety showed the opposite tendency (AOR = 1.47, q = 0.098), which is exploratory. Greater trust in online/AI information was associated with fewer

multiple-doctor consultations (AOR = 0.59). Neither score was associated with self-medication or with delaying care because of reassurance.

Table 11. Adjusted associations of health anxiety and cyberchondria with pre-consultation behaviours and the doctor–patient relationship

Outcome	Events	SHAI AOR (95% CI)	p	CSS AOR (95% CI)	p
Urgent visit/ER	66	0.91 (0.66–1.25)	0.551	1.51 (1.09–2.10)	0.014
Multiple doctors	76	1.06 (0.79–1.43)	0.688	1.20 (0.90–1.60)	0.208
Delay/skip care (reassured)	104	0.89 (0.66–1.19)	0.417	0.83 (0.63–1.10)	0.190
Delay care (terrified)	26	1.49 (0.94–2.36)	0.087	1.59 (1.09–2.30)	0.015
Self-medication	64	0.81 (0.56–1.16)	0.245	1.00 (0.71–1.41)	0.994
Ignore symptom	90	1.16 (0.85–1.58)	0.359	0.65 (0.48–0.89)	0.007
Never disclose to physician	95	1.47 (1.08–1.98)	0.013	0.57 (0.43–0.78)	<0.001
Negative effect on doctor relationship	104	1.03 (0.78–1.38)	0.824	1.35 (1.00–1.83)	0.049

AOR per 1 SD of each score, from logistic models containing both scores plus trust, AI use, age, gender, health background, chronic disease and language. FDR-adjusted q-values are given in the supplementary workbook.

3.9 Pathways: mediation and moderation

In exploratory mediation models (Table 12, Figure 9), excessive searching statistically accounted for part of the association between health anxiety and post-search distress (indirect = 0.16, 95% CI 0.11–0.23; direct c′ = 0.30), and post-search distress for almost all of the association between health anxiety and reassurance-seeking consultations (indirect = 0.22; direct c′ = 0.01, p = 0.932). The reverse ordering was equally significant (indirect = 0.15, 95% CI 0.09–0.22), so the data cannot distinguish direction. Participants with elevated health anxiety scored higher on all four dimensions, most markedly excessiveness and distress (Figure 10).

None of the tested moderators (trust, AI use, health background, gender, chronic disease, language or age) significantly altered the SHAI–CSS association after FDR correction (all q > 0.40; Table 12 note), and the correlation was moderate to strong in every subgroup of adequate size (r = 0.35 to 0.57).

Table 12. Exploratory mediation models decomposing the association between health anxiety (HA) and the dimensions of cyberchondria (standardised coefficients, n = 300)

Model	a	b	Total (c)	Direct (c′)	Indirect (95% boot CI)

A. HA → excessive searching → post-search distress	0.46	0.35	0.46	0.30	0.16 (0.11–0.23)
B. HA → excessive searching → compulsion	0.46	0.27	0.28	0.15	0.13 (0.07–0.20)
C. HA → post-search distress → reassurance seeking	0.46	0.48	0.22	0.01 (ns)	0.22 (0.16–0.29)
D. Excessive searching → distress → HA (reverse)	0.49	0.30	0.47	0.32	0.15 (0.09–0.22)
E. Serial: HA → excess → distress → reassurance				−0.04 (ns)	0.07 (0.04–0.11)

a: predictor → mediator; b: mediator → outcome adjusted for the predictor; indirect effect = a × b = c − c′. All models adjusted for age, gender, health background, chronic disease, trust and AI use; 5000 bootstrap resamples. All a, b and total paths p < 0.001. Moderation of the SHAI–CSS association (interaction terms): trust in online/ai data p = 0.34; ai chatbot use p = 0.47; health-related background p = 0.66; female gender p = 0.98; chronic condition p = 0.75; english-language responder p = 0.29; age group (ordinal) p = 0.07 (all q > 0.40).




Figure 9. *Mediation models: (A) excessive searching as a pathway from health anxiety to post-search distress; (B) post-search distress as a pathway from health anxiety to reassurance seeking.*




Figure 10. *Cyberchondria dimensions by health-anxiety status (mean ± 95% CI).*

3.10 Sensitivity analyses

All sensitivity analyses left the main findings unchanged (Table 13). The only result that depended on an analytic decision was the association between AI use and cyberchondria, which disappeared under the strict definition of AI use.

Table 13. Sensitivity analyses of the main findings

Analysis	Sample	Result	Conclusion
Main analysis (all participants)	n = 300	r = 0.47	Reference
Excluding the three annotated responses	n = 297	r = 0.46	Unchanged
Excluding straight-line responders	n = 291	r = 0.47	Unchanged
SHAI-14 without item 2 (reordered options)	13-item total	r = 0.48	Unchanged
Bootstrap of the SHAI coefficient (CSS model)	n = 300	B 95% CI 0.54–0.83	Stable
Quantile regression (25th–90th percentiles)	n = 300	B = 0.55 to 0.85	Consistent
Quadratic term for SHAI	n = 300	p = 0.245	Linear association
Multicollinearity (CSS model)		maximum VIF = 3.9	Low
Strict AI-use definition (ticked only)	n = 234 users	B = 1.49, p = 0.314	AI effect not robust
Early vs late responders	halves	CSS p = 0.46; SHAI p = 0.07	No difference
Regression residuals		Shapiro–Wilk p = 0.020; Breusch–Pagan p = 0.004	Robust SEs used

r: Pearson correlation between SHAI-14 and CSS-12. B: unstandardised coefficient of SHAI in the robust regression of CSS-12. Straight-line responders gave identical answers to every item of the SHAI-14 or of the CSS-12.

4. Discussion

4.1 Principal findings

This study is, to our knowledge, the first to apply the CSS-12 in a Baghdad-based sample, and among the first to place AI chatbots alongside search engines as a source of health information before consulting a doctor. Four findings stand out. First, AI chatbots were used by 78.0–88.7% of participants, depending on the definition, more than search engines, and about two in five used AI without search engines at all. Second, health anxiety and cyberchondria were moderately and robustly associated (r = 0.47), independently of all other measured factors, and health anxiety was the

dominant predictor of cyberchondria. Third, the association was concentrated in the excessiveness and distress dimensions; an exploratory decomposition showed that most of it ran statistically through excessive searching and post-search distress. Fourth, cyberchondria was reflected in behaviour towards physicians, most robustly in more frequent disclosure of searches.

4.2 Health anxiety and cyberchondria

The observed correlation ($r = 0.47$) supports H1 and H2 but is weaker than the pooled estimates of about 0.62–0.63 from meta-analyses [18,19]. Part of the gap reflects measurement error: corrected for the reliability of both scales, the correlation would be about 0.56, close to the meta-analytic range, although our observed estimate remains 0.47. Other explanations are also plausible. Our sample had relatively low health anxiety (mean SHAI 12.3) and a narrow range in which correlations are attenuated; recruitment through online channels may have favoured confident, habitual Internet users for whom searching is routine rather than alarming; and most previous studies used the full 33-item CSS, whose distress and reassurance items overlap more with health anxiety. The pattern across dimensions is also informative. Distress and excessiveness, the dimensions closest to the cognitive model of health anxiety, were most strongly related to SHAI, while compulsion (interference with work and social life) was rare and only weakly related, consistent with the subscale range reported previously [19]. At item level, it was the *interpretation* of sensations and the *fear* of serious illness, not mere awareness of bodily sensations, that tracked cyberchondria. This is consistent with the view that cyberchondria is more closely tied to catastrophic appraisal than to bodily vigilance itself [5,19].

The prevalence of elevated health anxiety (19% at SHAI ≥ 18) should be interpreted cautiously. The cut-off was derived against healthy controls in Sweden [42]; it is not a diagnosis, and prevalence ranged from 7% to 33%

across plausible cut-offs. Nevertheless, it indicates that a substantial minority of digitally active adults in Baghdad experience health worry at a level that would warrant clinical attention, consistent with high rates of health anxiety reported in Iraq during the pandemic [9].

4.3 Do health anxiety and cyberchondria fuel each other?

The mediation results are consistent with the reinforcing cycle proposed by cognitive-behavioural models and by the longitudinal Dutch study [5,21]: people with higher health anxiety search more, more searching goes with more distress, and distress goes with more reassurance seeking from doctors. That the direct effect of health anxiety on reassurance seeking became negligible once post-search distress was accounted for is compatible with online searching being one route by which health anxiety is turned into healthcare use, but it may equally reflect the strong overlap between the Distress and Reassurance subscales. However, these results must not be over-interpreted. In cross-sectional data, mediation tests are tests of a hypothesised ordering, not of causation, they can be seriously biased estimates of longitudinal processes [54], and a significant indirect effect cannot by itself identify the true mediator or the causal order [55]. Our reverse-direction model (searching → distress → health anxiety) was equally significant, which is exactly what one would expect if the relationship is bidirectional, but also what one would expect if both constructs simply share variance. In addition, the mediators and outcomes in models A–C are subscales of the same instrument, so part of their association reflects shared method variance and overlapping item content [56]. The appropriate conclusion is that our data are *compatible with*, but cannot *demonstrate*, a cycle in which health anxiety and cyberchondria fuel each other.

H3 is therefore supported statistically, but not causally.

4.4 The role of AI chatbots

The extent of AI chatbot use (78% ticked; 88.7% including those who named a chatbot) is a new finding and contrasts with a 2024 US survey in which about one in five respondents reported using ChatGPT for health information [28], and with a 2025 online survey in which 21% used LLM chatbots for health information against 98% using search engines [57]. Our sample was young and highly educated, which partly explains the difference: AI use fell from 96% in those aged 18–19 to 72% in those aged 40 and over. Under the inclusive definition, AI users had higher cyberchondria, mainly in excessiveness and reassurance seeking, but this association disappeared under the stricter definition, rested on a small and older comparison group (n = 34), and AI use did not modify the strength of the health anxiety–cyberchondria link. We therefore find no reliable evidence that AI chatbot use is associated with cyberchondria independently of health anxiety; this is an absence of evidence rather than evidence of absence. Because LLM chatbots answer in fluent, personalised and empathetic language [26,27], they could invite repeated, conversational querying; alternatively, people who already search heavily may simply adopt every available tool. The level of AI use in our sample resembles that of young Turkish dermatology outpatients (61%), in whom pre-visit AI consultation mainly confirmed an existing decision to seek care and was not associated with increased anxiety [31]; in a large Czech sample, health anxiety was linked to AI use only through trust [30], whereas in our data trust was unrelated to health anxiety. On current evidence, AI chatbots appear to be a new channel for, rather than a demonstrated driver of, cyberchondria, which is consistent with observations that health-anxious people are over-represented among users of digital symptom checkers [29].

Trust in online and AI information was mostly moderate and was associated with excessive searching but not with distress. Higher trust was

also associated with lower odds of consulting several doctors, suggesting that trusted AI answers may substitute for some "doctor shopping". This is potentially beneficial for health services, but it is also where false reassurance becomes a risk: higher trust showed a non-significant trend towards delaying care because of reassurance (AOR = 1.33, p = 0.088), echoing a multinational study from the Middle East and North Africa in which trust in online information strongly predicted delayed consultation [36].

4.5 Pre-consultation behaviour and the doctor–patient relationship

The most common response after searching, delaying or skipping care because the information was reassuring (35%), is a double-edged finding. For self-limiting symptoms it may save unnecessary visits; for serious ones it represents a risk of delayed diagnosis. Notably, this behaviour was not related to health anxiety or cyberchondria, meaning it occurs across the spectrum and cannot be dismissed as a feature of anxious users. Conversely, cyberchondria was associated with urgent visits and emergency attendance and with fear-driven delay, although these associations did not survive FDR correction and are exploratory, mirroring evidence that health anxiety and cyberchondria increase healthcare utilisation [6,16]. Thus online searching appears to push different people in opposite directions: towards complacency in some and towards over-use of services in others.

The disclosure findings are clinically important. Almost a third of participants said they never told their doctor about their searches for fear of being judged or dismissed; because the response option combined frequency with a reason, this figure describes non-disclosure attributed to fear of judgement rather than all non-disclosure. Higher cyberchondria was robustly associated with more disclosure (q = 0.012), whereas the tendency for higher health anxiety to go with concealment was nominal only (q = 0.098). One interpretation is that patients for whom searching is a dominant behaviour

bring their findings to the consultation as part of reassurance seeking, while health-anxious patients may anticipate being labelled as "worried well"; the second part of this interpretation needs confirmation. Concealment matters because online information discussed openly can strengthen the doctor–patient relationship, whereas undisclosed information is more likely to undermine it [33]. One in three participants reported that searching had complicated (22%) or significantly reduced (13%) their relationship with or trust in physicians, and this tended to be more likely at higher levels of cyberchondria (a nominal association).

H4 is therefore only partially supported: the only behavioural association that survived correction for multiple testing was between higher cyberchondria and more frequent disclosure of searches to physicians.

4.6 Medical and health-science students

Medical students had higher unadjusted health anxiety than non-health employees, but this difference disappeared after adjustment, largely reflecting their younger age; age was an independent predictor of elevated health anxiety. The SHAI–CSS correlation tended to be stronger among medical students. This is compatible with the idea of "medical student syndrome", but evidence for that concept is weak: in a recent Palestinian study using the same SHAI ≥ 18 threshold, elevated health anxiety was actually least common among medical students [58], and our data suggest that medical knowledge does not protect against cyberchondria (medical students' CSS scores were identical to those of other participants).

4.7 Comparison with regional studies

Our findings are in line with emerging Arab-region data. In Jeddah, cyberchondria was common and associated with smartphone addiction [34]; among Saudi primary-care patients, cyberchondria had a positive effect on

health anxiety in structural models [35]; and the Arabic CSS-12 showed a structure similar to ours in Lebanon [15]. Among 1435 Egyptian medical students, the population closest to ours, 85.8% showed moderate cyberchondria on the CSS-12, and worry about one's own or a family member's health predicted it [59]; a Saudi validation of the CSS-12 likewise found acceptable reliability and positive correlations with health anxiety [46]. The good fit of the four-factor CSS-12, its full scalar invariance across Arabic and English responders and the absence of item-level DIF add to the evidence that the instrument functions well in Arabic-speaking populations.

4.8 Strengths and limitations

Strengths of this study include the use of two established instruments with good reliability and confirmed structure; a bilingual questionnaire with formal tests of measurement invariance across languages; the separate assessment of AI chatbots; complete data; correction for multiple testing; robust regression, bootstrap inference and several sensitivity analyses; and the linking of psychological measures to concrete behaviours relevant to clinical practice.

The limitations are substantial and should be weighed carefully:

- **Design and causality:** the cross-sectional design cannot establish temporal order; mediation results describe statistical pathways, not causal mechanisms, and may be biased relative to longitudinal effects.
- **Sampling and generalisability:** an online convenience/snowball sample is subject to self-selection. The sample was young, highly educated and over-represented medical students (44.0%) and people with health backgrounds (63.3%), so prevalence estimates and the frequency of AI use cannot be generalised to the adult population of Baghdad. Residence in Baghdad was assured by the distribution channels rather than verified by a

question, and non-Internet users, older adults and less-educated people are under-represented by design.

- **Measurement:** the Arabic wording was produced by the research team without a formal forward–backward translation and cognitive pre-testing; formal tests supported full (scalar) invariance of the CSS-12 and metric invariance of the SHAI-14 across language groups, with no item showing DIF after correction, but SHAI-14 intercepts were not fully equivalent, the language groups were self-selected rather than randomised, and statistical invariance does not replace formal cross-cultural adaptation with cognitive pre-testing. Published Arabic versions of both scales exist [15,44,46] but were not used, so comparability with Arabic studies that used them is uncertain. On the form, the response options of one SHAI item (item 2) were displayed out of their original order; scoring followed the original key, and dropping the item did not change the main result, but the order may still have influenced answers. CSS item 3 refers only to search engines and may under-capture repeated querying of AI chatbots. "Language relied on" was self-selected, not randomised. The SHAI-14 one-factor model fitted poorly under maximum-likelihood estimation. The SHAI cut-off was derived in Sweden, and the CSS-12 has no validated clinical cut-off, so the "high cyberchondria" group is sample-relative.
- **Single-item and self-designed measures:** trust, motivations, behaviours, disclosure and relationship were assessed with single, non-validated items, some allowing multiple answers. The questionnaire asked for "most frequent" behaviours but allowed several answers; the trust item combined online tools and AI, so it cannot separate trust in search results from trust in chatbots; the disclosure option combined frequency with a reason (fear of judgement); and "somewhat complicated" is not clearly a negative effect on the relationship.

- **Self-report bias:** all measures were self-reported and may be affected by recall and social desirability bias; no clinical data or diagnoses were available, and chronic disease was self-reported.
- **Multiple testing and power:** despite FDR correction, the large number of exploratory tests means that some isolated associations may be false positives; several behavioural associations were nominally significant only. Some subgroups (non-AI users, n = 34; healthcare workers, n = 15; older age groups) were small, and the logistic model for clinical health anxiety had about five events per predictor.
- **Classification of AI use:** 32 respondents named an AI chatbot without ticking the AI option. They were classed as users, and the independent association between AI use and cyberchondria depended on this decision, so it should be regarded as tentative.
- **Data quality:** an online survey cannot exclude careless responding entirely; nine straight-line responders were identified, although their exclusion did not change the results.

5. Conclusions and Recommendations

5.1 Conclusions

1. AI chatbots, mainly ChatGPT, have become a leading source of health information before consulting a doctor among digitally active adults in Baghdad: 78.0% ticked them (88.7% including those who named a chatbot), more than used search engines.
2. About one in five participants (19%) had elevated health anxiety (SHAI ≥ 18, a screening threshold rather than a diagnosis; 0.7% at the stricter cut-off of 29), and younger people and those searching for instant reassurance were most affected.

3. Health anxiety and cyberchondria are moderately and robustly associated (r = 0.47); health anxiety is the strongest independent predictor of cyberchondria, especially of excessive searching and post-search distress.
4. The pattern of associations is compatible with a reinforcing cycle (health anxiety, excessive searching, distress, reassurance-seeking consultations), but the cross-sectional decomposition, based on subscales of one instrument, cannot establish direction.
5. There is no reliable evidence that AI-chatbot use is associated with cyberchondria independently of health anxiety: the association depended on how inconsistent answers about AI use were classified, and AI use did not change the strength of the health anxiety–cyberchondria link.
6. Online searching shapes the consultation: about one-third of participants delay care because they feel reassured, about one-third never disclose their searches to their doctor for fear of judgement, and about one-third say searching has complicated or reduced their trust in physicians. Higher cyberchondria was robustly linked with more disclosure; links with urgent care-seeking, concealment by health-anxious participants and a strained relationship were exploratory.

5.2 Recommendations

For clinicians and medical education:

- Routinely and non-judgementally ask patients what they have searched for or asked an AI chatbot, to bring hidden information into the consultation and reduce concealment.
- Recognise repeated searching, post-search distress and repeated reassurance seeking as possible markers of health anxiety, and screen with a brief tool such as the SHAI where appropriate; offer cognitive-behavioural strategies rather than repeated reassurance or investigations.

- Include digital health communication, AI literacy and the management of health anxiety in undergraduate medical curricula, and support medical students who may themselves be vulnerable.

For the Ministry of Health and public-health bodies:

- May consider campaigns on digital and AI health literacy that explain both the uses and the limits of online information, especially false reassurance that leads to delayed care.
- May consider providing or endorsing reliable Arabic-language health information sources and triage services so that people searching for "whether I need a doctor" are directed to safe advice.

For developers of AI tools:

- May consider designing health responses that recognise signs of health anxiety (e.g. repeated questions about the same symptom), avoid escalating rare diagnoses without context, and encourage appropriate professional consultation, in Arabic as well as English.

For future research:

- Develop and test interventions: ClinicalTrials.gov lists only nine registered studies on cyberchondria (October 2026), none in Iraq and six of them observational and three interventional [60], so trials of brief digital health-literacy or cognitive-behavioural interventions in Arabic are a clear gap.
- Conduct longitudinal or experience-sampling studies to test whether health anxiety and cyberchondria reinforce each other over time, and experimental studies comparing AI chatbot and search-engine answers on post-search anxiety.
- Use probability sampling of the general Baghdad population, including older and less-educated adults, and use formally adapted Arabic instruments (the published CSS-12-Ar and Arabic SHAI, or a new

forward–backward adaptation with measurement-invariance testing), with CSS item 3 worded to cover both search engines and AI chatbots.

- Link self-reported behaviours to objective healthcare utilisation and clinical outcomes, especially delayed diagnosis following false reassurance.

6. References

1. White RW, Horvitz E. Cyberchondria: studies of the escalation of medical concerns in Web search. ACM Trans Inf Syst. 2009;27(4):1-37. doi:10.1145/1629096.1629101
2. Starcevic V, Berle D. Cyberchondria: towards a better understanding of excessive health-related Internet use. Expert Rev Neurother. 2013;13(2):205-13. doi:10.1586/ern.12.162
3. Kemp S. Digital 2025: Iraq [Internet]. DataReportal; 2025 [cited 2026 Oct 15]. Available from: https://datareportal.com/reports/digital-2025-iraq
4. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed., text revision (DSM-5-TR). Washington (DC): American Psychiatric Association Publishing; 2022.
5. Salkovskis PM, Warwick HMC. Morbid preoccupations, health anxiety and reassurance: a cognitive-behavioural approach to hypochondriasis. Behav Res Ther. 1986;24(5):597-602. doi:10.1016/0005-7967(86)90041-0
6. Tyrer P. Recent advances in the understanding and treatment of health anxiety. Curr Psychiatry Rep. 2018;20(7):49. doi:10.1007/s11920-018-0912-0
7. Tyrer P. Why health anxiety needs to be recognised in hospital practice. Clin Med (Lond). 2020;20(3):339-42. doi:10.7861/clinmed.2019-0346
8. Kosic A, Lindholm P, Järvholm K, Hedman-Lagerlöf E, Axelsson E. Three decades of increase in health anxiety: systematic review and meta-analysis of birth cohort changes in university student samples from 1985 to 2017. J Anxiety Disord. 2020;71:102208. doi:10.1016/j.janxdis.2020.102208
9. Karim SK, Taha PH, Amin NMM, Ahmed HS, Yousif MK, Hallumy AM. COVID-19-related anxiety disorder in Iraq during the pandemic: an online cross-sectional study. Middle East Curr Psychiatry. 2020;27(1):55. doi:10.1186/s43045-020-00067-4
10. Taha PH. Home quarantine induced health anxiety during the beginning of the COVID-19 pandemic: evidence from Iraq. Disaster Med Public Health Prep. 2023;17:e26. Epub 2021 Jul 26. doi:10.1017/dmp.2021.242
11. Starcevic V, Berle D, Arnáez S. Recent insights into cyberchondria. Curr Psychiatry Rep. 2020;22(11):56. doi:10.1007/s11920-020-01179-8
12. Vismara M, Caricasole V, Starcevic V, Cinosi E, Dell'Osso B, Martinotti G, et al. Is cyberchondria a new transdiagnostic digital compulsive syndrome? A systematic

review of the evidence. Compr Psychiatry. 2020;99:152167. doi:10.1016/j.comppsych.2020.152167

13. McElroy E, Shevlin M. The development and initial validation of the cyberchondria severity scale (CSS). J Anxiety Disord. 2014;28(2):259-65. doi:10.1016/j.janxdis.2013.12.007
14. McElroy E, Kearney M, Touhey J, Evans J, Cooke Y, Shevlin M. The CSS-12: development and validation of a short-form version of the cyberchondria severity scale. Cyberpsychol Behav Soc Netw. 2019;22(5):330-5. doi:10.1089/cyber.2018.0624
15. Hallit S, Rogoza R, Abi Semaan C, Azzi V, Sawma T, Obeid S. Validation of the Arabic version of the cyberchondria severity scale 12 items (CSS-12-Ar) among a sample of Lebanese adults. BMC Psychiatry. 2023;23(1):618. doi:10.1186/s12888-023-05123-x
16. Mathes BM, Norr AM, Allan NP, Albanese BJ, Schmidt NB. Cyberchondria: overlap with health anxiety and unique relations with impairment, quality of life, and service utilization. Psychiatry Res. 2018;261:204-11. doi:10.1016/j.psychres.2018.01.002
17. Jungmann SM, Brand S, Kolb J, Witthöft M. Do Dr. Google and health apps have (comparable) side effects? An experimental study. Clin Psychol Sci. 2020;8(2):306-17. doi:10.1177/2167702619894904
18. McMullan RD, Berle D, Arnáez S, Starcevic V. The relationships between health anxiety, online health information seeking, and cyberchondria: systematic review and meta-analysis. J Affect Disord. 2019;245:270-8. doi:10.1016/j.jad.2018.11.037
19. Schenkel SK, Jungmann SM, Gropalis M, Witthöft M. Conceptualizations of cyberchondria and relations to the anxiety spectrum: systematic review and meta-analysis. J Med Internet Res. 2021;23(11):e27835. doi:10.2196/27835
20. Jungmann SM, Dessauer E. Health-related internet use and cyberchondria in adolescents: population-based cross-sectional survey. J Med Internet Res. 2025;27:e65792. doi:10.2196/65792
21. te Poel F, Baumgartner SE, Hartmann T, Tanis M. The curious case of cyberchondria: a longitudinal study on the reciprocal relationship between health anxiety and online health information seeking. J Anxiety Disord. 2016;43:32-40. doi:10.1016/j.janxdis.2016.07.009
22. Šoštarić M, Mikac U, Jokić-Begić N. Understanding cyberchondria in pregnant women: longitudinal assessment of risk factors, triggers, and outcomes. J Psychosom Obstet Gynaecol. 2023;44(1):2265050. doi:10.1080/0167482X.2023.2265050
23. Jokic-Begic N, Lauri Korajlija A, Mikac U. Cyberchondria in the age of COVID-19. PLoS One. 2020;15(12):e0243704. doi:10.1371/journal.pone.0243704
24. Kurcer MA, Erdogan Z, Cakir Kardes V. The effect of the COVID-19 pandemic on health anxiety and cyberchondria levels of university students. Perspect Psychiatr Care. 2022;58(1):132-40. doi:10.1111/ppc.12850

25. Miezah D, Adzakpa R, Bani EMA, Obeng P. Prevalence and associated factors of cyberchondria: a scoping review. ScientificWorldJournal. 2026;2026:9950027. doi:10.1155/tswj/9950027
26. Thirunavukarasu AJ, Ting DSJ, Elangovan K, Gutierrez L, Tan TF, Ting DSW. Large language models in medicine. Nat Med. 2023;29(8):1930-40. doi:10.1038/s41591-023-02448-8
27. Ayers JW, Poliak A, Dredze M, Leas EC, Zhu Z, Kelley JB, et al. Comparing physician and artificial intelligence chatbot responses to patient questions posted to a public social media forum. JAMA Intern Med. 2023;183(6):589-96. doi:10.1001/jamainternmed.2023.1838
28. Ayo-Ajibola O, Davis RJ, Lin ME, Riddell J, Kravitz RL. Characterizing the adoption and experiences of users of artificial intelligence-generated health information in the United States: cross-sectional questionnaire study. J Med Internet Res. 2024;26:e55138. doi:10.2196/55138
29. Wetzel AJ, Klemmt M, Müller R, Rieger MA, Joos S, Koch R. Only the anxious ones? Identifying characteristics of symptom checker app users: a cross-sectional survey. BMC Med Inform Decis Mak. 2024;24(1):21. doi:10.1186/s12911-024-02430-5
30. Svestkova A, Huang Y, Smahel D. Factors that influence trust and willingness to use generative AI for health information: a cross-sectional study. Digit Health. 2025;11:20552076251360973. doi:10.1177/20552076251360973
31. Dığış M, Kaya K, Demir B, Çiçek İN, Akdağ D, Akdağ Ş. Frequency of use and attitudes toward AI-based health counseling among patients attending the dermatology outpatient clinic: a cross-sectional study. Cutan Ocul Toxicol. 2026 Aug 6:1-11. Epub ahead of print. doi:10.1080/15569527.2026.2711643
32. Wang J, Ashvetiya T, Quaye E, Parakh K, Martin SS. Online health searches and their perceived effects on patients and patient-clinician relationships: a systematic review. Am J Med. 2018;131(10):1250.e1-1250.e10. doi:10.1016/j.amjmed.2018.04.019
33. Luo A, Qin L, Yuan Y, Yang Z, Liu F, Huang P, et al. The effect of online health information seeking on physician-patient relationships: systematic review. J Med Internet Res. 2022;24(2):e23354. doi:10.2196/23354
34. El-Zayat A, Namnkani SA, Alshareef NA, Mustfa MM, Eminaga NS, Algarni GA. Cyberchondria and its association with smartphone addiction and electronic health literacy among a Saudi population. Saudi J Med Med Sci. 2023;11(2):162-8. doi:10.4103/sjmms.sjmms_491_22
35. Pasay-An E, Alalawi RH, Shanmugam SR, Alshammari SA, Alreshidi MS, Alkubati S, et al. "Dr. Google" consultation and health anxiety among primary healthcare patients: the mediation effect of health literacy. PLoS One. 2025;20(10):e0325791. doi:10.1371/journal.pone.0325791
36. Dahshan A, Mahfouz Khalil MI, Abdelfatah D, Abu Ejheisheh M, Alshammari MSH, Mekdad AK, et al. Influence of online health information and artificial intelligence on

decision-making among migraine patients: a multinational cross-sectional study from the MENA region. Cephalalgia. 2026;46(4):3331024261435594. doi:10.1177/03331024261435594

37. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. Lancet. 2007;370(9596):1453-7. doi:10.1016/S0140-6736(07)61602-X
38. Eysenbach G. Improving the quality of Web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). J Med Internet Res. 2004;6(3):e34. doi:10.2196/jmir.6.3.e34
39. Serdar CC, Cihan M, Yücel D, Serdar MA. Sample size, power and effect size revisited: simplified and practical approaches in pre-clinical, clinical and laboratory studies. Biochem Med (Zagreb). 2021;31(1):010502. doi:10.11613/BM.2021.010502
40. Salkovskis PM, Rimes KA, Warwick HMC, Clark DM. The Health Anxiety Inventory: development and validation of scales for the measurement of health anxiety and hypochondriasis. Psychol Med. 2002;32(5):843-53. doi:10.1017/S0033291702005822
41. Arnáez S, García-Soriano G, López-Santiago J, Belloch A. The Spanish validation of the Short Health Anxiety Inventory: psychometric properties and clinical utility. Int J Clin Health Psychol. 2019;19(3):251-60. doi:10.1016/j.ijchp.2019.05.003
42. Österman S, Axelsson E, Lindefors N, Hedman-Lagerlöf E, Hedman-Lagerlöf M, Kern D, et al. The 14-item short health anxiety inventory (SHAI-14) used as a screening tool: appropriate interpretation and diagnostic accuracy of the Swedish version. BMC Psychiatry. 2022;22:701. doi:10.1186/s12888-022-04367-3
43. Poshtan MM, Aflakseir A, Witthöft M, Cherry MG, Ramzi M. Persian version of the Short Health Anxiety Inventory (SHAI) for patients with cancer: evaluation of psychometric properties, factor structure, and association with related constructs. J Psychosom Res. 2024;187:111958. doi:10.1016/j.jpsychores.2024.111958
44. Alshayea AK. Latent structure, measurement invariance, and reliability of an Arabic version of the Short Health Anxiety Inventory. J Exp Psychopathol. 2020;11(2). doi:10.1177/2043808720912629
45. Cruchinho P, López-Franco MD, Capelas ML, Almeida S, Bennett PM, Miranda da Silva M, et al. Translation, cross-cultural adaptation, and validation of measurement instruments: a practical guideline for novice researchers. J Multidiscip Healthc. 2024;17:2701-28. doi:10.2147/JMDH.S419714
46. Alshaibani M. Validity and reliability of the short form of Cyberchondria Severity Scale (CSS-12) in Saudi Arabia. J Consum Health Internet. 2023;27(3):282-93. doi:10.1080/15398285.2023.2237347
47. Hayes AF, Coutts JJ. Use omega rather than Cronbach's alpha for estimating reliability. But… Commun Methods Meas. 2020;14(1):1-24. doi:10.1080/19312458.2020.1718629

48. McNeish D, Wolf MG. Dynamic fit index cutoffs for confirmatory factor analysis models. Psychol Methods. 2023;28(1):61-88. doi:10.1037/met0000425
49. Lovik A, Nassiri V, Verbeke G, Molenberghs G. A modified Tucker's congruence coefficient for factor matching. Methodology. 2020;16(1):59-74. doi:10.5964/meth.2813
50. Leitgöb H, Seddig D, Asparouhov T, Behr D, Davidov E, De Roover K, et al. Measurement invariance in the social sciences: historical development, methodological challenges, state of the art, and future perspectives. Soc Sci Res. 2023;110:102805. doi:10.1016/j.ssresearch.2022.102805
51. Jin KY, Chen HF. MIMIC approach to assessing differential item functioning with control of extreme response style. Behav Res Methods. 2020;52(1):23-35. doi:10.3758/s13428-019-01198-1
52. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J R Stat Soc Series B Methodol. 1995;57(1):289-300. doi:10.1111/j.2517-6161.1995.tb02031.x
53. Hayes AF. Introduction to mediation, moderation, and conditional process analysis: a regression-based approach. 3rd ed. New York: Guilford Press; 2022.
54. O'Laughlin KD, Martin MJ, Ferrer E. Cross-sectional analysis of longitudinal mediation processes. Multivariate Behav Res. 2018;53(3):375-402. doi:10.1080/00273171.2018.1454822
55. Rohrer JM, Hünermund P, Arslan RC, Elson M. That's a lot to process! Pitfalls of popular path models. Adv Methods Pract Psychol Sci. 2022;5(2). doi:10.1177/25152459221095827
56. Podsakoff PM, Podsakoff NP, Williams LJ, Huang C, Yang J. Common method bias: it's bad, it's complex, it's widespread, and it's not easy to fix. Annu Rev Organ Psychol Organ Behav. 2024;11:17-61. doi:10.1146/annurev-orgpsych-110721-040030
57. Yun HS, Bickmore T. Online health information-seeking in the era of large language models: cross-sectional web-based survey study. J Med Internet Res. 2025;27:e68560. doi:10.2196/68560
58. Fasfoos A, Sunkrot MS, Jadallah I, Id'eas AH, Hoshieh M, Haytham T, et al. Health anxiety among medical and non-medical university students in Palestine: associations with intolerance of uncertainty, daily phone use, and field of study. BMC Psychol. 2026;14(1). doi:10.1186/s40359-026-04751-8
59. El-Zoghby SM, Zaghloul NM, Tawfik AM, Elsherbiny NM, Shehata SA, Soltan EM. Cyberchondria and smartphone addiction: a correlation survey among undergraduate medical students in Egypt. J Egypt Public Health Assoc. 2024;99(1):7. doi:10.1186/s42506-024-00154-y
60. ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). Search results for "cyberchondria"; [cited 2026 Oct 16]. Available from: https://clinicaltrials.gov/search?cond=cyberchondria

Appendices

Appendix A. Item-by-item correlations between the SHAI-14 and CSS-12

Item-by-item Spearman correlations between all 14 SHAI-14 items and all 12 CSS-12 items (168 pairs). Correction for multiple testing was applied within this family of 168 tests: 111 pairs were significant after Benjamini–Hochberg FDR correction and 58 after Holm–Bonferroni correction. Full values are given in the supplementary workbook (sheet T28_item_by_item).




Figure 11. *Item-by-item correlations between SHAI-14 and CSS-12 items (Spearman ρ). * significant after FDR correction across the 168 pairs; ** also significant after Holm–Bonferroni correction.*

Appendix B. Study questionnaire

The bilingual (English–Arabic) questionnaire as administered through Google Forms is reproduced below (41 questions).

Digital Health Information-Seeking, Health Anxiety, and Cyberchondria Questionnaire
استبيان البحث عن المعلومات الصحية الرقمية، والقلق الصحي، والسايبركوندريا (توهم المرض عبر الإنترنت)

* Indicates required question



1. **By checking the box below, I confirm that I am at least 18 years of age, have read the description of this study, and voluntarily agree to participate. I understand that I can withdraw from this survey at any point without penalty.** *

بالتأشير على المربع أدناه، أؤكد أن عمري لا يقل عن 18 عاماً، وأنني قرأت وصف هذه الدراسة، وأوافق طواعية على المشاركة فيها. كما أفهم أنه يمكنني الانسحاب من هذا الاستبيان في أي وقت دون أي تبعات.

Mark only one oval.

I explicitly agree and consent to participate in this study. / أوافق وأقبل المشاركة في هذه الدراسة البحثية.

2. **Age Group / الفئة العمرية** *

Mark only one oval.

- 18-19
- 20-29
- 30-39
- 40-49
- 50-59
- 60+

3. **Gender: الجنس** *

Mark only one oval.

- Male / ذكر
- Female / انثى

4. **What is the highest level of education you have completed?** *

ما هو أعلى مستوى تعليمي أكملته؟

Mark only one oval.

- Currently a student / لا أزال طالباً مستمراً بالدراسة
- Postgraduate degree (Master's or Ph.D.) / دراسات عليا (ماجستير أو دكتوراه)
- Bachelor's degree / بكالوريوس
- Diploma / Vocational training // دبلوم (معهد / تدريب مهني)
- High school or below / إعدادية أو أقل
- Other: _____

5. **What is the nature of your current study or work?** *

ما هي طبيعة دراستكِ أو عملكِ الحالي؟

Mark only one oval.

- Medical student (Medicine) / طالب في كلية الطب البشري
- Student in another healthcare field (Pharmacy, Dentistry, Nursing, Analysis, Medical Technologies, etc.) / طالب في مجال صحي آخر (صيدلة، أسنان، تمريض، تحليلات، تقنيات، وما شابه)
- Medical Doctor / Physician (General Practitioner, Resident, or Specialist) / طبيب بشري (ممارس، مقيم، أو اختصاص)
- Other healthcare professional (Pharmacist, Dentist, Nurse, Technician, etc.) / عامل في القطاع الصحي غير الأطباء (صيدلاني، طبيب أسنان، ممرض، تقني، وما شابه)
- Student in a non-healthcare field / طالب في تخصص غير صحي
- Employed/working in a non-healthcare field / موظف أو عامل في مجال غير صحي
- Other: _____

6. **Do you suffer from any diagnosed chronic medical condition** *
هل تعاني من اي امراض طبية مزمنة تم تشخيصها مسبقا

Mark only one oval.

- Yes / نعم
- No / لا

7. **If you answerd yes above please specify which chronic disease you have (Leave blank if you prefer not to say)**
اذا اجبت بنعم اعلاه يرجى كتابة اي مرض مزمن لديك
(اتركه فارغا اذا كنت تفضل عدم الذكر)

8. **Which tools do you use most frequently when looking up health or medical symptoms online?** *
(Select all that apply)
ما هي الادوات التي تستخدمها بشكل متكرر عند البحث عن الأعراض الصحية أو الطبية عبر الإنترنت؟
(يمكن اختيار اكثر من اجابة)

Tick all that apply.

- [ ] Search engine / محركات البحث (google)
- [ ] AI Chatbots / روبوتات الذكاء الاصطناعي
- [ ] Dedicated Medical Platforms / المنصات الطبية المتخصصة(WebMed, Mayo Clinic..)
- [ ] Social Media Networks / منصات التواصل الاجتماعي (Facebook ,Reddit ,Instagram)

9. **If you checked AI above**
specify which AI do you use most of the time
اذا قمت بتحديد الذكاء الاصطناعي
قم بتحديد اي برنامج غالبا

Mark only one oval.

- ChatGPT
- Gemini
- Claude
- Other: _____

10. **What is your primary motivation for searching medical information online? (Select all that apply):** *
ما هو دافعك الرئيسي للبحث عن معلومات طبية عبر الإنترنت؟
(يمكنك اختيار أكثر من إجابة)

Tick all that apply.

- [ ] Immediate convenience and easy accessibility / السهولة الفورية وإمكانية الوصول السريع في أي وقت
- [ ] High costs or logistical difficulty of professional doctor visits / التكاليف المرتفعة أو الصعوبات اللوجستية المرتبطة بزيارة الطبيب المختص
- [ ] Seeking instant reassurance regarding an unexplained bodily symptom / الرغبة في الحصول على طمأنينة فورية بشأن عرض جسدي غير مفسر
- [ ] Intending to self-diagnose or double-check a physician's prescription / نية التشخيص الذاتي أو التحقق من دقة وصفة الطبيب المعالج
- [ ] I just want to see if the symptom is serious enough to warrant a doctor's visit in the first place. / أريد فقط معرفة ما إذا كانت الأعراض خطيرة بدرجة كافية تستدعي زيارة الطبيب في المقام الأول
- [ ] Other: _____

11. **On a scale from 1 to 5, how much do you trust the medical data generated by online tools/AI?** *

على مقياس من 1 إلى 5، ما مدى ثقتك في البيانات الطبية الناتجة عن أدوات الإنترنت أو الذكاء الاصطناعي؟

Mark only one oval.

- Do not trust at all / لا اثق على الاطلاق
- Slightly trust/ واثق قليلا
- Moderately trust/ واثق بدرجة معتدلة
- Very much trust/ واثق جداً
- Completely trust / ثقة مطلقة

12. **Worrying about health** *
القلق بشأن الصحة

Mark only one oval.

- I do not worry about my health /أنا لا أقلق بشأن صحتي
- I occasionally worry about my health / ينتابني القلق أحياناً بشأن صحتي
- I spend much of my time worrying about my health / أقضي الكثير من وقتي في القلق بشأن صحتي
- I spend most of my time worrying about my health / أقضي معظم وقتي في القلق بشأن صحتي

13. **Body awareness / الوعي بآلام الجسد** *

Mark only one oval.

- I notice aches/pains as much as most other people (of my age) / ألاحظ الأوجاع/الآلام بنفس القدر مثل معظم الأشخاص الآخرين (من هم في عمري)
- I notice aches/pains less than most other people (of my age) / ألاحظ الأوجاع/الآلام بشكل أقل من معظم الأشخاص الآخرين (من هم في عمري)
- I notice aches/pains more than most other people (of my age) / ألاحظ الأوجاع/الآلام بشكل أكثر من معظم الأشخاص الآخرين (من هم في عمري)
- I am aware of aches/pains in my body all the time / أنا واعي ومدرك لوجود أوجاع/آلام في جسدي طوال الوقت

14. **Reaction to bodily sensations / رد الفعل تجاه الأحاسيس الجسدية** *

Mark only one oval.

- As a rule, I am not aware of bodily sensations or changes / كقاعدة عامة، أنا لا أنتبه للأحاسيس أو التغيرات الجسدية
- Sometimes I am aware of bodily sensations or changes / في بعض الأحيان أكون واعياً للأحاسيس أو التغيرات الجسدية
- I am often aware of bodily sensations or changes / غالباً ما أكون واعياً للأحاسيس أو التغيرات الجسدية
- I am constantly aware of bodily sensations or changes / أنا واعي ومدرك للأحاسيس أو التغيرات الجسدية باستمرار وبشكل دائم

15. **Thoughts of illness / الأفكار المتعلقة بالمرض** *

Mark only one oval.

- Resisting thoughts of illness is never a problem / لا تشكل مقاومة الأفكار المتعلقة بالمرض أي مشكلة بالنسبة لي على الإطلاق
- Most of the time I can resist thoughts of illness / في معظم الأوقات، يمكنني مقاومة الأفكار المتعلقة بالمرض
- I try to resist thoughts of illness but am often unable to do so / أحاول مقاومة الأفكار المتعلقة بالمرض ولكنني غالباً ما أكون عاجزاً عن ذلك
- Thoughts of illness are so strong that I no longer even try to resist them / الأفكار المتعلقة بالمرض قوية جداً لدرجة أنني لم أعد أحاول مقاومتها

16. **Fear of serious illness / الخوف من الإصابة بمرض خطير** *

Mark only one oval.

- As a rule, I am not afraid that I have a serious illness / كقاعدة عامة، لا يخالجني الخوف من أنني مصاب بمرض خطير
- I am sometimes afraid that I have a serious illness / ينتابني الخوف أحياناً من أنني مصاب بمرض خطير
- I am often afraid that I have a serious illness. / غالباً ما ينتابني الخوف من أنني مصاب بمرض خطير
- I am always afraid that I have a serious illness / أنا دائم الخوف من أنني مصاب بمرض خطير

17. **Images of illness / الصور الذهنية عن المرض** *

Mark only one oval.

- I do not have images (mental pictures) of myself being ill / لا تتكون لدي صور (تخيلات ذهنية) لنفسي وأنا مريض
- I occasionally have images of myself being ill / تتكون لدي أحياناً صور لنفسي وأنا مريض
- I frequently have images of myself being ill / تتكون لدي كثيراً صور لنفسي وأنا مريض
- I constantly have images of myself being ill / تتكون لدي باستمرار صور لنفسي وأنا مريض

18. **Distraction from health thoughts / تشتيت الانتباه عن الأفكار الصحية** *

Mark only one oval.

- I do not have any difficulty taking my mind off thoughts about my health / لا أجد أي صعوبة في صرف ذهني عن الأفكار المتعلقة بصحتي
- I sometimes have difficulty taking my mind off thoughts about my health / أجد أحياناً صعوبة في صرف ذهني عن الأفكار المتعلقة بصحتي
- I often have difficulty taking my mind off thoughts about my health / غالباً ما أجد صعوبة في صرف ذهني عن الأفكار المتعلقة بصحتي
- Nothing can take my mind off thoughts about my health / لا يوجد شيء على الإطلاق يمكنه صرف ذهني عن الأفكار المتعلقة بصحتي

19. **Medical reassurance / الطمأنينة الطبية** *

Mark only one oval.

- I am lastingly relieved if my doctor tells me there is nothing wrong / أشعر بارتياح دائم إذا أخبرني الطبيب أنه لا يوجد خطب ما
- I am initially relieved, but the worries sometimes return later / أشعر بالارتياح في البداية، لكن المخاوف تعود أحياناً لاحقاً
- I am initially relieved, but the worries always return later / أشعر بالارتياح في البداية، لكن المخاوف تعود دائماً لاحقاً
- I am not relieved if my doctor tells me there is nothing wrong / لا أشعر بالارتياح حتى لو اخبرني الطبيب انه لا يوجد خطب ما

20. **Hearing about illness / السماع عن الأمراض** *

Mark only one oval.

- If I hear about an illness, I never think I have it myself / إذا سمعت عن مرض ما، لا يخطر ببالي أبداً أنني مصاب به
- If I hear about an illness, I sometimes think I have it myself / إذا سمعت عن مرض ما، يخطر ببالي أحياناً أنني مصاب به
- If I hear about an illness, I often think I have it myself / إذا سمعت عن مرض ما، غالباً ما يخطر ببالي أنني مصاب به
- If I hear about an illness, I always think I have it myself / إذا سمعت عن مرض ما، يخطر ببالي دائماً أنني مصاب به

21. **Interpretation of sensations / تفسير الأحاسيس الجسدية** *

Mark only one oval.

- If I have a bodily sensation or change, I rarely wonder what it means / إذا شعرت بإحساس أو تغير جسدي، فنادراً ما أتساؤل عما يعنيه
- If I have a bodily sensation or change, I often wonder what it means / إذا شعرت بإحساس أو تغير جسدي، فغالباً ما أتساؤل عما يعنيه
- If I have a bodily sensation or change, I always wonder what it means / إذا شعرت بإحساس أو تغير جسدي، فأنا دائماً أتساؤل عما يعنيه
- If I have a bodily sensation or change, I must know what it means / إذا شعرت بإحساس أو تغير جسدي، فلا بد لي من معرفة ماذا يعني حتماً

22. **Perceived risk / تقدير خطر الإصابة** *

Mark only one oval.

- I usually feel at very low risk for developing a serious illness / أشعر عادةً بأن احتمالية إصابتي بمرض خطير منخفضة جداً
- I usually feel at fairly low risk for developing a serious illness / أشعر عادةً بأن احتمالية إصابتي بمرض خطير منخفضة إلى حد ما
- I usually feel at moderate risk for developing a serious illness / أشعر عادةً بأن احتمالية إصابتي بمرض خطير متوسطة
- I usually feel at high risk for developing a serious illness / أشعر عادةً بأن احتمالية إصابتي بمرض خطير عالية جداً

23. **Thinking you are ill / الاعتقاد بالمرض** *

Mark only one oval.

- I never think I have a serious illness / لا أظن أبداً أنني مصاب بمرض خطير
- I sometimes think I have a serious illness / أظن أحياناً أنني مصاب بمرض خطير
- I often think I have a serious illness / غالباً ما أظن أنني مصاب بمرض خطير
- I usually think that I am seriously ill / أظن عادةً أنني مريض جداً بالفعل

24. **Unexplained sensations / أحاسيس غير مفسرة** *

Mark only one oval.

- If I notice an unexplained bodily sensation, I don't find it difficult to think about other things / إذا لاحظت إحساساً جسدياً غير مفسر، لا أجد صعوبة في التفكير بأشياء أخرى
- If I notice an unexplained bodily sensation, I sometimes find it difficult to think about other things / إذا لاحظت إحساساً جسدياً غير مفسر، أجد أحياناً صعوبة في التفكير بأشياء أخرى
- If I notice an unexplained bodily sensation, I often find it difficult to think about other things / إذا لاحظت إحساساً جسدياً غير مفسر، أجد غالباً صعوبة في التفكير بأشياء أخرى
- If I notice an unexplained bodily sensation, I always find it difficult to think about other things / إذا لاحظت إحساساً جسدياً غير مفسر، أجد دائماً صعوبة في التفكير بأشياء أخرى

25. **Friends & family perspective / وجهة نظر العائلة والأصدقاء** *

Mark only one oval.

- My family/friends would say I do not worry enough about my health / قد يقول عائلتي/أصدقائي إنني لا أقلق كفاية بشأن صحتي
- My family/friends would say I have a normal attitude to my health / قد يقول عائلتي/أصدقائي إن لدي موقفاً طبيعياً ومتوازناً تجاه صحتي
- My family/friends would say I worry too much about my health / قد يقول عائلتي/أصدقائي إنني أقلق أكثر من اللازم بشأن صحتي
- My family/friends would say I am a hypochondriac / قد يقول عائلتي/أصدقائي إنني مصاب بـ "توهم المرض" (وسواس مرضي)

26. **If I notice an unexplained bodily sensation, I search for it on the internet.** *

إذا لاحظتُ عرضاً جسدياً غير مفسر، أقوم بالبحث عنه عبر الإنترنت.

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

27. **I read different web pages about the same health condition.** *

أقرأ صفحات ويب مختلفة ومتعددة حول نفس الحالة الصحية التي أشعر بها

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

28. **I enter the same symptoms into a search engine on more than one occasion.** *

أدخل نفس الأعراض في محرك البحث في أكثر من مناسبة أو بشكل متكرر.

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

29. **Researching health-related conditions or symptoms online distracts me from reading news, sports, or entertainment materials.** *

البحث عن الأعراض أو الحالات الطبية عبر الإنترنت يشتتني عن متابعة الأخبار أو الرياضة أو المواد الترفيهية.

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

30. **Researching health-related conditions or symptoms online interrupts my work (e.g. working on emails or documents).** *

البحث عن الأعراض أو الحالات الطبية عبر الإنترنت يقاطع عملي (مثل كتابة رسائل البريد الإلكتروني أو العمل على المستندات).

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

31. **Researching health-related conditions or symptoms online interrupts my offline social activities (e.g. decreasing time spent with friends/family).** *

البحث عن الأعراض أو الحالات الطبية عبر الإنترنت يقاطع أنشطتي الاجتماعية الواقعية (مثل تقليل الوقت المستقطع مع الأصدقاء أو العائلة).

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

32. **I panic when I read online that a symptom I have could be a sign of a rare or serious condition.** *

أشعر بالذعر عندما أقرأ عبر الإنترنت أن العرض الذي أعاني منه قد يكون مؤشراً لحالة نادرة أو خطيرة.

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

33. **I feel fine until I read about a serious medical condition online.** *

أشعر بأنني بخير إلى أن أقرأ عن حالة صحية خطيرة عبر الإنترنت.

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

34. **I feel more anxious or distressed after researching potential health-related conditions or symptoms online.** *

أشعر بمزيد من القلق أو الضيق بعد البحث عن الأعراض أو الحالات الطبية المحتملة عبر الإنترنت.

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

35. **Researching health-related conditions or symptoms online leads me to consult with my general practitioner.** *

البحث عن الأعراض أو الحالات الطبية عبر الإنترنت يدفعني لاستشارة طبيبي العام.

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

36. **I suggest to my doctor/medical professional that I may need a specific diagnostic procedure that I have read about online (e.g. a biopsy, a specific blood test).** *

أقترح على طبيبي/المختص الصحي أنني قد أحتاج إلى إجراء تشخيصي محدد قرأت عنه عبر الإنترنت (مثل خزعة أو تحليل دم معين).

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

37. **Researching health-related conditions or symptoms online leads me to consult with other medical specialists.** *

البحث عن الأعراض أو الحالات الطبية عبر الإنترنت يدفعني لاستشارة أخصائيين طبيين آخرين.

Mark only one oval.

- Never / أبداً
- Rarely / نادراً
- Sometimes / أحياناً
- Often / غالباً
- Always / دائماً

38. **After searching your health symptoms online, what are your most frequent behavioral responses? (Select all that apply):** *

بعد البحث عن أعراضك الصحية عبر الإنترنت، ما هي استجاباتك السلوكية الأكثر تكراراً؟ (يمكنك اختيار أكثر من إجابة):

Tick all that apply.

- [ ] I purchase self-prescribed medications or treatments without consulting a doctor / أقوم بشراء أدوية أو علاجات ذاتية دون استشارة الطبيب بناءً على نتائج البحث
- [ ] I delay or completely skip seeking professional medical help because the online data reassured me / أقوم بتأجيل أو إلغاء طلب المساعدة الطبية المتخصصة تماماً لأن المعلومات المتوفرة على الإنترنت طمأنتني.
- [ ] I delay professional help because the online data terrified or overwhelmed me / أقوم بتأجيل طلب المساعدة الطبية المتخصصة لأن المعلومات المتوفرة على الإنترنت أرعبتني أو أشعرتني بالقلق الشديد
- [ ] I immediately request an urgent appointment, go to the emergency room, or seek a fast diagnosis / أطلب فوراً موعداً عاجلاً، أو أذهب إلى ردهة الطوارئ، أو أبحث عن تشخيص سريع.
- [ ] I arrange visits with multiple different doctors to double-check my findings / أقوم بزيارة عدة أطباء مختلفين للتأكد من صحة استنتاجاتي (ما عثرت عليه من معلومات)
- [ ] Decided to ignore the symptom entirely. / أقرر تجاهل العَرَض تماماً

39. **During a consultation, do you explicitly share the online search results or AI outputs with your physician?** *

أثناء الاستشارة الطبية، هل تخبر طبيبك صراحةً بنتائج بحثك على الإنترنت أو بما قدمه لك الذكاء الاصطناعي؟

Mark only one oval.

- Yes, I always share my online findings transparently / نعم، أطلع الطبيب دائماً على نتائج بحثي عبر الإنترنت بكل شفافية.
- Sometimes, depending on how the doctor responds or listens / أحياناً، اعتماداً على كيفية استجابة الطبيب أو مدى إصغائه لي.
- No, I never mention my online searches out of fear of being judged or dismissed / لا، لا أذكر أبداً ما بحثت عنه عبر الإنترنت خوفاً من أن يُحكم علي (بالوسوسة مثلاً) أو أن يتم تجاهل كلامي.

40. **How has digital health information-seeking affected your relationship and trust with professional healthcare providers?** *

كيف أثر بحثك عن المعلومات الصحية الرقمية على علاقتك بمقدمي الرعاية الصحية المتخصصين (الاطباء) ومدى ثقتك بهم

Mark only one oval.

- It significantly reduced my trust in my doctor's diagnosis or treatment recommendations / قلل بشكل كبير من ثقتي في تشخيص طبيبي أو التوصيات العلاجية والدوائية المقدمة لي
- It somewhat complicated my relationship, causing me to question choices, but I still follow advice / عقد علاقتي بالطبيب إلى حد ما، مما جعلني أشكك في خياراته، لكني ما زلت ملتزماً باتباع نصيحته في النهاية
- It had no negative impact; it serves purely as helpful supplementary information to my doctor's care / لم يكن له أي تأثير سلبي؛ فهو يعمل كمجرد معلومات تكميلية مفيدة وداعمة للرعاية التي يقدمها طبيبي

41. **Which language did you rely on most while completing this questionnaire?** * **اللغة التي اعتمدت عليها بشكل أساسي أثناء تعبئة هذا الاستبيان؟**

Mark only one oval.

- Mostly English / بشكل أساسي اللغة الإنجليزية
- Mostly Arabic / بشكل أساسي اللغة العربية

This content is neither created nor endorsed by Google.

Google Forms