

UNIT 5

FUZZY SETS AND FUZZY LOGIC

Prof. V.Mary Rajam Vandana

Assistant Professor

Department of AIDS

Content

- Fuzzy Set Operations
- Membership Functions
- Fuzzy Logic
- Hedges
- Fuzzy Proposition
- Inference Rules
- Fuzzy Systems

Introduction

Uncertainty

When A is a fuzzy set and x is a relevant object, the proposition “ x is a member of A ” is not necessarily either true or false. It may be true only to some degree, the degree to which x is actually a member of A .

For example: the weather today

- Sunny: If we define any cloud cover of 25% or less is sunny.
- This means that a cloud cover of 26% is not sunny?
- “Vagueness” should be introduced.

CRISP SET - FUZZY SET

- The **crisp set** is defined in such a way as to partition the individuals in some given universe of discourse into two groups: **members and nonmembers**.

However, many classification concepts do not exhibit this characteristic.
For example, the set of tall people, expensive cars, or sunny days.

- A **fuzzy set** can be defined mathematically by assigning to each possible individual in the universe of discourse a value representing its **grade of membership** in the fuzzy set.

For example: a fuzzy set representing our concept of **sunny** might assign a degree of membership of 1 to a cloud cover of 0%, 0.8 to a cloud cover of 20%, 0.4 to a cloud cover of 30%, and 0 to a cloud cover of 75%.

2. FUZZY SETS: BASIC TYPES

MEMBERSHIP FUNCTION

- A characteristic function: the values assigned to the elements of the universal set fall within a specified range and indicate the membership grade of these elements in the set.
- Larger values denote higher degrees of set membership.
- A set defined by membership functions is a **fuzzy set**.
- The most commonly used range of values of membership functions is the **unit interval** $[0,1]$.

The universal set X is always a crisp set.

Notation:

The membership function of a fuzzy set A is denoted by μ_A

$$\mu_A: X \rightarrow [0,1]$$

Alternatively, the function can be denoted by A and has the form

$$A: X \rightarrow [0,1]$$

We use the **second notation**

FUZZY SETS: BASIC TYPES





Figure 1.2 Examples of membership functions that may be used in different contexts for characterizing fuzzy sets of real numbers close to 2.

ROUGH SET Lower and Upper Approximations



FUZZY SETS: BASIC TYPES

- An example:
 - Define the seven levels of education:

0 – no education
1 – elementary school
2 – high school
3 – two-year college degree
4 – bachelor's degree
5 – master's degree
6 – doctoral degree




Figure 1.3 Examples of fuzzy sets expressing the concepts of people that are little educated (○), highly educated (●), and very highly educated (□).

FUZZY VARIABLES

- Several fuzzy sets representing **linguistic concepts** such as low, medium, high, and so on are often employed to define states of a variable. Such a variable is usually called a **fuzzy variable**.




Figure 1.4 Temperature in the range $[T_1, T_2]$ conceived as: (a) a fuzzy variable; (b) a traditional (crisp) variable.

ORDINARY FUZZY SETS

Given a universal set X , a fuzzy set is defined by a function of the form

$$A: X \rightarrow [0,1]$$

This kind of fuzzy sets are called **ordinary fuzzy sets**.

INTERVAL-VALUED FUZZY SETS

- The membership functions of ordinary fuzzy sets are often overly precise. We may be able to identify appropriate membership functions only approximately.
- **Interval-valued fuzzy sets**: a fuzzy set whose membership functions does not assign to each element of the universal set one real number, but a closed interval of real numbers between the identified lower and upper bounds.

$$A: X \rightarrow \mathcal{E}([0,1]), \quad \mathcal{E}([0,1]) \subset \mathcal{P}([0,1]).$$

Power set




Figure 1.5 An example of an interval-valued fuzzy set ($A(a) = [\alpha_1, \alpha_2]$).

- **FUZZY SETS OF TYPE 2:**

- The set of all **ordinary fuzzy sets** that can be defined with the universal set $[0,1]$.
- It is also called a fuzzy power set of $[0,1]$.




Figure 1.6 Illustration of the concept of a fuzzy set of type 2.

Discussions:

- The primary disadvantage of **interval-value fuzzy sets**, compared with **ordinary fuzzy sets**, is computationally more demanding.
- The computational demands for dealing with **fuzzy sets of type 2** are even greater than those for dealing with interval-valued fuzzy sets.
- This is the primary reason why the fuzzy sets of type 2 have almost **never been utilized in any applications**.

- Consider three fuzzy sets that represent the concepts of a young, middle-aged, and old person. The membership functions are defined on the interval $[0,80]$ as follows:

$$A_1(x) = \begin{cases} 1 & \text{when } x \leq 20 \\ (35 - x)/15 & \text{when } 20 < x < 35 \\ 0 & \text{when } x \geq 35 \end{cases}$$

$$A_2(x) = \begin{cases} 0 & \text{when either } x \leq 20 \text{ or } \geq 60 \\ (x - 20)/15 & \text{when } 20 < x < 35 \\ (60 - x)/15 & \text{when } 45 < x < 60 \\ 1 & \text{when } 35 \leq x \leq 45 \end{cases}$$

$$A_3(x) = \begin{cases} 0 & \text{when } x \leq 45 \\ (x - 45)/15 & \text{when } 45 < x < 60 \\ 1 & \text{when } x \geq 60 \end{cases}$$

Find line passing through
 (x,y) and $(20,1)$:
 $1/[35-20] = y/[35-x]$

$$A_2(x) = \begin{cases} 0 & \text{when either } x \leq 20 \text{ or } \geq 60 \\ (x - 20)/15 & \text{when } 20 < x < 35 \\ (60 - x)/15 & \text{when } 45 < x < 60 \\ 1 & \text{when } 35 \leq x \leq 45 \end{cases}$$




Figure 1.7 Membership functions representing the concepts of a young, middle-aged, and old person. Shown discrete approximation D_2 of A_2 is defined numerically in Table 1.2.

TABLE 1.2 DISCRETE APPROXIMATION OF MEMBERSHIP FUNCTION A_2 (FIG. 1.7) BY FUNCTION D_2 OF THE FORM:
 $D_2 : [0, 2, 4, \dots, 80] \rightarrow [0, 1]$

x	$D_2(x)$
$x \notin \{22, 24, \dots, 58\}$	0.00
$x \in \{22, 58\}$	0.13
$x \in \{24, 56\}$	0.27
$x \in \{26, 54\}$	0.40
$x \in \{28, 52\}$	0.53
$x \in \{30, 50\}$	0.67
$x \in \{32, 48\}$	0.80
$x \in \{34, 46\}$	0.93
$x \in \{36, 38, \dots, 44\}$	1.00

- α -cut and strong α -cut

- Given a fuzzy set A defined on X and any number $\alpha \in [0,1]$, the α -cut and strong α -cut are the **crisp sets**:

$${}^{\alpha}A = \{x|A(x) \geq \alpha\}$$

$${}^{\alpha+}A = \{x|A(x) > \alpha\}.$$

- The α -cut of a fuzzy set A is the **crisp set** that contains all the elements of the universal set X whose membership grades in A are **greater than or equal to** the specified value of α .
- The **strong α -cut** of a fuzzy set A is the **crisp set** that contains all the elements of the universal set X whose membership grades in A are **only greater than** the specified value of α .

References

1. **Op-Amps & Linear ICs (TextBook)** By Ramakanth A. Gayakwad | PHI
2. **Operational Amplifiers and Linear IC's** By David A. Bell | Oxford University Press | 3rd edition
3. **Linear Integrated Circuits (TextBook)** By D. Roy Chowdhury | New Age International(p) Ltd
4. **Operational Amplifiers & Linear Integrated Circuits: Theory & Applications** By Denton J. Daibey | TMH

- For example:




Figure 1.7 Membership functions representing the concepts of a young, middle-aged, and old person. Shown discrete approximation D_2 of A_2 is defined numerically in Table 1.2.

$${}^0A_1 = {}^0A_2 = {}^0A_3 = [0, 80] = X;$$

$${}^\alpha A_1 = [0, 35 - 15\alpha], {}^\alpha A_2 = [15\alpha + 20, 60 - 15\alpha], {}^\alpha A_3 = [15\alpha + 45, 80] \text{ for all } \alpha \in (0, 1];$$

$${}^{\alpha+}A_1 = (0, 35 - 15\alpha), {}^{\alpha+}A_2 = (15\alpha + 20, 60 - 15\alpha), {}^{\alpha+}A_3 = (15\alpha + 45, 80) \text{ for all } \alpha \in [0, 1];$$

$${}^{1+}A_1 = {}^{1+}A_2 = {}^{1+}A_3 = \emptyset.$$

- A level set of A :
 - The set of all levels $\alpha \in [0,1]$ that represent distinct α -cuts of a given fuzzy set A .

$$\Lambda(A) = \{\alpha | A(x) = \alpha \text{ for some } x \in X\}.$$

- For example:

$$\Lambda(A_1) = \Lambda(A_2) = \Lambda(A_3) = [0, 1], \text{ and}$$

$$\Lambda(D_2) = \{0, 0.13, 0.27, 0.4, 0.53, 0.67, 0.8, 0.93, 1\}.$$

$$A_2(x) = \begin{cases} 0 & \text{when either } x \leq 20 \text{ or } \geq 60 \\ (x - 20)/15 & \text{when } 20 < x < 35 \\ (60 - x)/15 & \text{when } 45 < x < 60 \\ 1 & \text{when } 35 \leq x \leq 45 \end{cases}$$

TABLE 1.2 DISCRETE APPROXIMATION
OF MEMBERSHIP FUNCTION A_2 (FIG. 1.7)
BY FUNCTION D_2 OF THE FORM:
 $D_2 : \{0, 2, 4, \dots, 80\} \rightarrow [0, 1]$

x	$D_2(x)$
$x \notin \{22, 24, \dots, 58\}$	0.00
$x \in \{22, 58\}$	0.13
$x \in \{24, 56\}$	0.27
$x \in \{26, 54\}$	0.40
$x \in \{28, 52\}$	0.53
$x \in \{30, 50\}$	0.67
$x \in \{32, 48\}$	0.80
$x \in \{34, 46\}$	0.93
$x \in \{36, 38, \dots, 44\}$	1.00

- For example: consider the discrete approximation D_2 of fuzzy set A_2





TABLE 1.2 DISCRETE APPROXIMATION
OF MEMBERSHIP FUNCTION A_2 (FIG. 1.7)
BY FUNCTION D_2 OF THE FORM:
 $D_2 : [0, 2, 4, \dots, 80] \rightarrow [0, 1]$

x	$D_2(x)$
$x \notin \{22, 24, \dots, 58\}$	0.00
$x \in \{22, 58\}$	0.13
$x \in \{24, 56\}$	0.27
$x \in \{26, 54\}$	0.40
$x \in \{28, 52\}$	0.53
$x \in \{30, 50\}$	0.67
$x \in \{32, 48\}$	0.80
$x \in \{34, 46\}$	0.93
$x \in \{36, 38, \dots, 44\}$	1.00



- The **standard complement of fuzzy set A** with respect to the universal set X is defined for all $x \in X$ by the equation

$$\bar{A}(x) = 1 - A(x)$$

- Elements of X for which $\bar{A}(x) = 1 - A(x)$ are called equilibrium points of A .
- For example, the equilibrium points of A_2 in Fig. 1.7 are 27.5 and 52.5.





Figure 1.7 Membership functions representing the concepts of a young, middle-aged, and old person. Shown discrete approximation D_2 of A_2 is defined numerically in Table 1.2.






- Given two fuzzy sets, A and B , their **standard intersection and union** are defined for all $x \in X$ by the equations

$$(A \cap B)(x) = \min[A(x), B(x)],$$

$$(A \cup B)(x) = \max[A(x), B(x)],$$

where min and max denote the minimum operator and the maximum operator, respectively.



● Another example:

- A_1, A_2, A_3 are normal.
- B and C are subnormal.
- B and C are convex.
- $B \cup C$ and $\overline{B \cup C}$ are **not** convex.








Figure 1.7 Membership functions representing the concepts of a young, middle-aged, and old person. Shows discrete approximation D_2 of A_2 is defined numerically in Table 1.2.



Normality and convexity may be lost when we operate on fuzzy sets by the standard operations of intersection and complement.

Figure 1.13 Illustration of standard operation on fuzzy sets $B = A_1 \cap A_2$ and $C = A_2 \cap A_3$ (A_1, A_2, A_3 are given in Fig. 1.7).

- Discussions:
 - Normality** and **convexity** may be **lost** when we operate on fuzzy sets by the standard operations of **intersection and complement**.
 - The fuzzy intersection and fuzzy union will satisfies **all** the properties of the Boolean lattice listed in **Table 1.1** **except** the law of contradiction and the law of excluded middle.

TABLE 1.1 FUNDAMENTAL PROPERTIES OF CRISP SET OPERATIONS

Involution	$\overline{\overline{A}} = A$
Commutativity	$A \cup B = B \cup A$ $A \cap B = B \cap A$
Associativity	$(A \cup B) \cup C = A \cup (B \cup C)$ $(A \cap B) \cap C = A \cap (B \cap C)$
Distributivity	$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
Idempotence	$A \cup A = A$ $A \cap A = A$
Absorption	$A \cup (A \cap B) = A$ $A \cap (A \cup B) = A$
Absorption by X and $\emptyset$	$A \cup X = X$ $A \cap \emptyset = \emptyset$
Identity	$A \cup \emptyset = A$ $A \cap X = A$
Law of contradiction	$A \cap \overline{A} = \emptyset$
Law of excluded middle	$A \cup \overline{A} = X$
De Morgan's laws	$\overline{A \cap B} = \overline{A} \cup \overline{B}$ $\overline{A \cup B} = \overline{A} \cap \overline{B}$

- The law of contradiction

$$A \cap \bar{A} = \phi$$

- To verify that the law of contradiction is **violated** for fuzzy sets, we need only to show that

$$\min[A(x), 1 - A(x)] = 0$$

is **violated** for at least one $x \in X$

- This is easy since the equation is obviously violated for any value
- $A(x) \in (0,1)$, and is satisfied only for $A(x) \in \{0,1\}$.

- To verify the law of absorption,

$$A \cup (A \cap B) = A$$

- This requires showing that $\max[A(x), \min[A(x), B(x)]] = A(x)$
- is satisfied for all $x \in X$.

- Consider two cases:

- 1. $A(x) \leq B(x)$ $\longrightarrow \max[A(x), \min[A(x), B(x)]] = \max[A(x), A(x)] = A(x)$

- 2. $A(x) > B(x)$ $\longrightarrow \max[A(x), \min[A(x), B(x)]] = \max[A(x), B(x)] = A(x)$

$\longrightarrow \max[A(x), \min[A(x), B(x)]] = A(x)$

- Given two fuzzy set $A, B \in \mathcal{F}(X)$,
we say that A is a subset of B and write iff $A \subseteq B$
 $A(x) \leq B(x)$

For all $x \in X$

$A \subseteq B$ iff $A \cap B = A$ and $A \cup B = B$ for any $A, B \in \mathcal{F}(X)$.

- Let $s = [i(1), i(2), \dots, i(k)]$ be a subsequence of $[1, 2, \dots, n]$ and let $s^* = [i(k+1), i(k+2), \dots, i(n)]$ be the sequence complementary to $[i(1), i(2), \dots, i(k)]$.
- The projection of n -ary fuzzy relation R on $U(s) = U(i(1)) \times U(i(2)) \times \dots \times U(i(k))$ denoted $\text{Proj}[U(s)](R)$ is k -ary fuzzy relation

$$\{((u(i(1)), u(i(2)), \dots, u(i(k))), \sup_{u(i(k+1)), u(i(k+2)), \dots, u(i(n))} \mu[R](u(1), u(2), \dots, u(n)))\}$$

Example: Let's take relation R – less than (previous page).

$$\text{Proj}[U_1](R) = \{(0, 1), (10, 0.9), (20, 0.7), (30, 0.5), \dots\}$$

- The converse of the projection of n -ary relation is called a cylindrical extension.
 - Let R be k -ary fuzzy relation on $U(s) = U(i(1)) \times U(i(2)) \times \dots \times U(i(k))$.
 - A cylindrical extension of R in $U = U(1) \times U(2) \times \dots \times U(n)$ is
- $$C(R) = \{(u(1), u(2), \dots, u(n)) : \mu[R](u(i(1)), u(i(2)), \dots, u(i(k)))\}$$

Example: Fuzzy set Fast1 on U1, Fast 2 on U2.

$U1 = U2 = \{0, 10, 20, 30, 40, 50, 60, 70, 80\}$.

$Fast1 = Fast2 = \{(0, 0), (10, 0.01), (20, 0.02), (30, 0.05), (40, 0.1), (50, 0.4), (60, 0.8), (70, 0.9), (80, 1)\}$.

$C(Fast2)$ – cylindrical extension on U1

	0	10	20	30	40	50	60	70	80
0	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1
10	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1
20	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1
30	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1
40	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1
50	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1
60	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1
70	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1
80	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1

C(Fast1) – cylindrical extension on U2

	0	10	20	30	40	50	60	70	80
0	0	0	0	0	0	0	0	0	0
10	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
20	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
30	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
40	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
50	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
60	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
70	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9
80	1	1	1	1	1	1	1	1	1

- Let R be fuzzy relation on $U(1) \times U(2) \times \dots \times U(r)$ and S be fuzzy relation on $U(s) \times U(s+1) \times \dots \times U(n)$, where $1 \leq s \leq r \leq n$.
- The **join** of R and S is defined as $c(R) \cap c(S)$, where $c(R)$, $c(S)$ are cylindrical extensions.

The join of c(Fast1) and c(Fast2)

	0	10	20	30	40	50	60	70	80
0	0	0	0	0	0	0	0	0	0
10	0	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
20	0	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02
30	0	0.01	0.02	0.05	0.05	0.05	0.05	0.05	0.05
40	0	0.01	0.02	0.05	0.1	0.1	0.1	0.1	0.1
50	0	0.01	0.02	0.05	0.1	0.4	0.4	0.4	0.4
60	0	0.01	0.02	0.05	0.1	0.4	0.8	0.8	0.8
70	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	0.9
80	0	0.01	0.02	0.05	0.1	0.4	0.8	0.9	1

Different versions of composition exist.

Let R be fuzzy relation on $U(1) \times U(2) \times \dots \times U(r)$, and S be fuzzy relation on $U(s) \times U(s+1) \times \dots \times U(n)$.

Let $\{i_1, i_2, \dots, i_k\} = (\{1, 2, \dots, r\} - \{s, s+1, \dots, n\}) \cup (\{s, s+1, \dots, n\} - \{1, 2, \dots, r\})$

Symmetric difference 

The **composition** of R and S denoted by $R \circ S$ is defined as:

$$\text{Proj}[U(i_1), U(i_2), \dots, U(i_k)](c(R) \cap c(S)).$$

Example: $R = \text{Fast} \circ \text{Less_Than}$

CONCEPTION OF FUZZY LOGIC

- Many decision-making and problem-solving tasks are too complex to be defined precisely however, people succeed by using imprecise knowledge
- Fuzzy logic resembles human reasoning in its use of approximate information and uncertainty to generate decisions.

Natural Language

- Consider:
 - Joe is tall -- what is tall?
 - Joe is very tall -- what does this differ from tall?
- Natural language (like most other activities in life and indeed the universe) is not easily translated into the absolute terms of 0 and **1**.

“false” “true”




Fuzzy Logic

- An approach to uncertainty that combines real values $[0...1]$ and logic operations
- Fuzzy logic is based on the ideas of fuzzy set theory and fuzzy set membership often found in natural (e.g., spoken) language.

Example: “Young”


- Example:
 - Ann is 28, 0.8 in set “Young”
 - Bob is 35, 0.1 in set “Young”
 - Charlie is 23, 1.0 in set “Young”
- Unlike statistics and probabilities, the *degree* is not describing *probabilities* that the item is in the set, but instead describes *to what extent* the item is the set.

Membership function of fuzzy logic




Fuzzy values have associated degrees of membership in the set.

Crisp set vs. Fuzzy set





A traditional crisp set



A fuzzy set

Crisp set vs. Fuzzy set



BENEFITS OF FUZZY LOGIC

Fuzzy logic systems has several benefits which are as follows:

- 1.Handles imprecise or noisy data:** Whether the data is distorted or not completely accurate, it can still make sense of it.
- 2.Easy to design and understand:** Unlike many complex algorithms, fuzzy logic systems are straightforward and simple to implement.
- 3.Efficient for complex problems:** Fuzzy logic mirrors human reasoning, so it's great for solving problems in many areas like decision-making or control systems.
- 4.Low memory requirements:** The algorithms are simple and can operate with little data.
 - You want the value to switch gradually as *Young* becomes *Middle* and *Middle* becomes *Old*. This is the idea of fuzzy logic.

Fuzzy Set Operations

Fuzzy union ($\cup$): the union of two fuzzy sets is the maximum (MAX) of each element from two sets.

Ex

- $A = \{1.0, 0.20, 0.75\}$
- $B = \{0.2, 0.45, 0.50\}$
- $A \cup B = \{\text{MAX}(1.0, 0.2), \text{MAX}(0.20, 0.45), \text{MAX}(0.75, 0.50)\}$
 $= \{1.0, 0.45, 0.75\}$

Fuzzy intersection ($\cap$): the intersection of two fuzzy sets is just the MIN of each element from the two sets.

- E.g.
 - $A \cap B = \{\text{MIN}(1.0, 0.2), \text{MIN}(0.20, 0.45), \text{MIN}(0.75, 0.50)\}$
 $= \{0.2, 0.20, 0.50\}$

The *complement* of a fuzzy variable with DOM x is $(1-x)$.

Complement ($_c$): The *complement* of a fuzzy set is composed of all elements' *complement*.

Example.

$$\begin{aligned} | \quad A^c &= \{1 - 1.0, 1 - 0.2, 1 - 0.75\} \\ | \quad &= \{0.0, 0.8, 0.25\} \end{aligned}$$

Crisp Relations

- Ordered pairs showing connection between two sets:
(a,b): a is related to b
(2,3) are related with the relation "<"
- Relations are set themselves
 $< = \{(1,2), (2, 3), (2, 4), \dots\}$
- Relations can be expressed as matrices

<	1	2
1	×	☺
2	×	×

Fuzzy Relations

- Triples showing connection between two sets:
 $(a, b, \#)$: a is related to b with degree $\#$
- Fuzzy relations are set themselves
- Fuzzy relations can be expressed as matrices

Fuzzy Relations Matrices

- Example: Color-Ripeness relation for tomatoes

$R_1(x, y)$	unripe	semi ripe	ripe
green	1	0.5	0
yellow	0.3	1	0.4
Red	0	0.2	1


Where is Fuzzy Logic used?

- Natural Language Processing
- Industrial Automation
- Robotics
- Decision Support System
- Image Processing

Fuzzy Expert System

- Fuzzy expert system is a collection of membership functions and rules that are used to reason about data.
- Usually, the rules in a fuzzy expert system are have the following form: *“if x is low and y is high then z is medium”*

Operation of Fuzzy System



Building Fuzzy Systems

- Fuzzification
- Inference
- Composition
- Defuzzification

Fuzzification

- Establishes the fact base of the fuzzy system. It identifies the input and output of the system, defines appropriate IF THEN rules, and uses raw data to derive a membership function.
- Consider an air conditioning system that determine the best circulation level by sampling temperature and moisture levels. The inputs are the current temperature and moisture level. The fuzzy system outputs the best air circulation level: “none”, “low”, or “high”. **The following fuzzy rules are used:**
 1. If the room is hot, circulate the air a lot.
 2. If the room is cool, do not circulate the air.
 3. If the room is cool and moist, circulate the air slightly.
- A knowledge engineer determines membership functions that map temperatures to fuzzy values and map moisture measurements to fuzzy values.

Inference

- Evaluates all rules and determines their truth values. If an input does not precisely correspond to an IF THEN rule, partial matching of the input data is used to interpolate an answer.
- Continuing the example, suppose that the system has measured temperature and moisture levels and mapped them to the fuzzy values of .7 and .1 respectively.
- The system now infers the truth of each fuzzy rule. To do this a simple method called MAX-MIN is used.
- This method sets the fuzzy value of the THEN clause to the fuzzy value of the IF clause. Thus, the method infers fuzzy values of 0.7, 0.1, and 0.1 for rules 1, 2, and 3 respectively.

Composition

- Combines all fuzzy conclusions obtained by inference into a single conclusion. Since different fuzzy rules might have different conclusions, consider all rules.
- Continuing the example, each inference suggests a different action
 - rule 1 suggests a "high" circulation level
 - rule 2 suggests turning off air circulation
 - rule 3 suggests a "low" circulation level.
- A simple MAX-MIN method of selection is used where the maximum fuzzy value of the inferences is used as the final conclusion. So, composition selects a fuzzy value of 0.7 since this was the highest fuzzy value associated with the inference conclusions.

Defuzzification

- Convert the fuzzy value obtained from composition into a “crisp” value. This process is often complex since the fuzzy set might not translate directly into a crisp value.
- Defuzzification is necessary, since controllers of physical systems require discrete signals.
- Continuing the example, composition outputs a fuzzy value of 0.7. This imprecise value is not directly useful since the air circulation levels are “none”, “low”, and “high”.
- The defuzzification process converts the fuzzy output of 0.7 into one of the air circulation levels. In this case it is clear that a fuzzy output of 0.7 indicates that the circulation should be set to “high”.

Defuzzification


- There are many defuzzification methods. Two of the more common techniques are the centroid and maximum methods.
- In the centroid method, the crisp value of the output variable is computed by finding the variable value of the center of gravity of the membership function for the fuzzy value.
- In the maximum method, one of the variable values at which the fuzzy subset has its maximum truth value is chosen as the crisp value for the output variable.

Fuzzification

- Two Inputs (x, y) and one output (z)
- Membership functions:

$$\text{low}(t) = 1 - (t / 10)$$

$$\text{high}(t) = t / 10$$



$$\text{Low}(x) = 0.68, \quad \text{High}(x) = 0.32, \quad \text{Low}(y) = 0.39, \quad \text{High}(y) = 0.61$$

Create rule base


- Rule 1: If x is low AND y is low Then z is high
- Rule 2: If x is low AND y is high Then z is low
- Rule 3: If x is high AND y is low Then z is low
- Rule 4: If x is high AND y is high Then z is high

Inference

- Rule1: $\text{low}(x)=0.68, \text{low}(y)=0.39 \Rightarrow$
 $\text{high}(z)=\text{MIN}(0.68,0.39)=0.39$ Rule strength
- Rule2: $\text{low}(x)=0.68, \text{high}(y)=0.61 \Rightarrow$
 $\text{low}(z)=\text{MIN}(0.68,0.61)=0.61$
- Rule3: $\text{high}(x)=0.32, \text{low}(y)=0.39 \Rightarrow$
 $\text{low}(z)=\text{MIN}(0.32,0.39)=0.32$
- Rule4: $\text{high}(x)=0.32, \text{high}(y)=0.61 \Rightarrow$
 $\text{high}(z)=\text{MIN}(0.32,0.61)=0.32$

Composition


- $\text{Low}(z) = \text{MAX}(\text{rule2}, \text{rule3}) = \text{MAX}(0.61, 0.32) = 0.61$
- $\text{High}(z) = \text{MAX}(\text{rule1}, \text{rule4}) = \text{MAX}(0.39, 0.32) = 0.39$



Defuzzification

Center of Gravity

$$C = \frac{\int_{Min}^{Max} tf(t)dt}{\int_{Min}^{Max} f(t)dt}$$



A Real Fuzzy Logic System

- The subway in Sendai, Japan uses a fuzzy logic control system developed by Serji Yasunobu of Hitachi.
- It took 8 years to complete and was finally put into use in 1987.

Control System

- Based on rules of logic obtained from train drivers so as to model real human decisions as closely as possible
- **Task:** Controls the speed at which the train takes curves as well as the acceleration and braking systems of the train

- The results of the fuzzy logic controller for the Sendai subway are excellent!!
- The train movement is smoother than most other trains
- Even the skilled human operators who sometimes run the train cannot beat the automated system in terms of smoothness or accuracy of stopping

Fuzzy Logic

Interpretation Domain → Fuzzy Sets

u	μ_{Fast}
0	0
10	0.01
20	0.02
30	0.05
40	0.1
50	0.4
60	0.8
70	0.9
80	1

Fuzzy set Fast

u	$\mu_{\text{Dangerous}}$
0	0
10	0.05
20	0.1
30	0.15
40	0.2
50	0.3
60	0.7
70	1
80	1

Fuzzy set Dangerous

Homework:

Find the following fuzzy logic propositions:

-X is fast and Y is dangerous


-If X is fast then Y is dangerous

Example II

if temperature is cold and oil is cheap


then heating is high

Example II



Definition [Zadeh 1973]


A **linguistic variable** is characterized by a quintuple



Example

A **linguistic variable** is characterized by a quintuple

$$(x, T(x), U, G, M)$$



Example semantic rule:


$$M(\text{old}) = \left\{ (u, \mu_{\text{old}}(u)) \mid u \in [0, 100] \right\}$$

$$\mu_{\text{old}}(u) = \begin{cases} 0 & u \in [0, 50] \\ \left[1 + \left(\frac{u-50}{5} \right)^{-2} \right]^{-1} & u \in [50, 100] \end{cases}$$

Example II

Linguistic Variable : temperature

Linguistics Terms (Fuzzy Sets) : {cold, warm, hot}



Classical Implication

$$A \rightarrow B$$

A	B	$A \rightarrow B$
T	T	T
T	F	F
F	T	T
F	F	T

A	B	$A \rightarrow B$
1	1	1
1	0	0
0	1	1
0	0	1

$$\neg A \cup B$$

A	B	$\neg A \cup B$
T	T	T
T	F	F
F	T	T
F	F	T

A	B	$\neg A \cup B$
1	1	1
1	0	0
0	1	1
0	0	1

$$A \rightarrow B$$

$$\mu_{A \rightarrow B}(x, y) = \begin{cases} 1 & \mu_A(x) \leq \mu_B(y) \\ \mu_B(y) & \text{otherwise} \end{cases}$$

A	B	$A \rightarrow B$
1	1	1
1	0	0
0	1	1
0	0	1

$$\neg A \cup B$$

$$\mu_{\neg A \cup B}(x, y) = \max(1 - \mu_A(x), \mu_B(y))$$

A	B	$\neg A \cup B$
1	1	1
1	0	0
0	1	1
0	0	1

Modus Ponens

A	B	$A \rightarrow B$
1	1	1
1	0	0
0	1	1
0	0	1

$$\frac{A \rightarrow B \quad \wedge A}{B}$$

$$\frac{\neg A \cup B \quad \wedge A}{B}$$

$$\frac{\text{If } A \text{ then } B \quad \wedge A \text{ is true}}{B \text{ is true}}$$

FUZZY IF –THEN RULES

$A \rightarrow B \equiv$ If x is A then y is B .

$\underbrace{\hspace{10em}}$
antecedent
or
premise

$\underbrace{\hspace{10em}}$
consequence
or
conclusion

Examples

$A \rightarrow B \equiv$ If x is A then y is B .

- If *pressure is high*, then *volume is small*.
- If the *road is slippery*, then *driving is dangerous*.
- If a *tomato is red*, then *it is ripe*.
- If the *speed is high*, then *apply the brake a little*.

Fuzzy Rules as Relations

$$\underbrace{A \rightarrow B}_R \equiv \text{If } x \text{ is } A \text{ then } y \text{ is } B.$$



A fuzzy rule can be defined
as a binary relation with MF

$$\underbrace{\mu_R(x, y) = \mu_{A \rightarrow B}(x, y)}$$

Depends on how
to interpret $A \rightarrow B$


$$\mu_R(x, y) = \mu_{A \rightarrow B}(x, y) = ?$$

Interpretations of $A \rightarrow B$



$$\mu_R(x, y) = \mu_{A \rightarrow B}(x, y) = ?$$

Interpretations of $A \rightarrow B$



A coupled with B (A and B)

$$R = A \rightarrow B$$

$$= \int_{X \times Y} \mu_A(x) \tilde{*} \mu_B(y) / (x, y)$$

t -norm

$$\mu_R(x, y) = \min(\mu_A(x), \mu_B(y))$$

$$\mu_R(x, y) = \mu_{A \rightarrow B}(x, y) = ?$$

Interpretations of $A \rightarrow B$

A entails B (not A or B)

- Material implication

$$R = A \rightarrow B \equiv \neg A \cup B$$

- Propositional calculus


$$R = A \rightarrow B \equiv \neg A \cup (A \cap B)$$

- Extended propositional calculus

$$R = A \rightarrow B \equiv (\neg A \cap \neg B) \cup B$$

- Generalization of modus ponens

$$\mu_R(x, y) = \begin{cases} 1 & \mu_A(x) \leq \mu_B(y) \\ \mu_B(y) & \text{otherwise} \end{cases}$$



$$\mu_R(x, y) = \mu_{A \rightarrow B}(x, y) = ?$$

Interpretations of $A \rightarrow B$

A entails B (not A or B)

- Material implication

$$R = A \rightarrow B \equiv \neg A \cup B$$

$$\mu_R(x, y) = \max(1 - \mu_A(x), \mu_B(x))$$

- Propositional calculus

$$R = A \rightarrow B \equiv \neg A \cup (A \cap B)$$

$$\mu_R(x, y) = \max(1 - \mu_A(x), \min(\mu_A(x), \mu_B(x)))$$

- Extended propositional calculus

$$R = A \rightarrow B \equiv (\neg A \cap \neg B) \cup B$$

$$\mu_R(x, y) = \max(1 - \max(\mu_A(x), \mu_B(x)), \mu_B(x))$$

- Generalization of modus ponens

$$\mu_R(x, y) = \begin{cases} 1 & \mu_A(x) \leq \mu_B(y) \\ \mu_B(y) & \text{otherwise} \end{cases}$$

Generalized Modus Ponens

Single rule with single antecedent

Rule: if x is A then y is B

Fact: x is A'

Conclusion: y is B'



Fuzzy Reasoning

Single Rule with Single Antecedent

Rule: if x is A then y is B

Fact: x is A'

Conclusion: y is B'



$$\mu_R(x, y) = \mu_A(x) \wedge \mu_B(y)$$



Single Rule with Single Antecedent

Rule: if x is A then y is B

Fact: x is A'

Conclusion: y is B'

$$\begin{aligned}\mu_{B'}(y) &= \max_x \min(\mu_{A'}(x), \mu_R(x, y)) \\ &= \vee_x (\mu_{A'}(x) \wedge \mu_R(x, y)) \\ &= \vee_x (\mu_{A'}(x) \wedge \mu_A(x) \wedge \mu_B(y)) \\ &= \underbrace{\left[\vee_x (\mu_{A'}(x) \wedge \mu_A(x)) \right]}_{\text{Firing Strength}} \wedge \mu_B(y)\end{aligned}$$



$$\mu_R(x, y) = \mu_A(x) \wedge \mu_B(y)$$



Single Rule with Single Antecedent

Rule: if x is A then y is B

Fact: x is A'

Conclusion: y is B'

$$\begin{aligned}\mu_{B'}(y) &= \max_x \min(\mu_{A'}(x), \mu_R(x, y)) \\ &= \vee_x (\mu_{A'}(x) \wedge \mu_R(x, y)) \\ &= \vee_x (\mu_{A'}(x) \wedge \mu_A(x) \wedge \mu_B(y)) \\ &= [\vee_x (\mu_{A'}(x) \wedge \mu_A(x))] \wedge \mu_B(y)\end{aligned}$$



Fuzzy Reasoning

Single Rule with Multiple Antecedents

Rule: if x is A and y is B then z is C

Fact: x is A and y is B


Conclusion: z is C

Single Rule with Multiple Antecedents

Rule: if x is A and y is B then z is C

Fact: x is A' and y is B'

Conclusion: z is C'






Single Rule with Multiple Antecedents

Rule: if x is A and y is B then z is C

Fact: x is A' and y is B'

Conclusion: z is C'



Rule: if x is A and y is B then z is C

Fact: x is A' and y is B'

Conclusion: z is C'

$$R = A \times B \rightarrow C$$

$$\mu_R(x, y, z) = \mu_{(A \times B) \times C}(x, y, z)$$

$$= \mu_A(x) \wedge \mu_B(y) \wedge \mu_C(z)$$

Max-Min Composition

$$\mu_{C'}(y) = \max_{x,y} \min(\mu_{A',B'}(x, y), \mu_R(x, y, z))$$

$$= \vee_{x,y} (\mu_{A',B'}(x, y) \wedge \mu_R(x, y, z))$$


$$= \vee_{x,y} (\mu_{A'}(x) \wedge \mu_{B'}(y) \wedge \mu_A(x) \wedge \mu_B(y) \wedge \mu_C(z))$$

$$= \left[\vee_x (\mu_A(x) \wedge \mu_{A'}(x)) \right] \wedge \left[\vee_y (\mu_B(y) \wedge \mu_{B'}(y)) \right] \wedge \mu_C(z)$$

Firing Strength

Fuzzy Reasoning

Single Rule with Multiple Antecedents



$$R = A \times B \rightarrow C$$

Rule: if x is A and y is B then z is C

Fact: x is A' and y is B'

Conclusion: z is C'

$$\mu_R(x, y, z) = \mu_{(A \times B) \times C}(x, y, z)$$


$$= \mu_A(x) \wedge \mu_B(y) \wedge \mu_C(z)$$

$$\mu_{C'}(y) = \max_{x,y} \min(\mu_{A',B'}(x, y), \mu_R(x, y, z))$$

$$C' = (A' \times B') \circ (A \times B \rightarrow C)$$

$$= \underbrace{\left[\bigvee_x (\mu_A(x) \wedge \mu_{A'}(x)) \right] \wedge \left[\bigvee_y (\mu_B(y) \wedge \mu_{B'}(y)) \right]}_{\text{Firing Strength}} \wedge \mu_C(z)$$

Firing Strength



Fuzzy Reasoning Multiple Rules with Multiple Antecedents

Rule1: if x is A_1 and y is B_1 then z is C_1

Rule2: if x is A_2 and y is B_2 then z is C_2

Fact: x is A' and y is B'

Conclusion: z is C'







Rule1: if x is A_1 and y is B_1 then z is C_1

Rule2: if x is A_2 and y is B_2 then z is C_2

Fact: x is A' and y is B'

Conclusion: z is C'

Fuzzy Reasoning– Multiple Rules with Multiple Antecedents









$C' = ?$

Rule1: if x is A_1 and y is B_1 then z is C_1

Rule2: if x is A_2 and y is B_2 then z is C_2

Fact: x is A' and y is B'

Conclusion: z is C'





$C' = ?$

Rule1: if x is A_1 and y is B_1 then z is C_1


Rule2: if x is A_2 and y is B_2 then z is C_2

Fact: x is A' and y is B'

Conclusion: z is C'



$$\begin{aligned}
 C' &= (A' \times B') \circ (R_1 \cup R_2) \\
 &= [(A' \times B') \circ R_1] \cup [(A' \times B') \circ R_2] \\
 &= C'_1 \cup C'_2
 \end{aligned}$$



Parul[®]
University

NAAC
GRADE **A++**



<https://paruluniversity.ac.in/>

