



Erratum: Guidelines for the Use of Botulinum Toxin in Otolaryngology From the Korean Society of Laryngology, Phoniatrics and Logopedics Guideline Task Force

Korean Society of Laryngology, Phoniatrics, and Logopedics Guideline Task Force;

Myung Jin Ban^{1,*} · Chang Hwan Ryu^{2,*} · Joo Hyun Woo³ · Young Chan Lee⁴ · Dong Kun Lee⁵ · Minsu Kwon⁶
Yong Tae Hong⁷ · Gil Joon Lee⁸ · Hyung Kwon Byeon⁹ · Seung Ho Choi⁶ · Seung Won Lee¹⁰

Department of Otorhinolaryngology-Head and Neck Surgery,

¹Soonchunhyang University Cheonan Hospital, Soonchunhyang University College of Medicine, Cheonan, Korea

²National Cancer Center, Goyang, Korea

³Gil Medical Center, Gachon University College of Medicine, Incheon, Korea

⁴School of Medicine, Kyung Hee University, Seoul, Korea

⁵Dong-A University College of Medicine, Busan, Korea

⁶Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

⁷Jeonbuk National University Hospital, Jeonbuk National University Medical School, Jeonju, Korea

⁸School of Medicine, Kyungpook National University, Daegu, Korea

⁹Soonchunhyang University Seoul Hospital, Soonchunhyang University College of Medicine, Seoul, Korea

¹⁰Soonchunhyang University Bucheon Hospital, Soonchunhyang University College of Medicine, Bucheon, Korea

Clinical and Experimental Otorhinolaryngology 2023;16(4):291-307.

<https://doi.org/10.21053/ceo.2023.00458>

In the article entitled “Guidelines for the Use of Botulinum Toxin in Otolaryngology From the Korean Society of Laryngology, Phoniatrics and Logopedics Guideline Task Force,” published in *Clinical and Experimental Otorhinolaryngology*, the recommendation statement for KQ 8 was incorrectly published.

The correct recommendation statement is as follows:

KQ 8. Is BT injection effective for the treatment of patients with sialorrhea?

Recommendation

Direct BT injection to the salivary gland reduces saliva flow in patients with sialorrhea. (Strong recommendation, Moderate-quality evidence)