

GE23131-Programming Using C-2024

Quiz navigation

1

2

3

Show one page at a time

Finish review

Status	Finished
Started	Monday, 23 December 2024, 5:33 PM
Completed	Tuesday, 12 November 2024, 1:56 PM
Duration	41 days 3 hours

Question **1**

Correct

Marked out of 3.00

 [Flag question](#)

A set of N numbers (separated by one space) is passed as input to the program. The program must identify the count of numbers where the number is odd number.

Input Format:

The first line will contain the N numbers separated by one space.

Boundary Conditions:

$3 \leq N \leq 50$

The value of the numbers can be from -99999999 to 99999999

Output Format:

The count of numbers where the numbers are odd numbers.

Input:

5 10 15 20 25 30 35 40 45 50

Output:

5

Explanation:

The numbers meeting the criteria are 5, 15, 25, 35, 45.

Answer: (penalty regime: 0 %)

```
1 #include<stdio.h>
2 int main(){
3     int a , count = 0;
4     char ch;
5     while (ch != '\n'){
6         scanf("%d",&a);
7         scanf("%c",&ch);
8         if(a%2!=0){
9             count++;
10        }
11    }
12    printf("%d",count);
13    return 0;
14 }
```

	Input	Expected	Got	
✓	5 10 15 20 25 30 35 40 45 50	5	5	✓

Passed all tests! ✓

Question **2**

Correct

Marked out of 5.00

🚩 [Flag question](#)

Given a number N, return true if and only if it is a *confusing number*, which satisfies the following condition:

We can rotate digits by 180 degrees to form new digits. When 0, 1, 6, 8, 9 are rotated 180 degrees, they become 0, 1, 9, 8, 6 respectively. When 2, 3, 4, 5 and 7 are rotated 180 degrees, they become invalid. A *confusing number* is a number that when rotated 180 degrees becomes a **different** number with each digit valid.

Example 1:

6 -> 9

Input: 6

Output: true

Explanation:

We get 9 after rotating 6, 9 is a valid number and 9!=6.

Input: 89

Output: true

Explanation:

We get 68 after rotating 89, 86 is a valid number and $86 \neq 89$.

Example 3:

11 -> 11

Input: 11

Output: false

Explanation:

We get 11 after rotating 11, 11 is a valid number but the value remains the same, thus 11 is not a confusing number.

Note:

1. $0 \leq N \leq 10^9$
2. After the rotation we can ignore leading zeros, for example if after rotation we have 0008 then this number is considered as just 8.

Answer: (penalty regime: 0 %)

```
1 #include<stdio.h>
2 int main(){
3     int a, b, c = 0;
4     scanf("%d",&a);
5     while(a!=0){
6         b=a%10;
7         if(b==2 || b==3 || b==4 || b==5 || b==7){
8             c++;
9         }
10        a=a/10;
11    }
```

```
12  if (c>0){
13      printf("false");
14  }else {
15      printf("true");
16  }
17  return 0;
18 }
```

	Input	Expected	Got	
✓	6	true	true	✓
✓	89	true	true	✓
✓	25	false	false	✓

Passed all tests! ✓

Question **3**

Correct

Marked out of 7.00

Flag question

A nutritionist is labeling all the best power foods in the market. Every food item arranged in a single line, will have a value beginning from 1 and increasing by 1 for each, until all items have a value associated with them. An item's value is the same as the number of macronutrients it has. For example, food item with value 1 has 1 macronutrient, food item with value 2 has 2 macronutrients, and incrementing in this fashion.

The nutritionist has to recommend the best combination to patients, i.e. maximum total of macronutrients. However, the nutritionist must avoid prescribing a particular

macronutrients that can be prescribed to a patient, without the sum matching the given 'unhealthy' number.

Here's an illustration:

Given 4 food items (hence value: 1,2,3 and 4), and the unhealthy sum being 6 macronutrients, on choosing items 1, 2, 3 -> the sum is 6, which matches the 'unhealthy' sum. Hence, one of the three needs to be skipped. Thus, the best combination is from among:

- $2 + 3 + 4 = 9$
- $1 + 3 + 4 = 8$
- $1 + 2 + 4 = 7$

Since $2 + 3 + 4 = 9$, allows for maximum number of macronutrients, 9 is the right answer.

Complete the code in the editor below. It must return an integer that represents the maximum total of macronutrients, modulo 1000000007 ($10^9 + 7$).

It has the following:

n : an integer that denotes the number of food items

k : an integer that denotes the unhealthy number

Constraints

- $1 \leq n \leq 2 \times 10^9$
- $1 \leq k \leq 4 \times 10^{15}$

The first line contains an integer, n , that denotes the number of food items.

The second line contains an integer, k , that denotes the unhealthy number.

Sample Input 0

2

2

Sample Output 0

3

Explanation 0

The following sequence of $n = 2$ food items:

1. Item 1 has 1 macronutrients.
2. $1 + 2 = 3$; observe that this is the max total, and having avoided having exactly $k = 2$ macronutrients.

Sample Input 1

2

Sample Output 1

2

Explanation 1

- 1. Cannot use item 1 because $k = 1$ and $sum \equiv k$ has to be avoided at any time.
- 2. Hence, max total is achieved by $sum = 0 + 2 = 2$.

Sample Case 2

Sample Input For Custom Testing

Sample Input 2

3

3

Sample Output 2

5

Explanation 2

Answer: (penalty regime: 0 %)

```
1 #include<stdio.h>
2 int main(){
3     long n, k, i, sum = 0;
4     scanf("%ld %ld", &n, &k);
5     for( i = 0 ; i <= n ; i++ ){
6         sum += i;
7         if( sum == k ){
8             sum -= 1;
9         }
10    }
11    printf("%ld", sum%1000000007);
12    return 0;
13 }
```

	Input	Expected	Got	
✓	2 2	3	3	✓
✓	2 1	2	2	✓
✓	3 3	5	5	✓

Finish review