



# STAT 505 - Applied Multivariate Statistical

## Start Here!

- ▶ [Welcome to STAT 505!](#)
- ▶ [Faculty login \(PSU Access Account\)](#)

## Lessons

- ▶ [Lesson 0: Matrices and Vectors](#)
- ▶ [Lesson 1: Graphical Display of Multivariate Data](#)
- ▶ [Lesson 2: Measures of Central Tendency, Dispersion and Association](#)
- ▶ [Lesson 3: Linear Combinations of Random Variables](#)
- ▶ [Lesson 4: Multivariate Normal Distribution](#)
- ▶ [Lesson 5: Sample Mean Vector and Sample Correlation and Related Inference Problems](#)
- ▶ [Lesson 6: Multivariate Conditional Distribution and Partial Correlation](#)
- ▼ [Lesson 7: Principal Components Analysis \(PCA\)](#)
  - [7.1 - Principal Component Analysis \(PCA\) Procedure](#)
  - [7.2 - How do we find the coefficients?](#)
  - [7.3 - Example: Places Rated](#)
  - [7.4 - Interpretation of the Principal Components](#)
  - [7.5 - Alternative: Standardize the Variables](#)
  - [7.6 - Example: Places Rated after Standardization](#)
  - [7.7 - Once the Components Have Been Calculated](#)
  - [7.8 - Summary](#)
- ▶ [Lesson 8: Canonical Correlation Analysis](#)
- ▶ [Lesson 9: Factor Analysis](#)
- ▶ [Lesson 10: Discriminant Analysis](#)
- ▶ [Lesson 11: Inferences Regarding Multivariate Population Mean](#)
- ▶ [Lesson 12: Cluster Analysis](#)
- ▶ [Lesson 13: Multivariate Analysis of Variance \(MANOVA\)](#)

[Home](#) // [Lesson 7: Principal Components Analysis \(PCA\)](#)

## 7.1 - Principal Component Analysis (PCA) Procedure

[Printer-friendly version](#)

Suppose that we have a random vector  $\mathbf{X}$ .

$$\mathbf{X} = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_p \end{pmatrix}$$

with population variance-covariance matrix

$$\text{var}(\mathbf{X}) = \Sigma = \begin{pmatrix} \sigma_1^2 & \sigma_{12} & \cdots & \sigma_{1p} \\ \sigma_{21} & \sigma_2^2 & \cdots & \sigma_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{p1} & \sigma_{p2} & \cdots & \sigma_p^2 \end{pmatrix}$$

Consider the linear combinations

$$\begin{aligned} Y_1 &= e_{11}X_1 + e_{12}X_2 + \cdots + e_{1p}X_p \\ Y_2 &= e_{21}X_1 + e_{22}X_2 + \cdots + e_{2p}X_p \\ &\vdots \\ Y_p &= e_{p1}X_1 + e_{p2}X_2 + \cdots + e_{pp}X_p \end{aligned}$$

Each of these can be thought of as a linear regression, predicting  $Y_i$  from  $X_1, X_2, \dots, X_p$ . There is no intercept, but  $e_{i1}, e_{i2}, \dots, e_{ip}$  can be viewed as regression coefficients.

Note that  $Y_i$  is a function of our random data, and so is also random. Therefore it has a population variance

$$\text{var}(Y_i) = \sum_{k=1}^p \sum_{l=1}^p e_{ik}e_{il}\sigma_{kl} = \mathbf{e}_i' \Sigma \mathbf{e}_i$$

► [Lesson 14: Repeated Measures Analysis](#)

## Resources

- ◉ [SAS Tour](#)
- ◉ [SAS Interactive Data Analysis](#)
- ◉ [A Quick Introduction to Minitab](#)
- ◉ [Worked Examples from the Course That Use Software](#)

Moreover,  $Y_i$  and  $Y_j$  will have a population covariance

$$\text{cov}(Y_i, Y_j) = \sum_{k=1}^p \sum_{l=1}^p e_{ik} e_{jl} \sigma_{kl} = \mathbf{e}_i' \Sigma \mathbf{e}_j$$

Here the coefficients  $e_{ij}$  are collected into the vector

$$\mathbf{e}_i = \begin{pmatrix} e_{i1} \\ e_{i2} \\ \vdots \\ e_{ip} \end{pmatrix}$$

### ***First Principal Component (PCA1): $Y_1$***

The *first principal component* is the linear combination of x-variables that has maximum variance (among all linear combinations), so it accounts for as much variation in the data as possible.

Specifically we will define coefficients  $e_{11}, e_{12}, \dots, e_{1p}$  for that component in such a way that its variance is maximized, subject to the constraint that the sum of the squared coefficients is equal to one. This constraint is required so that a unique answer may be obtained.

More formally, select  $e_{11}, e_{12}, \dots, e_{1p}$  that maximizes

$$\text{var}(Y_1) = \sum_{k=1}^p \sum_{l=1}^p e_{1k} e_{1l} \sigma_{kl} = \mathbf{e}_1' \Sigma \mathbf{e}_1$$

subject to the constraint that

$$\mathbf{e}_1' \mathbf{e}_1 = \sum_{j=1}^p e_{1j}^2 = 1$$

### ***Second Principal Component (PCA2): $Y_2$***

The *second principal component* is the linear combination of  $x$ -variables that accounts for as much of the remaining variation as possible, with the constraint that the correlation between the first and second component is 0

Select  $\mathbf{e}_{21}, \mathbf{e}_{22}, \dots, \mathbf{e}_{2p}$  that maximizes the variance of this new component...

$$\text{var}(Y_2) = \sum_{k=1}^p \sum_{l=1}^p e_{2k} e_{2l} \sigma_{kl} = \mathbf{e}_2' \Sigma \mathbf{e}_2$$

subject to the constraint that the sums of squared coefficients add up to one,

$$\mathbf{e}_2' \mathbf{e}_2 = \sum_{j=1}^p e_{2j}^2 = 1$$

along with the additional constraint that these two components will be uncorrelated with one another.

$$\text{cov}(Y_1, Y_2) = \sum_{k=1}^p \sum_{l=1}^p e_{1k} e_{2l} \sigma_{kl} = \mathbf{e}_1' \Sigma \mathbf{e}_2 = 0$$

All subsequent principal components have this same property – they are linear combinations that account for as much of the remaining variation as possible and they are not correlated with the other principal components

We will do this in the same way with each additional component. For instance:

***i<sup>th</sup> Principal Component (PCA<sub>i</sub>):  $Y_i$***

We select  $\mathbf{e}_{i1}, \mathbf{e}_{i2}, \dots, \mathbf{e}_{ip}$  that maximizes

$$\text{var}(Y_i) = \sum_{k=1}^p \sum_{l=i}^p e_{ik} e_{il} \sigma_{kl} = \mathbf{e}_i' \Sigma \mathbf{e}_i$$

subject to the constraint that the sums of squared coefficients add up to one...along with the additional constraint that this new component will be uncorrelated with all the previously defined components.

$$\mathbf{e}_1' \mathbf{e}_1 = \sum_{j=1}^p e_{1j}^2 = 1$$

$$\text{textcov}(Y_1, Y_i) = \sum_{k=1}^p \sum_{l=i}^p e_{1k} e_{il} \sigma_{kl} = \mathbf{e}_1' \Sigma \mathbf{e}_i = 0,$$

$$\text{cov}(Y_2, Y_i) = \sum_{k=1}^p \sum_{l=i}^p e_{2k} e_{il} \sigma_{kl} = \mathbf{e}_2' \Sigma \mathbf{e}_i = 0,$$

$$\vdots$$

$$\text{cov}(Y_{i-1}, Y_i) = \sum_{k=1}^p \sum_{l=i}^p e_{i-1,k} e_{il} \sigma_{kl} = \mathbf{e}_{i-1}' \Sigma \mathbf{e}_i = 0$$

Therefore all principal components are uncorrelated with one another.

[◀ Lesson 7: Principal Components Analysis \(PCA\)](#)

[up](#)

[7.2 - How do we find the coefficients? ▶](#)

 [Printer-friendly version](#)



© 2014 The Pennsylvania State University. All rights reserved.