1. The probability of a leap year selected at random contain 53 Sunday is:				
(a) 53/ 366 (b) 1/7 (<mark>c) 2/7</mark> (d) 53/365				
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at				
random. The probability of drawing a black ball is:				
(a) 3/5 (b) 2/5 (<mark>c) 0/5</mark> (d) 1/5				
3. The probability that it will rain tomorrow is 0.85. What is the				
probability that it will not rain tomorrow				
(a) 0.25 (b) 0.145 (c) 3/20 (d) none of these 4.				
What is the probability that a number selected from the				
numbers (1, 2, 3,,15) is a multiple of 4?				
(<mark>a) 1/5</mark> (b) 4/5 (c) 2/15 (d) 1/3				
5. What are the total outcomes when we throw three coins?				
(a) 4 (b) 5 (<mark>c) 8</mark> (d) 7				
6. The probability that a prime number selected at random from the				
numbers (1,2,3,35) is :				
(a) 12/35 (b <mark>) 11/35</mark> (c) 13/35 (d) none of these 7.	The			
sum of the probability of an event and non event is:				
(a) 2 (<mark>b) 1</mark> (c) 0 (d) none of these.				
8. The following probabilities are given; choose the correct answer for that which is not possible.				
(a) 0.15 (b) 2/7 (<mark>c) 7/5</mark> (d) none of these. 9. If				
three coins are tossed simultaneously, than the probability of getting a	it			
least two heads, is:				
(a) 1/4 (b) 3/8 (c <mark>) ½</mark> (d) 1/8				
10. A letter is chosen at random from the letters of the word				
♦ SSASSINATION ♦ The probability that the letter chosen has:				
(a <mark>) 6/13</mark> (b) 7/13 (c) 1 (d) none of these.				
11. A dice is thrown. Find the probability of getting an even number.				
(A) 2/3 (B) 1 (C) 5/6 (D) 1/2				
12. Two coins are thrown at the same time. Find the probability of gett	ing			

both heads.

of 9 is: (A) 1/10 14. 100 car	ce are throw (B) 3/10	ın simultaneo			
(A) 1/10 14. 100 car	(B) 3/10	in Simultance	ously. The pro	bability of getti	ing a sum
14. 100 car	(B) 3/10				
	(b) 3/10) (C <mark>) 1/</mark>	<mark>/9</mark> (D)	4/9	
		pered from 1	to 100. Find t	he probability o	of getting
a prime nu	mber.				
(A) 3/4	(B) 27/50	(<mark>C) 1/4</mark>	l ([0) 29/100	
balls in a b			(D) 20	en the number	of blue
(A) 143/150) (B <mark>) 1</mark>	<mark>.47/150</mark>	(C) 1/25	(D) 1/50	
thorou	ughly. One c bility that th	ard is drawn ne number on	from this box	ced in a box an randomly, the fect square. (A <mark>)</mark>	n the
(D) 1/.					
. , .	is the proba	bility of getti	ng 53 Monda	ys in a leap yea	r?
. , .		bility of getti 5 (C <mark>) 2/7</mark>			r?

(A) 1/6	(B <mark>) 1/12</mark>	(C) 7/12	(D) 5/12			
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that						

(A) 3/4(B) 1/2 (C) 1 (D) 1/4

Aryan will lose the game.

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:

(A) 364/365 (B) 31/365 (C<mark>) 1/365</mark> (D) 1/133225

23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is?

(C<mark>) 3/5</mark> (A) 1/5(B) 2/5(D) 4/5

24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (C) 8 (D) 7

25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is:

(D) 2/25(A) 7/50 (B) 4/25 (C) 1/25

26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?

b) $\frac{5}{42}$ d) $\frac{4}{21}$ c)

27. Runs scored by batsman in 5 one day matches are 50, 70, 82, 93, and 20. The standard deviation is _____.

c) 25.29 a<mark>) 25.79</mark> b) 25.49 d) 25.69

days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
a) 13, 15	b) <mark>13, 18</mark>	c) 18, 15		d) 13, 16		
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is						
a) $^{1}/^{2}$	b) $^{1}/3$	c <mark>)</mark>	<mark>1_{/4}</mark>	d) $^{1}/6$		
30. X is a variate between 0 and 3. The value of E(X²) is						
a) 8	b) 7	c) 27	d <mark>) 9</mark>			
31. The random variables X and Y have variances 0.2 and 0.5 respectively.						

Let Z= 5X-2Y. The variance of Z is?

28. Find median and mode of the messages received on 9 consecutive

a) 3	b) 4	c) 5	d <mark>) 7</mark>		
32.Out of the following values, which one is not possible in probability?					
a) $P(x) = 1$	b) ∑ x	P(x) = 3			
c) $P(x) = 0.5$	5 d <mark>) P(</mark> 3	(x) = -0.5			
22.67()	.	I =/	٠		
	= 2 and E(z) = 4				
a) 2	b) 6	•	•	ufficient data	
		-	t random varia		
•	<u>-</u>	•	d) Und	defined	
35.If Σ P(x)	$= k^2 - 8 $ then		k is?		
a) 0	b) 1	c <mark>) 3</mark>	d) Ins	ufficient data	
36.If P(x) =	0.5 and x = 4,	then $E(x) = ?$			
a) 1	b) 0.5	c) 4	d) <mark>2</mark>		
37.In a discalways?	crete probabil	ity distributio	on, the sum of a	ll probabilities is	
a) 0	b) Infinite	c <mark>) 1</mark>	d) Und	defined	
38.If the p a) 0.4, 0.24			get is 0.4, find r c) 0.4, 0.16	nean and variance. d) 0.6, 0.16	
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6					
40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1					
•	s the mean ar	nd variance fo	or standard norr	mal distribution?	
a) Mean is	0 and variance	<mark>e is 1</mark> b) Mea	n is 1 and variar	nce is 0	
c) Mean is	0 and variance	e is ∞ d) Mea	an is ∞ and varia	ance is 0	

42. Variance of a random variable X is given by _ d) (E(X))2 a) E(X) c) E(X2) - (E(X))2b) E(X2) 43. Mean of a random variable X is given by ____ c) E(X2) - (E(X))2d) (E(X))2 a) E(X) b) E(X2) 44. Mean of a constant 'a' is _____ . c) a/2 a) 0 b) a d) 1 45. Variance of a constant 'a' is ______. c) a/2 a) 0 b) a d) 1 46. Find the mean and variance of X?

Х	0	1	2	3	4
f(x)	1/9	2/9	3/9	2/9	1/9


a) 2, 4/3

b) 3, 4/3

c) 2, 2/3

d) 3, 2/3

47. Find the expectation of a random variable X?



d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

a١	nn			

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use Binomial Distribution

a)
$$P(X = x) = nCx px qx$$

b)
$$P(X = x) = nCx px q(n-x)$$

c)
$$P(X = x) = xCn qx p(n-x)$$

d)
$$P(x = x) = xCn pn qx$$

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard Deviation?

a)
$$\sqrt{np}$$

a)
$$\sqrt{np}$$
 b) \sqrt{pq} c) (np)2 d) \sqrt{npq}

d)
$$\sqrt{npq}$$