Active sensing explains cortical variability and improves visual information

UCLA Stein Eye Institute · UC Berkeley

Declan P. Rowley[2], Ryan A. Ressmeyer[1], Tejasvi Kothapalli[3,4], Alexander C. Huk[2], Jude F. Mitchell[5], Daniel A. Butts[6], Jacob L. Yates[3,4]

[1]Department of Bioengineering, University of Washington; [2]Fuster Laboratory for Cognitive Neuroscience, Departments of Psychiatry & Biobehavioral Sciences and Ophthalmology, UCLA
[3]Redwood Center for Theoretical Neuroscience, UC Berkeley; [4]Herbert Wertheim School of Optometry and Vision Science, UC Berkeley
[5]Brain and Cognitive Science, University of Rochester; [6]Department of Biology, University of Maryland, College Park

Read the preprint


Fixational eye movements: noise or signal?

Even during fixation the eyes drift and make microsaccades, so each repeat of an image lands on a different part of the foveal receptive fields.

Noise view
A repeated stimulus is a repeated input, so trial-to-trial variability is noise to be averaged away.


Active-sensing view
Eye motion turns space into time and is part of how the fovea encodes fine detail[5,6].

Q1 How much V1 "noise" do fixational eye movements drive? → 1–5
Q2 Does that activity carry visual information? → 6–8



Median gaze on 88 repeats (navy: one path) vs a typical foveal V1 RF (0.42°)


1 Gaze shifts within an RF change population responses



Gaze shift 0.33° — smaller than a typical receptive field (median diameter 0.42°, 96 units)


Dots: each trial's median gaze in the shaded window. Circles: population RF at each group's mean gaze.
One session, 132 units; same image sequence every trial. Lines: mean rate per unit.
Dataset: 2 marmosets · 1,022 foveal V1 units · 19 sessions · 20 Hz natural images

2 Small eye movements structure foveal V1 activity




Colors: eye position projected onto the RF axis.

3 FEMs dominate conditional rate variance




4 FEM correction reduces variability and shared noise




19 sessions; 25-ms bins. Points: mean ± SD. Gray bands: 95% shuffle null; violins: pairwise correlations.
Corrected mean noise correlation: 95% CI −0.010 to 0.029. Both tests compare correction with shuffle.

5 FEM covariance is compact and aligned with stimulus-driven dimensions



25-ms counts; alignment uses four leading dimensions per component.
Dots: sessions · large dots: mean ± SD
Gray bands: 95% shuffle-null intervals

6 Gaze-dependent retinal input predicts single-trial responses




Train: free viewing · Test: held-out flashed images
Same fitted weights in all conditions

Median fraction of conditional rate variance explained (962 units · 19 sessions)	Trial average	Full	Retinal only	Stabilized
	0.139	0.235	0.226	0.042


Triangles: medians; band: empirical ±0.1.

7 Fixational eye movements sharpen spatial coding in the model



8 Fixational motion engages tuning and sharpens spatial coding

725 units



Tuning engagement: within-unit percentile of motion-induced tuning-relevant power.
All quintiles/stages > 0 (Holm-adjusted p < 0.05).
Gray: median / IQR, 5th–95th whiskers. Purple: median and 95% bootstrap CI.

References
[1] McFarland 2016 · [2] Yates 2023 · [3] Wu 2023
[4] Ressmeyer 2026 · [5] Rucci & Victor 2015 · [6] Kuang 2012

Funding
NEI K99-EY037839-01 (D.P.R.); R01-EY030998 (J.F.M.); R01-EY018363 (M.R.).
Fuster Endowment (A.C.H.); NEI K99/R00-EY032179 (J.L.Y.); NIH/NEI T32-EY007043 (T.K.).

Contact
Declan P. Rowley · UCLA
declanrowley@g.ucla.edu