

Member ID: _____

Time: _____

Rank: _____



ICD-10-CM MEDICAL DIAGNOSTIC CODING (600)

NATIONAL 2022

CONCEPT KNOWLEDGE:

Multiple Choice (15 @ 3 points each) _____ (45 points)

APPLICATION KNOWLEDGE:

Coding (10 @ 10 points each) _____ (100 points)

TOTAL POINTS _____ **(145 points)**

Test Time: 60 minutes

GENERAL GUIDELINES:

Failure to adhere to any of the following rules will result in disqualification:

1. Member must hand in this test booklet and all printouts if any. Failure to do so will result in disqualification.
2. No equipment, supplies, or materials other than those specified for this event are allowed in the testing area. No previous BPA tests and/or sample tests (handwritten, photocopied, or keyed) are allowed in the testing area.
3. Electronic devices will be monitored according to ACT standards.

Directions: Select the letter that best answers the question.

1. A combination code is a single code used to classify _____.
 - A. two diagnoses
 - B. a diagnosis with an associated secondary process
 - C. a diagnosis with an associated complication
 - D. all of these

2. When a pregnant woman with asymptomatic HIV presents for a health care encounter, the correct sequence is:
 - A. O98.7-, B20
 - B. O98.7-, Z21
 - C. B20, O98.7-
 - D. Z21, O98.7-

3. The coding of severe sepsis requires a minimum of _____ codes.
 - A. 4
 - B. 3
 - C. 2
 - D. 1

4. The correct sequence of codes for a patient treated for anemia associated with an adverse effect from chemotherapy is _____.
 - A. Anemia, Neoplasm, Adverse Effect
 - B. Adverse Effect, Neoplasm, Anemia
 - C. Neoplasm, Anemia, Adverse Effect
 - D. Anemia, Adverse Effect, Neoplasm

5. If the type of diabetes mellitus is not documented in the medical record the default is _____.
 - A. E10.-
 - B. E08.-
 - C. E13.-
 - D. E11.-

6. When coding disorders related to psychoactive substance use, if both use and dependence are documented, assign _____.
 - A. only the code for abuse
 - B. only the code for dependence
 - C. both use and dependence
 - D. only the code of substance use

7. The time frame that defines when pain becomes chronic and classifiable to G89.2 is _____.
 - A. six months
 - B. one year
 - C. no time frame
 - D. after the acute phase
8. A pressure ulcer (L89.-) with full thickness skin loss involving damage or necrosis of subcutaneous tissue would be classified as _____.
 - A. Stage 4
 - B. Stage 3
 - C. Stage 2
 - D. Stage 1
9. Chapter 20 External Causes of Morbidity codes may be sequenced as the first-listed or principal diagnosis.
 - A. True
 - B. False
10. The appropriate 7th character for subsequent care of complications of fractures involving malunion is:
 - A. P, Q, or R
 - B. K, M, or N
 - C. A, B, or C
 - D. B, E, or H
11. The ICD-10-CM classifies Chronic Kidney Disease based on _____.
 - A. severity
 - B. type
 - C. category
 - D. duration
12. The second trimester of pregnancy includes weeks 14 to _____.
 - A. 20
 - B. 26
 - C. 28
 - D. 24
13. The perinatal period is defined as before birth through the _____ day following birth.
 - A. 14th
 - B. 21st
 - C. 30th
 - D. 28th

14. Signs or symptoms that are associated routinely with a disease process should be assigned as additional codes, unless otherwise instructed by the classification.
- A. True
 - B. False
15. Burns are classified by depth as first, second, and third degree. Second degree burns involve _____.
- A. blistering
 - B. full-thickness
 - C. erythema
 - D. all are correct

Directions: Using the ICD-10-CM manual, assign codes to the following:

16. Allergic reaction to ibuprofen that has induced Stevens-Johnson Syndrome. _____
17. Second degree burn of the right forearm, total body surface area is 4%. _____
18. 37-year-old mother of two, in her second trimester, is seen for a regular prenatal visit. _____
19. Acute cholangitis with gallstones. _____
20. Patient experiencing febrile seizures. After exam and laboratory results, patient it is determined that the febrile seizures are due to Pneumonia due to *Staphylococcus Aureus*. _____
21. Patient is seen by cardiology for surgical clearance for a Type 3 Salter-Harris fracture of the left humerus. The patient is known to have atrial fibrillation that is controlled by anticoagulants. _____
22. Facial droop due to intracerebral hemorrhage. _____
23. Acute myelomonocytic leukemia, in remission. _____
24. Cystitis, urethritis, and periurethral abscess due to gonorrhea. _____
25. Spondylitis due to brucellosis of the sacrococcygeal region. _____