

Member ID: \_\_\_\_\_

Time: \_\_\_\_\_

Rank: \_\_\_\_\_



# ICD-10-CM MEDICAL DIAGNOSTIC CODING (600) **NATIONAL 2023**

**Multiple Choice and Practical Application:**

Multiple Choice (25 @ 4 points each) \_\_\_\_\_ (100 points)

Practical Application (25 @ 4 points each) \_\_\_\_\_ (100 points)

*Total Points* \_\_\_\_\_ (200 points)

**Test Time: 60 minutes**

**GENERAL GUIDELINES:**

*Failure to adhere to any of the following rules will result in disqualification:*

1. Member must hand in this test booklet and all printouts if any. Failure to do so will result in disqualification.
2. No equipment, supplies, or materials other than those specified for this event are allowed in the testing area. No previous BPA tests and/or sample tests (handwritten, photocopied, or keyed) are allowed in the testing area.
3. Electronic devices will be monitored according to ACT standards.

**Directions:** Identify the letter of the choice that *best* completes the statement or answers the question.

1. Which punctuation is used to indicate nonessential modifiers?
  - A. Brackets
  - B. Colons
  - C. Semicolons
  - D. Parentheses
2. A subsequent acute myocardial infarction code (category I22) is reported when the patient has a new type 1 acute myocardial infarction within what time period from the original infarction?
  - A. 48 hours
  - B. 2 weeks
  - C. 4 weeks
  - D. 8 weeks
3. Each ICD-10-PCS code contains nine digits.
  - A. True
  - B. False
4. Which code set is used to report procedures performed in acute inpatient hospitals?
  - A. CPT
  - B. HCPCS
  - C. ICD-10-PCS
  - D. ICD-10-CM
5. What type of codes are used when the medical record provides detail of the condition, and an ICD-10-CM code does not exist?
  - A. "Other" codes
  - B. "And" codes
  - C. "Excludes" codes
  - D. "Unlisted" codes
6. A diagnosis-related group is used to determine payment to hospitals for treatment of a patient's principal diagnosis and comorbidities.
  - A. True
  - B. False
7. What does ICD-10-CM Guideline I.C.1.a.2.b. refer to?
  - A. Selection of codes for a patient with inconclusive HIV serology.
  - B. Sequencing of codes for a patient admitted for a HIV-related condition.
  - C. Selection of codes for a patient with a HIV infection during pregnancy.
  - D. Sequencing of codes for a patient with HIV disease admitted for an unrelated condition.
8. Codes from category O09, Supervision of high-risk pregnancy, are used during which type of encounter?
  - A. During the postnatal period.
  - B. During the prenatal period.
  - C. During the delivery.
  - D. During the puerperium period.

9. Select the type of code which describes how the injury or health condition happened to the patient.
  - A. Diagnostic codes
  - B. Radiologic codes
  - C. Procedure codes
  - D. External cause codes
10. When a patient takes a medication as prescribed and has a side effect, which column would you use in the Table of Drugs and Chemicals to select the appropriate code?
  - A. Poisoning, Accidental
  - B. Poisoning, Intentional
  - C. Adverse Effect
  - D. Underdosing
11. What is selected when a patient has sepsis and associated acute organ dysfunction?
  - A. Severe sepsis
  - B. Urosepsis
  - C. Septic shock
  - D. Sepsis with organ dysfunction
12. Select the most appropriate codes for a pathologic fracture of distal left femur due to osteosarcoma of femur.
  - A. M84.552G, C40.10
  - B. M84.551A, C40.22
  - C. M84.352A, C40.21
  - D. M84.552A, C40.22
13. Select the most appropriate codes for a patient with a family history of rheumatoid arthritis is being screened for rheumatoid arthritis.
  - A. Z13.820, Z82.69
  - B. Z13.89, Z82.62
  - C. Z13.828, Z82.61
  - D. Z13.828
14. The coder should rely solely on the discharge summary to capture all diagnoses and procedures that were treated and performed.
  - A. True
  - B. False

15. The patient presented to the emergency department with a chief complaint of hematuria, sharp and sudden pain in the lower abdomen, and vomiting for 24 hours. After examination, labs and abdominal ultrasound, the patient was admitted to the hospital to undergo lithotripsy for renal calculi. Select the appropriate diagnoses which would be submitted for this patient stay.
  - A. Hematuria
  - B. Renal calculi
  - C. Hematuria, abdominal pain, and vomiting
  - D. Renal calculi and lithotripsy
16. A gravid patient at 11 weeks' gestation presents for a routine prenatal visit. This patient had preeclampsia in her previous pregnancy, and the physician documents high-risk pregnancy. She has blood work done on this visit and is found to be anemic. The physician prescribes an iron supplement.
  - A. O09.891, O99.019, D50.9, Z3A.11
  - B. O09.91, O99.011, D64.9, Z3A.11
  - C. O09.891, O99.011, D64.9, Z3A.11
  - D. O09.91, O99.019, D50.9, Z3A.11
17. An infant is born by cesarean section. On exam, the infant is noted to have neonatal teeth and a strawberry nevus on the left leg.
  - A. Z38.01, K00.9, Q82.5
  - B. Z38.00, K00.9, Q82.8
  - C. Z38.01, K00.6, Q82.5
  - D. Z38.00, K00.6, Q82.8
18. A patient with known metastatic cancer to the ribs is admitted for pain control. The patient has a history of lung cancer, which was surgically excised 3 years ago. The patient is administered fentanyl.
  - A. C79.51, G89.3, Z85.118
  - B. G89.3, C79.51, Z85.118
  - C. C79.51, G89.3, C34.90
  - D. G89.3, G89.3, C34.90
19. A patient combined prescription Percodan with beer and became stuporous and drowsy.
  - A. T40.2x1A, T51.0x1A
  - B. T40.2x1D, T51.0x1D, R40.1, R40.0
  - C. T40.2x1S, T51.0x1S, R40.1, R40.0
  - D. T40.2x1A, T51.0x1A, R40.1, R40.0
20. How are codes under subcategory T86 are used?
  - A. Only for complications of transplanted organs
  - B. Only for rejection of transplanted organs
  - C. For both complications and rejection of transplanted organs
  - D. For neither complications nor rejection of transplanted organs

21. An obstetric patient is 13 weeks 6.5 days. Select the appropriate weeks of gestation code.
- A. Z3A.00
  - B. Z3A.13
  - C. Z3A.14
  - D. Z3A.23
22. When does the perinatal period end?
- A. 28 days after birth.
  - B. 42 days after birth.
  - C. 30 days after birth.
  - D. 90 days after birth.
23. A quadruplet child is delivered by cesarian section. Select the code to be applied to the child's record.
- A. Z38.01
  - B. Z38.66
  - C. Z38.64
  - D. Z37.53
24. When coding sequela of cerebrovascular disease which includes paralysis, what is the correct coding selection when the affected side is documented but not specified as dominant or nondominant?
- A. If the left side is affected, the default is dominant.
  - B. If the right side is affected, the default is nondominant.
  - C. If both sides are affected, the default is dominant.
  - D. If the right side is affected, the default is dominant.
25. Codes from Chapter 17 (Q00-Q99) should be identified and reported only time at the birth of the child and not during the life of the child.
- A. True
  - B. False

## PRACTICAL APPLICATION

**Directions:** Use ICD-10-CM coding manuals to answer the following questions. Do not code external cause codes. All answers must be legible.

Format your answers as follows:

Questions requiring more than one code:     Code 1, Code 2

1. Patient is diagnosed with juvenile pauciarticular rheumatoid arthritis in the right knee related to left-sided ulcerative colitis.

---

2. Patient's mother brings patient to pediatrician because of fever, severe cough, and wheezing. After examination, physician documents a diagnosis of bronchiolitis due to RSV virus.

---

3. After extensive examination and testing, female patient is diagnosed with lower-inner quadrant left breast cancer with metastasis to parietal lobe of brain.

---

4. Patient with toxic shock syndrome due to *Staphylococcus aureus*.

---

5. Diabetic patient, on oral antidiabetic medication, has proliferative retinopathy and macular edema in right eye.

---

6. Child presents with hives on skin after being given mother's leftover amoxicillin.

---

7. Patient presents to clinic for repair of skin lacerations to left proximal posterior forearm.

---

8. Patient undergoes coronary artery bypass using two arterial grafts due to coronary artery disease with unstable angina.

---

9. Patient with filarial orchitis.

---

10. During a diagnostic arthroscopy of the right knee after a sports injury, the surgeon repairs a peripheral tear in the lateral meniscus.

---

11. Patient is admitted for radiation therapy to treat cancer of the tail of the pancreas.

---

12. Patient with alcoholic cirrhosis of the liver with bleeding esophageal varices was admitted to hospital. There was no evidence of withdrawal syndromes due to the patient's alcohol dependence.

---

13. Patient is diagnosed with hypertensive chronic systolic and diastolic heart failure and Stage 5 chronic kidney disease. Patient requires regular hemodialysis.

---