

# Decentralized Online Optimization

December 23, 2018

## Abstract

ddd

## 1 Notations and assumptions

$$\mathcal{R}_T^{DOG} = \sum_{i=1}^n \sum_{t=1}^T \mathbb{E}_{\xi_{i,t} \sim D_{i,t}} f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - \mathbb{E}_{\xi_{i,t} \sim D_{i,t}} f_{i,t}(\mathbf{x}_t^*; \xi_{i,t}),$$

where, for any  $\mathbf{x}$ ,

$$f_{i,t}(\mathbf{x}; \xi_{i,t}) := \beta g_{i,t}(\mathbf{x}) + (1 - \beta) h_t(\mathbf{x}; \xi_{i,t})$$

with  $0 < \beta < 1$

Define the dynamic regret as

$$\mathcal{R}_T^{DOG} = \sum_{i=1}^n \sum_{t=1}^T \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} f_{i,t}(\mathbf{x}_t^*; \xi),$$

where, for any  $\mathbf{x}$ ,

$$f_{i,t}(\mathbf{x}; \xi) := \beta g_{i,t}(\mathbf{x}) + (1 - \beta) h_t(\mathbf{x}; \xi)$$

with  $0 < \beta < 1$ , and  $\xi$  is a random variable drawn from an unknown distribution.  $g_{i,t}$  is an adversary loss function, and  $f$  is a given loss function.

The budget of the dynamics is defined as

$$\sum_{t=1}^T \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\| \leq D.$$

**Assumption 1.** *We make the following assumptions.*

- For any  $i \in [n]$  and  $t \in [T]$ , we assume  $\max \left\{ \|\nabla g_{i,t}(\mathbf{x})\|^2, \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \|\nabla h_t(\mathbf{x}; \xi_{i,t})\|^2, \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \|\nabla f_{i,t}(\mathbf{x}; \xi_{i,t})\|^2 \right\} \leq G$ , and  $\mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \|\nabla h_t(\mathbf{x}; \xi_{i,t}) - \nabla h_t(\mathbf{x})\|^2 \leq \sigma_{i,t}^2$ .
- For any  $\mathbf{x}$  and  $\mathbf{y}$ , we assume  $\|\mathbf{x} - \mathbf{y}\|^2 \leq R$ .
- For any  $i \in [n]$  and  $t \in [T]$ , we assume the function  $f_{i,t}$  is convex and differentiable, and the function  $h_t$  has  $L$  Lipschitz gradients.

---

**Algorithm 1** DOG: Decentralized Online Gradient.

---

**Require:** The learning rate  $\eta$ , number of iterations  $T$ , and the confusion matrix  $\mathbf{W}$ .

- 1: **for**  $t = 1, 2, \dots, T$  **do**  
    For the  $i$ -th node with  $i \in [n]$ :
  - 2:     Predict  $\mathbf{x}_{i,t}$ .
  - 3:     Observe the loss function  $f_{i,t}$ ,  
        and suffer loss  $f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})$ .
  - Update:
  - 4:     Query the gradient  $\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})$ .
  - 5:      $\mathbf{x}_{i,t+1} = \sum_{j=1}^n \mathbf{W}_{i,j} \mathbf{x}_{j,t} - \eta \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})$ .
- 

## 2 Algorithm

**Theorem 1.** Using Assumptions 1, and choosing  $\eta > 0$  in Algorithm 1, we have

$$\begin{aligned} & \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \sum_{t=1}^T \sum_{i=1}^n f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - f_t(\mathbf{x}_t^*; \xi_{i,t}) \\ & \leq \eta n T G \left( 21\beta + \frac{6\eta}{(1-\rho)^2} \left( L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) + 3L\eta \left( \frac{1}{\beta} + 4 \right) \right) + \left( \frac{1}{\beta} + 4 \right) n (h_t(\bar{\mathbf{x}}_1) - h_t(\bar{\mathbf{x}}_{T+1})) \\ & \quad + \left( \left( L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) \frac{4\eta^2}{(1-\rho)^2} + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n\beta} \right) \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{RD} + R). \end{aligned}$$

**Corollary 1.** Using Assumptions 1, and choosing

$$\eta = \sqrt{\frac{nD}{T(1+n\beta)}}$$

in Algorithm 1, we have

$$\mathcal{R}_T^{DOG} \leq \mathcal{O} \left( \sqrt{(1+n\beta)nDT} \right).$$

## Appendix

**Proof to Theorem 1:**

*Proof.*

$$\begin{aligned} & \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \frac{1}{n} \sum_{i=1}^n f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - f_t(\mathbf{x}_t^*; \xi_{i,t}) \\ & = \frac{1}{n} \sum_{i=1}^n \beta (g_{i,t}(\mathbf{x}_{i,t}) - g_{i,t}(\mathbf{x}_t^*)) + (1-\beta) \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} (h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - h_t(\mathbf{x}_t^*; \xi_{i,t})) \\ & \leq \frac{1}{n} \sum_{i=1}^n \beta \langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \mathbf{x}_{i,t} - \mathbf{x}_t^* \rangle + (1-\beta) \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{x}_{i,t} - \mathbf{x}_t^* \rangle \\ & = \frac{1}{n} \sum_{i=1}^n \beta (\langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \rangle + \langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \bar{\mathbf{x}}_{t+1} - \mathbf{x}_t^* \rangle) \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{n} \sum_{i=1}^n (1-\beta) \left( \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \rangle \right) \\
& + \frac{1}{n} \sum_{i=1}^n (1-\beta) \left( \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_{t+1} - \mathbf{x}_t^* \rangle \right) \\
& = \underbrace{\frac{1}{n} \sum_{i=1}^n \beta (\langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \rangle)}_{I_1(t)} \\
& + \underbrace{\frac{1}{n} \sum_{i=1}^n (1-\beta) \left( \langle \nabla h_t(\mathbf{x}_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \rangle \right)}_{I_2(t)} \\
& + \underbrace{\mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_{t+1} - \mathbf{x}_t^* \right\rangle}_{I_3(t)}
\end{aligned}$$

Now, we begin to bound  $I_1(t)$ .

$$\begin{aligned}
I_1(t) & \leq \frac{\beta}{n} \sum_{i=1}^n \left( \frac{\eta}{2} \|\nabla g_{i,t}(\mathbf{x}_{i,t})\|^2 + \frac{1}{2\eta} \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{\eta}{2} \|\nabla f_{i,t}(\mathbf{x}_{i,t})\|^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \right) \\
& \leq \beta G \eta + \frac{\beta}{2n\eta} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{\beta}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2.
\end{aligned}$$

Now, we begin to bound  $I_2(t)$ .

$$I_2(t) = (1-\beta) \left( \underbrace{\frac{1}{n} \sum_{i=1}^n \langle \nabla h_t(\mathbf{x}_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle}_{J_1(t)} + \underbrace{\mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \left\langle \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \right\rangle}_{J_2(t)} \right).$$

For  $J_1(t)$  and  $\nu > 0$ , we have

$$\begin{aligned}
J_1(t) & = \frac{1}{n} \sum_{i=1}^n \langle \nabla h_t(\mathbf{x}_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle \\
& = \frac{1}{n} \sum_{i=1}^n \langle \nabla h_t(\mathbf{x}_{i,t}) - \nabla h_t(\bar{\mathbf{x}}_t), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \frac{1}{n} \sum_{i=1}^n \langle \nabla h_t(\bar{\mathbf{x}}_t), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle \\
& \stackrel{\textcircled{1}}{\leq} \frac{L}{n} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{1}{n} \sum_{i=1}^n \langle \nabla h_t(\bar{\mathbf{x}}_t), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle \\
& \leq \frac{L}{n} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{1}{n} \sum_{i=1}^n \left( \frac{\eta}{2\nu} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 + \frac{\nu}{2\eta} \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \right) \\
& \leq \frac{L}{n} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{\eta}{2\nu} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 + \frac{\nu}{2\eta n} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2. \tag{1}
\end{aligned}$$

① holds due to  $h_t$  has  $L$ -Lipischitz gradients.

According to Lemma 2, we have

$$\begin{aligned}
& \frac{\eta}{2} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 \\
& \leq \frac{\eta}{2} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 + \frac{\eta}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 \\
& \leq \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} (h_t(\bar{\mathbf{x}}_t) - h_t(\bar{\mathbf{x}}_{t+1})) + 4G\eta\beta^2 + \frac{\eta L^2(1-\beta)^2}{n} \sum_{i=1}^n \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2. \quad (2)
\end{aligned}$$

Substituting (2) into (1), we obtain

$$\begin{aligned}
& J_1(t) \\
& \leq \frac{L}{n} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \left( \frac{1}{\nu} \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} (h_t(\bar{\mathbf{x}}_t) - h_t(\bar{\mathbf{x}}_{t+1})) + \frac{4G\eta\beta^2}{\nu} + \frac{\eta L^2(1-\beta)^2}{n\nu} \sum_{i=1}^n \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 \right) \\
& \quad + \frac{3GL\eta^2}{\nu} + \frac{2L\eta^2}{n\nu} \sum_{i=1}^n \sigma_{i,t}^2 + \frac{\nu}{2n\eta} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\
& = \left( \frac{L}{n} + \frac{\eta L^2(1-\beta)^2}{n\nu} + \frac{\nu}{2n\eta} \right) \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{1}{\nu} \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} (h_t(\bar{\mathbf{x}}_t) - h_t(\bar{\mathbf{x}}_{t+1})) \\
& \quad + \frac{G\eta(4\beta^2 + 3L\eta)}{\nu} + \frac{2L\eta^2}{n\nu} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

For  $J_2(t)$ , we have

$$\begin{aligned}
J_2(t) &= \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \left\langle \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \right\rangle \\
&\leq \left( \frac{\eta}{2} \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \left\| \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) \right\|^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \right) \\
&\leq \left( \frac{\eta}{2} \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \left\| \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - \nabla h_t(\mathbf{x}_{i,t}) + \nabla h_t(\mathbf{x}_{i,t}) \right\|^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \right) \\
&\leq \left( \eta \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \left\| \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - \nabla h_t(\mathbf{x}_{i,t}) \right\|^2 + \eta \left\| \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}) \right\|^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \right) \\
&\leq \frac{\eta}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + \eta \left\| \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}) - \nabla h_t(\bar{\mathbf{x}}_t) + \nabla h_t(\bar{\mathbf{x}}_t) \right\|^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\leq \frac{\eta}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + 2\eta \left\| \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}) - \nabla h_t(\bar{\mathbf{x}}_t) \right\|^2 + 2\eta \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\leq \frac{\eta}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + \frac{2\eta L^2}{n} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + 2\eta \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2.
\end{aligned}$$

Recall Lemma 2, and we have

$$J_2(t)$$

$$\begin{aligned}
&\leq \frac{\eta}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + \frac{2\eta L^2}{n} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + 4 \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} (h_t(\bar{\mathbf{x}}_t) - h_t(\bar{\mathbf{x}}_{t+1})) + 16G\eta\beta^2 + \frac{4\eta L^2(1-\beta)^2}{n} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\
&\quad + 12GL\eta^2 + \frac{8L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\leq \frac{\eta + 8nL\eta^2}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + (2\eta L^2 + 4\eta L^2(1-\beta)^2) \frac{1}{n} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + 4 \left( \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_t) - \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_{t+1}) \right) \\
&\quad + 16G\eta\beta^2 + 12GL\eta^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2.
\end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
&I_2(t) \\
&= (1-\beta)(J_1(t) + J_2(t)) \\
&\leq \left( \frac{L}{n} + \frac{\eta L^2(1-\beta)^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} \right) \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \left( \frac{1}{\nu} + 4 \right) \left( \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_t) - \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_{t+1}) \right) \\
&\quad + (1-\beta) \left( 4G\eta\beta^2 \left( \frac{1}{\nu} + 4 \right) + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \right) + \frac{(1-\beta)3LG\eta^2}{\nu} \left( \frac{1}{\nu} + 4 \right) + (1-\beta) \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{i=1}^n \sigma_{i,t}^2 \\
&\leq \left( \frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} \right) \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \left( \frac{1}{\nu} + 4 \right) \left( \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_t) - \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_{t+1}) \right) \\
&\quad + 4G\eta\beta^2 \left( \frac{1}{\nu} + 4 \right) + \frac{1-\beta}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 + 3LG\eta^2 \left( \frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

Combine those bounds of  $I_1(t)$  and  $I_2(t)$ . We thus have

$$\begin{aligned}
&I_1(t) + I_2(t) \\
&\leq \beta G\eta + \frac{\beta}{2n\eta} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{\beta}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\quad + \left( \frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} \right) \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \left( \frac{1}{\nu} + 4 \right) \left( \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_t) - \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_{t+1}) \right) \\
&\quad + 4G\eta\beta^2 \left( \frac{1}{\nu} + 4 \right) + \frac{1-\beta}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 + 3LG\eta^2 \left( \frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{i=1}^n \sigma_{i,t}^2 \\
&\leq \left( 1 + 4\beta \left( \frac{1}{\nu} + 4 \right) \right) \beta G\eta + \left( \frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} + \frac{\beta}{2n\eta} \right) \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\
&\quad + \left( \frac{1}{\nu} + 4 \right) \left( \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_t) - \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_{t+1}) \right) + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\quad + 3LG\eta^2 \left( \frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

Therefore, we have

$$\sum_{t=1}^T (I_1(t) + I_2(t))$$

$$\begin{aligned}
&\leq \left(1 + 4\beta \left(\frac{1}{\nu} + 4\right)\right) T\beta G\eta + \left(\frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} + \frac{\beta}{2n\eta}\right) \sum_{t=1}^T \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\
&\quad + \left(\frac{1}{\nu} + 4\right) (h_1(\bar{\mathbf{x}}_1) - h_{T+1}(\bar{\mathbf{x}}_{T+1})) + \frac{1}{2\eta} \sum_{t=1}^T \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\quad + 3TLG\eta^2 \left(\frac{1}{\nu} + 4\right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

Now, we begin to bound  $I_3(t)$ . Recall that the update rule is

$$\mathbf{x}_{i,t+1} = \sum_{j=1}^n \mathbf{W}_{ij} \mathbf{x}_{j,t} - \eta \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}).$$

By taking average over  $i \in [n]$  on both sides, we have

$$\bar{\mathbf{x}}_{t+1} = \bar{\mathbf{x}}_t - \eta \left( \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right). \quad (3)$$

Denote a new auxiliary function  $h(\mathbf{z})$  as

$$h(\mathbf{z}) = \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{z} \right\rangle + \frac{1}{2\eta} \|\mathbf{z} - \bar{\mathbf{x}}_t\|^2.$$

Note that (3) is equivalent to

$$\begin{aligned}
\bar{\mathbf{x}}_{t+1} &= \underset{\mathbf{z} \in \mathbb{R}^d}{\operatorname{argmin}} h(\mathbf{z}) \\
&= \underset{\mathbf{z} \in \mathbb{R}^d}{\operatorname{argmin}} \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{z} \right\rangle + \frac{1}{2\eta} \|\mathbf{z} - \bar{\mathbf{x}}_t\|^2.
\end{aligned}$$

Furthermore, denote a new auxiliary variable  $\bar{\mathbf{x}}_\tau$  as

$$\bar{\mathbf{x}}_\tau = \bar{\mathbf{x}}_{t+1} + \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}),$$

where  $0 \leq \tau \leq 1$ . According to the optimality of  $\bar{\mathbf{x}}_{t+1}$ , we have

$$\begin{aligned}
0 &\leq h(\bar{\mathbf{x}}_\tau) - h(\bar{\mathbf{x}}_{t+1}) \\
&= \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_\tau - \bar{\mathbf{x}}_{t+1} \right\rangle + \frac{1}{2\eta} \left( \|\bar{\mathbf{x}}_\tau - \bar{\mathbf{x}}_t\|^2 - \|\bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t\|^2 \right) \\
&= \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}) \right\rangle + \frac{1}{2\eta} \left( \|\bar{\mathbf{x}}_{t+1} + \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}) - \bar{\mathbf{x}}_t\|^2 - \|\bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t\|^2 \right) \\
&= \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}) \right\rangle + \frac{1}{2\eta} \left( \|\tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1})\|^2 + 2 \langle \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}), \bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t \rangle \right).
\end{aligned}$$

Dividing  $\tau$  on both sides, and letting  $\tau$  be close to 0, we have

$$I_3(t) = \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_{t+1} - \mathbf{x}_t^* \right\rangle$$

$$\begin{aligned}
&\leq \frac{1}{2\eta} \left( \lim_{\tau \rightarrow 0} \tau \|(\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1})\|^2 + 2 \langle \mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}, \bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t \rangle \right) \\
&= \frac{1}{\eta} \langle \mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}, \bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t \rangle \\
&= \frac{1}{2\eta} \left( \|\mathbf{x}_t^* - \bar{\mathbf{x}}_t\|^2 - \|\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}\|^2 - \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \right). \tag{4}
\end{aligned}$$

Besides, we have

$$\begin{aligned}
&\|\mathbf{x}_{t+1}^* - \bar{\mathbf{x}}_{t+1}\|^2 - \|\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}\|^2 \\
&= \|\mathbf{x}_{t+1}^*\|^2 - \|\mathbf{x}_t^*\|^2 - 2 \langle \bar{\mathbf{x}}_{t+1}, -\mathbf{x}_t^* + \mathbf{x}_{t+1}^* \rangle \\
&= (\|\mathbf{x}_{t+1}^*\| - \|\mathbf{x}_t^*\|) (\|\mathbf{x}_{t+1}^*\| + \|\mathbf{x}_t^*\|) - 2 \langle \bar{\mathbf{x}}_{t+1}, -\mathbf{x}_t^* + \mathbf{x}_{t+1}^* \rangle \\
&\leq \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\| (\|\mathbf{x}_{t+1}^*\| + \|\mathbf{x}_t^*\|) - 2 \|\bar{\mathbf{x}}_{t+1}\| \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\| \\
&\leq 4\sqrt{R} \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\|. \quad (\text{due to } \|\mathbf{x} - \mathbf{y}\|^2 \leq R, \forall \mathbf{x}, \mathbf{y} \in \mathcal{X})
\end{aligned}$$

Thus, telescoping  $I_3(t)$  over  $t \in [T]$ , we have

$$\begin{aligned}
\sum_{t=1}^T I_3(t) &\leq \frac{1}{2\eta} \left( 4\sqrt{R} \sum_{t=1}^T \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\| + \|\bar{\mathbf{x}}_1^* - \bar{\mathbf{x}}_1\|^2 - \|\bar{\mathbf{x}}_T^* - \bar{\mathbf{x}}_{T+1}\|^2 \right) - \frac{1}{2\eta} \sum_{t=1}^T \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\leq \frac{1}{2\eta} \left( 4\sqrt{R} \sum_{t=1}^T D + R \right) - \frac{1}{2\eta} \sum_{t=1}^T \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2.
\end{aligned}$$

Combining those bounds of  $I_1(t)$ ,  $I_2(t)$  and  $I_3(t)$  together, we finally obtain

$$\begin{aligned}
&\mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1, t=1}^{n, T}} \sum_{t=1}^T \sum_{i=1}^n f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - f_t(\mathbf{x}_t^*; \xi_{i,t}) \\
&\leq n \sum_{t=1}^T (I_1(t) + I_2(t) + I_3(t)) \\
&\leq \left( 1 + 4\beta \left( \frac{1}{\nu} + 4 \right) \right) nT\beta G\eta + \left( L + \frac{\eta L^2}{\nu} + \frac{\nu}{2\eta} + 6\eta L^2 + \frac{\beta}{2\eta} \right) \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1, t=1}^{n, T}} \sum_{t=1}^T \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\
&\quad + \left( \frac{1}{\nu} + 4 \right) n (h_t(\bar{\mathbf{x}}_1) - h_t(\bar{\mathbf{x}}_{T+1})) \\
&\quad + 3nTLG\eta^2 \left( \frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n\nu} \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{R}D + R) \\
&\stackrel{\textcircled{1}}{\leq} \left( 1 + 4\beta \left( \frac{1}{\nu} + 4 \right) \right) nT\beta G\eta + \left( L + \frac{\eta L^2}{\nu} + \frac{\nu}{2\eta} + 6\eta L^2 + \frac{\beta}{2\eta} \right) \frac{\eta^2}{(1-\rho)^2} \left( 6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right) \\
&\quad + \left( \frac{1}{\nu} + 4 \right) n (h_t(\bar{\mathbf{x}}_1) - h_t(\bar{\mathbf{x}}_{T+1})) \\
&\quad + 3nTLG\eta^2 \left( \frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n\nu} \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{R}D + R).
\end{aligned}$$

① holds due to Lemma 3, that is,

$$\mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1, t=1}^{n, T}} \sum_{i=1}^n \sum_{t=1}^T \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \leq \frac{\eta^2}{(1-\rho)^2} \left( 6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right).$$

When  $\nu = \beta$ , we have

$$\begin{aligned}
& \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \sum_{t=1}^T \sum_{i=1}^n f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - f_t(\mathbf{x}_t^*; \xi_{i,t}) \\
& \leq 21nT\beta G\eta + \left( L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) \frac{\eta^2}{(1-\rho)^2} \left( 6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right) \\
& \quad + \left( \frac{1}{\beta} + 4 \right) n (h_t(\bar{\mathbf{x}}_1) - h_t(\bar{\mathbf{x}}_{T+1})) + 3nTLG\eta^2 \left( \frac{1}{\beta} + 4 \right) \\
& \quad + \frac{\beta(\eta + 8nL\eta^2) + 2nL\eta^2}{n\beta} \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{RD} + R) \\
& = \eta nTG \left( 21\beta + \frac{6\eta}{(1-\rho)^2} \left( L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) + 3L\eta \left( \frac{1}{\beta} + 4 \right) \right) + \left( \frac{1}{\beta} + 4 \right) n (h_t(\bar{\mathbf{x}}_1) - h_t(\bar{\mathbf{x}}_{T+1})) \\
& \quad + \left( \left( L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) \frac{4\eta^2}{(1-\rho)^2} + \frac{\beta(\eta + 8nL\eta^2) + 2nL\eta^2}{n\beta} \right) \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{RD} + R).
\end{aligned}$$

It completes the proof.  $\square$

**Lemma 1.** For any  $\beta$  with  $0 \leq \beta \leq 1$ , we have

$$\mathbb{E}_{\xi_{i,t} \sim D_{i,t}} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \leq 6G + 4\sigma_{i,t}^2.$$

*Proof.*

$$\begin{aligned}
& \mathbb{E}_{\xi_{i,t} \sim D_{i,t}} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\
& = \mathbb{E}_{\xi_{i,t} \sim D_{i,t}} \|\beta \nabla g_{i,t}(\mathbf{x}_{i,t}) + (1-\beta) \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\
& \leq 2\beta^2 \|\nabla g_{i,t}(\mathbf{x}_{i,t})\|^2 + 2(1-\beta)^2 \mathbb{E}_{\xi_{i,t} \sim D_{i,t}} \|\nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\
& \leq 2G\beta^2 + 2(1-\beta)^2 \mathbb{E}_{\xi_{i,t} \sim D_{i,t}} \|\nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - \nabla h_t(\mathbf{x}_{i,t}) + \nabla h_t(\mathbf{x}_{i,t})\|^2 \\
& \leq 2G\beta^2 + 4(1-\beta)^2 \mathbb{E}_{\xi_{i,t} \sim D_{i,t}} \|\nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - \nabla h_t(\mathbf{x}_{i,t})\|^2 + 4(1-\beta)^2 \|\nabla h_t(\mathbf{x}_{i,t})\|^2 \\
& \leq 2G\beta^2 + 4(1-\beta)^2 \sigma_{i,t}^2 + 4(1-\beta)^2 G \\
& \leq 6G + 4\sigma_{i,t}^2.
\end{aligned}$$

$\square$

**Lemma 2.**

$$\begin{aligned}
& \frac{\eta}{2} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 + \frac{\eta}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 \\
& \leq \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} (h_t(\bar{\mathbf{x}}_t) - h_t(\bar{\mathbf{x}}_{t+1})) + 4G\eta\beta^2 + \frac{\eta L^2(1-\beta)^2}{n} \sum_{i=1}^n \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$



*Proof.*

$$\begin{aligned}
& \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} h_t(\bar{\mathbf{x}}_{t+1}) \\
& \leq h_t(\bar{\mathbf{x}}_t) + \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \langle \nabla h_t(\bar{\mathbf{x}}_t), \bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t \rangle + \frac{L}{2} \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \|\bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t\|^2 \\
& = h_t(\bar{\mathbf{x}}_t) + \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \left\langle \nabla h_t(\bar{\mathbf{x}}_t), -\frac{\eta}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right\rangle + \frac{L}{2} \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \left\| \frac{\eta}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right\|^2 \\
& = h_t(\bar{\mathbf{x}}_t) + \left\langle \nabla h_t(\bar{\mathbf{x}}_t), -\frac{\eta}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\rangle + \frac{L\eta^2}{2n} \sum_{i=1}^n \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\
& = h_t(\bar{\mathbf{x}}_t) + \frac{\eta}{2} \left( \left\| \nabla h_t(\bar{\mathbf{x}}_t) - \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 - \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 - \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 \right) \\
& \quad + \frac{L\eta^2}{2n} \sum_{i=1}^n \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\
& \stackrel{\textcircled{1}}{\leq} h_t(\bar{\mathbf{x}}_t) + \frac{\eta}{2} \left\| \nabla h_t(\bar{\mathbf{x}}_t) - \frac{1}{n} \sum_{i=1}^n (\beta \nabla g_{i,t}(\mathbf{x}_{i,t}) + (1-\beta) \nabla h_t(\mathbf{x}_{i,t})) \right\|^2 - \frac{\eta}{2} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 - \frac{\eta}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 \\
& \quad + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2 \\
& \leq h_t(\bar{\mathbf{x}}_t) + \frac{\eta}{2} \left( 2\beta^2 \left\| \nabla h_t(\bar{\mathbf{x}}_t) - \frac{1}{n} \sum_{i=1}^n \nabla g_{i,t}(\mathbf{x}_{i,t}) \right\|^2 + 2(1-\beta)^2 \left\| \nabla h_t(\bar{\mathbf{x}}_t) - \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}) \right\|^2 \right) \\
& \quad - \frac{\eta}{2} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 - \frac{\eta}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2 \\
& \leq h_t(\bar{\mathbf{x}}_t) + \frac{\eta}{2} \left( 4\beta^2 G + 2(1-\beta)^2 L^2 \frac{1}{n} \sum_{i=1}^n \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 \right) - \frac{\eta}{2} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 - \frac{\eta}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2 \\
& \leq h_t(\bar{\mathbf{x}}_t) + 2\eta\beta^2 G + \eta(1-\beta)^2 L^2 \left( \frac{1}{n} \sum_{i=1}^n \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 \right) - \frac{\eta}{2} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 - \frac{\eta}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

① holds due to Lemma 1. Equivalently, we obtain

$$\begin{aligned}
& \frac{\eta}{2} \|\nabla h_t(\bar{\mathbf{x}}_t)\|^2 + \frac{\eta}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 \\
& \leq \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1}^n} (h_t(\bar{\mathbf{x}}_t) - h_t(\bar{\mathbf{x}}_{t+1})) + 2G\eta\beta^2 + \frac{\eta L^2(1-\beta)^2}{n} \sum_{i=1}^n \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

It completes the proof.  $\square$

**Lemma 3.**

$$\mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1, t=1}^{n,T}} \sum_{i=1}^n \sum_{t=1}^T \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \leq \frac{\eta^2}{(1-\rho)^2} \left( 6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right).$$

*Proof.* Recall that

$$\mathbf{x}_{i,t+1} = \sum_{j=1}^n \mathbf{W}_{ij} \mathbf{x}_{j,t} - \eta \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}),$$

and

$$\bar{\mathbf{x}}_{t+1} = \bar{\mathbf{x}}_t - \eta \left( \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right).$$

Denote

$$\begin{aligned} \mathbf{X}_t &= [\mathbf{x}_{1,t}, \mathbf{x}_{2,t}, \dots, \mathbf{x}_{n,t}] \in \mathbb{R}^{d \times n}, \\ \mathbf{G}_t &= [\nabla f_{1,t}(\mathbf{x}_{1,t}; \xi), \nabla f_{2,t}(\mathbf{x}_{2,t}; \xi), \dots, \nabla f_{n,t}(\mathbf{x}_{n,t}; \xi)] \in \mathbb{R}^{d \times n}. \end{aligned}$$

By letting  $\mathbf{x}_{i,1} = \mathbf{0}$  for any  $i \in [n]$ , the update rule is re-formulated as

$$\mathbf{X}_{t+1} = \mathbf{X}_t \mathbf{W} - \eta \mathbf{G}_t = - \sum_{s=1}^t \eta \mathbf{G}_s \mathbf{W}^{t-s}.$$

Similarly, denote  $\bar{\mathbf{G}}_t = \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})$ , and we have

$$\bar{\mathbf{x}}_{t+1} = \bar{\mathbf{x}}_t - \eta \left( \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right) = - \sum_{s=1}^t \eta \bar{\mathbf{G}}_s. \quad (5)$$

Therefore,

$$\begin{aligned} & \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\ & \stackrel{\textcircled{1}}{=} \sum_{i=1}^n \left\| \sum_{s=1}^{t-1} \eta \bar{\mathbf{G}}_s - \eta \mathbf{G}_s \mathbf{W}^{t-s-1} \mathbf{e}_i \right\|^2 \\ & \stackrel{\textcircled{2}}{=} \left\| \sum_{s=1}^{t-1} \eta \mathbf{G}_s \mathbf{v}_1 \mathbf{v}_1^T - \eta \mathbf{G}_s \mathbf{W}^{t-s-1} \right\|_F^2 \\ & \stackrel{\textcircled{3}}{\leq} \left( \eta \rho^{t-s-1} \left\| \sum_{s=1}^{t-1} \mathbf{G}_s \right\|_F \right)^2 \\ & \leq \left( \sum_{s=1}^{t-1} \eta \rho^{t-s-1} \|\mathbf{G}_s\|_F \right)^2. \end{aligned}$$

① holds due to  $\mathbf{e}_i$  is a unit basis vector, whose  $i$ -th element is 1 and other elements are 0s. ② holds due to  $\mathbf{v}_1 = \frac{1}{\sqrt{n}}$ . ③ holds due to Lemma 4.

According to Lemma 5, we have

$$\begin{aligned} & \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1,t=1}^{n,T}} \sum_{i=1}^n \sum_{t=1}^T \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\ & \leq \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1,t=1}^{n,T}} \sum_{t=1}^T \left( \sum_{s=1}^{t-1} \eta \rho^{t-s-1} \|\mathbf{G}_s\|_F \right)^2 \end{aligned}$$

$$\begin{aligned}
&\leq \frac{\eta^2}{(1-\rho)^2} \mathbb{E}_{\{\xi_{i,t} \sim D_{i,t}\}_{i=1, t=1}^{n,T}} \left( \sum_{t=1}^T \|\mathbf{G}_t\|_F^2 \right) \\
&\leq \frac{\eta^2}{(1-\rho)^2} \left( \sum_{t=1}^T \sum_{i=1}^n \mathbb{E}_{\xi_{i,t} \sim D_{i,t}} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \right) \\
&\leq \frac{\eta^2}{(1-\rho)^2} \left( 6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right) \\
&\leq \frac{\eta^2}{(1-\rho)^2} \left( 6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right).
\end{aligned}$$

It completes the proof.  $\square$

**Lemma 4** (Appeared in Lemma 5 in [Tang et al., 2018]). *For any matrix  $\mathbf{X}_t \in \mathbb{R}^{d \times n}$ , decompose the confusion matrix  $\mathbf{W}$  as  $\mathbf{W} = \sum_{i=1}^n \lambda_i \mathbf{v}_i \mathbf{v}_i^T = \mathbf{P} \mathbf{\Lambda} \mathbf{P}^T$ , where  $\mathbf{P} = [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n] \in \mathbb{R}^{n \times n}$ ,  $\mathbf{v}_i$  is the normalized eigenvector of  $\lambda_i$ .  $\mathbf{\Lambda}$  is a diagonal matrix, and  $\lambda_i$  be its  $i$ -th element. We have*

$$\|\mathbf{X}_t \mathbf{W}^t - \mathbf{X}_t \mathbf{v}_1 \mathbf{v}_1^T\|_F^2 \leq \|\rho^t \mathbf{X}_t\|_F^2,$$

where  $\rho = \max\{|\lambda_2(\mathbf{W})|, |\lambda_n(\mathbf{W})|\}$ .

**Lemma 5** (Appeared in Lemma 6 in [Tang et al., 2018]). *Given two non-negative sequences  $\{a_t\}_{t=1}^\infty$  and  $\{b_t\}_{t=1}^\infty$  that satisfying*

$$a_t = \sum_{s=1}^t \rho^{t-s} b_s,$$

with  $\rho \in [0, 1)$ , we have

$$\sum_{t=1}^k a_t^2 \leq \frac{1}{(1-\rho)^2} \sum_{s=1}^k b_s^2.$$

## References

H. Tang, S. Gan, C. Zhang, T. Zhang, and J. Liu. Communication Compression for Decentralized Training. *arXiv.org*, Mar. 2018.