

Decentralized Online Optimization

December 26, 2018

Abstract

ddd

1 Notations and assumptions

For any $i \in [n]$ and $t \in [T]$, the random variable $\xi_{i,t}$ is subject to a distribution $D_{i,t}$, that is,

$$\xi_{i,t} \sim D_{i,t}.$$

Besides, a set of random variables $\Xi_{n,T}$ and the corresponding set of distributions are defined by

$$\Xi_{n,T} = \{\xi_{i,t}\}_{1 \leq i \leq n, 1 \leq t \leq T}, \text{ and } \mathcal{D}_{n,T} = \{D_{i,t}\}_{1 \leq i \leq n, 1 \leq t \leq T},$$

respectively. For math brevity, we use the notation $\Xi_{n,T} \sim \mathcal{D}_{n,T}$ to represent that $\xi_{i,t} \sim D_{i,t}$ holds for any $i \in [n]$ and $t \in [T]$.

For any online algorithm $A \in \mathcal{A}$, define its dynamic regret as

$$\mathcal{R}_T^A = \mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \left(\sum_{i=1}^n \sum_{t=1}^T f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - f_{i,t}(\mathbf{x}_t^*; \xi_{i,t}) \right),$$

where, for any \mathbf{x} ,

$$f_{i,t}(\mathbf{x}; \xi_{i,t}) := \beta g_{i,t}(\mathbf{x}) + (1 - \beta) h_{i,t}(\mathbf{x}; \xi_{i,t})$$

with $0 < \beta < 1$, and $\xi_{i,t}$ is a random variable drawn from an unknown distribution $D_{i,t}$. $g_{i,t}$ is an adversary loss function. $h_{i,t}(\cdot, \xi_{i,t})$ is a given loss function depending on the random variable $\xi_{i,t}$. Besides, we denote

$$H_{i,t}(\cdot) = \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} h_{i,t}(\cdot; \xi_{i,t}).$$

The budget of the dynamics is defined as

$$\sum_{t=1}^T \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\| \leq M.$$

Assumption 1. *We make the following assumptions.*

- For any $i \in [n]$ and $t \in [T]$, there exists a constant G such that

$$\max \left\{ \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla h_t(\mathbf{x}; \xi_{i,t})\|^2, \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla f_{i,t}(\mathbf{x}; \xi_{i,t})\|^2 \right\} \leq G,$$

and

$$\mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla h_t(\mathbf{x}; \xi_{i,t}) - \nabla H_t(\mathbf{x})\|^2 \leq \sigma_{i,t}^2.$$

- For any \mathbf{x} and \mathbf{y} , we assume $\|\mathbf{x} - \mathbf{y}\|^2 \leq R$.
- For any $i \in [n]$ and $t \in [T]$, we assume the function $f_{i,t}$ is convex and differentiable, and the function H_t has L -Lipschitz gradients.

2 Algorithm

Algorithm 1 DOG: Decentralized Online Gradient.

Require: The learning rate η , number of iterations T , and the confusion matrix \mathbf{W} .

```

1: for  $t = 1, 2, \dots, T$  do
    For the  $i$ -th node with  $i \in [n]$ :
2:     Predict  $\mathbf{x}_{i,t}$ .
3:     Observe the loss function  $f_{i,t}$ ,
        and suffer loss  $f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})$ .
    Update:
4:     Query the gradient  $\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})$ .
5:      $\mathbf{x}_{i,t+1} = \sum_{j=1}^n \mathbf{W}_{i,j} \mathbf{x}_{j,t} - \eta \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})$ .
```

Theorem 1. For any $0 \leq \beta \leq 1$ and $\eta > 0$, we denote

$$C_1 = 11\beta + \frac{6\eta}{(1-\rho)^2} \left(L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) + 3L\eta \left(\frac{1}{\beta} + 4 \right),$$

$$C_2 = \left(L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) \frac{4\eta^2}{(1-\rho)^2} + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n\beta}.$$

Use Assumption 1, and choose $\eta > 0$ in Algorithm 1. We have

$$\mathcal{R}_T^{DOG} \leq \eta n T G C_1 + \left(\frac{1}{\beta} + 4 \right) n (H_t(\bar{\mathbf{x}}_1) - H_t(\bar{\mathbf{x}}_{t+1})) + C_2 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{RM} + R).$$

Corollary 1. Using Assumption 1, and choosing

$$\eta = \sqrt{\frac{nM}{T(1+n\beta)}}$$

in Algorithm 1, we have

$$\mathcal{R}_T^{DOG} \lesssim \mathcal{O} \left(\sqrt{(1+n\beta)nMT} \right).$$

Appendix

Proof to Theorem 1:

Proof.

$$\begin{aligned}
& \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \frac{1}{n} \sum_{i=1}^n f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - f_t(\mathbf{x}_t^*; \xi_{i,t}) \\
&= \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \frac{1}{n} \sum_{i=1}^n \beta (g_{i,t}(\mathbf{x}_{i,t}) - g_{i,t}(\mathbf{x}_t^*)) + (1-\beta) \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - h_t(\mathbf{x}_t^*; \xi_{i,t}))
\end{aligned}$$

$$\begin{aligned}
&\leq \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \frac{1}{n} \sum_{i=1}^n \beta \langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \mathbf{x}_{i,t} - \mathbf{x}_t^* \rangle + (1-\beta) \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{x}_{i,t} - \mathbf{x}_t^* \rangle \\
&= \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \frac{1}{n} \sum_{i=1}^n \beta (\langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \rangle + \langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \bar{\mathbf{x}}_{t+1} - \mathbf{x}_t^* \rangle) \\
&\quad + \frac{1}{n} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \sum_{i=1}^n (1-\beta) (\langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \rangle) \\
&\quad + \frac{1}{n} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \sum_{i=1}^n (1-\beta) (\langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_{t+1} - \mathbf{x}_t^* \rangle) \\
&= \underbrace{\mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \frac{1}{n} \sum_{i=1}^n \beta (\langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \langle \nabla g_{i,t}(\mathbf{x}_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \rangle)}_{I_1(t)} \\
&\quad + \underbrace{\mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \frac{1}{n} \sum_{i=1}^n (1-\beta) (\langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \rangle)}_{I_2(t)} \\
&\quad + \underbrace{\mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_{t+1} - \mathbf{x}_t^* \right\rangle}_{I_3(t)}
\end{aligned}$$

Now, we begin to bound $I_1(t)$.

$$\begin{aligned}
I_1(t) &\leq \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \frac{\beta}{n} \sum_{i=1}^n \left(\frac{\eta}{2} \|\nabla g_{i,t}(\mathbf{x}_{i,t})\|^2 + \frac{1}{2\eta} \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{\eta}{2} \|\nabla f_{i,t}(\mathbf{x}_{i,t})\|^2 + \frac{1}{2\eta} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \right) \\
&\leq \beta G\eta + \frac{\beta}{2n\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{\beta}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2.
\end{aligned}$$

Now, we begin to bound $I_2(t)$.

$$I_2(t) = (1-\beta) \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \left(\underbrace{\frac{1}{n} \sum_{i=1}^n \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle}_{J_1(t)} + \underbrace{\left\langle \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \right\rangle}_{J_2(t)} \right).$$

For $J_1(t)$ and $\nu > 0$, we have

$$\begin{aligned}
&J_1(t) \\
&= \frac{1}{n} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \sum_{i=1}^n \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle \\
&= \frac{1}{n} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \sum_{i=1}^n \langle \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - \nabla H_t(\bar{\mathbf{x}}_t), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \frac{1}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \langle \nabla H_t(\bar{\mathbf{x}}_t), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle \\
&= \frac{1}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \langle \nabla H_t(\mathbf{x}_{i,t}) - \nabla H_t(\bar{\mathbf{x}}_t), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle + \frac{1}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \langle \nabla H_t(\bar{\mathbf{x}}_t), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle
\end{aligned}$$

$$\begin{aligned}
& \stackrel{\textcircled{1}}{\leq} \frac{L}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{1}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \langle \nabla H_t(\bar{\mathbf{x}}_t), \mathbf{x}_{i,t} - \bar{\mathbf{x}}_t \rangle \\
& \leq \frac{L}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{1}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \left(\frac{\eta}{2\nu} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 + \frac{\nu}{2\eta} \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \right) \\
& \leq \frac{L}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{\eta}{2\nu} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 + \frac{\nu}{2\eta n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2. \quad (1)
\end{aligned}$$

① holds due to H_t has L -Lipischitz gradients.

According to Lemma 2, we have

$$\begin{aligned}
& \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 \\
& \leq \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) + 2G\eta\beta^2 + \frac{\eta L^2(1-\beta)^2}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2. \quad (2)
\end{aligned}$$

Substituting (2) into (1), we obtain

$$\begin{aligned}
& J_1(t) \\
& \leq \frac{L}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{1}{\nu} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) \\
& \quad + \frac{2G\eta\beta^2}{\nu} + \frac{\eta L^2(1-\beta)^2}{n\nu} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 \\
& \quad + \frac{3GL\eta^2}{\nu} + \frac{2L\eta^2}{n\nu} \sum_{i=1}^n \sigma_{i,t}^2 + \frac{\nu}{2n\eta} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\
& = \left(\frac{L}{n} + \frac{\eta L^2(1-\beta)^2}{n\nu} + \frac{\nu}{2n\eta} \right) \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{1}{\nu} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) \\
& \quad + \frac{G\eta(2\beta^2 + 3L\eta)}{\nu} + \frac{2L\eta^2}{n\nu} \sum_{i=1}^n \sigma_{i,t}^2 \\
& \leq \left(\frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} \right) \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{1}{\nu} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) \\
& \quad + \frac{G\eta(2\beta^2 + 3L\eta)}{\nu} + \frac{2L\eta^2}{n\nu} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

For $J_2(t)$, we have

$$\begin{aligned}
& J_2(t) \\
& = \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \left\langle \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1} \right\rangle \\
& \leq \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) \right\|^2 + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
& \leq \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - \nabla H_t(\bar{\mathbf{x}}_t) + \nabla H_t(\bar{\mathbf{x}}_t) \right\|^2 + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2
\end{aligned}$$

$$\begin{aligned}
&\leq \eta \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - \nabla H_t(\mathbf{x}_{i,t}) \right\|^2 + \eta \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla H_t(\mathbf{x}_{i,t}) \right\|^2 \\
&\quad + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\leq \frac{\eta}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + \eta \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla H_t(\mathbf{x}_{i,t}) - \nabla H_t(\bar{\mathbf{x}}_t) + \nabla H_t(\bar{\mathbf{x}}_t) \right\|^2 + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\leq \frac{\eta}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + 2\eta \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla H_t(\mathbf{x}_{i,t}) - \nabla H_t(\bar{\mathbf{x}}_t) \right\|^2 \\
&\quad + 2\eta \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\leq \frac{\eta}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + \frac{2\eta L^2}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\
&\quad + 2\eta \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2. \tag{3}
\end{aligned}$$

The last inequality holds due to H_t has L Lipschitz gradients.

Substituting (2) into (3), and we have

$$\begin{aligned}
&J_2(t) \\
&\leq \frac{\eta}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + \frac{2\eta L^2}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + 4 \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) + 8G\eta\beta^2 \\
&\quad + \frac{4\eta L^2(1-\beta)^2}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + 12GL\eta^2 + \frac{8L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2 + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\leq \frac{\eta + 8nL\eta^2}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + (2\eta L^2 + 4\eta L^2(1-\beta)^2) \frac{1}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + 4 \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) \\
&\quad + 8G\eta\beta^2 + 12GL\eta^2 + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\leq \frac{\eta + 8nL\eta^2}{n^2} \sum_{i=1}^n \sigma_{i,t}^2 + \frac{6\eta L^2}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + 4 \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) \\
&\quad + 8G\eta\beta^2 + 12GL\eta^2 + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2.
\end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
&I_2(t) \\
&= (1-\beta)(J_1(t) + J_2(t)) \\
&= (1-\beta) \left(\frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} \right) \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + (1-\beta) \left(\frac{1}{\nu} + 4 \right) \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) \\
&\quad + (1-\beta) \left(2G\eta\beta^2 \left(\frac{1}{\nu} + 4 \right) + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \right) \\
&\quad + \frac{(1-\beta)3LG\eta^2}{\nu} \left(\frac{1}{\nu} + 4 \right) + (1-\beta) \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{i=1}^n \sigma_{i,t}^2
\end{aligned}$$

$$\begin{aligned}
&\leq \left(\frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} \right) \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \left(\frac{1}{\nu} + 4 \right) \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) \\
&\quad + 2G\eta\beta^2 \left(\frac{1}{\nu} + 4 \right) + \frac{1-\beta}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 + 3LG\eta^2 \left(\frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

Combine those bounds of $I_1(t)$ and $I_2(t)$. We thus have

$$\begin{aligned}
&I_1(t) + I_2(t) \\
&\leq \beta G\eta + \frac{\beta}{2n\eta} \sum_{i=1}^n \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \frac{\beta}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\quad + \left(\frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} \right) \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 + \left(\frac{1}{\nu} + 4 \right) \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) \\
&\quad + 2G\eta\beta^2 \left(\frac{1}{\nu} + 4 \right) + \frac{1-\beta}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 + 3LG\eta^2 \left(\frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{i=1}^n \sigma_{i,t}^2 \\
&\leq \left(1 + 2\beta \left(\frac{1}{\nu} + 4 \right) \right) \beta G\eta + \left(\frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} + \frac{\beta}{2n\eta} \right) \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\
&\quad + \left(\frac{1}{\nu} + 4 \right) \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\quad + 3LG\eta^2 \left(\frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
&\sum_{t=1}^T (I_1(t) + I_2(t)) \\
&\leq \left(1 + 2\beta \left(\frac{1}{\nu} + 4 \right) \right) T\beta G\eta + \left(\frac{L}{n} + \frac{\eta L^2}{n\nu} + \frac{\nu}{2n\eta} + \frac{6\eta L^2}{n} + \frac{\beta}{2n\eta} \right) \mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{t=1}^T \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\
&\quad + \left(\frac{1}{\nu} + 4 \right) (H_1(\bar{\mathbf{x}}_1) - H_{T+1}(\bar{\mathbf{x}}_{T+1})) + \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{t=1}^T \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \\
&\quad + 3TLG\eta^2 \left(\frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n^2\nu} \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

Now, we begin to bound $I_3(t)$. Recall that the update rule is

$$\mathbf{x}_{i,t+1} = \sum_{j=1}^n \mathbf{W}_{ij} \mathbf{x}_{j,t} - \eta \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}).$$

By taking average over $i \in [n]$ on both sides, we have

$$\bar{\mathbf{x}}_{t+1} = \bar{\mathbf{x}}_t - \eta \left(\frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right). \quad (4)$$

Denote a new auxiliary function $h(\mathbf{z})$ as

$$\phi(\mathbf{z}) = \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{z} \right\rangle + \frac{1}{2\eta} \|\mathbf{z} - \bar{\mathbf{x}}_t\|^2.$$

Note that (4) is equivalent to

$$\begin{aligned}\bar{\mathbf{x}}_{t+1} &= \underset{\mathbf{z} \in \mathbb{R}^d}{\operatorname{argmin}} \phi(\mathbf{z}) \\ &= \underset{\mathbf{z} \in \mathbb{R}^d}{\operatorname{argmin}} \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \mathbf{z} \right\rangle + \frac{1}{2\eta} \|\mathbf{z} - \bar{\mathbf{x}}_t\|^2.\end{aligned}$$

Furthermore, denote a new auxiliary variable $\bar{\mathbf{x}}_\tau$ as

$$\bar{\mathbf{x}}_\tau = \bar{\mathbf{x}}_{t+1} + \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}),$$

where $0 \leq \tau \leq 1$. According to the optimality of $\bar{\mathbf{x}}_{t+1}$, we have

$$\begin{aligned}0 &\leq \phi(\bar{\mathbf{x}}_\tau) - \phi(\bar{\mathbf{x}}_{t+1}) \\ &= \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_\tau - \bar{\mathbf{x}}_{t+1} \right\rangle + \frac{1}{2\eta} (\|\bar{\mathbf{x}}_\tau - \bar{\mathbf{x}}_t\|^2 - \|\bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t\|^2) \\ &= \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}) \right\rangle + \frac{1}{2\eta} (\|\bar{\mathbf{x}}_{t+1} + \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}) - \bar{\mathbf{x}}_t\|^2 - \|\bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t\|^2) \\ &= \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}) \right\rangle + \frac{1}{2\eta} (\|\tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1})\|^2 + 2 \langle \tau (\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}), \bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t \rangle).\end{aligned}$$

Dividing τ on both sides, and letting τ be close to 0, we have

$$\begin{aligned}I_3(t) &= \underset{\Xi_{n,t} \sim \mathcal{D}_{n,t}}{\mathbb{E}} \left\langle \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}), \bar{\mathbf{x}}_{t+1} - \mathbf{x}_t^* \right\rangle \\ &\leq \frac{1}{2\eta} \underset{\Xi_{n,t} \sim \mathcal{D}_{n,t}}{\mathbb{E}} \left(\lim_{\tau \rightarrow 0} \tau \|\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}\|^2 + 2 \langle \mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}, \bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t \rangle \right) \\ &= \frac{1}{\eta} \underset{\Xi_{n,t} \sim \mathcal{D}_{n,t}}{\mathbb{E}} \langle \mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}, \bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t \rangle \\ &= \frac{1}{2\eta} \underset{\Xi_{n,t} \sim \mathcal{D}_{n,t}}{\mathbb{E}} \left(\|\mathbf{x}_t^* - \bar{\mathbf{x}}_t\|^2 - \|\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}\|^2 - \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2 \right).\end{aligned}\tag{5}$$

Besides, we have

$$\begin{aligned}&\|\mathbf{x}_{t+1}^* - \bar{\mathbf{x}}_{t+1}\|^2 - \|\mathbf{x}_t^* - \bar{\mathbf{x}}_{t+1}\|^2 \\ &= \|\mathbf{x}_{t+1}^*\|^2 - \|\mathbf{x}_t^*\|^2 - 2 \langle \bar{\mathbf{x}}_{t+1}, -\mathbf{x}_t^* + \mathbf{x}_{t+1}^* \rangle \\ &= (\|\mathbf{x}_{t+1}^*\| - \|\mathbf{x}_t^*\|) (\|\mathbf{x}_{t+1}^*\| + \|\mathbf{x}_t^*\|) - 2 \langle \bar{\mathbf{x}}_{t+1}, -\mathbf{x}_t^* + \mathbf{x}_{t+1}^* \rangle \\ &\leq \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\| (\|\mathbf{x}_{t+1}^*\| + \|\mathbf{x}_t^*\|) - 2 \|\bar{\mathbf{x}}_{t+1}\| \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\| \\ &\leq 4\sqrt{R} \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\|. \quad (\text{due to } \|\mathbf{x} - \mathbf{y}\|^2 \leq R, \forall \mathbf{x}, \mathbf{y} \in \mathcal{X})\end{aligned}$$

Thus, telescoping $I_3(t)$ over $t \in [T]$, we have

$$\begin{aligned}&\sum_{t=1}^T I_3(t) \\ &\leq \frac{1}{2\eta} \underset{\Xi_{n,T} \sim \mathcal{D}_{n,T}}{\mathbb{E}} \left(4\sqrt{R} \sum_{t=1}^T \|\mathbf{x}_{t+1}^* - \mathbf{x}_t^*\| + \|\bar{\mathbf{x}}_1^* - \bar{\mathbf{x}}_1\|^2 - \|\bar{\mathbf{x}}_T^* - \bar{\mathbf{x}}_{T+1}\|^2 \right) - \frac{1}{2\eta} \underset{\Xi_{n,T} \sim \mathcal{D}_{n,T}}{\mathbb{E}} \sum_{t=1}^T \underset{\Xi_{n,t} \sim \mathcal{D}_{n,t}}{\mathbb{E}} \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2\end{aligned}$$

$$\leq \frac{1}{2\eta} \left(4\sqrt{RM} + R \right) - \frac{1}{2\eta} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \sum_{t=1}^T \|\bar{\mathbf{x}}_t - \bar{\mathbf{x}}_{t+1}\|^2.$$

Combining those bounds of $I_1(t)$, $I_2(t)$ and $I_3(t)$ together, we finally obtain

$$\begin{aligned} & \mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{t=1}^T \sum_{i=1}^n f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - f_t(\mathbf{x}_t^*; \xi_{i,t}) \\ & \leq n \sum_{t=1}^T (I_1(t) + I_2(t) + I_3(t)) \\ & \leq \left(1 + 2\beta \left(\frac{1}{\nu} + 4 \right) \right) nT\beta G\eta + \left(L + \frac{\eta L^2}{\nu} + \frac{\nu}{2\eta} + 6\eta L^2 + \frac{\beta}{2\eta} \right) \mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{t=1}^T \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\ & \quad + \left(\frac{1}{\nu} + 4 \right) n (H_t(\bar{\mathbf{x}}_1) - H_t(\bar{\mathbf{x}}_{t+1})) \\ & \quad + 3nTLG\eta^2 \left(\frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n\nu} \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{RM} + R) \\ & \stackrel{\textcircled{1}}{\leq} \left(1 + 2\beta \left(\frac{1}{\nu} + 4 \right) \right) nT\beta G\eta + \left(L + \frac{\eta L^2}{\nu} + \frac{\nu}{2\eta} + 6\eta L^2 + \frac{\beta}{2\eta} \right) \frac{\eta^2}{(1-\rho)^2} \left(6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right) \\ & \quad + \left(\frac{1}{\nu} + 4 \right) n (H_t(\bar{\mathbf{x}}_1) - H_t(\bar{\mathbf{x}}_{t+1})) \\ & \quad + 3nTLG\eta^2 \left(\frac{1}{\nu} + 4 \right) + \frac{\nu(\eta + 8nL\eta^2) + 2nL\eta^2}{n\nu} \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{RM} + R). \end{aligned}$$

① holds due to Lemma 3, that is,

$$\mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{i=1}^n \sum_{t=1}^T \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \leq \frac{\eta^2}{(1-\rho)^2} \left(6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right).$$

When $\nu = \beta$, we finally have

$$\begin{aligned} & \mathcal{R}_T^{DOG} \\ & = \mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{t=1}^T \sum_{i=1}^n f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) - f_t(\mathbf{x}_t^*; \xi_{i,t}) \\ & \leq 11nT\beta G\eta + \left(L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) \frac{\eta^2}{(1-\rho)^2} \left(6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right) \\ & \quad + \left(\frac{1}{\beta} + 4 \right) n (H_t(\bar{\mathbf{x}}_1) - H_t(\bar{\mathbf{x}}_{t+1})) + 3nTLG\eta^2 \left(\frac{1}{\beta} + 4 \right) \\ & \quad + \frac{\beta(\eta + 8nL\eta^2) + 2nL\eta^2}{n\beta} \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{RM} + R) \\ & = \eta nTG \left(11\beta + \frac{6\eta}{(1-\rho)^2} \left(L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) + 3L\eta \left(\frac{1}{\beta} + 4 \right) \right) + \left(\frac{1}{\beta} + 4 \right) n (H_t(\bar{\mathbf{x}}_1) - H_t(\bar{\mathbf{x}}_{t+1})) \\ & \quad + \left(\left(L + \frac{\eta L^2}{\beta} + 6\eta L^2 + \frac{\beta}{\eta} \right) \frac{4\eta^2}{(1-\rho)^2} + \frac{\beta(\eta + 8nL\eta^2) + 2nL\eta^2}{n\beta} \right) \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 + \frac{n}{2\eta} (4\sqrt{RM} + R). \end{aligned}$$

It completes the proof. \square

Lemma 1. For any β with $0 \leq \beta \leq 1$, we have

$$\mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \leq 6G + 4\sigma_{i,t}^2.$$

Proof.

$$\begin{aligned} & \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\ &= \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\beta \nabla g_{i,t}(\mathbf{x}_{i,t}) + (1 - \beta) \nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\ &\leq 2\beta^2 \|\nabla g_{i,t}(\mathbf{x}_{i,t})\|^2 + 2(1 - \beta)^2 \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\ &\leq 2G\beta^2 + 2(1 - \beta)^2 \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - \nabla H_t(\mathbf{x}_{i,t}) + \nabla H_t(\mathbf{x}_{i,t})\|^2 \\ &\leq 2G\beta^2 + 4(1 - \beta)^2 \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla h_t(\mathbf{x}_{i,t}; \xi_{i,t}) - \nabla H_t(\mathbf{x}_{i,t})\|^2 + 4(1 - \beta)^2 \|\nabla H_t(\mathbf{x}_{i,t})\|^2 \\ &\leq 2G\beta^2 + 4(1 - \beta)^2 \sigma_{i,t}^2 + 4(1 - \beta)^2 G \\ &\leq 6G + 4\sigma_{i,t}^2. \end{aligned}$$

\square

Lemma 2.

$$\begin{aligned} & \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 \\ &\leq \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) + 2G\eta\beta^2 + \frac{\eta L^2(1 - \beta)^2}{n} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \sum_{i=1}^n \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2. \end{aligned} \quad (6)$$

Proof.

$$\begin{aligned} & \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} H_t(\bar{\mathbf{x}}_{t+1}) \\ &\leq \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} H_t(\bar{\mathbf{x}}_t) + \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \langle \nabla H_t(\bar{\mathbf{x}}_t), \bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t \rangle + \frac{L}{2} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\bar{\mathbf{x}}_{t+1} - \bar{\mathbf{x}}_t\|^2 \\ &= \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} H_t(\bar{\mathbf{x}}_t) + \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \left\langle \nabla H_t(\bar{\mathbf{x}}_t), -\frac{\eta}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right\rangle + \frac{L}{2} \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \left\| \frac{\eta}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right\|^2 \\ &\leq \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} H_t(\bar{\mathbf{x}}_t) + \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\langle \nabla H_t(\bar{\mathbf{x}}_t), -\frac{\eta}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\rangle + \frac{L\eta^2}{2n} \sum_{i=1}^n \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\ &= \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} H_t(\bar{\mathbf{x}}_t) + \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left(\left\| \nabla H_t(\bar{\mathbf{x}}_t) - \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 - \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 - \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 \right) \\ &\quad + \frac{L\eta^2}{2n} \sum_{i=1}^n \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \\ &\stackrel{\textcircled{1}}{\leq} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} H_t(\bar{\mathbf{x}}_t) + \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \nabla H_t(\bar{\mathbf{x}}_t) - \frac{1}{n} \sum_{i=1}^n (\beta \nabla g_{i,t}(\mathbf{x}_{i,t}) + (1 - \beta) \nabla h_t(\mathbf{x}_{i,t})) \right\|^2 \end{aligned}$$

$$\begin{aligned}
& -\frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 - \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2 \\
& \leq \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} H_t(\bar{\mathbf{x}}_t) - \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 - \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2 \\
& \quad + \frac{\eta}{2} \left(2\beta^2 \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \nabla H_t(\bar{\mathbf{x}}_t) - \frac{1}{n} \sum_{i=1}^n \nabla g_{i,t}(\mathbf{x}_{i,t}) \right\|^2 + 2(1-\beta)^2 \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \nabla H_t(\bar{\mathbf{x}}_t) - \frac{1}{n} \sum_{i=1}^n \nabla H_t(\mathbf{x}_{i,t}) \right\|^2 \right) \\
& \leq \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} H_t(\bar{\mathbf{x}}_t) + \frac{\eta}{2} \left(4\beta^2 G + 2(1-\beta)^2 L^2 \frac{1}{n} \sum_{i=1}^n \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 \right) - \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 \\
& \quad - \frac{\eta}{2} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2 \\
& \leq \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} H_t(\bar{\mathbf{x}}_t) + 2\eta\beta^2 G + \eta(1-\beta)^2 L^2 \left(\frac{1}{n} \sum_{i=1}^n \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 \right) \\
& \quad - \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 - \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

① holds due to Lemma 1. Finally, we obtain

$$\begin{aligned}
& \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 \\
& \leq \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\nabla H_t(\bar{\mathbf{x}}_t)\|^2 + \frac{\eta}{2} \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \left\| \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}) \right\|^2 \\
& \leq \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} (H_t(\bar{\mathbf{x}}_t) - H_t(\bar{\mathbf{x}}_{t+1})) + 2G\eta\beta^2 + \frac{\eta L^2(1-\beta)^2}{n} \sum_{i=1}^n \mathbb{E}_{\Xi_{n,t-1} \sim \mathcal{D}_{n,t-1}} \|\bar{\mathbf{x}}_t - \mathbf{x}_{i,t}\|^2 + 3GL\eta^2 + \frac{2L\eta^2}{n} \sum_{i=1}^n \sigma_{i,t}^2.
\end{aligned}$$

It completes the proof. \square

Lemma 3.

$$\mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{i=1}^n \sum_{t=1}^T \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \leq \frac{\eta^2}{(1-\rho)^2} \left(6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right).$$

Proof. Recall that

$$\mathbf{x}_{i,t+1} = \sum_{j=1}^n \mathbf{W}_{ij} \mathbf{x}_{j,t} - \eta \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}),$$

and

$$\bar{\mathbf{x}}_{t+1} = \bar{\mathbf{x}}_t - \eta \left(\frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right).$$

Denote

$$\mathbf{X}_t = [\mathbf{x}_{1,t}, \mathbf{x}_{2,t}, \dots, \mathbf{x}_{n,t}] \in \mathbb{R}^{d \times n},$$

$$\mathbf{G}_t = [\nabla f_{1,t}(\mathbf{x}_{1,t}; \xi_{1,t}), \nabla f_{2,t}(\mathbf{x}_{2,t}; \xi_{2,t}), \dots, \nabla f_{n,t}(\mathbf{x}_{n,t}; \xi_{n,t})] \in \mathbb{R}^{d \times n}.$$

By letting $\mathbf{x}_{i,1} = \mathbf{0}$ for any $i \in [n]$, the update rule is re-formulated as

$$\mathbf{X}_{t+1} = \mathbf{X}_t \mathbf{W} - \eta \mathbf{G}_t = - \sum_{s=1}^t \eta \mathbf{G}_s \mathbf{W}^{t-s}.$$

Similarly, denote $\bar{\mathbf{G}}_t = \frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})$, and we have

$$\bar{\mathbf{x}}_{t+1} = \bar{\mathbf{x}}_t - \eta \left(\frac{1}{n} \sum_{i=1}^n \nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t}) \right) = - \sum_{s=1}^t \eta \bar{\mathbf{G}}_s. \quad (7)$$

Therefore,

$$\begin{aligned} & \sum_{i=1}^n \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\ & \stackrel{\textcircled{1}}{=} \sum_{i=1}^n \left\| \sum_{s=1}^{t-1} \eta \bar{\mathbf{G}}_s - \eta \mathbf{G}_s \mathbf{W}^{t-s-1} \mathbf{e}_i \right\|^2 \\ & \stackrel{\textcircled{2}}{=} \left\| \sum_{s=1}^{t-1} \eta \mathbf{G}_s \mathbf{v}_1 \mathbf{v}_1^T - \eta \mathbf{G}_s \mathbf{W}^{t-s-1} \right\|_F^2 \\ & \stackrel{\textcircled{3}}{\leq} \left(\eta \rho^{t-s-1} \left\| \sum_{s=1}^{t-1} \mathbf{G}_s \right\|_F \right)^2 \\ & \leq \left(\sum_{s=1}^{t-1} \eta \rho^{t-s-1} \|\mathbf{G}_s\|_F \right)^2. \end{aligned}$$

① holds due to \mathbf{e}_i is a unit basis vector, whose i -th element is 1 and other elements are 0s. ② holds due to $\mathbf{v}_1 = \frac{1}{\sqrt{n}}$. ③ holds due to Lemma 4.

Thus, we have

$$\begin{aligned} & \mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{i=1}^n \sum_{t=1}^T \|\mathbf{x}_{i,t} - \bar{\mathbf{x}}_t\|^2 \\ & \leq \mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{t=1}^T \left(\sum_{s=1}^{t-1} \eta \rho^{t-s-1} \|\mathbf{G}_s\|_F \right)^2 \\ & \stackrel{\textcircled{1}}{\leq} \frac{\eta^2}{(1-\rho)^2} \mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \left(\sum_{t=1}^T \|\mathbf{G}_t\|_F^2 \right) \\ & = \frac{\eta^2}{(1-\rho)^2} \left(\mathbb{E}_{\Xi_{n,T} \sim \mathcal{D}_{n,T}} \sum_{t=1}^T \sum_{i=1}^n \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \right) \\ & = \frac{\eta^2}{(1-\rho)^2} \left(\sum_{t=1}^T \sum_{i=1}^n \mathbb{E}_{\Xi_{n,t} \sim \mathcal{D}_{n,t}} \|\nabla f_{i,t}(\mathbf{x}_{i,t}; \xi_{i,t})\|^2 \right) \\ & \stackrel{\textcircled{2}}{\leq} \frac{\eta^2}{(1-\rho)^2} \left(6nTG + 4 \sum_{t=1}^T \sum_{i=1}^n \sigma_{i,t}^2 \right). \end{aligned}$$

① holds due to Lemma 5. ② holds due to Lemma 1. It completes the proof. \square

Lemma 4 (Appeared in Lemma 5 in [Tang et al., 2018]). *For any matrix $\mathbf{X}_t \in \mathbb{R}^{d \times n}$, decompose the confusion matrix \mathbf{W} as $\mathbf{W} = \sum_{i=1}^n \lambda_i \mathbf{v}_i \mathbf{v}_i^\top = \mathbf{P} \mathbf{\Lambda} \mathbf{P}^\top$, where $\mathbf{P} = [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n] \in \mathbb{R}^{n \times n}$, \mathbf{v}_i is the normalized eigenvector of λ_i . $\mathbf{\Lambda}$ is a diagonal matrix, and λ_i be its i -th element. We have*

$$\|\mathbf{X}_t \mathbf{W}^t - \mathbf{X}_t \mathbf{v}_1 \mathbf{v}_1^\top\|_F^2 \leq \|\rho^t \mathbf{X}_t\|_F^2,$$

where $\rho = \max\{|\lambda_2(\mathbf{W})|, |\lambda_n(\mathbf{W})|\}$.

Lemma 5 (Appeared in Lemma 6 in [Tang et al., 2018]). *Given two non-negative sequences $\{a_t\}_{t=1}^\infty$ and $\{b_t\}_{t=1}^\infty$ that satisfying*

$$a_t = \sum_{s=1}^t \rho^{t-s} b_s,$$

with $\rho \in [0, 1)$, we have

$$\sum_{t=1}^k a_t^2 \leq \frac{1}{(1-\rho)^2} \sum_{s=1}^k b_s^2.$$

References

H. Tang, S. Gan, C. Zhang, T. Zhang, and J. Liu. Communication Compression for Decentralized Training. *arXiv.org*, Mar. 2018.