

Shared feature: Linear constant coefficient (LCC) equations (general N -th order and 2nd order example)

$$\sum_{n=0}^N a_n \frac{d^n \phi}{dx^n} = 0 \quad \xrightarrow{N=2} \quad a_2 \frac{d^2 \phi}{dx^2} + a_1 \frac{d\phi}{dx} + a_0 \phi = 0 \quad (1)$$

and Cauchy-Euler (CE) equations

$$\sum_{n=0}^N b_n x^n \frac{d^n \phi}{dx^n} = 0 \quad \xrightarrow{N=2} \quad b_2 x^2 \frac{d^2 \phi}{dx^2} + b_1 x \frac{d\phi}{dx} + b_0 \phi = 0 \quad (2)$$

These are homogeneous equations, whose general solutions can be written as linear combinations of N linearly independent fundamental solutions:

$$\phi_{\text{gen}}(x) = \sum_{\ell=1}^N c_{\ell} \phi_{\ell}(x) \quad \xrightarrow{N=2} \quad \phi_{\text{gen}}(x) = c_1 \phi_1(x) + c_2 \phi_2(x) \quad (3)$$

where each $\phi_{\ell}(x)$ can be determined using appropriate trial solutions that reduce the ODE to algebra for finding roots of a characteristic polynomial. The c_{ℓ} coeffs get selected by additional BC or IC conditions.

1 LCC solutions

For LCC equations, the trial solution is $\boxed{\phi(x) = e^{mx}}$. The values for the exponents m are roots of the characteristic polynomial equation:

$$a_n m^n + a_{n-1} m^{n-1} + \cdots + a_2 m^2 + a_1 m + a_0 = 0 \quad (a_2 m^2 + a_1 m + a_0 = 0) \quad (4)$$

- If m_* is a multiple root (order k) then it yields k distinct $\phi_{\ell}(x)$ solutions in the form:

$$e^{m_* x} \quad x e^{m_* x} \quad x^2 e^{m_* x} \quad \cdots \quad x^{k-1} e^{m_* x} \quad (5)$$

- If all the coefficients in the ODE are real, then if $m_* = \alpha + i\beta$ is a complex root then the complex-conjugate $\overline{m_*} = \alpha - i\beta$ is always also a root. Therefore, two ϕ_{ℓ} solutions are

$$c_1 e^{(\alpha+i\beta)x} + c_2 e^{(\alpha-i\beta)x} = c_1 e^{\alpha x} e^{i\beta x} + c_2 e^{\alpha x} e^{-i\beta x} \quad (6)$$

using $e^{c+d} = e^c e^d$ and Euler's formula, $e^{i\theta} = \cos \theta + i \sin \theta$, the solutions can always be re-written in terms of only real functions:

$$= c_1 e^{\alpha x} (\cos(\beta x) + i \sin(\beta x)) + c_2 e^{\alpha x} (\cos(\beta x) - i \sin(\beta x)) = c_3 \boxed{e^{\alpha x} \cos(\beta x)} + c_4 \boxed{e^{\alpha x} \sin(\beta x)} \quad (7)$$

For real-valued problems, always use form (7), not (6)!

2 CE solutions

For CE equations, the trial solution is $\boxed{\phi(x) = x^m}$. The powers m are roots of the characteristic polynomial equation (for an example 4th order CE ODE):

$$b_4 m(m-1)(m-2)(m-3) + b_3 m(m-1)(m-2) + b_2 m(m-1) + b_1 m + b_0 = 0 \quad (8)$$

- If m_* is a k^{th} order multiple root, then solutions are $x^{m_*}, x^{m_*} \ln(x), x^{m_*} (\ln(x))^2, \dots, x^{m_*} (\ln(x))^{k-1}$.
- If $m_* = \alpha + i\beta$ (and $\overline{m_*} = \alpha - i\beta$) are complex roots, then using $x^r = (e^{\ln(x)})^r = e^{r \ln(x)}$ and $e^{i\theta} = \cos \theta + i \sin \theta$:

$$\begin{aligned} c_1 x^{\alpha+i\beta} + c_2 x^{\alpha-i\beta} &= c_1 e^{\alpha \ln(x)} (\cos(\beta \ln(x)) + i \sin(\beta \ln(x))) + c_2 e^{\alpha \ln(x)} (\cos(\beta \ln(x)) - i \sin(\beta \ln(x))) \\ &= c_3 \boxed{x^{\alpha} \cos(\beta \ln(x))} + c_4 \boxed{x^{\alpha} \sin(\beta \ln(x))} \quad (\text{use this form!}) \end{aligned}$$

Inhomogeneous LCC or CE equations $Ly = f$ can be solved in terms of sums of homogeneous and particular solutions (see summary sheet on the method of Un-determined Coefficients) or using variation of parameters or eigenfunction expansions.