

Linear algebra, Exercise 8

Luqing Ye*

August 16, 2014

Exercise (Tao). For this question, the field of scalars will be the complex numbers instead of the reals (i.e., all matrices, etc. are allowed to have complex entries). Let θ be a real number, and let A be the 2×2 rotational matrix

$$A = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

- Show that A has eigenvalues $e^{i\theta}, e^{-i\theta}$. What are the eigenvectors corresponding to $e^{i\theta}$ and $e^{-i\theta}$?
- Write $A = QDQ^{-1}$ for some invertible matrix Q and diagonal matrix D .

Proof.

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \lambda \begin{pmatrix} x \\ y \end{pmatrix},$$

so

$$\begin{pmatrix} \cos \theta - \lambda & -\sin \theta \\ \sin \theta & \cos \theta - \lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

where $(x, y) \neq (0, 0)$. So

$$(\cos \theta - \lambda)^2 + \sin^2 \theta = 0 \iff \lambda^2 - 2\lambda \cos \theta + 1 = 0 \iff \lambda = \cos \theta \pm i \sin \theta.$$

When $\lambda = \cos \theta + i \sin \theta$, then

$$\begin{pmatrix} -i \sin \theta & -\sin \theta \\ \sin \theta & -i \sin \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

So when $\theta = \pi k, k \in \mathbf{Z}$, the eigenvector corresponding to $e^{i\theta}$ is arbitrary, otherwise, the eigenvector is in the form of $(iy, y), y \in \mathbf{C} \setminus \{0\}$.

When $\lambda = \cos \theta - i \sin \theta$, then $ix \sin \theta - y \sin \theta = 0$. When $\theta = \pi k, k \in \mathbf{Z}$, then the eigenvector corresponding to $e^{i\theta}$ is arbitrary. Otherwise, the eigenvector is in the form of $(x, ix), x \in \mathbf{C} \setminus \{0\}$.

- Denote the linear transformation corresponding to the matrix A by L_A . The eigenvector of $A, (i, 1)$ and $(1, i)$, are linearly independent in \mathbf{C} . Let $\alpha = ((i, 1), (1, i)) = (w_1, w_2)$ be an ordered basis, let $\beta = ((1, 0), (0, 1)) = (v_1, v_2)$ be another ordered basis. Then

$$[L_A]_{\alpha}^{\alpha} = \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix}.$$

And

$$\begin{cases} w_1 = iv_1 + v_2, \\ w_2 = v_1 + iv_2. \end{cases}$$

So

$$[I]_{\alpha}^{\beta} = \begin{pmatrix} i & 1 \\ 1 & i \end{pmatrix}, [I]_{\beta}^{\alpha} = \begin{pmatrix} -i & \frac{1}{2} \\ \frac{1}{2} & -i \end{pmatrix}.$$

*叶卢庆 (1992—), E-mail: yeluqingmathematics@gmail.com

So

$$A = [L_A]^\beta_\beta = [I]^\beta_\alpha [L_A]^\alpha_\alpha [I]^\alpha_\beta.$$

Done.

□

<http://blog.sciencenet.cn/u/Yaleking>