

Course outline

① Fundamentals

- ▶ Notation
- ▶ Functions
- ▶ Approximation

② Series

- ▶ Summation
- ▶ Taylor series

③ Linear algebra


- ▶ Representing big, complex, data
- ▶ Systems of equations
- ▶ Dimension reduction

④ Probability


- ▶ Discrete random variables
- ▶ Continuous random variables & integration

⑤ Optimisation

Taylor series approximation




Linear approximation



Orange line is the tangent line at $a = 0.5$:

$$P_1(x) = f(a) + f'(x_0)(x - a)$$


Quadratic approximation



Green line is a quadratic function:

$$P_2(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2$$

Taylor polynomial approximation



In general:

$$P_n(x) = \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (x - a)^i$$

Linear approximation

How do we get to Taylor polynomials?

Let's find a degree 1 polynomial P_1 such that the value of $P_1(a)$ and $P_1'(a)$ agree with f at $x = a$.

Linear approximation

Example

Find the first order Taylor polynomial for $f(x) = \ln(x)$ at $a = 1$ and use it to approximate $\ln(1.1)$.

Quadratic approximation

Find a degree 2 polynomial P_2 such that $P_2(a)$, $P_2'(a)$, and $P_2''(a)$ agree with f at a .

Quadratic approximation

Example

Approximate $\ln 1.1$ using the quadratic $P_2(x)$.

General polynomial approximation

n th degree polynomial approximation:

$$P_n(x) = c_0 + c_1(x - a) + c_2(x - a)^2 + \cdots + c_n(x - a)^n$$

Demand $P_n(a)$, $P'_n(a)$, ..., $P_n^{(n)}(a)$ agree with f at a .

Taylor and Maclaurin polynomials

Definition (Taylor Polynomial of degree n for $f(x)$ at $x = a$)

$$P_n(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \cdots \\ + \frac{f^{(n)}(a)}{n!}(x - a)^n$$

Taylor and Maclaurin polynomials

Definition (Taylor Polynomial of degree n for $f(x)$ at $x = a$)

$$P_n(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \cdots \\ + \frac{f^{(n)}(a)}{n!}(x - a)^n$$

Definition (Maclaurin Polynomial of degree n for $f(x)$)

$$P_n(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \cdots + \frac{f^{(n)}(0)}{n!}x^n$$

Taylor and Maclaurin polynomials

Example

Find the Taylor polynomial of degree n for $\ln x$ with $a = 1$. Use $P_4(x)$ to estimate $\ln 1.1$.

Theorem (Taylor's Theorem)

Suppose the function f has $(n + 1)$ derivatives on some interval containing a and x . Then if

$$P_n(x) = f(a) + f'(a)(x - a) + \cdots + \frac{f^{(n)}(a)}{n!}(x - a)^n$$

is the n th Taylor polynomial of f at a , $f(x) = P_n(x) + R_n(x)$ where the remainder

$$R_n(x) = \frac{f^{(n+1)}(z)}{(n + 1)!}(x - a)^{n+1}$$

for some number z between a and x .

Proof by reference: Stewart, pg. 787-9

Taylor's Theorem

The point of this is:

$$\begin{aligned}\text{error} &= |f(x) - P_n(x)| = |R_n(x)| \\ &= \left| \frac{f^{(n+1)}(z)}{(n+1)!} (x-a)^{n+1} \right|.\end{aligned}$$

If we can find C such that

$$|f^{(n+1)}(z)| \leq C, \quad \forall z \in [a, x]$$

then

$$|f(x) - P_n(x)| \leq \frac{C}{(n+1)!} |x-a|^{n+1}.$$

Taylor's Theorem

Example

Determine the accuracy of the use of $P_4(x)$ to estimate $\ln(1.1)$.

Taylor's Theorem

Example

Determine the accuracy of the use of $P_4(x)$ to estimate $\ln(1.1)$.

Example

If we use the Maclaurin polynomial $P_n(x)$ for e^x , find the smallest value of n which gives $e = e^1$ to within the accuracy of 0.000005 (i.e 5-dp's).

Taylor's Theorem

Example

Determine the accuracy of the use of $P_4(x)$ to estimate $\ln(1.1)$.

Example

If we use the Maclaurin polynomial $P_n(x)$ for e^x , find the smallest value of n which gives $e = e^1$ to within the accuracy of 0.000005 (i.e 5-dp's).

Example

Use a Maclaurin polynomial of degree 3 for $(1+x)^{1/2}$ to approximate $\sqrt{5}$. Estimate the error.

Taylor's Theorem

Example

Determine the accuracy of the use of $P_4(x)$ to estimate $\ln(1.1)$.

Example

If we use the Maclaurin polynomial $P_n(x)$ for e^x , find the smallest value of n which gives $e = e^1$ to within the accuracy of 0.000005 (i.e 5-dp's).

Example

Use a Maclaurin polynomial of degree 3 for $(1+x)^{1/2}$ to approximate $\sqrt{5}$. Estimate the error.

Example

Find the Maclaurin polynomial of degree $n = 2k$ for $f(x) = \cos x$.

Taylor series

Definition (Taylor series of f at a)

If the function f has derivatives of all orders, then

$$f(x) = \lim_{n \rightarrow \infty} P_n(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x - a)^k.$$

If $a = 0$ this is the *Maclaurin series*.

Some important Taylor series

$$e^x : \quad 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$\sin x : \quad x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

$$\cos x : \quad 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

$$\frac{1}{1-x} : \quad 1 + x + x^2 + x^3 + \cdots = \sum_{n=0}^{\infty} x^n$$

$$(1+x)^k = 1 + \sum_{n=1}^{\infty} \binom{k}{n} x^n$$

Taylor series

Theorem (Convergence of Power Series)

Consider the power series

$$\sum_{n=0}^{\infty} a_n x^n.$$

Then either

- (1) the series converges for all values of x , or*
- (2) the series converges only for $x = 0$, or*
- (3) there exists a number $R > 0$ such that $\sum_{n=0}^{\infty} a_n x^n$ converges for all x with $|x| < R$ and diverges for all x with $|x| > R$.*

The number R is called the radius of convergence. In case (1) we often write " $R = \infty$ " and in case (2), $R = 0$.


Taylor series

Example


Find the intervals of convergence for

- $\cos(x)$
- $1/(1 - x)$

Who cares about Taylor series?



Who cares about Taylor series?



Who cares about Taylor series?

Define a *loss function* $f(m)$:

$$\begin{aligned} f(m) &= \sum_{i \in \{\text{misclassifications}\}} d((x_i, y_i) \text{ from line } y = -mx) \\ &= \sum_{i \in \{\text{misclassifications}\}} \frac{|mx_i + y_i|}{\sqrt{m^2 + 1}} \end{aligned}$$

This function $f(m)$ looks gross! No fun at all to differentiate.

Who cares about Taylor series?

Find the (approximate) minimum using *gradient descent*:

- 1 Guess a solution m
- 2 Change m by some amount h , $m \rightarrow m + h$, such that $f(m + h) < f(m)$
- 3 If h is very small, STOP. Otherwise GOTO 2.

But how do we choose h ?

Who cares about Taylor series?


Let $x = m + h$, $a = m$. Taylor polynomial ($n = 1$):

$$f(m + h) \approx f(m) + f'(m)h$$


Choose $h = -\eta f'(m)$, so

$$f(m + h) = f(m) - \eta (f'(m))^2 < f(m)$$


Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?




Who cares about Taylor series?



Who cares about Taylor series?



Who cares about Taylor series?



Who cares about Taylor series?

