

Mathematical Physics Studies

Walter D. van Suijlekom

# Noncommutative Geometry and Particle Physics

*Second Edition*

OPEN ACCESS



Springer

# **Mathematical Physics Studies**

## **Series Editors**

Giuseppe Dito, Institut de Mathématiques de Bourgogne, Université de Bourgogne, Dijon, France

Edward Frenkel, Department of Mathematics, University of California at Berkley, Berkeley, CA, USA

Sergei Gukov, California Institute of Technology, Pasadena, CA, USA

Yasuyuki Kawahigashi, Department of Mathematical Sciences, The University of Tokyo, Tokyo, Japan

Maxim Kontsevich, Institut des Hautes Etudes Scientifiques, Bures-sur-Yvette, France

Nicolaas P. Landsman, Chair of Mathematical Physics, Radboud Universiteit Nijmegen, Nijmegen, Gelderland, The Netherlands

Bruno Nachtergaele, Department of Mathematics, University of California, Davis, CA, USA

Hal Tasaki, Department of Physics, Gakushuin University, Tokyo, Japan

The series publishes research monographs that address physics problems within a rigorous mathematical framework, highlighting recent developments at the intersection of mathematics and theoretical physics. Research areas covered include dynamical systems, equilibrium and nonequilibrium statistical mechanics, general relativity, classical and integrable systems, many-body quantum systems and condensed matter physics, partial differential equations, probability and random structures, particle physics, quantum field theory, quantum information, quantum mechanics and spectral theory, and string theory and quantum gravity. Each monograph delves into these areas, emphasizing their mathematical foundations and applications in modern physics.

Walter D. van Suijlekom

# Noncommutative Geometry and Particle Physics

Second Edition



Springer

Walter D. van Suijlekom  
Department of Astrophysics, Institute for  
Mathematics, Astrophysics and Particle  
Physics (IMAPP)  
Radboud University  
Nijmegen, The Netherlands



ISSN 0921-3767  
Mathematical Physics Studies  
ISBN 978-3-031-59119-8  
<https://doi.org/10.1007/978-3-031-59120-4>

ISSN 2352-3905 (electronic)  
ISBN 978-3-031-59120-4 (eBook)

1<sup>st</sup> edition: © Springer Science+Business Media Dordrecht 2015  
2<sup>nd</sup> edition: © The Editor(s) (if applicable) and The Author(s) 2025. This book is an open access publication.

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG  
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

If disposing of this product, please recycle the paper.

# Preface to the Second Edition

After the appearance of the first edition of this book there have been numerous developments in the field that urged me to update the book. On the mathematical side, new concepts, techniques and results have been found during the last few years, which are now included in our treatment. This includes the perturbation semigroup and cyclic cocycles in the Taylor expansion of the spectral action. On the particle physics side, the step to go Beyond the Standard Model with Pati–Salam unification now forms a prominent part of our book, while the first attempts towards quantization of the theory are included in the final chapter.

I have also revised the introductory part I, aiming for a more complete treatment of noncommutative differential geometry. There are more details on the analytical properties of the Dirac operator on a compact Riemannian  $\text{spin}^c$  manifold, and also more noncommutative examples to better illustrate the concept of a spectral triple. For example, the noncommutative torus that formed one of the first examples of a noncommutative Riemannian spin manifold now makes its appearance in Sect. 5.3.1. It reappears for illustrative examples of even and odd cyclic cocycles in Sect. 6.2.1.

I would like to further thank Calum Beck, Rui Dong, Eva-Maria Hekkelman, Teije Kuijper, Malte Leimbach, Teun van Nuland, Leo Polak and Berend Visser for sending me numerous typos in the first edition and in a draft version of the second edition.

I thank Leonora, Joris, Daniël and Mathilde for their continuing patience and love.

Nijmegen, The Netherlands  
December 2023

Walter D. van Suijlekom

# Preface to the First Edition

The seeds of this book have been planted in the far east, where I wrote lecture notes for international schools in Tianjin, China in 2007 and in Bangkok, Thailand in 2011. I then realized that an up-to-date text for beginning noncommutative geometers on the applications of this rather new mathematical field to particle physics was missing in the literature.

This made me decide to transform my notes into the form of a book. Besides the given challenge inherent in such a project, this was not made easy because of recent, rapid developments in the field, making it difficult to choose what to include and to decide where to stop in my treatment. The current state of affairs is at least touched upon in the final chapter of this book, where I discuss the latest particle physics models in noncommutative geometry, and compare them to the latest experimental findings. With this, I hope to have provided a path that starts with the basic principles of noncommutative geometry and leads to the forefront of research in noncommutative geometry and particle physics.

The intended audience consists of mathematicians with some knowledge of particle physics, and of theoretical physicists with some mathematical background. Concerning the level of this textbook, for mathematicians I assume prerequisites on gauge theories at the level of e.g. [1, 2], and recommend to first read the book [3] to really appreciate the last few chapters of this book on particle physics/the Standard Model. For physicists, I assume knowledge of some basic algebra, Hilbert space and operator theory (e.g. [4, Chapter 2]), and Riemannian geometry (e.g. [5, 6]). This makes the book particularly suitable for a starting PhD student, after a master degree in mathematical/theoretical physics including the above background.

I would like to thank the organizers and participants of the aforementioned schools for their involvement and their feedback. This also applies to the MRI-Masterclass in Utrecht in 2010 and the Conference on index theory in Bogotá in 2008, where Chapter 6 finds its roots. Much feedback on previous drafts was gratefully received from students in my class on noncommutative geometry in Nijmegen: Bas Jordans, Joey van der Leer and Sander Uijlen. I thank my students and co-authors Jord Boeinink, Thijs van den Broek and Koen van den Dungen for allowing me to transcribe part of our results in the present book form. Simon Brain, Alan Carey and Adam

Rennie are gratefully acknowledged for their feedback and suggested corrections. Strong motivation to writing this book was given to me by my co-author Matilde Marcolli. I thank Gerard Bäuerle, Gianni Landi and Klaas Landsman for having been my main tutors in writing, and Klaas in particular for a careful final proofreading. I also thank Aldo Rampioni at Springer for his help and guidance. I thank Alain Connes for his inspiration and enthusiasm for the field, without whose work this book could of course not have been written.

I am thankful to my family and friends for their continuous love and support. My deepest gratitude goes to Mathilde for being my companion in life, and to Daniël for making sure that the final stages of writing were frequently, and happily, interrupted.

April 2014

Walter D. van Suijlekom

## References

1. D. Bleecker. *Gauge theory and variational principles*. Addison-wesley, Reading, 1981.
2. J. Baez and J. P. Muniain. *Gauge fields, knots and gravity*, volume 4 of *Series on Knots and Everything*. World Scientific Publishing Co. Inc., River Edge, NJ, 1994.
3. W. N. Cottingham and D. A. Greenwood. *An Introduction to the Standard Model of Particle Physics*. Cambridge University Press, Cambridge, second edition, 2007.
4. W. Thirring. *Quantum mathematical physics*. Springer-Verlag, Berlin, second edition, 2002. Atoms, molecules and large systems, Translated from the 1979 and 1980 German originals by Evans M. Harrell II.
5. J. Jost. *Riemannian Geometry and Geometric Analysis*. Universitext. Springer-Verlag, Berlin, third edition, 2002.
6. M. Nakahara. *Geometry, Topology and Physics*. IOP Publishing, 1990.

# Contents

<b>1</b>	<b>Introduction</b>	1
	References	5
 <b>Part I Noncommutative Geometric Spaces</b>		
<b>2</b>	<b>Finite Noncommutative Spaces</b>	9
2.1	Finite Spaces and Matrix Algebras	9
2.1.1	Commutative Matrix Algebras	13
2.1.2	Finite Spaces and Matrix Algebras	13
2.2	Noncommutative Geometric Finite Spaces	18
2.2.1	Morphisms Between Finite Spectral Triples	23
2.3	Classification of Finite Spectral Triples	25
	Notes	28
	References	29
<b>3</b>	<b>Finite Real Noncommutative Spaces</b>	31
3.1	Finite Real Spectral Triples	31
3.1.1	Morphisms Between Finite Real Spectral Triples	33
3.2	Classification of Finite Real Spectral Triples	35
3.3	Real Algebras and Krajewski Diagrams	43
3.4	Classification of Irreducible Geometries	45
	Notes	46
	References	47
<b>4</b>	<b>Riemannian Spin Manifolds</b>	49
4.1	Clifford Algebras	49
4.1.1	Representation Theory of Clifford Algebras	53
4.2	Riemannian Spin Geometry	55
4.2.1	Spin Manifolds	55
4.2.2	Clifford Connections, Spin Connections and the Dirac Operator	59
4.2.3	Lichnerowicz Formula	63

4.3	The Dirac Operator: Analytical Aspects .....	64
4.3.1	Bounded Commutators .....	64
4.3.2	Essential Self-adjointness .....	66
4.3.3	Compact Resolvent .....	68
Notes .....		71
References .....		71
<b>5</b>	<b>Noncommutative Riemannian Spin Manifolds .....</b>	<b>73</b>
5.1	Gelfand Duality .....	73
5.2	Spectral Triples .....	75
5.3	Examples of Noncommutative Manifolds .....	80
5.3.1	The Noncommutative Torus .....	80
5.3.2	Generalization to Toric Manifolds .....	82
Notes .....		84
References .....		86
<b>6</b>	<b>The Local Index Formula in Noncommutative Geometry .....</b>	<b>89</b>
6.1	Local Index Formula on the Circle and on the Torus .....	89
6.1.1	The Winding Number on the Circle .....	89
6.1.2	The Winding Number on the Torus .....	91
6.2	Hochschild and Cyclic Cohomology .....	95
6.2.1	Cyclic Cocycles for the Noncommutative Torus .....	99
6.3	Abstract Differential Calculus .....	101
6.4	Residues and the Local $(b, B)$ -Cocycle .....	106
6.5	The Local Index Formula .....	110
6.6	The Local Index Formula for Toric Noncommutative Manifolds .....	113
Notes .....		114
References .....		116
<b>Part II Noncommutative Geometry and Gauge Theories</b>		
<b>7</b>	<b>Gauge Theories from Noncommutative Manifolds .....</b>	<b>121</b>
7.1	'Inner' Unitary Equivalences as the Gauge Group .....	121
7.1.1	The Gauge Algebra .....	124
7.2	Morita Self-equivalences as Gauge Fields .....	125
7.2.1	Morita Equivalence .....	125
7.2.2	Morita Equivalence and Spectral Triples .....	130
7.3	Inner Fluctuations Without the First-Order Condition .....	133
7.3.1	Special Case $\mathcal{E} = \mathcal{A}$ and Inner Fluctuations .....	133
7.3.2	The Semi-group of Inner Perturbations .....	136
Notes .....		143
References .....		144

<b>8 Localization of Gauge Theories from Noncommutative Geometry</b>	145
8.1 Commutative Subalgebra and $C^*$ -Bundles	145
8.2 Localization of the Gauge Group	148
8.3 Localization of Gauge Fields	150
8.4 Localization of Toric Noncommutative Manifolds	151
Notes	154
References	155
<b>9 Spectral Invariants</b>	157
9.1 Spectral Action Functional	157
9.2 Asymptotic Expansion of the Spectral Action	159
9.3 Perturbative Expansion in the Gauge Field	161
9.3.1 Taylor Expansion of the Spectral Action	164
9.3.2 Cyclic Cocycles Underlying the Spectral Action	166
9.3.3 Brackets and Noncommutative Integrals Over Universal Forms	168
Notes	173
References	176
<b>10 Almost-Commutative Manifolds and Gauge Theories</b>	179
10.1 Gauge Symmetries of AC Manifolds	179
10.1.1 Unimodularity	181
10.2 Gauge Fields and Scalar Fields	183
10.2.1 Gauge Transformations	185
10.3 The Heat Expansion of the Spectral Action	186
10.3.1 A Generalized Lichnerowicz Formula	186
10.3.2 The Heat Expansion	190
10.4 The Spectral Action on AC Manifolds	193
Notes	199
References	200
<b>11 The Noncommutative Geometry of Electrodynamics</b>	203
11.1 The Two-Point Space	203
11.1.1 The Product Space	204
11.1.2 $U(1)$ Gauge Theory	206
11.2 Electrodynamics	207
11.2.1 The Finite Space	208
11.2.2 A Non-trivial Finite Dirac Operator	209
11.2.3 The Almost-Commutative Manifold	210
11.2.4 The Spectral Action	211
11.2.5 The Fermionic Action	212
11.2.6 Fermionic Degrees of Freedom	215
11.3 Grassmann Variables, Grassmann Integration and Pfaffians	216
Notes	217
References	218

<b>12</b>	<b>The Noncommutative Geometry of Yang–Mills Fields</b>	219
12.1	Spectral Triple Obtained from an Algebra Bundle	219
12.2	Yang–Mills Theory as a Noncommutative Manifold	223
12.2.1	From Algebra Bundles to Principal Bundles	223
12.2.2	Inner Fluctuations and Spectral Action	224
12.2.3	Topological Spectral Action	227
Notes	.....	227
References	.....	228
<b>13</b>	<b>The Noncommutative Geometry of the Standard Model</b>	229
13.1	The Finite Space	229
13.2	The Gauge Theory	233
13.2.1	The Gauge Group	233
13.2.2	The Gauge and Scalar Fields	235
13.3	The Spectral Action	239
13.3.1	Coupling Constants and Unification	244
13.3.2	The Higgs Mechanism	245
13.4	The Fermionic Action	249
Notes	.....	255
References	.....	255
<b>14</b>	<b>Phenomenology of the Noncommutative Standard Model</b>	257
14.1	Mass Relations	257
14.1.1	Fermion Masses	257
14.1.2	The Higgs Mass	258
14.1.3	The Seesaw Mechanism	259
14.2	Renormalization Group Flow	260
14.2.1	Coupling Constants	260
14.2.2	Renormalization Group Equations	262
14.2.3	Running Masses	263
14.2.4	Higgs Mass: Comparison to Experimental Results	267
Notes	.....	268
References	.....	269
<b>15</b>	<b>Beyond the Standard Model: Pati–Salam Unification</b>	271
15.1	The Finite Noncommutative Space of the Pati–Salam Model	271
15.2	The Gauge and Scalar Field Contents	273
15.3	Truncation to the Standard Model	276
15.4	Phenomenology of the Noncommutative Pati–Salam Model	278
15.4.1	Grand Unification of the Gauge Couplings	279
15.4.2	Running of the Higgs Mass	281
Notes	.....	284
References	.....	285

Contents	xiii
<b>16 Towards a Quantum Theory</b> .....	287
16.1 Second Quantization of Spectral Triples .....	287
16.1.1 KMS and a Dynamical System .....	288
16.1.2 Fermionic Second Quantization .....	289
16.1.3 von Neumann Information Theoretic Entropy .....	292
16.2 One-Loop Corrections to the Spectral Action .....	293
16.2.1 Ward Identity for the Gauge Propagator .....	295
16.2.2 Two-Point Functions at One-Loop .....	296
16.2.3 One-Loop Counterterms to the Spectral Action .....	298
Notes .....	300
References .....	300
<b>Appendix: Alphabetic Index, and Notation Index</b> .....	303
<b>Index</b> .....	311

# Chapter 1

## Introduction



Ever since the early days of noncommutative geometry it has become clear that this field of mathematics has close ties with physics, and with gauge theories in particular. In fact, non-abelian gauge theories, and even more prominently, the Standard Model of particle physics, were a guiding principle in the formulation of noncommutative manifolds in [1, 2].

For one thing, noncommuting operators appear naturally in quantum mechanics. As a matter of fact, there is a rather direct path from experimentally measured atomic spectra to Heisenberg's matrix mechanics which is one of the motivating examples of noncommutative geometry [3, Sect. I.1].

In the other direction, it turns out that the main technical device in noncommutative geometry, a *spectral triple*, naturally gives rise to a gauge theory. This holds in full generality, but the great potential of the noncommutative approach, at least in particle physics, becomes really visible when specific examples are considered that in fact correspond to familiar gauge theories arising in physics. This is crowned by the derivation [4] of the full Standard Model of particle physics together with all its subtleties, including the Higgs field, the spontaneous symmetry breaking mechanism, neutrino mixing, see-saw mechanism, *et cetera*.

It is the goal of this book to explore this path, and, starting with the basics, to work towards applications in particle physics, notably to the Standard Model of elementary particles.

The first ingredient of a spectral triple is an *involutive* or *\*-algebra*  $\mathcal{A}$  of operators in a Hilbert space  $\mathcal{H}$ , with the involution given by the hermitian adjoint of an operator. This immediately gives rise to a *gauge group*  $\mathfrak{G}$  determined by the unitary elements in  $\mathcal{A}$ . In general, if  $\mathcal{A}$  is noncommutative, then this group is non-abelian.

The *gauge fields* arise from a second, purely spectral data, in the guise of a *self-adjoint operator*  $D$  in  $\mathcal{H}$ , satisfying suitable conditions (cf. Definition 5.9 below). The operator  $D$  is modeled on the Dirac operator on a Riemannian spin manifold  $M$ , an elliptic first-order differential operator whose square coincides, up to a scalar term, with the Laplacian.

A key role will be played by the spectrum of  $D$ , assumed discrete; we will list its eigenvalues (with multiplicities) as  $\{\lambda_n\}_{n \in \mathbb{Z}}$ . The gauge group  $\mathfrak{G}$  acts on  $D$  by conjugation with a unitary operator,  $D \mapsto UDU^*$ . Unitarity guarantees invariance of the spectrum under such a gauge transformation.

Hence a *spectral invariant* is in particular *gauge invariant*, and it is natural to define the so-called *spectral action* as [5, 6]

$$\sum_{n \in \mathbb{Z}} f\left(\frac{\lambda_n}{\Lambda}\right).$$

Here the function  $f$  is a suitable cutoff function that makes the outcome of the sum finite, and  $\Lambda$  is a real cutoff parameter. The spectral action is interpreted as an action functional that describes the dynamics and interactions of the gauge fields constituting  $D$ .

The *fermionic fields* that are associated to a spectral triple are simply vectors  $\psi$  in the given Hilbert space, and their natural invariant is the *fermionic action*:

$$(\psi, D\psi).$$

The previous paragraphs sketch the derivation of a generalized gauge theory from any spectral triple. When one restrict to a particular class of spectral triples, this leads to ordinary gauge theory defined on a manifold  $M$  in terms of vector bundles and connections. The idea is very simple, essentially dating back to [7]: one considers the noncommutative space  $M \times F$  given by the product of  $M$  with a finite, noncommutative space  $F$ . The space  $F$  gives rise to the internal, gauge degrees of freedom. In fact, it is described by a finite-dimensional algebra of matrices, for which the gauge group becomes a matrix Lie group, such as  $SU(N)$ . The self-adjoint operator  $D_F$  is given by a hermitian matrix. Combined with the background manifold  $M$ , these objects are turned into global ones:  $\mathcal{A}$  consists of the sections of a bundle of matrix algebras, and  $D$  is a combination of  $D_F$  and the Dirac operator on  $M$  (assumed to be a Riemannian spin manifold). The operator  $D$  is found to be parametrized by gauge fields and scalar fields in suitable representations of the gauge group  $\mathfrak{G}$ . The fermionic fields  $\psi$  are sections of a spinor bundle on which  $D$  acts as a linear differential operator, minimally coupled to the gauge fields.

As we already said, the spectral action is manifestly gauge invariant, and for this latter class of examples it describes a scalar gauge theory for the group  $\mathfrak{G}$ . As a bonus, it is minimally coupled to (Euclidean) gravity, in that the gravitational degrees of freedom are present as a background field in the Dirac operator on  $M$ . Moreover, the fermionic action then gives the usual coupling of the fermionic fields to the gauge, scalar and gravitational fields.

In this respect, one of the great achievements of noncommutative geometry is the derivation of the full Standard Model of particle physics from a noncommutative space  $M \times F_{SM}$  [4]. In fact, from this geometric Ansatz one obtains the Standard Model gauge fields, the scalar Higgs field, and the full fermionic content of the Standard Model. Moreover, the spectral and fermionic action on  $M \times F_{SM}$  give the

full Lagrangian of the Standard Model, including (amongst other benefits) both the Higgs spontaneous symmetry breaking mechanism and minimal coupling to gravity. In addition, the spectral action introduces relations between the coupling constants and the masses of the Standard Model. This allows one to derive physical predictions such as the Higgs mass, and also indicates how to go beyond the Standard Model, finally bringing us back to experiment.

This book is divided into two parts. Part 1 presents the mathematical basics of noncommutative geometry and discusses the local index formula as a mathematical application. As a stand alone, it may be used as a first introduction to noncommutative geometry.

The second part starts in the same mathematical style, where in the first two chapters we analyze the structure of a gauge theory associated to any spectral triple. Comparable to a kaleidoscope, we then focus on a specific class of examples, and within this class select the physically relevant models. In the last two chapters this culminates in the derivation of the full Standard Model of particle physics. All these examples heavily exploit the results from Part 1. Hence the reader who is already somewhat familiar with noncommutative geometry, but is interested in the gauge-theoretical aspects, may want to skip Part 1 and jump immediately to the second part.

Let us quickly go through the contents of each of the chapters. Chapters 2 and 3 present a ‘light’ version of noncommutative geometry, restricting ourselves to *finite* noncommutative spaces. In other words, we here only consider finite-dimensional spectral triples and avoid technical complications that arise in the general case. Besides the pedagogical advantage, these finite spaces will in fact turn out to be crucial to the physical applications of the later chapters, where they describe the aforementioned internal space  $F$ .

Thus, in Chap. 2 we start with finite discrete topological spaces and replace them by matrix algebras. The question whether this procedure can be reversed leads naturally to the notion of Morita equivalence between matrix algebras. The next step is the translation of a metric structure into a symmetric matrix, motivating the definition of a finite spectral triple. We discuss Morita equivalence for spectral triples and conclude with a diagrammatic classification of finite spectral triples.

In Chap. 3 we enrich finite spectral triples with a real structure and discuss Morita equivalences in this context. We give a classification of finite real spectral triples based on Krajewski diagrams [8] and relate this to the classification of irreducible geometries in [9].

Chapter 4 starts with a concise background on Riemannian spin geometry, leading to a treatment of the Dirac operator, including its analytical aspects.

Chapter 5 then introduces noncommutative Riemannian spin manifolds (aka spectral triples) in full generality, exemplified by toric noncommutative manifolds.

As a first application of spectral triples, we present a proof of the local index formula of Connes and Moscovici [10] in Chap. 6, following Higson’s proof [11].

In the second part of this book we start to build gauge theories from (real) spectral triples. Chapter 7 takes a very general approach and associates a gauge group and a semigroup of gauge fields to any real spectral triple. An intriguing localization result

can be formulated (Chap. 8) in terms of a bundle of  $C^*$ -algebras on a background topological space. The gauge group acts fiberwise on this bundle and the gauge fields appear as sections thereof.

Maintaining the same level of generality, we introduce gauge invariant quantities in Chap. 9, to wit the spectral action, the topological spectral action (which is closely related to the above index), and the fermionic action [5, 6]. We discuss two possible ways to expand the spectral action, either asymptotically in terms of the cutoff  $\Lambda$ , or perturbatively in terms of the gauge fields parametrizing  $D$ .

In Chap. 10 we introduce the important class of examples alluded to before, i.e., noncommutative spaces of the form  $M \times F$  with  $F$  finite. Here, Chaps. 2 and 3 prove their value in the description of  $F$ . Following [12] we analyze the structure of the gauge group  $\mathfrak{G}_F$  for this class of examples, and determine the gauge fields and scalar fields as well as the corresponding gauge transformations. Using heat kernel methods, we obtain an asymptotic expansion for the spectral action on  $M \times F$  in terms of local formulas (on  $M$ ). We conclude that the spectral action describes the dynamics and interactions of a scalar gauge theory for the group  $\mathfrak{G}_F$ , minimally coupled to gravity. This general form of the spectral action on  $M \times F$  will be heavily used in the remainder of this book.

As a first simple example we treat abelian gauge theory in Chap. 11, for which the gauge group  $\mathfrak{G}_F \simeq U(1)$ . Following [13] we describe how to obtain the Lagrangian of electrodynamics from the spectral action.

The next step is the derivation of non-abelian Yang–Mills gauge theory from noncommutative geometry, which we discuss in Chap. 12. We obtain topologically non-trivial gauge configurations by working with algebra bundles, essentially replacing the above direct product  $M \times F$  by a fibered product [14].

Chapter 13 contains the derivation of the Standard Model of particle physics from a noncommutative manifold  $M \times F_{SM}$ , first obtained in [4]. We apply our results from Chap. 10 to obtain the Standard Model gauge group and gauge fields, and the scalar Higgs field. Moreover, the computation of the spectral action can be applied to this example and yields the full Lagrangian of the Standard Model, including Higgs spontaneous symmetry breaking and minimally coupled to gravity. We also give a detailed discussion on the fermionic action.

The phenomenology of the noncommutative Standard Model is discussed in Chap. 14. Indeed, the spectral action yields relations between the coupling constants and masses of the Standard Model, from which physical predictions can be derived. Here, we adopt the well-known renormalization group equations of the Standard Model to run the couplings to the relevant energy scale. This gives the notorious prediction for the Higgs mass at the order of 170 GeV. As this is at odds with the experiments at the Large Hadron Collider at CERN, we give a careful analysis of the hypotheses used in the derivation of the Standard Model Lagrangian from noncommutative geometry.

In Chap. 15 we use these observations to go beyond the Standard Model with non-commutative geometry. In particular, we will discuss a Pati–Salam model [15–18] that enlarges the particle content of the Standard Model. We show that this noncommutative model is compatible with the experimentally measured Higgs mass.

We end this book in Chap. 16 with an overview of recent and ongoing work searching for a quantum theory for noncommutative geometry. Indeed, as we realize the applications of conventional quantum field theory methods to the noncommutative models of Chaps. 14 and 15 cannot be the end of the story. Indeed, a more intrinsically defined quantum theory should be developed, and we indicate the first steps in this direction.

In order not to interrupt the text too much, I have chosen to collect background information and references to the literature as ‘Notes’ at the end of each chapter.

## References

1. Connes, A.: Noncommutative geometry and reality. *J. Math. Phys.* **36**(11), 6194–6231 (1995)
2. Connes, A.: Gravity coupled with matter and the foundation of non-commutative geometry. *Commun. Math. Phys.* **182**, 155–176 (1996)
3. Connes, A.: Noncommutative Geometry. Academic, San Diego (1994)
4. Chamseddine, A.H., Connes, A., Marcolli, M.: Gravity and the Standard Model with neutrino mixing. *Adv. Theor. Math. Phys.* **11**, 991–1089 (2007)
5. Chamseddine, A.H., Connes, A.: Universal formula for noncommutative geometry actions: unifications of gravity and the Standard Model. *Phys. Rev. Lett.* **77**, 4868–4871 (1996)
6. Chamseddine, A.H., Connes, A.: The spectral action principle. *Commun. Math. Phys.* **186**, 731–750 (1997)
7. Connes, A.: Essay on physics and noncommutative geometry. In: The Interface of Mathematics and Particle Physics (Oxford, 1988). Institute of Mathematics and its Applications Conference Series New Series, vol. 24 pages 9–48. Oxford University Press, New York (1990)
8. Krajewski, T.: Classification of finite spectral triples. *J. Geom. Phys.* **28**, 1–30 (1998)
9. Chamseddine, A.H., Connes, A.: Why the standard model. *J. Geom. Phys.* **58**, 38–47 (2008)
10. Connes, A., Moscovici, H.: The local index formula in noncommutative geometry. *Geom. Funct. Anal.* **5**, 174–243 (1995)
11. Higson, N.: The residue index theorem of Connes and Moscovici. In: Surveys in Noncommutative Geometry. Clay Mathematics Proceedings, vol. 6, pp. 71–126. American Mathematical Society, Providence, RI (2006)
12. van den Dungen, K., van Suijlekom, W.D.: Particle physics from almost commutative space-times. *Rev. Math. Phys.* **24**, 1230004 (2012)
13. van den Dungen, K., van Suijlekom, W.D.: Electrodynamics from noncommutative geometry. *J. Noncommut. Geom.* **7**, 433–456 (2013)
14. Boeijink, J., van Suijlekom, W.D.: The noncommutative geometry of Yang-Mills fields. *J. Geom. Phys.* **61**, 1122–1134 (2011)
15. Chamseddine, A.H., Connes, A.: Resilience of the spectral standard model. *JHEP* **1209**, 104 (2012)
16. Chamseddine, A.H., Connes, A., Van Suijlekom, W.D.: Inner fluctuations in noncommutative geometry without the first order condition. *J. Geom. Phys.* **73**, 222–234 (2013)
17. Chamseddine, A.H., Connes, A., van Suijlekom, W.D.: Beyond the spectral standard model: emergence of Pati-Salam unification. *JHEP* **1311**, 132 (2013)
18. Chamseddine, A.H., Connes, A., van Suijlekom, W.D.: Grand unification in the spectral Pati-Salam model. *JHEP* **11**, 011 (2015)

**Open Access** This chapter is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



**Part I**

**Noncommutative Geometric Spaces**

# Chapter 2

## Finite Noncommutative Spaces



In this chapter (and the next) we consider finite discrete topological spaces. However, we will stretch their usual definition, which is perhaps geometrically not so interesting, to include the more intriguing finite *noncommutative* spaces. Intuitively, this means that each point has some internal structure, described by a particular noncommutative algebra. With such a notion of finite noncommutative spaces, we search for the appropriate notion of maps between, and (geo)metric structure on such spaces, and arrive at a diagrammatic classification of such finite noncommutative geometric spaces. Our exposition of the finite case already gives a good first impression of what noncommutative geometry has in store, whilst having the advantage that it avoids technical complications that might obscure such a first tour through noncommutative geometry. The general case is subsequently treated in Chap. 5.

### 2.1 Finite Spaces and Matrix Algebras

Consider a finite topological space  $X$  consisting of  $N$  points (equipped with the discrete topology):

$$1 \bullet \quad 2 \bullet \quad \dots \dots \quad N \bullet$$

The first step towards a noncommutative geometrical description is to trade spaces for their corresponding function algebras.

**Definition 2.1** A (complex, unital) *algebra* is a vector space  $A$  (over  $\mathbb{C}$ ) with a bilinear associative product  $A \times A \rightarrow A$  denoted by  $(a, b) \mapsto ab$  (and a unit 1 satisfying  $1a = a1 = a$  for all  $a \in A$ ).

A  *$*$ -algebra* (or, *involutive algebra*) is an algebra  $A$  together with a conjugate-linear map (the *involution*)  $* : A \rightarrow A$  such that  $(ab)^* = b^*a^*$  and  $(a^*)^* = a$  for all  $a, b \in A$ .

In this book, we restrict to unital algebras, and simply refer to them as algebras.

In the present case, we consider the  $*$ -algebra  $C(X)$  of  $\mathbb{C}$ -valued functions on the above finite space  $X$ . It is equipped with a pointwise linear structure,

$$(f + g)(x) = f(x) + g(x), \quad (\lambda f)(x) = \lambda(f(x)),$$

for any  $f, g \in C(X)$ ,  $\lambda \in \mathbb{C}$  and for any point  $x \in X$ , and with pointwise multiplication

$$fg(x) = f(x)g(x).$$

There is an involution given by complex conjugation at each point:

$$f^*(x) = \overline{f(x)}.$$

The  $C$  in  $C(X)$  stands for continuous and, indeed, any  $\mathbb{C}$ -valued function on a finite space  $X$  with the discrete topology is automatically continuous.

The  $*$ -algebra  $C(X)$  has a rather simple structure: it is isomorphic to the  $*$ -algebra  $\mathbb{C}^N$  with each complex entry labeling the value the function takes at the corresponding point, with the involution given by complex conjugation of each entry. A convenient way to encode the algebra  $C(X) \simeq \mathbb{C}^N$  is in terms of diagonal  $N \times N$  matrices, representing a function  $f : X \rightarrow \mathbb{C}$  as

$$f \rightsquigarrow \begin{pmatrix} f(1) & 0 & \cdots & 0 \\ 0 & f(2) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & f(N) \end{pmatrix}.$$

Hence, pointwise multiplication then simply becomes matrix multiplication, and the involution is given by hermitian conjugation.


If  $\phi : X_1 \rightarrow X_2$  is a map of finite discrete spaces, then there is a corresponding map from  $C(X_2) \rightarrow C(X_1)$  given by pullback:

$$\phi^* f = f \circ \phi \in C(X_1); \quad (f \in C(X_2)).$$

Note that the pullback  $\phi^*$  is a  $*$ -homomorphism (or,  $*$ -algebra map) under the pointwise product, in that

$$\phi^*(fg) = \phi^*(f)\phi^*(g), \quad \phi^*(\overline{f}) = \overline{\phi^*(f)}, \quad \phi^*(\lambda f + g) = \lambda\phi^*(f) + \phi^*(g).$$

For example, let  $X_1$  be the space consisting of three points, and  $X_2$  the space consisting of two points. If a map  $\phi : X_1 \rightarrow X_2$  is defined according to the following diagram,



then

$$\phi^* : \mathbb{C}^2 \simeq C(X_2) \rightarrow \mathbb{C}^3 \simeq C(X_1)$$

is given by

$$(\lambda_1, \lambda_2) \mapsto (\lambda_1, \lambda_2, \lambda_2).$$

**Exercise 2.1** Show that  $\phi : X_1 \rightarrow X_2$  is an injective (surjective) map of finite spaces if and only if  $\phi^* : C(X_2) \rightarrow C(X_1)$  is surjective (injective).

**Definition 2.2** A (complex) *matrix algebra*  $A$  is a direct sum

$$A = \bigoplus_{i=1}^N M_{n_i}(\mathbb{C}),$$

for some positive integers  $n_i$  and  $N$ . The involution on  $A$  is given by hermitian conjugation, and we simply refer to the  $*$ -algebra  $A$  with this involution as a matrix algebra.

Hence, we have associated a matrix algebra  $C(X)$  to the finite space  $X$ , which behaves naturally with respect to maps between topological spaces and  $*$ -algebras. A natural question is whether this procedure can be inverted. In other words, given a matrix algebra  $A$ , can we obtain a finite discrete space  $X$  such that  $A \simeq C(X)$ ? Since  $C(X)$  is always commutative but matrix algebras need not be, we quickly arrive at the conclusion that the answer is negative. This can be resolved in two ways:

- (1) Restrict to *commutative* matrix algebras.
- (2) Allow for more morphisms (and consequently, more isomorphisms) between matrix algebras, e.g., by generalizing  $*$ -homomorphisms.

Before explaining each of these options, let us introduce some useful definitions concerning representations of finite-dimensional  $*$ -algebras (which are not necessarily commutative) which moreover extend in a straightforward manner to the infinite-dimensional case (cf. Definitions 5.5 and 5.6). We first need the prototypical example of a  $*$ -algebra.

**Example 2.3** Let  $H$  be an (finite-dimensional) inner product space, with inner product  $(\cdot, \cdot) \rightarrow \mathbb{C}$ . We denote by  $L(H)$  the  $*$ -algebra of operators on  $H$  with product

given by composition and the involution is given by mapping an operator  $T$  to its adjoint  $T^*$ .

Note that  $L(H)$  is a normed vector space: for  $T \in L(H)$  we set

$$\|T\|^2 = \sup_{h \in H} \{(Th, Th) : (h, h) \leq 1\}.$$

Equivalently,  $\|T\|$  is given by the square root of the largest eigenvalue of  $T^*T$ .

**Definition 2.4** A *representation* of a finite-dimensional  $*$ -algebra  $A$  is a pair  $(H, \pi)$  where  $H$  is a (finite-dimensional, complex) inner product space and  $\pi$  is a  $*$ -algebra map

$$\pi : A \rightarrow L(H).$$

A representation  $(H, \pi)$  is called irreducible if  $H \neq 0$  and the only subspaces in  $H$  that are left invariant under the action of  $A$  are  $\{0\}$  or  $H$ .

We will also refer to a finite-dimensional inner product space as a *finite-dimensional Hilbert space*.

**Example 2.5** Consider  $A = M_n(\mathbb{C})$ . The defining representation is given by  $H = \mathbb{C}^n$  on which  $A$  acts by left matrix multiplication; hence it is irreducible. An example of a reducible representation is  $H = \mathbb{C}^n \oplus \mathbb{C}^n$ , with  $a \in M_n(\mathbb{C})$  acting in block-form:

$$a \in M_n(\mathbb{C}) \mapsto \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \in L(\mathbb{C}^n \oplus \mathbb{C}^n) \simeq M_{2n}(\mathbb{C})$$

which therefore decomposes as the direct sum of two copies of the defining representation. See also Lemma 2.15 below.

**Exercise 2.2** Given a representation  $(H, \pi)$  of a  $*$ -algebra  $A$ , the **commutant**  $\pi(A)'$  of  $\pi(A)$  is defined as

$$\pi(A)' = \{T \in L(H) : \pi(a)T = T\pi(a) \text{ for all } a \in A\}.$$

- (1) Show that  $\pi(A)'$  is also a  $*$ -algebra.
- (2) Show that a representation  $(H, \pi)$  of  $A$  is irreducible if and only if the commutant  $\pi(A)'$  of  $\pi(A)$  consists of multiples of the identity.

**Definition 2.6** Two representations  $(H_1, \pi_1)$  and  $(H_2, \pi_2)$  of a  $*$ -algebra  $A$  are *unitarily equivalent* if there exists a unitary map  $U : H_1 \rightarrow H_2$  such that

$$\pi_1(a) = U^*\pi_2(a)U.$$

**Definition 2.7** The *structure space*  $\widehat{A}$  of  $A$  is the set of all unitary equivalence classes of irreducible representations of  $A$ .

We end this subsection with an illustrative exercise on passing from representations of a  $*$ -algebra to matrices over that  $*$ -algebra.

- Exercise 2.3** (1) If  $A$  is a unital  $*$ -algebra, show that the  $n \times n$ -matrices  $M_n(A)$  with entries in  $A$  form a unital  $*$ -algebra.  
 (2) Let  $\pi : A \rightarrow L(H)$  be a representation of a  $*$ -algebra  $A$  and set  $H^n = H \oplus \cdots \oplus H$  ( $n$  copies). Show that the following defines a representation  $\tilde{\pi} : M_n(A) \rightarrow L(H^n)$  of  $M_n(A)$ :

$$\tilde{\pi}((a_{ij})) = (\pi(a_{ij})); \quad ((a_{ij}) \in M_n(A)).$$

- (3) Let  $\tilde{\pi} : M_n(A) \rightarrow L(H^n)$  be a representation of the  $*$ -algebra  $M_n(A)$ . Show that the following defines a representation  $\pi : A \rightarrow L(H^n)$  of the  $*$ -algebra  $A$ :

$$\pi(a) = \tilde{\pi}(a\mathbb{I}_n)$$

where  $\mathbb{I}_n$  is the identity in  $M_n(A)$ .

### 2.1.1 Commutative Matrix Algebras

We now explain how option (1) on Sect. 2.1 above resolves the question raised by constructing a space from a commutative matrix algebra  $A$ . A natural candidate for such a space is, of course, the structure space  $\widehat{A}$ , which we now determine. Note that any commutative matrix algebra is of the form  $A \simeq \mathbb{C}^N$ , for which by Exercise 2.2(2) any irreducible representation is given by a map of the form

$$\pi_i : (\lambda_1, \dots, \lambda_N) \in \mathbb{C}^N \mapsto \lambda_i \in \mathbb{C}$$

for some  $i = 1, \dots, N$ . We conclude that  $\widehat{A} \simeq \{1, \dots, N\}$ .

We conclude that there is a *duality* between finite spaces and commutative matrix algebras. This is nothing but a finite-dimensional version of *Gelfand duality* (see Theorem 5.7 below) between compact Hausdorff topological spaces and unital commutative  $C^*$ -algebras. In fact, we will see later (Proposition 5.4) that any finite-dimensional  $C^*$ -algebra is a matrix algebra, which reduces Gelfand duality to the present finite-dimensional duality.

### 2.1.2 Finite Spaces and Matrix Algebras

The above trade of finite discrete spaces for finite-dimensional commutative  $*$ -algebras does not really make them any more interesting, for the  $*$ -algebra is always of the form  $\mathbb{C}^N$ . A more interesting perspective is given by the noncommutative alternative, viz. option (2) below Definition 2.2 in Sect. 2.1. We thus aim for

a duality between finite spaces and *equivalence classes* of matrix algebras. These equivalence classes are described by a generalized notion of isomorphisms between matrix algebras, also known as Morita equivalence.

Let us first recall the notion of an algebra (bi)module.

**Definition 2.8** Let  $A, B$  be algebras (not necessarily matrix algebras). A left  $A$ -module is a vector space  $E$  that carries a left representation of  $A$ , i.e., there is a bilinear map  $A \times E \ni (a, e) \mapsto a \cdot e \in E$  such that

$$(a_1 a_2) \cdot e = a_1 \cdot (a_2 \cdot e); \quad (a_1, a_2 \in A, e \in E).$$

Similarly, a right  $B$ -module is a vector space  $F$  that carries a right representation of  $B$ , i.e., there is a bilinear map  $F \times B \ni (f, b) \mapsto f \cdot b \in F$  such that

$$f \cdot (b_1 b_2) = (f \cdot b_1) \cdot b_2; \quad (b_1, b_2 \in B, f \in F).$$

Finally, an  $A - B$ -bimodule  $E$  is both a left  $A$ -module and a right  $B$ -module, with mutually commuting actions:

$$a \cdot (e \cdot b) = (a \cdot e) \cdot b; \quad (a \in A, b \in B, e \in E).$$

When no confusion can arise, we will also write  $ae$  instead of  $a \cdot e$  to denote the left module action, and similarly for the right action.

There is a natural notion of (left)  **$A$ -module homomorphism** as a linear map  $\phi : E \rightarrow F$  that respect the representation of  $A$ :

$$\phi(a \cdot e) = a \cdot \phi(e); \quad (a \in A, e \in E).$$

Similarly for right modules and bimodules.

We introduce the following notation:

- $_A E$  for a left  $A$ -module  $E$ ;
- $F_B$  for a right  $B$ -module  $F$ ;
- $_A E_B$  for an  $A - B$ -bimodule  $E$ .

**Exercise 2.4** Check that a representation  $\pi : A \rightarrow L(H)$  of a  $*$ -algebra  $A$  (cf. Definition 2.4) turns  $H$  into a left  $A$ -module  $_A H$ .

**Exercise 2.5** Show that  $A$  is itself an  $A - A$ -bimodule  $_A A_A$ , with left and right actions given by the product in  $A$ .

If  $E$  is a right  $A$ -module, and  $F$  is a left  $A$ -module, we can form the *balanced tensor product*:

$$E \otimes_A F := E \otimes F / \left\{ \sum_i e_i a_i \otimes f_i - e_i \otimes a_i f_i : \quad a_i \in A, e_i \in E, f_i \in F \right\}.$$

In other words, the quotient imposes  $A$ -linearity of the tensor product, i.e., in  $E \otimes_A F$  we have

$$ea \otimes_A f = e \otimes_A af; \quad (a \in A, e \in E, f \in F).$$

**Definition 2.9** Let  $A, B$  be matrix algebras. A *Hilbert bimodule* for the pair  $(A, B)$  is given by an  $A - B$ -bimodule  $E$  together with a  $B$ -valued inner product  $\langle \cdot, \cdot \rangle_E : E \times E \rightarrow B$  satisfying

$$\begin{aligned} \langle e_1, a \cdot e_2 \rangle_E &= \langle a^* \cdot e_1, e_2 \rangle_E; \quad (e_1, e_2 \in E, a \in A), \\ \langle e_1, e_2 \cdot b \rangle_E &= \langle e_1, e_2 \rangle_E b; \quad \langle e_1, e_2 \rangle_E^* = \langle e_2, e_1 \rangle_E; \quad (e_1, e_2 \in E, b \in B), \\ \langle e, e \rangle_E &\geq 0 \text{ with equality if and only if } e = 0; \quad (e \in E). \end{aligned}$$

The set of Hilbert bimodules for  $(A, B)$  will be denoted by  $\mathbf{KK}_f(A, B)$ .

In the following, we will also write  $\langle \cdot, \cdot \rangle$  instead of  $\langle \cdot, \cdot \rangle_E$ , unless confusion might arise.

**Exercise 2.6** Check that a representation  $\pi : A \rightarrow L(H)$  (cf. Definition 2.4 and Exercise 2.4) of a matrix algebra  $A$  turns  $H$  into a Hilbert bimodule for  $(A, \mathbb{C})$ .

**Exercise 2.7** Show that the  $A - A$ -bimodule given by  $A$  itself (cf. Exercise 2.5) is an element in  $\mathbf{KK}_f(A, A)$  by establishing that the following formula defines an  $A$ -valued inner product  $\langle \cdot, \cdot \rangle_A : A \times A \rightarrow A$ :

$$\langle a, a' \rangle_A = a^* a'; \quad (a, a' \in A).$$

**Example 2.10** More generally, let  $\phi : A \rightarrow B$  be a  $*$ -algebra homomorphism between matrix algebras  $A$  and  $B$ . From it, we can construct a Hilbert bimodule  $E_\phi$  in  $\mathbf{KK}_f(A, B)$  as follows. Let  $E_\phi$  be  $B$  as a vector space with the natural right  $B$ -module structure and inner product (cf. Exercise 2.7), but with  $A$  acting on the left via the homomorphism  $\phi$ :

$$a \cdot b = \phi(a)b; \quad (a \in A, b \in E_\phi).$$

**Definition 2.11** The *Kasparov product*  $F \circ E$  between Hilbert bimodules  $E \in \mathbf{KK}_f(A, B)$  and  $F \in \mathbf{KK}_f(B, C)$  is given by the balanced tensor product

$$F \circ E := E \otimes_B F; \quad (E \in \mathbf{KK}_f(A, B), F \in \mathbf{KK}_f(B, C)),$$

so that  $F \circ E \in \mathbf{KK}_f(A, C)$ , with  $C$ -valued inner product given on elementary tensors by

$$\langle e_1 \otimes f_1, e_2 \otimes f_2 \rangle_{E \otimes_B F} = \langle f_1, \langle e_1, e_2 \rangle_E f_2 \rangle_F, \tag{2.1.1}$$

and extended linearly to all of  $E \otimes F$ .

Note that this product is associative up to isomorphism.

**Exercise 2.8** Show that the association  $\phi \rightsquigarrow E_\phi$  from Example 2.10 is *natural* in the sense that

- (1)  $E_{\text{id}_A} \simeq A \in \mathbf{KK}_f(A, A)$ ,
- (2) for  $*$ -algebra homomorphisms  $\phi : A \rightarrow B$  and  $\psi : B \rightarrow C$  we have an isomorphism

$$E_\psi \circ E_\phi \equiv E_\phi \otimes_B E_\psi \simeq E_{\psi \circ \phi} \in \mathbf{KK}_f(A, C),$$

that is, as  $A - C$ -bimodules.

**Exercise 2.9** In the above definition:

- (1) Check that  $E \otimes_B F$  is an  $A - C$ -bimodule.
- (2) Check that  $\langle \cdot, \cdot \rangle_{E \otimes_B F}$  defines a  $C$ -valued inner product.
- (3) Check that  $\langle a^*(e_1 \otimes f_1), e_2 \otimes f_2 \rangle_{E \otimes_B F} = \langle e_1 \otimes f_1, a(e_2 \otimes f_2) \rangle_{E \otimes_B F}$ .

Conclude that  $F \circ E$  is indeed an element of  $\mathbf{KK}_f(A, C)$ .

Let us consider the Kasparov product with the Hilbert bimodule for  $(A, A)$  given by  $A$  itself (cf. Exercise 2.7). Then, since for  $E \in \mathbf{KK}_f(A, B)$  we have  $E \circ A = A \otimes_A E \simeq E$ , the bimodule  ${}_A A_A$  is the identity element with respect to the Kasparov product (up to isomorphism). This motivates the following definition.

**Definition 2.12** Two matrix algebras  $A$  and  $B$  are called *Morita equivalent* if there exist elements  $E \in \mathbf{KK}_f(A, B)$  and  $F \in \mathbf{KK}_f(B, A)$  such that

$$E \otimes_B F \simeq A, \quad F \otimes_A E \simeq B,$$

where  $\simeq$  denotes isomorphism as Hilbert bimodules.

If  $A$  and  $B$  are Morita equivalent, then the representation theories of both matrix algebras are equivalent. More precisely, if  $A$  and  $B$  are Morita equivalent, then a right  $A$ -module is sent to a right  $B$ -module by tensoring with  $\_ \otimes_A E$  for an invertible element  $E$  in  $\mathbf{KK}_f(A, B)$ .

**Example 2.13** As seen in Exercises 2.4 and 2.6, the vector space  $E = \mathbb{C}^n$  is an  $M_n(\mathbb{C}) - \mathbb{C}$ -bimodule; with the standard  $\mathbb{C}$ -valued inner product it becomes a Hilbert module for  $(M_n(\mathbb{C}), \mathbb{C})$ . Similarly, the vector space  $F = \mathbb{C}^n$  is a  $\mathbb{C} - M_n(\mathbb{C})$ -bimodule by right matrix multiplication. An  $M_n(\mathbb{C})$ -valued inner product is given by

$$\langle v_1, v_2 \rangle = \bar{v}_1 v_2^t \in M_n(\mathbb{C}).$$

We determine the Kasparov products of these Hilbert bimodules as

$$E \otimes_{\mathbb{C}} F \simeq M_n(\mathbb{C}); \quad F \otimes_{M_n(\mathbb{C})} E \simeq \mathbb{C}.$$

In other words,  $E \in \mathbf{KK}_f(M_n(\mathbb{C}), \mathbb{C})$  and  $F \in \mathbf{KK}_f(\mathbb{C}, M_n(\mathbb{C}))$  are each other's inverse with respect to the Kasparov product. We conclude that  $M_n(\mathbb{C})$  and  $\mathbb{C}$  are Morita equivalent.

This observation leads us to our first little result.

**Theorem 2.14** *Two matrix algebras are Morita equivalent if and only if their structure spaces are isomorphic as finite discrete spaces, i.e., have the same cardinality.*

**Proof** Let  $A$  and  $B$  be Morita equivalent. Thus there exists Hilbert bimodules  ${}_A E_B$  and  ${}_B F_A$  such that

$$E \otimes_B F \simeq A, \quad F \otimes_A E \simeq B.$$

If  $[(\pi_B, H)] \in \widehat{B}$  then we can define a representation  $\pi_A$  by setting

$$\pi_A : A \rightarrow L(E \otimes_B H); \quad \pi_A(a)(e \otimes v) = ae \otimes v. \quad (2.1.2)$$

Vice versa, we construct  $\pi_B : B \rightarrow L(F \otimes_A W)$  from  $[(\pi_A, W)] \in \widehat{A}$  by setting  $\pi_B(b)(f \otimes w) = bf \otimes w$  and these two maps are one another's inverse. Thus,  $\widehat{A} \simeq \widehat{B}$  (see Exercise 2.10 below).

For the converse, we start with a basic result on irreducible representations of  $M_n(\mathbb{C})$ .

**Lemma 2.15** *The matrix algebra  $M_n(\mathbb{C})$  has a unique irreducible representation (up to isomorphism) given by the defining representation on  $\mathbb{C}^n$ .*

**Proof** It is clear from Exercise 2.2 that  $\mathbb{C}^n$  is an irreducible representation of  $A = M_n(\mathbb{C})$ . Suppose  $H$  is irreducible and of dimension  $K$ , and define a linear map

$$\phi : \underbrace{A \oplus \cdots \oplus A}_{K \text{ copies}} \rightarrow H^*; \quad \phi(a_1, \dots, a_K) \rightarrow e^1 \circ a_1^t + \cdots + e^K \circ a_K^t$$

in terms of a basis  $\{e^1, \dots, e^K\}$  of the dual vector space  $H^*$ . Here  $v \circ a$  denotes pre-composition of  $v \in H^*$  with  $a \in A$ , acting on  $H$ . This is a morphism of  $M_n(\mathbb{C})$ -modules, provided a matrix  $a$  acts on the dual vector space  $H^*$  by sending  $v \mapsto v \circ a^t$ . It is also surjective, so that the dual map  $\phi^* : H \rightarrow (A^K)^*$  is injective. Upon identifying  $(A^K)^*$  with  $A^K$  as  $A$ -modules, and noting that  $A = M_n(\mathbb{C}) \simeq \bigoplus^n \mathbb{C}^n$  as  $A$ -modules, it follows that  $H$  is a submodule of  $A^K \simeq \bigoplus^{nK} \mathbb{C}^n$ . By irreducibility  $H \simeq \mathbb{C}^n$ .  $\square$

Now, if  $A, B$  are matrix algebras of the following form

$$A = \bigoplus_{i=1}^N M_{n_i}(\mathbb{C}), \quad B = \bigoplus_{j=1}^M M_{m_j}(\mathbb{C}),$$

then  $\widehat{A} \simeq \widehat{B}$  implies that  $N = M$ . Then, define

$$E := \bigoplus_{i=1}^N \mathbb{C}^{n_i} \otimes \mathbb{C}^{m_i},$$

with  $A$  acting by block-diagonal matrices on the first tensor and  $B$  acting in a similar way by right matrix multiplication on the second leg of the tensor product. Also, set

$$F := \bigoplus_{i=1}^N \mathbb{C}^{m_i} \otimes \mathbb{C}^{n_i},$$

with  $B$  now acting on the left and  $A$  on the right. Then, as above,

$$\begin{aligned} E \otimes_B F &\simeq \bigoplus_{i=1}^N (\mathbb{C}^{n_i} \otimes \mathbb{C}^{m_i}) \otimes_{M_{m_i}(\mathbb{C})} (\mathbb{C}^{m_i} \otimes \mathbb{C}^{n_i}) \\ &\simeq \bigoplus_{i=1}^N \mathbb{C}^{n_i} \otimes (\mathbb{C}^{m_i} \otimes_{M_{m_i}(\mathbb{C})} \mathbb{C}^{m_i}) \otimes \mathbb{C}^{n_i} \\ &\simeq \bigoplus_{i=1}^N \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_i} \simeq A, \end{aligned}$$

and similarly we obtain  $F \otimes_A E \simeq B$ , as required.  $\square$

**Exercise 2.10** Fill in the gaps in the above proof:

- (a) Show that the representation  $\pi_A$  defined by (2.1.2) is irreducible if and only if  $\pi_B$  is.
- (b) Show that the association of the class  $[\pi_A]$  to  $[\pi_B]$  through (2.1.2) is independent of the choice of representatives  $\pi_A$  and  $\pi_B$ .


We conclude that there is a duality between finite spaces and Morita equivalence classes of matrix algebras. By replacing  $*$ -homomorphisms  $A \rightarrow B$  by Hilbert bimodules for  $(A, B)$ , we introduce a much richer structure at the level of morphisms between matrix algebras. For example, any finite-dimensional inner product space defines an element in  $\mathrm{KK}_f(\mathbb{C}, \mathbb{C})$ , whereas there is only one map from the corresponding structure space consisting of one point to itself. When combined with Exercise 2.10 we conclude that Hilbert bimodules form a proper extension of the  $*$ -morphisms between matrix algebras.

## 2.2 Noncommutative Geometric Finite Spaces

Consider again a finite space  $X$ , described as the structure space of a matrix algebra  $A$ . We would like to introduce some geometry on  $X$  and, in particular, a notion of a metric on  $X$ .

Thus, the question we want to address is how we can (algebraically) describe distances between the points in  $X$ , say, as embedded in a metric space. Recall that a metric on a finite discrete space  $X$  is given by an array  $\{d_{ij}\}_{i,j \in X}$  of real non-negative

entries, indexed by a pair of elements in  $X$  and requiring that  $d_{ij} = d_{ji}$ ,  $d_{ij} \leq d_{ik} + d_{kj}$ , and  $d_{ij} = 0$  if and only if  $i = j$ :



**Example 2.16** If  $X$  is embedded in a metric space (e.g., Euclidean space), it can be equipped with the induced metric.

**Example 2.17** The **discrete metric** on the discrete space  $X$  is given by:

$$d_{ij} = \begin{cases} 1 & \text{if } i \neq j \\ 0 & \text{if } i = j. \end{cases}$$

In the commutative case, we have the following remarkable result, which completely characterizes the metric on  $X$  in terms of linear algebraic data. It is the key result towards a *spectral* description of finite geometric spaces.

**Theorem 2.18** Let  $d_{ij}$  be a metric on the space  $X$  of  $N$  points, and set  $A = \mathbb{C}^N$  with elements  $a = (a(i))_{i=1}^N$ , so that  $\widehat{A} \simeq X$ . Then there exists a representation  $\pi$  of  $A$  on a finite-dimensional inner product space  $H$  and a symmetric operator  $D$  on  $H$  such that

$$d_{ij} = \sup_{a \in A} \{|a(i) - a(j)| : \| [D, \pi(a)] \| \leq 1\}. \quad (2.2.1)$$

**Proof** We claim that this would follow from the equality

$$\| [D, \pi(a)] \| = \max_{k \neq l} \left\{ \frac{1}{d_{kl}} |a(k) - a(l)| \right\}. \quad (*)$$

Indeed, if this holds, then

$$\sup_a \{|a(i) - a(j)| : \| [D, a] \| \leq 1\} \leq d_{ij}.$$

The reverse inequality follows by taking  $a \in A$  for fixed  $i, j$  to be  $a(k) = d_{ik}$ . Then, we find  $|a(i) - a(j)| = d_{ij}$ , while  $\| [D, \pi(a)] \| \leq 1$  for this  $a$  follows from the reverse triangle inequality for  $d_{ij}$ :

$$\frac{1}{d_{kl}} |a(k) - a(l)| = \frac{1}{d_{kl}} |d_{ik} - d_{il}| \leq 1.$$

We prove  $(*)$  by induction on  $N$ . If  $N = 2$ , then on  $H = \mathbb{C}^2$  we define a representation  $\pi : A \rightarrow L(H)$  and a hermitian matrix  $D$  by

$$\pi(a) = \begin{pmatrix} a(1) & 0 \\ 0 & a(2) \end{pmatrix}, \quad D = \begin{pmatrix} 0 & (d_{12})^{-1} \\ (d_{12})^{-1} & 0 \end{pmatrix}.$$

It follows that  $\|[D, a]\| = (d_{12})^{-1}|a(1) - a(2)|$ .

Suppose then that  $(*)$  holds for  $N$ , with representation  $\pi_N$  of  $\mathbb{C}^N$  on an inner product space  $H_N$  and symmetric operator  $D_N$ ; we will show that it also holds for  $N + 1$ . We define

$$H_{N+1} = H_N \oplus \bigoplus_{i=1}^N H_N^i$$

with  $H_N^i := \mathbb{C}^2$ . Imitating the above construction in the case  $N = 2$ , we define the representation  $\pi_{N+1}$  by

$$\begin{aligned} \pi_{N+1}(a(1), \dots, a(N+1)) &= \pi_N(a(1), \dots, a(N)) \\ &\oplus \begin{pmatrix} a(1) & 0 \\ 0 & a(N+1) \end{pmatrix} \oplus \dots \oplus \begin{pmatrix} a(N) & 0 \\ 0 & a(N+1) \end{pmatrix}, \end{aligned}$$

and define the operator  $D_{N+1}$  by

$$\begin{aligned} D_{N+1} &= D_N \oplus \begin{pmatrix} 0 & (d_{1(N+1)})^{-1} \\ (d_{1(N+1)})^{-1} & 0 \end{pmatrix} \\ &\oplus \dots \oplus \begin{pmatrix} 0 & (d_{N(N+1)})^{-1} \\ (d_{N(N+1)})^{-1} & 0 \end{pmatrix}. \end{aligned}$$

It follows by the induction hypothesis that  $(*)$  holds for  $N + 1$ .  $\square$

**Exercise 2.11** Make the above proof explicit for the case  $N = 3$ . In other words, compute the metric of (2.2.1) on the space of three points from the set of data  $A = \mathbb{C}^3$ ,  $H = (\mathbb{C}^2)^{\oplus 3}$  with representation  $\pi : A \rightarrow L(H)$  given by

$$\pi(a(1), a(2), a(3)) = \begin{pmatrix} a(1) & 0 \\ 0 & a(2) \end{pmatrix} \oplus \begin{pmatrix} a(1) & 0 \\ 0 & a(3) \end{pmatrix} \oplus \begin{pmatrix} a(2) & 0 \\ 0 & a(3) \end{pmatrix},$$

and hermitian matrix

$$D = \begin{pmatrix} 0 & x_1 \\ x_1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & x_2 \\ x_2 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & x_3 \\ x_3 & 0 \end{pmatrix},$$

with  $x_1, x_2, x_3 \in \mathbb{R}$ .

**Exercise 2.12** Compute the metric on the space of three points given by formula (2.2.1) for the set of data  $A = \mathbb{C}^3$  acting in the defining representation on  $H = \mathbb{C}^3$ , and

$$D = \begin{pmatrix} 0 & d^{-1} & 0 \\ d^{-1} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

for some non-zero  $d \in \mathbb{R}$ .

Even though the above translation of the metric on  $X$  into algebraic data assumes commutativity of  $A$ , the distance formula itself can be extended to the case of a noncommutative matrix algebra  $A$ . In fact, suppose we are given a  $*$ -algebra representation of  $A$  on an inner product space, together with a symmetric operator  $D$  on  $H$ . Then we can define a metric on the structure space  $\widehat{A}$  by

$$d_{ij} = \sup_{a \in A} \{ |\operatorname{Tr} a(i) - \operatorname{Tr} a(j)| : \| [D, a] \| \leq 1 \}, \quad (2.2.2)$$

where  $i$  labels the matrix algebra  $M_{n_i}(\mathbb{C})$  in the decomposition of  $A$ . This distance formula is a special case of Connes' distance formula (see Note 5 on Chap. 5) on the structure space of  $A$ .

**Exercise 2.13** Show that the  $d_{ij}$  in (2.2.2) is a metric (actually, an **extended metric**, taking values in  $[0, \infty]$ ) on  $\widehat{A}$  by establishing that

$$d_{ij} = 0 \iff i = j, \quad d_{ij} = d_{ji}, \quad d_{ij} \leq d_{ik} + d_{kj}.$$

This suggests that the above structure consisting of a matrix algebra  $A$ , a finite-dimensional representation space  $H$ , and a hermitian matrix  $D$  provides the data needed to capture a metric structure on the finite space  $X = \widehat{A}$ . In fact, in the case that  $A$  is commutative, the above argument combined with our finite-dimensional Gelfand duality of Sect. 2.1.1 is a reconstruction theorem. Indeed, we reconstruct a given metric space  $(X, d)$  from the data  $(A, H, D)$  associated to it.

We arrive at the following definition , adapted to our finite-dimensional setting.

**Definition 2.19** A finite spectral triple is a triple  $(A, H, D)$  consisting of a unital  $*$ -algebra  $A$  represented faithfully on a finite-dimensional Hilbert space  $H$ , together with a symmetric operator  $D : H \rightarrow H$ .

We do not demand that  $A$  is a matrix algebra, since this turns out to be automatic:

**Lemma 2.20** *If  $A$  is a unital  $*$ -algebra that acts faithfully on a finite-dimensional Hilbert space, then  $A$  is a matrix algebra of the form*

$$A \simeq \bigoplus_{i=1}^N M_{n_i}(\mathbb{C}).$$

**Proof** Since  $A$  acts faithfully on a Hilbert space it is a  $*$ -subalgebra of a matrix algebra  $L(H) = M_{\dim(H)}(\mathbb{C})$ ; the only such subalgebras are themselves matrix algebras.  $\square$

Unless we want to distinguish different representations of  $A$  on  $H$ , the above representation will usually be implicitly assumed, thus considering elements  $a \in A$  as operators on  $H$ .

**Example 2.21** Let  $A = M_n(\mathbb{C})$  act on  $H = \mathbb{C}^n$  by matrix multiplication, with the standard inner product. A symmetric operator on  $H$  is represented by a hermitian  $n \times n$  matrix.

We will loosely refer to  $D$  as a **finite Dirac operator**, as its infinite-dimensional analogue on Riemannian spin manifolds is the usual Dirac operator (see Chap. 5). In the present case, we can use it to introduce a ‘differential geometric structure’ on the finite space  $X$  that is related to the notion of **divided difference**. The latter is given, for each pair of points  $i, j \in X$ , by

$$\frac{a(i) - a(j)}{d_{ij}}.$$

Indeed, these divided differences appear precisely as the entries of the commutator  $[D, a]$  for the operator  $D$  as in Theorem 2.18.

**Exercise 2.14** Use the explicit form of  $D$  in Theorem 2.18 to confirm that the commutator of  $D$  with  $a \in C(X)$  is expressed in terms of the above divided differences.

We will see later that in the continuum case, the commutator  $[D, \cdot]$  corresponds to taking derivatives of functions on a manifold.

**Definition 2.22** Let  $(A, H, D)$  be a finite spectral triple. The  $A$ -bimodule of *Connes’ differential one-forms* is given by

$$\Omega_D^1(A) := \left\{ \sum_k a_k [D, b_k] : a_k, b_k \in A \right\}.$$

Consequently, there is a map  $d : A \rightarrow \Omega_D^1(A)$ , given by  $d(\cdot) = [D, \cdot]$ .

**Exercise 2.15** Verify that  $d$  is a derivation of a  $*$ -algebra, in that:

$$d(ab) = d(a)b + ad(b); \quad d(a^*) = -d(a)^*.$$

**Exercise 2.16** Verify that  $\Omega_D^1(A)$  is an  $A$ -bimodule by rewriting the operator  $\sum_k a(a_k [D, b_k])b$  ( $a, b, a_k, b_k \in A$ ) as  $\sum_k a'_k [D, b'_k]$  for some  $a'_k, b'_k \in A$ .

As a first little result—though with an actual application to matrix models in physics—we compute Connes’ differential one-forms for the above Example 2.21.

**Lemma 2.23** Let  $(A, H, D) = (M_n(\mathbb{C}), \mathbb{C}^n, D)$  be the finite spectral triple of Example 2.21 with  $D$  a hermitian  $n \times n$  matrix. If  $D$  is not a multiple of the identity, then  $\Omega_D^1(A) \simeq M_n(\mathbb{C})$ .

**Proof** We may assume that  $D$  is a diagonal matrix:  $D = \sum_i \lambda_i e_{ii}$  in terms of real numbers  $\lambda_i$  (not all equal) and the standard basis  $\{e_{ij}\}$  of  $M_n(\mathbb{C})$ . For fixed  $i, j$  choose  $k$  such that  $\lambda_k \neq \lambda_j$ . Then

$$\left( \frac{1}{\lambda_k - \lambda_j} e_{ik} \right) [D, e_{kj}] = e_{ij}.$$

Hence, since  $e_{ik}, e_{kj} \in M_n(\mathbb{C})$ , any basis vector  $e_{ij} \in \Omega_D^1(A)$ . Since also  $\Omega_D^1(A) \subset L(\mathbb{C}^n) \simeq M_n(\mathbb{C})$ , the result follows.  $\square$

**Exercise 2.17** Consider the following finite spectral triple:

$$\left( A = \mathbb{C}^2, H = \mathbb{C}^2, D = \begin{pmatrix} 0 & \lambda \\ \bar{\lambda} & 0 \end{pmatrix} \right),$$

with  $\lambda \neq 0$ . Show that the corresponding space of differential one-forms  $\Omega_D^1(A)$  is isomorphic to the vector space of all off-diagonal  $2 \times 2$  matrices.

### 2.2.1 Morphisms Between Finite Spectral Triples

In a spectral triple  $(A, H, D)$  both the  $*$ -algebra  $A$  and a finite Dirac operator  $D$  act on the inner product space  $H$ . Hence, the most natural notion of equivalence between spectral triples is that of unitary equivalence.

**Definition 2.24** Two finite spectral triples  $(A_1, H_1, D_1)$  and  $(A_2, H_2, D_2)$  are called unitarily equivalent if  $A_1 = A_2$  and if there exists a unitary operator  $U : H_1 \rightarrow H_2$  such that

$$\begin{aligned} U\pi_1(a)U^* &= \pi_2(a); & (a \in A_1), \\ UD_1U^* &= D_2. \end{aligned}$$

**Exercise 2.18** Show that unitary equivalence of spectral triples is an equivalence relation.

**Remark 2.25** A special type of unitary equivalence is given by the unitaries in the matrix algebra  $A$  itself. Indeed, for any such unitary element  $u$  the spectral triples  $(A, H, D)$  and  $(A, H, uDu^*)$  are unitarily equivalent. Another way of writing  $uDu^*$  is  $D + u[D, u^*]$ , so that this type of unitary equivalence effectively adds a differential one-form to  $D$ .

Following the spirit of our extended notion of morphisms between algebras, we might also deduce a notion of “equivalence” coming from Morita equivalence of the corresponding matrix algebras. Namely, given a Hilbert bimodule  $E$  in  $\mathbf{KK}_f(B, A)$ , we can try to construct a finite spectral triple on  $B$  starting from a finite spectral triple

on  $A$ . This transfer of metric structure is accomplished as follows. Let  $(A, H, D)$  be a spectral triple; we construct a new spectral triple  $(B, H', D')$ . First, we define a vector space

$$H' = E \otimes_A H,$$

which inherits a left action of  $B$  from the  $B$ -module structure of  $E$ . Also, it is an inner product space, with  $\mathbb{C}$ -valued inner product given as in (2.1.1).

The naive choice of a symmetric operator  $D'$  given by  $D'(e \otimes \xi) = e \otimes D\xi$  will not do, because it does not respect the ideal defining the balanced tensor product over  $A$ , being generated by elements of the form

$$ea \otimes \xi - e \otimes a\xi; \quad (e \in E, a \in A, \xi \in H).$$

A better definition is

$$D'(e \otimes \xi) = e \otimes D\xi + \nabla(e)\xi, \quad (2.2.3)$$

where  $\nabla : E \rightarrow E \otimes_A \Omega_D^1(A)$  is some map that satisfies the **Leibniz rule**

$$\nabla(ea) = \nabla(e)a + e \otimes [D, a]; \quad (e \in E, a \in A). \quad (2.2.4)$$

Indeed, this is precisely the property that is needed to make  $D'$  a well-defined operator on the balanced tensor product  $E \otimes_A H$ :

$$D'(ea \otimes \xi - e \otimes a\xi) = ea \otimes D\xi + \nabla(ea)\xi - e \otimes D(a\xi) - \nabla(e)a\xi = 0.$$

A map  $\nabla : E \rightarrow E \otimes_A \Omega_D^1(A)$  that satisfies Eq. (2.2.4) is called a **connection** on the right  $A$ -module  $E$  associated to the derivation  $d : a \mapsto [D, a]$  ( $a \in A$ ).

**Theorem 2.26** *If  $(A, H, D)$  is a finite spectral triple and  $E \in \mathbf{KK}_f(B, A)$ , then (in the above notation)  $(B, E \otimes_A H, D')$  is a finite spectral triple, provided that  $\nabla$  satisfies the compatibility condition*

$$\langle e_1, \nabla e_2 \rangle_E - \langle \nabla e_1, e_2 \rangle_E = d\langle e_1, e_2 \rangle_E; \quad (e_1, e_2 \in E). \quad (2.2.5)$$

**Proof** We only need to show that  $D'$  is a symmetric operator. Indeed, for  $e_1, e_2 \in E$  and  $\xi_1, \xi_2 \in H$  we compute

$$\begin{aligned} \langle e_1 \otimes \xi_1, D'(e_2 \otimes \xi_2) \rangle_{E \otimes_A H} &= \langle \xi_1, \langle e_1, \nabla e_2 \rangle_E \xi_2 \rangle_H + \langle \xi_1, \langle e_1, e_2 \rangle_E D\xi_2 \rangle_H \\ &= \langle \xi_1, \langle \nabla e_1, e_2 \rangle_E \xi_2 \rangle_H + \langle \xi_1, d\langle e_1, e_2 \rangle_E \xi_2 \rangle_H \\ &\quad + \langle D\xi_1, \langle e_1, e_2 \rangle_E \xi_2 \rangle_H - \langle \xi_1, [D, \langle e_1, e_2 \rangle_E] \xi_2 \rangle_H \\ &= \langle D'(e_1 \otimes \xi_1), e_2 \otimes \xi_2 \rangle_{E \otimes_A H}, \end{aligned}$$

using the stated compatibility condition and the fact that  $D$  is symmetric.  $\square$

Theorem 2.26 is our finite-dimensional analogue of Theorem 7.15, to be obtained below.

**Exercise 2.19** Let  $\nabla$  and  $\nabla'$  be two connections on a right  $A$ -module  $E$ . Show that their difference  $\nabla - \nabla'$  is a right  $A$ -linear map  $E \rightarrow E \otimes_A \Omega_D^1(A)$ .

**Exercise 2.20** In this exercise, we consider the case that  $B = A$  and also  $E = A$ . Let  $(A, H, D)$  be a spectral triple, we determine  $(A, H', D')$ .

- (1) Show that the derivation  $d(\cdot) = [D, \cdot] : A \rightarrow A \otimes_A \Omega_D^1(A) = \Omega_D^1(A)$  is a connection on  $A$  considered a right  $A$ -module.
- (2) Upon identifying  $A \otimes_A H \simeq H$ , what is the operator  $D'$  of Eq. (2.2.3) when the connection  $\nabla$  on  $A$  is given by  $d$  as in (1)?
- (3) Use (1) and (2) of this exercise to show that any connection  $\nabla : A \rightarrow A \otimes_A \Omega_D^1(A)$  is given by

$$\nabla = d + \omega,$$

with  $\omega \in \Omega_D^1(A)$ .

- (4) Upon identifying  $A \otimes_A H \simeq H$ , what is the operator  $D'$  of Eq. (2.2.3) with the connection on  $A$  given as  $\nabla = d + \omega$ .

If we combine the above Exercise 2.20 with Lemma 2.23, we see that  $\nabla = d - D$  is an example of a connection on  $M_N(\mathbb{C})$  (as a module over itself and with  $\omega = -D$ ), since  $\Omega_D^1(A) \simeq M_N(\mathbb{C})$ . Hence, for this choice of connection the new finite spectral triple as constructed in Theorem 2.26 is given by  $(M_N(\mathbb{C}), \mathbb{C}^N, D' = 0)$ . So, Morita equivalence of algebras does not carry over to an equivalence relation on spectral triples. Indeed, we now have  $\Omega_{D'}^1(M_N(\mathbb{C})) = 0$ , so that no non-zero  $D$  can be generated from this spectral triple and the symmetry of this relation fails.

## 2.3 Classification of Finite Spectral Triples

Here we classify finite spectral triples on  $A$  modulo unitary equivalence, in terms of so-called **decorated graphs**.

**Definition 2.27** A *graph* is an ordered pair  $(\Gamma^{(0)}, \Gamma^{(1)})$  consisting of a set  $\Gamma^{(0)}$  of *vertices* and a set  $\Gamma^{(1)}$  of pairs of vertices (called *edges*).

We allow edges of the form  $e = (v, v)$  for any vertex  $v$ , that is, we allow loops at any vertex.

Consider then a finite spectral triple  $(A, H, D)$ ; let us determine the structure of all three ingredients and construct a graph from it.

**The algebra:** We have already seen in Lemma 2.20 that

$$A \simeq \bigoplus_{i=1}^N M_{n_i}(\mathbb{C}),$$

for some  $n_1, \dots, n_N$ . The structure space of  $A$  is given by  $\widehat{A} \simeq \{1, \dots, N\}$  with each integer  $i \in \widehat{A}$  corresponding to the equivalence classes of the representation of  $A$  on  $\mathbb{C}^{n_i}$ . If we label the latter equivalence class by  $\mathbf{n}_i$  we can also identify  $\widehat{A} \simeq \{\mathbf{n}_1, \dots, \mathbf{n}_N\}$ .

**The Hilbert space:** Any finite-dimensional faithful representation  $H$  of such a matrix algebra  $A$  is completely reducible (i.e., a direct sum of irreducible representations).

**Exercise 2.21** Prove this result for any  $*$ -algebra by establishing that the complement  $W^\perp$  of an  $A$ -submodule  $W \subset H$  is also an  $A$ -submodule of  $H$ .

Combining this with the proof of Lemma 2.15, we conclude that the finite-dimensional Hilbert space representation  $H$  of  $A$  has a decomposition into irreducible representations, which we write as

$$H \simeq \bigoplus_{i=1}^N \mathbb{C}^{n_i} \otimes V_i,$$

with each  $V_i$  a vector space; we will refer to the dimension of  $V_i$  as the **multiplicity** of the representation labeled by  $\mathbf{n}_i$  and to  $V_i$  itself as the **multiplicity space**. The above isomorphism is given by a unitary map.

To begin the construction of our decorated graph, we indicate the presence of a summand  $\mathbf{n}_i$  in  $H$  by drawing a node at position  $\mathbf{n}_i \in \widehat{A}$  in a diagram based on the structure space  $\widehat{A}$  of the matrix algebra  $A$  (see Fig. 2.1 for an example). Multiple nodes at the same position represent multiplicities of the representations in  $H$ .

**The finite Dirac operator:** Corresponding to the above decomposition of  $H$  we can write  $D$  as a sum of matrices

$$D_{ij} : \mathbb{C}^{n_i} \otimes V_i \rightarrow \mathbb{C}^{n_j} \otimes V_j,$$

restricted to these subspaces. The condition that  $D$  is symmetric implies that  $D_{ij} = D_{ji}^*$ . In terms of the above diagrammatic representation of  $H$ , we express a non-zero  $D_{ij}$  and  $D_{ji}$  as a (multiple) edge between the nodes  $\mathbf{n}_i$  and  $\mathbf{n}_j$  (see Fig. 2.2 for an example).

Another way of putting this is as follows, in terms of decorated graphs.

**Definition 2.28** A  $\Lambda$ -decorated graph is given by an ordered pair  $(\Gamma, \Lambda)$  of a finite graph  $\Gamma$  and a finite set  $\Lambda$  of positive integers, with a labeling:

$$\begin{array}{ccccccc} \mathbf{n}_1 & \cdots & \mathbf{n}_i & \cdots & \mathbf{n}_j & \cdots & \mathbf{n}_N \\ \circ & & \circ & & \circ & & \circ \end{array}$$

**Fig. 2.1** A node at  $\mathbf{n}_i$  indicates the presence of the summand  $\mathbb{C}^{n_i}$ ; the double node at  $\mathbf{n}_j$  indicates the presence of the summand  $\mathbb{C}^{n_j} \oplus \mathbb{C}^{n_j}$  in  $H$



**Fig. 2.2** The edges between the nodes  $\mathbf{n}_i$  and  $\mathbf{n}_j$ , and  $\mathbf{n}_i$  and  $\mathbf{n}_N$  represent non-zero operators  $D_{ij} : \mathbb{C}^{n_i} \rightarrow \mathbb{C}^{n_j} \otimes \mathbb{C}^2$  (multiplicity 2) and  $D_{iN} : \mathbb{C}^{n_i} \rightarrow \mathbb{C}^{n_N}$ , respectively. Their adjoints give the operators  $D_{ji}$  and  $D_{Ni}$

- of the vertices  $v \in \Gamma^{(0)}$  by elements  $n(v) \in \Lambda$ ;
- of the edges  $e = (v_1, v_2) \in \Gamma^{(1)}$  by operators  $D_e : \mathbb{C}^{n(v_1)} \rightarrow \mathbb{C}^{n(v_2)}$  and its conjugate-transpose  $D_e^* : \mathbb{C}^{n(v_2)} \rightarrow \mathbb{C}^{n(v_1)}$ ,

so that  $n(\Gamma^{(0)}) = \Lambda$ .

The operators  $D_e$  between vertices that are labeled by  $\mathbf{n}_i$  and  $\mathbf{n}_j$ , respectively, add up to the above  $D_{ij}$ . Explicitly,


$$D_{ij} = \sum_{\substack{e=(v_1, v_2) \\ n(v_1)=\mathbf{n}_i \\ n(v_2)=\mathbf{n}_j}} D_e,$$

so that also  $D_{ij}^* = D_{ji}$ . Thus we have proved the following result.

**Theorem 2.29** *There is a one-to-one correspondence between finite spectral triples modulo unitary equivalence and  $\Lambda$ -decorated graphs, given by associating a finite spectral triple  $(A, H, D)$  to a  $\Lambda$ -decorated graph  $(\Gamma, \Lambda)$  in the following way:*

$$A = \bigoplus_{n \in \Lambda} M_n(\mathbb{C}), \quad H = \bigoplus_{v \in \Gamma^{(0)}} \mathbb{C}^{n(v)}, \quad D = \sum_{e \in \Gamma^{(1)}} D_e + D_e^*.$$

**Example 2.30** The following  $\Lambda$ -decorated graph



corresponds to the spectral triple  $(M_n(\mathbb{C}), \mathbb{C}^n, D = D_e + D_e^*)$  of Example 2.21.

**Exercise 2.22** Draw the  $\Lambda$ -decorated graph corresponding to the spectral triple

$$\left( A = \mathbb{C}^3, H = \mathbb{C}^3, D = \begin{pmatrix} 0 & \lambda & 0 \\ \bar{\lambda} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right); \quad (\lambda \neq 0).$$

**Exercise 2.23** Use  $\Lambda$ -decorated graphs to classify all finite spectral triples (modulo unitary equivalence) on the matrix algebra  $A = \mathbb{C} \oplus M_2(\mathbb{C})$ .

**Exercise 2.24** Suppose that  $(A_1, H_1, D_1)$  and  $(A_2, H_2, D_2)$  are two finite spectral triples. We consider their direct sum and tensor product and give the corresponding  $\Lambda$ -decorated graphs.

- (1) Show that  $(A_1 \oplus A_2, H_1 \oplus H_2, (D_1, D_2))$  is a finite spectral triple.
- (2) Describe the  $\Lambda$ -decorated graph of this *direct sum spectral triple* in terms of the  $\Lambda$ -decorated graphs of the original spectral triples.
- (3) Show that  $(A_1 \otimes A_2, H_1 \otimes H_2, D_1 \otimes 1 + 1 \otimes D_2)$  is a finite spectral triple.
- (4) Describe the  $\Lambda$ -decorated graph of this *tensor product spectral triple* in terms of the  $\Lambda$ -decorated graphs of the original spectral triples.

## Notes

### Section 2.1. Finite Spaces and Matrix Algebras

1. The notation  $\text{KK}_f$  in Definition 2.9 is chosen to suggest a close connection to Kasparov's bivariant KK-theory [1], here restricted to the finite-dimensional case. In fact, in the case of matrix algebras the notion of a *Kasparov module* for a pair of  $C^*$ -algebras  $(A, B)$  (cf. [2, Sect. 17.1] for a definition) coincides (up to homotopy) with that of a Hilbert bimodule for  $(A, B)$  (cf. [3, Sect. IV.2.1] for a definition).
2. Definition 2.12 agrees with the notion of equivalence between arbitrary rings introduced by Morita [4]. Moreover, it is a special case of strong Morita equivalence between  $C^*$ -algebras as introduced by Rieffel [5].
3. Theorem 2.14 is a special case of a more general result on the structure spaces of Morita equivalent  $C^*$ -algebras (see e.g., [6, Sect. 3.3]).

### Section 2.2. Noncommutative Geometric Finite Spaces

4. Theorem 2.18 can be found in [7].
5. The reconstruction theorem mentioned in the text before Definition 2.19 is a special case, to wit the finite-dimensional case, of a result by Connes [8] on a reconstruction of Riemannian (spin) manifolds from so-called spectral triples (cf. Definition 5.9 and Note 6 on Chap. 5 below).
6. A complete proof of Lemma 2.20 can be found in [9, Theorem 3.5.4].
7. For a complete exposition on differential algebras, connections on modules, *et cetera*, we refer to [10, Chap. 8] and [11] and references therein.
8. The failure of Morita equivalence to induce an equivalence between spectral triples was noted in [12, Remark 1.143] (see also [13, Remark 5.1.2]). This suggests that it is better to consider Hilbert bimodules as *correspondences* rather than equivalences, as was already suggested by Connes and Skandalis in [14] and also appeared in the applications of noncommutative geometry to number theory (cf. [12, Chap. 4.3]) and quantization [15]. This forms the starting point for a categorical description of (finite) spectral triples themselves. As objects the category has finite spectral triples  $(A, H, D)$ , and as morphisms it has pairs  $(E, \nabla)$  as above. This category is the topic of for instance [16–21], working in the more general setting of spectral triples, hence requiring much more analysis as compared to our finite-dimensional case. The category of finite spectral triples plays a crucial role in the noncommutative generalization of spin networks in [22].

## References

1. Kasparov, G.G.: The operator  $K$ -functor and extensions of  $C^*$ -algebras. Izv. Akad. Nauk SSSR Ser. Mat. **44**, 571–636, 719 (1980)
2. Blackadar, B.: K-Theory for Operator Algebras. Mathematical Sciences Research Institute Publications. Cambridge University Press (1998)
3. Landsman, N.P.: Mathematical Topics between Classical and Quantum Mechanics. Springer, New York (1998)
4. Morita, K.: Duality for modules and its applications to the theory of rings with minimum condition. Sci. Rep. Tokyo Kyoiku Daigaku Sect. A **6**, 83–142 (1958)
5. Rieffel, M.A.: Morita equivalence for  $C^*$ -algebras and  $W^*$ -algebras. J. Pure Appl. Algebra **5**, 51–96 (1974)
6. Raeburn, I., Williams, D.P.: Morita equivalence and continuous-trace  $C^*$ -algebras. Mathematical Surveys and Monographs, vol. 60. American Mathematical Society, Providence, RI (1998)
7. Iochum, B., Krajewski, T., Martinetti, P.: Distances in finite spaces from noncommutative geometry. J. Geom. Phys. **37**, 100–125 (2001)
8. Connes, A.: On the spectral characterization of manifolds. J. Noncommut. Geom. **7**, 1–82 (2013)
9. Etingof, P., Golberg, O., Hensel, S., Liu, T., Schwendner, A., Vaintrob, D., Yudovina, E.: Introduction to representation theory. Student Mathematical Library, vol. 59. American Mathematical Society, Providence, RI, 2011. With historical interludes by Slava Gerovitch
10. Landi, G.: An Introduction to Noncommutative Spaces and their Geometry. Springer (1997)
11. André, Y.: Différentielles non commutatives et théorie de Galois différentielle ou aux différences. Ann. Sci. École Norm. Sup. (4) **34**, 685–739 (2001)
12. Connes, A., Marcolli, M.: Noncommutative Geometry. Quantum Fields and Motives. AMS, Providence (2008)
13. Venselaar, J.: Classification and equivalences of noncommutative tori and quantum lens spaces. Ph.D. thesis, Utrecht University (2012)
14. Connes, A., Skandalis, G.: The longitudinal index theorem for foliations. Publ. Res. Inst. Math. Sci. **20**, 1139–1183 (1984)
15. Landsman, N.P.: Quantized reduction as a tensor product. In: Quantization of Singular Symplectic Quotients. Progress in Mathematics, vol. 198, pp. 137–180. Birkhäuser, Basel (2001)
16. Mesland, B.: Bivariant  $K$ -theory of groupoids and the noncommutative geometry of limit sets. Ph.D. thesis, Universität Bonn (2009)
17. Mesland, B.: Unbounded bivariant  $K$ -theory and correspondences in noncommutative geometry. J. Reine Angew. Math. **691**, 101–172 (2014)
18. Kaad, J., Lesch, M.: A local global principle for regular operators in Hilbert  $C^*$ -modules. J. Funct. Anal. **262**, 4540–4569 (2012)
19. Kaad, J., Lesch, M.: Spectral flow and the unbounded Kasparov product. Adv. Math. **248**, 495–530 (2013)
20. Mesland, B., Rennie, A.: Nonunital spectral triples and metric completeness in unbounded  $KK$ -theory. J. Funct. Anal. **271**, 2460–2538 (2016)
21. van den Dungen, K., Mesland, B.: Homotopy equivalence in unbounded  $KK$ -theory. Ann. K-Theory **5**, 501–537 (2020)
22. Marcolli, M., van Suijlekom, W.D.: Gauge networks in noncommutative geometry. J. Geom. Phys. **75**, 71–91 (2014)

**Open Access** This chapter is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 3

## Finite Real Noncommutative Spaces



In this chapter, we will enrich the finite noncommutative spaces as analyzed in the previous chapter with a *real structure*. For one thing, this makes the definition of a finite spectral triple more symmetric by demanding the inner product space  $H$  be an  $A - A$ -bimodule, rather than just a left  $A$ -module. The implementation of this bimodule structure by an anti-unitary operator has close ties with the Tomita-Takesaki theory of Von Neumann algebras, as well as with physics through charge conjugation, as will become clear in the applications in the later chapters of this book. Our exposition includes a diagrammatic classification of finite real spectral triples for all so-called KO-dimensions, and also identifies the irreducible finite geometries among them.

### 3.1 Finite Real Spectral Triples

First, the structure of a finite spectral triple can be enriched by introducing a  $\mathbb{Z}_2$ -grading  $\gamma$  on  $H$ , i.e.,  $\gamma^* = \gamma$ ,  $\gamma^2 = 1$ , demanding that  $A$  is *even* and  $D$  is *odd* with respect to this grading:

$$\gamma D = -D\gamma, \quad \gamma a = a\gamma; \quad (a \in A).$$

Next, there is a more symmetric refinement of the notion of finite spectral triple in which  $H$  is an  $A - A$ -bimodule, rather than just a left  $A$ -module. Recall that an anti-unitary operator is an invertible operator  $J : H \rightarrow H$  that satisfies  $\langle J\xi_1, J\xi_2 \rangle = \langle \xi_2, \xi_1 \rangle$  for all  $\xi_1, \xi_2 \in H$ .

**Definition 3.1** A *finite real spectral triple* is given by a finite spectral triple  $(A, H, D)$  and an anti-unitary operator  $J : H \rightarrow H$  called *real structure*, such that  $a^\circ := Ja^*J^{-1}$  is a right representation of  $A$  on  $H$ , i.e.,  $(ab)^\circ = b^\circ a^\circ$ . We also require that

$$[a, b^\circ] = 0, \quad [[D, a], b^\circ] = 0, \tag{3.1.1}$$

**Table 3.1** The KO-dimension  $k$  of a real spectral triple is determined by the signs  $\{\epsilon, \epsilon', \epsilon''\}$  appearing in  $J^2 = \epsilon$ ,  $JD = \epsilon'DJ$  and  $J\gamma = \epsilon''\gamma J$

$k$	0	1	2	3	4	5	6	7
$\epsilon$	1	1	-1	-1	-1	-1	1	1
$\epsilon'$	1	-1	1	1	1	-1	1	1
$\epsilon''$	1		-1		1		-1	

for all  $a, b \in A$ . Moreover, we demand that  $J$ ,  $D$  and (in the even case)  $\gamma$  satisfy the commutation relations:

$$J^2 = \epsilon, \quad JD = \epsilon'DJ, \quad J\gamma = \epsilon''\gamma J.$$

for numbers  $\epsilon, \epsilon', \epsilon'' \in \{-1, 1\}$ . These signs determine the *KO-dimension*  $k$  (modulo 8) of the finite real spectral triple  $(A, H, D; J, \gamma)$  defined according to Table 3.1.

The signs in Table 3.1 are motivated by the classification of Clifford algebras, see Sect. 4.1 below. The two conditions in (3.1.1) are called the **commutant property**, and the **first-order or order one condition**, respectively. They imply that the left action of an element in  $A$  and  $\Omega_D^1(A)$  commutes with the right action of  $A$ . This is equivalent to the commutation between the right action of  $A$  and  $\Omega_D^1(A)$  with the left action of  $A$ .

**Remark 3.2** The so-called *opposite algebra*  $A^\circ$  is defined to be equal to  $A$  as a vector space but with opposite product  $\circ$ :

$$a \circ b := ba.$$

Thus,  $a^\circ = Ja^*J^{-1}$  defines a left representation of  $A^\circ$  on  $H$ :  $(a \circ b)^\circ = a^\circ b^\circ$ .

**Example 3.3** Consider the matrix algebra  $M_N(\mathbb{C})$ , acting on the inner product space  $H = M_N(\mathbb{C})$  by left matrix multiplication, and with inner product given by the Hilbert–Schmidt inner product:

$$\langle a, b \rangle = \text{Tr } a^* b.$$

Define

$$\gamma(a) = a, \quad J(a) = a^*; \quad (a \in H).$$

Since  $D$  must be odd with respect to the grading  $\gamma$ , it vanishes identically.

**Exercise 3.1** In the previous example, show that the right action of  $M_N(\mathbb{C})$  on  $H = M_N(\mathbb{C})$  as defined by  $a \mapsto a^\circ$  is given by right matrix multiplication.

The following exercises are inspired by Tomita–Takesaki theory of Von Neumann algebras.

**Exercise 3.2** Let  $A = \bigoplus_i M_{n_i}(\mathbb{C})$  be a matrix algebra, which is represented on a vector space  $H = \bigoplus_i \mathbb{C}^{n_i} \otimes \mathbb{C}^{m_i}$ , i.e., is such that the irreducible representation  $\mathbf{n}_i$  has multiplicity  $m_i$ .

- (1) Show that the commutant  $A'$  of  $A$  is isomorphic to  $\bigoplus_i M_{m_i}(\mathbb{C})$ . As a consequence, the double commutant coincides with  $A$ , that is to say  $A'' \cong A$ .

We say that  $\xi \in H$  is a **cyclic vector** for  $A$  if

$$A\xi := \{a\xi : a \in A\} = H.$$

We call  $\xi \in H$  a **separating vector** for  $A$  if

$$a\xi = 0 \implies a = 0; \quad (a \in A).$$

- (2) Show that if  $\xi$  is a separating vector for the action of  $A$ , it is cyclic for the action of  $A'$ . (Hint: Assume  $\xi$  is not cyclic for the action of  $A'$  and try to derive a contradiction).

**Exercise 3.3** Suppose that  $(A, H, D = 0)$  is a finite spectral triple such that  $H$  possesses a cyclic and separating vector  $\xi$  for  $A$ .

- (1) Show that the formula  $S(a\xi) = a^*\xi$  defines an anti-linear operator  $S : H \rightarrow H$ .
- (2) Show that  $S$  is invertible.
- (3) Let  $J : H \rightarrow H$  be the operator appearing in the polar decomposition  $S = J\Delta^{1/2}$  of  $S$  with  $\Delta = S^*S$ . Show that  $J$  is an anti-unitary operator.

Conclude that  $(A, H, D = 0; J)$  is a finite real spectral triple. Can you find such an operator  $J$  in the case of Exercise 3.2?

### 3.1.1 Morphisms Between Finite Real Spectral Triples

We are now going to extend the notion of unitary equivalence (cf. Definition 2.24) to finite real spectral triples.

**Definition 3.4** We call two finite real spectral triples  $(A_1, H_1, D_1; J_1, \gamma_1)$  and  $(A_2, H_2, D_2; J_2, \gamma_2)$  *unitarily equivalent* if  $A_1 = A_2$  and if there exists a unitary operator  $U : H_1 \rightarrow H_2$  such that

$$\begin{aligned} U\pi_1(a)U^* &= \pi_2(a); \quad (a \in A_1), \\ UD_1U^* &= D_2, \quad U\gamma_1U^* = \gamma_2, \quad UJ_1U^* = J_2. \end{aligned}$$

Building on our discussion in Sect. 2.2.1, we can also extend Morita equivalence to finite real spectral triples. Namely, given a Hilbert bimodule  $E$  for  $(B, A)$ , we will construct a finite real spectral triple  $(B, H', D'; J', \gamma')$  on  $B$ , starting from a finite real spectral triple  $(A, H, D; J, \gamma)$  on  $A$ .

**Definition 3.5** Let  $E$  be a  $B - A$ -bimodule. The *conjugate module*  $E^\circ$  is given by the  $A - B$ -bimodule

$$E^\circ = \{\bar{e} : e \in E\},$$

with  $a \cdot \bar{e} \cdot b = \overline{b^* \cdot e \cdot a^*}$  for any  $a \in A, b \in B$ .

This implies for any  $\lambda \in \mathbb{C}$  that  $\bar{\lambda e} = \bar{\lambda} \bar{e}$ , which explains the suggestive notation  $\bar{e}$  for the elements of  $E^\circ$ . The bimodule  $E^\circ$  is not quite a Hilbert bimodule for  $(A, B)$ , since we do not have a natural  $B$ -valued inner product. However, there is a  $A$ -valued inner product on the left  $A$ -module  $E^\circ$  given by

$$\langle \bar{e}_1, \bar{e}_2 \rangle = \langle e_2, e_1 \rangle; \quad (e_1, e_2 \in E).$$

As opposed to the inner product in Definition 2.9, this inner product is left  $A$ -linear:  $\langle \bar{e}_1, a\bar{e}_2 \rangle = a \langle \bar{e}_1, \bar{e}_2 \rangle$  for all  $a \in A$ , as can be easily checked.

**Exercise 3.4** Show that  $E^\circ$  is a Hilbert bimodule for  $(B^\circ, A^\circ)$ .

Let us then start the construction of a finite real spectral triple on  $B$  by setting

$$H' := E \otimes_A H \otimes_A E^\circ.$$

There is a ( $\mathbb{C}$ -valued) inner product on  $H'$  given by combining the  $A$ -valued inner products on  $E, E^\circ$  with the  $\mathbb{C}$ -valued inner product on  $H$ , much as in (2.1.1). The action of  $B$  on  $H'$  is given by

$$b(e_1 \otimes \xi \otimes \bar{e}_2) = (be_1) \otimes \xi \otimes \bar{e}_2, \tag{3.1.2}$$

using just the  $B - A$ -bimodule structure of  $E$ . In addition, there is a right action of  $B$  on  $H'$  defined by acting on the right on the component  $E^\circ$ . In fact, it is implemented by the following anti-unitary,

$$J'(e_1 \otimes \xi \otimes \bar{e}_2) = e_2 \otimes J\xi \otimes \bar{e}_1,$$

i.e.,  $b^\circ = J'b^*(J')^{-1}$  with  $b^* \in B$  acting on  $H'$  according to (3.1.2).

Moreover, there is a finite Dirac operator given in terms of the connection  $\nabla : E \rightarrow E \otimes_A \Omega_D^1(A)$  as in Sect. 2.2.1. First, we need the result of the following exercise.

**Exercise 3.5** Let  $\nabla : E \rightarrow E \otimes_A \Omega_D^1(A)$  be a right connection on  $E$  and consider the following anti-linear map

$$\begin{aligned} \tau : E \otimes_A \Omega_D^1(A) &\rightarrow \Omega_D^1(A) \otimes_A E^\circ; \\ e \otimes \omega &\mapsto -\omega^* \otimes \bar{e}. \end{aligned}$$

Show that the map  $\bar{\nabla} : E^\circ \rightarrow \Omega_D^1(A) \otimes_A E^\circ$  defined by  $\bar{\nabla}(\bar{e}) = \tau \circ \nabla(e)$  is a left connection, i.e., show that it satisfies the left Leibniz rule:

$$\bar{\nabla}(a\bar{e}) = [D, a] \otimes \bar{e} + a\bar{\nabla}(\bar{e}).$$

The connections  $\nabla$  and  $\bar{\nabla}$  give rise to a Dirac operator on  $E \otimes_A H \otimes_A E^\circ$ :

$$D'(e_1 \otimes \xi \otimes \bar{e}_2) = (\nabla e_1)\xi \otimes \bar{e}_2 + e_1 \otimes D\xi \otimes \bar{e}_2 + e_1 \otimes \xi(\bar{\nabla}\bar{e}_2).$$

The right action of  $\omega \in \Omega_D^1(A)$  on  $\xi \in H$  is then defined by  $\xi \mapsto \epsilon' J \omega^* J^{-1} \xi$ .

Finally, for even spectral triples one defines a grading on  $E \otimes_A H \otimes_A E^\circ$  by  $\gamma' = 1 \otimes \gamma \otimes 1$ .

**Theorem 3.6** Suppose  $(A, H, D; J, \gamma)$  is a finite real spectral triple of KO-dimension  $k$ , and let  $\nabla : E \rightarrow E \otimes_A \Omega_D^1(A)$  be a compatible connection (cf. Eq.(2.2.5)). Then  $(B, H', D'; J', \gamma')$  is a finite real spectral triple of KO-dimension  $k$ .

**Proof** The only non-trivial thing to check is that the KO-dimension is preserved. In fact, one readily checks that  $(J')^2 = 1 \otimes J^2 \otimes 1 = \epsilon$  and  $J'\gamma' = \epsilon''\gamma'J'$ . Also,

$$\begin{aligned} J'D'(e_1 \otimes \xi \otimes \bar{e}_2) &= J'((\nabla e_1)\xi \otimes \bar{e}_2 + e_1 \otimes D\xi \otimes \bar{e}_2 + e_1 \otimes \xi(\tau\nabla e_2)) \\ &= \epsilon'D'(e_2 \otimes J\xi \otimes \bar{e}_1) \equiv \epsilon'D'J'(e_1 \otimes \xi \otimes \bar{e}_2), \end{aligned}$$

where we have used  $J'(e_1 \otimes J\omega J^{-1}\xi \otimes \bar{e}_2) = e_2 \otimes \omega J\xi \otimes \bar{e}_1$ .  $\square$

## 3.2 Classification of Finite Real Spectral Triples

In this section, we classify all finite real spectral triples  $(A, H, D; J, \gamma)$  modulo unitary equivalence using **Krajewski diagrams**. These play a similar role for finite real spectral triples as Dynkin diagrams do for simple Lie algebras. Moreover, they extend our  $\Lambda$ -decorated graphs of the previous chapter to the case of real spectral triples.

**The algebra:** First, we already know from our classification of finite spectral triples in Sect. 2.3 that

$$A \simeq \bigoplus_{i=1}^N M_{n_i}(\mathbb{C}),$$

for some  $n_1, \dots, n_N$ . Thus, the structure space of  $A$  is again given by  $\widehat{A} = \{\mathbf{n}_1, \dots, \mathbf{n}_N\}$  where  $\mathbf{n}_i$  denotes the irreducible representation of  $A$  on  $\mathbb{C}^{n_i}$ .

**The Hilbert space:** As before, the irreducible, faithful representations of  $A = \bigoplus_{i=1}^N M_{n_i}(\mathbb{C})$  are given by corresponding direct sums:

$$\bigoplus_{i=1}^N \mathbb{C}^{n_i}$$

on which  $A$  acts by left block-diagonal matrix multiplication.

Now, besides the representation of  $A$ , there should also be a representation of  $A^\circ$  on  $H$  which commutes with that of  $A$ . In other words, we are looking for the irreducible

representations of  $A \otimes A^\circ$ . If we denote the unique irreducible representation of  $M_n(\mathbb{C})^\circ$  by  $\mathbb{C}^{n^\circ}$ , this implies that any irreducible representation of  $A \otimes A^\circ$  is given by a summand in

$$\bigoplus_{i,j=1}^N \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ}.$$


Consequently, any finite-dimensional Hilbert space representation of  $A$  has a decomposition into irreducible representations

$$H = \bigoplus_{i,j=1}^N \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij},$$

with  $V_{ij}$  a vector space; we will refer to the dimension of  $V_{ij}$  as the **multiplicity** of the representation  $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ}$ .

The integers  $\mathbf{n}_i$  and  $\mathbf{n}_j^\circ$  form the grid of a diagram (cf. Fig. 3.1 for an example). Whenever there is a node at the coordinates  $(\mathbf{n}_i, \mathbf{n}_j^\circ)$ , the representation  $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ}$  is present in the direct sum decomposition of  $H$ . Multiplicities are indicated by multiple nodes.

**Example 3.7** Consider the algebra  $A = \mathbb{C} \oplus M_2(\mathbb{C})$ . The irreducible representations of  $A$  are given by **1** and **2**. The two diagrams



**Fig. 3.1** A node at  $(\mathbf{n}_i, \mathbf{n}_j^\circ)$  indicates the presence of the summand  $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ}$  in  $H$ ; the double node indicates the presence of  $(\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_i \circ}) \oplus (\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_i \circ})$  in  $H$

correspond to  $H_1 = \mathbb{C} \oplus M_2(\mathbb{C})$  and  $H_2 = \mathbb{C} \oplus \mathbb{C}^2$ , respectively. We have used the fact that  $\mathbb{C}^2 \otimes \mathbb{C}^{2\circ} \simeq M_2(\mathbb{C})$ . The left action of  $A$  on  $H_1$  is given by the matrix

$$\begin{pmatrix} \lambda & 0 \\ 0 & a \end{pmatrix},$$

with  $a \in M_2(\mathbb{C})$  acting on  $M_2(\mathbb{C}) \subset H_1$  by left matrix multiplication. The right action of  $A$  on  $H_1$  corresponds to the same matrix acting by right matrix multiplication.

On  $H_2$ , the left action of  $A$  is given by matrix multiplication by the above matrix on vectors in  $\mathbb{C} \oplus \mathbb{C}^2$ . However, the right action of  $(\lambda, a) \in A$  is given by scalar multiplication with  $\lambda$  on all of  $H_2$ .

**The real structure:** Before turning to the finite Dirac operator  $D$ , we exploit the presence of a real structure  $J : H \rightarrow H$  in the diagrammatic approach started above.

**Exercise 3.6** Let  $J$  be an anti-unitary operator on a finite-dimensional Hilbert space. Show that  $J^2$  is a unitary operator.

**Lemma 3.8** *Let  $J$  be an anti-unitary operator on a finite-dimensional Hilbert space  $H$  with  $J^2 = \pm 1$ .*

- (1) *If  $J^2 = 1$  then there is an orthonormal basis  $\{e_k\}$  of  $H$  such that  $Je_k = e_k$ .*
- (2) *If  $J^2 = -1$  then there is an orthonormal basis  $\{e_k, f_k\}$  of  $H$  such that  $Je_k = f_k$  (and, consequently,  $Jf_k = -e_k$ ).*

**Proof** (1) Take any  $v \in H$  and set

$$e_1 := \begin{cases} c(v + Jv) & \text{if } Jv \neq -v \\ iv & \text{if } Jv = -v, \end{cases}$$

with  $c$  a normalization constant. Then  $J(v + Jv) = Jv + J^2v = v + Jv$  and  $J(iv) = -iJv = iv$  in the two respective cases, so that  $Je_1 = e_1$ .

Next, take a vector  $v'$  that is orthogonal to  $e_1$ . Then

$$(e_1, Jv') = (J^2v', Je_1) = (v', Je_1) = (v', e_1) = 0,$$

so that also  $Jv' \perp e_1$ . As before, we set


$$e_2 := \begin{cases} c(v' + Jv') & \text{if } Jv' \neq -v' \\ iv' & \text{if } Jv' = -v', \end{cases}$$

which by the above is orthogonal to  $e_1$ . Continuing in this way gives a basis  $\{e_k\}$  for  $H$  with  $Je_k = e_k$ .

(2) Take any  $v \in H$  and set  $e_1 = cv$  with  $c$  a normalization constant. Then  $f_1 = Je_1$  is orthogonal to  $e_1$ , since

$$(f_1, e_1) = (Je_1, e_1) = -(Je_1, J^2e_1) = -(Je_1, e_1) = -(f_1, e_1).$$

**Fig. 3.2** The presence of the real structure  $J$  implies a symmetry in the diagram along the diagonal



Next, take another  $v' \perp e_1, f_1$  and set  $e_2 = c'v'$ . As before,  $f_2 := Je_2$  is orthogonal to  $e_2$ , and also to  $e_1$  and  $f_1$ :

$$\begin{aligned} (e_1, f_2) &= (e_1, Je_2) = -(J^2 e_1, Je_2) = -(e_2, Je_1) = -(e_2, f_1) = 0, \\ (f_1, f_2) &= (Je_1, Je_2) = (e_2, e_1) = 0. \end{aligned}$$

Continuing in this way gives a basis  $\{e_k, f_k\}$  for  $H$  with  $Je_k = f_k$ . □

We will now apply these results to the anti-unitary operator given by a real structure on a spectral triple. Recall that in this case,  $J : H \rightarrow H$  implements a right action of  $A$  on  $H$ , via

$$a^\circ = Ja^*J^{-1}$$

satisfying  $[a, b^\circ] = 0$ . Together with the block-form of  $A$ , this implies that

$$J(a_1^* \oplus \cdots \oplus a_N^*) = (a_1^\circ \oplus \cdots \oplus a_N^\circ)J.$$

We conclude that the Krajewski diagram for a real spectral triple must be symmetric along the diagonal,  $J$  mapping each subspace  $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij}$  bijectively to  $\mathbb{C}^{n_j} \otimes \mathbb{C}^{n_i \circ} \otimes V_{ji}$ .

**Proposition 3.9** *Let  $J$  be a real structure on a finite real spectral triple  $(A, H, D; J)$ .*

- (1) *If  $J^2 = 1$  (KO-dimension 0, 1, 6, 7) then there is an orthonormal basis  $\{e_k^{(ij)}\}$  ( $i, j = 1, \dots, N, k = 1, \dots, \dim V_{ij}\}$ ) with  $e_k^{(ij)} \in \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij}$  such that*

$$Je_k^{(ij)} = e_k^{(ji)}; \quad (i, j = 1, \dots, N; k = 1, \dots, \dim V_{ij}).$$

- (2) If  $J^2 = -1$  (KO-dimension 2, 3, 4, 5) then there is an orthonormal basis  $\{e_k^{(ij)}, f_k^{(ji)}\}$  ( $i \leq j = 1, \dots, N, k = 1, \dots, \dim V_{ij}$ ) with  $e_k^{(ij)} \in \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij}$ ,  $f_k^{(ji)} \in \mathbb{C}^{n_j} \otimes \mathbb{C}^{n_i \circ} \otimes V_{ji}$  and such that

$$Je_k^{(ij)} = f_k^{(ji)}; \quad (i \leq j = 1, \dots, N; k = 1, \dots, \dim V_{ij}).$$

**Proof** We imitate the proof Lemma 3.8.

- (1) If  $i \neq j$ , take  $v \in \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij}$  and set  $e_1^{(ij)} = cv$ . Then, by the above observation,  $e_1^{(ji)} = Je_1^{(ij)}$  is an element in  $\mathbb{C}^{n_j} \otimes \mathbb{C}^{n_i \circ} \otimes V_{ji}$ . Next, take  $v' \in \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij}$  with  $v' \perp v$  and apply the same procedure to obtain  $e_2^{(ij)}$  and  $e_2^{(ji)}$ . Continuing in this way gives an orthonormal basis  $\{e_k^{(ij)}\}$  for  $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij}$ , and an orthonormal basis  $\{e_k^{(ji)}\}$  for  $\mathbb{C}^{n_j} \otimes \mathbb{C}^{n_i \circ} \otimes V_{ji}$  which satisfy  $Je_k^{(ij)} = e_k^{(ji)}$ .

If  $i = j$ , then Lemma 3.8(1) applies directly to the anti-unitary operator given by  $J$  restricted to  $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_i \circ} \otimes V_{ii}$ .

- (2) can be proved along the same lines.  $\square$

Note that this result implies that in the case of KO-dimension 2, 3, 4 and 5, the diagonal  $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_i \circ} \otimes V_{ii}$  needs to have even multiplicity.

**The finite Dirac operator:** Corresponding to the above decomposition of  $H$  we can write  $D$  as a sum of matrices

$$D_{ij,kl} : \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij} \rightarrow \mathbb{C}^{n_k} \otimes \mathbb{C}^{n_l \circ} \otimes V_{kl},$$

restricted to these subspaces. The condition  $D^* = D$  implies that  $D_{kl,ij} = D_{ij,kl}^*$ . In terms of the above diagrammatic representation of  $H$ , we express a non-zero  $D_{ij,kl}$  as a line between the nodes  $(\mathbf{n}_i, \mathbf{n}_j^\circ)$  and  $(\mathbf{n}_k, \mathbf{n}_l^\circ)$ . Instead of drawing directed lines, we draw a single undirected line, capturing both  $D_{ij,kl}$  and its adjoint  $D_{kl,ij}$ .


**Lemma 3.10** *The condition  $JD = \pm DJ$  and the order one condition given by  $[[D, a], b^\circ] = 0$  forces the lines in the diagram to run only vertically or horizontally (or between the same node), thereby maintaining the diagonal symmetry between the nodes in the diagram.*

**Proof** The condition  $JD = \pm DJ$  easily translates into a commuting diagram:

$$\begin{array}{ccc} \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij} & \xrightarrow{D} & \mathbb{C}^{n_k} \otimes \mathbb{C}^{n_l \circ} \otimes V_{kl} \\ J \downarrow & & J \downarrow \\ \mathbb{C}^{n_j} \otimes \mathbb{C}^{n_i \circ} \otimes V_{ji} & \xrightarrow{\pm D} & \mathbb{C}^{n_l} \otimes \mathbb{C}^{n_k \circ} \otimes V_{lk} \end{array}$$

thus relating  $D_{ij,kl}$  to  $D_{ji,lk}$ , maintaining the diagonal symmetry (Fig. 3.3).

**Fig. 3.3** The lines between two nodes represent a non-zero  $D_{li,ji}$ :  
 $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_i^\circ} \rightarrow \mathbb{C}^{n_j} \otimes \mathbb{C}^{n_j^\circ}$ , as well as its adjoint  $D_{ji,ii}$ :  
 $\mathbb{C}^{n_j} \otimes \mathbb{C}^{n_j^\circ} \rightarrow \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_i^\circ}$ . The non-zero components  $D_{ii,ij}$  and  $D_{ij,ii}$  are related to  $\pm D_{li,ji}$  and  $\pm D_{ji,ii}$ , respectively, according to  $JD = \pm DJ$



If we write the order one condition  $[[D, a], b^\circ] = 0$  for diagonal elements  $a = \lambda_1 \mathbb{I}_{n_1} \oplus \cdots \oplus \lambda_N \mathbb{I}_{n_N} \in A$  and  $b = \mu_1 \mathbb{I}_{n_1} \oplus \cdots \oplus \mu_N \mathbb{I}_{n_N} \in A$  with  $\lambda_i, \mu_i \in \mathbb{C}$ , we compute

$$D_{ij,kl}(\lambda_i - \lambda_k)(\bar{\mu}_j - \bar{\mu}_l) = 0,$$

for all  $\lambda_i, \mu_j \in \mathbb{C}$ . As a consequence,  $D_{ij,kl} = 0$  whenever  $i \neq k$  or  $j \neq l$ .  $\square$

**Grading:** Finally, if there is a grading  $\gamma : H \rightarrow H$ , then each node in the diagram gets labeled by a plus or minus sign. The rules are that:

- $D$  connects nodes with different signs;
- If the node  $(\mathbf{n}_i, \mathbf{n}_j^\circ)$  has sign  $\pm$ , then the node  $(\mathbf{n}_j, \mathbf{n}_i^\circ)$  has sign  $\pm \epsilon''$ , according to  $J\gamma = \epsilon''\gamma J$ .

Finally, we arrive at a diagrammatic classification of finite real spectral triples of any KO-dimension.

**Definition 3.11** A Krajewski diagram of KO-dimension  $k$  is given by an ordered pair  $(\Gamma, \Lambda)$  of a finite graph  $\Gamma$  and a finite set  $\Lambda$  of positive integers with a labeling:

- of the vertices  $v \in \Gamma^{(0)}$  by elements  $\iota(v) = (n(v), m(v)) \in \Lambda \times \Lambda$ , where the existence of an edge from  $v$  to  $v'$  implies that either  $n(v) = n(v')$ ,  $m(v) = m(v')$ , or both;
- of the edges  $e = (v_1, v_2) \in \Gamma^{(1)}$  by non-zero operators:

$$\begin{aligned} D_e : \mathbb{C}^{n(v_1)} &\rightarrow \mathbb{C}^{n(v_2)} && \text{if } m(v_1) = m(v_2); \\ D_e : \mathbb{C}^{m(v_1)} &\rightarrow \mathbb{C}^{m(v_2)} && \text{if } n(v_1) = n(v_2), \end{aligned}$$

and their adjoints  $D_e^*$ ,

together with an involutive graph automorphism  $j : \Gamma \rightarrow \Gamma$  so that the following conditions hold:

- (1) every row or column in  $\Lambda \times \Lambda$  has non-empty intersection with  $\iota(\Gamma)$ ;
- (2) for each vertex  $v$  we have  $(n(j(v))) = m(v)$ ;
- (3) for each edge  $e$  we have  $D_e = \epsilon' D_{j(e)}$ ;
- (4) if the KO-dimension  $k$  is even, then the vertices are additionally labeled by  $\pm 1$  and the edges only connect opposite signs. The signs at  $v$  and  $j(v)$  differ by a factor  $\epsilon$ , according to the table of Definition 3.1;
- (5) if the KO-dimension is 2, 3, 4, 5 then the inverse image under  $\iota$  of the diagonal elements in  $\Lambda \times \Lambda$  contains an even number of vertices of  $\Gamma$ .

Note that this definition allows for different vertices of  $\Gamma$  to be labeled by the same element in  $\Lambda \times \Lambda$ ; this accounts for the multiplicities appearing in  $V_{ij}$  that we have encountered before.

This indeed gives rise to a diagram of the above type, by putting a node at position  $(\mathbf{n}_i, \mathbf{n}_j^\circ)$  for each vertex carrying the label  $(\mathbf{n}_i, \mathbf{n}_j) \in \Lambda \times \Lambda$ . The notation  $\mathbf{n}_j^\circ$  instead of  $\mathbf{n}_j$  is just for a convenient diagrammatic exposition. The operators  $D_e$  between vertices that are labeled by  $(\mathbf{n}_i, \mathbf{n}_j)$  and  $(\mathbf{n}_k, \mathbf{n}_l)$ , respectively, add up to the above  $D_{ij,kl}$ . Explicitly,

$$D_{ij,kl} = \sum_{\substack{e=(v_1,v_2) \in \Gamma^{(1)} \\ \iota(v_1)=(\mathbf{n}_i,\mathbf{n}_j) \\ \iota(v_2)=(\mathbf{n}_k,\mathbf{n}_l)}} D_e,$$


so that indeed  $D_{ij,kl}^* = D_{kl,ij}$ . Moreover, the only non-zero entries  $D_{ij,kl}$  will appear when  $i = k$ , or  $j = l$ , or both. Thus, we have shown

**Theorem 3.12** *There is a one-to-one correspondence between finite real spectral triples of KO-dimension  $k$  modulo unitary equivalence and Krajewski diagrams of KO-dimension  $k$ . Specifically, one associates a real spectral triple  $(A, H, D; J, \gamma)$  to a Krajewski diagram in the following way:*

$$\begin{aligned} A &= \bigoplus_{n \in \Lambda} M_n(\mathbb{C}); \\ H &= \bigoplus_{v \in \Gamma^{(0)}} \mathbb{C}^{n(v)} \otimes \mathbb{C}^{m(v)\circ}; \\ D &= \sum_{e \in \Gamma^{(1)}} D_e + D_e^*. \end{aligned}$$

Moreover, the real structure  $J : H \rightarrow H$  is given as in Proposition 3.9, with the basis dictated by the graph automorphism  $j : \Gamma \rightarrow \Gamma$ . Finally, a grading  $\gamma$  on  $H$  is defined by setting  $\gamma$  to be  $\pm 1$  on  $\mathbb{C}^{n(v)} \otimes \mathbb{C}^{m(v)\circ} \subset H$  according to the labeling by  $\pm 1$  of the vertex  $v$ .

**Example 3.13** Consider the case  $A = \mathbb{C} \oplus \mathbb{C}$ . There are ten possible Krajewski diagrams in KO-dimension 0 with multiplicities less than or equal to 1: in terms of  $\widehat{A} = \{\mathbf{1}_1, \mathbf{1}_2\}$ , we have



where the diagonal vertices are labeled with a plus sign, and the off-diagonal vertices with a minus sign.

Let us consider the last diagram in the top row in more detail and give the corresponding spectral triple:

$$\begin{array}{cc} \mathbf{1}_1 & \mathbf{1}_2 \\ \mathbf{1}_1^\circ & \text{---} \\ \mathbf{1}_2^\circ & \text{---} \end{array}$$

First, the inner product space is  $H = \mathbb{C}^3$ , where we choose the middle copy of  $\mathbb{C}$  to correspond to the node on the diagonal. The edges indicate that there are non-zero components of  $D$  that map between the first two copies of  $\mathbb{C}$  in  $H$  and between the second and third copy of  $\mathbb{C}$ . In other words,

$$D = \begin{pmatrix} 0 & \lambda & 0 \\ \bar{\lambda} & 0 & \mu \\ 0 & \bar{\mu} & 0 \end{pmatrix}$$

for some  $\lambda, \mu \in \text{Hom}(\mathbb{C}, \mathbb{C}) \simeq \mathbb{C}$  that are the given labels on the two edges. In this basis,

$$\gamma = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

Finally,  $J$  is given by the matrix  $K$  composed with complex conjugation on  $H$ , where

$$K = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

From this it is clear that we indeed have

$$D\gamma = -\gamma D; \quad DJ = JD; \quad J\gamma = \gamma J.$$

**Exercise 3.7** Use the ten Krajewski diagrams of the previous example to show that on  $A = \mathbb{C} \oplus \mathbb{C}$  a finite real spectral triple of  $KO$ -dimension 6 with  $\dim H \leq 4$  must have vanishing finite Dirac operator.

**Example 3.14** Consider  $A = M_n(\mathbb{C})$  so that  $\widehat{A} = \{\mathbf{n}\}$ . We then have a Krajewski diagram

$$\begin{array}{c} \mathbf{n} \\ \mathbf{n}^\circ \quad \circ \end{array}$$

The node can be labeled only by either plus or minus one, the choice being irrelevant. This means that  $H = \mathbb{C}^n \otimes \mathbb{C}^{n^\circ} \simeq M_n(\mathbb{C})$  with  $\gamma$  the trivial grading. The operator  $J$  is a combination of complex conjugation and the flip on  $\mathbf{n} \otimes \mathbf{n}^\circ$ : this translates to  $M_n(\mathbb{C})$  as taking the matrix adjoint. Moreover, since the single node has label  $\pm 1$ , there are no non-zero Dirac operators. Hence, the finite real spectral triple of this diagram corresponds to

$$(A = M_n(\mathbb{C}), H = M_n(\mathbb{C}), D = 0; J = (\cdot)^*, \gamma = 1),$$

and was encountered already in Exercise 3.3.

### 3.3 Real Algebras and Krajewski Diagrams

Thus far, we have considered finite spectral triples on complex algebras. In practice, it is useful to allow real  $*$ -algebras in Definition 2.19 as well.

**Definition 3.15** A *real algebra* is a vector space  $A$  over  $\mathbb{R}$  with a bilinear associative product  $A \times A \rightarrow A$  denoted by  $(a, b) \mapsto ab$  and a unit  $1$  satisfying  $1a = a1 = a$  for all  $a \in A$ .

A *real  $*$ -algebra* (or, *involutive algebra*) is a real algebra  $A$  together with a real linear map (the *involution*)  $* : A \rightarrow A$  such that  $(ab)^* = b^*a^*$  and  $(a^*)^* = a$  for all  $a, b \in A$ .

**Example 3.16** A particularly interesting example in this context is given by  $\mathbb{H}$ , the real  $*$ -algebra of quaternions, defined as a real subalgebra of  $M_2(\mathbb{C})$ :

$$\mathbb{H} = \left\{ \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} : \alpha, \beta \in \mathbb{C} \right\}.$$

This is indeed closed under multiplication. As a matter of fact,  $\mathbb{H}$  consists of those matrices in  $M_2(\mathbb{C})$  that commute with the operator  $I$  defined by

$$I \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} -\bar{v}_2 \\ \bar{v}_1 \end{pmatrix}.$$

The involution is inherited from  $M_2(\mathbb{C})$  and is given by hermitian conjugation.

- Exercise 3.8** (1) Show that  $\mathbb{H}$  is a real  $*$ -algebra which contains a real subalgebra isomorphic to  $\mathbb{C}$ .  
 (2) Show that  $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{C} \simeq M_2(\mathbb{C})$  as complex  $*$ -algebras.  
 (3) Show that  $M_k(\mathbb{H})$  is a real  $*$ -algebra for any integer  $k$ .  
 (4) Show that  $M_k(\mathbb{H}) \otimes_{\mathbb{R}} \mathbb{C} \simeq M_{2k}(\mathbb{C})$  as complex  $*$ -algebras.

When considering Hilbert space representations of a real  $*$ -algebra, one must be careful, because the Hilbert space will be assumed to be a complex space.

**Definition 3.17** A *representation* of a finite-dimensional real  $*$ -algebra  $A$  is a pair  $(H, \pi)$  where  $H$  is a (finite-dimensional, complex) Hilbert space and  $\pi$  is a real-linear  $*$ -algebra map

$$\pi : A \rightarrow L(H).$$

Also, although there is a great deal of similarity, we stress that the definition of the real structure  $J$  in Definition 2.19 is not related to the algebra  $A$  being real or complex.

**Exercise 3.9** Show that there is a one-to-one correspondence between Hilbert space representations of a real  $*$ -algebra  $A$  and complex representations of its complexification  $A \otimes_{\mathbb{R}} \mathbb{C}$ . Conclude that the unique irreducible (Hilbert space) representation of  $M_k(\mathbb{H})$  is given by  $\mathbb{C}^{2k}$ .

**Lemma 3.18** Suppose that a real  $*$ -algebra  $A$  is represented faithfully on a finite-dimensional Hilbert space  $H$  through a real-linear  $*$ -algebra map  $\pi : A \rightarrow L(H)$ . Then  $A$  is a matrix algebra:

$$A \simeq \bigoplus_{i=1}^N M_{n_i}(\mathbb{F}_i),$$

where  $\mathbb{F}_i = \mathbb{R}, \mathbb{C}$  or  $\mathbb{H}$ , depending on  $i$ .

**Proof** The representation  $\pi$  allows to consider  $A$  as a real  $*$ -subalgebra of  $M_{\dim H}(\mathbb{C})$ , hence  $A + iA$  can be considered a complex  $*$ -subalgebra of  $M_{\dim H}(\mathbb{C})$ . Thus  $A + iA$  is a matrix algebra, and we may restrict to the case  $A + iA = M_k(\mathbb{C})$  for some  $k \geq 1$ . Note that  $A \cap iA$  is a two-sided  $*$ -ideal in  $M_k(\mathbb{C})$ . As such, it must be either the whole of  $M_k(\mathbb{C})$ , or zero. In the first case,  $A + iA = A \cap iA$  so that  $A = M_k(\mathbb{C})$ . If  $A \cap iA = \{0\}$ , then we can uniquely write any element in  $M_k(\mathbb{C})$  as  $a + ib$  with  $a, b \in A$ . Moreover,  $A$  is the fixed point algebra of the anti-linear automorphism  $\alpha$  of  $M_k(\mathbb{C})$  given by  $\alpha(a + ib) = a - ib$  ( $a, b \in A$ ). We can implement  $\alpha$  by an anti-linear isometry  $I$  on  $\mathbb{C}^k$  such that  $\alpha(x) = IxI^{-1}$  for all  $x \in M_k(\mathbb{C})$ . Since  $\alpha^2 = 1$ , the operator  $I^2$  commutes with  $M_k(\mathbb{C})$  and is therefore proportional to a complex scalar. Together with  $I^2$  being an isometry, this implies that  $I^2 = \pm 1$  and that  $A$  is precisely the commutant of  $I$ . We now once again use Lemma 3.8 to conclude that

- If  $I^2 = 1$ , then there is a basis  $\{e_i\}$  of  $\mathbb{C}^k$  such that  $Ie_i = e_i$ . Since a matrix in  $M_k(\mathbb{C})$  that commutes with  $I$  must have real entries, this gives

$$A = M_k(\mathbb{R}).$$

- If  $I^2 = -1$ , then there is a basis  $\{e_i, f_i\}$  of  $\mathbb{C}^k$  such that  $Ie_i = f_i$  (and thus  $k$  is even). Since a matrix in  $M_k(\mathbb{C})$  that commutes with  $I$  must be a  $k/2 \times k/2$ -matrix with quaternionic entries, we obtain

$$A = M_{k/2}(\mathbb{H}). \quad \square$$

We now reconsider the diagrammatic classification of finite spectral triples, with real  $*$ -algebras represented faithfully on a Hilbert space. In fact, as far as the decomposition of  $H$  into irreducible representations is concerned, we can replace  $A$  by the complex  $*$ -algebra

$$A + iA \simeq \bigoplus_{i=1}^N M_{n_i}(\mathbb{C}).$$

Thus, the Krajewski diagrams in Definition 3.11 classify such finite real spectral triples as well as long as we take the  $\mathbb{F}_i$  for each  $i$  into account. That is, we enhance the set  $\Lambda$  to be

$$\Lambda = \{\mathbf{n}_1 \mathbb{F}_1, \dots, \mathbf{n}_N \mathbb{F}_N\},$$

reducing to the previously defined  $\Lambda$  when all  $\mathbb{F}_i = \mathbb{C}$ .

## 3.4 Classification of Irreducible Geometries

We now classify *irreducible* finite real spectral triples of KO-dimension 6. This leads to a remarkably concise list of spectral triples, based on the matrix algebras  $M_N(\mathbb{C}) \oplus M_N(\mathbb{C})$  for some  $N$ .

**Definition 3.19** A finite real spectral triple  $(A, H, D; J, \gamma)$  is called *irreducible* if the triple  $(A, H, J)$  is irreducible. More precisely, we demand that:

- (1) The representations of  $A$  and  $J$  in  $H$  are irreducible;
- (2) The action of  $A$  on  $H$  has a separating vector (cf. Exercise 3.2).

**Theorem 3.20** Let  $(A, H, D; J, \gamma)$  be an irreducible finite real spectral triple of KO-dimension 6. Then there exists a positive integer  $N$  such that  $A \simeq M_N(\mathbb{C}) \oplus M_N(\mathbb{C})$ .


**Proof** Let  $(A, H, D; J, \gamma)$  be an arbitrary finite real spectral triple, corresponding to e.g., the Krajewski diagram of Fig. 3.2. Thus, as in Sect. 2.3 we have

$$A = \bigoplus_{i=1}^N M_{n_i}(\mathbb{C}), \quad H = \bigoplus_{i,j=1}^N \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \otimes V_{ij},$$

with  $V_{ij}$  corresponding to the multiplicities as before. Now each  $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j}$  is an irreducible representation of  $A$ , but in order for  $H$  to support a real structure  $J : H \rightarrow H$  we need both  $\mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j}$  and  $\mathbb{C}^{n_j} \otimes \mathbb{C}^{n_i}$  to be present in  $H$ . Moreover, Lemma 3.8 with  $J^2 = 1$  assures that already with multiplicities  $\dim V_{ij} = 1$  there exists such a real structure. Hence, the irreducibility condition (1) above yields

$$H = \mathbb{C}^{n_i} \otimes \mathbb{C}^{n_j \circ} \oplus \mathbb{C}^{n_j} \otimes \mathbb{C}^{n_i \circ},$$

for some  $i, j \in \{1, \dots, N\}$ . Or, as a Krajewski diagram:



Then, let us consider condition (2) on the existence of a separating vector. Note first that the representation of  $A$  in  $H$  is faithful only if  $A = M_{n_i}(\mathbb{C}) \oplus M_{n_j}(\mathbb{C})$ . Second, the stronger condition of a separating vector  $\xi$  then implies  $n_i = n_j$ , as it is equivalent to  $A'\xi = H$  for the commutant  $A'$  of  $A$  in  $H$  (see Exercise 3.2). Namely, since  $A' = M_{n_j}(\mathbb{C}) \oplus M_{n_i}(\mathbb{C})$  with  $\dim A' = n_i^2 + n_j^2$ , and  $\dim H = 2n_i n_j$  we find the desired equality  $n_i = n_j$ .  $\square$

With the complex finite-dimensional algebras  $A$  given by  $M_N(\mathbb{C}) \oplus M_N(\mathbb{C})$ , the additional demand that  $H$  carries a symplectic structure  $I^2 = -1$  yields real algebras of which  $A$  is the complexification (as in the proof of Lemma 3.18). In view of Exercise 3.8(4) we see that this requires  $N = 2k$  so that one naturally considers triples  $(A, H, J)$  for which  $A = M_k(\mathbb{H}) \oplus M_{2k}(\mathbb{C})$  and  $H = \mathbb{C}^{2(2k)^2}$ . The case  $k = 2$  will come back in the final Chap. 13 as the relevant one to consider in particle physics applications that go beyond the Standard Model.

## Notes

### Section 3.1. Finite Real Spectral Triples

1. The operator  $D$  in Definition 3.1 is a first-order differential operator on the bimodule  $H$  in the sense of [1].

2. Exercises 3.2 and 3.3 develop Tomita–Takesaki theory for matrix algebras, considered as finite-dimensional Von Neumann algebras. For a complete treatment of this theory for general Von Neumann algebras, we refer to e.g., [2].

### Section 3.2. Classification of Finite Real Spectral Triples

3. Krajewski’s work on the classification of all finite real spectral triples  $(A, H, D; J, \gamma)$  modulo unitary equivalence (based on a suggestion in [3]) is published in [4]. Similar results were obtained independently in [5]. We have extended Krajewski’s work—which is in KO-dimension 0—to any KO-dimension. The classification of finite real spectral triples (but without Krajewski diagrams) is also the subject of [6]. The KO-dimension 6 case—which is of direct physical interest as we will see below in Chap. 13—was also handled in [7].
4. Lemma 3.8 is based on [8], where Wigner showed that anti-unitary operators on finite-dimensional Hilbert spaces can be written in a normal form. His crucial observation is that  $J^2$  is unitary, allowing for a systematic study of a normal form of  $J$  for each of the eigenvalues of  $J^2$  (these eigenvalues form a discrete subset of the complex numbers of modulus one). In our case of interest,  $J$  is a real structure on a spectral triple (as in Definition 3.1), so that  $J^2 = \pm 1$ .
5. In the labelling of the nodes in a Krajewski diagram with  $\pm$ -signs, it is important whether or not we adopt the so-called *orientation axiom* [3]. In the finite-dimensional case, this axiom demands that the grading  $\gamma$  can be implemented by elements  $x_i, y_i \in A$  as  $\gamma = \sum_i x_i y_i^\circ$ . Hence, this is completely dictated by the operator  $J$  and the representation of  $A$ . In terms of our diagrams, this translates to the fact that the grading of a node only depends on the label  $(\mathbf{n}_i, \mathbf{n}_j^\circ)$ . In this book, we will not assume the orientation axiom.

### Section 3.4. Classification of Irreducible Geometries

6. Finite irreducible geometries have been classified by Chamseddine and Connes in [9], using different methods. We here confront their result with the above approach to finite spectral triples using Krajewski diagrams and find that they are compatible.

## References

1. Dubois-Violette, M., Masson, T.: On the first-order operators in bimodules. *Lett. Math. Phys.* **37**, 467–474 (1996)
2. Takesaki, M.: Theory of operator algebras. II. Encyclopaedia of Mathematical Sciences, vol. 125. Springer, Berlin (2003). Operator Algebras and Non-commutative Geometry, 6
3. Connes, A.: Gravity coupled with matter and the foundation of non-commutative geometry. *Commun. Math. Phys.* **182**, 155–176 (1996)
4. Krajewski, T.: Classification of finite spectral triples. *J. Geom. Phys.* **28**, 1–30 (1998)
5. Paschke, M., Sitarz, A.: Discrete spectral triples and their symmetries. *J. Math. Phys.* **39**, 6191–6205 (1998)
6. Ćačić, B.: Moduli spaces of Dirac operators for finite spectral triples. In: Marcolli, M., Parashar, D. (eds.) Quantum Groups and Noncommutative Spaces: Perspectives on Quantum Geometry. Vieweg Verlag (2010)
7. Jureit, J.-H., Stephan, C.A.: On a classification of irreducible almost-commutative geometries IV. *J. Math. Phys.* **49**, 033502 (2008)
8. Wigner, E.P.: Normal form of antiunitary operators. *J. Math. Phys.* **1**, 409–413 (1960)
9. Chamseddine, A.H., Connes, A.: Why the standard model. *J. Geom. Phys.* **58**, 38–47 (2008)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 4

## Riemannian Spin Manifolds



We now extend our treatment of noncommutative geometric spaces from the finite case to the continuum. This generalizes spin manifolds to the noncommutative world. The resulting spectral triples (Chap. 5) form the key technical device in noncommutative geometry, and in the physical applications of Part 2 of this book in particular.

We start with a treatment of Clifford algebras, as a preparation for the definition of a spin structure on a Riemannian manifold, and end with the analytical aspects of the Dirac operator.

### 4.1 Clifford Algebras

Let  $V$  be a vector space over a field  $\mathbb{F}$  ( $= \mathbb{R}, \mathbb{C}$  or  $\mathbb{H}$ ), equipped with a quadratic form  $Q : V \rightarrow \mathbb{F}$ , i.e.

$$Q(\lambda v) = \lambda^2 Q(v); \quad (\lambda \in \mathbb{F}, v \in V),$$
$$Q(v + w) + Q(v - w) = 2Q(v) + 2Q(w); \quad (v, w \in V).$$

**Definition 4.1** For a quadratic form  $Q$  on  $V$ , the *Clifford algebra*  $\text{Cl}(V, Q)$  is the algebra generated (over  $\mathbb{F}$ ) by the vectors  $v \in V$  and with unit 1 subject to the relation

$$v^2 = Q(v)1. \tag{4.1.1}$$

Note that the Clifford algebra  $\text{Cl}(V, Q)$  is  $\mathbb{Z}_2$ -graded, with grading  $\chi$  given by

$$\chi(v_1 \cdots v_k) = (-1)^k v_1 \cdots v_k,$$

which is indeed compatible with relation (4.1.1). Accordingly, we decompose

$$\text{Cl}(V, Q) =: \text{Cl}^0(V, Q) \oplus \text{Cl}^1(V, Q)$$

into an even and odd part.

**Exercise 4.1** Show that in  $\text{Cl}(V, Q)$  we have

$$vw + wv = 2g_Q(v, w),$$

where  $g_Q$  is the pairing  $V \times V \rightarrow \mathbb{F}$  associated to  $Q$ , given by

$$g_Q(v, w) = \frac{1}{2} (Q(v + w) - Q(v) - Q(w)).$$

We also introduce the following convenient notation for the Clifford algebras for the vector spaces  $\mathbb{R}^n$  and  $\mathbb{C}^n$  equipped with the standard quadratic form  $Q_n(x_1, \dots, x_n) = x_1^2 + \dots + x_n^2$ :

$$\begin{aligned}\text{Cl}_n^+ &:= \text{Cl}(\mathbb{R}^n, Q_n); \\ \text{Cl}_n^- &:= \text{Cl}(\mathbb{R}^n, -Q_n); \\ \text{Cl}_n &:= \text{Cl}(\mathbb{C}^n, Q_n).\end{aligned}$$

Both  $\text{Cl}_n^+$  and  $\text{Cl}_n^-$  are algebras over  $\mathbb{R}$  generated by  $e_1, \dots, e_n$  with relations

$$e_i e_j + e_j e_i = \pm 2\delta_{ij}, \quad (4.1.2)$$

for all  $i, j = 1, \dots, n$ . Moreover, the even part  $(\text{Cl}_n^\pm)^0$  of  $\text{Cl}_n^\pm$  consists of products of an even number of  $e_i$ 's, and the odd part  $(\text{Cl}_n^\pm)^1$  of products of an odd number of  $e_i$ 's.

The Clifford algebra  $\text{Cl}_n$  is the complexification of both  $\text{Cl}_n^+$  and  $\text{Cl}_n^-$ , and is therefore generated over  $\mathbb{C}$  by the same  $e_1, \dots, e_n$  satisfying (4.1.2).

- Exercise 4.2**
- (1) Check that Eq. (4.1.2) indeed corresponds to the defining relations in  $\text{Cl}_n^\pm$ .
  - (2) Show that the elements  $e_{i_1} \cdots e_{i_r}$  with  $1 \leq i_1 < i_2 < \dots < i_r \leq n$  form a basis for  $\text{Cl}_n^\pm$ .
  - (3) Conclude that  $\dim_{\mathbb{R}} \text{Cl}_n^\pm = 2^n$  and, accordingly,  $\dim_{\mathbb{C}} \text{Cl}_n = 2^n$ .
  - (4) Find an isomorphism  $\text{Cl}(\mathbb{C}^n, Q_n) \simeq \text{Cl}(\mathbb{C}^n, -Q_n)$  as Clifford algebras.

**Proposition 4.2** *The even part  $(\text{Cl}_{n+1}^-)^0$  of  $\text{Cl}_{n+1}^-$  is isomorphic to  $\text{Cl}_n^-$ .*

**Proof** We construct a map  $\Psi : \text{Cl}_n^- \mapsto (\text{Cl}_{n+1}^-)^0$  given on generators by

$$\Psi(e_i) = e_{n+1} e_i. \quad (4.1.3)$$

Indeed, for  $i, j = 1, \dots, n$  we have

$$\Psi(e_i)\Psi(e_j) + \Psi(e_j)\Psi(e_i) = e_i e_j + e_j e_i = -2\delta_{ij} = \Psi(-2\delta_{ij}),$$

using  $e_i e_{n+1} = -e_{n+1} e_i$  and  $e_{n+1} e_{n+1} = -1$ . Thus,  $\Psi$  extends to a homomorphism  $\text{Cl}_n^- \mapsto (\text{Cl}_{n+1}^-)^0$ . Moreover, since  $\Psi$  sends basis vectors in  $\text{Cl}_n^-$  to basis vectors in  $(\text{Cl}_{n+1}^-)^0$  and the dimensions of  $\text{Cl}_n^-$  and  $(\text{Cl}_{n+1}^-)^0$  coincide, it is an isomorphism.  $\square$

**Exercise 4.3** Show that the same expression (4.1.3) induces an isomorphism from  $\text{Cl}_n^-$  to the even part  $(\text{Cl}_{n+1}^+)^0$  and conclude that  $(\text{Cl}_{n+1}^+)^0 \simeq (\text{Cl}_{n+1}^-)^0$ .

Next, we compute the Clifford algebras  $\text{Cl}_n^\pm$  and  $\text{Cl}_n$ . We start with a recursion relation:

**Proposition 4.3** *For any  $k \geq 1$  we have*

$$\begin{aligned}\text{Cl}_k^+ \otimes_{\mathbb{R}} \text{Cl}_2^- &\simeq \text{Cl}_{k+2}^-, \\ \text{Cl}_k^- \otimes_{\mathbb{R}} \text{Cl}_2^+ &\simeq \text{Cl}_{k+2}^+.\end{aligned}$$

**Proof** The map  $\Psi : \text{Cl}_{k+2}^- \rightarrow \text{Cl}_k^+ \otimes_{\mathbb{R}} \text{Cl}_2^-$  given on generators by

$$\Psi(e_i) = \begin{cases} 1 \otimes e_i & i = 1, 2 \\ e_{i-2} \otimes e_1 e_2 & i = 3, \dots, n \end{cases}$$

extends to the desired isomorphism.  $\square$

Let us compute some of the Clifford algebras in lowest dimensions.

**Proposition 4.4**

$$\begin{aligned}\text{Cl}_1^+ &\simeq \mathbb{R} \oplus \mathbb{R}, & \text{Cl}_1^- &\simeq \mathbb{C}, \\ \text{Cl}_2^+ &\simeq M_2(\mathbb{R}), & \text{Cl}_2^- &\simeq \mathbb{H}.\end{aligned}$$

**Proof** The Clifford algebra  $\text{Cl}_1^+$  is generated (over  $\mathbb{R}$ ) by 1 and  $e_1$  with relation  $e_1^2 = 1$ . We map  $\text{Cl}_1^+$  linearly to the algebra  $\mathbb{R} \oplus \mathbb{R}$  by sending

$$1 \mapsto (1, 1), \quad e_1 \mapsto (1, -1).$$

A dimension count shows that this map is a bijection.

The Clifford algebra  $\text{Cl}_2^+$  is generated by 1,  $e_1$ ,  $e_2$  with relations

$$e_1^2 = 1, \quad e_2^2 = 1, \quad e_1 e_2 = -e_2 e_1.$$

A bijective map  $\text{Cl}_2^+ \xrightarrow{\sim} M_2(\mathbb{R})$  is given on generators by

$$1 \mapsto \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad e_1 \mapsto \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad e_2 \mapsto \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

We leave the remaining  $\text{Cl}_1^-$  and  $\text{Cl}_2^-$  as an illustrative exercise to the reader.  $\square$

**Exercise 4.4** Show that  $\text{Cl}_1^- \simeq \mathbb{C}$  and  $\text{Cl}_2^- \simeq \mathbb{H}$ .

**Table 4.1** Clifford algebras  $\text{Cl}_n^\pm$  and their complexifications  $\mathbb{C}\text{Cl}_n$  for  $n = 1, \dots, 8$ 

$n$	$\text{Cl}_n^+$	$\text{Cl}_n^-$	$\mathbb{C}\text{Cl}_n$
1	$\mathbb{R} \oplus \mathbb{R}$	$\mathbb{C}$	$\mathbb{C} \oplus \mathbb{C}$
2	$M_2(\mathbb{R})$	$\mathbb{H}$	$M_2(\mathbb{C})$
3	$M_2(\mathbb{C})$	$\mathbb{H} \oplus \mathbb{H}$	$M_2(\mathbb{C}) \oplus M_2(\mathbb{C})$
4	$M_2(\mathbb{H})$	$M_2(\mathbb{H})$	$M_4(\mathbb{C})$
5	$M_2(\mathbb{H}) \oplus M_2(\mathbb{H})$	$M_4(\mathbb{C})$	$M_4(\mathbb{C}) \oplus M_4(\mathbb{C})$
6	$M_4(\mathbb{H})$	$M_8(\mathbb{R})$	$M_8(\mathbb{C})$
7	$M_8(\mathbb{C})$	$M_8(\mathbb{R}) \oplus M_8(\mathbb{R})$	$M_8(\mathbb{C}) \oplus M_8(\mathbb{C})$
8	$M_{16}(\mathbb{R})$	$M_{16}(\mathbb{R})$	$M_{16}(\mathbb{C})$

Combining the above two Propositions, we derive Table 4.1 for the Clifford algebras  $\text{Cl}_n^\pm$  and  $\mathbb{C}\text{Cl}_n$  for  $n = 1, \dots, 8$ . For instance,

$$\text{Cl}_3^+ \simeq \text{Cl}_1^- \otimes_{\mathbb{R}} \text{Cl}_2^+ \simeq \mathbb{C} \otimes_{\mathbb{R}} M_2(\mathbb{R}) \simeq M_2(\mathbb{C})$$

and

$$\text{Cl}_4^+ \simeq \text{Cl}_2^- \otimes_{\mathbb{R}} \text{Cl}_2^+ \simeq \mathbb{H} \otimes_{\mathbb{R}} M_2(\mathbb{R}) \simeq M_2(\mathbb{H})$$

and so on. In particular, we have

$$\text{Cl}_n^+ \otimes \text{Cl}_4^+ \simeq \text{Cl}_{n+4}^+$$

and

$$\text{Cl}_{n+8}^+ \simeq \text{Cl}_n^+ \otimes \text{Cl}_8^+.$$

With  $\text{Cl}_8^+ \simeq M_{16}(\mathbb{R})$  we conclude that  $\text{Cl}_{k+8}^+$  is *Morita equivalent* to  $\text{Cl}_k^+$  (cf. Theorem 2.14). Similarly,  $\text{Cl}_{k+8}^-$  is Morita equivalent to  $\text{Cl}_k^-$ . Thus, in this sense Table 4.1 has periodicity eight and we have determined  $\text{Cl}_n^\pm$  for all  $n$ .

For the complex Clifford algebras, there is a periodicity of two:

$$\mathbb{C}\text{Cl}_n \otimes_{\mathbb{C}} \mathbb{C}\text{Cl}_2 \simeq \mathbb{C}\text{Cl}_{n+2},$$

so that with  $\mathbb{C}\text{Cl}_2 \simeq M_2(\mathbb{C})$  we find that  $\mathbb{C}\text{Cl}_n$  is Morita equivalent to  $\mathbb{C}\text{Cl}_{n+2}$ .

The (semi)simple structure of  $\mathbb{C}\text{Cl}_n$  is further clarified by

**Definition 4.5** The *chirality operator*  $\gamma_{n+1}$  in  $\mathbb{C}\text{Cl}_n$  is defined as the element

$$\gamma_{n+1} = (-i)^m e_1 \cdots e_n,$$

where  $n = 2m$  or  $n = 2m + 1$ , depending on whether  $n$  is even or odd.

**Exercise 4.5** Show that

- (1) if  $n = 2m$  is even, then  $\gamma_{n+1}$  generates the center of  $\mathbb{C}\text{Cl}_n^0 = \mathbb{C}\text{Cl}_{n-1}$ ,
- (2) if  $n = 2m + 1$  is odd, then  $\gamma_{n+1}$  lies in the odd part  $\mathbb{C}\text{Cl}_{2m+1}^1$ , and the center of  $\mathbb{C}\text{Cl}_n$  is spanned by 1 and  $\gamma_{n+1}$ .

### 4.1.1 Representation Theory of Clifford Algebras

We determine the irreducible representations of the Clifford algebras  $\mathbb{C}\text{Cl}_n^\pm$  and  $\mathbb{C}\text{Cl}_n$ . Let us start with the complex Clifford algebras.

**Proposition 4.6** *The irreducible representations of  $\mathbb{C}\text{Cl}_n$  are given as*

$$\begin{aligned} & \mathbb{C}^{2^m}; \quad (n = 2m), \\ & \mathbb{C}^{2^m}, \mathbb{C}^{2^m}; \quad (n = 2m + 1). \end{aligned}$$

**Proof** Since the  $\mathbb{C}\text{Cl}_n$  are matrix algebras we can invoke Lemma 2.15 to conclude that in the even-dimensional case the irreducible representation of  $\mathbb{C}\text{Cl}_{2m} \simeq M_{2^m}(\mathbb{C})$  is given by the defining representation  $\mathbb{C}^{2^m}$ . In the odd-dimensional case we have

$$\mathbb{C}\text{Cl}_{2m+1} \simeq M_{2^m}(\mathbb{C}) \oplus M_{2^m}(\mathbb{C}),$$

so that the irreducible representations are given by two copies of  $\mathbb{C}^{2^m}$ , corresponding to the two summands in this matrix algebra.  $\square$

For the real Clifford algebras  $\mathbb{C}\text{Cl}_n^\pm$  we would like to obtain the irreducible representations from those just obtained for the complexification  $\mathbb{C}\text{Cl}_n \simeq \mathbb{C}\text{Cl}_n^\pm \otimes_{\mathbb{R}} \mathbb{C}$ . As  $\mathbb{C}\text{Cl}_n^\pm$  are matrix algebras over  $\mathbb{R}$  and  $\mathbb{H}$ , this leads us to the following possibilities:

- (1) Restrict an (irreducible) representation of  $\mathbb{C}\text{Cl}_n$  to a real subspace, stable under  $\mathbb{C}\text{Cl}_n^\pm$ ;
- (2) Extend an (irreducible) representation of  $\mathbb{C}\text{Cl}_n$  to a quaternionic space, carrying a representation of  $\mathbb{C}\text{Cl}_n^\pm$ .

This is very similar to our approach to real algebras in Sect. 3.3. In fact, we will use an anti-linear map  $J_n^\pm$  on the representation space, furnishing it with a real ( $(J_n^\pm)^2 = 1$ ) or quaternionic structure ( $(J_n^\pm)^2 = -1$ ) to select the real subalgebra  $\mathbb{C}\text{Cl}_n^\pm \subset \mathbb{C}\text{Cl}_n$ . For the even-dimensional case we search for operators  $J_{2m}^\pm$  such that on the irreducible  $\mathbb{C}\text{Cl}_{2m}$ -representations  $\mathbb{C}^{2^m}$  we have

$$\mathbb{C}\text{Cl}_{2m}^\pm \simeq \{a \in \mathbb{C}\text{Cl}_{2m} : [J_{2m}^\pm, a] = 0\}. \quad (4.1.4)$$

The odd case is slightly more subtle, as only the even part  $(\mathbb{C}\text{Cl}_n^\pm)^0$  of  $\mathbb{C}\text{Cl}_n^\pm$  can be recovered in this way:

$$(\mathbb{C}\text{Cl}_{2m+1}^\pm)^0 \simeq \{a \in \mathbb{C}\text{Cl}_{2m+1}^0 : [J_{2m+1}^\pm, a] = 0\}. \quad (4.1.5)$$

**Proposition 4.7** For any  $m \geq 1$  there exist anti-linear operators  $J_{2m}^\pm : \mathbb{C}^{2^m} \rightarrow \mathbb{C}^{2^m}$  and  $J_{2m+1}^\pm : \mathbb{C}^{2^m} \rightarrow \mathbb{C}^{2^m}$  such that the Eqs. (4.1.4) and (4.1.5) hold.

**Proof** From Proposition 4.2 and Exercise 4.3 we see that  $(\text{Cl}_{2m+1}^\pm)^0 \simeq \text{Cl}_{2m}^-$  and  $(\text{Cl}_{2m+1})^0 \simeq \text{Cl}_{2m}$  so that the odd case follows from the even case.

By periodicity we can further restrict to construct only  $J_{2m}^\pm$  for  $m = 1, 2, 3, 4$ . For  $m = 1$  we select the real form  $\text{Cl}_2^+ \simeq M_2(\mathbb{R})$  in  $\text{Cl}_2 \simeq M_2(\mathbb{C})$  as the commutant of  $J_2^+$  with

$$\begin{aligned} J_2^+ : \mathbb{C}^2 &\rightarrow \mathbb{C}^2; \\ \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} &\mapsto \begin{pmatrix} \bar{v}_1 \\ \bar{v}_2 \end{pmatrix}. \end{aligned}$$

Instead, as in Example 3.16,  $\text{Cl}_2^- \simeq \mathbb{H}$  can be identified as a real subalgebra  $\text{Cl}_2 \simeq M_2(\mathbb{C})$  with the commutant of  $J_2^-$ , where

$$\begin{aligned} J_2^- : \mathbb{C}^2 &\rightarrow \mathbb{C}^2; \\ \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} &\mapsto \begin{pmatrix} -\bar{v}_2 \\ \bar{v}_1 \end{pmatrix}. \end{aligned}$$

For  $m = 2$  the sought-for operator  $J_4^+ \equiv J_4^-$  on  $\mathbb{C}^4$  is given by  $J_2^- \oplus J_2^-$ .

For  $m = 3$  we set  $J_6^+ = (J_2^-)^{\oplus 4}$  to select  $\text{Cl}_6^+ \simeq M_4(\mathbb{H})$  inside  $\text{Cl}_6$ , and  $J_6^- = (J_2^+)^{\oplus 4}$  to select  $\text{Cl}_6^- \simeq M_8(\mathbb{R})$ .

Finally, for  $m = 4$  the operator  $J_8^+ \equiv J_8^- := (J_2^+)^{\oplus 8}$  selects the two isomorphic real forms  $\text{Cl}_8^\pm \subset \text{Cl}_8$ .  $\square$

The signs for the squares  $(J_n^\pm)^2$  are listed in Table 4.2. The isomorphisms between the odd- and even-dimensional cases are illustrated by the fact that

$$(J_{2m+1}^\pm)^2 = (J_{2m}^-)^2.$$

with periodicity eight. We also indicated the commutation between  $J_n^\pm$  and odd elements in  $\text{Cl}_n^\pm$  and between  $J_n^\pm$  and the chirality operator  $\gamma_{n+1}$ . For the derivation

**Table 4.2** The real and quaternionic structures on the irreducible representations of  $\text{Cl}_n$  that select  $\text{Cl}_n^\pm$  via (4.1.4) for  $n$  even and  $(\text{Cl}_n^\pm)^0$  via (4.1.5) for  $n$  odd. For later reference, we also indicated the commutation or anti-commutation of  $J_n^-$  with the chirality operator  $\gamma_{n+1}$  defined in Definition 4.5 and odd elements in  $(\text{Cl}_n^\pm)^1 \subset \text{Cl}_n^\pm$

$n$	1	2	3	4	5	6	7	8
$(J_n^+)^2 = \pm 1$	1	1	-1	-1	-1	-1	1	1
$(J_n^-)^2 = \pm 1$	1	-1	-1	-1	-1	1	1	1
$J_n^- x = (\pm 1)x J_n^-$ , $x$ odd	-1	1	1	1	-1	1	1	1
$J_n^- \gamma_{n+1} = (\pm 1)\gamma_{n+1} J_n^-$		-1		1		-1		1

of the former note that for  $n$  even  $J_n^\pm$  commutes with all elements in  $\text{Cl}_n^\pm$ , whereas for  $n$  odd we follow the proof of Proposition 4.7:

- $n = 1$ :  $J_1^-$  is equal to  $J_0^-$ , which is given by  $J_0^-(z) = \bar{z}$  for  $z \in \mathbb{C}$ , and (4.1.5) selects  $(\text{Cl}_1^-)^0 \simeq \mathbb{R}$  in  $\text{Cl}_1^- \simeq \mathbb{C}$ . Thus, the remaining part  $(\text{Cl}_1^-)^1 \simeq i\mathbb{R}$  so that odd elements  $x \in (\text{Cl}_1^-)^1$  anti-commute with  $J_1^-$ .
- $n = 3$ :  $J_3^-$  is equal to  $J_2^-$ , which is given by the standard quaternionic structure on  $\mathbb{C}^2$ . It then follows that all of  $\text{Cl}_3^- \simeq \mathbb{H} \oplus \mathbb{H}$  commutes with  $J_3^-$ .
- $n = 5$ : in this case  $J_5^-$  is equal to  $J_4^-$ , which is two copies of  $J_2^-$ . This selects  $(\text{Cl}_5^-)^0 \simeq M_2(\mathbb{H})$  in  $\text{Cl}_5^- \simeq M_4(\mathbb{C})$ . Again, the remaining part  $(\text{Cl}_5^-)^1 \simeq iM_2(\mathbb{H})$  so that odd elements  $x \in (\text{Cl}_5^-)^0$  anti-commute with  $J_5^-$ .
- $n = 7$ :  $J_7^-$  is equal to  $J_6^-$ , which is given by component-wise complex conjugation of vectors in  $\mathbb{C}^8$ . It follows that all of  $\text{Cl}_7^- \simeq M_8(\mathbb{R}) \oplus M_8(\mathbb{R})$  commutes with  $J_6^-$ .

Finally, in the even case  $n = 2m$  the (anti)-commutation between the chirality operator  $\gamma_{n+1}$  and the *anti-linear* operator  $J_n^-$  depends only on the power of the factor  $i^m$ . Indeed, the even product of  $e_i$ 's in Definition 4.5 already commutes with  $J_n^-$ , so that the signs  $(-1)^m$  for  $n = 2m$  follow from

$$J_n^- i^m = (-i)^m J_n^-.$$

The last three rows of Table 4.2 give precisely the sign table that appears for real spectral triples below, where  $n$  is the corresponding KO-dimension, and hence coincide with Table 3.1 of Definition 3.1. We will now slowly move to the spin manifold case, tracing KO-dimension back to its historical roots.

## 4.2 Riemannian Spin Geometry

We here give a concise introduction to Riemannian spin manifolds and work towards a Dirac operator. For convenience, we restrict to compact manifolds.

### 4.2.1 Spin Manifolds

The definition of Clifford algebras can be extended to Riemannian manifolds, as we will now explain. First, for completeness we recall the definition of a Riemannian metric on a manifold.

**Definition 4.8** A *Riemannian metric* on a manifold  $M$  is a symmetric bilinear form on (smooth) vector fields  $\Gamma^\infty(TM)$

$$g : \Gamma^\infty(TM) \times \Gamma^\infty(TM) \rightarrow C^\infty(M)$$

such that

- (1)  $g(X, Y)$  is a real function if  $X$  and  $Y$  are real vector fields;
- (2)  $g$  is  $C^\infty(M)$ -bilinear:

$$g(fX, Y) = g(X, fY) = fg(X, Y); \quad (f \in C^\infty(M));$$

- (3)  $g(X, X) \geq 0$  for all real vector fields  $X$  and  $g(X, X) = 0$  if and only if  $X = 0$ .

The non-degeneracy condition (3) allows us to identify  $\Gamma^\infty(TM)$  with  $\Omega_{\text{dR}}^1(M) = \Gamma^\infty(T^*M)$ .

A Riemannian metric  $g$  on  $M$  gives rise to a **distance function** on  $M$ , given as an infimum of path lengths

$$d_g(x, y) = \inf_{\gamma} \left\{ \int_0^1 \sqrt{g(\dot{\gamma}(t), \dot{\gamma}(t))} dt : \gamma(0) = x, \gamma(1) = y \right\}. \quad (4.2.1)$$

Moreover, the inner product that  $g$  defines on the fibers  $T_x M$  of the tangent bundle allows us to define Clifford algebras at each point in  $M$  as follows. With the inner product at  $x \in M$  given explicitly by  $g_x(X_x, Y_x) := g(X, Y)|_x$  we consider the quadratic form on  $T_x M$  defined by

$$Q_g(X_x) = g_x(X_x, X_x).$$

We can then apply the construction of the Clifford algebra of the previous section to each fiber of the tangent bundle. At each point  $x \in M$  this gives rise to  $\text{Cl}(T_x M, Q_g)$  and its complexification  $\mathbb{C}\text{I}(T_x M, Q_g)$ . When  $x$  varies, these Clifford algebras combine to give a bundle of algebras.

**Definition 4.9** The *Clifford algebra bundle*  $\text{Cl}^+(TM)$  is the bundle of algebras  $\text{Cl}(T_x M, Q_g)$ , with the transition functions inherited from  $TM$ . Namely, transition functions on the tangent bundle are given for open  $U, V \subset M$  by  $t_{UV} : U \cap V \rightarrow SO(n)$  where  $n = \dim M$ . Their action on each fiber  $T_x M$  can be extended to  $\text{Cl}(T_x M, Q_g)$  by

$$v_1 v_2 \cdots v_k \mapsto t_{UV}(v_1) \cdots t_{UV}(v_k); \quad (v_1, \dots, v_k \in T_x M).$$

The algebra of smooth real-valued sections of  $\text{Cl}^+(TM)$  will be denoted by  $\text{Cliff}^+(M) = \Gamma^\infty(\text{Cl}^+(TM))$ .

Similarly, replacing  $Q_g$  by  $-Q_g$ , we define  $\text{Cliff}^-(M)$  as the space of sections of  $\text{Cl}^-(TM)$ .

Finally, we define the complexified algebra

$$\mathbb{C}\text{Iff}(M) := \text{Cliff}^+(M) \otimes_{\mathbb{R}} \mathbb{C},$$

consisting of smooth sections of the bundle of complexified algebras  $\mathbb{C}\text{I}(TM)$ , which is defined in a similar manner.

Let us determine local expressions for the algebra  $\text{Cliff}^+(M)$ . If  $\{x^\mu\}_{\mu=1}^n$  are local coordinates on a chart  $U$  of  $M$ , the algebra of sections of  $\text{Cliff}^+(M)|_U$  is generated by  $\gamma_\mu$  with relations

$$\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu = 2g_{\mu\nu}, \quad (4.2.2)$$

with  $g_{\mu\nu} = g(\partial_\mu, \partial_\nu)$ . After choosing an orthonormal frame for  $\Gamma^\infty(TM)|_U$  with respect to the metric  $g$ , at a point of  $U$  this relation reduces precisely to the relation (4.1.2).

Let us see if we can import more of the structure for Clifford algebras explored so far to the setting of a Riemannian manifold. First, recall that

$$\mathbb{C}\text{l}_{2m} \cong M_{2^m}(\mathbb{C}), \quad \mathbb{C}\text{l}_{2m+1}^0 \cong M_{2^m}(\mathbb{C}).$$

Another way of phrasing this is to say that the (even parts of the) Clifford algebras  $\mathbb{C}\text{l}_n$  are endomorphism algebras  $\text{End}(\mathbb{C}^{2^m})$ . The natural question that arises in the setting of Riemannian manifolds is whether or not this holds for all fibers of the Clifford algebra bundle, in which case it would extend to a global isomorphism of algebra bundles.

**Definition 4.10** A Riemannian manifold is called *spin<sup>c</sup>* if there exists a vector bundle  $S \rightarrow M$  such that there is an algebra bundle isomorphism

$$\begin{aligned} \mathbb{C}\text{l}(TM) &\simeq \text{End}(S) \quad (M \text{ even-dimensional}), \\ \mathbb{C}\text{l}(TM)^0 &\simeq \text{End}(S) \quad (M \text{ odd-dimensional}). \end{aligned}$$

The pair  $(M, S)$  is called a *spin<sup>c</sup> structure* on  $M$ .

If a spin<sup>c</sup> structure  $(M, S)$  exists we refer to  $S$  as the **spinor bundle** and the sections in  $\Gamma^\infty(S)$  as **spinors**. Using the metric and the action of  $\text{Cliff}^+(M)$  by endomorphisms on  $\Gamma^\infty(S)$  we introduce the following notion.

**Definition 4.11** Let  $(M, S)$  be a spin<sup>c</sup> structure on  $M$ . *Clifford multiplication* is defined by the linear map

$$\begin{aligned} c : \Omega_{\text{dR}}^1(M) \times \Gamma^\infty(S) &\rightarrow \Gamma^\infty(S); \\ (\omega, \psi) &\mapsto \omega^\# \cdot \psi, \end{aligned}$$

where  $\omega^\#$  is the vector field in  $\Gamma^\infty(TM)$  corresponding to the one-form  $\omega \in \Omega_{\text{dR}}^1(M)$  via the metric  $g$ . This vector field acts as an endomorphism on  $\Gamma^\infty(S)$  via the embedding  $\Gamma^\infty(TM) \hookrightarrow \text{Cliff}^+(M) \subset \Gamma^\infty \text{End}(S)$ .

In local coordinates on  $U \subset M$ , we can write  $\omega|_U = \omega_\mu dx^\mu$  with  $\omega_\mu \in C^\infty(U)$  so that Clifford multiplication can be written as

$$c(\omega)\psi|_U \equiv c(\omega, \psi)|_U = \omega_\mu (\gamma^\mu \psi)|_U; \quad (\psi \in \Gamma^\infty(S)),$$

with  $\gamma^\mu = g^{\mu\nu}\gamma_\nu$  and  $\gamma_\nu$  as in (4.2.2) but now represented as endomorphisms on the fibers of  $S$ . The appearance of  $\gamma^\mu$  comes from the identification of the basis covector  $dx^\mu \in \Omega_{\text{dr}}^1(M)|_U$  with the basis vector  $\partial_\mu \in \Gamma^\infty(TM)|_U$  using the metric, which is then embedded in  $\text{Cliff}^+(M)$ . That is, we have

$$dx_p^\mu = g(\partial_\mu, \cdot)_p$$

as (non-degenerate) maps from  $T_p M$  to  $\mathbb{C}$  with  $p \in U \subset M$ .

Recall that if  $M$  is compact, then any vector bundle carries a smoothly varying inner product on its fibers,

$$\langle \cdot, \cdot \rangle : \Gamma^\infty(S) \times \Gamma^\infty(S) \rightarrow C^\infty(M).$$

**Exercise 4.6** Use a partition of unity argument to show that any vector bundle on a compact manifold  $M$  admits a smoothly varying inner product on its fibers.

**Definition 4.12** The *Hilbert space of square-integrable spinors*  $L^2(S)$  is defined as the completion of  $\Gamma^\infty(S)$  in the norm corresponding to the inner product

$$(\psi_1, \psi_2) = \int_M \langle \psi_1, \psi_2 \rangle(x) \sqrt{\det g} dx,$$

where  $\sqrt{\det g} dx$  is the Riemannian volume form.

Recall that in the previous subsection we selected the real Clifford algebras  $\text{Cl}_n^\pm$  as subalgebras in  $\text{Cl}_n$  that commute with a certain anti-linear operator  $J_n^\pm$ . We now try to select  $\text{Cliff}^\pm(M) \subset \text{Cliff}(M)$ , considered as endomorphisms on  $\Gamma^\infty(S)$ , through a globally-defined operator  $J_M : \Gamma^\infty(S) \rightarrow \Gamma^\infty(S)$ , so that

$$(J_M \psi)(x) = J_n^\pm(\psi(x)),$$

for any section  $\psi \in \Gamma^\infty(S)$ , where  $n = \dim M$ . Such a global operator does not always exist: this gives rise to the notion of a spin manifold. It is conventional to work with  $J_n^-$  to select  $\text{Cliff}^-(M) \subset \text{Cliff}(M)$ , making our sign Table 4.2 fit with the usual definition of KO-dimension in noncommutative geometry.

**Definition 4.13** A Riemannian  $\text{spin}^c$  manifold is called *spin* if there exists an anti-unitary operator  $J_M : \Gamma^\infty(S) \rightarrow \Gamma^\infty(S)$  such that:

- (1)  $J_M$  commutes with the action of real-valued smooth functions on  $\Gamma^\infty(S)$ ;
- (2)  $J_M$  commutes with  $\text{Cliff}^-(M)$  (or with  $\text{Cliff}^-(M)^0$  in the odd case).

We call the pair  $(S, J_M)$  a *spin structure* on  $M$  and refer to the operator  $J_M$  as the *charge conjugation*.

If the manifold  $M$  is even dimensional, we can define a grading

$$(\gamma_M \psi)(x) = \gamma_{n+1}(\psi(x)); \quad (\psi \in \Gamma^\infty(S)).$$

Then, the sign rules of Table 4.2 for the square of  $J_n^-$  and the (anti)-commutation of  $J_n^-$  with  $\gamma_{n+1}$  and odd elements in  $\text{Cl}_n^-$  hold in each fiber of  $\Gamma^\infty(S)$ . Hence, we find that also globally

$$J_M^2 = \epsilon, \quad J_M x = \epsilon' x J_M; \quad (x \in (\text{Cliff}^-(M))^1), \quad J_M \gamma_M = \epsilon'' \gamma_M J_M,$$

with  $\epsilon, \epsilon', \epsilon'' \in \{\pm 1\}$  being the signs in Table 4.2 with  $n = \dim M$  modulo eight. This will be crucial for our definition of a real spectral triple in the next section, where these signs determine the KO-dimension of a noncommutative Riemannian spin manifold.

#### 4.2.2 Clifford Connections, Spin Connections and the Dirac Operator

The presence of a spin structure on a Riemannian manifold allows for the construction of a first-order differential operator that up to a scalar term squares to the Laplacian associated to  $g$ . This is the same operator that Dirac searched for (with success) in his attempt to replace the Schrödinger equation by a more general covariant differential equation in Minkowski space. The Dirac operator that we will describe below is the analogue for Riemannian spin manifolds of Dirac's operator on flat Minkowski space.

**Definition 4.14** A *connection* on a vector bundle  $E \rightarrow M$  is given by a  $\mathbb{C}$ -linear map on the space of smooth sections:

$$\nabla : \Gamma^\infty(E) \rightarrow \Omega_{\text{dR}}^1(M) \otimes_{C^\infty(M)} \Gamma^\infty(E)$$

that satisfies the Leibniz rule

$$\nabla(f\eta) = f\nabla(\eta) + df \otimes \eta; \quad (f \in C^\infty(M), \eta \in \Gamma^\infty(E)).$$

The *curvature*  $\Omega^E$  of  $\nabla$  is defined by the  $C^\infty(M)$ -linear map

$$\Omega^E := \nabla^2 : \Gamma^\infty(E) \rightarrow \Omega^2(M) \otimes_{C^\infty(M)} \Gamma^\infty(E).$$

Finally, if  $\langle \cdot, \cdot \rangle$  is a smoothly varying (i.e.,  $C^\infty(M)$ -valued) inner product on  $\Gamma^\infty(E)$ , a connection is said to be *hermitian*, or *compatible* if

$$-\langle \nabla \eta, \eta' \rangle + \langle \eta, \nabla \eta' \rangle = d\langle \eta, \eta' \rangle; \quad (\eta, \eta' \in \Gamma^\infty(E)).$$

Equivalently, when evaluated on a vector field  $X \in \Gamma^\infty(TM)$  a connection gives rise to a map

$$\nabla_X : \Gamma^\infty(E) \rightarrow \Gamma^\infty(E).$$

More precisely, the relation with the above definition is given by

$$\nabla_X(\eta) := \nabla(\eta)(X),$$

for all  $X \in \Gamma^\infty(TM)$  and  $\eta \in \Gamma^\infty(E)$ . The corresponding curvature then becomes

$$\Omega^E(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X, Y]}, \quad (X, Y \in \Gamma^\infty(TM)), \quad (4.2.3)$$

i.e., it is a measure of the defect of  $\nabla$  to be a Lie algebra map.

**Example 4.15** Consider the tangent bundle  $TM \rightarrow M$  on a Riemannian manifold  $(M, g)$ . A classical result is that there is a unique connection on  $TM$  that is compatible with the inner product  $g$  on  $\Gamma(TM)$ , i.e.

$$\langle \nabla_X Y, Z \rangle + \langle Y, \nabla_X Z \rangle = X(\langle Y, Z \rangle)$$

and that is torsion-free, i.e.

$$\nabla_X Y - \nabla_Y X = [X, Y]; \quad (X, Y \in \Gamma^\infty(TM)).$$

This connection is called the *Levi–Civita connection* and can be written in local coordinates  $\{x^\mu\}_{\mu=1}^n$  on a chart  $U \subset M$  as  $\nabla(\partial_\nu) = \Gamma_{\mu\nu}^\kappa dx^\mu \otimes \partial_\kappa$ , or

$$\nabla_{\partial_\mu}(\partial_\nu) = \Gamma_{\mu\nu}^\kappa \partial_\kappa.$$

The  $C^\infty(U)$ -valued coefficients  $\Gamma_{\mu\nu}^\kappa$  are the so-called *Christoffel symbols* and torsion-freeness corresponds to the symmetry  $\Gamma_{\mu\nu}^\kappa = \Gamma_{\nu\mu}^\kappa$ .

Recall also the definition of the *Riemannian curvature tensor* on  $(M, g)$  as the curvature of the Levi–Civita connection, i.e.

$$R(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X, Y]} \in \Gamma(\text{End } TM),$$

which is indeed a  $C^\infty(M)$ -linear map. Locally, we have for its components

$$R_{\mu\nu\kappa\lambda} := g(\partial_\mu, R(\partial_\kappa, \partial_\lambda)\partial_\nu).$$

The contraction  $R_{\nu\lambda} := g^{\mu\kappa} R_{\mu\nu\kappa\lambda}$  is called the *Ricci tensor*, and the subsequent contraction  $s := g^{\nu\lambda} R_{\nu\lambda} \in C^\infty(M)$  is the *scalar curvature*.

Similar results hold for the cotangent bundle, with the unique, compatible, torsion-free connection thereon related to the above via the metric  $g$ .

**Definition 4.16** If  $\nabla^E$  is a connection on a vector bundle  $E$ , the *Laplacian* associated to  $\nabla^E$  is the second order differential operator on  $E$  defined by

$$\Delta^E := -\text{Tr}_g(\nabla \otimes 1 + 1 \otimes \nabla^E) \circ \nabla^E : \Gamma^\infty(E) \rightarrow \Gamma^\infty(E),$$

where

$$\begin{aligned} \nabla \otimes 1 + 1 \otimes \nabla^E : \Omega_{\text{dR}}^1(M) \otimes_{C^\infty(M)} \Gamma^\infty(E) \\ \rightarrow \Omega_{\text{dR}}^1(M) \otimes_{C^\infty(M)} \Omega_{\text{dR}}^1(M) \otimes_{C^\infty(M)} \Gamma^\infty(E) \end{aligned}$$

is the combination of the Levi–Civita connection on the cotangent bundle with the connection  $\nabla^E$  and  $\text{Tr}_g$  is the trace associated to  $g$  mapping  $\Omega_{\text{dR}}^1(M) \otimes_{C^\infty(M)} \Omega_{\text{dR}}^1(M) \rightarrow C^\infty(M)$ .

Locally, we find

$$\Delta^E = -g^{\mu\nu}(\nabla_\mu^E \nabla_\nu^E - \Gamma_{\mu\nu}^\kappa \nabla_\kappa^E).$$

If  $M$  is a Riemannian spin<sup>c</sup> manifold, then the above Levi–Civita connection can be lifted to the spinor bundle. First, choose a local orthonormal frame for  $TM|_U$ :

$$\{E_1, \dots, E_n\} \text{ for } \Gamma(TM)|_U : \quad g(E_a, E_b) = \delta_{ab}.$$

The corresponding dual orthonormal frame of  $T^*M|_U$  is denoted by  $\theta^a$ . We can then write the Christoffel symbols in this basis, namely by

$$\nabla E_a =: \tilde{\Gamma}_{\mu a}^b dx^\mu \otimes E_b$$

on vector fields, and on one-forms by

$$\nabla \theta^b = -\tilde{\Gamma}_{\mu a}^b dx^\mu \otimes \theta^a.$$

The compatibility of  $\nabla$  with the inner product  $g$  implies the skew-symmetry of  $\tilde{\Gamma}_{\mu a}^b$  under the exchange of  $a$  and  $b$ .

Also note that the local orthonormal frame for  $TM|_U$  allows us to write Clifford relations for (globally) fixed matrices  $\gamma^a$ :

$$\gamma^a \gamma^b + \gamma^b \gamma^a = 2\delta^{ab}; \quad (a, b = 1, \dots, n). \quad (4.2.4)$$

We now come to lift this structure from the tangent bundle to the spinor bundle. More precisely, one requires the following compatibility between the Levi–Civita connection, Clifford multiplication, and the connection on the spinor bundle.

**Definition 4.17** Let  $M$  be a spin<sup>c</sup> manifold. A *Clifford connection*  $\nabla^S$  on the spinor bundle  $S \rightarrow M$  is a hermitian connection  $\nabla^S$  on the spinor bundle  $S \rightarrow M$  such that

$$\nabla_X^S(c(\omega)\psi) = c(\nabla_X(\omega))\psi + c(\omega)\nabla_X^S(\psi); \quad (4.2.5)$$

for any  $X \in \mathfrak{X}(M)$ ,  $\omega \in \Omega_{\text{dR}}^1(M)$ ,  $\psi \in \Gamma^\infty(S)$ . Here  $\nabla$  is the Levi–Civita connection on the cotangent bundle.

We also have the following concrete formula.

**Lemma 4.18** *Let  $M$  be a  $\text{spin}^c$  manifold. Then the following local formula defines a Clifford connection on the spinor bundle:*

$$\nabla_\mu^S \psi(x) = \left( \partial_\mu - \frac{1}{4} \tilde{\Gamma}_{\mu a}^b \gamma^a \gamma_b \right) \psi(x).$$

*Any other Clifford connection on  $S$  is of the form  $\nabla^S + \alpha$  where  $\alpha = -\alpha^*$  is a purely imaginary one-form.*

**Proof** Take  $X = \partial_\mu$  in local coordinates  $\{x^\mu\}$  on  $U$  and take  $\omega = \theta^c$  with respect to an orthonormal frame  $\{\theta^c\}$  for  $T^*M|_U$ . Then  $c(\omega) = \gamma^c$  and we find for any  $\psi \in \Gamma^\infty(S)$  that

$$\begin{aligned} \nabla_{\partial_\mu}^S (\gamma^c \psi) - \gamma^c \nabla_{\partial_\mu}^S \psi &= -\frac{1}{4} \tilde{\Gamma}_{\mu a}^b \gamma^a \gamma_b \gamma^c \psi + \frac{1}{4} \tilde{\Gamma}_{\mu a}^b \gamma^c \gamma^a \gamma_b \psi \\ &= -\frac{1}{4} \tilde{\Gamma}_{\mu a}^b \gamma^a \gamma_b \gamma^c \psi + \frac{1}{4} \tilde{\Gamma}_{\mu a}^b (2\delta^{ca} \gamma_b - 2\delta_b^c \gamma^a + \gamma^a \gamma_b \gamma^c) \psi \\ &= \left( \frac{1}{2} \tilde{\Gamma}_{\mu c}^b - \frac{1}{2} \tilde{\Gamma}_{\mu a}^c \gamma^a \right) \psi \\ &= -\tilde{\Gamma}_{\mu a}^c \gamma^a \psi \end{aligned}$$

using the relations (4.2.4) and skew-symmetry of  $\tilde{\Gamma}_{\mu c}^b$  under the exchange of  $b$  and  $c$ . By definition of the Christoffel symbols  $\tilde{\Gamma}$  we also have that  $c(\nabla_{\partial_\mu}(\theta^c))\psi = -\tilde{\Gamma}_{\mu a}^c \gamma^a \psi$  so that the compatibility (4.2.5) is satisfied. The skew-symmetric property of  $\tilde{\Gamma}_{\mu c}^b$  combines with hermiticity of  $\gamma^a$  to yield hermiticity of  $\nabla^S$  and this completes the proof of the first statement.

If  $\tilde{\nabla}$  is another connection on  $S$  we can always write  $\tilde{\nabla} = \nabla + \alpha$  where  $\alpha \in \text{End}_{C^\infty(M)}(\Gamma^\infty(S)) \otimes_{C^\infty(M)} \Omega_{\text{dR}}^1(M)$ . For this connection to be Clifford we need  $\alpha(\partial_\mu) \in \text{End}_{C^\infty(M)}(\Gamma^\infty(S))$  to commute with all  $c(\omega)$ . Since  $M$  is  $\text{spin}^c$  we have  $\text{End}_{C^\infty(M)}(\Gamma^\infty(S)) \cong \mathbb{C}\text{liff}(M)^{(0)}$ . From this we derive that at each point  $x \in M$  the linear map  $\alpha(\partial_\mu)_x$  should be a scalar multiple of the identity. Hermiticity of  $\tilde{\nabla}$  then implies that  $\alpha(\partial_\mu) \in C^\infty(M, i\mathbb{R})$ .  $\square$

We will call the above Clifford connection  $\nabla^S$  on  $S$  the **spin connection**.

**Proposition 4.19** *If  $M$  is a spin manifold and  $J_M$  is the corresponding anti-unitary operator on  $\Gamma(S)$ , then the spin connection  $\nabla^S$  is the unique Clifford connection that commutes with  $J_M$ .*

**Proof** Observe that the product  $\gamma^a \gamma^b = -(i\gamma^a)(i\gamma^b)$  is in the even part of the Clifford algebra  $\text{Cl}_n^-$ , since

$$(i\gamma^a)(i\gamma^b) + (i\gamma^b)(i\gamma^a) = -2\delta^{ab}.$$

Since by definition the operator  $J_n^-$  commutes with the even elements in  $\text{Cl}_n^-$  acting fiberwise on the spinor bundle, it follows that  $\nabla^S$  commutes with  $J_M$ .

Since any other Clifford connection differs from  $\nabla^S$  by the addition of a purely imaginary one-form  $\alpha$ , commutation with  $J_M$  implies that  $\alpha = 0$ .  $\square$

All of the above structure culminates in the following definition

**Definition 4.20** Let  $M$  be a spin manifold, with spin structure  $(S, J_M)$ . The *Dirac operator*  $D_M$  is the composition of the spin connection on  $S$  with Clifford multiplication of Definition 4.11:

$$D_M : \Gamma^\infty(S) \xrightarrow{\nabla^S} \Omega_{\text{dR}}^1(M) \otimes_{C^\infty(M)} \Gamma^\infty(S) \xrightarrow{-ic} \Gamma^\infty(S).$$

In local coordinates, we have

$$D_M \psi(x) = -i\gamma^\mu \left( \partial_\mu - \frac{1}{4} \tilde{\Gamma}_{\mu a}^b \gamma^a \gamma_b \right) \psi(x).$$

### 4.2.3 Lichnerowicz Formula

Let us come back to the original motivation of Dirac, which was to find an operator whose square is the Laplacian. Up to a scalar this continues to hold for the Dirac operator on a Riemannian spin manifold, a result that will turn out to be very useful later on in our physical applications. For this reason we include it here with proof.

**Theorem 4.21** Let  $(M, g)$  be a Riemannian spin manifold with Dirac operator  $D_M$ . Then

$$D_M^2 = \Delta^S + \frac{1}{4}s,$$

in terms of the Laplacian  $\Delta^S$  associated to the spin connection  $\nabla^S$  and the scalar curvature  $s$ .

**Proof** We exploit the local expressions for  $D_M$ ,  $\Delta^S$  and  $s$ , as the above formula is supposed to hold in each chart that trivializes  $S$ . With  $D_M = -i\gamma^\mu \nabla_\mu^S$  we compute

$$\begin{aligned} D_M^2 &= -\gamma^\mu \nabla_\mu^S \gamma^\nu \nabla_\nu^S = -\gamma^\mu \gamma^\nu \nabla_\mu^S \nabla_\nu^S - \gamma^\mu c(\nabla_\mu dx^\kappa) \nabla_\kappa^S \\ &= -\gamma^\mu \gamma^\nu (\nabla_\mu^S \nabla_\nu^S - \Gamma_{\mu\nu}^\kappa \nabla_\kappa^S). \end{aligned}$$

We then use the Clifford relations (4.2.2) to write  $\gamma^\mu \gamma^\nu = \frac{1}{2}[\gamma^\mu, \gamma^\nu] + g^{\mu\nu}$ , and combine this with torsion freedom  $\Gamma_{\mu\nu}^\kappa = \Gamma_{\nu\mu}^\kappa$  to obtain

$$D_M^2 = -g^{\mu\nu} (\nabla_\mu^S \nabla_\nu^S - \Gamma_{\mu\nu}^\kappa \nabla_\kappa^S) - \frac{1}{2} [\gamma^\mu, \gamma^\nu] \nabla_\mu^S \nabla_\nu^S \equiv \Delta^S - \frac{1}{2} \gamma^\mu \gamma^\nu R^S(\partial_\mu, \partial_\nu),$$

in terms of the Laplacian for  $\nabla^S$  on  $S$  and the curvature  $R^S$  thereof. The latter is given by  $-\frac{1}{4}R_{\kappa\lambda,\mu\nu}\gamma^\kappa\gamma^\lambda$ , as one can easily compute from the explicit local form of  $\nabla^S$  in Definition 4.17. Thus,

$$D_M^2 = \Delta^S - \frac{1}{8}R_{\mu\nu\kappa\lambda}\gamma^\mu\gamma^\nu\gamma^\kappa\gamma^\lambda.$$

Using the cyclic symmetry of the Riemann curvature tensor in the last three indices, and the Clifford relations (4.2.2) we find that the second term on the right-hand side is equal to  $\frac{1}{4}R_{\nu\lambda}g^{\nu\lambda} = \frac{1}{4}s$ , in terms of the scalar curvature defined in Example 4.15.  $\square$

### 4.3 The Dirac Operator: Analytical Aspects

In this section we will establish a series of key results that forms the starting point for an operator-algebraic formulation of noncommutative Riemannian spin manifolds.

**Theorem 4.22** *Let  $M$  be a compact Riemannian spin manifold (without boundary). The Dirac operator  $D_M$  is essentially self-adjoint on  $\Gamma^\infty(S) \subset L^2(S)$  with compact resolvent  $(i + D_M)^{-1}$ , and has bounded commutators with elements in  $C^\infty(M)$ . In fact*

$$[D_M, f] = -ic(df),$$

so that  $\|[D_M, f]\| = \|f\|_{\text{Lip}}$  is the Lipschitz (semi)-norm of  $f$ :

$$\|f\|_{\text{Lip}} = \sup_{x \neq y} \left\{ \frac{|f(x) - f(y)|}{d_g(x, y)} \right\}.$$

We divide the proof of this Theorem into three parts which we treat in the subsequent subsections: bounded commutators, essential self-adjointness, and compact resolvent.

#### 4.3.1 Bounded Commutators

**Proposition 4.23** *The commutator  $[D_M, f]$  defined on  $\Gamma^\infty(S)$  extends to a bounded operator on  $L^2(S)$ . More precisely, we have  $[D_M, f] = -ic(df)$  and  $\|[D_M, f]\| = \|f\|_{\text{Lip}}$ .*

**Proof** It follows from the Leibniz rule that

$$[D_M, f](\psi) = -ic(dx^\mu)[\nabla_\mu^S, f]\psi = -ic(dx^\mu)(\partial_\mu f) \cdot \psi = -ic(df)\psi$$

where we have assumed that  $\text{supp}(\psi)$  is contained in a chart that trivializes the spinor bundle  $S \rightarrow M$  so we can use the local formula for  $D_M$ .

For the norm of the commutator we compute

$$\|[D_M, f]\|^2 = \|c(df^*)c(df)\| = \sup_{x \in M} |g^{-1}(df^*, df)(x)|.$$

We may write this as  $\|\text{grad } f\|_\infty^2 := \sup_{x \in M} \|g(\text{grad}_x f^*, \text{grad}_x f)\|$  in terms of the gradient vector field defined by  $\text{grad } f := (df)^\sharp$ . We claim that  $\|\text{grad } f\|_\infty = \|f\|_{\text{Lip}}$ .

First, consider a smooth path  $\gamma : [0, 1] \rightarrow M$  such that  $\gamma(0) = x, \gamma(1) = y$ . Then

$$\begin{aligned} f(x) - f(y) &= \int_0^1 \frac{d}{dt} f(\gamma(t)) dt \\ &= \int_0^1 (d_{\gamma(t)} f)(\dot{\gamma}(t)) dt. \\ &= \int_0^1 g_{\gamma(t)}(\text{grad}_{\gamma(t)} f, \dot{\gamma}(t)) dt. \end{aligned}$$

By Cauchy–Schwartz inequality we then have

$$|f(x) - f(y)| \leq \|\text{grad } f\|_\infty l(\gamma)$$

in terms of the Riemannian length  $l(\gamma)$  of the path  $\gamma$ . If we take an infimum over all such paths  $\gamma$  we find that  $|f(x) - f(y)| \leq \|\text{grad } f\|_\infty d_g(x, y)$  so that  $\|f\|_{\text{Lip}} \leq \|\text{grad } f\|_\infty$ .

For the other inequality, suppose instead that there exists  $x \in M$  so that  $\|\text{grad}_x f\| > \|f\|_{\text{Lip}} + \epsilon$  for some  $\epsilon > 0$ . Consider again a smooth path  $\gamma : [0, 1] \rightarrow M$  such that  $\gamma(0) = x$ . Then there exists  $\delta > 0$  such that for all  $0 < t < \delta$  we have

$$\left| \frac{1}{t} (f(\gamma(t)) - f(\gamma(0))) - g_x(\text{grad}_x f, \dot{\gamma}(0)) \right| < \frac{\epsilon}{2}$$

which implies that

$$\left| \frac{1}{t} (f(\gamma(t)) - f(\gamma(0))) \right| > \left| g_x(\text{grad}_x f, \dot{\gamma}(0)) \right| - \frac{\epsilon}{2}$$

Now take a normalized  $\dot{\gamma}(0) = \frac{\text{grad}_x f}{\|\text{grad}_x f\|}$  and parametrize  $\gamma$  naturally so that  $l(\gamma(0)) \rightarrow \gamma(t)) = t$ . Then the above inequality yields

$$\begin{aligned}
|f(\gamma(t)) - f(\gamma(0))| &> \left( \|\text{grad}_x f\| - \frac{\epsilon}{2} \right) t \\
&> \left( \|f\|_{\text{Lip}} + \frac{\epsilon}{2} \right) l(\gamma(0) \rightarrow \gamma(t)) \\
&> \left( \|f\|_{\text{Lip}} + \frac{\epsilon}{2} \right) d_g(\gamma(0), \gamma(t))
\end{aligned}$$

But this implies that  $\|f\|_{\text{Lip}} > \|f\|_{\text{Lip}} + \frac{\epsilon}{2}$  which is a contradiction.  $\square$

### 4.3.2 Essential Self-adjointness

We first show that  $D_M$  is a symmetric operator on  $\Gamma^\infty(S)$  and then apply a general result on essential self-adjointness for symmetric differential operators on compact manifolds without boundary.

**Proposition 4.24** *For all  $\psi_1, \psi_2 \in \Gamma^\infty(S)$  we have*

$$(D_M(\psi_1), \psi_2) = (\psi_1, D_M(\psi_2)).$$

**Proof** First, an application of the hermiticity of the spinor connection yields

$$\begin{aligned}
(D_M(\psi_1), \psi_2) &= +i \int_M \langle c(dx^\mu) \nabla_\mu^S(\psi_1), \psi_2 \rangle \sqrt{\det g} \cdot dx^1 \wedge \cdots \wedge dx^n \\
&= (\psi_1, D_M(\psi_2)) - i \int_M \langle \psi_1, c(\nabla_\mu(dx^\mu)) \psi_2 \rangle \sqrt{\det g} \cdot dx^1 \wedge \cdots \wedge dx^n \\
&\quad + i \int_M \partial_\mu (\langle \psi_1, c(dx^\mu) \psi_2 \rangle) \sqrt{\det g} \cdot dx^1 \wedge \cdots \wedge dx^n
\end{aligned}$$

An argument based on integration by parts shows that it is now sufficient to establish the following expression

$$\nabla_\mu(dx^\mu) \sqrt{\det g} = -\partial_\mu(\sqrt{\det g}) dx^\mu \quad (4.3.1)$$

For the right-hand side we use the det/log relationship:

$$\partial_\mu(\sqrt{\det g}) = \frac{1}{2} \text{Tr}((\partial_\mu g) g^{-1}) \sqrt{\det g}$$

which follows from basic linear algebra and the chain rule. We compute for the left-hand side of (4.3.1) that

$$\nabla_\mu(dx^\mu) = -\Gamma_{\mu\nu}^\mu dx^\nu$$

where the Christoffel symbols are given locally by

$$\Gamma_{\mu\nu}^\kappa = \frac{1}{2} g^{\kappa\lambda} (\partial_\mu g_{\nu\lambda} + \partial_\nu g_{\mu\lambda} - \partial_\lambda g_{\mu\nu}).$$

In other words, we have

$$\Gamma_{\mu\nu}^\mu = \frac{1}{2} g^{\mu\lambda} \partial_\nu (g_{\mu\lambda}) = \frac{1}{2} \text{Tr}(g^{-1} \partial_\nu g),$$

from which it follows that

$$\nabla_\mu(dx^\mu) = -\frac{1}{2} \text{Tr}(g^{-1} \partial_\mu g) dx^\mu$$

so that validity of Eq. (4.3.1) follows.  $\square$

We proceed with the following well-known result (see Note 7 of this chapter below), which is valid for any closable operator  $T$  on a Hilbert space  $\mathcal{H}$  (and so in particular to the symmetric operator  $D_M$ ).

**Lemma 4.25** *Let  $T$  be a closable operator on a Hilbert space  $\mathcal{H}$ . Then  $u \in \mathcal{H}$  belongs to the domain of the closure  $\bar{T}$  of  $T$  if and only if there exists a sequence  $\{u_j\}$  in the domain of  $T$  such that  $u_j \rightarrow u$  and  $\|Tu_j\|$  is bounded.*

**Proof** Let  $\xi \in \text{Dom}(T^*)$ . Then

$$|(u, T^*\xi)| = \lim_{n \rightarrow \infty} |(u_n, T^*\xi)| = \lim_{n \rightarrow \infty} |(Tu_n, \xi)| \leq \lim_{n \rightarrow \infty} \|Tu_n\| \|\xi\|.$$

using Cauchy–Schwartz inequality. Since  $\|Tu_n\|$  is bounded, it follows that  $\xi \rightarrow (u, T^*\xi)$  is a bounded functional on  $\text{Dom}(T^*)$ .  $\square$

We now apply this to symmetric first-order differential operators on compact manifolds without boundary. Recall that a *first-order differential operator*  $D$  on a vector bundle  $E \rightarrow M$  has the following local expression:

$$D = \sum_{\mu} A^\mu(x) \frac{\partial}{\partial x^\mu} + B(x)$$

where  $A^\mu(x), B(x) : E_x \rightarrow E_x$  are endomorphisms acting on the fibers of  $E$ . We will always act with  $D$  on smooth sections  $\Gamma^\infty(E)$  of the bundle  $E \rightarrow M$  and we furthermore fix an inner product on  $\Gamma^\infty(E)$ .

**Proposition 4.26** *Every symmetric first-order differential operator  $D$  on a vector bundle  $E$  over a compact manifold without boundary is essentially self-adjoint.*

**Proof** The proof uses so-called *Friedrichs' mollifiers* on  $M$ . For all sufficiently small  $t > 0$  there exist (cf. Exercise below) self-adjoint operators  $F_t : L^2(E) \rightarrow L^2(E)$  such that

- (i)  $\|F_t\| \leq 1$ ;
- (ii) for each  $u \in L^2(E)$ ,  $F_t u \rightarrow u$  in  $L^2(E)$  as  $t \rightarrow 0$ ;
- (iii) for each  $u \in L^2(E)$ ,  $F_t u$  is smooth;

- (iv) the commutator  $[D, F_t]$  extends to a bounded operator on  $L^2(E)$ , whose norm is bounded independent of  $t$ .

Then, let  $u \in \text{Dom}(D^*)$ . In order to conclude that  $D$  is essentially self-adjoint we will show that  $u \in \text{Dom}(\overline{D})$ . Note that  $F_t u$  is smooth and tends to  $u$  as  $t \rightarrow 0$ . Moreover,

$$D(F_t u) = F_t D^* u + [D^*, F_t] u$$

On  $\text{Dom}(D^*)$  we have  $[D^*, F_t] = [F_t, D]^*$ , which is the adjoint of a bounded operator (with norm bound independent of  $t$ ). Thus,  $D(F_t u)$  is uniformly bounded so that  $u \in \text{Dom}(\overline{D})$  by the previous Lemma.  $\square$

**Corollary 4.27** *The Dirac operator  $D_M$  on a compact Riemannian spin<sup>c</sup> manifold without boundary is essentially self-adjoint.*

**Exercise 4.7** Let  $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$  be a smooth, positive function with compact support and with total mass 1. Define an operator  $F_t$  on  $L^2(\mathbb{R}^n)$  by

$$(F_t u)(x) = t^{-n} \int_{\mathbb{R}^n} \varphi\left(\frac{x-y}{t}\right) u(y) dy.$$

Show that  $\{F_t\}$  is a family of Friedrichs' mollifiers on  $L^2(\mathbb{R}^n)$ , i.e., a family satisfying (i)–(iv) in the Proof of Proposition 4.26. Using local coordinates and partitions of unity, graft this family onto an arbitrary compact manifold  $M$  to construct a family of Friedrichs' mollifiers on  $M$ .

### 4.3.3 Compact Resolvent

Here we will rely on a crucial embedding result on Sobolev spaces, namely the Rellich Lemma, which we state without proof.

To start, recall the definition of the first Sobolev space  $H^1(\mathbb{R}^n)$  on Euclidean space  $\mathbb{R}^n$ : it is the completion of the compactly supported smooth functions  $C_c^\infty(\mathbb{R}^n)$  in the norm coming from the inner product

$$(f_1, f_2)_{H^1(\mathbb{R}^n)} = (f_1, f_2)_{L^2} + \sum_{\mu=1}^n (\partial_\mu f_1, \partial_\mu f_2)_{L^2}; \quad (f_1, f_2 \in C_c^\infty(\mathbb{R}^n)).$$

More generally, for a compact manifold  $M$  we can use partition of unity to extend this definition to give  $H^1(M)$ . So, let  $\chi_\alpha$  be a partition of unity subordinate to an atlas  $(U_\alpha, \phi_\alpha)$  of  $M$ . We define  $H^1(M)$  to be the completion of  $C^\infty(M)$  with respect to the inner product

$$(f_1, f_2)_{H^1(M)} = \sum_{\alpha} ((\chi_\alpha \cdot f_1) \circ \phi_\alpha^{-1}, (\chi_\alpha \cdot f_2) \circ \phi_\alpha^{-1})_{H^1(\mathbb{R}^n)}$$

This definition turns out to be independent of the choice of an atlas (see Note 9 of this chapter below). Moreover, by using trivializing charts it can easily be extended to give rise to the Sobolev spaces  $H^1(M, E)$  of sections of a bundle  $E \rightarrow M$ .

**Lemma 4.28** (Rellich) *Let  $M$  be a compact manifold. Then the inclusion map  $H^1(M, E)$  into  $L^2(M, E)$  is a compact map.*

We will not prove it in full generality here (see Note 10 of this chapter below), but give a proof for the case of the circle in Exercise 4.8 below.

This result can be used to show the compact resolvent property for the Dirac operator  $D_M$  in the following way. First of all, the norm coming from the inner product on  $H^1(M, E)$  is equivalent to the graph norm of  $\overline{D_M}$  (see Note 9 of this chapter below), so that Rellich Lemma then implies that the inclusion map

$$\iota : \text{Dom}(\overline{D_M}) \rightarrow L^2(S)$$

is a compact map.

**Proposition 4.29** *The adjoint  $\iota^* : L^2(S) \rightarrow \text{Dom}(\overline{D_M}) \subseteq L^2(S)$  is given by  $\iota^* = (1 + \overline{D_M}^2)^{-1}$ .*

**Proof** For  $\psi_1 \in L^2(S)$  and  $\psi_2 \in \text{Dom}(\overline{D_M})$  we have in terms of the inner product on the graph  $G(\overline{D_M})$  of  $\overline{D_M}$ :

$$\begin{aligned} \left( (1 + \overline{D_M}^2)^{-1}(\psi_1), \psi_2 \right)_{G(\overline{D_M})} &= \left( (1 + \overline{D_M}^2)^{-1}\psi_1, \psi_2 \right)_{L^2} \\ &\quad + \left( \overline{D_M}(1 + \overline{D_M}^2)^{-1}\psi_1, \overline{D_M}\psi_2 \right)_{L^2} \\ &= (\psi_1, \psi_2)_{L^2} = (\psi_1, \iota(\psi_2))_{L^2}. \end{aligned}$$

□

**Corollary 4.30** *The resolvent  $(i + \overline{D_M})^{-1}$  of the Dirac operator on a compact Riemannian spin<sup>c</sup> manifold without boundary is a compact operator.*

**Proof** We start by writing

$$(i + \overline{D_M})^{-1} = ((i + \overline{D_M})^{-1}(1 + \overline{D_M}^2)^{1/2}(1 + \overline{D_M}^2)^{-1/2}).$$

The operator  $(i + \overline{D_M})^{-1}(1 + \overline{D_M}^2)^{1/2}$  is a bounded operator. In fact, using the functional calculus on self-adjoint operators we find that

$$\| (i + \overline{D_M})^{-1}(1 + \overline{D_M}^2)^{1/2} \| \leq \sup_{t \in \mathbb{R}} \left\{ \frac{\sqrt{1 + t^2}}{|i + t|} \right\} = 1.$$

Also, observe that  $(1 + \overline{D_M}^2)^{-1/2}$  is a square root of the positive operator  $(1 + \overline{D_M}^2)^{-1}$ . When we consider the latter as an operator on  $L^2(S)$  it is compact by the above Proposition. This is enough to conclude that  $(i + \overline{D_M})^{-1}$  is compact. □

In the following exercise we consider Rellich Lemma on the circle.

**Exercise 4.8** Write a function  $f \in L^2(S^1)$  as a Fourier series:

$$f = \sum_{n \in \mathbb{Z}} f_n e_n$$

where  $e_n(t) = e^{2\pi i n t}$  and the sequence  $\{f_n\}$  is in  $l^2(\mathbb{Z})$ . The Sobolev space  $H^1(S^1)$  consists of those  $L^2$ -functions for which

$$\|f\|_{H^1}^2 = \sum_{n \in \mathbb{Z}} (1 + n^2) |f_n|^2$$

is finite.

(1) Establish the following estimate

$$\sup_t \left| \sum_{|n| \leq N} f_n e_n \right| \leq \left( \sum_{|n| \leq N} |f_n|^2 (1 + n^2) \right)^{1/2} \left( \sum_{|n| \leq N} \frac{1}{1 + n^2} \right)^{1/2}$$

and derive from this that  $H^1(S^1) \subset C(S^1)$ .

(2) Show that the inclusion  $H^1(S^1) \rightarrow L^2(S^1)$  is given by the norm limit of the sequence  $P_N$  of finite-rank operators that send a function  $f$  to the  $N$ 'th partial sum

$$P_N f = \sum_{|n| \leq N} f_n e_n$$

In some cases one can derive the above analytical properties (such as essential self-adjointness and compact resolvent) from the knowledge of the spectrum of eigenvalues, as the following exercise shows.

**Exercise 4.9** Let  $\{\lambda_n\}_{n \in \mathbb{Z}}$  be a sequence of real numbers, possibly with degeneracies but ordered such that  $\lambda_n \leq \lambda_{n+1}$ . Furthermore, assume that  $\pm\infty$  are the only accumulation points and that  $\lambda_{\pm n} \rightarrow \pm\infty$  as  $n \rightarrow \infty$ . We define a dense subspace in  $l^2(\mathbb{Z})$  by

$$\text{Dom}(D) = \text{span}_{\mathbb{C}}\{e_n : n \in \mathbb{Z}\}$$

and introduce an unbounded linear operator  $D$  on  $\text{Dom}(D) \subset l^2(\mathbb{Z})$  by setting

$$De_n = \lambda_n e_n.$$

- (1) Show that  $D$  is an essentially self-adjoint operator.
- (2) Show that the resolvent  $(i + D)^{-1}$  is a compact operator on  $l^2(\mathbb{Z})$ .

## Notes

### Section 4.1. Clifford Algebras

1. In our treatment of Clifford algebras, we stay close to the seminal paper by Atiyah, Bott and Shapiro [1], but also refer to the standard textbook [2] and the book [3, Chap. 5]. We also take inspiration from the lecture notes [4, 5].
2. The definition of a quadratic form given here is equivalent with the usual definition, which states that  $Q$  is a quadratic form if  $Q(v) = S(v, v)$  for some symmetric bilinear form  $S$  (cf. Exercise 4.1). This is shown by Jordan and von Neumann in [6].
3. The periodicity eight encountered for the real Clifford algebras  $\text{Cl}_k^\pm$  is closely related to the eightfold periodicity of KO-theory [7]. The periodicity two encountered for the complex Clifford algebras  $\mathbb{C}\text{l}_n$  is closely related to Bott periodicity in K-theory [8].

### Section 4.2 . Riemannian Spin Geometry

4. A standard textbook on Riemannian geometry is [9]. For a complete treatment of Riemannian spin manifolds we refer to e.g., [2, 10]. The noncommutative approach to (commutative) spin geometry that we adopt here can also be found in [3, Chap. 9] or [4, 11].
5. In Definition 4.10 a Riemannian manifold is said to be  $\text{spin}^c$  if  $\mathbb{C}\text{l}(TM) \simeq \text{End}(S)$  (even case). Glancing back at Chap. 2 we see that  $\mathbb{C}\text{l}_n$  is Morita equivalent to  $\mathbb{C}$  ( $n$  even). With Definition 7.9 of the next Chapter, we conclude that a manifold is  $\text{spin}^c$  precisely if (the  $C^*$ -completion of)  $\text{Cliff}(M)$  is Morita equivalent to  $C(M)$ . This is the algebraic approach to  $\text{spin}^c$  manifolds laid out in [3, Sect. 9.2].
6. A first reference to Theorem 4.22 is [12, Sect. VI.1].
7. Lemma 4.25 can be found as [13, Lemma 1.8.1])
8. The proof of Proposition 4.26 is based on [13, Lemma 10.2.5]
9. The fact that the definition of the inner product for  $H^1(M)$  is independent of the choice of an atlas is shown in for instance [14, Sect. 1.3.4]. The relation of this inner product to the graph norm of  $\overline{D_M}$  is a deep result which is a consequence of ellipticity of the Dirac operator, see for instance [14, Lemma 1.3.6].
10. A proof of the Rellich Lemma 4.28 can be found in [15, Corollary II.1.2] and in [16, Lemma 1.7].

## References

1. Atiyah, M.F., Bott, R., Shapiro, A.: Clifford modules. Topology **3**, 3–38 (1964)
2. Lawson, H.B., Michelsohn, M.-L.: Spin Geometry. Princeton University Press (1989)
3. Gracia-Bondía, J.M., Várilly, J.C., Figueroa, H.: Elements of Noncommutative Geometry. Birkhäuser, Boston (2001)
4. Várilly, J.C.: Dirac operators and spectral geometry. Notes taken by Paweł Witkowski (2006)
5. Landsman, N.P.: Notes on noncommutative geometry. <http://www.math.ru.nl/~landsman/ncg2010.html>
6. Jordan, P., Von Neumann, J.: On inner products in linear, metric spaces. Ann. Math. (2) **36**, 719–723 (1935)
7. Atiyah, M.F.: K-theory and reality. Quart. J. Math. Oxford Ser. (2) **17**, 367–386 (1966)
8. Atiyah, M., Bott, R.: On the periodicity theorem for complex vector bundles. Acta Math. **112**, 229–247 (1964)

9. Jost, J.: Riemannian Geometry and Geometric Analysis, 3rd edn. Universitext. Springer-Verlag, Berlin (2002)
10. Berline, N., Getzler, E., Vergne, M.: Heat Kernels and Dirac Operators. Springer, Berlin (1992)
11. Várilly, J.C.: An Introduction to Noncommutative Geometry. Publishing House (EMS Series of Lectures in Mathematics) European Mathematical Society (2006)
12. Connes, A.: Noncommutative Geometry. Academic, San Diego (1994)
13. Higson, N., Roe, J.: Analytic  $K$ -homology. Oxford Mathematical Monographs. Oxford University Press, Oxford (2000). Oxford Science Publications
14. Gilkey, P.B.: Invariance theory, the heat equation, and the Atiyah-Singer index theorem. Mathematics Lecture Series, vol. 11. Publish or Perish Inc., Wilmington, DE (1984)
15. Treves, F.: Introduction to Pseudodifferential and Fourier Integral Operators, vol. 1. Plenum Press (1980)
16. Higson, N.: The local index formula in noncommutative geometry. In: Contemporary Developments in Algebraic  $K$ -Theory, ICTP Lectures Notes, XV, pp. 443–536. Abdus Salam International Centre for Theoretical Physics, Trieste (2004)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 5

## Noncommutative Riemannian Spin Manifolds



This chapter introduces the main technical device—spectral triples—that generalizes Riemannian spin geometry to the noncommutative world. We exemplify this by means of toric noncommutative manifolds; this includes the noncommutative torus.

### 5.1 Gelfand Duality

The first step towards noncommutative manifolds is to arrive at an algebraic characterization of topological spaces. This is accomplished by **Gelfand duality**, giving a one-to-one correspondence between compact Hausdorff topological spaces and commutative  $C^*$ -algebras. Let us recall some definitions.

**Definition 5.1** A  $C^*$ -algebra  $A$  is a (complex)  $*$ -algebra (Definition 2.1) that is complete with respect to a multiplicative norm (i.e.  $\|ab\| \leq \|a\|\|b\|$  for all  $a, b \in A$ ) that satisfies the  $C^*$ -property:

$$\|a^*a\| = \|a\|^2.$$

**Example 5.2** The key example of a commutative  $C^*$ -algebra is the algebra  $C(X)$  for a compact topological space  $X$ . Indeed, uniform continuity is captured by the norm

$$\|f\| = \sup\{|f(x)| : x \in X\}$$

and involution defined by  $f^*(x) = f(\bar{x})$ . This indeed satisfies  $\|f^*f\| = \|f\|^2$ .

**Example 5.3** Another key example where  $A$  is noncommutative is given by the  $*$ -algebra of bounded operators  $\mathcal{B}(\mathcal{H})$  on a Hilbert space  $\mathcal{H}$ , equipped with the operator norm.

The following result connects with the matrix algebras of Chap. 2.

**Proposition 5.4** *If  $A$  is a finite-dimensional  $C^*$ -algebra, then it is isomorphic to a matrix algebra:*

$$A \simeq \bigoplus_{i=1}^N M_{n_i}(\mathbb{C}).$$

**Proof** See Note 2 in this chapter. □

In Chap. 2 we defined the structure space of a  $*$ -algebra  $A$  to consist of (equivalence classes of) irreducible representations of  $A$ . Let us extend this definition to  $C^*$ -algebras.

**Definition 5.5** A *representation* of a  $C^*$ -algebra  $A$  is a pair  $(\mathcal{H}, \pi)$  where  $\mathcal{H}$  is a Hilbert space and  $\pi$  is a  $*$ -algebra map

$$\pi : A \rightarrow \mathcal{B}(\mathcal{H}).$$

A representation  $(\mathcal{H}, \pi)$  is irreducible if  $\mathcal{H} \neq 0$  and the only closed subspaces in  $\mathcal{H}$  that are left invariant under the action of  $A$  are  $\{0\}$  and  $\mathcal{H}$ .

Two representations  $(\mathcal{H}_1, \pi_1)$  and  $(\mathcal{H}_2, \pi_2)$  of a  $C^*$ -algebra  $A$  are *unitarily equivalent* if there exists a unitary map  $U : \mathcal{H}_1 \rightarrow \mathcal{H}_2$  such that

$$\pi_1(a) = U^* \pi_2(a) U.$$

**Definition 5.6** The *structure space*  $\widehat{A}$  of a  $C^*$ -algebra  $A$  is the set of all unitary equivalence classes of irreducible representations of  $A$ .

In Chap. 2 we considered the commutative matrix algebra  $\mathbb{C}^N$  whose structure space was the finite topological space consisting of  $N$  points. Let us sketch the generalization to compact Hausdorff topological spaces, building towards **Gelfand duality**. As a motivating example, we consider the  $C^*$ -algebra  $C(X)$  for a compact Hausdorff topological space  $X$  (cf. Example 5.2). As this  $C^*$ -algebra is commutative, a standard argument shows that any irreducible representation  $\pi$  of  $C(X)$  is one-dimensional. In fact, any such  $\pi$  is equivalent to the evaluation map  $\text{ev}_x$  at some point  $x$  of  $X$ , given by

$$\begin{aligned} \text{ev}_x : C(X) &\rightarrow \mathbb{C}; \\ f &\mapsto f(x). \end{aligned}$$

Being a one-dimensional representation,  $\text{ev}_x$  is automatically an irreducible representation. It follows that the structure space of  $C(X)$  is given by the set of points of  $X$ . But more is true, as the topology of  $X$  is also captured by the structure space. Namely, since in the commutative case the irreducible representations are one-dimensional  $\pi : A \rightarrow \mathbb{C}$  the structure space can be equipped with the **weak  $*$ -topology**. That is to say, for a sequence  $\{\pi_n\}_n$  in  $\widehat{A}$ ,  $\pi_n$  converges weakly to  $\pi$  if  $\pi_n(a) \rightarrow \pi(a)$  for all  $a \in A$ .

We state the main result, generalizing our finite-dimensional version of Sect. 2.1.1 to the infinite-dimensional setting.

**Theorem 5.7** (Gelfand duality) *The structure space  $\widehat{A}$  of a commutative unital  $C^*$ -algebra  $A$  is a compact Hausdorff topological space, and  $A \simeq C(\widehat{A})$  via the Gelfand transform*

$$a \in A \mapsto \widehat{a} \in \widehat{A}; \quad \widehat{a}(\pi) = \pi(a).$$

Moreover, for any compact Hausdorff topological space  $X$  we have

$$\widehat{C(X)} \simeq X.$$

**Proof** See Note 3 in this chapter. □


## 5.2 Spectral Triples

The next milestone which we need to reach noncommutative Riemannian spin geometry is the translation of the Riemannian distance (4.2.1) on a compact Riemannian spin manifold into functional analytic data. Indeed, we will give an alternative formula as a *supremum* over functions in  $C^\infty(M)$ . The translation from points in  $M$  to functions on  $M$  is accomplished by imposing that the gradient of the functions is less than 1 (see Fig. 5.1). This is the continuum analogue of Theorem 2.18.

**Proposition 5.8** *Let  $M$  be a Riemannian spin<sup>c</sup>-manifold with Dirac operator  $D_M$ . The following formula defines a distance between points in  $\widehat{C(M)} \simeq M$ :*

$$d(x, y) = \sup_{f \in C^\infty(M)} \{ |f(x) - f(y)| : \| [D_M, f] \| \leq 1 \}.$$

Moreover, this distance function  $d$  coincides with the Riemannian distance function  $d_g$ .



**Fig. 5.1** The translation of the distance between points  $x, y$  in  $M$  to a formulation in terms of functions of slope  $\leq 1$

**Proof** First, note that the relation  $\|f\|_{\text{Lip}} = \|[D_M, f]\| \leq 1$  (cf. Theorem 4.20) already ensures that  $d(x, y) \leq d_g(x, y)$ . For the opposite inequality we fix  $y \in M$  and consider the function  $f_{g,y}(z) = d_g(z, y)$ . Then  $\|f_{g,y}\|_{\text{Lip}} \leq 1$  and

$$d(x, y) \geq |f_{g,y}(x) - f_{g,y}(y)| = d_g(x, y),$$

as required.  $\square$

Thus, we have reconstructed the Riemannian distance on  $M$  from the algebra  $C^\infty(M)$  of functions on  $M$  and the Dirac operator  $D_M$ , both acting in the Hilbert space  $L^2(S)$  of square-integrable operators. Note that the triple  $(C^\infty(M), L^2(S), D_M)$  consists of mere functional analytical, or ‘spectral’ objects, instead of geometrical. Upon allowing for noncommutative algebras as well, we arrive at the following spectral data required to describe a noncommutative Riemannian spin manifold.

**Definition 5.9** A *spectral triple*  $(\mathcal{A}, \mathcal{H}, D)$  is given by a unital  $*$ -algebra  $\mathcal{A}$  represented as bounded operators on a Hilbert space  $\mathcal{H}$  and a self-adjoint operator  $D$  in  $\mathcal{H}$  such that the resolvent  $(i + D)^{-1}$  is a compact operator and  $[D, a]$  extends to a bounded operator for each  $a \in \mathcal{A}$ .

A spectral triple is *even* if the Hilbert space  $\mathcal{H}$  is endowed with a  $\mathbb{Z}_2$ -grading  $\gamma$  such that  $\gamma a = a\gamma$  and  $\gamma D = -D\gamma$ .

A *real structure* of KO-dimension  $n \in \mathbb{Z}/8\mathbb{Z}$  on a spectral triple is an anti-linear isometry  $J : \mathcal{H} \rightarrow \mathcal{H}$  such that

$$J^2 = \varepsilon, \quad JD = \varepsilon'DJ, \quad J\gamma = \varepsilon''\gamma J \quad (\text{even case}),$$

where the numbers  $\varepsilon, \varepsilon', \varepsilon'' \in \{-1, 1\}$  are given as a function of  $n$  modulo 8, as they appear in Table 5.1.

Moreover, with  $b^0 = Jb^*J^{-1}$  we impose the *commutant property* and the *order one condition*:

$$[a, b^0] = 0, \quad [[D, a], b^0] = 0; \quad (a, b \in \mathcal{A}). \quad (5.2.1)$$

A spectral triple with a real structure is called a *real spectral triple*.

**Table 5.1** The KO-dimension  $n$  of a real spectral triple is determined by the signs  $\{\varepsilon, \varepsilon', \varepsilon''\}$  appearing in  $J^2 = \varepsilon$ ,  $JD = \varepsilon'DJ$  and  $J\gamma = \varepsilon''\gamma J$

$n$	0	1	2	3	4	5	6	7
$\varepsilon$	1	1	-1	-1	-1	-1	1	1
$\varepsilon'$	1	-1	1	1	1	-1	1	1
$\varepsilon''$	1		-1		1		-1	

**Remark 5.10** The notation  $(\mathcal{A}, \mathcal{H}, D)$  is chosen to distinguish a general spectral triple from the finite spectral triples considered in Chaps. 2 and 3, which were denoted as  $(A, H, D)$ .

The basic example of a spectral triple is the **canonical triple** associated to a compact Riemannian spin manifold:

- $\mathcal{A} = C^\infty(M)$ , the algebra of smooth functions on  $M$ ;
- $\mathcal{H} = L^2(S)$ , the Hilbert space of square integrable sections of a spinor bundle  $S \rightarrow M$ ;
- $D = D_M$ , the Dirac operator associated to the Levi–Civita connection lifted to the spinor bundle.

The real structure  $J$  is given by the charge conjugation  $J_M$  of Definition 4.13. If the manifold is even dimensional then there is a grading on  $\mathcal{H}$ , defined just below Definition 4.13. Since the signs in the above table coincide with those in Table 4.2, the KO-dimension of the canonical triple coincides with the dimension of  $M$ .

**Example 5.11** The tangent bundle of the circle  $\mathbb{S}^1$  is trivial and has one-dimensional fibers, so that spinors are given by ordinary functions on  $\mathbb{S}^1$ . Moreover, the Dirac operator  $D_{\mathbb{S}^1}$  is given by  $-id/dt$  where  $t \in [0, 2\pi]$ , acting on  $C^\infty(\mathbb{S}^1)$  (which is a core for  $D_{\mathbb{S}^1}$ ). The eigenfunction of  $D_{\mathbb{S}^1}$  are the exponential function  $e^{int}$  with eigenvalues  $n \in \mathbb{Z}$ . As such,  $(i + D_{\mathbb{S}^1})^{-1}$  is a compact operator. Moreover  $[D_{\mathbb{S}^1}, f] = -idf/dt$  is bounded. Summarizing, we have the following spectral triple:

$$\left( C^\infty(\mathbb{S}^1), L^2(\mathbb{S}^1), -i \frac{d}{dt} \right).$$

Note that the supremum norm of a function  $f \in C^\infty(\mathbb{S}^1)$  coincides with the operator norm of  $f$  considered as multiplication operator on  $L^2(\mathbb{S}^1)$ . A real structure is given by complex conjugation on  $L^2(\mathbb{S}^1)$ , making the above a real spectral triple of KO-dimension 1.

**Example 5.12** Since the tangent bundle of the torus  $\mathbb{T}^2$  is trivial, we have  $\text{Cliff}(\mathbb{T}^2) \simeq C(\mathbb{T}^2) \otimes \text{Cl}_2$ . As a consequence, the spinor bundle is trivial,  $S = \mathbb{T}^2 \times \mathbb{C}^2$ , and  $L^2(S) = L^2(\mathbb{T}^2) \otimes \mathbb{C}^2$ . The generators  $\gamma^1$  and  $\gamma^2$  are given by

$$\gamma^1 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \gamma^2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

which satisfy (4.2.4). The chirality operator is then given by

$$\gamma_{\mathbb{T}^2} = -i\gamma^1\gamma^2 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix},$$

and the real structure  $J_{\mathbb{T}^2}$  that selects  $\text{Cl}_2^- \subset \mathbb{C}\text{Cl}_2$  is

$$J_{\mathbb{T}^2} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} -\bar{v}_2 \\ \bar{v}_1 \end{pmatrix}.$$

Finally, the Dirac operator on  $\mathbb{T}^2$  is

$$D_{\mathbb{T}^2} = -i\gamma^\mu \partial_\mu = \begin{pmatrix} 0 & -\partial_1 - i\partial_2 \\ \partial_1 - i\partial_2 & 0 \end{pmatrix}.$$

The eigenspinors of  $D_{\mathbb{T}^2}$  are given by the vectors

$$\phi_{n_1, n_2}^\pm(t_1, t_2) := \frac{1}{\sqrt{2}} \begin{pmatrix} e^{i(n_1 t_1 + n_2 t_2)} \\ \pm \frac{i n_1 + n_2}{\sqrt{n_1^2 + n_2^2}} e^{i(n_1 t_1 + n_2 t_2)} \end{pmatrix}; \quad (n_1, n_2 \in \mathbb{Z}),$$

with eigenvalues  $\pm\sqrt{n_1^2 + n_2^2}$ . Again, this ensures that  $(i + D_{\mathbb{T}^2})^{-1}$  is a compact operator. For the commutator with a function  $f \in C^\infty(\mathbb{T}^2)$  we compute

$$[D_{\mathbb{T}^2}, f] = \begin{pmatrix} 0 & -\partial_1 f - i\partial_2 f \\ \partial_1 f - i\partial_2 f & 0 \end{pmatrix},$$

which is bounded because  $\partial_1 f$  and  $\partial_2 f$  are bounded. The signs in the commutation between  $J_{\mathbb{T}^2}$ ,  $D_{\mathbb{T}^2}$  and  $\gamma_{\mathbb{T}^2}$  makes the following a spectral triple of KO-dimension 2:

$$(C^\infty(\mathbb{T}^2), L^2(\mathbb{T}^2) \otimes \mathbb{C}^2, D_{\mathbb{T}^2}; J_{\mathbb{T}^2}, \gamma_{\mathbb{T}^2}).$$

Other examples are given by finite spectral triples, discussed at length—and classified—in Chap. 2. Indeed, the compact resolvent condition is automatic in finite-dimensional Hilbert spaces; similarly, any operator such as  $[D, a]$  is bounded as in this case also  $D$  is a bounded operator. More serious noncommutative examples are presented below in Sect. 5.3.

Corresponding to the direct product of manifolds, one can take the product of spectral triples as follows (see also Exercise 2.24). Suppose that  $(\mathcal{A}_1, \mathcal{H}_1, D_1; \gamma_1, J_1)$  and  $(\mathcal{A}_2, \mathcal{H}_2, D_2; \gamma_2, J_2)$  are even real spectral triples, then we define the **product spectral triple** by

$$\begin{aligned} \mathcal{A} &= \mathcal{A}_1 \otimes \mathcal{A}_2; \\ \mathcal{H} &= \mathcal{H}_1 \otimes \mathcal{H}_2; \\ D &= D_1 \otimes 1 + \gamma_1 \otimes D_2; \\ \gamma &= \gamma_1 \otimes \gamma_2; \\ J &= J_1 \otimes J_2. \end{aligned}$$

If  $(\mathcal{A}_2, \mathcal{H}_2, D_2; J_2)$  is odd, then we can still form the product when we leave out  $\gamma$ . Note that  $D^2 = D_1^2 \otimes 1 + 1 \otimes D_2^2$ , since the cross-terms vanish due to the fact that  $\gamma_1 D_1 = -D_1 \gamma_1$ .

**Example 5.13** In the physical applications later in this book (Chap. 10 and afterwards) we are mainly interested in almost-commutative manifolds which are defined as products of a Riemannian spin manifold  $M$  with a finite noncommutative space  $F$ . More precisely, we will consider

$$M \times F := (C^\infty(M), L^2(S), \mathcal{D}; J_M, \gamma_M) \otimes (A_F, H_F, D_F; J_F, \gamma_F),$$

with  $(A_F, H_F, D_F; J_F, \gamma_F)$  as in Definition 2.19. Note that this can be identified with:

$$M \times F = (C^\infty(M, A_F), L^2(S \otimes (M \times H_F)), \mathcal{D} \otimes 1 + \gamma_M \otimes D_F; J_M \otimes J_F, \gamma_M \otimes \gamma_F),$$

in terms of the trivial vector bundle  $M \times H_F$  on  $M$ .

Returning to the general case, Definition 2.24 encountered before in the context of finite spectral triples can be translated *verbatim* to the general case:

**Definition 5.14** Two spectral triples  $(\mathcal{A}_1, \mathcal{H}_1, D_1)$  and  $(\mathcal{A}_2, \mathcal{H}_2, D_2)$  are called *unitarily equivalent* if  $\mathcal{A}_1 = \mathcal{A}_2$  and if there exists a unitary operator  $U : \mathcal{H}_1 \rightarrow \mathcal{H}_2$  such that

$$\begin{aligned} U\pi_1(a)U^* &= \pi_2(a); & (a \in \mathcal{A}_1), \\ UD_1U^* &= D_2, \end{aligned}$$

where we have explicitly indicated the representations  $\pi_i$  of  $\mathcal{A}_i$  on  $\mathcal{H}_i$  ( $i = 1, 2$ ).

Moreover, any spectral triple gives rise to a differential calculus. This generalizes our previous Definition 2.22 for the finite-dimensional case. Again, we focus only on differential one-forms, as this is sufficient for our applications to gauge theory later on.

**Definition 5.15** The  $\mathcal{A}$ -bimodule of *Connes' differential one-forms* is given by

$$\Omega_D^1(\mathcal{A}) := \left\{ \sum_k a_k [D, b_k] : a_k, b_k \in \mathcal{A} \right\},$$

and the corresponding derivation  $d : \mathcal{A} \rightarrow \Omega^1(\mathcal{A})$  is given by  $d = [D, \cdot]$ .

**Exercise 5.1** (1) In the case of a Riemannian spin manifold  $M$ , verify that we can identify  $\Omega_{\mathcal{D}}^1(C^\infty(M)) \simeq \Omega_{\text{dR}}^1(M)$ , the usual De Rham differential one-forms.  
(2) In the case of an almost-commutative manifold  $M \times F$ , verify that we have

$$\Omega_{\mathcal{D} \otimes 1 + \gamma_M \otimes D_F}^1(C^\infty(M, A_F)) \simeq \Omega_{\text{dR}}^1(M, A_F) \oplus C^\infty(M, \Omega_{D_F}^1(A_F)).$$

### 5.3 Examples of Noncommutative Manifolds

#### 5.3.1 The Noncommutative Torus

We now give a detailed exposition of a noncommutative example of a real spectral triple, namely, we describe the *noncommutative torus*. Let us start with the noncommutative topological, i.e. the  $C^*$ -algebraic aspects.

**Definition 5.16** Let  $\theta$  be a real number. We define the *noncommutative torus  $C^*$ -algebra*  $A_\theta$  to be the unital  $C^*$ -algebra generated by  $u, v$  subject to the relations

$$u^*u = uu^* = 1; \quad v^*v = vv^* = 1; \quad vu = \lambda uv; \quad (\lambda = e^{2\pi i\theta})$$

The *smooth noncommutative torus algebra* is then given by

$$\mathcal{A}_\theta := \{a = \sum_{r,s} a_{rs} u^r v^s : (a_{rs}) \in \mathcal{S}(\mathbb{Z}^2)\}$$

where the space of Schwartz sequences is defined as

$$\mathcal{S}(\mathbb{Z}^2) := \{(a_{rs}) : \sup_{r,s \in \mathbb{Z}} (1 + r^2 + s^2)^k |a_{rs}|^2 < \infty \text{ for any } k \geq 0\}$$

Note that for the  $*$ -algebra structure in  $\mathcal{A}_\theta$  we have for  $a, b \in \mathcal{A}_\theta$ :

$$ab = \sum_{r,s,n,m} \lambda^{mn} a_{r-n,m} b_{n,s-m} u^r v^s; \quad a^* = \sum_{r,s} \lambda^{rs} \overline{a_{-r,-s}} u^r v^s. \quad (5.3.1)$$

Also, if  $\theta = 0$  we may identify  $u, v$  with multiplication operators by  $z_1, z_2 \in \mathbb{S}^1$ , so that  $\mathcal{A}_0 \cong C^\infty(\mathbb{T}^2)$  identifying the series  $\sum_{r,s} a_{rs} u^r v^s$  as a Fourier series in two variables. This should explain the terminology noncommutative torus whenever  $\theta \neq 0$ .

Next, consider the normalized faithful trace  $\tau : \mathcal{A}_\theta \rightarrow \mathbb{C}$  given by

$$\tau(a) = a_{00},$$

so that  $\tau(a^*a) = \sum_{r,s} |a_{rs}|^2 > 0$  for  $a \neq 0$ . Also  $\tau(1) = 1$  and  $\tau(ab) = \tau(ba)$  as follows readily from Eq. (5.3.1).

**Lemma 5.17** *The trace  $\tau$  extends to a continuous trace  $\tau : A_\theta \rightarrow \mathbb{C}$  and in fact  $|\tau(a)| \leq \|a\|$ .*

**Proof** Continuity of  $\tau : A_\theta \rightarrow \mathbb{C}$  follows since we may write on polynomials  $a \in \mathbb{C}\langle u, v \rangle$ :

$$\tau(a) \cdot 1 = \int_{\mathbb{T}^2} \alpha_{t_1, t_2}(a) dt$$

where  $\alpha_{t_1, t_2}(u^r v^s) = e^{i(rt_1 + st_2)}$  defines an action of  $\mathbb{T}^2$  on  $\mathcal{A}_\theta$ . It thus follows that  $|\tau(a)| \leq \|a\|$ .  $\square$

Using this faithful trace we may apply the GNS-construction. This gives a Hilbert space

$$\mathcal{H}_\tau := L^2(A_\theta, \tau)$$

which—since  $\tau$  is faithful—is given by the closure of  $A_\theta$  with respect to the Hilbert space norm  $\|a\|_2 = \sqrt{\tau(a^*a)}$ . The representation  $\pi_\tau : A_\theta \rightarrow \mathcal{B}(\mathcal{H}_\tau)$  is given by

$$\pi_\tau(a)b = ab.$$

As in Exercise 3.2 there is a Tomita involution with respect to the cyclic and separating vector  $1 \in \mathcal{H}_\tau$ . It is given explicitly by

$$J_\tau(a) = a^*.$$

This anti-unitary operator can be used to define a right representation of  $A_\theta$  on  $\mathcal{H}_\tau$  by setting  $\pi_\tau^\circ(a) = J_\tau \pi(a^*) J_\tau^{-1}$ . In fact, we then have

$$\pi_\tau^\circ(a)b = J_\tau a^* b^* = ba$$

for all  $a, b \in A_\theta$ . It then follows that  $[\pi_\tau^\circ(a), \pi(b)] = 0$ , so that  $\mathcal{H}_\tau$  becomes a  $A_\theta$ -bimodule.

We now prepare for spinors by doubling our Hilbert space:

$$\mathcal{H} = \mathcal{H}_\tau \oplus \mathcal{H}_\tau.$$

The algebra representation of  $A_\theta$  on  $\mathcal{H}$  is the diagonal representation  $\pi = \pi_\tau \oplus \pi$  while we set

$$J = \begin{pmatrix} 0 & -J_\tau \\ J_\tau & 0 \end{pmatrix}; \quad \gamma = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

As such, we have  $J^2 = -1$  and  $J\gamma = -\gamma J$ , suggesting the KO-dimension to be 2 as can be read off from Table 5.1.

The final step in the description of the noncommutative torus in terms of a real spectral triple is the introduction of a noncommutative analogue of a Dirac operator. For this, we first consider the basic derivations  $\delta_j : \mathcal{A}_\theta \rightarrow \mathcal{A}_\theta$  given by

$$\delta_1 \left( \sum_{r,s} a_{rs} u^r v^s \right) = \sum_{r,s} i r a_{rs} u^r v^s; \quad \delta_2 \left( \sum_{r,s} a_{rs} u^r v^s \right) = \sum_{r,s} i s a_{rs} u^r v^s. \quad (5.3.2)$$

Indeed, it follows from Eq. (5.3.1) that

$$\delta_j(ab) = \delta_j(a)b + a\delta_j(b); \quad (\delta_j a)^* = \delta_j a^*. \quad (5.3.3)$$

One should consider the derivations  $\delta_1, \delta_2$  as the noncommutative analogues of the partial derivatives  $\partial/\partial t_j$  on  $\mathcal{A}_0 \cong C^\infty(\mathbb{T}^2)$ . Also note that there is a noncommutative ‘‘Stokes’ Theorem’’ in the sense that  $\tau(\delta_j(a)) = 0$ .

**Lemma 5.18** *The map  $a \mapsto \delta_j a$  for  $a \in \mathcal{A}_\theta$  extends to a closed (unbounded) skew-adjoint operator on the Hilbert space  $\mathcal{H}_\tau$ .*

**Proof** For this it is sufficient to note that

$$\tau((\delta_j a)^* b) = \tau(\delta_j(a^* b)) - \tau(a^* \delta_j(b)) = -\tau(a^* \delta_j(b)). \quad \square$$

This result allows us to give the following definition, which is inspired of course by the commutative case of Example 5.12.

**Definition 5.19** *The Dirac operator on the noncommutative torus* is the symmetric operator  $D_{\mathcal{A}_\theta} : \mathcal{A}_\theta \otimes \mathbb{C}^2 \rightarrow \mathcal{H}$  defined by

$$D_{\mathcal{A}_\theta} = \begin{pmatrix} 0 & -\delta_1 - i\delta_2 \\ \delta_1 - i\delta_2 & 0 \end{pmatrix}$$

One computes (as in Example 5.12) that an orthonormal eigenbasis of  $D_{\mathcal{A}_\theta}$  is given by

$$\psi_{rs} = \frac{1}{\sqrt{2}} \left( \pm \frac{u^r v^s}{\sqrt{r^2 + s^2}} u^r v^s \right); \quad (r, s \in \mathbb{Z}),$$

with eigenvalues  $\pm\sqrt{r^2 + s^2}$ . Since this coincides with the classical spectrum of  $D_{\mathbb{T}^2}$  on  $L^2(\mathbb{T}^2) \otimes \mathbb{C}^2$  we may conclude that the resolvent of  $D_{\mathcal{A}_\theta}$  is compact. Note that this would also follow from Exercise 4.9. Finally, we compute for any  $a \in \mathcal{A}_\theta$  that

$$[D_{\mathcal{A}_\theta}, \pi(a)] = \begin{pmatrix} 0 & \pi(-\delta_1(a) - i\delta_2(a)) \\ \pi(\delta_1(a) - i\delta_2(a)) & 0 \end{pmatrix}$$

which extends to a bounded operator on  $\mathcal{H}$ . We have thus proven the following result.

**Proposition 5.20** *The data  $(\mathcal{A}_\theta, \mathcal{H}, D_{\mathcal{A}_\theta}; J, \gamma)$  defined above is a real spectral triple of KO-dimension 2.*

### 5.3.2 Generalization to Toric Manifolds

Let  $M$  be an  $m$  dimensional compact Riemannian manifold equipped with an isometric smooth action of an  $n$ -torus  $\mathbb{T}^n$ ,  $n \geq 2$ . We denote by  $\sigma$  the corresponding action of  $\mathbb{T}^n$  by automorphisms—obtained by pull-backs—on the algebra  $C^\infty(M)$  of smooth functions on  $M$ .

The algebra  $C^\infty(M)$  may be decomposed into spectral subspaces which are indexed by the dual group  $\mathbb{Z}^n = \widehat{\mathbb{T}}^n$ . Now, with  $s = (s_1, \dots, s_n) \in \mathbb{T}^n$ , each  $r \in \mathbb{Z}^n$  yields a character of  $\mathbb{T}^n$ ,  $s \mapsto e^{2\pi i r \cdot s}$ , with the scalar product  $r \cdot s := r_1 s_1 + \dots + r_n s_n$ . The  $r$ th spectral subspace for the action  $\sigma$  of  $\mathbb{T}^n$  on  $C^\infty(M)$  consists of those smooth functions  $f_r$  for which

$$\sigma_s(f_r) = e^{2\pi i r \cdot s} f_r, \quad (5.3.4)$$

and each  $f \in C^\infty(M)$  is the sum of a unique series  $f = \sum_{r \in \mathbb{Z}^n} f_r$ , which is rapidly convergent in the Fréchet topology of  $C^\infty(M)$  (see reference in Note 12 below for more details).

Let now  $\theta = (\theta_{jk} = -\theta_{kj})$  be a real antisymmetric  $n \times n$  matrix. The  $\theta$ -deformation of  $C^\infty(M)$  may be defined by replacing the ordinary product by a deformed product, given on spectral subspaces by

$$f_r \times_\theta g_{r'} := f_r \sigma_{\frac{1}{2}r \cdot \theta}(g_{r'}) = e^{\pi i r \cdot \theta \cdot r'} f_r g_{r'}, \quad (5.3.5)$$

where  $r \cdot \theta$  is the element in  $\mathbb{R}^n$  with components  $(r \cdot \theta)_k = \sum r_j \theta_{jk}$  for  $k = 1, \dots, n$ . The product in (5.3.5) is then extended linearly to all functions in  $C^\infty(M)$ . We denote the space  $C^\infty(M)$  endowed with the product  $\times_\theta$  by  $C^\infty(M_\theta)$ . The action  $\sigma$  of  $\mathbb{T}^n$  on  $C^\infty(M)$  extends to an action on  $C^\infty(M_\theta)$  given again by (5.3.4) on the homogeneous elements.

Next, let us take  $M$  to be a spin manifold with  $L^2(S)$  the Hilbert space of spinors and  $D_M$  the usual Dirac operator of the metric of  $M$ . Smooth functions act on spinors by pointwise multiplication thus giving a representation  $\pi : C^\infty(M) \rightarrow \mathcal{B}(L^2(S))$ .

We assume that there is a double cover  $c : \tilde{\mathbb{T}}^n \rightarrow \mathbb{T}^n$  and a representation of  $\tilde{\mathbb{T}}^n$  on  $L^2(S)$  by unitary operators  $U(s)$ ,  $s \in \tilde{\mathbb{T}}^n$ , so that

$$U(s) D_M U(s)^{-1} = D_M, \quad (5.3.6)$$

since the torus action is assumed to be isometric, and such that for all  $f \in C^\infty(M)$ ,

$$U(s) \pi(f) U(s)^{-1} = \pi(\sigma_{c(s)}(f)). \quad (5.3.7)$$

We say that an element  $T \in \mathcal{B}(L^2(S))$  is called smooth for the action of  $\tilde{\mathbb{T}}^n$  if the map

$$\tilde{\mathbb{T}}^n \ni s \mapsto \alpha_s(T) := U(s) T U(s)^{-1},$$

is smooth for the norm topology. From its very definition,  $\alpha_s$  coincides on  $\pi(C^\infty(M)) \subset \mathcal{B}(L^2(S))$  with the automorphism  $\sigma_{c(s)}$ . Moreover, much as it was done before for the smooth functions, we shall use the torus action to give a spectral decomposition of smooth elements of  $\mathcal{B}(L^2(S))$ . Any such a smooth element  $T$  is written as a (rapidly convergent) series  $T = \sum T_r$  with  $r \in \mathbb{Z}^n$  and each  $T_r$  is homogeneous of degree  $r$  under the action of  $\tilde{\mathbb{T}}^n$ , i.e.

$$\alpha_s(T_r) = e^{2\pi i r \cdot s} T_r, \quad \forall s \in \tilde{\mathbb{T}}^n. \quad (5.3.8)$$

Let  $(P_1, P_2, \dots, P_n)$  be the infinitesimal generators of the action of  $\tilde{\mathbb{T}}^n$  so that we can write  $U(s) = \exp 2\pi i s \cdot P$ . Now, with  $\theta$  a real  $n \times n$  anti-symmetric matrix as above, one defines a twisted representation of the smooth elements of  $\mathcal{B}(L^2(S))$  on  $L^2(S)$  by

$$L_\theta(T) := \sum_r T_r U(\frac{1}{2}r \cdot \theta) = \sum_r T_r \exp \{ \pi i r_j \theta_{jk} P_k \}, \quad (5.3.9)$$

Taking smooth functions on  $M$  as elements of  $\mathcal{B}(L^2(S))$ , via the representation  $\pi$ , the previous definition gives an algebra  $L_\theta(C^\infty(M))$  which we may think of as a representation of the algebra  $C^\infty(M_\theta)$ . Indeed, by the very definition of the product  $\times_\theta$  in (5.3.5) one establishes that

$$L_\theta(f \times_\theta g) = L_\theta(f)L_\theta(g), \quad (5.3.10)$$

proving that the algebra  $C^\infty(M_\theta)$  (i.e.  $C^\infty(M)$  equipped with the product  $\times_\theta$ ) is isomorphic to the algebra  $L_\theta(C^\infty(M))$ .

**Theorem 5.21** *The datum  $(C^\infty(M_\theta), L^2(S), D_M)$  is a spectral triple.*

**Proof** The resolvent of  $D_M$  is compact by assumption. The boundedness of the commutators  $[D_M, L_\theta(f)]$  for  $f \in C^\infty(M)$  follows from the relation  $[D_M, L_\theta(f)] = L_\theta([D_M, f])$ ,  $D_M$  being of degree 0 since  $\mathbb{T}^n$  acts by isometries, so that each  $P_k$  commutes with  $D_M$ . See also Note 13 in this chapter.  $\square$

This noncommutative Riemannian spin manifold is a so-called isospectral deformation of the classical Riemannian geometry of  $M$ , in that the spectrum of the operator  $D_M$  coincides with that of the classical Dirac operator on  $M$ . Moreover, if  $M$  is even and spin then there is a grading  $\gamma_M$  and operator  $J_M$  that make  $(C^\infty(M_\theta), L^2(S), D_M; J_M, \gamma_M)$  a real spectral triple.

## Notes

### Section 5.1. Gelfand Duality

1. A complete treatment of  $C^*$ -algebras, their representation theory and Gelfand duality can be found in [1, Sect. II.2.2] or [2, Sect. I.4].
2. A proof of Proposition 5.4 can be found in [2, Theorem 11.2].
3. A proof of Theorem 5.7 can be found in e.g. [1, Theorem II.2.2.4] or [2, Theorem 3.11].
4. Spectral triples were introduced by Connes in the early 1980s. See [3, Sect. IV.2.8] (where they were called unbounded  $K$ -cycles) and [4].
5. The distance formula appearing in Proposition 5.8, as well as the proof of this Proposition can be found in [3, Sect. VI.1]. Moreover, it extends to a distance formula on the state space  $S(A)$  of a  $C^*$ -algebra  $A$  as follows. Recall that a linear functional  $\omega : A \rightarrow \mathbb{C}$  is a state if it

is positive  $\omega(a^*a) > 0$  for all non-zero  $a \in A$ , and such that  $\omega(1) = 1$ . One then defines a distance function on  $S(A)$  by [4]

$$d(\omega_1, \omega_2) = \sup_{a \in A} \{|\omega_1(a) - \omega_2(a)| : \| [D, a] \| \leq 1\}.$$

It is noted in [5, 6] that this distance formula, in the case of locally compact complete manifolds, is in fact a reformulation of the Wasserstein distance in the theory of optimal transport. We also refer to [7–9].

- 6. Proposition 5.8 establishes that from the canonical triple on a Riemannian spin manifold  $M$  one can reconstruct the Riemannian distance on  $M$ . As a matter of fact, there is a reconstruction theorem for the smooth manifold structure of  $M$  as well [10]. It states that if  $(\mathcal{A}, \mathcal{H}, D; J, \gamma)$  is a real spectral triple with  $\mathcal{A}$  commutative, then under suitable conditions [11] there is a Riemannian spin manifold  $(M, g)$  with spin structure  $(S, J_M)$  such that  $(\mathcal{A}, \mathcal{H}, D; J, \gamma)$  is given by  $(C^\infty(M), L^2(S), \mathcal{D}; J_M, \gamma_M)$  (see also the discussion in [12, Sect. 11.4]).
- 7. Real spectral triples as defined in Definition 5.9 are noncommutative generalization of Riemannian spin manifolds. An immediate question that arises is whether noncommutative generalizations of Riemannian spin<sup>c</sup> manifolds, or even just Riemannian manifolds can be defined. In fact, building on the algebraic approach to defining spin<sup>c</sup> manifolds as in [12] (as also adopted above) the authors [13] introduce such noncommutative analogues. For earlier attempts, refer to [14].
- 8. Products of spectral triples are described in detail in [15], and generalized to include the odd case as well in [16].
- 9. The differential calculi that are associated to any spectral triple are explained in [3, Sect. VI.1] (see also [17, Chap. 7]).

### Section 5.3. Examples of Noncommutative Manifolds

- 10. The noncommutative torus formed the guiding example for noncommutative manifolds in the early days of noncommutative geometry, and already appears in [18]. The deformation quantization aspects of the noncommutative torus were analyzed early on as well, by Rieffel in [19].

The  $C^*$ -algebra  $A_\theta$  is also called the rotation algebra. This is because of the following realization as operators on  $L^2(\mathbb{S}^1)$ . We consider for  $\psi \in L^2(\mathbb{S}^1)$ :

$$U\psi(z) = z\psi(z); \quad V\psi(z) = \psi(\lambda z).$$

Thus,  $U$  generated the  $C^*$ -algebras  $C(S^1)$  and conjugation by  $V$  gives an automorphism  $\alpha$  of  $C(S^1)$  by rotation with the angle  $\theta$ . The association  $u \mapsto U, v \mapsto V$  is a representation of the  $C^*$ -algebra  $A_\theta$  on  $L^2(\mathbb{S}^1)$ , which in fact implements an isomorphism  $A_\theta \cong C(S^1) \rtimes_\alpha \mathbb{Z}$  with a crossed product algebra associated to the rotation  $\alpha$ .

- 11. The Gelfand–Naimark–Segal (or, GNS) construction is a general procedure that constructs a Hilbert space representation of a  $C^*$ -algebra, starting with a given state on it. More details can be found e.g. in [1, Sect. II.6.4].
- 12. It is shown in [20] that there is a natural completion of the algebra  $C^\infty(M_\theta)$  to a  $C^*$ -algebra  $C(M_\theta)$  whose smooth subalgebra—under the extended action of  $\mathbb{T}^n$ —is precisely  $C^\infty(M_\theta)$ . Thus, we can understand  $L_\theta$  as a *quantization map* from

$$L_\theta : C^\infty(M) \rightarrow C^\infty(M_\theta), \tag{5.3.11}$$

which provides a strict deformation quantization in the sense of Rieffel. More generally, he considers a (not necessarily commutative)  $C^*$ -algebra  $A$  carrying an action of  $\mathbb{R}^n$ . For an anti-symmetric  $n \times n$  matrix  $\theta$ , one defines a star product  $\times_\theta$  between elements in  $A$  much as we did before. The algebra  $A$  equipped with the product  $\times_\theta$  gives rise to a  $C^*$ -algebra denoted by  $A_\theta$ . Then the collection  $\{A_{\hbar\theta}\}_{\hbar \in [0,1]}$  is a continuous family of  $C^*$ -algebras providing a strict deformation quantization in the direction of the Poisson structure on  $A$  defined by the matrix  $\theta$ .

13. Theorem 5.21 was obtained in [21]; see also [22].

## References

1. Blackadar, B.: Operator algebras, Theory of  $C^*$ -algebras and von Neumann algebras. Operator Algebras and Non-commutative Geometry, III, Encyclopaedia of Mathematical Sciences, vol. 122. Springer, Berlin (2006)
2. Takesaki, M.: Theory of operator algebras. I, Encyclopaedia of Mathematical Sciences, vol. 22. Springer, Berlin (2002). Reprint of the first (1979) edition, Operator Algebras and Non-commutative Geometry, 5
3. Connes, A.: Noncommutative Geometry. Academic, San Diego (1994)
4. Connes, A.: Noncommutative geometry and reality. J. Math. Phys. **36**(11), 6194–6231 (1995)
5. Rieffel, M.A.: Metrics on state spaces. Doc. Math. **4**, 559–600 (1999)
6. D’Andrea, F., Martinetti, P.: A view on optimal transport from noncommutative geometry. SIGMA Symmetry Integrability Geom. Methods Appl. **6** (2010) Paper 057, 24
7. D’Andrea, F., Martinetti, P.: On Pythagoras theorem for products of spectral triples. Lett. Math. Phys. **103**, 469–492 (2013)
8. Martinetti, P.: Carnot-Carathéodory metric and gauge fluctuation in noncommutative geometry. Commun. Math. Phys. **265**, 585–616 (2006)
9. Martinetti, P., Wulkenhaar, R.: Discrete Kaluza-Klein from scalar fluctuations in noncommutative geometry. J. Math. Phys. **43**, 182–204 (2002)
10. Connes, A.: On the spectral characterization of manifolds. J. Noncommut. Geom. **7**, 1–82 (2013)
11. Connes, A.: Gravity coupled with matter and the foundation of non-commutative geometry. Commun. Math. Phys. **182**, 155–176 (1996)
12. Gracia-Bondía, J.M., Várilly, J.C., Figueroa, H.: Elements of Noncommutative Geometry. Birkhäuser, Boston (2001)
13. Lord, S., Rennie, A., Várilly, J.C.: Riemannian manifolds in noncommutative geometry. J. Geom. Phys. **62**, 1611–1638 (2012)
14. Frohlich, J., Grandjean, O., Recknagel, A.: Supersymmetric Quantum Theory and Non-Commutative Geometry. Commun. Math. Phys. **193** (1998)
15. Vanhecke, F.J.: On the product of real spectral triples. Lett. Math. Phys. **50**, 157–162 (1999)
16. Dabrowski, L., Dossena, G.: Product of real spectral triples. Int. J. Geom. Meth. Mod. Phys. **8**, 1833–1848 (2011)
17. Landi, G.: An Introduction to Noncommutative Spaces and their Geometry. Springer (1997)
18. Connes, A.:  $C^*$ -algèbres et géométrie différentielle. C.R. Acad. Sci. Paris Ser. A-B **290**, A599–A604 (1980)
19. Rieffel, M.A.:  $C^*$ -algebras associated with irrational rotations. Pac. J. Math. **93**, 415–429 (1981)
20. Rieffel, M.A.: Deformation quantization for actions of  $\mathbb{R}^d$ . Mem. Am. Math. Soc. **506** (1993)
21. Connes, A., Landi, G.: Noncommutative manifolds: The instanton algebra and isospectral deformations. Commun. Math. Phys. **221**, 141–159 (2001)
22. Connes, A., Dubois-Violette, M.: Noncommutative finite-dimensional manifolds. I. Spherical manifolds and related examples. Commun. Math. Phys. **230**, 539–579 (2002)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 6

## The Local Index Formula in Noncommutative Geometry



In this chapter we present a proof of the Connes–Moscovici index formula, expressing the index of a (twisted) operator  $D$  in a spectral triple  $(\mathcal{A}, \mathcal{H}, D)$  by a local formula. First, we illustrate the contents of this chapter in the context of two examples in the odd and even case: the index on the circle and on the torus.

### 6.1 Local Index Formula on the Circle and on the Torus

#### 6.1.1 The Winding Number on the Circle

Consider the canonical triple on the circle (Example 5.11):

$$\left( C^\infty(\mathbb{S}^1), L^2(\mathbb{S}^1), D_{\mathbb{S}^1} = -i \frac{d}{dt} \right).$$

The eigenfunctions of  $D_{\mathbb{S}^1}$  are given for any  $n \in \mathbb{Z}$  by  $e_n(t) = e^{int}$ , where  $t \in [0, 2\pi]$ . Indeed,  $D_{\mathbb{S}^1} e_n = n e_n$  and  $\{e_n\}_{n \in \mathbb{Z}}$  forms an orthonormal basis for  $L^2(\mathbb{S}^1)$ . We denote the projection onto the non-negative eigenspace of  $D_{\mathbb{S}^1}$  by  $P$ , i.e.

$$P e_n = \begin{cases} e_n & \text{if } n \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

This is equivalent to defining  $P = (1 + F)/2$ , where  $F = D_{\mathbb{S}^1} |D_{\mathbb{S}^1}|^{-1}$  (defined to be  $+1$  on  $\ker D_{\mathbb{S}^1}$ ). Concretely,  $F$  is the **Hilbert transform**:

$$F \left( \sum_{n \in \mathbb{Z}} \psi_n e_n(t) \right) = - \sum_{n < 0} \psi_n e_n + \sum_{n \geq 0} \psi_n e_n,$$

with complex coefficients  $\psi_n$  ( $n \in \mathbb{Z}$ ).

Let  $u$  be a unitary in  $C^\infty(\mathbb{S}^1)$ , say  $u = e_m$  for some  $m \in \mathbb{Z}$ . The index we are interested in is given by the difference between the dimensions of the kernel and cokernel of  $PuP : PL^2(\mathbb{S}^1) \rightarrow PL^2(\mathbb{S}^1)$ :

$$\text{index } PuP = \dim \ker PuP - \dim \ker Pu^*P.$$

Indeed,  $\text{Im } T^\perp = \ker T^*$  for any bounded operator. We wish to write this index as a local, integral expression. First, we check that the index is well defined by noting that  $PuP$  has finite-dimensional kernel and cokernel. In fact, the kernel of  $PuP$  (with  $u = e_m$ ) consists of  $\psi = \sum_{n \geq 0} \psi_n e_n \in PL^2(\mathbb{S}^1)$  such that

$$P \left( \sum_{n \geq 0} \psi_n e_{m+n} \right) = 0.$$

In other words, the kernel of  $PuP$  consists of linear combinations of the vectors  $e_0, \dots, e_{-m-1}$  for  $m < 0$ . We conclude that  $\dim \ker PuP = -m$  if  $m < 0$ . If  $m > 0$  then this dimension is zero, but in that case  $\dim \ker Pu^*P = m$ . In both cases, and also in the remaining case  $m = 0$ , for  $u = e_m$  we find that

$$\text{index } PuP = -m.$$

**Exercise 6.1** In this exercise we show that  $\text{index } PuP$  is well defined for any unitary  $u \in C^\infty(\mathbb{S}^1)$ .

1. Show that  $[F, e_m]$  is a compact operator for any  $m \in \mathbb{Z}$ .
2. Show that  $[F, f]$  is a compact operator for any function  $f = \sum_n f_n e_n \in C^\infty(\mathbb{S}^1)$  (convergence is in sup-norm).
3. Atkinson's Theorem states that an operator is Fredholm (*i.e.* has finite kernel and cokernel) if it is invertible modulo compact operators. Use this to show that  $PuP$  is a Fredholm operator.


On the other hand, we can compute the following zeta function given by the trace (taken for simplicity over the complement of  $\ker D_{\mathbb{S}^1}$ ):

$$\text{Tr} (u^*[D_{\mathbb{S}^1}, u] |D_{\mathbb{S}^1}|^{-2z-1}) = m \text{Tr} |D_{\mathbb{S}^1}|^{-2z-1} = 2m\zeta(1+2z),$$

since  $[D_{\mathbb{S}^1}, u] = mu$  for  $u = e_m$ . Here  $\zeta(s)$  is the well-known *Riemann zeta function*. Since  $\zeta(s)$  has a pole at  $s = 1$ , we conclude that

$$\text{index } PuP = -\text{res}_{z=0} \text{Tr} (u^*[D_{\mathbb{S}^1}, u] |D_{\mathbb{S}^1}|^{-2z-1}).$$

**Fig. 6.1** The map  
 $e_m : t \in [0, 2\pi) \mapsto e^{imt}$   
winds  $m$  times around the  
circle; this winding number  
is (minus) the index of the  
operator  $P e_m P$



This is a manifestation of the noncommutative index formula in the simple case of the circle, expressing the winding number  $m$  (cf. Fig. 6.1) of the unitary  $u = e_m$  as a ‘local’ expression. In fact,

$$\text{res}_{z=0} \text{Tr} (u^*[D_{\mathbb{S}^1}, u] |D_{\mathbb{S}^1}|^{-2z-1}) = \frac{1}{2\pi i} \int_{\mathbb{S}^1} u^* du,$$

as one can easily check. The right-hand side is indeed a local integral expression for the (global) index of  $P u P$ .

In this chapter, we generalize this formula to any (odd) spectral triple, translating this locality to the appropriate algebraic notion, namely, in terms of cyclic cocycles.

**Exercise 6.2** Prove the following index formula, for a unitary  $u = e_m$ , say, with  $m < 0$ :

$$\text{index } P u P = -\frac{1}{4} \text{Tr } F[F, u^*][F, u].$$

### 6.1.2 The Winding Number on the Torus

The same winding number—now in one of the two circle directions—can also be obtained as an index on the two-dimensional torus, as we will now explain.

Consider the even canonical triple on the 2-dimensional torus (Example 5.12):

$$\left( C^\infty(\mathbb{T}^2), L^2(\mathbb{T}^2) \otimes \mathbb{C}^2, D_{\mathbb{T}^2} = \begin{pmatrix} 0 & -\partial_1 - i\partial_2 \\ \partial_1 - i\partial_2 & 0 \end{pmatrix} \right).$$

The eigenspinors of  $D_{\mathbb{T}^2}$  are given by the vectors

$$\phi_{n_1, n_2}^\pm(t_1, t_2) := \frac{1}{\sqrt{2}} \left( \begin{array}{c} e^{i(n_1 t_1 + n_2 t_2)} \\ \pm \frac{i n_1 + n_2}{\sqrt{n_1^2 + n_2^2}} e^{i(n_1 t_1 + n_2 t_2)} \end{array} \right); \quad (n_1, n_2 \in \mathbb{Z}),$$

with eigenvalues  $\pm \sqrt{n_1^2 + n_2^2}$ .

Instead of unitaries, we now consider orthogonal projections  $p \in C^\infty(\mathbb{T}^2)$  or rather, projections in matrix algebras with entries in  $C^\infty(\mathbb{T}^2)$ . Indeed, there are no non-trivial projections  $p$  in  $C(\mathbb{T}^2)$ : a continuous function with the property  $p^2 = p$  is automatically 0 or 1. Thus, we consider the following class of orthogonal projections in  $M_2(C^\infty(\mathbb{T}^2))$ :

$$p = \begin{pmatrix} f & g + hU^* \\ g + hU & 1 - f \end{pmatrix}, \quad (6.1.1)$$

where  $f, g, h$  are real-valued (periodic) functions of the first variable  $t_1$ , and  $U$  is a unitary depending only on the second variable  $t_2$ , say  $U(t_2) = e_m(t_2)$ . The projection property  $p^2 = p$  translates into the two conditions

$$gh = 0, \quad g^2 + h^2 = f - f^2.$$

A possible solution of these relations is given by

$$0 \leq f \leq 1 \text{ such that } f(0) = 1, \quad f(\pi) = 0,$$

and then  $g = \chi_{[0, \pi]} \sqrt{f - f^2}$  and  $h = \chi_{[\pi, 2\pi]} \sqrt{f - f^2}$ , where  $\chi_X$  is the indicator function for the set  $X$  (see Fig. 6.2).

The Fredholm operator we would like to compute the index of is  $p(D_{\mathbb{T}^2} \otimes \mathbb{I}_2)p$ , acting on the doubled spinor Hilbert space  $L^2(S) \otimes \mathbb{C}^2 \simeq L^2(\mathbb{T}^2) \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ . This doubling is due to the fact that we take a  $2 \times 2$  matricial projection. To avoid notation cluttery, we will simply write  $D_{\mathbb{T}^2}$  for  $D_{\mathbb{T}^2} \otimes \mathbb{I}_2$ .

The local index formula which we would like to illustrate on the torus is

$$\text{index } p D_{\mathbb{T}^2} p = -\text{res}_{z=0} \text{Tr} \left( \gamma \left( p - \frac{1}{2} \right) [D_{\mathbb{T}^2}, p] [D_{\mathbb{T}^2}, p] |D_{\mathbb{T}^2}|^{-2-2z} \right),$$


where the trace is both over the matrix indices of  $p$  and over the spinor indices.

**Proposition 6.1** *With  $U(t_2) = e_m(t_2)$  and  $p$  of the above form, we have*

$$\text{res}_{z=0} \text{Tr} \left( \gamma \left( p - \frac{1}{2} \right) [D_{\mathbb{T}^2}, p] [D_{\mathbb{T}^2}, p] |D_{\mathbb{T}^2}|^{-2-2z} \right) = m.$$

**Proof** We use the following formula from Exercise 6.3, which holds for any  $F \in C^\infty(\mathbb{T}^2)$ :

$$\text{Tr } F |D_{\mathbb{T}^2}|^{-2s} = \frac{\zeta_E(s)}{\pi} \int_{\mathbb{T}^2} F, \quad (6.1.2)$$



**Fig. 6.2** Functions  $f, g, h$  that ensure that  $p$  in (6.1.1) is a projection

where the trace is over spinor indices, and where  $\zeta_E$  is the Epstein zeta function, defined by

$$\zeta_E(s) = \sum_{n_1, n_2 \in \mathbb{Z}} (n_1^2 + n_2^2)^{-s}.$$

Since  $\zeta_E$  has a pole at  $s = 1$  with residue  $\pi$ , we conclude that

$$\text{res}_{z=0} \text{Tr } F |D|^{-2-2z} = \int_{\mathbb{T}^2} F.$$

Returning to the claimed equality, we compute the trace over spinor indices:

$$\begin{aligned} \text{Tr } \gamma(p - \frac{1}{2}) [D_{\mathbb{T}^2}, p]^2 &= \text{Tr}(p - \frac{1}{2}) \begin{pmatrix} 0 & -\partial_1 p - i\partial_2 p \\ \partial_1 p - i\partial_2 p & 0 \end{pmatrix}^2 \\ &= 2i(p - \frac{1}{2})(\partial_1 p \partial_2 p - \partial_2 p \partial_1 p). \end{aligned}$$

Since  $g$  and  $h$  in (6.1.1) have disjoint support,  $g'h = 0$ , we have

$$\partial_1 p \partial_2 p = -\partial_2 p \partial_1 p = -im \begin{pmatrix} -hh' & f'hU^* \\ f'hU & hh' \end{pmatrix}.$$

Hence, taking the remaining trace over the indices of the projection, we find

$$\text{Tr } 2i(p - \frac{1}{2})(\partial_1 p \partial_2 p - \partial_2 p \partial_1 p) = 4m(-2fhh' + hh' + 2f'h^2).$$

Inserting this back in (6.1.2) we see that we have to integrate the right-hand side over the circle. A series of partial integrations yields

$$\frac{1}{2\pi} \int -2fhh' + hh' + 2f'h^2 = \frac{1}{2\pi} \int 3f'h^2.$$

Inserting the explicit expression of  $h$ , we easily determine

$$\int f'h^2 = \int_{\pi}^{2\pi} (f - f^2)f' = \int_0^1 (x - x^2)dx = \frac{1}{6}.$$


Combining all coefficients, including the residue of Epstein's zeta function, we finally find

$$\text{res}_{z=0} \text{Tr} \left( \gamma \left( p - \frac{1}{2} \right) [D_{\mathbb{T}^2}, p]^2 |D_{\mathbb{T}^2}|^{-2-2z} \right) = 4m \frac{3}{2\pi} \frac{1}{6}\pi = m,$$

as required.  $\square$

Thus, we recover the winding number of the unitary  $U$ , winding  $m$  times around one of the circle directions in  $\mathbb{T}^2$ , just as in the previous subsection. The case  $m = 2$  is depicted in Fig. 6.3; it shows the winding of the range of  $p$  in  $\mathbb{C}^2$  at  $t_1 = 3\pi/4$  and with  $t_2$  varying from 0 to  $2\pi$ .

The fact that the index of  $pD_{\mathbb{T}^2}p$  is also equal to (minus) this winding number is highly non-trivial and much more difficult to prove. Therefore, already this simple example illustrates the power of the Connes–Moscovici index formula, expressing the index by a local formula. We will now proceed and give a proof of the local index formula for any spectral triple.



**Fig. 6.3** Winding twice around one of the circle directions on the torus. Let the range of the projection  $p$  be  $v(t_1, t_2)s$  with  $s \in \mathbb{C}$  and  $v(t_1, t_2) \in \mathbb{C}^2$  varies with  $(t_1, t_2) \in \mathbb{T}^2$ . We have drawn the real and imaginary parts of the first component  $v_1(t_1 = 3\pi/4, t_2)s$  with  $0 \leq t_2 \leq 2\pi$  and  $-1 \leq s \leq 1$ . The other component  $v_2(t_1 = 3\pi/4, t_2)$  is constant

**Exercise 6.3** Prove Eq. (6.1.2), *i.e.* show that for any function  $F \in C^\infty(\mathbb{T}^2)$  we have

$$\mathrm{Tr} F |D_{\mathbb{T}^2}|^{-2s} = \frac{\zeta_E(s)}{\pi} \int_{\mathbb{T}^2} F.$$

## 6.2 Hochschild and Cyclic Cohomology

We introduce cyclic cohomology, which can be seen as a noncommutative generalization of De Rham homology.

**Definition 6.2** If  $\mathcal{A}$  is an algebra, we define the space of *n-cochains*, denoted by  $C^n(\mathcal{A})$ , as the space of  $(n+1)$ -linear functionals on  $\mathcal{A}$  with the property that if  $a^j = 1$  for some  $j \geq 1$ , then  $\phi(a^0, \dots, a^n) = 0$ . Define operators  $b : C^n(\mathcal{A}) \rightarrow C^{n+1}(\mathcal{A})$  and  $B : C^{n+1}(\mathcal{A}) \rightarrow C^n(\mathcal{A})$  by

$$\begin{aligned} b\phi(a^0, a^1, \dots, a^{n+1}) &:= \sum_{j=0}^n (-1)^j \phi(a^0, \dots, a^j a^{j+1}, \dots, a^{n+1}) \\ &\quad + (-1)^{n+1} \phi(a^{n+1} a^0, a^1, \dots, a^n), \\ B\phi(a^0, a^1, \dots, a^n) &:= \sum_{j=0}^n (-1)^{nj} \phi(1, a^j, a^{j+1}, \dots, a^{j-1}). \end{aligned}$$

**Exercise 6.4** Show that  $b^2 = 0$ ,  $B^2 = 0$ , and  $bB + Bb = 0$ .

This means that a cochain which is in the image of  $b$  is also in the kernel of  $b$ , and similarly for  $B$ . We say that  $b$  and  $B$  define *complexes* of cochains

$$\begin{array}{ccccccc} \cdots & \xrightarrow{b} & C^n(\mathcal{A}) & \xrightarrow{b} & C^{n+1}(\mathcal{A}) & \xrightarrow{b} & \cdots \\ \cdots & \xleftarrow{B} & C^n(\mathcal{A}) & \xleftarrow{B} & C^{n+1}(\mathcal{A}) & \xleftarrow{B} & \cdots, \end{array}$$

where the maps have the (complex) defining property that composing them gives zero:  $b \circ b = 0 = B \circ B$ . This property of  $b$  and  $B$  being a differential is a crucial ingredient in *cohomology*, where so-called *cohomology groups* are defined as the quotients of the kernel by the image of the differential. In our case, we have

**Definition 6.3** The *Hochschild cohomology* of  $\mathcal{A}$  is given by the quotients

$$HH^n(\mathcal{A}) = \frac{\ker b : C^n(\mathcal{A}) \rightarrow C^{n+1}(\mathcal{A})}{\mathrm{Im} b : C^{n-1}(\mathcal{A}) \rightarrow C^n(\mathcal{A})}; \quad (n \geq 0).$$

Elements in  $\ker b : C^n(\mathcal{A}) \rightarrow C^{n+1}(\mathcal{A})$  are called *Hochschild n-cocycles*, and elements in  $\mathrm{Im} b : C^{n-1}(\mathcal{A}) \rightarrow C^n(\mathcal{A})$  are called *Hochschild n-coboundaries*.

- Exercise 6.5** (1) Characterize the cohomology group  $HH^0(\mathcal{A})$  for any algebra  $\mathcal{A}$ .  
 (2) Compute  $HH^n(\mathbb{C})$  for any  $n \geq 0$ .  
 (3) Establish the following functorial property of  $HH^n$ : if  $\psi : \mathcal{A} \rightarrow \mathcal{B}$  is an algebra map, then there is a homomorphism of groups  $\psi^* : HH^n(\mathcal{B}) \rightarrow HH^n(\mathcal{A})$ .

**Example 6.4** Let  $M$  be a compact  $n$ -dimensional manifold without boundary. The following expression defines an  $n$ -cochain on  $\mathcal{A} = C^\infty(M)$ :

$$\phi(f_0, f_1, \dots, f_n) = \int_M f_0 df_1 \cdots df_n.$$

In fact, one can compute that  $b\phi = 0$  so that this is an  $n$ -cocycle which defines a class in the Hochschild cohomology group  $HH^n(C^\infty(M))$ .

**Exercise 6.6** Check that  $b\phi = 0$  in the above example.

Next, we turn our attention to the differential  $B$ , and its compatibility with  $b$ . Namely,  $b$  and  $B$  define a so-called *double complex*:

$$\begin{array}{ccccccc} & \vdots & & \vdots & & \vdots & \\ & b \uparrow & & b \uparrow & & b \uparrow & \\ \dots & \xrightarrow{B} & C^3(\mathcal{A}) & \xrightarrow{B} & C^2(\mathcal{A}) & \xrightarrow{B} & C^1(\mathcal{A}) \xrightarrow{B} C^0(\mathcal{A}) \\ & b \uparrow & & b \uparrow & & b \uparrow & \\ & \vdots & & \vdots & & \vdots & \\ \dots & \xrightarrow{B} & C^2(\mathcal{A}) & \xrightarrow{B} & C^1(\mathcal{A}) & \xrightarrow{B} & C^0(\mathcal{A}) \\ & b \uparrow & & b \uparrow & & b \uparrow & \\ & \vdots & & \vdots & & \vdots & \\ \dots & \xrightarrow{B} & C^1(\mathcal{A}) & \xrightarrow{B} & C^0(\mathcal{A}) & & \\ & b \uparrow & & & & & \\ & \vdots & & & & & \\ \dots & \xrightarrow{B} & C^0(\mathcal{A}) & & & & \end{array}$$

The *totalization* of this double complex by definition consists of the even and odd cochains:

$$\begin{aligned} C^{\text{ev}}(\mathcal{A}) &= \bigoplus_k C^{2k}(\mathcal{A}); \\ C^{\text{odd}}(\mathcal{A}) &= \bigoplus_k C^{2k+1}(\mathcal{A}), \end{aligned}$$

and these also form a complex, now with differential  $b + B$ :

$$\dots \xrightarrow{b+B} C^{\text{ev}}(\mathcal{A}) \xrightarrow{b+B} C^{\text{odd}}(\mathcal{A}) \xrightarrow{b+B} C^{\text{ev}}(\mathcal{A}) \xrightarrow{b+B} \dots$$

**Definition 6.5** The *periodic cyclic cohomology* of  $\mathcal{A}$  is the cohomology of the totalization of this complex. That is, the even and odd cyclic cohomology groups are given by

$$\begin{aligned} HCP^{\text{ev}}(\mathcal{A}) &= \frac{\ker b + B : C^{\text{ev}}(\mathcal{A}) \rightarrow C^{\text{odd}}(\mathcal{A})}{\text{Im } b + B : C^{\text{odd}}(\mathcal{A}) \rightarrow C^{\text{ev}}(\mathcal{A})}, \\ HCP^{\text{odd}}(\mathcal{A}) &= \frac{\ker b + B : C^{\text{odd}}(\mathcal{A}) \rightarrow C^{\text{ev}}(\mathcal{A})}{\text{Im } b + B : C^{\text{ev}}(\mathcal{A}) \rightarrow C^{\text{odd}}(\mathcal{A})}. \end{aligned}$$

Elements in  $\ker b + B$  are called (even or odd)  $(b, B)$ -cocycles, and elements in  $\text{Im } b + B$  are called (even or odd)  $(b, B)$ -coboundaries.

Explicitly, an even  $(b, B)$ -cocycle is given by a sequence

$$(\phi_0, \phi_2, \phi_4, \dots),$$

where  $\phi_{2k} \in C^{2k}(\mathcal{A})$ , and

$$b\phi_{2k} + B\phi_{2k+2} = 0,$$

for all  $k \geq 0$ . Note that only finitely many  $\phi_{2k}$  are non-zero.

Similarly, an odd  $(b, B)$ -cocycle is given by a sequence

$$(\phi_1, \phi_3, \phi_5, \dots),$$

where  $\phi_{2k+1} \in C^{2k+1}(\mathcal{A})$  and

$$b\phi_{2k+1} + B\phi_{2k+3} = 0,$$

for all  $k \geq 0$ , and also  $B\phi_1 = 0$ . Again, only finitely many  $\phi_{2k+1}$  are non-zero.

The following result allows us to evaluate an even (odd)  $(b, B)$ -cocycle on a projection (unitary) in a given  $*$ -algebra  $\mathcal{A}$ .

**Proposition 6.6** *Let  $\mathcal{A}$  be a unital  $*$ -algebra.*

- If  $\phi = (\phi_1, \phi_3, \dots)$  is an odd  $(b, B)$ -cocycle for  $\mathcal{A}$ , and  $u$  is an unitary in  $\mathcal{A}$ , then the quantity

$$\langle \phi, u \rangle := \frac{1}{\sqrt{\pi}} \sum_{k=0}^{\infty} (-1)^{k+1} k! \phi_{2k+1}(u^*, u, \dots, u^*, u)$$

only depends on the class of  $\phi$  in  $HCP^{\text{odd}}(\mathcal{A})$ .

- If  $\phi = (\phi_0, \phi_2, \dots)$  is an even  $(b, B)$ -cocycle for  $\mathcal{A}$ , and  $p$  is an projection in  $\mathcal{A}$ , then the quantity

$$\langle \phi, p \rangle := \phi_0(p) + \sum_{k=1}^{\infty} (-1)^k \frac{(2k)!}{k!} \phi_{2k}(p - \frac{1}{2}, p, p, \dots, p)$$

only depends on the class of  $\phi$  in  $HCP^{\text{ev}}(\mathcal{A})$ .

**Proof** We show that  $\langle (b + B)\Theta, u \rangle = 0$  for any even cochain  $(\Theta_0, \Theta_2, \dots)$  and that  $\langle (b + B)\Theta, e \rangle = 0$  for any odd cochain  $(\Theta_1, \Theta_3, \dots)$ .

The former equation would follow from

$$(-1)^{k+1} k! b \Theta_{2k}(u^*, u, \dots, u^*, u) + (-1)^k (k-1)! B \Theta_{2k}(u^*, u, \dots, u^*, u) = 0,$$

for any  $k \geq 0$ . Using the definition of  $b$  and  $B$ , we compute that indeed:

$$\begin{aligned} & (-1)^{k+1} k! [\Theta_{2k}(1, u^*, u, \dots, u^*, u) + (-1)^{2k+1} \Theta_{2k}(1, u, u^*, \dots, u, u^*)] \\ & + (-1)^k (k-1)! [k \Theta_{2k}(1, u^*, u, \dots, u^*, u) - k \Theta_{2k}(1, u, u^*, \dots, u, u^*)] = 0. \end{aligned}$$

The second claim would follow from

$$(-1)^{k+1} \frac{(2k+2)!}{(k+1)!} b \Theta_{2k+1}(p - \frac{1}{2}, p, \dots, p) + (-1)^k \frac{(2k)!}{k!} B \Theta_{2k+1}(p - \frac{1}{2}, p, \dots, p) = 0,$$

for any  $k \geq 1$ , and indeed

$$-2b\Theta_1(p - \frac{1}{2}, p, p) + B\Theta_1(p) = 0.$$

Let us start with the latter, for which we compute

$$\begin{aligned} & -2[2\Theta_1(p - \frac{1}{2}p, p) - \Theta_1(p - \frac{1}{2}, p)] + \Theta_1(1, p) = \\ & -2\Theta_1(p, p) + 2\Theta_1(p, p) - \Theta_1(1, p) + \Theta_1(1, p) = 0. \end{aligned}$$

The same trick applies also to the first expression, for any  $k \geq 1$ :

$$\begin{aligned} & (-1)^{k+1} \frac{(2k+2)!}{(k+1)!} [2\Theta_{2k+1}(p - \frac{1}{2}p, p, \dots, p) - \Theta_{2k+1}(p - \frac{1}{2}, p, \dots, p)] \\ & + (-1)^k \frac{(2k)!}{k!} [(2k+1)\Theta_{2k+1}(1, p, \dots, p)] = 0, \end{aligned}$$

which follows directly from the identity

$$\frac{1}{2} \frac{(2k+2)!}{(k+1)!} - (2k+1) \frac{(2k)!}{k!} = 0.$$

□

**Exercise 6.7** Let  $\phi \in C^k(\mathcal{A})$  be a  $b$ -cocycle (i.e.  $b\phi = 0$ ) that also satisfies the following condition of being *cyclic*:

$$\phi(a^0, a^1, \dots, a^k) = (-1)^k \phi(a^k, a^0, a^1, \dots, a^{k-1}),$$

for all  $a^0, a^1, \dots, a^k \in \mathcal{A}$ . Show that  $(0, \dots, 0, \phi, 0, \dots)$  (with  $\phi$  at the  $k$ 'th position) is a  $(b, B)$ -cocycle.

**Exercise 6.8** In the example of the circle, show that the odd cochain  $(\phi^1, 0, \dots)$  on  $C^\infty(\mathbb{S}^1)$  with (cf. Exc. (6.2))

$$\phi^1(f^0, f^1) = \text{Tr } F[F, f^0][F, f^1]; \quad (f^0, f^1 \in C^\infty(\mathbb{S}^1)),$$

is an odd  $(b, B)$ -cocycle.

### 6.2.1 Cyclic Cocycles for the Noncommutative Torus

We will illustrate the above periodic cyclic cohomology and the evaluation on projections and unitaries with the noncommutative torus. Recall its structure from Sect. 5.3.1.

In view of Exercise 6.5(1) it is immediate that  $\tau$  defines a Hochschild cocycle:  $b\tau = 0$ . In other words,  $\tau$  defines an element in  $HH^0(\mathcal{A}_\theta)$ . In fact, it also defines an element in  $HCP^{\text{ev}}(\mathcal{A}_\theta)$ , to wit,  $(\tau, 0, \dots)$ . Let us check that it does not represent the zero class by exploiting the evaluation from Proposition 6.6 on the trivial projection  $p = 1$ . Indeed, since  $\langle \tau, 1 \rangle = \tau(1) = 1$  and since the evaluation does not depend on the representative of  $\tau$  (as is shown in Proposition 6.6), we find that  $\tau$  cannot be cohomologous to zero.

In order to construct odd cyclic cocycles we need the derivations  $\delta_1, \delta_2 : \mathcal{A}_\theta \rightarrow \mathcal{A}_\theta$  from Eq. (5.3.2). We then define two 1-cochains  $\psi_1, \psi_2 \in C^1(\mathcal{A}_\theta)$  by

$$\psi_1(a^0, a^1) = \tau(a^0 \delta_1(a^1)); \quad \psi_2(a^0, a^1) = \tau(a^0 \delta_2(a^1)).$$

Let us check that they are Hochschild cocycles:

$$\begin{aligned} b\psi_1(a^0, a^1, a^2) &= \psi_1(a^0 a^1, a^2) - \psi_1(a^0, a^1 a^2) + \psi_1(a^2 a^0, a^1) \\ &= \tau(a^0 a^1 \delta_1(a^2)) - \tau(a^0 \delta_1(a^1 a^2)) + \tau(a^2 a^0 \delta_1(a^1)) = 0, \end{aligned}$$

by using the Leibniz property (5.3.3) for  $\delta_1$ . The same argument also shows that  $b\psi_2 = 0$ .

Using Stokes theorem we may also show that

$$B\psi_1(a^0) = \tau(\delta_1(a^0)) = 0, \quad B\psi_2(a^0) = 0.$$

Hence,  $(0, \psi_1, 0, \dots)$  and  $(0, \psi_2, 0, \dots)$  are odd  $(b, B)$ -cocycles and define class in odd periodic cyclic cohomology  $HCP^{\text{odd}}(\mathcal{A}_\theta)$ . Again by exploiting the evaluation

on unitaries we may check that  $\psi_1$  and  $\psi_2$  define non-trivial and different classes in  $HCP^{\text{odd}}(\mathcal{A}_\theta)$ . Indeed,

$$\langle \psi_1, u \rangle = -\frac{1}{\sqrt{\pi}}\psi_1(u^*, u) = -\frac{i}{\sqrt{\pi}}; \quad \langle \psi_1, v \rangle = 0$$

while

$$\langle \psi_2, u \rangle = 0; \quad \langle \psi_2, v \rangle = -\frac{i}{\sqrt{\pi}}.$$

Finally, there is a two-cochain defined by

$$\phi(a^0, a^1, a^2) = \tau(a^0(\delta_1(a^1)\delta_2(a^2) - \delta_2(a^1)\delta_1(a^2)))$$

**Exercise 6.9** Show that  $b\phi = 0$  and that  $B\phi = 0$ .

We conclude that  $\phi$  defines a class in even periodic cyclic cohomology  $HCP^{\text{ev}}(\mathcal{A}_\theta)$ . Since the evaluation of  $\phi$  on the projection 1 vanishes, we already know that  $\phi$  defines a different class in  $HCP^{\text{ev}}(\mathcal{A}_\theta)$  than  $\tau$ . Moreover, one may evaluate  $\phi$  non-trivially on a projection  $p$ . This will be worked out in Exercise 6.10 below (*cf.* Note 10 in this chapter).

It turns out that the above four  $(b, B)$ -cocycles fully describe the periodic cyclic cohomology of  $\mathcal{A}_\theta$  in the sense that

$$HCP^{\text{ev}}(\mathcal{A}_\theta) = \mathbb{C}[\tau] \oplus \mathbb{C}[\phi]; \quad HCP^{\text{odd}}(\mathcal{A}_\theta) = \mathbb{C}[\psi_1] \oplus \mathbb{C}[\psi_2].$$

This should be considered as the noncommutative analogue of the cell decomposition of the torus: one 0-cell (a point), two 1-cells (two circles) and one 2-cell (the torus).


**Exercise 6.10** We consider the pairing between the  $(b, B)$ -cocycle  $\phi$  and a class of projections on the noncommutative torus. Consider the following element  $p \in \mathcal{A}_\theta$ :

$$p = \left( \sum_{n \in \mathbb{Z}} a_n v^n \right) u + \left( \sum_{n \in \mathbb{Z}} b_n v^n \right) + u^* \left( \sum_{n \in \mathbb{Z}} c_n v^n \right) \quad (6.2.1)$$

for some Schwartz sequences  $(a_n)$ ,  $(b_n)$ ,  $(c_n)$  (*cf.* Definition 5.16). We also write  $g$ ,  $f$ ,  $h$  for the corresponding functions on  $\mathbb{S}^1$ :

$$g(t) = \sum_{n \in \mathbb{Z}} a_n e^{int}; \quad f(t) = \sum_{n \in \mathbb{Z}} b_n e^{int}; \quad h(t) = \sum_{n \in \mathbb{Z}} c_n e^{int}.$$

- (1) Show that  $p$  is an orthogonal projection, *i.e.*  $p^2 = p = p^*$  if and only if  $h = g^*$  while  $f$  and  $g$  satisfy ( $\forall t \in \mathbb{S}^1$ ):



**Fig. 6.4** Functions  $f$  and  $g = h^*$  that ensure that  $p$  defined in (6.2.1) is a projection in  $\mathcal{A}_\theta$

- $g(t)g(t - 2\pi\theta) = 0$ ;
- $(f(t) + f(t - 2\pi\theta))g(t) = g(t)$ ;
- $f(t) - f(t^2) = |g(t)|^2 + |g(t + 2\pi\theta)|^2$ .

- (2) Verify that a class of examples of functions  $f, g$  that satisfy the above three conditions can be given as follows. Take any  $\varepsilon$  such that  $\varepsilon < 2\pi\theta$  and  $2\pi\theta + \varepsilon < \pi$ . On  $[0, \varepsilon]$  let  $f$  be any smooth function with values between 0 and 1 such that  $f(0) = 0$  and  $f(\varepsilon) = 1$ . On  $[2\pi\theta, 2\pi\theta + \varepsilon]$  define  $f(t) = 1 - f(t - 2\pi\theta)$ , while on  $[\varepsilon, 2\pi\theta]$  and  $[2\pi\theta + \varepsilon, 2\pi]$  let  $f$  take values 1 and 0, respectively. Finally, let  $g$  be defined on  $[2\pi\theta, 2\pi\theta + \varepsilon]$  by  $g(t) = \sqrt{f(t)(1 - f(t))}$  and equal to zero elsewhere on  $[0, 2\pi]$  (see Fig. 6.4).
- (3) Returning to the general case of functions  $f, g$  satisfying the conditions in (1), show that  $\tau(p) = \frac{1}{2\pi} \int_{S^1} f(t) dt$ .
- (4) Compute  $\delta_1 p \delta_2 p, \delta_2 p \delta_1 p$  and show that

$$\tau(p \delta_1 p \delta_2 p - p \delta_2 p \delta_1 p) = \frac{3i}{2\pi} \int_{S^1} |g(t)|^2 (f'(t - 2\pi\theta) - f'(t)) dt$$

- (5) Show that for the explicit choice of  $f, g$  from (2) this reduces to

$$\tau(p \delta_1 p \delta_2 p - p \delta_2 p \delta_1 p) = \frac{3i}{2\pi} \int_{2\pi\theta}^{2\pi\theta+\varepsilon} f(t)(1 - f(t))(-2f'(t)) dt = \frac{i}{2\pi}.$$

HINT: Here one may want to take inspiration from the proof of Proposition 6.1.

- (6) Deduce from this that the evaluation  $\langle \phi, p \rangle = -i/\pi$  for these projections.

## 6.3 Abstract Differential Calculus

We return to the general case. Starting with a spectral triple, we now introduce a differential calculus. In the case of the canonical triple of a spin manifold  $M$ , this will agree with the usual differential calculus on  $M$ .

Let  $(\mathcal{A}, \mathcal{H}, D)$  be a spectral triple; we assume that  $D$  is invertible. We introduce *Sobolev spaces*  $\mathcal{H}^s$  as follows:

$$\mathcal{H}^s := \text{Dom } |D|^s; \quad (s \in \mathbb{R}).$$

These spaces are naturally normed by

$$\|\xi\|_s^2 = \| |D|^s \xi \|^2,$$

and are complete in this norm. Moreover, for  $s > t$  the inclusion  $\mathcal{H}^s \rightarrow \mathcal{H}^t$  is continuous.

**Exercise 6.11** Prove this last statement.

Obviously  $\mathcal{H}^0 = \mathcal{H}$ , while at the other extreme we have the intersection

$$\mathcal{H}^\infty := \bigcap_{s \geq 0} \mathcal{H}^s.$$

**Definition 6.7** For each  $r \in \mathbb{R}$  we define operators of *analytic order*  $\leq r$  to be operators in  $\mathcal{H}^\infty$  that extend to bounded operators from  $\mathcal{H}^s$  to  $\mathcal{H}^{s-r}$  for all  $s \in \mathbb{R}$ . We denote the space of such operators by  $\text{op}^r$ .

In order to find interesting differential operators coming from our spectral triple, we introduce some smoothness conditions. The first is that the spectral triple is **finitely summable**, *i.e.* there exists  $p$  so that  $|D|^{-p}$  is a trace class operator.

**Definition 6.8** A spectral triple  $(\mathcal{A}, \mathcal{H}, D)$  is called *regular* if  $\mathcal{A}$  and  $[D, \mathcal{A}] = \{[D, a] : a \in \mathcal{A}\}$  belong to the smooth domain of  $\delta(\cdot) = [|D|, \cdot]$ . That is, for each  $k \geq 0$  the operators  $\delta^k(a)$  and  $\delta^k([D, a])$  are bounded.

We will denote by  $\mathcal{B}$  the algebra generated by  $\delta^k(a)$ ,  $\delta^k([D, a])$  for all  $a \in \mathcal{A}$  and  $k \geq 0$ .

**Definition 6.9** Let  $(\mathcal{A}, \mathcal{H}, D)$  be a finitely-summable regular spectral triple. The *dimension spectrum*  $\text{Sd}$  is the subset of  $\{z \in \mathbb{C} : \Re(z) \geq 0\}$  of singularities of the analytic functions

$$\zeta_b(z) = \text{Tr } b |D|^{-z}; \quad (b \in \mathcal{B}).$$

We say the dimension spectrum is *simple* when the functions  $\zeta_b$  have at most simple poles.

In our treatment we restrict to finitely-summable, regular spectral triples with simple dimension spectrum and for which there is a finite number of poles in  $\text{Sd}$ .

**Lemma 6.10** *The algebra  $\mathcal{B}$  maps  $\mathcal{H}^\infty$  to itself.*

**Proof** This follows by induction from the identity

$$\begin{aligned}\|T\xi\|_s^2 &= \|T\xi\|^2 + \||D|^s T\xi\|^2 \\ &= \|T\|^2 \|\xi\|^2 + (\||D|^{s-1} \delta(T)\xi\| + \||D|^{s-1} T|D|\xi\|)^2,\end{aligned}$$

for any operator  $T$  in the smooth domain of  $\delta$  and any  $s \geq 0$ .  $\square$

We will regard the elements in  $\mathcal{B}$  as pseudodifferential operators of order 0, according to the following definition.

**Definition 6.11** A *pseudodifferential operator of order  $k \in \mathbb{Z}$*  associated to a regular spectral triple  $(\mathcal{A}, \mathcal{H}, D)$  is given by a finite sum:

$$b_k |D|^k + b_{k-1} |D|^{k-1} + \dots,$$

where  $b_k, b_{k-1}, \dots \in \mathcal{B}$ . We denote the space of pseudodifferential operators of order  $k$  by  $\Psi^k(\mathcal{A}, \mathcal{H}, D)$ , or simply  $\Psi^k(\mathcal{A})$ .

**Lemma 6.12** The subspaces  $\Psi^k(\mathcal{A})$  ( $k \in \mathbb{Z}$ ) furnish a  $\mathbb{Z}$ -filtration on the algebra  $\Psi(\mathcal{A})$  of pseudodifferential operators.

**Proof** This follows directly from the expression:

$$b_1 |D|^{k_1} \cdot b_2 |D|^{k_2} = \sum_{j=0}^{k_1} \binom{k_1}{j} b_1 \delta^j(b_2) |D|^{k_1+k_2-j}. \quad \square$$

On this algebra, the map  $\delta(\cdot) = [|D|, \cdot]$  acts as a derivation, preserving the filtration. For any operator  $T$  in  $\mathcal{H}$  we also define the following (iterated) derivation,

$$\nabla(T) = [D^2, T]; \quad T^{(k)} := \nabla^k(T).$$

**Exercise 6.12** Prove that for any  $P \in \Psi(\mathcal{A})$  we have

$$\nabla(P) = 2\delta(P)|D| + \delta^2(P).$$

Conclude that  $\nabla : \Psi^k(\mathcal{A}) \rightarrow \Psi^{k+1}(\mathcal{A})$ .

**Proposition 6.13** Let  $P \in \Psi^k(\mathcal{A})$ . Then  $P : \mathcal{H}^{s+k} \rightarrow \mathcal{H}^s$  is a continuous map. Hence, such a  $P$  has analytic order  $\leq k$  and we have  $\Psi^k(\mathcal{A}) \subset \text{op}^k$ .

Using this abstract pseudodifferential calculus, we now introduce the functionals of relevance for the index formula.

**Definition 6.14** Let  $(\mathcal{A}, \mathcal{H}, D; \gamma)$  be a regular spectral triple. For pseudodifferential operators  $X^0, X^1, \dots, X^p \in \Psi(\mathcal{A})$  and  $\Re(z) \gg 0$  define

$$\langle X^0, X^1, \dots, X^p \rangle_z = (-1)^p \frac{\Gamma(z)}{2\pi i} \text{Tr} \left( \int \lambda^{-z} \gamma X^0 (\lambda - D^2)^{-1} X^1 (\lambda - D^2)^{-1} \cdots X^p (\lambda - D^2)^{-1} d\lambda \right).$$

Let us show that this expression is well defined, *i.e.* that the integral is actually trace class. We first practice with this expression in a special case.

**Exercise 6.13** Assume that  $X^j \in \Psi^{k_j}(\mathcal{A})$  commutes with  $D$  for all  $j = 0, \dots, p$ .

(1) Use Cauchy's integral formula to show that

$$\langle X^0, X^1, \dots, X^p \rangle_z = \frac{\Gamma(z + p)}{p!} \text{Tr}(\gamma X^0 \cdots X^p |D|^{-2z-2p}).$$

(2) Show that this expression extends to a meromorphic function on  $\mathbb{C}$ .

This exercise suggests that, in the general case, we move all terms  $(\lambda - D^2)^{-1}$  in  $\langle X^0, X^1, \dots, X^p \rangle_z$  to the right. This we will do in the remainder of this section. First, we need the following result.

**Lemma 6.15** Let  $X \in \Psi^q(\mathcal{A})$  and let  $n > 0$ . Then for any positive integer  $k$ , we have

$$\begin{aligned} (\lambda - D^2)^{-n} X &= X(\lambda - D^2)^{-n} + nX^{(1)}(\lambda - D^2)^{-(n+1)} \\ &\quad + \frac{n(n+1)}{2} X^{(2)}(\lambda - D^2)^{-(n+2)} + \dots \\ &\quad + \frac{n(n+1) \cdots (n+k)}{k!} X^{(k)}(\lambda - D^2)^{-(n+k)} + R_k, \end{aligned}$$

where the remainder  $R_k$  is of analytic order  $q - 2n - k - 1$  or less.

**Proof** This follows by repeatedly applying the formula

$$\begin{aligned} (\lambda - D^2)^{-1} X &= X(\lambda - D^2)^{-1} + [(\lambda - D^2)^{-1}, X] \\ &= X(\lambda - D^2)^{-1} + (\lambda - D^2)^{-1}[D^2, X](\lambda - D^2)^{-1}. \end{aligned}$$

This yields an asymptotic expansion

$$(\lambda - D^2)^{-1} X \sim \sum_{i \geq 0} X^{(i)}(\lambda - D^2)^{-1-i},$$

so that for each  $m \ll 0$  every sufficiently large finite partial sum agrees with the left-hand side up to an operator of analytic order  $m$  or less. Indeed, truncating the above sum at  $i = k$ , we find that the remainder is

$$(\lambda - D^2)^{-1} X^{(k+1)} (\lambda - D^2)^{-1-k},$$

which is of analytic order  $-2 + (q + k + 1) - 2(k + 1) = q - k - 3$  or less.

More generally for any positive integer  $n$  one has:

$$(\lambda - D^2)^{-n} X \sim \sum_{k \geq 0} (-1)^k \binom{-n}{k} X^{(k)} (\lambda - D^2)^{-n-k}.$$

Estimates similar to those above show that the remainder has the claimed analytic order.  $\square$

We now arrive at the final result of this section which will form the main ingredient in the next section, where we will introduce the  $(b, B)$ -cocycles relevant for the index formula.

**Proposition 6.16** *The expression  $\langle X^0, \dots, X^p \rangle_z$  in Definition 6.14 seen as a function of  $z$  extends meromorphically to  $\mathbb{C}$ .*

**Proof** We use Lemma 6.15 to bring all  $(\lambda - D^2)^{-1}$  to the right. We first introduce the combinatorial quantities:

$$c(k_1, \dots, k_j) = \frac{(k_1 + \dots + k_j + j)!}{k_1! \dots k_j! (k_1 + 1) \dots (k_1 + \dots + k_j + j)},$$

for non-negative integers  $k_1, \dots, k_j$ . These satisfy

$$c(k_1, \dots, k_j) = c(k_1, \dots, k_{j-1}) \frac{(k_1 + \dots + k_{j-1} + j) \dots (k_1 + \dots + k_j + j - 1)}{k_j!},$$

while  $c(k_1) = 1$  for all  $k_1$ .

From Lemma 6.15 we know that there is the following asymptotic expansion:

$$(\lambda - D^2)^{-1} X^1 \sim \sum_{k_1 \geq 0} c(k_1) X^{1^{(k_1)}} (\lambda - D^2)^{-k_1}.$$

Then, in the subsequent step we find

$$\begin{aligned} (\lambda - D^2)^{-1} X^1 (\lambda - D^2)^{-1} X^2 &\sim \sum_{k_1 \geq 0} c(k_1) X^{1^{(k_1)}} (\lambda - D^2)^{-(k_1+2)} X^2 \\ &\sim \sum_{k_1, k_2 \geq 0} c(k_1, k_2) X^{1^{(k_1)}} X^{2^{(k_2)}} (\lambda - D^2)^{-(k_1+k_2+2)}, \end{aligned}$$

and finally

$$(\lambda - D^2)^{-1} X^1 \cdots (\lambda - D^2)^{-1} X^p \sim \sum_{k \geq 0} c(k) X^{1(k_1)} \cdots X^{p(k_p)} (\lambda - D^2)^{-(|k|+p)},$$

where  $k = (k_1, \dots, k_p)$  is a multi-index and  $|k| = k_1 + \dots + k_p$ .

Multiplying this with  $\gamma X_0$  and integrating as in Definition 6.14, this yields

$$\begin{aligned} & (-1)^p \frac{\Gamma(z)}{2\pi i} \int \lambda^{-z} \gamma X^0 (\lambda - D^2)^{-1} X^1 \cdots (\lambda - D^2)^{-1} X^p (\lambda - D^2)^{-1} d\lambda \\ & \sim \sum_{k \geq 0} c(k) \gamma X^0 X^{1(k_1)} \cdots X^{p(k_p)} (-1)^p \frac{\Gamma(z)}{2\pi i} \int \lambda^{-z} (\lambda - D^2)^{-(|k|+p+1)} d\lambda \\ & = \sum_{k \geq 0} c(k) \gamma X^0 X^{1(k_1)} \cdots X^{p(k_p)} (-1)^p \Gamma(z) \binom{-z}{|k|+p} |D|^{-2(z+|k|+p)}, \end{aligned}$$

where we have used the integral formula, valid for real  $\lambda_0$ :

$$\frac{1}{2\pi i} \int \frac{\lambda^{-z}}{(\lambda - \lambda_0)^{N+1}} d\lambda = \binom{-z}{N} \lambda_0^{-(N+z)}. \quad (6.3.1)$$

Finally, using the functional equation for the gamma function,

$$(-1)^p \Gamma(z) \binom{-z}{|k|+p} = (-1)^{|k|} \frac{\Gamma(z+p+|k|)}{(|k|+p)!},$$

we obtain an asymptotic expansion

$$\begin{aligned} \langle X_0, \dots, X_p \rangle_z & \sim \sum_{k \geq 0} (-1)^{|k|} \frac{\Gamma(z+p+|k|)}{(|k|+p)!} c(k) \\ & \times \text{Tr} \left( \gamma X^0 X^{1(k_1)} \cdots X^{p(k_p)} |D|^{-2(z+|k|+p)} \right). \end{aligned} \quad (6.3.2)$$

As  $|k|$  becomes large the remainder in the truncated expansion on the right-hand side becomes trace class.  $\square$

**Exercise 6.14** Use Cauchy's integral formula to prove Eq. (6.3.1).

## 6.4 Residues and the Local $(b, B)$ -Cocycle

In this section we derive even and odd  $(b, B)$ -cocycles on a given algebra  $\mathcal{A}$  from the functionals  $\langle X^0, X^1, \dots, X^p \rangle_z$  defined in the previous section. First, we derive some useful relations between them. We denote the  $\mathbb{Z}_2$ -grading of an operator  $X$

by  $(-1)^X$ , according to the grading  $\gamma$  on  $\mathcal{H}$ . Moreover, for such an operator  $X$  we denote the graded commutator by  $[D, X] = DX - (-1)^X XD$ . Note that with these conventions we have

$$[D, [D, T]] = [D^2, T] \equiv \nabla(T),$$

for any even operator  $T$ .

**Lemma 6.17** *The meromorphic functions  $\langle X^0, \dots, X^p \rangle_z$  satisfy the following functional equations:*

$$\langle X^0, \dots, X^p \rangle_z = (-1)^{X^p} \langle X^p, X^0, \dots, X^{p-1} \rangle_z; \quad (\text{a})$$

$$\langle X^0, \dots, X^p \rangle_{z+1} = \sum_{j=0}^p \langle X^0, \dots, X^{j-1}, 1, X^j, \dots, X^p \rangle_z; \quad (\text{b})$$

$$\begin{aligned} \langle X^0, \dots, [D^2, X^j], \dots, X^p \rangle_z &= \langle X^0, \dots, X^{j-1} X^j, \dots, X^p \rangle_z \\ &\quad - \langle X^0, \dots, X^j X^{j+1}, \dots, X^p \rangle_z; \end{aligned} \quad (\text{c})$$

$$\sum_{j=0}^p (-1)^{X^0 \dots X^{j-1}} \langle X^0, \dots, [D, X^j], \dots, X^p \rangle_z = 0. \quad (\text{d})$$

**Proof** (a) follows directly from the property of the trace in  $\langle X^0, \dots, X^p \rangle_z$ , taking into account the commutation of  $X^p$  with the grading  $\gamma$ . For (b), note that the integral of the following expression vanishes:

$$\begin{aligned} \frac{d}{d\lambda} &(\lambda^{-z} X^0 (\lambda - D^2)^{-1} \dots X^p (\lambda - D^2)^{-1}) \\ &= -z \lambda^{-z-1} X^0 (\lambda - D^2)^{-1} \dots X^p (\lambda - D^2)^{-1} \\ &\quad - \sum_{j=0}^p \lambda^{-z} X^0 (\lambda - D^2)^{-1} \dots (\lambda - D^2)^{-1} X^j (\lambda - D^2)^{-1} \dots X^p (\lambda - D^2)^{-1}. \end{aligned}$$

Equation (c) follows from

$$(\lambda - D^2)^{-1} [D^2, X^j] (\lambda - D^2)^{-1} = -(\lambda - D^2)^{-1} X^j + X^j (\lambda - D^2)^{-1}.$$

Finally, (d) is equivalent to

$$\text{Tr } \gamma \left[ D, \int \lambda^{-z} X^0 (\lambda - D^2)^{-1} \dots X^p (\lambda - D^2)^{-1} d\lambda \right] = 0,$$

which is the supertrace of a (graded) commutator.  $\square$

**Definition 6.18** For any  $p \geq 0$ , define a  $(p+1)$ -linear functional on  $\mathcal{A}$  with values in the meromorphic functions on  $\mathbb{C}$  by

$$\Psi_p(a^0, \dots, a^p) = \langle a^0, [D, a^1], \dots, [D, a^p] \rangle_{s-\frac{p}{2}}.$$

**Proposition 6.19** *The even  $(b, B)$ -cochain  $\Psi = (\Psi_0, \Psi_2, \dots)$  is an (improper) even  $(b, B)$ -cocycle in the sense that*

$$b\Psi_{2k} + B\Psi_{2k+2} = 0.$$

*Similarly, the odd  $(b, B)$ -cochain  $\Psi = (\Psi_1, \Psi_3, \dots)$  is an (improper) odd  $(b, B)$ -cocycle.*

**Proof** It follows from the definition of  $B$  and a subsequent application of (a) and (b) of Lemma 6.17 that

$$\begin{aligned} B\Psi_{2k+2}(a^0, \dots, a^{2k+1}) &= \sum_{j=0}^{2k+1} (-1)^j \langle 1, [D, a^j], \dots, [D, a^{j-1}] \rangle_{s-(k+1)} \\ &= \sum_{j=0}^{2k+1} \langle [D, a^0], \dots, [D, a^{j-1}], 1, [D, a^j], \dots, [D, a^{2k+1}] \rangle_{s-(k+1)} \\ &= \langle [D, a^0], \dots, [D, a^{2k+1}] \rangle_{s-k}. \end{aligned}$$

Also, from the definition of  $b$  and the Leibniz rule

$$[D, a^j a^{j+1}] = a^j [D, a^{j+1}] + [D, a^j] a^{j+1}$$

it follows that

$$\begin{aligned} b\Psi_{2k}(a^0, \dots, a^{2k+1}) &= \langle a^0 a^1, [D, a^2], \dots, [D, a^{2k+1}] \rangle_{s-k} \\ &\quad - \langle a^0, a^1 [D, a^2], \dots, [D, a^{2k+1}] \rangle_{s-k} \\ &\quad - \langle a^0, [D, a^1] a^2, \dots, [D, a^{2k+1}] \rangle_{s-k} \\ &\quad + \langle a^0, [D, a^1], a^2 [D, a^3], \dots, [D, a^{2k+1}] \rangle_{s-k} \\ &\quad + \langle a^0, [D, a^1], [D, a^2] a^3, \dots, [D, a^{2k+1}] \rangle_{s-k} \\ &\quad - \dots \\ &\quad - \langle a^{2k+1} a^0, [D, a^1], \dots, [D, a^{2k}] \rangle_{s-k}, \end{aligned}$$

which, by Lemma 6.17(c), becomes

$$\sum_{j=0}^{2k+1} (-1)^{j-1} \langle a^0, [D, a^1], \dots, [D^2, a^j], \dots, [D, a^{2k+1}] \rangle_{s-k}.$$

Combining these expressions for  $B\Psi_{2k+2}$  and  $b\Psi_{2k}$  and writing  $X^0 = a^0$ , and  $X^j = [D, a^j]$  for  $j \geq 1$ , we obtain

$$\begin{aligned} B\Psi_{2k+2}(a^0, \dots, a^{2k+1}) + b\Psi_{2k}(a^0, \dots, a^{2k+1}) \\ = \sum_{j=0}^{2k+1} (-1)^{X^0 \dots X^j} \langle X^0, \dots, [D, X^j], \dots, X^{2k+1} \rangle_{s-k}, \end{aligned}$$

which vanishes because of Lemma 6.17(d).

In the odd case, a similar argument shows that  $b\Psi_{2k-1} + B\Psi_{2k+1} = 0$ .  $\square$

The above cocycles have been termed *improper* because all  $\Psi_p$  might be non-zero, on top of which (rather than in  $\mathbb{C}$ ) they take values in the field of meromorphic functions on  $\mathbb{C}$ . By taking residues of the meromorphic functions  $\Psi_p$  we obtain a *proper* even or odd  $(b, B)$ -cocycle. This is the residue cocycle that was introduced by Connes and Moscovici.

**Theorem 6.20** *For any  $p \geq 0$  and all  $a^0, \dots, a^p \in \mathcal{A}$  the following formulas define an even or odd  $(b, B)$ -cocycle:*

$$\text{res}_{s=0} \Psi_0(a^0) = \text{Tr } \gamma a^0 |D|^{-2s}|_{s=0},$$

and

$$\begin{aligned} \text{res}_{s=0} \Psi_p(a^0, \dots, a^p) \\ = \sum_{k \geq 0} c_{p,k} \text{res}_{s=0} \text{Tr} \left( \gamma a^0 [D, a^1]^{(k_1)} \cdots [D, a^p]^{(k_p)} |D|^{-p-2|k|-2s} \right), \end{aligned}$$

for  $p \geq 1$ , where the constants  $c_{p,k}$  are given in terms of the (non-negative) multi-indices  $(k_1, \dots, k_p)$  by

$$c_{p,k} := \frac{(-1)^{|k|}}{k!} \frac{\Gamma(|k| + \frac{p}{2})}{(k_1 + 1)(k_1 + k_2 + 2) \cdots (k_1 + \cdots k_p + p)}.$$

**Proof** We use the asymptotic expansion (a). Indeed, setting  $z = s - \frac{p}{2}$  in that expression and taking residues at  $s = 0$  gives the desired expansion, with the coefficients  $c_{p,k}$  appearing because

$$c_{p,k} \equiv (-1)^{|k|} \Gamma\left(|k| + \frac{p}{2}\right) \frac{c(k)}{(p + |k|)!}.$$

$\square$

## 6.5 The Local Index Formula

Let  $(\mathcal{A}, \mathcal{H}, D)$  be a regular spectral triple, as above. The local index formula expresses the index of twisted Dirac operators in terms of cocycles in the  $(b, B)$  bicomplex, which are easier to compute. We are interested in the indices of the following two Fredholm operators.

Suppose that  $(\mathcal{A}, \mathcal{H}, D)$  is even. If  $p \in \mathcal{A}$  is a projection, then  $D_p = pDp$  is a Fredholm operator on the Hilbert space  $\mathcal{H}$ . This follows from the fact that  $D_p$  is essentially a finite-dimensional extension of the Fredholm operator  $D$ . We are interested in the index of this so-called twisted Dirac operator  $D_p$ .

In case that  $(\mathcal{A}, \mathcal{H}, D)$  is an odd spectral triple, we take a unitary  $u \in \mathcal{A}$  and define  $D_u = PuP$ , where  $P = \frac{1}{2}(1 + \text{Sign } D)$ . Again,  $D_u$  is a Fredholm operator on  $\mathcal{H}$  and we are interested in the index of  $D_u$ .

**Theorem 6.21** *Let  $(\mathcal{A}, \mathcal{H}, D)$  be a regular spectral triple with simple and finite dimension spectrum  $\text{Sd}$  and let  $\text{res}_{s=0}\Psi$  be the (even or odd)  $(b, B)$ -cocycle derived previously.*

- *If  $(\mathcal{A}, \mathcal{H}, D)$  is even and  $p$  is a projection in  $\mathcal{A}$ , then*

$$\text{index } D_p = \langle \text{res}_{s=0}\Psi, p \rangle.$$

- *If  $(\mathcal{A}, \mathcal{H}, D)$  is odd and  $u$  is a unitary in  $\mathcal{A}$ , then*

$$\text{index } D_u = \langle \text{res}_{s=0}\Psi, u \rangle.$$

**Remark 6.22** Sometimes a projection or a unitary is given in  $M_N(\mathcal{A})$  instead of  $\mathcal{A}$ . The above result can be extended easily to this case, namely by constructing a spectral triple on  $M_N(\mathcal{A})$  and doing the index computation there. Indeed, it would follow from Theorem 7.15 that if  $(\mathcal{A}, \mathcal{H}, D)$  is a spectral triple, then so is  $(M_N(\mathcal{A}), \mathcal{H} \otimes \mathbb{C}^N, D \otimes \mathbb{I}_N)$ .

**Proof of Theorem 6.21.** We will prove the even case in two steps (for the odd case see Note 15 in this chapter),

- (1) the Atiyah–Bott formula for the index:

$$\text{index } D_p = \text{res}_{s=0}\Gamma(s) \text{Tr } \gamma|D_p|^{-2s}.$$

- (2) Change the representative of the class  $\text{res}_{s=0}\Psi$  in  $HCP^{\text{ev}}(\mathcal{A})$  to reduce to the case that  $D$  commutes with  $p$ , so that

$$\langle \text{res}_{s=0}\Psi, p \rangle = \text{res}_{s=0}\Gamma(s) \text{Tr } \gamma p|D|^{-2s}.$$

For (1) let us first prove another well-known formula.

**Lemma 6.23** (McKean–Singer formula) *Let  $(\mathcal{A}, \mathcal{H}, D)$  be an even spectral triple. Then*

$$\text{index } D = \text{Tr } \gamma e^{-tD^2}.$$

**Proof** Since  $D$  is odd with respect to  $\gamma$ , its spectrum lies symmetrically around 0 in  $\mathbb{R}$ , including multiplicities. If we denote the  $\lambda$ -eigenspace in  $\mathcal{H}$  by  $\mathcal{H}_\lambda$  we therefore have  $\dim \mathcal{H}_\lambda = \dim \mathcal{H}_{-\lambda}$  for any non-zero eigenvalue  $\lambda$ . Including also the kernel of  $D$ , we have

$$\text{Tr } \gamma e^{-tD^2} = \sum_{\lambda > 0} (\dim \mathcal{H}_\lambda - \dim \mathcal{H}_{-\lambda}) e^{-t\lambda^2} + \text{Tr}_{\mathcal{H}_0} \gamma = \text{Tr}_{\ker D} \gamma,$$

which is nothing but the index of  $D$ .  $\square$

Note that the McKean–Singer formula tells us in particular that  $\text{Tr } \gamma e^{-tD^2}$  does not depend on  $t$ . Using the integral formula of the gamma function, we can write:

$$\text{Tr } \gamma |D|^{-2s} = \frac{1}{\Gamma(s)} \int_0^\infty \text{Tr } \gamma e^{-tD^2} t^{s-1} dt. \quad (6.5.1)$$

We analyze the behaviour of the right-hand side as  $s \rightarrow 0$ . For this, we use

$$\frac{1}{\Gamma(s)} \sim s, \quad s \rightarrow 0.$$

Thus, only the pole part of the above integral contributes to the zeta function evaluated at  $s = 0$ . This is given by

$$\int_0^1 \text{Tr } \gamma e^{-tD^2} t^{s-1} dt = \frac{1}{s} \text{index } D,$$

where we have used the McKean–Singer formula. The remaining integral from 1 to  $\infty$  gives an entire function of  $s$ , because by finite summability the eigenvalues of  $D$  grow as  $j^{1/p}$  for some  $p > 0$ . In other words,

$$\text{index } D = \text{Tr } \gamma |D|^{-2s}|_{s=0},$$

which proves (1).

Let us then continue with (2). Consider the family of operators

$$D_t = D + t[p, [D, p]]; \quad (t \in [0, 1]).$$

We have  $D_0 = D$  and  $D_1 = pDp + (1-p)D(1-p)$  so that  $[D_1, p] = 0$ . Moreover,  $\text{index } D_t$  depends continuously on  $t$ , and (being an integer) it is therefore constant in  $t$ .

Next, we consider a family of improper cocycles  $\Psi^t$  which are defined by replacing  $D$  by  $D_t$  in Definition 6.18.

**Lemma 6.24** *The derivative of  $\Psi^t$  is an (improper) even cyclic coboundary, i.e. there exists a cochain  $\Theta^t$  such that*

$$\frac{d}{dt} \Psi_p^t + B\Theta_{p+1}^t + b\Theta_{p-1}^t = 0,$$

which is explicitly given by

$$\Theta_p^t(a^0, \dots, a^p) = \sum_{j=0}^p (-1)^{j-1} \langle a^0, \dots [D, a^j], \dot{D}, [D, a^{j+1}], \dots [D, a^p] \rangle_{s-\frac{p+1}{2}},$$

with  $\dot{D} = \frac{d}{dt} D_t \equiv [p, [D, p]]$ .

**Proof** Imitating the proof of Proposition 6.19 one can show the following identity (see also Note 17 in this chapter).

$$\begin{aligned} & B\Theta_{2k+1}^t(a^0, \dots, a^{2k}) + b\Theta_{2k-1}^t(a^0, \dots, a^{2k}) \\ &= - \sum_{j=0}^{2k} \langle a^0, [D, a^1], \dots [D, a^j], [D, \dot{D}], \dots, [D, a^{2k}] \rangle_{s-k} \\ &\quad - \sum_{j=1}^{2k} \langle a^0, [D, a^1], \dots [\dot{D}, a^j], \dots, [D, a^{2k}] \rangle_{s-k}. \end{aligned}$$

The fact that  $\frac{d}{dt} \Psi^t$  coincides with the right-hand side follows from

$$\frac{d}{dt} (\lambda - D_t^2)^{-1} = (\lambda - D_t^2)^{-1} (D\dot{D} + \dot{D}D) (\lambda - D_t^2)^{-1}. \quad \square$$

Continuing the proof of the theorem, we integrate the resulting coboundary to obtain

$$B \int_0^1 \Theta_{2k+1}^t dt + b \int_0^1 \Theta_{2k-1}^t dt = \Psi_{2k}^0 - \Psi_{2k}^1.$$

In other words,  $\text{res}_{s=0} \Psi^0$  and  $\text{res}_{s=0} \Psi^1$  define the same class in even cyclic cohomology  $HCP^{\text{ev}}(\mathcal{A})$ . So, with the help of Proposition 6.6, we can compute  $\langle \text{res}_{s=0} \Psi, p \rangle$  using  $\Psi^1$  instead of  $\Psi^0 \equiv \Psi$ , with the advantage that  $D_1$  commutes with  $p$ . Indeed, this implies that

$$\Psi_{2k}^1(p - \frac{1}{2}, p, \dots, p) = 0,$$

for all  $k \geq 1$ , so that

$$\begin{aligned}
\langle \text{res}_{s=0} \Psi^1, p \rangle &\equiv \text{res}_{s=0} \Psi_0^1(p) + \sum_{k \geq 1} (-1)^k \frac{(2k)!}{k!} \text{res}_{s=0} \Psi_{2k}^1(p - \tfrac{1}{2}, p, \dots, p) \\
&= \text{res}_{s=0} \Psi_0^1(p) \\
&= \text{res}_{s=0} \Gamma(s) \text{Tr } \gamma p |D_1|^{-2s}.
\end{aligned}$$

This completes the proof of Theorem 6.21, as by the Atiyah–Bott formula the latter expression is the index of  $D_p$ .  $\square$

## 6.6 The Local Index Formula for Toric Noncommutative Manifolds

We here illustrate the local index formula for the class of toric noncommutative manifolds  $M_\theta$  that were described in Sect. 5.3.2. It turns out that the index formula simplifies drastically in this case.

**Theorem 6.25** *For a projection  $p \in M_N(C^\infty(M_\theta))$ , we have*

$$\text{index } D_p = \text{Tr } \gamma p |D|^{-2s} \Big|_{s=0} + \sum_{k \geq 1} c_k \text{res}_{s=0} \text{Tr} \left( \gamma \left( p - \frac{1}{2} \right) [D, p]^{2k} |D|^{-2(k+s)} \right)$$

where  $c_k = (k-1)!/(2k)!$ .

**Proof** First of all, note that the twist  $L_\theta$  commutes with the action  $\alpha_s$  of  $\widetilde{\mathbb{T}}^n$  on an operator  $T$ . Indeed, if  $T$  is homogeneous of degree  $r$ , then  $L_\theta(T)$  is of degree  $r$ ,

$$\alpha_s(L_\theta(T)) = \mathcal{U}(s) T \mathcal{U}(r') \mathcal{U}(s)^{-1} = \mathcal{U}(s) T \mathcal{U}(s)^{-1} \mathcal{U}(r') = e^{2\pi i s_\mu r_\mu} L_\theta(T).$$

with  $r'_\nu = r_\mu \theta_{\mu\nu}$  so that  $r' \in \widetilde{\mathbb{T}}^n$ .

We write the cocycles  $\text{res}_{s=0} \Psi_{2k}$  appearing in the local index formula in terms of the twist  $L_\theta$  as

$$\begin{aligned}
\text{res}_{s=0} \Psi_{2k} (L_\theta(f^0), L_\theta(f^1), \dots, L_\theta(f^{2k})) &= \\
\text{res}_{s=0} \text{Tr} (\gamma L_\theta(f^0) \times_\theta [D_M, f^1]^{(\alpha_1)} \times_\theta \cdots \times_\theta [D_M, f^{2k}]^{(\alpha_{2k})} |D_M|^{-2(|\alpha|+k+s)}) &,
\end{aligned} \tag{6.6.1}$$

where we extended the  $\times_\theta$ -product to  $C^\infty(M_\theta) \bigcup [D_M, C^\infty(M_\theta)]$  which can be done unambiguously since  $D_M$  is of degree 0. Suppose now that  $f^0, \dots, f^{2k} \in C^\infty(M)$  are homogeneous of degree  $r^0, \dots, r^{2k}$ , respectively, under the action of  $\mathbb{T}^n$ , so that the operator  $f^0 \times_\theta [D_M, f^1] \times_\theta \cdots \times_\theta [D_M, f^{2k}]$  is a homogeneous element of degree  $r$  (a simple expression in terms of the  $r^i$ ). By working out the  $\times_\theta$ -product one finds a multiple of  $f^0 [D_M, f^1] \cdots [D_M, f^{2k}]$ , with a factor which is a power of the deformation parameter  $\lambda$ . Forgetting about this factor we obtain from (5.3.9) that

$$L_\theta(f^0[D_M, f^1] \cdots [D_M, f^{2k}]) = f^0[D_M, f^1] \cdots [D_M, f^{2k}]U(r_\mu\theta_{\mu 1}, \dots, r_\mu\theta_{\mu n}).$$

After applying a Mellin transform (6.5.1) one finds that each term in the local index formula for  $(C^\infty(M_\theta), \mathcal{H}, D_M)$  then takes the form

$$\begin{aligned} & \text{res}_{s=0} \text{Tr} (\gamma f^0[D_M, f^1]^{(\alpha_1)} \cdots [D_M, f^{2k}]^{(\alpha_{2k})} |D_M|^{-2(|\alpha|+k+s)} U(s)) \\ &= \Gamma(|\alpha| + k) \lim_{t \rightarrow 0} t^{|\alpha|+k} \text{Tr} (\gamma f^0[D_M, f^1]^{(\alpha_1)} \cdots [D_M, f^{2k}]^{(\alpha_{2k})} e^{-t D_M^2} U(s)), \end{aligned} \quad (6.6.2)$$

for every  $s \in \mathbb{T}^n$ . It turns out that this limit vanishes when  $|\alpha| \neq 0$  (see Note 20 below) and this completes the proof.  $\square$

## Notes

1. The local index formula was obtained by Connes and Moscovici in [1]. In our proof of the local index formula, we closely follow Higson [2]. More general proofs have been obtained in [3–5], see Note 14 in this chapter.

### Section 6.1. Local Index Formula on the Circle and on the Torus

2. The Theorem of Atkinson that appears in Exercise 6.1 can be found in [6, Proposition 3.3.11].
3. The index formula on the circle of Exercise 6.2 is a special case of [7, Theorem 5].
4. In Sect. 6.1.2 we follow [8], where a class of projections on the torus was constructed, much inspired by the so-called Powers–Rieffel projections on the noncommutative torus [9].
5. The zeta function  $\zeta_E$  that appears in (6.1.2) is a special case of an Epstein zeta function, introduced and analyzed in [10]. It turns out that  $\zeta_E$  has a pole at  $s = 1$  with residue  $\pi$ . That (6.1.2) holds also follows from the general result [1, Theorem I.2].

### Section 6.2. Hochschild and cyclic cohomology

6. In [7] Connes introduced cyclic cohomology as a noncommutative generalization of De Rham homology, and showed that for the algebra  $C^\infty(M)$  cyclic cohomology indeed reduces to De Rham homology. Besides the original article there are many texts in which this is worked out in full detail (e.g. [11–14]).
7. Example 6.4 is a special case of the fact that  $HH^k(C^\infty(M)) \simeq \Omega_k(M)$ , the space of De Rham  $k$ -currents. The latter are by definition continuous linear forms on the space of De Rham differential  $k$ -forms  $\Omega_{\text{dR}}^k(M)$ . This isomorphism is proved in [7].
8. Proposition 6.6 was established in [11]. The statement can be slightly enhanced. Namely, the quantities in Proposition 6.6 also only depend on the classes of  $u$  and  $p$  in the (odd and even) K-theory of  $\mathcal{A}$ . We refer to [11, Sect. IV.1.γ] for more details.
9. Originally, Connes introduced cyclic cohomology by means of cocycles satisfying such a *cyclic* condition, explaining the terminology. It turns out that this is equivalent to taking an even/odd cocycle in the  $(b, B)$ -bicomplex. For more details we refer to [7, Theorem II.40] (or [11, Theorem III.1.29]).

- 10.** The non-trivial evaluation of  $\phi$  on a projection in the noncommutative torus algebra plays a crucial role in the noncommutative geometric description of the integer quantum Hall effect. More details can be found in [15] and [11, Sect. IV.6. $\gamma$ ]. Exercise 6.10 gives an alternative way to show that  $\phi$  is non-trivial, and is very close to the computation that led to Proposition 6.1 dealing with the commutative case. It is based on the Powers–Rieffel projections introduced in [9].

### Section 6.3. Abstract differential calculus

- 11.** In our development of an abstract differential calculus we closely follow Connes and Moscovici [1]. In the case of the canonical triple of a spin manifold  $M$ , this will reproduce (part of) the usual differential calculus on  $M$ . We refer to [2] for a more detailed treatment. Note that the hypothesis that  $D$  is invertible can be removed, as described in [2, Sect. 6.1].
- 12.** The Sobolev spaces  $\mathcal{H}^s$  have appeared in the literature (including the first edition of this book) with a defining norm  $\|\xi\|^2 + \||D|^s \xi\|^2$ , while then operators of analytic order  $\leq r$  were defined as operators that extend to bounded maps from  $\mathcal{H}^s$  to  $\mathcal{H}^{s-r}$  for all  $s \geq 0$ . However, then one runs into the problem that  $|D|$  itself does not extend to a bounded operator from  $\mathcal{H}^0 \rightarrow \mathcal{H}^{-1}$ . This has been corrected in the current version, according to [16–18].
- 13.** The notion of finite summability for spectral triples was introduced in [11, Sect. IV.2. $\gamma$ ] (see also [12, Definition 10.8]).
- 14.** Even though we restrict to finitely-summable, regular spectral triples with simple dimension spectrum and for which there is a finite number of poles in  $S_d$ , the index formula can be proved in the presence of essential and infinitely many singularities as well [3–5].

### Section 6.5. The local index formula

- 15.** In our proof of Theorem 6.21 we follow Higson [2]. For the odd case, we refer to the original paper by Connes and Moscovici [1] (see also the more general [3]).
- 16.** The McKean–Singer formula is due to [19].
- 17.** For more details on the ‘transgression formula’ that is essential in the proof of Lemma 6.24 we refer to the discussion resulting in [12, Eq. 10.40].
- 18.** It is noted in [1, Remark II.1] that if  $(\mathcal{A}, \mathcal{H}, D)$  is the canonical triple associated to a Riemannian spin manifold  $M$ , then the local index formula of Connes and Moscovici reduces to the celebrated Atiyah–Singer index theorem for the Dirac operator [20, 21]. Namely, the operator  $D_p$  is then the Dirac operator with coefficients in a vector bundle  $E \rightarrow M$ . The latter is defined as a subbundle of the trivial bundle  $M \times \mathbb{C}^N$  using the projection  $p \in M_N(C(M))$ : one sets the fiber to be  $E_x = p(x)\mathbb{C}^N$  at each point  $x \in M$ . We then have

$$\text{index } D_p = (2\pi i)^{-\frac{n}{2}} \int_M \hat{A}(R) \wedge \text{ch}(E),$$

where  $\hat{A}(R)$  is the  $\hat{A}$ -form of the Riemannian curvature of  $M$  and  $\text{ch}(E)$  is the Chern character of the vector bundle  $E$  (cf. [22]). The proof exploits Getzler’s symbol calculus [23–25], as in [26]. See also [27].

## Section 6.6. The local index formula for toric noncommutative manifolds

19. Sect. 6.6 is based on [28, 29].
20. The appearance of  $U(s)$  in the proof of Theorem 6.25 is a consequence of the close relation with the index formula for a  $\mathbb{T}^n$ -equivariant Dirac spectral triple on  $M$ . In [30] Chern and Hu considered an even dimensional compact spin manifold  $M$  on which a (connected compact) Lie group  $G$  acts by isometries. The equivariant Chern character was defined as an equivariant version of the JLO-cocycle, the latter being an element in equivariant entire cyclic cohomology. The essential point is that they obtained an explicit formula for the above residues. Moreover, the vanishing of the term in Eq. (6.6.2) for  $|\alpha| \neq 0$  is [30, Theorem 2].

## References

1. Connes, A., Moscovici, H.: The local index formula in noncommutative geometry. *Geom. Funct. Anal.* **5**, 174–243 (1995)
2. N. Higson. The residue index theorem of Connes and Moscovici. In: Surveys In Noncommutative Geometry. Clay Mathematics Proceedings, pp. 71–126. American Mathematical Society, Providence, RI (2006)
3. Carey, A.L., Phillips, J., Rennie, A., Sukochev, F.A.: The local index formula in semifinite von Neumann algebras. I. Spectral flow. *Adv. Math.* **202**, 451–516 (2006)
4. Carey, A.L., Phillips, J., Rennie, A., Sukochev, F.A.: The local index formula in semifinite von Neumann algebras. II. The even case. *Adv. Math.* **202**, 517–554 (2006)
5. Carey, A.L., Phillips, J., Rennie, A., Sukochev, F.A.: The Chern character of semifinite spectral triples. *J. Noncommut. Geom.* **2**, 141–193 (2008)
6. Pedersen, G.K.: Analysis Now. Springer, New York (1989)
7. Connes, A.: Noncommutative differential geometry. *Publ. Math. IHES* **39**, 257–360 (1985)
8. T.A. Loring. *The Torus and Noncommutative Topology*. ProQuest LLC, Ann Arbor, MI. Thesis (Ph.D.)—University of California, Berkeley (1986)
9. Rieffel, M.A.:  $C^*$ -algebras associated with irrational rotations. *Pac. J. Math.* **93**, 415–429 (1981)
10. Epstein, P.: Zur Theorie allgemeiner Zetafunktionen. *Math. Ann.* **56**, 615–644 (1903)
11. Connes, A.: Noncommutative Geometry. Academic, San Diego (1994)
12. Gracia-Bondía, J.M., Várilly, J.C., Figueroa, H.: Elements of Noncommutative Geometry. Birkhäuser, Boston (2001)
13. Khalkhali, M.: Basic Noncommutative Geometry. EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich (2009)
14. Loday, J.-L.: Cyclic Homology. Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin (1992)
15. Bellissard, J.:  $K$ -theory of  $C^*$ -algebras in solid state physics. In: Statistical Mechanics And Field Theory: Mathematical Aspects (Groningen, 1985), Lecture Notes in Physics, vol. 257. pp. 99–156. Springer, Berlin (1986)
16. Uuye, O.: Pseudo-differential operators and regularity of spectral triples. In: Perspectives on Noncommutative Geometry, Fields Institute Communications, vol. 61, pp. 153–163. American Mathematical Society, Providence, RI (2011)
17. Hekkelman, E.-M., van Nuland, T.: Private communication
18. Multiple operator integrals, pseudodifferential calculus, and asymptotic expansions Eva-Maria Hekkelman, Edward McDonald, Teun D. H. van Nuland. [ArXiv:2404.16338](https://arxiv.org/abs/2404.16338)
19. McKean, H.P., Jr., Singer, I.M.: Curvature and the eigenvalues of the Laplacian. *J. Differ. Geometry* **1**, 43–69 (1967)

20. Atiyah, M.F., Singer, I.M.: The index of elliptic operators I. *Ann. Math.* **87**, 484–530 (1968)
21. Atiyah, M.F., Singer, I.M.: The index of elliptic operators III. *Ann. Math.* **87**, 546–604 (1968)
22. Berline, N., Getzler, E., Vergne, M.: Heat Kernels and Dirac Operators. Springer, Berlin (1992)
23. Getzler, E.: Pseudodifferential operators on supermanifolds and the Atiyah-Singer index theorem. *Commun. Math. Phys.* **92**, 163–178 (1983)
24. Getzler, E.: A short proof of the local Atiyah-Singer index theorem. *Topology* **25**, 111–117 (1986)
25. Getzler, E.: The odd Chern character in cyclic homology and spectral flow. *Topology* **32**, 489–507 (1993)
26. Block, J., Fox, J.: Asymptotic pseudodifferential operators and index theory. In: Geometric and Topological Invariants of Elliptic Operators (Brunswick, ME, 1988), Contemporary Mathematics, vol. 105, pp. 1–32. American Mathematical Society, Providence, RI (1990)
27. Ponge, R.: A new short proof of the local index formula and some of its applications. *Commun. Math. Phys.* **241**, 215–234 (2003)
28. Landi, G., van Suijlekom, W.D.: Principal fibrations from noncommutative spheres. *Commun. Math. Phys.* **260**, 203–225 (2005)
29. Landi, G., van Suijlekom, W.D.: Noncommutative instantons from twisted conformal symmetries. *Commun. Math. Phys.* **271**, 591–634 (2007)
30. Chern, S., Hu, X.: Equivariant Chern character for the invariant Dirac operator. *Michigan Math. J.* **44**, 451–473 (1997)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



**Part II**

**Noncommutative Geometry and Gauge  
Theories**

# Chapter 7

## Gauge Theories from Noncommutative Manifolds



In this chapter we demonstrate how every noncommutative (Riemannian spin) manifold, *viz.* every spectral triple, gives rise to a gauge theory in a generalized sense. We derive so-called inner fluctuations via Morita equivalences and interpret these as generalized gauge fields. This is quite similar to the construction in the finite case in Chaps. 2 and 3.

### 7.1 ‘Inner’ Unitary Equivalences as the Gauge Group

In Chap. 2 we already noticed the special role played by the unitary elements in the matrix algebras, and how they give rise to equivalences of finite noncommutative spaces (*cf.* Remark 2.25). We now extend this to general real spectral triples  $(\mathcal{A}, \mathcal{H}, D; J, \gamma)$ .

**Definition 7.1** A  $*$ -automorphism of a  $*$ -algebra  $\mathcal{A}$  is a linear invertible map  $\alpha : \mathcal{A} \rightarrow \mathcal{A}$  that satisfies

$$\alpha(ab) = \alpha(a)\alpha(b), \quad \alpha(a^*) = \alpha(a)^*.$$

We denote the group of automorphisms of the  $*$ -algebra  $\mathcal{A}$  by  $\text{Aut}(\mathcal{A})$ .

An automorphism  $\alpha$  is called *inner* if it is of the form  $\alpha(a) = uau^*$  for some element  $u \in \mathcal{U}(\mathcal{A})$  where

$$\mathcal{U}(\mathcal{A}) = \{u \in \mathcal{A} : uu^* = u^*u = 1\}$$

is the group of unitary elements in  $\mathcal{A}$ . The group of inner automorphisms is denoted by  $\text{Inn}(\mathcal{A})$ .

The group of *outer automorphisms* of  $\mathcal{A}$  is defined by the quotient

$$\text{Out}(\mathcal{A}) := \text{Aut}(\mathcal{A}) / \text{Inn}(\mathcal{A}).$$

Note that  $\text{Inn}(\mathcal{A})$  is indeed a normal subgroup of  $\text{Aut}(\mathcal{A})$  since

$$\beta \circ \alpha_u \circ \beta^{-1}(a) = \beta(u\beta^{-1}(a)u^*) = \beta(u)a\beta(u)^* = \alpha_{\beta(u)}(a),$$

for any  $\beta \in \text{Aut}(\mathcal{A})$ .

An inner automorphism  $\alpha_u$  is completely determined by the unitary element  $u \in \mathcal{U}(\mathcal{A})$ , but not in a unique manner. In other words, the map  $\phi: \mathcal{U}(\mathcal{A}) \rightarrow \text{Inn}(\mathcal{A})$  given by  $u \mapsto \alpha_u$  is surjective, but not injective. The kernel is given by  $\ker(\phi) = \{u \in \mathcal{U}(\mathcal{A}) \mid uau^* = a, a \in \mathcal{A}\}$ . In other words,  $\ker \phi = \mathcal{U}(Z(\mathcal{A}))$  where  $Z(\mathcal{A})$  is the center of  $\mathcal{A}$ . We conclude that the group of inner automorphisms is given by the quotient

$$\text{Inn}(\mathcal{A}) \simeq \mathcal{U}(\mathcal{A})/\mathcal{U}(Z(\mathcal{A})). \quad (7.1.1)$$

This can be summarized by the following exact sequences:

$$\begin{aligned} 1 &\longrightarrow \text{Inn}(\mathcal{A}) \longrightarrow \text{Aut}(\mathcal{A}) \longrightarrow \text{Out}(\mathcal{A}) \longrightarrow 1, \\ 1 &\longrightarrow \mathcal{U}(Z(\mathcal{A})) \longrightarrow \mathcal{U}(\mathcal{A}) \longrightarrow \text{Inn}(\mathcal{A}) \longrightarrow 1. \end{aligned}$$

**Example 7.2** If  $\mathcal{A}$  is a commutative  $*$ -algebra, then there are no non-trivial inner automorphisms since  $Z(\mathcal{A}) = \mathcal{A}$ . Moreover, if  $\mathcal{A} = C^\infty(X)$  with  $X$  a smooth compact manifold, then  $\text{Aut}(\mathcal{A}) \simeq \text{Diff}(X)$ , the group of diffeomorphisms of  $X$ . Explicitly, a diffeomorphism  $\phi: X \rightarrow X$  yields an automorphisms by pullback of a function  $f$ :

$$\phi^*(f)(x) = f(\phi(x)); \quad (x \in X).$$

Compare this with the discussion in the case of finite discrete topological spaces in Sect. 2.1. More generally, there is a continuous version of the above group isomorphism, relating  $\text{Aut}(C(X))$  one-to-one to homeomorphisms of  $X$ . This follows from functoriality of Gelfand duality. Namely, the Gelfand transform in Theorem 5.7 naturally extends to homomorphisms between commutative unital  $C^*$ -algebras, mapping these to homeomorphism between the corresponding structure spaces.

The fact that all automorphisms of  $C^\infty(X)$  come from a diffeomorphism of  $X$  can be seen as follows. Consider a smooth family  $\{\alpha_t\}_{t \in [0,1]}$  of automorphisms of  $C^\infty(X)$  from  $\alpha_{t=0} = \text{id}$  to  $\alpha_{t=1} = \alpha$ . The derivative at  $t = 0$  of this family,  $\dot{\alpha} := d\alpha_t/dt|_{t=0}$ , is a  $*$ -algebra derivation, since

$$\dot{\alpha}(f_1 f_2) = \frac{d}{dt} \alpha_t(f_1 f_2)|_{t=0} = \frac{d}{dt} \alpha_t(f_1) \alpha_t(f_2)|_{t=0} = \dot{\alpha}(f_1) f_2 + f_1 \dot{\alpha}(f_2).$$

As such,  $\dot{\alpha}$  corresponds to a smooth vector field on  $X$  and the end point  $\phi_{t=1}$  of the flow  $\phi_t$  of this vector field is the sought-for diffeomorphism of  $X$ . Its pullback  $\phi_{t=1}^*$  on smooth functions coincides with the automorphism  $\alpha_{t=1} = \alpha$ .

**Example 7.3** At the other extreme, we consider an example where all automorphisms are inner. Let  $\mathcal{A} = M_N(\mathbb{C})$  and let  $u$  be an element in the unitary group  $U(N)$ . Then  $u$  acts as an automorphism on  $a \in M_N(\mathbb{C})$  by sending  $a \mapsto uau^*$ . If  $u = \lambda \mathbb{I}_N$  is a multiple of the identity with  $\lambda \in U(1)$ , this action is trivial, hence the group of automorphisms of  $\mathcal{A}$  is the projective unitary group  $PU(N) = U(N)/U(1)$ , in concordance with (7.1.1).

The fact that all automorphisms are inner follows from the following observation. First, any  $*$ -algebra map  $\alpha : M_N(\mathbb{C}) \rightarrow M_N(\mathbb{C})$  can be considered a representation of  $\mathcal{A}$  on  $\mathbb{C}^N$ . As the unique irreducible representation space of  $M_N(\mathbb{C})$  is given by the defining representation (Lemma 2.15) we conclude that the representation  $\alpha$  is unitarily equivalent to the defining representation on  $\mathbb{C}^N$ . Hence,  $\alpha(a) = uau^*$  with  $u \in U(N)$ .

**Exercise 7.1** Show that  $\text{Aut}(M_N(\mathbb{C}) \oplus M_N(\mathbb{C})) \simeq (PU(N) \times PU(N)) \rtimes S_2$  with the symmetric group  $S_2$  acting by permutation on the two copies of  $PU(N)$ .

Inner automorphisms  $\alpha_u$  not only act on the  $*$ -algebra  $\mathcal{A}$ , via the representation  $\pi : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$  they also act on the Hilbert space  $\mathcal{H}$  present in the spectral triple. In fact, with  $U = \pi(u)J\pi(u)J^{-1}$ , the unitary  $u$  induces a unitary equivalence of real spectral triples in the sense of Definition 3.4, as the following exercise shows.

**Exercise 7.2** Use Definition 5.9 to establish the following transformation rules for a unitary  $U = \pi(u)J\pi(u)J^{-1}$  with  $u \in \mathcal{U}(\mathcal{A})$ :

$$\begin{aligned} U\pi(a)U^* &= \pi \circ \alpha_u(a); \\ U\gamma &= \gamma U; \\ UJu^* &= J. \end{aligned} \tag{7.1.2}$$

We conclude that an inner automorphism  $\alpha_u$  of  $\mathcal{A}$  induces a unitary equivalent spectral triple  $(\mathcal{A}, \mathcal{H}, UDU^*; J, \gamma)$ , where the action of the  $*$ -algebra is given by  $\pi \circ \alpha_u$ . Note that the grading and the real structure are left unchanged under these ‘inner’ unitary equivalences; only the operator  $D$  is affected by the unitary transformation. For the latter, we compute, using (5.2.1),

$$D \mapsto UDU^* = D + u[D, u^*] + \epsilon' Ju[D, u^*]J^{-1}, \tag{7.1.3}$$

where as before we have suppressed the representation  $\pi$ . We recognize the extra terms as *pure gauge* fields  $udu^*$  in the space of Connes’ differential one-forms  $\Omega_D^1(\mathcal{A})$  of Definition 5.15. This motivates the following definition

**Definition 7.4** The *gauge group*  $\mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$  of the spectral triple is

$$\mathfrak{G}(\mathcal{A}, \mathcal{H}; J) := \{U = uJuJ^{-1} \mid u \in \mathcal{U}(\mathcal{A})\}.$$

Recall (from Sect. 8.1) the construction of a complex subalgebra  $\mathcal{A}_J$  in the center of  $\mathcal{A}$  from a real spectral triple  $(\mathcal{A}, \mathcal{H}, D; J)$ , given by

$$\mathcal{A}_J := \{a \in \mathcal{A} : aJ = Ja^*\}.$$

**Proposition 7.5** *There is a short exact sequence of groups*

$$1 \rightarrow \mathcal{U}(\mathcal{A}_J) \rightarrow \mathcal{U}(\mathcal{A}) \rightarrow \mathfrak{G}(\mathcal{A}, \mathcal{H}; J) \rightarrow 1.$$

Moreover, there is a surjective map  $\mathfrak{G}(\mathcal{A}, \mathcal{H}; J) \rightarrow \text{Inn}(\mathcal{A})$ .

**Proof** Consider the map  $\text{Ad}: \mathcal{U}(\mathcal{A}) \rightarrow \mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$  given by  $u \mapsto uJuJ^{-1}$ . This map  $\text{Ad}$  is a group homomorphism, since the commutation relation  $[u, JvJ^{-1}] = 0$  of (5.2.1) implies that

$$\text{Ad}(v)\text{Ad}(u) = vJuJ^{-1}uJuJ^{-1} = vuJuJ^{-1} = \text{Ad}(vu).$$

By definition  $\text{Ad}$  is surjective, and  $\ker(\text{Ad}) = \{u \in \mathcal{U}(\mathcal{A}) \mid uJuJ^{-1} = 1\}$ . The relation  $uJuJ^{-1} = 1$  is equivalent to  $uJ = Ju^*$  which is the defining relation of the commutative subalgebra  $\mathcal{A}_J$ . This proves that  $\ker(\text{Ad}) = \mathcal{U}(\mathcal{A}_J)$ . The map  $\mathfrak{G}(\mathcal{A}, \mathcal{H}; J) \rightarrow \text{Inn}(\mathcal{A})$  is given by (7.1.2), from which surjectivity readily follows.  $\square$

**Corollary 7.6** *If  $\mathcal{U}(\mathcal{A}_J) = \mathcal{U}(Z(\mathcal{A}))$ , then  $\mathfrak{G}(\mathcal{A}, \mathcal{H}; J) \simeq \text{Inn}(\mathcal{A})$ .*

**Proof** This is immediate from the above Proposition and (7.1.1).  $\square$

We summarize this by the following sequence, which is exact in the horizontal direction:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathcal{U}(\mathcal{A}_J) & \longrightarrow & \mathcal{U}(\mathcal{A}) & \longrightarrow & \mathfrak{G}(\mathcal{A}, \mathcal{H}; J) \longrightarrow 1 \\ & & \downarrow & & \parallel & & \downarrow \\ 1 & \longrightarrow & \mathcal{U}(Z(\mathcal{A})) & \longrightarrow & \mathcal{U}(\mathcal{A}) & \longrightarrow & \text{Inn}(\mathcal{A}) \longrightarrow 1 \end{array}$$

### 7.1.1 The Gauge Algebra

A completely analogous discussion applies to the definition of a gauge Lie algebra, where instead of automorphisms we now take (inner and outer) **derivations** of  $\mathcal{A}$ . The following definition essentially gives the infinitesimal version of  $\mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$ .

**Definition 7.7** The *gauge Lie algebra*  $\mathfrak{g}(\mathcal{A}, \mathcal{H}; J)$  of the spectral triple is

$$\mathfrak{g}(\mathcal{A}, \mathcal{H}; J) := \{T = X + JXJ^{-1} \mid X \in \mathfrak{u}(\mathcal{A})\},$$

where  $\mathfrak{u}(\mathcal{A})$  consists of the skew-hermitian elements in  $\mathcal{A}$ .

One easily checks using the commutant property,

$$[T, T'] = [X, X'] + J[X, X']J^{-1},$$

so that  $\mathfrak{g}(\mathcal{A}, \mathcal{H}; J)$  is indeed a Lie algebra.

**Proposition 7.8** *There is a short exact sequence of Lie algebras*

$$0 \rightarrow \mathfrak{u}(\mathcal{A}_J) \rightarrow \mathfrak{u}(\mathcal{A}) \rightarrow \mathfrak{g}(\mathcal{A}, \mathcal{H}; J) \rightarrow 0.$$

There are also inner derivations of  $\mathcal{A}$  that are of the form  $a \rightarrow [X, a]$ ; these form a Lie subalgebra  $\text{Der}_{\text{Inn}}(\mathcal{A})$  of the Lie algebra of all derivations  $\text{Der}(\mathcal{A})$ . If  $\mathfrak{u}(\mathcal{A}_J) = \mathfrak{u}(Z(\mathcal{A}))$  then

$$\mathfrak{g}(\mathcal{A}, \mathcal{H}; J) \simeq \text{Der}_{\text{Inn}}(\mathcal{A}),$$

which essentially is the infinitesimal version of Corollary 7.6.

**Exercise 7.3** Show that  $\text{Der}(M_N(\mathbb{C})) \simeq su(N)$  as Lie algebras.

## 7.2 Morita Self-equivalences as Gauge Fields

We have seen that a non-abelian gauge group appears naturally when the unital  $*$ -algebra  $\mathcal{A}$  in a real spectral triple is noncommutative. Moreover, noncommutative algebras allow for a more general—and in fact more natural—notion of equivalence than automorphic equivalence, namely Morita equivalence. We have already seen this in Chap. 2. Indeed, let us imitate the construction in Theorem 2.26 and 3.6 and see if we can lift Morita equivalence to the level of spectral triples in this more general setting.

Let us first recall some of the basic definitions. We keep working in the setting of unital algebras, which greatly simplifies matters (See Note 4 in this chapter).

### 7.2.1 Morita Equivalence

Recall Definition 2.8 of algebra modules. For two right  $\mathcal{A}$ -modules  $\mathcal{E}$  and  $\mathcal{F}$  we denote the space of right  **$\mathcal{A}$ -module homomorphisms** by  $\text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{F})$ , i.e.

$$\text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{F}) := \{\phi : \mathcal{E} \rightarrow \mathcal{F} : \phi(\eta a) = \phi(\eta)a \text{ for all } \eta \in \mathcal{E}, a \in A\}. \quad (7.2.1)$$

We also write  $\text{End}_{\mathcal{A}}(\mathcal{E}) := \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{E})$  for the **algebra of right  $\mathcal{A}$ -module endomorphisms** of  $\mathcal{E}$ .

**Definition 7.9** Two unital algebras  $\mathcal{A}$  and  $\mathcal{B}$  are called *Morita equivalent* if there exists a  $\mathcal{B} - \mathcal{A}$ -bimodule  $\mathcal{E}$  and an  $\mathcal{A} - \mathcal{B}$ -bimodule  $\mathcal{F}$  such that

$$\mathcal{E} \otimes_{\mathcal{A}} \mathcal{F} \simeq \mathcal{B}, \quad \mathcal{F} \otimes_{\mathcal{B}} \mathcal{E} \simeq \mathcal{A},$$

as  $\mathcal{B}$  and  $\mathcal{A}$ -bimodules, respectively.

**Exercise 7.4** Taking inspiration from Exercise 2.9, show that Morita equivalence is an equivalence relation.

**Exercise 7.5** Define  $\mathcal{A}^N = \mathcal{A} \oplus \cdots \oplus \mathcal{A}$  ( $N$  copies) as an  $\mathcal{A} - M_N(\mathcal{A})$ -bimodule.

1. Show that  $\mathcal{A}^N \otimes_{\mathcal{A}} \mathcal{A}^N \simeq M_N(\mathcal{A})$ , as  $M_N(\mathcal{A}) - M_N(\mathcal{A})$ -bimodules.
2. Show that  $\mathcal{A}^N \otimes_{M_N(\mathcal{A})} \mathcal{A}^N \simeq \mathcal{A}$ , so that  $M_N(\mathcal{A})$  is Morita equivalent to  $\mathcal{A}$ .

A convenient characterisation of Morita equivalent algebras is given by the concept of endomorphism algebras of so-called finitely generated projective modules, as we now explain.

**Definition 7.10** A right  $\mathcal{A}$ -module is called *finitely generated projective* (or, briefly, *finite projective*) if there is an idempotent  $p = p^2$  in  $M_N(\mathcal{A})$  for some  $N$  such that  $\mathcal{E} \simeq p\mathcal{A}^N$ .

**Lemma 7.11** A right  $\mathcal{A}$ -module is finitely generated projective if and only if

$$\text{End}_{\mathcal{A}}(\mathcal{E}) \simeq \mathcal{E} \otimes_{\mathcal{A}} \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A}).$$

**Proof** First note that the right-hand side can be considered to be a two-sided ideal in  $\text{End}_{\mathcal{A}}(\mathcal{E})$ . Namely, we consider an element  $\eta \otimes_{\mathcal{A}} \phi$  in  $\mathcal{E} \otimes_{\mathcal{A}} \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A})$  as an element in  $\text{End}_{\mathcal{A}}(\mathcal{E})$  by mapping

$$\xi \mapsto \eta\phi(\xi); \quad (\xi \in \mathcal{E}).$$

That this map is injective and that its image forms an ideal in  $\text{End}_{\mathcal{A}}(\mathcal{E})$  is readily checked. Hence, the above isomorphism is equivalent to the existence of an element in  $\mathcal{E} \otimes_{\mathcal{A}} \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A})$  that acts as the identity map  $\text{id}_{\mathcal{E}}$  on  $\mathcal{E}$ .

Suppose that  $\mathcal{E}$  is finite projective,  $\mathcal{E} \simeq p\mathcal{A}^N$  for some idempotent  $p \in M_N(\mathcal{A})$ . We identify two maps

$$\begin{aligned} \lambda : \mathcal{E} &\rightarrow \mathcal{A}^N, \\ \rho : \mathcal{A}^N &\rightarrow \mathcal{E}, \end{aligned}$$

which are injective and surjective, respectively. These maps are related to the identification of  $\mathcal{E}$  with a direct summand of  $\mathcal{A}^N$ , via the obvious direct sum decomposition  $\mathcal{A}^N = p\mathcal{A}^N \oplus (1-p)\mathcal{A}^N$ . Namely,  $\lambda$  identifies  $\mathcal{E}$  with  $p\mathcal{A}^N \subset \mathcal{A}^N$ , whereas  $\rho$  projects  $\mathcal{A}^N$  onto the direct summand  $p\mathcal{A}^N$  and then identifies it with  $\mathcal{E}$ . Let us write  $\lambda_k$  for the  $k$ th component of  $\lambda$  mapping  $\mathcal{E}$  to  $\mathcal{A}^N$ ; thus,  $\lambda_k : \mathcal{E} \rightarrow \mathcal{A}$  is right  $\mathcal{A}$ -linear

for any  $k = 1, \dots, N$ . We write  $\rho_k := \rho(e_k) \in \mathcal{E}$ , where  $\{e_k\}_{k=1}^N$  is the standard basis of  $\mathcal{A}^N$ . The composition  $\sum_{k=1}^N \rho_k \otimes \lambda_k$  then acts as the identity operator on  $\mathcal{E}$ .

Conversely, suppose  $\text{id}_{\mathcal{E}}$  can be written as a finite sum

$$\sum_{k=1}^N \rho_k \otimes \lambda_k \in \mathcal{E} \otimes_{\mathcal{A}} \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A}). \quad (7.2.2)$$

Reversing the construction in the previous paragraph, we are now going to define an idempotent  $p \in M_N(\mathcal{A})$  such that  $\mathcal{E} \simeq p\mathcal{A}^N$ . Thus, we define maps

$$\begin{aligned} \lambda : \mathcal{E} &\rightarrow \mathcal{A}^N; & \eta &\mapsto (\lambda_1(\eta), \dots, \lambda_N(\eta)), \\ \rho : \mathcal{A}^N &\rightarrow \mathcal{E}; & (a_1, \dots, a_N) &\mapsto \rho_1 a_1 + \dots + \rho_N a_N. \end{aligned}$$

From their very definition, these maps satisfy  $\rho \circ \lambda = \text{id}_{\mathcal{E}}$ , so that  $p = \lambda \circ \rho$  is the sought-for idempotent in  $M_N(\mathcal{A})$ .  $\square$

**Exercise 7.6** In this exercise we are going to analyze the ambiguity due to the balanced tensor product that appears in the decomposition (7.2.2) of  $\text{id}_{\mathcal{E}}$ .

1. If  $\mathcal{E} = \mathcal{A}$  then  $\text{id}_{\mathcal{E}} = 1 \otimes 1 \subset \mathcal{E} \otimes_{\mathcal{A}} \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A})$  but also

$$\text{id}_{\mathcal{E}} = 1 \otimes 1 + a \otimes 1 + 1 \otimes (-a),$$

for any  $a \in \mathcal{A}$ . Show that the projection corresponding to the latter decomposition of  $\text{id}_{\mathcal{E}}$  is

$$p = \begin{pmatrix} 1 & 1 & -a \\ a & a & -a^2 \\ 1 & 1 & -a \end{pmatrix}.$$

2. Show that there is a similarity transformation  $S$  such that

$$SpS^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Therefore, the projection corresponding to  $\text{id}_{\mathcal{E}} = 1 \otimes 1$  appears as the first diagonal entry, and we can conclude that both decompositions give isomorphic projective modules  $p\mathcal{A}^3 \simeq \mathcal{A}$ .

3. Extend this argument to any finite projective  $\mathcal{E}$  to show that the construction of a projection  $p$  from (7.2.2) is well defined.

**Proposition 7.12** *Two unital algebras  $\mathcal{A}$  and  $\mathcal{B}$  are Morita equivalent if and only if  $\mathcal{B} \simeq \text{End}_{\mathcal{A}}(\mathcal{E})$ , with  $\mathcal{E}$  a finite projective  $\mathcal{A}$ -module.*

**Proof** If  $\mathcal{B} \simeq \text{End}_{\mathcal{A}}(\mathcal{E})$  for some finite projective  $\mathcal{E}$ , then  $\mathcal{F} = \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A})$  is the required  $\mathcal{A} - \mathcal{B}$ -bimodule implementing the desired Morita equivalence, with bimodule structure given by

$$(a \cdot \phi \cdot b)(\eta) = a\phi(b \cdot \eta); \quad (\phi \in \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A})). \quad (7.2.3)$$

The property  $\mathcal{E} \otimes_{\mathcal{A}} \mathcal{F} \simeq \mathcal{B}$  follows from Lemma 7.11, and the isomorphism  $\mathcal{F} \otimes_{\mathcal{B}} \mathcal{E} \simeq \mathcal{A}$  is implemented by the evaluation map, that is,

$$(\phi \otimes \eta) \in \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A}) \otimes_{\mathcal{B}} \mathcal{E} \mapsto \phi(\eta) \in \mathcal{A}.$$

Conversely, suppose  $\mathcal{A}$  and  $\mathcal{B}$  are Morita equivalent. If  $\mathcal{B} \simeq \mathcal{E} \otimes_{\mathcal{A}} \mathcal{F}$ , then  $\mathcal{B} \simeq \text{End}_{\mathcal{B}}(\mathcal{B}) \simeq \text{End}_{\mathcal{B}}(\mathcal{E} \otimes_{\mathcal{A}} \mathcal{F})$ , and there is an algebra map

$$\begin{aligned} \text{End}_{\mathcal{A}}(\mathcal{E}) &\rightarrow \text{End}_{\mathcal{B}}(\mathcal{E} \otimes_{\mathcal{A}} \mathcal{F}); \\ \phi &\mapsto \phi \otimes 1_{\mathcal{F}}. \end{aligned}$$

On the other hand,  $\text{End}_{\mathcal{A}}(\mathcal{B} \otimes_{\mathcal{B}} \mathcal{E}) \simeq \text{End}_{\mathcal{A}}(\mathcal{E})$ , and there is an algebra map

$$\begin{aligned} \text{End}_{\mathcal{B}}(\mathcal{B}) &\rightarrow \text{End}_{\mathcal{A}}(\mathcal{B} \otimes_{\mathcal{B}} \mathcal{E}); \\ \phi' &\mapsto \phi' \otimes 1_{\mathcal{E}}. \end{aligned}$$

Identifying  $\mathcal{E} \otimes_{\mathcal{A}} \mathcal{F} \simeq \mathcal{B}$  and  $\mathcal{F} \otimes_{\mathcal{B}} \mathcal{E} \simeq \mathcal{A}$ , one readily checks that these two maps are each other's inverses. This shows that  $\mathcal{B} \simeq \text{End}_{\mathcal{A}}(\mathcal{E})$ .

Finally, the fact that the right  $\mathcal{A}$ -module  $\mathcal{E}$  is finitely generated and projective follows *mutatis mutandis* from the proof of Lemma 7.11, after realizing that the isomorphism  $\mathcal{F} \otimes_{\mathcal{B}} \mathcal{E} \simeq \mathcal{A}$  associates an element in  $\text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A})$  to any element in  $\mathcal{F}$ .  $\square$

**Exercise 7.7** Show that (7.2.3) is a well-defined  $\mathcal{A} - \mathcal{B}$ -bimodule structure on  $\text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A})$ , i.e. show that it respects the right  $\mathcal{A}$ -linearity of the map  $\phi : \mathcal{E} \rightarrow \mathcal{A}$ .

We conclude this subsection by specializing from algebras to  $*$ -algebras. The above results on Morita equivalence still hold, with the additional requirement that in the definition of finite projectivity the idempotent  $p \in M_N(\mathbb{C})$  needs to be self-adjoint:  $p^* = p$ . That is to say,  $p$  is an **orthogonal projection**.

As in Definition 3.5, we define the **conjugate module**  $\mathcal{E}^\circ$  to a right  $\mathcal{A}$ -module  $\mathcal{E}$  as

$$\mathcal{E}^\circ = \{\bar{\xi} : \xi \in \mathcal{E}\},$$

equipped with a left  $\mathcal{A}$  action defined by  $a\bar{\xi} = \overline{\xi a^*}$  for any  $a \in \mathcal{A}$ .

**Proposition 7.13** *If  $\mathcal{A}$  is a  $*$ -algebra and  $\mathcal{E}$  is a finite projective right  $\mathcal{A}$ -module, then we can identify  $\text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A})$  as a left  $\mathcal{A}$ -module with the conjugate module  $\mathcal{E}^\circ$ ,*

**Proof** If  $\mathcal{E} \simeq p\mathcal{A}^N$  then  $\text{End}_{\mathcal{A}}(\mathcal{E}) \simeq pM_N(\mathcal{A})p$ , as one can easily show using the maps  $\lambda$  and  $\rho$  from the first part of the proof of Lemma 7.11. Hence  $\mathcal{E} \otimes_{\mathcal{A}} \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A}) \simeq pM_N(\mathcal{A})p$ . But also  $p\mathcal{A}^N \otimes_{\mathcal{A}} \mathcal{A}^N p \simeq pM_N(\mathcal{A})p$  (cf. Exercise 7.5), so  $\text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{A}) \simeq \mathcal{A}^N p$  as left  $\mathcal{A}$ -modules. We now show that  $\mathcal{E}^\circ \simeq \mathcal{A}^N p$  as well.

For that, write  $\xi \in \mathcal{E} \simeq p\mathcal{A}^N$  as a column vector:

$$\xi = \begin{pmatrix} \sum_{j=1}^N p_{1j} a_j \\ \vdots \\ \sum_{j=1}^N p_{Nj} a_j \end{pmatrix}.$$

The corresponding element  $\bar{\xi}$  in  $\mathcal{E}^\circ$  is identified with

$$\left( \sum_{j=1}^N a_j^* p_{j1} \quad \cdots \quad \sum_{j=1}^N a_j^* p_{jN} \right),$$

written as a row vector in  $\mathcal{A}^N p$ . Note that the relation between  $\xi$  and this row vector is essentially given by the involution on  $\mathcal{A}^N$ , exploiting the self-adjointness of  $p$ , that is,  $p_{ji}^* = p_{ij}$ . Consequently, the element  $a\bar{\xi} = \bar{\xi}a^*$  is mapped to

$$a \left( \sum_{j=1}^N a_j^* p_{j1} \quad \cdots \quad \sum_{j=1}^N a_j^* p_{jN} \right),$$

as required.  $\square$

**Proposition 7.14** *Let  $\mathcal{A}$  be a  $*$ -algebra and  $\mathcal{E}$  a finite projective right  $\mathcal{A}$ -module. Then there exists a hermitian structure on  $\mathcal{E}$ , that is to say, there is a pairing  $\langle \cdot, \cdot \rangle_{\mathcal{E}} : \mathcal{E} \times \mathcal{E} \rightarrow \mathcal{A}$  on  $\mathcal{E}$  that satisfies (as in Definition 2.9)*

$$\begin{aligned} \langle \eta_1, \eta_2 \cdot a \rangle_{\mathcal{E}} &= \langle \eta_1, \eta_2 \rangle_{\mathcal{E}} a; \quad (\eta_1, \eta_2 \in \mathcal{E}, a \in \mathcal{A}), \\ \langle \eta_1, \eta_2 \rangle_{\mathcal{E}}^* &= \langle \eta_2, \eta_1 \rangle_{\mathcal{E}}; \quad (\eta_1, \eta_2 \in \mathcal{E}), \\ \langle \eta, \eta \rangle_{\mathcal{E}} &\geq 0, \text{ with equality if and only if } \eta = 0; \quad (\eta \in \mathcal{E}). \end{aligned}$$

**Proof** On  $\mathcal{A}^N$  we have a hermitian structure given by

$$\langle \eta, \xi \rangle = \sum_{j=1}^N \eta_j^* \xi_j,$$

which satisfies the above properties. By restriction to  $p\mathcal{A}^N$  we then obtain a hermitian structure on  $\mathcal{E} \simeq p\mathcal{A}^N$ .  $\square$

### 7.2.2 Morita Equivalence and Spectral Triples

For a given spectral triple  $(\mathcal{A}, \mathcal{H}, D)$  and for a given finite projective right  $\mathcal{A}$ -module  $\mathcal{E}$ , we try to construct another spectral triple  $(\mathcal{B}, \mathcal{H}', D')$  where  $\mathcal{B} = \text{End}_{\mathcal{A}}(\mathcal{E})$ . This generalizes the finite-dimensional constructions of Chaps. 2 and 3. Naturally,

$$\mathcal{H}' := \mathcal{E} \otimes_{\mathcal{A}} \mathcal{H}$$

carries an action of  $\phi \in \mathcal{B}$ :

$$\phi(\eta \otimes \psi) = \phi(\eta) \otimes \psi; \quad (\eta \in \mathcal{E}, \psi \in \mathcal{H}).$$

Moreover, by finite projectivity of  $\mathcal{E}$ ,  $\mathcal{H}'$  is a Hilbert space. Indeed, we have

$$\mathcal{H}' \simeq p\mathcal{A}^N \otimes_{\mathcal{A}} \mathcal{H} \simeq p\mathcal{H}^N,$$

and since  $p$  is an orthogonal projection it has closed range.

However, the naive choice of an operator  $D'$  by  $D'(\eta \otimes \psi) = \eta \otimes D\psi$  will not do, because it does not respect the ideal defining the tensor product over  $\mathcal{A}$ , which is generated by elements of the form

$$\eta a \otimes \psi - \eta \otimes a\psi; \quad (\eta \in \mathcal{E}, a \in \mathcal{A}, \psi \in \mathcal{H}).$$

A better definition is

$$(1 \otimes_{\nabla} D)(\eta \otimes \psi) = \eta \otimes D\psi + \nabla(\eta)\psi.$$

where  $\nabla : \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \Omega_D^1(\mathcal{A})$  is a **connection** associated to the derivation  $d : a \mapsto [D, a]$  ( $a \in \mathcal{A}$ ). This means that  $\nabla$  is a linear map that satisfies the Leibniz rule:

$$\nabla(\eta a) = (\nabla\eta)a + \eta \otimes_{\mathcal{A}} da; \quad (\eta \in \mathcal{E}, a \in \mathcal{A}).$$

- Exercise 7.8** (1) Let  $\nabla$  and  $\nabla'$  be two connections on a right  $\mathcal{A}$ -module  $\mathcal{E}$ . Show that their difference  $\nabla - \nabla'$  is a right  $\mathcal{A}$ -linear map  $\mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \Omega_D^1(\mathcal{A})$ .  
(2) Show that the following map defines a connection on  $\mathcal{E} = p\mathcal{A}^N$ :

$$\nabla = p \circ d,$$

with  $d$  acting on each copy of  $\mathcal{A}$  as the commutator  $[D, \cdot]$ . This connection is referred to as the **Grassmann connection** on  $\mathcal{E}$ .

**Theorem 7.15** *If  $(\mathcal{A}, \mathcal{H}, D)$  is a spectral triple and  $\nabla$  is a connection on a finite projective right  $\mathcal{A}$ -module  $\mathcal{E}$ , then  $(\mathcal{B}, \mathcal{H}', 1 \otimes_{\nabla} D)$  is a spectral triple, provided that  $\nabla$  is a hermitian connection, i.e. provided that*

$$\langle \eta_1, \nabla \eta_2 \rangle_{\mathcal{E}} - \langle \nabla \eta_1, \eta_2 \rangle_{\mathcal{E}} = d\langle \eta_1, \eta_2 \rangle_{\mathcal{E}}; \quad (\eta_1, \eta_2 \in \mathcal{E}). \quad (7.2.4)$$

**Proof** Suppose  $\mathcal{E} = p\mathcal{A}^N$ , so that  $\mathcal{B} = \text{End}_{\mathcal{A}}(\mathcal{E}) \simeq pM_N(\mathcal{A})p$  and  $\mathcal{E} \otimes_{\mathcal{A}} \mathcal{H} \simeq p\mathcal{H}^N$ . The boundedness of the action of  $\mathcal{B}$  on  $\mathcal{E} \otimes_{\mathcal{A}} \mathcal{H}$  then follows directly from the boundedness of the action of  $\mathcal{A}$  on  $\mathcal{H}$ . Similarly, for  $\phi \in \mathcal{B}$  the commutator  $[D, \phi]$  can be regarded as a matrix with entries of the form  $[D, a]$  with  $a \in \mathcal{A}$ . These commutators are all bounded, so that  $[D, \phi]$  is bounded. Let us prove compactness of the resolvent. By Exercise 7.8 any connection can be written as  $\nabla = p \circ [D, \cdot] + \omega$  for a right  $\mathcal{A}$ -linear map  $\omega : \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \Omega_D^1(\mathcal{A})$ . Hence, after making the above identifications we see that the operator  $\nabla \otimes 1 + 1 \otimes D$  coincides with  $pDp + \omega$ . The action of  $\omega$  is as a bounded operator, which by (7.2.4) is self-adjoint. Moreover, it is given by a matrix acting on  $p\mathcal{H}^N$  with entries in  $\Omega_D^1(\mathcal{A})$ . Since for any self-adjoint operator  $T$  we have

$$(i + T + \omega)^{-1} = (i + T)^{-1} (1 - \omega(i + T + \omega)^{-1}),$$

with  $(1 - \omega(i + T + \omega)^{-1})$  bounded, compactness of the resolvent of  $pDp + \omega$  would follow from compactness of  $(ip + pDp)^{-1}$  (note that  $p$  is the identity on the Hilbert space  $p\mathcal{H}^N$ ). The required compactness property is a consequence of the identity

$$(ip + pDp)p(i + D)^{-1}p = p[i + D, p](i + D)^{-1}p + p.$$

Indeed, when multiplied on the left with  $(ip + pDp)^{-1}$  we find that on  $p\mathcal{H}^N$ :

$$(ip + pDp)^{-1} = p(i + D)^{-1}p - (ip + pDp)^{-1}p[D, p](i + D)^{-1}p,$$

which is compact since  $(i + D)^{-1}$  is compact by definition of a spectral triple.  $\square$

Analogously, for a given real spectral triple  $(\mathcal{A}, \mathcal{H}, D, J)$  we define another real spectral triple  $(\mathcal{B}, \mathcal{H}', D' = (1 \otimes_{\nabla} D) \otimes_{\nabla} 1; J')$  by setting

$$\mathcal{H}' := \mathcal{E} \otimes_{\mathcal{A}} \mathcal{H} \otimes_{\mathcal{A}} \mathcal{E}^{\circ}.$$

Then,  $\phi \in \mathcal{B}$  acts on  $\mathcal{H}'$  by

$$\phi(\eta \otimes \psi \otimes \bar{\xi}) = \phi(\eta) \otimes \psi \otimes \bar{\xi},$$

and the operator  $D'$  may be defined to be  $(1 \otimes_{\nabla} D) \otimes_{\nabla} 1$ , i.e.

$$D'(\eta \otimes \psi \otimes \bar{\xi}) = (\nabla\eta)\psi \otimes \bar{\xi} + \eta \otimes D\psi \otimes \bar{\xi} + \eta \otimes \psi(\bar{\nabla}\bar{\xi}),$$

while for  $J'$  we set

$$J'(\eta \otimes \psi \otimes \bar{\xi}) = \xi \otimes J\psi \otimes \bar{\eta}.$$

Finally, for even spectral triples one defines a grading  $\gamma'$  on  $\mathcal{E} \otimes_{\mathcal{A}} \mathcal{H} \otimes_{\mathcal{A}} \mathcal{E}^{\circ}$  by  $\gamma' = 1 \otimes \gamma \otimes 1$ . We have therefore proved:

**Theorem 7.16** *If  $(\mathcal{A}, \mathcal{H}, D; J, \gamma)$  is a real spectral triple and  $\nabla$  is a hermitian connection, then  $(\mathcal{B}, \mathcal{H}', D'; J', \gamma')$  is a real spectral triple.*

We now focus on **Morita self-equivalences**, for which  $\mathcal{B} = \mathcal{A}$  and  $\mathcal{E} = \mathcal{A}$  so that  $\text{End}_{\mathcal{A}}(\mathcal{E}) \simeq \mathcal{A}$ . Let us look at connections

$$\nabla : \mathcal{A} \rightarrow \Omega_D^1(\mathcal{A}).$$

Clearly, by the Leibniz rule we must have  $\nabla = d + \omega$  (see also Exercise 7.8), where  $\omega = \nabla(1) = \sum_j a_j [D, b_j]$  is a generic element in  $\Omega_D^1(\mathcal{A})$  acting as a bounded operator on  $\mathcal{H}$ . Similarly,  $\psi \bar{\nabla} \bar{a} = (\epsilon' J da J^{-1} + \epsilon' J \omega a J^{-1}) \psi$ . Since  $\mathcal{H}' \simeq \mathcal{H}$ , under this identification we have,

$$D'(\psi) \equiv D'(1 \otimes \psi \otimes \bar{1}) = \nabla(1)\psi + \psi \bar{\nabla}(\bar{1}) + D\psi = D\psi + \omega\psi + \epsilon' J \omega J^{-1} \psi.$$

In other words,  $D$  is ‘innerly perturbed’ by the given Morita self-equivalence to

$$D_{\omega} := D + \omega + \epsilon' J \omega J^{-1}, \quad (7.2.5)$$

where  $\omega^* = \omega \in \Omega_D^1(\mathcal{A})$  is called a **gauge field**, alternatively called an **inner fluctuation** of the operator  $D$ , since it is the algebra  $\mathcal{A}$  that—through Morita self-equivalences—generates the field  $\omega$ .

**Proposition 7.17** *A unitary equivalence of a real spectral triple  $(\mathcal{A}, \mathcal{H}, D; J)$  as implemented by  $U = u Ju J^{-1}$  with  $u \in \mathcal{U}(\mathcal{A})$  (discussed before Definition 7.4) is a special case of a Morita self-equivalence, arising by taking  $\omega = u[D, u^*]$ .*

**Proof** This follows upon inserting  $\omega = u[D, u^*]$  in the above formula for  $D_{\omega}$ , yielding (7.1.3).  $\square$

In the same way there is an action of the unitary group  $\mathcal{U}(\mathcal{A})$  on the new spectral triple  $(\mathcal{A}, \mathcal{H}, D_{\omega})$  by unitary equivalences. Recall that  $U = u Ju J^{-1}$  acts on  $D_{\omega}$  by conjugation:

$$D_{\omega} \mapsto U D_{\omega} U^*. \quad (7.2.6)$$

This is equivalent to

$$\omega \mapsto u \omega u^* + u[D, u^*],$$

which is the usual rule for a gauge transformation on a gauge field.

### 7.3 Inner Fluctuations Without the First-Order Condition

We now generalize inner fluctuations to real spectral triples that fail on the first-order condition. This will be used in the applications to particle physics beyond the Standard Model in Chap. 15.

Let us start with the following general result on Morita equivalence for spectral triples  $(\mathcal{A}, \mathcal{H}, D; J)$  that possibly do not satisfy the first-order condition. It turns out to be necessary to work with the  $\mathcal{A}$ -bimodule of universal differential one-forms  $\Omega^1(\mathcal{A})$ , instead of the Connes' differential one-forms  $\Omega_D^1(\mathcal{A})$ , see Note 7 in this chapter for a quick review on universal differential forms.

Just as in Theorem 7.16 of the previous section we now consider the operators induced on  $\mathcal{E} \otimes_{\mathcal{A}} \mathcal{H}$  by the operator  $D$  on  $\mathcal{H}$  and a (universal) connection  $\nabla$  on  $\mathcal{E}$ . We still exploit the same notation  $1 \otimes_{\nabla} D$ :

$$(1 \otimes_{\nabla} D)(\eta \otimes \psi) = (\nabla \eta)\psi + \eta \otimes D\psi,$$

with the universal one-form in the first term on the right-hand side acting via the representation  $\delta a \mapsto da = [D, a]$  on  $\mathcal{H}$ .

**Proposition 7.18** *Let  $(\mathcal{A}, \mathcal{H}, D; J)$  be a real spectral triple, possibly not fulfilling the first-order condition. Let  $\mathcal{E}$  be a finitely generated projective right  $\mathcal{A}$ -module, equipped with a universal connection  $\nabla : \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \Omega^1(\mathcal{A})$ . Then*

$$(1 \otimes_{\nabla} D) \otimes_{\overline{\nabla}} 1 = 1 \otimes_{\nabla} (D \otimes_{\overline{\nabla}} 1) \tag{7.3.1}$$

Moreover, the triple  $(\text{End}_{\mathcal{A}}(\mathcal{E}), \mathcal{E} \otimes_{\mathcal{A}} \mathcal{H} \otimes_{\mathcal{A}} \overline{\mathcal{E}}, D'; J')$  is a real spectral triple where  $D' = (1 \otimes_{\nabla} D) \otimes_{\overline{\nabla}} 1$  and the real structure is given by

$$J'(v_1 \otimes \xi \otimes \bar{v}_2) = (v_2 \otimes J\xi \otimes \bar{v}_1); \quad (v_1, v_2 \in \mathcal{E}, \xi \in \mathcal{H}).$$

**Proof** See Note 8 in this chapter. □

**Corollary 7.19** *If  $(\mathcal{A}, \mathcal{H}, D; J)$  satisfies the first-order condition, then so does  $(\text{End}_{\mathcal{A}}(\mathcal{E}), \mathcal{E} \otimes_{\mathcal{A}} \mathcal{H} \otimes_{\mathcal{A}} \overline{\mathcal{E}}, D'; J')$  and in that case the above inner fluctuation reduces to the usual one, given in terms of a connection  $\nabla : \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \Omega_D^1(\mathcal{A})$  (i.e. representing all universal connections using  $\delta \mapsto [D, \cdot]$ ).*

#### 7.3.1 Special Case $\mathcal{E} = \mathcal{A}$ and Inner Fluctuations

As a special case we take  $\mathcal{E} = \mathcal{A}$  and  $\nabla = \delta + A$  where  $A \in \Omega^1(\mathcal{A})$  is a self-adjoint, universal one-form

$$A = \sum_j a_j \delta(b_j); \quad (a_j, b_j \in \mathcal{A}). \tag{7.3.2}$$

Under the respective identifications  $\mathcal{H} = \mathcal{A} \otimes_{\mathcal{A}} \mathcal{H}$  and  $\mathcal{H} = \mathcal{H} \otimes_{\mathcal{A}} \mathcal{A}$ , we have

$$\begin{aligned} 1 \otimes_{\nabla} D &\simeq D + \sum_j a_j [D, b_j], \\ D \otimes_{\nabla} 1 &\simeq D + \sum_j \hat{a}_j [D, \hat{b}_j]. \end{aligned}$$

This then gives rise to the following Dirac operator

$$\begin{aligned} D' &= D + \sum_j a_j [D, b_j] + \sum_j \hat{a}_j [D, \hat{b}_j] + \sum_j \hat{a}_j [\omega_{(1)}, \hat{b}_j] \\ &=: D + \omega_{(1)} + \tilde{\omega}_{(1)} + \omega_{(2)} \end{aligned} \tag{7.3.3}$$

where we have defined

$$\begin{aligned} \omega_{(1)} &:= \sum_j a_j [D, b_j]; \\ \tilde{\omega}_{(1)} &:= \sum_j \hat{a}_j [D, \hat{b}_j]; \\ \omega_{(2)} &:= \sum_j \hat{a}_j [\omega_{(1)}, \hat{b}_j] \\ &= \sum_{j,k} \hat{a}_j a_k [[D, b_k], \hat{b}_j] \end{aligned}$$

The commutant property (5.2.1) shows that

$$\sum_j \hat{a}_j [\omega_{(1)}, \hat{b}_j] = \sum_{j,k} \hat{a}_j a_k [[D, b_k], \hat{b}_j] = \sum_{j,k} a_k \hat{a}_j [[D, \hat{b}_j], b_k] = \sum_k a_k [\tilde{\omega}_{(1)}, b_k]$$

which checks (7.3.1). Note that, with  $\epsilon = \pm 1$  such that  $J D J^{-1} = \epsilon D$  one has

$$\tilde{\omega}_{(1)} = \epsilon J \omega_{(1)} J^{-1}, \quad \omega_{(2)} = \epsilon J \omega_{(2)} J^{-1}$$

which follows from the commutant property (5.2.1).

It is clear from these formulas that  $\omega_{(2)}$  vanishes if  $(\mathcal{A}, \mathcal{H}, D; J)$  satisfies the first-order condition, thus reducing to formula (7.2.5) above. We will interpret the terms  $\omega_{(2)}$  as non-linear corrections to the *first-order*, linear inner fluctuations  $\omega_{(1)}$  of  $(\mathcal{A}, \mathcal{H}, D; J)$ . It is clear that the first order condition is equivalent to the linearity of the map from 1-forms to fluctuations. Let us check that the gauge transformations operate in the correct manner thanks to the quadratic correction term  $\omega_{(2)}$ . We shall understand this direct computation in a more conceptual manner in Sect. 7.3.2.

**Lemma 7.20** Let  $A \in \Omega^1(\mathcal{A})$  be a universal one form as in (7.3.2), and  $D' = D(A)$  be given by (7.3.3). Let  $u \in \mathcal{U}(\mathcal{A})$  and  $U = uJuJ^{-1}$ . Then one has

$$UD(A)U^* = D(\gamma_u(A)), \quad \gamma_u(A) = u\delta(u^*) + uAu^* \in \Omega^1(\mathcal{A}) \quad (7.3.4)$$

**Proof** Let  $A = \sum_1^n a_j \delta(b_j) \in \Omega^1(\mathcal{A})$ , one has

$$\gamma_u(A) = u(1 - \sum_1^n a_j b_j)\delta(u^*) + \sum_1^n ua_j \delta(b_j u^*) = \sum_0^n a'_j \delta(b'_j)$$

where  $a'_0 = u(1 - \sum_1^n a_j b_j)$  and  $b'_0 = u^*$ , while  $a'_j = ua_j$  and  $b'_j = b_j u^*$  for  $j > 0$ . What matters is the following, valid for any inclusion  $\mathcal{A} \subset \mathcal{B}$ , and  $T \in \mathcal{B}$

$$\sum_0^n a'_j [T, b'_j] = u[T, u^*] + u \left( \sum_1^n a_j [T, b_j] \right) u^* \quad (7.3.5)$$

We use the notation  $\widehat{T} = JTJ^{-1}$  for any operator  $T$  in  $\mathcal{H}$ , so that

$$\begin{aligned} \omega_{(1)} &:= \sum_j a_j [D, b_j]; \\ \omega_{(2)} &:= \sum_j \hat{a}_j [\omega_{(1)}, \hat{b}_j] \\ &= \sum_{j,k} \hat{a}_j [a_k [D, b_k], \hat{b}_j] \end{aligned}$$

We now apply these formulas using  $\gamma_u(A) = \sum_0^n a'_j \delta(b'_j)$  and obtain using (7.3.5),

$$\omega'_{(1)} = u[D, u^*] + u \left( \sum_1^n a_j [D, b_j] \right) u^* = u[D, u^*] + u\omega_{(1)}u^* \quad (7.3.6)$$

and

$$\omega'_{(2)} = \sum_j \hat{a}'_j [\omega'_{(1)}, \hat{b}'_j] = \hat{u}[\omega'_{(1)}, \hat{u}^*] + \hat{u} \left( \sum_j \hat{a}_j [\omega'_{(1)}, \hat{b}_j] \right) \hat{u}^* \quad (7.3.7)$$

So, using (7.3.6), we get (assuming to simplify that  $\epsilon = 1$  so  $\hat{D} = D$ )

$$\sum_j \hat{a}_j [\omega'_{(1)}, \hat{b}_j] = \sum_j \hat{a}_j [u[D, u^*], \hat{b}_j] + \sum_j \hat{a}_j [u\omega_{(1)}u^*, \hat{b}_j]$$

and the commutation of the  $\hat{x}$  with the  $y$ , for  $x, y \in \mathcal{A}$  gives

$$\sum_j \hat{a}_j [u\omega_{(1)} u^*, \hat{b}_j] = u \left( \sum_j \hat{a}_j [\omega_{(1)}, \hat{b}_j] \right) u^* = u\omega_{(2)} u^*$$

and using  $u[D, u^*] = uDu^* - D$ ,

$$\sum_j \hat{a}_j [u[D, u^*], \hat{b}_j] = u \left( \sum_j \hat{a}_j [D, \hat{b}_j] \right) u^* - \sum_j \hat{a}_j [D, \hat{b}_j] = u\hat{\omega}_{(1)} u^* - \hat{\omega}_{(1)}$$

so that we get:

$$\hat{u} \left( \sum_j \hat{a}_j [\omega'_{(1)}, \hat{b}_j] \right) \hat{u}^* = \hat{u}u\hat{\omega}_{(1)} u^* \hat{u}^* - \hat{u}\hat{\omega}_{(1)} \hat{u}^* + \hat{u}u\omega_{(2)} u^* \hat{u}^* \quad (7.3.8)$$

Next one has

$$\begin{aligned} \hat{u}[\omega'_{(1)}, \hat{u}^*] &= \hat{u}[u[D, u^*], \hat{u}^*] + \hat{u}[u\omega_{(1)} u^*, \hat{u}^*] \\ &= \hat{u}[u[D, u^*], \hat{u}^*] + \hat{u}u\omega_{(1)} u^* \hat{u}^* - u\omega_{(1)} u^* \end{aligned}$$

so that, using (7.3.7) we obtain

$$\omega'_{(2)} = \hat{u}[u[D, u^*], \hat{u}^*] + U\omega_{(1)} U^* - u\omega_{(1)} u^* + U\hat{\omega}_{(1)} U^* - \hat{u}\hat{\omega}_{(1)} \hat{u}^* + U\omega_{(2)} U^* \quad (7.3.9)$$

We then obtain

$$\begin{aligned} \omega'_{(1)} + \hat{\omega}'_{(1)} + \omega'_{(2)} &= u[D, u^*] + \hat{u}[D, \hat{u}^*] + \hat{u}[u[D, u^*], \hat{u}^*] \\ &\quad + U\omega_{(1)} U^* + U\hat{\omega}_{(1)} U^* + U\omega_{(2)} U^* \end{aligned}$$

and the result follows using

$$UDU^* = D + u[D, u^*] + \hat{u}[D, \hat{u}^*] + \hat{u}[u[D, u^*], \hat{u}^*]. \quad \square$$

### 7.3.2 The Semi-group of Inner Perturbations

We show that inner fluctuations come from the action on operators in Hilbert space of a semi-group  $\text{Pert}(\mathcal{A})$  of *inner perturbations* which only depends on the involutive algebra  $\mathcal{A}$  and extends the unitary group of  $\mathcal{A}$ . This covers both cases of ordinary spectral triples and real spectral triples (i.e. those which are equipped with the operator  $J$ ). In the latter case one simply uses the natural homomorphism of semi-groups  $\mu : \text{Pert}(\mathcal{A}) \rightarrow \text{Pert}(\mathcal{A} \otimes \hat{\mathcal{A}})$  given by  $\mu(A) = A \otimes \hat{A}$ . This implies in particular that

inner fluctuations of inner fluctuations are still inner fluctuations and that the corresponding algebraic rules are unchanged by passing from ordinary spectral triples to real spectral triples.

We first show that the formulas of the previous sections can be greatly simplified by representing the universal 1-forms as follows, where  $\mathcal{A}^\circ$  denotes the opposite algebra of  $\mathcal{A}$  and  $x \mapsto x^\circ$  the canonical anti-isomorphism  $\mathcal{A} \mapsto \mathcal{A}^\circ$ ,

**Lemma 7.21** (i) *The following map  $\eta$  is a surjection*

$$\eta : \{\sum a_j \otimes b_j^\circ \in \mathcal{A} \otimes \mathcal{A}^\circ \mid \sum a_j b_j = 1\} \rightarrow \Omega^1(\mathcal{A}), \quad \eta(\sum a_j \otimes b_j^\circ) = \sum a_j \delta(b_j).$$

(ii) *One has*

$$\eta \left( \sum b_j^* \otimes a_j^{*\circ} \right) = \left( \eta \left( \sum a_j \otimes b_j^\circ \right) \right)^*$$

(iii) *One has, for any unitary  $u \in \mathcal{A}$ ,*

$$\eta \left( \sum u a_j \otimes (b_j u^*)^\circ \right) = \gamma_u \left( \eta \left( \sum a_j \otimes b_j^\circ \right) \right)$$

where  $\gamma_u$  is the gauge transformation of potentials.

**Proof** (i) Let us start from an element  $\omega = \sum x_i \delta(y_i) \in \Omega^1(\mathcal{A})$ . Then since  $\delta(1) = 0$  it is the same as

$$(1 - \sum x_i y_i) \delta(1) + \sum x_i \delta(y_i)$$

and one checks that the normalization condition is now fulfilled.

(ii) The normalization condition is fulfilled by  $\sum b_j^* \otimes a_j^{*\circ}$  since  $\sum b_j^* a_j^* = (\sum a_j b_j)^*$ . Thus one gets the equality using  $\delta(x)^* = -\delta(x^*)$  and

$$\sum b_j^* \delta(a_j^*) = - \left( \sum \delta(a_j) b_j \right)^* = \left( \sum a_j \delta(b_j) \right)^*$$

(iii) The normalization condition is fulfilled by  $\sum u a_j \otimes (b_j u^*)^{*\circ}$  since  $\sum u a_j b_j u^* = 1$ . Moreover one has, using  $\delta(b_j u^*) = \delta(b_j) u^* + b_j \delta(u^*)$

$$\sum u a_j \delta(b_j u^*) = u \left( \sum a_j \delta(b_j) \right) u^* + u \delta(u^*)$$

□

**Proposition 7.22** (i) *Let  $A = \sum a_j \otimes b_j^\circ \in \mathcal{A} \otimes \mathcal{A}^\circ$  normalized by the condition  $\sum a_j b_j = 1$ . Then the operator  $D' = D(\eta(A))$  is equal to the inner fluctuation of  $D$  with respect to the algebra  $\mathcal{A} \otimes \hat{\mathcal{A}}$  and the 1-form  $\eta(A \otimes \hat{A})$ , that is*

$$D' = D + \sum a_i \hat{a}_j [D, b_i \hat{b}_j]$$

(ii) An inner fluctuation of an inner fluctuation of  $D$  is still an inner fluctuation of  $D$ , and more precisely one has, with  $A$  and  $A'$  normalized elements of  $\mathcal{A} \otimes \mathcal{A}^\circ$  as above,

$$(D(\eta(A))) (\eta(A')) = D(\eta(A' A))$$

where the product  $A' A$  is taken in the tensor product algebra  $\mathcal{A} \otimes \mathcal{A}^\circ$ .

**Proof** (i) One has, in  $\Omega^1(\mathcal{A} \otimes \hat{\mathcal{A}})$

$$[\delta(b_i), \hat{b}_j] = \delta(b_i \hat{b}_j) - b_i \delta(\hat{b}_j) - \hat{b}_j \delta(b_i)$$

and thus, using the normalization condition and the commutation of  $\mathcal{A}$  with  $\hat{\mathcal{A}}$ ,

$$\sum a_i \hat{a}_j [\delta(b_i), \hat{b}_j] = \sum a_i \hat{a}_j \delta(b_i \hat{b}_j) - \sum a_i \delta(b_i) - \sum \hat{a}_j \delta(\hat{b}_j)$$

Applying this with the derivation  $[D, .]$  instead of  $\delta$  one sees that, in the formula for  $D'$ , the terms in  $\omega_{(1)}$  and  $\hat{\omega}_{(1)}$  combine with  $\omega_{(2)}$  to give the required result.

(ii) We let  $A = \sum a_j \otimes b_j^\circ$  and  $A' = \sum x_s \otimes y_s^\circ$ , both being normalized. We let

$$a_{ij} = a_i \hat{a}_j, \quad b_{ij} = b_i \hat{b}_j, \quad x_{st} = x_s \hat{x}_t, \quad y_{st} = y_s \hat{y}_t$$

and we have

$$D' = D(\eta(A)) = D + \sum a_{ij} [D, b_{ij}]$$

and similarly

$$D'' = D'(\eta(A')) = (D(\eta(A))) (\eta(A')) = D(\eta(A)) + \sum x_{st} [D(\eta(A)), y_{st}]$$

which gives

$$D'' = D + \sum a_{ij} [D, b_{ij}] + \sum x_{st} [D, y_{st}] + \sum \sum x_{st} [a_{ij} [D, b_{ij}], y_{st}]$$

Now one has

$$x_{st} [a_{ij} [D, b_{ij}], y_{st}] = x_{st} (a_{ij} [D, b_{ij}] y_{st} - y_{st} a_{ij} [D, b_{ij}])$$

and the terms on the right sum up to

$$- \sum \sum x_{st} y_{st} a_{ij} [D, b_{ij}] = - \sum a_{ij} [D, b_{ij}]$$

Moreover one has

$$x_{st} a_{ij} [D, b_{ij}] y_{st} = x_{st} a_{ij} [D, b_{ij} y_{st}] - x_{st} a_{ij} b_{ij} [D, y_{st}]$$

and the terms on the right sum up to

$$-\sum \sum x_{st} a_{ij} b_{ij} [D, y_{st}] = -\sum x_{st} [D, y_{st}]$$

Thus we have shown that

$$D'' = D + \sum x_{st} a_{ij} [D, b_{ij} y_{st}]$$

which gives the required result using

$$x_{st} a_{ij} = x_s \hat{x}_t a_i \hat{a}_j = x_s a_i \hat{x}_t \hat{a}_j = x_s a_i \widehat{(x_t a_j)}$$

$$b_{ij} y_{st} = b_i \hat{b}_j y_s \hat{y}_t = b_i y_s \hat{b}_j \hat{y}_t = b_i y_s \widehat{(b_j y_t)}$$

and

$$\left( \sum x_s \otimes y_s^\circ \right) \left( \sum a_i \otimes b_i^\circ \right) = \sum x_s a_i \otimes (b_i y_s)^\circ$$

taking place in the algebra  $\mathcal{A} \otimes \mathcal{A}^\circ$ .  $\square$

Note that the normalization and self-adjointness conditions are preserved by the product of normalized elements in  $\mathcal{A} \otimes \mathcal{A}^\circ$ , since

$$\sum x_s a_i b_i y_s = \sum x_s y_s = 1$$

and moreover the following operation is an antilinear automorphism of  $\mathcal{A} \otimes \mathcal{A}^\circ$

$$\sum a_j \otimes b_j^\circ \mapsto \sum b_j^* \otimes a_j^{*\circ}$$

while the self-adjointness condition means to be in the fixed points of this automorphism. It is thus natural to introduce the following semi-group:

**Proposition 7.23** (i) *The self-adjoint normalized elements of  $\mathcal{A} \otimes \mathcal{A}^\circ$  form a semi-group  $\text{Pert}(\mathcal{A})$  under multiplication.*

(ii) *The transitivity of inner fluctuations (i.e. the fact that inner fluctuations of inner fluctuations are inner fluctuations) corresponds to the semi-group law in the semi-group  $\text{Pert}(\mathcal{A})$ .*

(iii) *The semi-group  $\text{Pert}(\mathcal{A})$  acts on real spectral triples through the homomorphism  $\mu : \text{Pert}(\mathcal{A}) \rightarrow \text{Pert}(\mathcal{A} \otimes \hat{\mathcal{A}})$  given by*

$$A \in \mathcal{A} \otimes \mathcal{A}^\circ \mapsto \mu(A) = A \otimes \hat{A} \in (\mathcal{A} \otimes \hat{\mathcal{A}}) \otimes (\mathcal{A} \otimes \hat{\mathcal{A}})^\circ \quad (7.3.10)$$

**Proof** We have shown above that  $\text{Pert}(\mathcal{A})$  is a semi-group. Using its action on operators in  $\mathcal{H}$  by  $T \mapsto \sum a_i T b_i$  one gets (ii). Proposition 7.22 gives (iii). One checks the multiplicativity of the map  $\mu$  as follows. Let  $A = \sum a_j \otimes b_j^\circ$ ,  $A' = \sum x_s \otimes y_s^\circ$ ,  $a_{ij} = a_i \hat{a}_j$ ,  $b_{ij} = b_i \hat{b}_j$ ,  $x_{st} = x_s \hat{x}_t$ ,  $y_{st} = y_s \hat{y}_t$  so that

$$\mu(A) = \sum a_{ij} \otimes b_{ij}^\circ, \quad \mu(A') = \sum x_{st} \otimes y_{st}^\circ$$

Then one has  $A'A = \sum x_s a_i \otimes (b_i y_s)^\circ$  and

$$\mu(A'A) = \sum x_s a_i \widehat{(x_t a_j)} \otimes \left( b_i y_s \widehat{(b_j y_t)} \right)^\circ = \sum x_{st} a_{ij} \otimes (b_{ij} y_{st})^\circ = \mu(A')\mu(A)$$

which completes the proof of (iii).  $\square$

Note that as a subset of  $\mathcal{A} \otimes \mathcal{A}^\circ$  the subset  $\text{Pert}(\mathcal{A})$  is stable under affine combinations  $\alpha A + \beta A'$  for  $\alpha, \beta \in \mathbb{R}$  and  $\alpha + \beta = 1$ . The map  $\mu$  is quadratic.

To summarize the above discussion we see that the inner fluctuations come from the action of the semi-group  $\text{Pert}(\mathcal{A})$  in a way which parallels the action of inner automorphisms and which, for real spectral triples, combines  $\mathcal{A}$  with  $\hat{\mathcal{A}}$ . Passing from the ordinary formalism of inner fluctuations for spectral triples to the case of real spectral triples is given by the homomorphism  $\mu : \text{Pert}(\mathcal{A}) \rightarrow \text{Pert}(\mathcal{A} \otimes \hat{\mathcal{A}})$  on the semi-groups of inner perturbations. The unitary group  $\mathcal{U}(\mathcal{A})$  maps to the semi-group  $\text{Pert}(\mathcal{A})$  by the homomorphism  $u \in \mathcal{U}(\mathcal{A}) \mapsto u \otimes (u^*)^\circ \in \text{Pert}(\mathcal{A})$ , and this homomorphism is compatible with  $\mu$ .

We end this section by determining the perturbation semi-group of the direct sum of  $*$ -algebras.

**Proposition 7.24** *Let  $\mathcal{A}, \mathcal{B}$  be  $*$ -algebras, then*

$$\text{Pert}(\mathcal{A} \oplus \mathcal{B}) \cong \text{Pert}(\mathcal{A}) \times \text{Pert}(\mathcal{B}) \times (\mathcal{A} \otimes \mathcal{B}^\circ \oplus \mathcal{B} \otimes \mathcal{A}^\circ)^{\text{sa}} \quad (7.3.11)$$

where sa stands for self-adjoint elements, i.e. those of the form  $\sum a_i \otimes b_i^\circ + b_i^* \otimes a_i^{*\circ}$ .

**Proof** We start with the following isomorphism of  $*$ -algebras:

$$(\mathcal{A} \oplus \mathcal{B}) \otimes (\mathcal{A} \oplus \mathcal{B})^\circ \cong \mathcal{A} \otimes \mathcal{A}^\circ \oplus \mathcal{B} \otimes \mathcal{B}^\circ \oplus \mathcal{A} \otimes \mathcal{B}^\circ \oplus \mathcal{B} \otimes \mathcal{A}^\circ.$$

Imposing the normalization and self-adjointness condition to obtain  $\text{Pert}(\mathcal{A} \oplus \mathcal{B})$  on the left-hand side translates on the right-hand side to give  $\text{Pert}(\mathcal{A}) \times \text{Pert}(\mathcal{B}) \times (\mathcal{A} \otimes \mathcal{B}^\circ \oplus \mathcal{B} \otimes \mathcal{A}^\circ)^{\text{sa}}$ . Indeed, normalization only affects the first two terms  $\mathcal{A} \otimes \mathcal{A}^\circ \oplus \mathcal{B} \otimes \mathcal{B}^\circ$  where, together with the self-adjointness condition it gives rise to  $\text{Pert}(\mathcal{A}) \times \text{Pert}(\mathcal{B})$ . The self-adjointness condition on  $\mathcal{A} \otimes \mathcal{B}^\circ \oplus \mathcal{B} \otimes \mathcal{A}^\circ$  gives rise to elements of the form stated above.

### 7.3.2.1 Examples of the Perturbation Semi-group

For commutative matrix algebras we have

**Proposition 7.25** *For any  $N \geq 1$  we have*

$$\text{Pert}(\mathbb{C}^N) \cong \mathbb{C}^{N(N-1)/2}$$

*with the semi-group structure given by componentwise multiplication.*

**Proof** Since  $\text{Pert}(\mathbb{C}) = \{1\}$ , Proposition 7.24 implies that

$$\text{Pert}(\mathbb{C}^N) \cong \text{Pert}(\mathbb{C}^{N-1}) \times \mathbb{C}^{N-1},$$

from which the proof follows.  $\square$

As a next example we determine the perturbation semi-group of  $M_2(\mathbb{C})$ .

Note that we have four basis elements for which the normalization condition becomes

$$(C_{11,11} + C_{12,21})e_{11} + (C_{11,12} + C_{12,22})e_{12} \\ + (C_{21,11} + C_{22,21})e_{21} + (C_{21,12} + C_{22,22})e_{22} = e_{11} + e_{22}.$$

This amounts to the conditions

$$C_{11,11} + C_{12,21} = 1, \quad C_{21,12} + C_{22,22} = 1, \\ C_{11,12} + C_{12,22} = 0, \quad C_{21,11} + C_{22,21} = 0.$$

The self-adjointness condition reads  $C_{ij,kl} = \overline{C_{lk,ji}}$ .

Using the transpose map we may identify

$$M_2(\mathbb{C}) \otimes M_2(\mathbb{C})^\circ \rightarrow M_4(\mathbb{C}), \quad e_{ij} \otimes e_{kl}^\circ \mapsto e_{ij} \otimes e_{lk},$$

The normalization and self-adjointness conditions on  $C_{ij,kl}$  translate to  $4 \times 4$ -matrices to arrive at the following general form for an element  $A \in \text{Pert}(M_2(\mathbb{C}))$ :

$$A = \begin{pmatrix} x_1 & z_3 & \overline{z_3} & 1 - x_1 \\ z_1 & z_2 & \overline{z_5} & -z_1 \\ \overline{z_1} & z_5 & \overline{z_2} & -\overline{z_1} \\ x_2 & z_4 & \overline{z_4} & 1 - x_2 \end{pmatrix}, \quad z_1, \dots, z_5 \in \mathbb{C}, x_1, x_2 \in \mathbb{R}. \quad (7.3.12)$$

The semi-group law ensures that the product of two such matrices again has this general form, something which is not immediately clear. Let us make this point more transparent and establish conditions on  $4 \times 4$  matrices that give rise to the above form.

For an element  $A \in M_4(\mathbb{C})$  to be of the form (7.3.12) is equivalent to demanding that

$$A(f_1 + f_4) = (f_1 + f_4),$$

$$\widehat{\Omega} \overline{A} = A \widehat{\Omega}, \quad \text{where } \widehat{\Omega} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

in terms of the standard basis  $\{f_i\}$  for  $\mathbb{C}^4$ . Equivalently, the matrix  $\widehat{\Omega}$  can be rewritten as a block matrix

$$\widehat{\Omega} = \begin{pmatrix} e_{11} & e_{21} \\ e_{12} & e_{22} \end{pmatrix} = \begin{pmatrix} e_{11}^\top & e_{12}^\top \\ e_{21}^\top & e_{22}^\top \end{pmatrix} = \sum_{i,j} e_{ij} \otimes e_{ji}.$$

Especially the last identity is useful, since we see that the eigenvectors of  $\widehat{\Omega}$  are given by  $e_1 \otimes e_1 \pm e_2 \otimes e_2$ , with eigenvalue 1, and  $e_1 \otimes e_2 \pm e_2 \otimes e_1$ , with eigenvalue 1 and  $-1$  depending on the  $+$  or  $-$  sign. Hence, upon changing to the basis

$$\{e_1 \otimes e_1 + e_2 \otimes e_2, e_1 \otimes e_1 - e_2 \otimes e_2, e_1 \otimes e_2 + e_2 \otimes e_1, e_1 \otimes e_2 - e_2 \otimes e_1\} \quad (7.3.13)$$

of eigenvectors we will get

$$\Omega = \begin{pmatrix} I_3 & 0 \\ 0 & -1 \end{pmatrix}. \quad (7.3.14)$$

Moreover, the vector  $f_1 + f_4$  which is left invariant by  $A$  is given by  $e_1 \otimes e_1 + e_2 \otimes e_2 \in \mathbb{C}^2 \otimes \mathbb{C}^2$ , which is also an eigenvector of  $\widehat{\Omega}$ . Hence with respect to the basis (7.3.13) we arrive at the following characterization of  $\text{Pert}(M_2(\mathbb{C}))$ :

### Proposition 7.26

$$\text{Pert}(M_2(\mathbb{C})) \cong \left\{ A \in M_4(\mathbb{C}) \mid A\omega = \omega, \ \Omega \overline{A} = A\Omega \right\},$$

with

$$\omega = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad \Omega = \begin{pmatrix} I_3 & 0 \\ 0 & -1 \end{pmatrix}.$$

This analysis extends to arbitrary matrix algebras, see Note 9 in this chapter.

## Notes

### Section 7.1. ‘Inner’ unitary equivalences as the gauge group

1. The interpretation of the inner automorphism group as the gauge group is presented in [1].
2. For a precise proof of the isomorphism between  $\text{Aut}(C(X))$  and the group of homeomorphisms of  $X$ , we refer to [2, Theorem II.2.2.6]. For a more detailed treatment of the smooth analogue, we refer to [3, Sect. 1.3].

### Section 7.2. Morita self-equivalences as gauge fields

3. The gauge group  $\mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$  introduced in Definition 7.4 (following [1, 4, 5]) is a natural lift of the group of inner automorphisms of the algebra  $\mathcal{A}$ , as is proved in Proposition 7.5. Another approach to lifting  $\text{Inn}(\mathcal{A})$  to be represented on  $\mathcal{H}$  is by *central extensions*; this is described in [6].
4. For unital algebras algebraic Morita equivalence [7] coincides with Rieffel’s notion of strong Morita equivalence for  $C^*$ -algebras [8]. This is proved in [9] and explains why we can safely work with algebraic tensor products. We also refer for a more general treatment to *e.g.* [3, Sect. 4.5] and [10, Sects. A.3 and A.4].
5. Besides Morita equivalence, also the more general notion of KK-equivalence can be lifted to spectral triples, but this requires much more analysis [11–13].
6. Theorem 7.15 and Theorem 7.16 are due to Connes in [1].

### Section 7.3. Inner fluctuations without the first-order condition

7. Universal differential forms are defined in terms of the graded differential algebra  $\Omega^\bullet(\mathcal{A})$  that is freely generated by  $a$  and  $\delta b$  for any  $a, b \in \mathcal{A}$ . In other words, a universal differential  $n$ -form  $\eta$  is given by an expression of the form

$$\eta = \sum_j a_0^j \delta a_1^j \cdots \delta a_n^j,$$

and its differential is

$$\delta\eta = \sum_j \delta a_0^j \delta a_1^j \cdots \delta a_n^j.$$

There will be no commutation relations imposed between the  $a_0, \delta a_0$  and the  $\delta a_j$ ’s, but we do have the Leibniz rule stating that

$$\delta(ab) = \delta(a)b + a\delta b$$

for any  $a, b \in \mathcal{A}$ . For an overview on universal differential forms, we refer to [10, Sect. 7.1].

8. Sect. 7.3 is based on [14]. That paper also contains a proof of Proposition 7.18.
9. Sect. 7.3.2.1 is based on [15]. This contains a description of the perturbation semi-group for all (real and complex) matrix algebras.

## References

1. Connes, A.: Gravity coupled with matter and the foundation of non-commutative geometry. *Commun. Math. Phys.* **182**, 155–176 (1996)
2. Blackadar, B.: Operator algebras, Encyclopaedia of Mathematical Sciences, vol. 122. Springer, Berlin. Theory of  $C^*$ -algebras and von Neumann algebras. Operator Algebras and Non-commutative Geometry, III (2006)
3. Gracia-Bondía, J.M., Várilly, J.C., Figueroa, H.: Elements of Noncommutative Geometry. Birkhäuser, Boston (2001)
4. Connes, A., Marcolli, M.: Noncommutative Geometry Quantum Fields and Motives. AMS, Providence (2008)
5. van den Dungen, K., van Suijlekom, W.D.: Particle physics from almost commutative space-times. *Rev. Math. Phys.* **24**, 1230004 (2012)
6. Lazzarini, S., Schucker, T.: A Farewell to unimodularity. *Phys. Lett. B* **510**, 277–284 (2001)
7. Morita, K.: Duality for modules and its applications to the theory of rings with minimum condition. *Sci. Rep. Tokyo Kyoiku Daigaku Sect. A* **6**, 83–142 (1958)
8. Rieffel, M.A.: Morita equivalence for  $C^*$ -algebras and  $W^*$ -algebras. *J. Pure Appl. Algebra* **5**, 51–96 (1974)
9. Beer, W.: On Morita equivalence of nuclear  $C^*$ -algebras. *J. Pure Appl. Algebra* **26**, 249–267 (1982)
10. Landi, G.: An Introduction to Noncommutative Spaces and their Geometry. Springer (1997)
11. Baaj, S., Julg, P.: Théorie bivariante de Kasparov et opérateurs non bornés dans les  $C^*$ -modules hilbertiens. *C. R. Acad. Sci. Paris Sér. I Math.* **296**, 875–878 (1983)
12. Kucerovsky, D.: The  $KK$ -product of unbounded modules. *K-Theory* **11**, 17–34 (1997)
13. Mesland, B.: Unbounded bivariant  $K$ -theory and correspondences in noncommutative geometry. *J. Reine Angew. Math.* **691**, 101–172 (2014)
14. Chamseddine, A.H., Connes, A., Van Suijlekom, W.D.: Inner fluctuations in noncommutative geometry without the first order condition. *J. Geom. Phys.* **73**, 222–234 (2013)
15. Neumann, N., van Suijlekom, W.D.: Perturbation semigroup of matrix algebras. *J. Noncommut. Geom.* **10**, 245–264 (2016)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 8

## Localization of Gauge Theories from Noncommutative Geometry



We ‘localize’ the generalized gauge theory derived from any spectral triple by constructing a  $C^*$ -bundle on which the gauge group acts by vertical automorphisms. This will be exemplified for toric noncommutative manifolds.

### 8.1 Commutative Subalgebra and $C^*$ -Bundles

Given a real spectral triple  $(\mathcal{A}, \mathcal{H}, D; J)$  we can construct a spectral triple on some commutative subalgebra of  $\mathcal{A}$ , derived from this data. Indeed, set

$$\mathcal{A}_J := \{a \in \mathcal{A} : aJ = Ja^*\}.$$

As we will see shortly, this is a complex subalgebra, contained in the center of  $\mathcal{A}$  (and hence commutative). Later, in Chap. 10, this subalgebra will turn out to be very useful in the description of the gauge group associated to any real spectral triple.

**Proposition 8.1** *Let  $(\mathcal{A}, \mathcal{H}, D; J)$  be a real spectral triple. Then*

- (1)  *$\mathcal{A}_J$  defines an involutive commutative complex subalgebra of the center of  $\mathcal{A}$ .*
- (2)  *$(\mathcal{A}_J, \mathcal{H}, D; J)$  is a real spectral triple.*
- (3) *Any  $a \in \mathcal{A}_J$  commutes with the algebra generated by the sums  $\sum_j a_j [D, b_j] \in \Omega_D^1(\mathcal{A})$  with  $a_j, b_j \in \mathcal{A}$ .*

**Proof** (1) If  $a \in \mathcal{A}_J$  then also  $Ja^*J^{-1} = (JaJ^{-1})^* = a$ , since  $J$  is isometric. Hence,  $\mathcal{A}_J$  is involutive. Moreover, for all  $a \in \mathcal{A}_J$  and  $b \in \mathcal{A}$  we have  $[a, b] = [Ja^*J^{-1}, b] = 0$  by the commutant property (5.2.1). Thus,  $\mathcal{A}_J$  is in the center of  $\mathcal{A}$ .

(2) Since  $\mathcal{A}_J$  is a subalgebra of  $\mathcal{A}$ , all conditions for a spectral triple are automatically satisfied.

(3) This follows from the order-one condition (5.2.1):

$$[a, [D, b]] = [Ja^*J^{-1}, [D, b]] = 0,$$

for  $a \in \mathcal{A}_J$  and  $b \in \mathcal{A}$ .  $\square$

**Example 8.2** In the case of a Riemannian spin manifold  $M$  with real structure  $J_M$  given by charge conjugation, one checks that

$$C^\infty(M)_{J_M} = C^\infty(M, \mathbb{R}).$$

More generally, under suitable conditions on the triple  $(\mathcal{A}, \mathcal{H}, D)$  the spectral triple  $(\mathcal{A}_J, \mathcal{H}, D)$  is a so-called commutative spin geometry. Then, Connes' Reconstruction Theorem (*cf.* Note 6 on Chap. 5) establishes the existence of a compact Riemannian spin manifold  $M$  such that there is an isomorphism  $(\mathcal{A}_J, \mathcal{H}, D) \cong (C^\infty(M), L^2(S \otimes E), D_E)$ . The spinor bundle  $S \rightarrow M$  is twisted by a vector bundle  $E \rightarrow M$  and the twisted Dirac operator is of the form  $D_E = D + \rho$  with  $\rho \in \Gamma^\infty(\text{End}(S \otimes E))$ .

In any case, as  $\mathcal{A}_J$  is commutative, Gelfand duality (Theorem 5.7) ensures the existence of a compact Hausdorff space such that  $\mathcal{A}_J \subset C(X)$  as a dense  $*$ -subalgebra. Indeed, the  $C^*$ -completion of  $\mathcal{A}_J$  in  $\mathcal{B}(\mathcal{H})$  is commutative and hence isomorphic to such a  $C(X)$ . We consider this space  $X$  to be the ‘background space’ on which  $(\mathcal{A}, \mathcal{H}, D; J, \gamma)$  describes a gauge theory, as we now work out in detail.

Heuristically speaking, the gauge group  $\mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$  introduced in Definition 7.4 considers only transformations that are ‘vertical’, or ‘purely non-commutative’ with respect to  $X$ , quotienting out the unitary transformations of the commutative subalgebra  $\mathcal{A}_J$ . In this chapter we will make this more precise by identifying a bundle  $\mathfrak{B} \rightarrow X$  of  $C^*$ -algebras such that:

- the space of continuous sections  $\Gamma(X, \mathfrak{B})$  forms a  $C^*$ -algebra isomorphic to  $A = \overline{\mathcal{A}}$ , the  $C^*$ -completion of  $\mathcal{A}$ ;
- the gauge group acts as bundle automorphisms covering the identity.

Moreover, we search for a bundle of  $C^*$ -algebras of which the gauge fields  $\omega \in \Omega_D^1(\mathcal{A})$  are sections and on which the gauge group again acts by bundle automorphisms.

We avoid technical complications that might arise from working with dense subalgebras of  $C^*$ -algebras, and work with the  $C^*$ -algebras  $\mathcal{A}_J$  and  $A$  themselves, as completions of  $\mathcal{A}_J$  and  $\mathcal{A}$ , respectively. First, note that there is an inclusion map  $C(X) \simeq \mathcal{A}_J \hookrightarrow A$ . This means that  $A$  is a so-called  $C(X)$ -algebra, which by definition is a  $C^*$ -algebra  $A$  with a map from  $C(X)$  to the center of  $A$ . Indeed, it follows from Proposition 8.1 that  $\mathcal{A}_J$  is contained in the center of  $A$ .

In such a case  $A$  is the  $C^*$ -algebra of continuous sections of an upper semi-continuous  $C^*$ -bundle over  $X$ . We will briefly sketch the setup (see Note 3 in this chapter). Recall that a function  $f : A \rightarrow \mathbb{C}$  is **upper semi-continuous** at  $a_0 \in A$  if  $\limsup_{a \rightarrow a_0} \|f(a)\| \leq \|f(a_0)\|$ .

**Definition 8.3** An *upper semi-continuous  $C^*$ -bundle* over a compact topological space  $X$  is a continuous, open, surjection  $\pi : \mathfrak{B} \rightarrow X$  together with operations and norms that turn each fiber  $\mathfrak{B}_x = \pi^{-1}(x)$  into a  $C^*$ -algebra, such that (1) the map  $a \mapsto \|a\|$  is upper semi-continuous, (2) all algebraic operations are continuous on  $\mathfrak{B}$ , (3) if  $\{a_i\}$  is a net in  $\mathfrak{B}$  such that  $\|a_i\| \rightarrow 0$  and  $\pi(a_i) \rightarrow x$  in  $X$ , then  $a_i \rightarrow 0_x$ , where  $0_x$  is the zero element in  $\mathfrak{B}_x$ .

A (continuous) *section* of  $\mathfrak{B}$  is a (continuous) map  $s : X \rightarrow \mathfrak{B}$  such that  $\pi(s(x)) = x$ .

A base for the topology on  $\mathfrak{B}$  is given by the following collection of open sets:

$$W(s, \mathcal{O}, \epsilon) := \{b \in \mathfrak{B} : \pi(b) \in \mathcal{O} \text{ and } \|b - s(\pi(b))\| < \epsilon\}, \quad (8.1.1)$$

indexed by continuous sections  $s \in \Gamma(X, \mathfrak{B})$ , open subsets  $\mathcal{O} \subset X$  and  $\epsilon > 0$ .

**Proposition 8.4** *The space  $\Gamma(X, \mathfrak{B})$  of continuous sections forms a  $C^*$ -algebra when it is equipped with the norm*

$$\|s\| := \sup_{x \in X} \|s(x)\|_{\mathfrak{B}_x}.$$

**Proof** See Note 3 in this chapter. □

In our case, after identifying  $C(X)$  with  $A_J$ , we can define a closed two-sided ideal in  $A$  by

$$I_x := \{fa : a \in A, f \in C(X), f(x) = 0\}^- . \quad (8.1.2)$$

We think of the quotient  $C^*$ -algebra  $\mathfrak{B}_x := A/I_x$  as the fiber of  $A$  over  $x$  and set

$$\mathfrak{B} := \coprod_{x \in X} \mathfrak{B}_x, \quad (8.1.3)$$

with an obvious surjective map  $\pi : \mathfrak{B} \rightarrow X$ . If  $a \in A$ , then we write  $a(x)$  for the image  $a + I_x$  of  $a$  in  $\mathfrak{B}_x$ , and we think of  $a$  as a section of  $\mathfrak{B}$ . The fact that all these sections are continuous and that elements in  $A$  can be obtained in this way is guaranteed by the following result.

**Theorem 8.5** *The above map  $\pi : \mathfrak{B} \rightarrow X$  with  $\mathfrak{B}$  as in (8.1.3) defines an upper semi-continuous  $C^*$ -bundle over  $X$ . Moreover, there is a  $C(X)$ -linear isomorphism of  $A$  onto  $\Gamma(X, \mathfrak{B})$ .*

**Proof** See Note 3 in this chapter. □

Having obtained the  $C^*$ -algebra  $A$  as the space of sections of a  $C^*$ -bundle, we are ready to analyze the action of the gauge group on  $A$ . Staying at the  $C^*$ -algebraic level, we consider the **continuous gauge group**

$$\mathfrak{G}(A, \mathcal{H}; J) \simeq \frac{\mathcal{U}(A)}{\mathcal{U}(A_J)}.$$

This contains the gauge group  $\mathfrak{G}(A, \mathcal{H}; J)$  of Definition 7.4 as a dense subgroup in the topology induced by the  $C^*$ -norm on  $A$ . The next result realizes the gauge group as a group of vertical bundle automorphisms of  $\mathfrak{B}$ .

**Proposition 8.6** *The action  $\alpha$  of  $\mathfrak{G}(A, \mathcal{H}; J)$  on  $A$  by inner  $C^*$ -algebra automorphisms induces an action  $\tilde{\alpha}$  of  $\mathfrak{G}(A, \mathcal{H}; J)$  on  $\mathfrak{B}$  by continuous bundle automorphisms that cover the identity. In other words, for  $g \in \mathfrak{G}(A, \mathcal{H}; J)$  we have*

$$\pi(\tilde{\alpha}_g(b)) = \pi(b); \quad (b \in \mathfrak{B}).$$

Moreover, under the identification of Theorem 8.5 the induced action  $\tilde{\alpha}^*$  on  $\Gamma(X, \mathfrak{B})$  given by

$$\tilde{\alpha}_g^*(s)(x) = \tilde{\alpha}_g(s(x))$$

coincides with the action  $\alpha$  on  $A$ .

**Proof** The action  $\alpha$  induces an action on  $A/I_x = \pi^{-1}(x)$ , since  $\alpha_g(I_x) \subset I_x$  for all  $g \in \mathfrak{G}(A, \mathcal{H}; J)$ . We denote the corresponding action of  $\mathfrak{G}(A, \mathcal{H}; J)$  on  $\mathfrak{B}$  by  $\tilde{\alpha}$ , so that, indeed,

$$\pi(\tilde{\alpha}_g(b)) = \pi(b); \quad (b \in \pi^{-1}(x)).$$

Let us also check continuity of this action. In terms of the base  $W(s, \mathcal{O}, \epsilon)$  of (8.1.1), we find that

$$\tilde{\alpha}_g(W(s, \mathcal{O}, \epsilon)) = W(\tilde{\alpha}_g^*(s), \mathcal{O}, \epsilon),$$

mapping open subsets one-to-one and onto open subsets.

For the second claim, it is enough to check that the action  $\tilde{\alpha}^*$  on the section  $s : x \mapsto a + I_x \in \mathfrak{B}_x$ , defined by an element  $a \in A$ , corresponds to the action  $\alpha$  on that  $a$ . In fact,

$$\tilde{\alpha}_g^*(s)(x) = \tilde{\alpha}_g(s(x)) = \alpha_g(a + I_x) = \alpha_g(a) + I_x,$$

which completes the proof.  $\square$

At the infinitesimal level, the derivations in the gauge algebra  $\mathfrak{g}(A, \mathcal{H}; J)$  also act vertically on the  $C^*$ -bundle  $\mathfrak{B}$  defined in (8.1.3), and the induced action on the sections  $\Gamma(X, \mathfrak{B})$  agrees with the action of  $\mathfrak{g}(A, \mathcal{H}; J)$  on  $A$ .

## 8.2 Localization of the Gauge Group

We now investigate whether or when  $\mathfrak{G}(A, \mathcal{H}; J)$  can be considered as the group of continuous sections of a group bundle on the same base space  $X$ . Set-theoretically, one expects the group bundle that corresponds to  $\Gamma(X, \mathfrak{B})$  to be given by

$$\mathfrak{GB} := \coprod_{x \in X} \frac{\mathcal{U}(\mathfrak{B}_x)}{\mathcal{U}(\mathbb{C})}.$$

We define a topology on  $\mathfrak{GB}$  as follows. First, the group bundle

$$\mathcal{U}\mathcal{B} := \coprod_{x \in X} \mathcal{U}(\mathfrak{B}_x)$$

is equipped with the induced topology from  $\mathcal{B}$ . Since each  $\mathfrak{B}_x$  is a complex unital algebra, we have  $\mathcal{U}(\mathbb{C}) \subset \mathcal{U}(\mathfrak{B}_x)$  so that we have a group subbundle  $\coprod_{x \in X} \mathcal{U}(\mathbb{C}) \subset \mathcal{U}\mathfrak{B}$ . We write  $\mathcal{U}\mathbb{C}$  for this group subbundle. The topology of  $\mathfrak{GB}$  is then the quotient topology of the bundle  $\mathcal{U}\mathcal{B}$  by the fiberwise action of the group bundle  $\mathcal{U}\mathbb{C}$ .

Before stating our main result on the structure of the gauge group, we consider the spaces of continuous sections of the group bundles  $\mathcal{U}\mathbb{C}$  and  $\mathcal{U}\mathcal{B}$ .

**Proposition 8.7** *We have the following group isomorphisms:*

$$\begin{aligned}\Gamma(X, \mathcal{U}\mathbb{C}) &\cong \mathcal{U}(A_J), \\ \Gamma(X, \mathcal{U}\mathcal{B}) &\cong \mathcal{U}(A).\end{aligned}$$

**Proof** Firstly, a continuous map from  $X$  to  $\mathcal{U}(\mathbb{C})$  is simply given by a unitary continuous function on  $X$ . Secondly, since  $\Gamma(X, \mathcal{B}) \cong A$ , unitarity translates from the product in  $A$  to the fiberwise product in  $\mathcal{B}$ , hence proving the result.  $\square$

We also need the following well-known result on covering spaces:

**Proposition 8.8** *Suppose given a covering space  $p : (\tilde{Y}, \tilde{y}_0) \rightarrow (Y, y_0)$  and a map  $f : (X, x_0) \rightarrow (Y, y_0)$  with  $X$  path-connected and locally path-connected. Then a lift  $\tilde{f} : (X, x_0) \rightarrow (\tilde{Y}, \tilde{y}_0)$  of  $f$  exists if and only if  $f_*(\pi_1(X, x_0)) \subset p_*(\pi_1(\tilde{Y}, \tilde{y}_0))$ .*

**Theorem 8.9** *If  $X$  is simply connected and if there exists a subbundle  $\tilde{\mathfrak{GB}} \subset \mathcal{U}\mathcal{B}$  that is a covering space of  $\mathfrak{GB}$  (via the quotient map  $\mathcal{U}\mathcal{B} \rightarrow \mathfrak{GB}$ ), then there is the following short exact sequence of groups*

$$1 \longrightarrow \Gamma(X, \mathcal{U}\mathbb{C}) \longrightarrow \Gamma(X, \mathcal{U}\mathcal{B}) \longrightarrow \Gamma(X, \mathfrak{GB}) \longrightarrow 1. \quad (8.2.1)$$

Consequently, in this case the gauge group is given as the space of continuous sections of the group bundle  $\mathfrak{GB}$ , i.e.

$$\mathfrak{G}(A, \mathcal{H}; J) \cong \Gamma(X, \mathfrak{GB}).$$

**Proof** Exactness of (8.2.1) is clear from the very definition of the group bundle  $\mathfrak{GB}$ , except perhaps for the claim of surjectivity of the map  $\Gamma(X, \mathcal{U}\mathcal{B}) \rightarrow \Gamma(X, \mathfrak{GB})$ . This follows from Proposition 8.8, applied to a continuous section  $g \in \Gamma(X, \mathfrak{GB})$ . Indeed, since  $\pi_1(X)$  is trivial, there always exists a lift  $\tilde{g} : X \rightarrow \tilde{\mathfrak{GB}} \subset \mathcal{U}\mathcal{B}$ , thus proving surjectivity.

For the second statement, exactness of the sequence implies that

$$\Gamma(X, \mathfrak{GB}) \cong \frac{\Gamma(X, \mathcal{U}\mathcal{B})}{\Gamma(X, \mathcal{U}\mathbb{C})} \cong \frac{\mathcal{U}(A)}{\mathcal{U}(A_J)}$$

using Proposition 8.7. But this is precisely the definition of the group  $\mathfrak{G}(A, \mathcal{H}; J)$ .  $\square$

This result allows for the following refinement of Proposition 8.6.

**Corollary 8.10** *Under the same conditions as in Theorem 8.9, the action of the gauge group  $\mathfrak{G}(A, \mathcal{H}; J)$  on  $A$  is induced by the action of the fibers  $\mathfrak{GB}_x := \mathcal{U}(\mathcal{B}_x)/\mathcal{U}(\mathbb{C})$  on the fibers  $\mathfrak{B}_x$  of  $\mathfrak{B}$  by inner automorphisms.*

**Proof** Let  $g \in \mathfrak{G}(A, \mathcal{H}; J)$  with pre-image  $u \in \mathcal{U}(A)$ , i.e. so that  $\alpha_g(a) = uau^*$ . Then  $g, u$  and  $a$  can be considered as continuous sections of bundles  $\mathfrak{GB}, \mathcal{U}\mathcal{B}$  and  $\mathcal{B}$  on  $X$ , respectively. At a point  $x \in X$  we have  $g(x) \in \mathfrak{GB}_x = \mathcal{U}(\mathcal{B}_x)/\mathcal{U}(\mathbb{C})$  with pre-image  $u(x) \in \mathcal{U}(\mathcal{B}_x)$  and we compute as sections of  $\mathcal{B} \rightarrow X$ :

$$(\alpha_g(a))(x) = u(x)a(x)u(x)^*,$$

thus establishing the result.  $\square$

Note that Theorem 8.5 also gives a bundle description of  $\text{Inn}(A)$  if  $Z(A) = A_J$ . Indeed, in combination with Corollary 8.10 we find that then  $\text{Inn}(A) \cong \Gamma(X, \mathfrak{GB})$ , realizing the group of inner automorphisms of  $A$  as the space of continuous sections of a group bundle.

### 8.3 Localization of Gauge Fields

Also the gauge fields  $\omega$  that enter as inner fluctuations of  $D$  can be parametrized by sections of some bundle of  $C^*$ -algebras. In order for this to be compatible with the vertical action of the gauge group found above, we will write any connection in the form,

$$\nabla = d + \omega_0 + \omega,$$

where  $d = [D, \cdot]$  and  $\omega_0, \omega \in \Omega_D^1(\mathcal{A})$ . The action of a gauge transformation on  $\nabla$  then induces the following transformation:

$$\omega_0 \mapsto u\omega_0u^* + u[D, u^*]; \quad \omega \mapsto u\omega u^*.$$

The  $C^*$ -algebra generated by  $\mathcal{A}$  and  $[D, \mathcal{A}]$  is a  $C(X)$ -algebra, since  $C(X) \cong A_J$ , which according to Proposition 8.1 commutes with both  $\mathcal{A}$  and  $[D, \mathcal{A}]$ . Thus, a similar construction as in the previous subsection establishes the existence of an upper semi-continuous  $C^*$ -bundle  $\mathfrak{B}_\Omega$  over  $X$ , explicitly given by

$$\mathfrak{B}_\Omega = \prod_{x \in X} C^*(\mathcal{A}, [D, \mathcal{A}]) / I'_x,$$

where  $C^*(\mathcal{A}, [D, \mathcal{A}])$  is the  $C^*$ -algebra generated by  $a$  and  $[D, b]$  for  $a, b \in \mathcal{A}$ , and  $I'_x$  is the two-sided ideal in  $C^*(\mathcal{A}, [D, \mathcal{A}])$  generated by  $I_x$  that has been defined before (see Equation (8.1.2)). Again, one can show that  $\Gamma(X, \mathfrak{B}_\Omega)$  is isomorphic to this  $C^*$ -algebra and establish the following result.

**Proposition 8.11** *Let  $\pi : \mathfrak{B}_\Omega \rightarrow X$  be as above.*

- (1) *The gauge field  $\omega$  defines a continuous section of  $\mathfrak{B}_\Omega$ .*
- (2) *The gauge group  $\mathfrak{G}(A, \mathcal{H}; J)$  acts fiberwise on this bundle, and the induced action on  $\Gamma(X, \mathfrak{B}_\Omega)$  agrees with the action on  $C^*(\mathcal{A}, [D, \mathcal{A}])$ .*

Consequently, if we regard  $\omega \in \Omega_D^1(\mathcal{A})$  as a continuous section  $\omega(x)$  of  $\mathfrak{B}_\Omega$ , an element  $uJuJ^{-1} \in \mathfrak{G}(A, \mathcal{H}; J)$  acts as

$$\omega(x) \mapsto (u\omega u^*)(x) \equiv u\omega(x)u^*.$$

## 8.4 Localization of Toric Noncommutative Manifolds

In Chap. 12 we will see a concrete example of the above localization for Yang–Mills gauge theories, phrased in the language of principal bundles. We will here give another illustrative example, given by toric noncommutative manifolds introduced in Sect. 5.3.2.

Thus, we consider an arbitrary compact Riemannian spin manifold  $M$  that carries a (smooth) action of a 2-torus by isometries. We then have a real spectral triple  $(C^\infty(M_\theta), \mathcal{H}, D_M; J_M)$  so let us determine the gauge theory corresponding to it. We distinguish two cases corresponding to  $\theta$  being rational or irrational. These two cases require completely different techniques and yield entirely different results.

For  $\theta$  rational we have the following result. If  $p, q$  are coprime and  $\theta = p/q$ , we set  $\Gamma_\theta = \mathbb{Z}/q\mathbb{Z}$ .

**Theorem 8.12** *We have the following equivalence of spectral triples:*

$$(C^\infty(M_\theta), L^2(M, \mathcal{S}), D) \cong (\Gamma^\infty(M/\Gamma_\theta, B), L^2(M/\Gamma_\theta, \pi_*\mathcal{S} \otimes B), \pi_*D)$$

in terms of the projection map  $\pi : M \rightarrow M/\Gamma_\theta$  and a  $*$ -algebra bundle  $B := M \times_{\Gamma_\theta} M_q(\mathbb{C})$  with base space  $M/\Gamma_\theta$ , for a suitable action of  $\Gamma_\theta$  on  $M_q(\mathbb{C})$

**Proof** See Note 7 in this chapter. □

Let us relate this to our gauge theory description using the commutative subalgebra  $C(M_\theta)_{J_M}$  in  $C(M)$ .

**Proposition 8.13** *For the real spectral triple  $(C^\infty(M_\theta), \mathcal{H}, D; J_M)$  we have for  $\theta$  rational that*

$$C(M_\theta)_{J_M} = Z(C(M_\theta)).$$

Moreover, in this case  $C(M_\theta)_{J_M} \cong C(M/\Gamma_\theta)$ .

**Proof** First,  $C(M_\theta)_{J_M} \subset Z(C(M_\theta))$  by Proposition 8.1. The converse inclusion is obtained as follows. We have  $Z(C(M_\theta)) \cong C(M/\Gamma_\theta)$  because  $Z(M_q(\mathbb{C})) = \mathbb{C}$  for all fibers. Moreover,  $C(M/\Gamma_\theta) = C(M)^{\Gamma_\theta}$  is a subalgebra of  $C(M)$ , all of whose elements satisfy the commutation relation  $aJ_M = J_M a^*$  (cf. Example 8.2).  $\square$

Hence, the bundle  $B = M \times_{\Gamma_\theta} M_q(\mathbb{C}) \rightarrow M/\Gamma_\theta$  is the sought-for  $C^*$ -bundle on which to define our gauge theory. Theorem 8.12 tells us that the  $C^*$ -algebra  $C(M_\theta)$  is isomorphic to the space of continuous sections of  $B$  —in concordance with our Theorem 8.5—and for the gauge group we actually have the following result:

$$\mathfrak{G}(C(M_\theta), \mathcal{H}; J_M) \cong \Gamma(M/\Gamma_\theta, M \times_{\Gamma_\theta} PU(q)),$$

if  $M/\Gamma_\theta$  is simply connected. In other words, we are considering a  $PU(q)$ -gauge theory, in the usual sense. This Lie group acts on the fiber  $M_q(\mathbb{C})$  of  $B$  in the adjoint representation.

Let us now proceed with the case that  $\theta$  is irrational.

**Proposition 8.14** *For the real spectral triple  $(C^\infty(M_\theta), \mathcal{H}, D; J_M)$  we have for  $\theta$  irrational*

$$C(M_\theta)_{J_M} = Z(C(M_\theta)).$$

Moreover, in this case  $C(M_\theta)_{J_M} \cong C(M/\mathbb{T}^2)$ .

**Proof** First note that  $Z(C(M_\theta)) = C(M_\theta)^{\mathbb{T}^2}$ , essentially because the center of  $A_\theta$  is trivial if  $\theta$  is irrational (see Note 9 below). Moreover, since  $C(M)^{\mathbb{T}^2}$  is unchanged under the deformation, as well as  $J_M$ , we find that  $C(M_\theta)^{\mathbb{T}^2} \cong C(M)^{\mathbb{T}^2}$  is contained in  $C(M_\theta)_{J_M}$  which also proves the second statement.  $\square$

This allows us to conclude with Theorem 8.5 that  $C(M_\theta)$  is isomorphic to the  $C^*$ -algebra  $\Gamma(M/\mathbb{T}^2, \mathfrak{B}^{M_\theta})$  of continuous sections of an upper semi-continuous  $C^*$ -bundle  $\mathfrak{B}^{M_\theta} \rightarrow M/\mathbb{T}^2$  and that  $\mathfrak{G}(C(M_\theta), \mathcal{H}; J_M)$  acts by vertical automorphisms on  $\mathfrak{B}^{M_\theta}$ . In fact, even more can be said in this case.

**Theorem 8.15** *The above  $C^*$ -bundle  $\mathfrak{B}^{M_\theta} \rightarrow M/\mathbb{T}^2$  is a continuous  $C^*$ -bundle. Moreover, its fibers are given by the following  $C^*$ -algebras:*

$$\mathfrak{B}_x^{M_\theta} \cong C(\mathbb{T}^2/\mathbb{T}_x^2, A_\theta)^{\mathbb{T}^2},$$

for  $x \in M/\mathbb{T}^2$  having isotropy group  $\mathbb{T}_x^2 \subseteq \mathbb{T}^2$ .

**Proof** See Note 11 in this chapter.  $\square$

Hence, the spectral triple  $(C(M_\theta), \mathcal{H}, D; J_M)$  yields a gauge theory defined in terms of a  $C^*$ -bundle  $\mathfrak{B}^{M_\theta} \rightarrow M/\mathbb{T}^2$ . The gauge group  $\mathfrak{G}(C(M_\theta), \mathcal{H}; J_M)$  is parametrized by unitaries in  $C(M_\theta)$  and acts vertically on the bundle  $\mathfrak{B}^{M_\theta}$ . We now determine the bundle structure of the gauge group, thereby making use of Theorem 8.9 above.

**Proposition 8.16** *There exists a subbundle  $\tilde{\mathfrak{GB}}^{M_\theta} \subset \mathcal{U}\mathfrak{B}^{M_\theta}$  that is a covering space of  $\mathfrak{GB}^{M_\theta}$  for the quotient map  $\mathcal{U}\mathfrak{B}^{M_\theta} \rightarrow \mathfrak{GB}^{M_\theta}$ . Consequently, if  $M/\mathbb{T}^2$  is simply connected we have*

$$\mathfrak{G}(C(M_\theta), \mathcal{H}; J_M) \cong \Gamma(M/\mathbb{T}^2, \mathfrak{GB}^{M_\theta}),$$

where the fibers of  $\mathfrak{GB}^{M_\theta}$  are given by

$$\mathfrak{GB}_x^{M_\theta} \cong \frac{\mathcal{U}(C(\mathbb{T}^2/\mathbb{T}_x^2, A_\theta)^{\mathbb{T}^2})}{\mathcal{U}(\mathbb{C})}; \quad (x \in M/\mathbb{T}^2).$$

**Proof** From Theorem 8.15 it follows that the fibers  $\mathcal{U}\mathfrak{B}_x^{M_\theta}$  of  $\mathcal{U}\mathfrak{B}^{M_\theta}$  are given by the topological groups  $\mathcal{U}(C(\mathbb{T}^2/\mathbb{T}_x^2, A_\theta)^{\mathbb{T}^2})$ . We define a subbundle of  $\mathcal{U}\mathfrak{B}^{M_\theta}$  using the unique tracial state  $\tau$  on  $A_\theta$ . First, consider the phase map  $\varphi : A_\theta \rightarrow U(1)$  given by

$$\varphi(a) = \frac{\tau(a)}{|\tau(a)|}; \quad (a \in A_\theta).$$

It induces a phase map on the fibers of  $\mathfrak{B}^{M_\theta}$  by composition:

$$\begin{aligned} \tilde{\varphi} : C(\mathbb{T}^2/\mathbb{T}_x^2, A_\theta)^{\mathbb{T}^2} &\rightarrow U(1), \\ f &\mapsto \varphi \circ f. \end{aligned}$$

We then define a subbundle  $\tilde{\mathfrak{GB}}^{M_\theta} \subset \mathcal{U}\mathfrak{B}^{M_\theta}$  by giving its fibers:

$$\tilde{\mathfrak{GB}}^{M_\theta}_x = \{u \in \mathcal{U}\mathfrak{B}^{M_\theta} : \tilde{\varphi}(u) = 1\}.$$

For  $\tilde{\mathfrak{GB}}^{M_\theta}$  to be a covering space of  $\mathfrak{GB}^{M_\theta}$ , we determine the kernel of the quotient map  $\mathcal{U}\mathfrak{B}^{M_\theta} \rightarrow \mathfrak{GB}^{M_\theta}$ , intersected with  $\tilde{\mathfrak{GB}}^{M_\theta}$ . In fact, being in the kernel amounts to  $u \in \mathcal{U}(\mathbb{C})$  so that  $\varphi(u) = 1$  implies that then  $u = 1$ . Hence,  $\tilde{\mathfrak{GB}}^{M_\theta}$  is a one-fold covering of  $\mathfrak{GB}^{M_\theta}$ .

If  $M/\mathbb{T}^2$  is simply connected, then Theorems 8.9 and 8.15 combine to prove the second statement.  $\square$

The above result allows for the following explicit bundle description of the group of inner automorphisms of  $C(M_\theta)$ . Note that  $M/\mathbb{T}^2$  is simply connected when  $M$  is (see Note 12 below).

**Corollary 8.17** *If  $M/\mathbb{T}^2$  is simply connected, then*

$$\text{Inn}(C(M_\theta)) \cong \Gamma(M/\mathbb{T}^2, \mathfrak{GB}^{M_\theta}).$$

**Proof** In Proposition 8.13 we have already established that  $Z(C(M_\theta)) \cong C(M_\theta)_{J_M}$ . Hence Corollary 7.6 applies and gives the group isomorphism  $\text{Inn}(C(M_\theta)) \cong \mathfrak{G}(C(M_\theta), \mathcal{H}; J_M)$ . Combining this with Proposition 8.16 yields the desired result.  $\square$

## Notes

### Section 8.1. Commutative subalgebra and $C^*$ -bundles

1. Chapter 8 is based on [1].
2. The definition of the commutative subalgebra  $\mathcal{A}_J$  in Sect. 8.1 is quite similar to the definition of a subalgebra of  $\mathcal{A}$  defined in [2, Proposition 3.3] (cf. [3, Proposition 1.125]), which is the *real* commutative subalgebra in the center of  $\mathcal{A}$  consisting of elements for which  $aJ = Ja$ . Following [4] we propose a similar but different definition, since this subalgebra will turn out to be very useful for the description of the gauge group associated to any real spectral triple.
3. The notion of  $C(X)$ -algebra was introduced by Kasparov in [5]. Proposition 8.4 and Theorem 8.5 are proved in [6, 7] (see also Appendix C in [8]). Note that the bundles are in general only upper semi-continuous, and not necessarily continuous. For a discussion of this point, see [7].

### Section 8.2. Localization of the gauge group

4. For a proof of Proposition 8.8 see [9, Proposition 1.33].
5. Theorem 8.9 generalizes a result of [10] on Lie group bundles to the general setting of group bundles.

### Section 8.3. Localization of gauge fields

6. Later, in Chaps. 10–13 we will work towards physical applications in which the above  $C^*$ -bundle is a locally trivial (or, even a globally trivial)  $*$ -algebra bundle with finite-dimensional fiber. The above generalized gauge theories then become ordinary gauge theories, defined in terms of vector bundles and connections. It would be interesting to study the gauge theories corresponding to the intermediate cases, such as continuous trace  $C^*$ -algebras (cf. [11] for a definition), or the more general KK-fibrations that were introduced in [12].

### Section 8.4. Localization of toric noncommutative manifolds

7. Theorem 8.12 is due to Čaćić in [13, Theorem 4.28].
8. The fact that the spectral triple  $(C^\infty(M_\theta), \mathcal{H}, D; J_M)$  for  $\theta$  is an example of an almost-commutative spectral triple in the sense of [10, 14, 15] was already noticed in [13].
9. A proof of the fact that the center of  $A_\theta$  is trivial if  $\theta$  is irrational can be found in e.g. [16, Proposition 3].
10. The fact that for  $\theta$  irrational the  $C^*$ -algebra  $C(M_\theta)$  is isomorphic to the  $C^*$ -algebra  $\Gamma(M/\mathbb{T}^2, \mathfrak{B}^{M_\theta})$  of continuous sections of an upper semi-continuous  $C^*$ -bundle  $\mathfrak{B}^{M_\theta} \rightarrow M/\mathbb{T}^2$ . This also follows from the more general results of [17] showing that torus-covariant  $C(X)$ -algebras are deformed to torus-covariant  $C(X)$ -algebras. Here a torus-covariant algebra is a  $C(X)$ -algebra which carries an action of  $\mathbb{T}^2$  that commutes with  $C(X)$ . In particular, this applies to the  $C(M/\mathbb{T}^2)$ -algebra  $C(M)$ , deforming to the  $C(M/\mathbb{T}^2)$ -algebra  $C(M_\theta)$ .

11. For the proof of Theorem 8.15 we may argue as follows: in addition to upper semi-continuity, in [17, Proposition 5.1] lower semi-continuity is shown to hold under some additional conditions. In fact, since the  $\mathbb{T}^2$ -orbit space of  $M$  is Hausdorff, Corollary 5.3 in *loc. cit.* implies that the Rieffel deformation  $C(M_\theta)$  of  $C(M)$  can be expressed as a continuous field of  $C^*$ -algebras over this orbit space. In other words, it is the  $C^*$ -algebra of sections of a continuous  $C^*$ -bundle over  $M/\mathbb{T}^2$ . The second claim follows from [17, Corollary 6.2].
12. A proof of the fact that  $M/\mathbb{T}^2$  is simply connected when  $M$  is can be found in [18, Corollary 6.3].
13. Even though we have restricted the discussion in Sect. 8.4 to an action of a 2-torus on a manifold, this can be generalized in a straightforward manner to actions of higher-dimensional tori. The appropriate notion of irrationality for the higher-dimensional noncommutative tori has been discussed in [19].
14. In addition to the topological factorization of toric noncommutative manifolds in a horizontal and vertical part obtained in Sect. 8.4, also the factorization of the geometric structure has been studied in [20–23]. These works involve the unbounded external Kasparov product, allowing for a tensor-sum decomposition of the Dirac operator on  $M_\theta$  (cf. Sect. 5.3.2) into a vertical operator and a Dirac operator on  $M/\mathbb{T}^n$ .

## References

1. van Suijlekom, W.D.: Localizing gauge theories from noncommutative geometry. *Adv. Math.* **290**, 682–708 (2016)
2. Chamseddine, A.H., Connes, A., Marcolli, M.: Gravity and the standard model with neutrino mixing. *Adv. Theor. Math. Phys.* **11**, 991–1089 (2007)
3. Connes, A., Marcolli, M.: Noncommutative Geometry Quantum Fields and Motives. AMS, Providence (2008)
4. van den Dungen, K., van Suijlekom, W.D.: Particle physics from almost commutative space-times. *Rev. Math. Phys.* **24**, 1230004 (2012)
5. Kasparov, G.G.: Equivariant  $KK$ -theory and the Novikov conjecture. *Invent. Math.* **91**, 147–201 (1988)
6. Kirchberg, E., Wassermann, S.: Operations on continuous bundles of  $C^*$ -algebras. *Math. Ann.* **303**, 677–697 (1995)
7. Nilsen, M.:  $C^*$ -bundles and  $C_0(X)$ -algebras. *Indiana Univ. Math. J.* **45**, 463–477 (1996)
8. W. D. P. Crossed products of  $C^*$ -algebras, Mathematical Surveys and Monographs, vol. 134. American Mathematical Society, Providence, RI (2007)
9. Hatcher, A.: Algebraic Topology. Cambridge University Press, Cambridge (2002)
10. Boeijink, J., van den Dungen, K.: On globally non-trivial almost-commutative manifolds. *J. Math. Phys.* **55**, 103508 (2014)
11. Raeburn, I., Williams, D.P.: Morita equivalence and continuous-trace  $C^*$ -algebras. Mathematical Surveys and Monographs, vol. 60. American Mathematical Society, Providence, RI (1998)
12. Echterhoff, S., Nest, R., Oyono-Oyono, H.: An analogue of Serre fibrations for  $C^*$ -algebra bundles. In: Noncommutativity and Singularities, Advanced Studies Pure Mathematics, vol. 55, pp. 209–221. Mathematical Society, Japan, Tokyo (2009)
13. Čaćić, B.: A reconstruction theorem for Connes-Landi deformations of commutative spectral triples. *J. Geom. Phys.* **98**, 82–109 (2015)
14. Boeijink, J., van Suijlekom, W.D.: The noncommutative geometry of Yang-Mills fields. *J. Geom. Phys.* **61**, 1122–1134 (2011)
15. Čaćić, B.: A Reconstruction theorem for almost-commutative spectral triples. *Lett. Math. Phys.* **100**, 181–202 (2012)

16. Connes, A., Dubois-Violette, M.: Noncommutative finite-dimensional manifolds. I. Spherical manifolds and related examples. *Commun. Math. Phys.* **230**, 539–579 (2002)
17. Belmonte, F., Mantoiu, M.: Covariant fields of  $C^*$ -algebras under Rieffel deformation. *SIGMA Symmetry Integrability Geom. Methods Appl.* **8** (2012) Paper 091, 12
18. Bredon, G.E.: *Introduction to Compact Transformation Groups*, Pure and Applied Mathematics, vol. 46. Academic, New York-London (1972)
19. Rieffel, M.A.: Non-commutative tori - A case study of non-commutative differentiable manifolds. *Contemp. Math.* **105**, 191–212 (1990)
20. Brain, S., Mesland, B., van Suijlekom, W.D.: Gauge theory for spectral triples and the unbounded Kasparov product. *J. Noncommut. Geom.* **10**, 135–206 (2016)
21. Kaad, J., van Suijlekom, W.D.: Factorization of Dirac operators on almost-regular fibrations of  $\text{spin}^c$  manifolds. *Doc. Math.* **25**, 2049–2084 (2020)
22. Kaad, J., van Suijlekom, W.D.: Factorization of Dirac operators on toric noncommutative manifolds. *J. Geom. Phys.* **132**, 282–300 (2018)
23. Čaćić, B., Mesland, B.: Gauge theory on noncommutative Riemannian principal bundles. *Commun. Math. Phys.* **388**, 107–198 (2021)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 9

## Spectral Invariants



In the previous chapter we have identified the gauge group canonically associated to any spectral triple and have derived the generalized gauge fields that carry an action of that gauge group. In this chapter we take the next step and search for gauge invariants of these gauge fields, to wit, the spectral action, the topological spectral action and the fermionic action. We derive (asymptotic) expansions of the spectral action.

### 9.1 Spectral Action Functional

The simplest *spectral invariant* associated to a spectral triple  $(\mathcal{A}, \mathcal{H}, D)$  is given by the trace of some function of  $D$ . We also allow for inner fluctuations, and more generally consider the operators  $D_\omega = D + \omega + \epsilon' J \omega J^{-1}$  with  $\omega = \omega^* \in \Omega_D^1(\mathcal{A})$ .

**Definition 9.1** Let  $f$  be a suitable positive and even function from  $\mathbb{R}$  to  $\mathbb{R}$ . The *spectral action* is defined by

$$S_b[\omega] := \text{Tr } f(D_\omega/\Lambda), \quad (9.1.1)$$

where  $\Lambda$  is a real cutoff parameter. The minimal condition on the function  $f$  is that it makes  $f(D_\omega/\Lambda)$  a traceclass operator, requiring sufficiently rapid decay at  $\pm\infty$ .

The subscript  $b$  refers to *bosonic* since in the later physical applications  $\omega$  will describe bosonic fields.

There is also a *topological spectral action*, which is defined in terms of the grading  $\gamma$  by

$$S_{\text{top}}[\omega] = \text{Tr } \gamma f(D_\omega/\Lambda). \quad (9.1.2)$$

The term ‘topological’ will be justified below. First, we prove gauge invariance of these functionals.

**Theorem 9.2** *The spectral action and the topological spectral action are gauge invariant functionals of the gauge field  $\omega \in \Omega_D^1(\mathcal{A})$ , assumed to transform under  $\text{Ad } u = uJuJ^{-1} \in \mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$  as*

$$\omega \mapsto u\omega u^* + u[D, u^*].$$

**Proof** By (7.2.6) this is equivalent to  $D_\omega \mapsto UD_\omega U^*$  with  $U = uJuJ^{-1}$ . Since the eigenvalues of  $UD_\omega U^*$  coincide with those of  $D_\omega$  and the (topological) spectral action is defined on the spectrum of  $D_\omega$ , the result follows.  $\square$

Another gauge invariant one can naturally associate to a spectral triple is of a *fermionic* nature, as opposed to the above bosonic spectral action functional. This invariant is given by combining the operator  $D_\omega$  with a Grassmann vector in the Hilbert space (cf. Appendix 11.3), as follows.

**Definition 9.3** The *fermionic action* is defined by

$$S_f[\omega, \psi] = (J\tilde{\psi}, D_\omega\tilde{\psi})$$

with  $\tilde{\psi} \in \mathcal{H}_{\text{cl}}^+$  where

$$\mathcal{H}_{\text{cl}}^+ = \left\{ \tilde{\psi} : \psi \in \mathcal{H}^+ \right\}$$

is the set of Grassmann variables in  $\mathcal{H}$  in the  $+1$ -eigenspace of the grading  $\gamma$ .

**Theorem 9.4** *The fermionic action is a gauge invariant functional of the gauge field  $\omega$  and the fermion field  $\psi$ , the latter transforming under  $\text{Ad } u \in \mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$  as*

$$\psi \mapsto uJuJ^{-1}\psi.$$

Moreover, if the KO-dimension of  $(\mathcal{A}, \mathcal{H}, D; \gamma, J)$  is 2 modulo 8, then  $(\psi, \psi') \mapsto \langle J\psi, D_\omega\psi' \rangle$  defines a skew-symmetric form on the  $+1$ -eigenspace of  $\gamma$  in  $\mathcal{H}$ .

**Proof** Again,  $D_\omega \mapsto UD_\omega U^*$  with  $U = uJuJ^{-1}$ , whilst  $\psi \mapsto U\psi$ . The claim then follows from the observation  $UJ = JU$ .

Skew-symmetry follows from a small computation:

$$\langle J\psi, D\psi' \rangle = -\langle J\psi, J^2D\psi' \rangle = -\langle JD\psi', \psi \rangle = -\langle DJ\psi', \psi \rangle = -\langle J\psi', D\psi \rangle.$$

where we used Table 4.2 for  $DJ = JD$  in KO-dimension 2 modulo 8.  $\square$

The above skew-symmetry is in concordance with the Grassmann nature of fermionic fields  $\tilde{\psi}$ , guaranteeing that  $S_f$  as defined above is in fact non-zero.

## 9.2 Asymptotic Expansion of the Spectral Action

The first type of expansion of the spectral action is an asymptotic series in powers of  $\Lambda$ ; the perturbative expansion in powers of the gauge field  $\omega$  will be considered in the next section.

We assume that  $f$  is given by a Laplace–Stieltjes transform:

$$f(x) = \int_{t>0} e^{-tx^2} d\mu(t),$$

with  $\mu$  a suitable measure on  $\mathbb{R}^+$ . This assumption allows us to find the following expression for the topological spectral action.

**Proposition 9.5** *Suppose  $f$  is of the above form. Then,*

$$S_{\text{top}}[\omega] = f(0) \text{ index } D_\omega.$$

**Proof** This follows from the McKean–Singer formula (Lemma 6.23):

$$\text{index } D_\omega = \text{Tr } \gamma e^{-tD_\omega^2/\Lambda^2}.$$

Since this expression is independent of  $\Lambda$  and  $t$ , an integration over  $t$  yields

$$\int_{t>0} d\mu(t) = f(0).$$

□

The asymptotic expansion of  $S$  can be derived from the existence of a **heat kernel expansion** of the form

$$\text{Tr } e^{-tD^2} = \sum_{\alpha} t^\alpha c_\alpha, \quad (9.2.1)$$

as  $t \rightarrow 0$ . Note that this is written down here for the unperturbed operator  $D$ , but similar expressions hold for any bounded perturbation of  $D$ , such as  $D_\omega$ .

**Lemma 9.6** *If  $(\mathcal{A}, \mathcal{H}, D)$  is a regular spectral triple with simple dimension spectrum (see Definition 6.9), then the heat kernel expansion (9.2.1) is valid as an asymptotic expansion as  $t \rightarrow 0$ . Moreover, for  $\alpha < 0$  we have*

$$\text{res}_{z=-2\alpha} \zeta_1(z) = \frac{2c_\alpha}{\Gamma(-\alpha)},$$

with  $\zeta_b(z) = \text{Tr } b|D|^{-z}$ .

**Proof** This follows from the Mellin transform:

$$|D|^{-z} = \frac{1}{\Gamma(z/2)} \int_0^\infty e^{-tD^2} t^{z/2-1} dt, \quad (9.2.2)$$

or, after inserting the heat kernel expansion,

$$\begin{aligned} \text{Tr } |D|^{-z} &= \frac{1}{\Gamma(z/2)} \sum_\alpha \int_0^\infty c_\alpha t^{\alpha+z/2-1} dt \\ &= \frac{1}{\Gamma(z/2)} \sum_\alpha \int_0^1 c_\alpha t^{\alpha+z/2-1} dt + \text{holomorphic} \\ &= \sum_\alpha \frac{c_\alpha}{\Gamma(z/2)(\alpha+z/2)} + \text{holomorphic}. \end{aligned}$$

Taking residues at  $z = -2\alpha$  on both sides gives the desired result.  $\square$

Using the Laplace–Stieltjes transform, we now derive an asymptotic expansion of the spectral action in terms of the heat coefficients  $c_\alpha$ .

**Proposition 9.7** *Under the above conditions, the spectral action is given asymptotically (as  $\Lambda \rightarrow \infty$ ) by*

$$\text{Tr } f(D/\Lambda) = \sum_{\beta \in \text{Sd}} f_\beta \Lambda^\beta \frac{2}{\Gamma(\beta/2)} c_{-\frac{1}{2}\beta} + f(0)c_0 + \mathcal{O}(\Lambda^{-1}), \quad (9.2.3)$$

where  $f_\beta := \int f(v)v^{\beta-1}dv$  and  $\text{Sd}$  is the dimension spectrum of  $(\mathcal{A}, \mathcal{H}, D)$ .

**Proof** This follows directly after inserting the heat expansion in the Laplace–Stieltjes transform:

$$\text{Tr } f(D/\Lambda) = \sum_\alpha \int_{t>0} t^\alpha \Lambda^\alpha c_\alpha d\mu(t). \quad (9.2.4)$$

The terms with  $\alpha > 0$  are of order  $\Lambda^{-1}$ ; if  $\alpha < 0$ , then

$$t^\alpha = \frac{1}{\Gamma(-\alpha)} \int_{v>0} e^{-tv} v^{-\alpha-1} dv.$$

Applying this to the integral (9.2.4) gives

$$\begin{aligned} \Lambda^{-2\alpha} c_\alpha \int_{t>0} t^\alpha d\mu(t) &= \Lambda^{-2\alpha} c_\alpha \int_{t>0} \int_{v>0} e^{-tv} v^{-\alpha-1} dv d\mu(t) \\ &= 2\Lambda^{-2\alpha} c_\alpha \int_{t>0} \int_{v>0} e^{-tv^2} v^{-2\alpha-1} dv d\mu(t) \\ &= 2\Lambda^{-2\alpha} c_\alpha \int_{v>0} f(v) v^{-2\alpha-1} dv \equiv 2\Lambda^{-2\alpha} c_\alpha f_{-2\alpha}, \end{aligned}$$

substituting  $v \mapsto v^2$  in going to the second line. Since  $c_\alpha = 0$  unless  $-2\alpha \in \text{Sd}$ , we substitute  $\beta = -2\alpha$  to obtain (9.2.3).  $\square$

**Corollary 9.8** *For the perturbed operator  $D_\omega$  we have*

$$S_b[\omega] = \sum_{\beta \in \text{Sd}} f_\beta \Lambda^\beta \text{res}_{z=\beta} \text{Tr} |D_\omega|^{-z} + f(0) \text{Tr} |D_\omega|^{-z}|_{z=0} + \mathcal{O}(\Lambda^{-1}).$$

### 9.3 Perturbative Expansion in the Gauge Field

Another approach to analyze  $S_b$  is given by expanding as a Taylor series in powers of  $\omega$ , rather than in  $\Lambda$ . We first take a closer look at the heat operator  $e^{-tD^2}$  and its perturbations.

**Lemma 9.9** *Let  $\omega$  be a bounded operator and denote  $D_\omega = D + \omega$ . Then*

$$e^{-t(D_\omega)^2} = e^{-tD^2} - t \int_0^1 ds e^{-st(D_\omega)^2} P(\omega) e^{-(1-s)tD^2},$$

with  $P(\omega) = D\omega + \omega D + \omega^2$ .

**Proof** Note that  $e^{-tD_\omega^2}$  is the unique solution of the Cauchy problem

$$\begin{cases} (d_t + D_\omega) u(t) = 0 \\ u(0) = 1, \end{cases}$$

with  $d_t = d/dt$ . Using the fundamental theorem of calculus, we find

$$\begin{aligned} d_t & \left[ e^{-tD^2} - \int_0^t dt' e^{-(t-t')D_\omega^2} P(\omega) e^{-t'D^2} \right] \\ &= -D_\omega^2 \left( e^{-tD^2} - \int_0^t dt' e^{-(t-t')D_\omega^2} P(\omega) e^{-t'D^2} \right), \end{aligned}$$

showing that the bounded operator  $e^{-tD^2} - \int_0^t dt' e^{-(t-t')D_\omega^2} P(\omega) e^{-t'D^2}$  also solves the above Cauchy problem.  $\square$

In what follows, we will repeatedly apply this Lemma to obtain a perturbative expansion for  $e^{-t(D_\omega)^2}$  in powers of  $\omega$  in terms of multiple integrals of heat operators. We introduce the following convenient notation, valid for operators  $X_0, \dots, X_n$ :

$$\langle X_0, \dots, X_n \rangle_{t,n} := t^n \text{Tr} \int_{\Delta_n} X_0 e^{-s_0 t D^2} X_1 e^{-s_1 t D^2} \cdots X_n e^{-s_n t D^2} d^n s.$$

Here, the standard  $n$ -simplex  $\Delta_n$  is the set of all  $n$ -tuples  $(t_1, \dots, t_n)$  satisfying  $0 \leq t_1 \leq \dots \leq t_n \leq 1$ . Equivalently,  $\Delta_n$  can be given as the set of  $n+1$ -tuples

$(s_0, s_1, \dots, s_n)$  such that  $s_0 + \dots + s_n = 1$  and  $0 \leq s_i \leq 1$  for any  $i = 0, \dots, n$ . Indeed, we have  $s_0 = t_1$ ,  $s_i = t_{i+1} - t_i$  and  $s_n = 1 - t_n$  and, vice versa,  $t_k = s_0 + s_1 + \dots + s_{k-1}$ .

We recall the notion of Gâteaux derivatives.

**Definition 9.10** The *Gâteaux derivative* of a map  $F : X \rightarrow Y$  (between locally convex topological vector spaces) at  $x \in X$  is defined for  $h \in X$  by

$$F'(x)(h) = \lim_{u \rightarrow 0} \frac{F(x + uh) - F(x)}{u}.$$

In general, the map  $F'(x)(\cdot)$  is not linear, in contrast with the Fréchet derivative. However, if  $X$  and  $Y$  are Fréchet spaces, then the Gâteaux derivatives actually defines a linear map  $F'(x)(\cdot)$  for any  $x \in X$ . In this case, higher order derivatives are denoted as  $F''$ ,  $F'''$ , et cetera, or more conveniently as  $F^{(k)}$  for the  $k$ 'th order derivative. The latter will be understood as a bounded operator from  $X \times \dots \times X$  ( $k+1$  copies) to  $Y$ , which is linear in the  $k$  last variables.

**Theorem 9.11** (Taylor's formula with integral remainder) *For a Gâteaux  $k+1$ -differentiable map  $F : X \rightarrow Y$  between Fréchet spaces  $X$  and  $Y$ ,*

$$\begin{aligned} F(x) &= F(a) + F'(a)(x - a) + \frac{1}{2!} F''(a)(x - a, x - a) + \dots \\ &\quad + \frac{1}{n!} F^{(k)}(a)(x - a, \dots, x - a) + R_k(x), \end{aligned}$$

for  $x, a \in X$ , with remainder given by

$$R_k(x) = \frac{1}{k!} \int_0^1 F^{(k+1)}(a + t(x - a))((1-t)h, \dots, (1-t)h, h) dt.$$

In view of this Theorem, we have the following asymptotic Taylor expansion (around 0) in  $\omega \in \Omega_D^1(\mathcal{A})$  for the spectral action  $S_b[\omega]$ :

$$S_b[\omega] = \sum_{n=0}^{\infty} \frac{1}{n!} S_b^{(n)}(0)(\omega, \dots, \omega), \tag{9.3.1}$$

provided we make the following

**Assumption 1** For all  $\alpha > 0$ ,  $\beta > 0$ ,  $\gamma > 0$  and  $0 \leq \epsilon < 1$ , there exist constants  $C_{\alpha\beta\gamma\epsilon}$  such that

$$\int_{t>0} \mathrm{Tr} t^\alpha |D|^\beta e^{-t(\epsilon D^2 - \beta)} |d\mu(t)| < C_{\alpha\beta\gamma\epsilon}.$$

**Proposition 9.12** If  $n = 0, 1, \dots$  and  $\omega \in \Omega_D^1(\mathcal{A})$ , then  $S_b^{(n)}(0)(\omega, \dots, \omega)$  exists, and

$$S_b^{(n)}(0)(\omega, \dots, \omega) = n! \sum_{k=0}^n (-1)^k \sum_{\varepsilon_1, \dots, \varepsilon_k} \langle 1, (1 - \varepsilon_1)\{D, \omega\} + \varepsilon_1 \omega^2, \dots, \\ (1 - \varepsilon_k)\{D, \omega\} + \varepsilon_k \omega^2 \rangle_{t,k} d\mu(t),$$

where the sum is over multi-indices  $(\varepsilon_1, \dots, \varepsilon_k) \in \{0, 1\}^k$  such that  $\sum_{i=1}^k (1 + \varepsilon_i) = n$ .

**Proof** We prove this by induction on  $n$ , the case  $n = 0$  being trivial. By definition of the Gâteaux derivative and using Lemma 9.9,

$$S_b^{(n+1)}(0)(\omega, \dots, \omega) = n! \sum_{k=0}^n \sum_{\varepsilon_1, \dots, \varepsilon_k} \left[ \sum_{i=1}^k (-1)^{k+1} \langle 1, (1 - \varepsilon_1)\{D, \omega\} + \varepsilon_1 \omega^2, \dots, \right. \\ \left. \{D, \omega\}, \dots, (1 - \varepsilon_k)\{D, \omega\} + \varepsilon_k \omega^2 \rangle_{t,k+1} \right. \\ \left. + \sum_{i=1}^k (-1)^k \langle 1, (1 - \varepsilon_1)\{D, \omega\} + \varepsilon_1 \omega^2, \dots, 2(1 - \varepsilon_i)\omega^2, \dots, \right. \\ \left. (1 - \varepsilon_k)\{D, \omega\} + \varepsilon_k \omega^2 \rangle_{t,k} \right] d\mu(t).$$

The first sum corresponds to a multi-index  $\vec{\varepsilon}' = (\varepsilon_1, \dots, \varepsilon_{i-1}, 0, \varepsilon_i, \dots, \varepsilon_k)$ , the second corresponds to  $\vec{\varepsilon}' = (\varepsilon_1, \dots, \varepsilon_i + 1, \dots, \varepsilon_k)$  if  $\varepsilon_i = 0$ , counted with a factor of 2. In both cases, we compute that  $\sum_j (1 + \varepsilon'_j) = n + 1$ . In other words, the induction step from  $n$  to  $n + 1$  corresponds to inserting in a sequence of 0's and 1's (of, say, length  $k$ ) either a zero at any of the  $k + 1$  places, or replacing a 0 by a 1 (with the latter counted twice). In order to arrive at the right combinatorial coefficient  $(n + 1)!$ , we have to show that any  $\vec{\varepsilon}'$  satisfying  $\sum_i (1 + \varepsilon'_i) = n + 1$  appears in precisely  $n + 1$  ways from  $\vec{\varepsilon}$  that satisfy  $\sum_i (1 + \varepsilon_i) = n$ . If  $\vec{\varepsilon}'$  has length  $k$ , it contains  $n + 1 - k$  times 1 as an entry and, consequently,  $2k - n - 1$  a 0. This gives (with the double counting for the 1's) for the number of possible  $\vec{\varepsilon}'$ :

$$2(n + 1 - k) + 2k - n - 1 = n + 1,$$

as claimed. This completes the proof.  $\square$

**Example 9.13**

$$\begin{aligned} S_b^{(1)}(0)(\omega) &= \int \left( -\langle 1, \{D, \omega\} \rangle_{t,1} \right) d\mu(t), \\ S_b^{(2)}(0)(\omega, \omega) &= 2 \int \left( -\langle 1, \omega^2 \rangle_{t,1} + \langle 1, \{D, \omega\}, \{D, \omega\} \rangle_{t,2} \right) d\mu(t), \\ S_b^{(3)}(0)(\omega, \omega, \omega) &= 3! \int \left( \langle 1, \omega^2, \{D, \omega\} \rangle_{t,2} + \langle 1, \{D, \omega\}, \omega^2 \rangle_{t,2} \right. \\ &\quad \left. - \langle 1, \{D, \omega\}, \{D, \omega\}, \{D, \omega\} \rangle_{t,3} \right) d\mu(t). \end{aligned}$$

**9.3.1 Taylor Expansion of the Spectral Action**

We fix a complete set of eigenvectors  $\{\psi_j\}_j$  of  $D$  with eigenvalues  $\lambda_j \in \mathbb{R}$ , respectively, forming an orthonormal basis for  $\mathcal{H}$ . We also write  $\omega_{ij} := (\psi_i, \omega \psi_j)$  for the matrix coefficients of  $\omega$  with respect to this orthonormal basis. Recall from Appendix 9.A the notion of divided difference  $f[x_0, x_1, \dots, x_n]$  of a function  $f : \mathbb{R} \rightarrow \mathbb{R}$ .

**Theorem 9.14** *If  $f$  satisfies Assumption 1 and  $\omega \in \Omega_D^1(\mathcal{A})$ , then*

$$S_b^{(n)}(0)(\omega, \dots, \omega) = n! \sum_{i_1, \dots, i_n} \omega_{i_n i_1} \omega_{i_1 i_2} \cdots \omega_{i_{n-1} i_n} f[\lambda_{i_p}, \lambda_{i_1}, \dots, \lambda_{i_n}].$$

**Proof** Proposition 9.12 gives us an expression for  $S_b^{(n)}$  in terms of the brackets  $\langle \dots \rangle_t$ . For these we compute:

$$\begin{aligned} &(-1)^k \langle 1, (1 - \varepsilon_1)\{D, \omega\} + \varepsilon_1 \omega^2, \dots, (1 - \varepsilon_k)\{D, \omega\} + \varepsilon_k \omega^2 \rangle_{t,k} d\mu(t) \\ &= (-1)^k \sum_{i_0=i_k, i_1, \dots, i_k} \int_{\Delta_k} \left( \prod_{j=1}^k ((1 - \varepsilon_j)(\lambda_{i_{j-1}} - \lambda_{i_j})\omega + \varepsilon_j \omega^2)_{i_{j-1} i_j} \right) \\ &\quad \times e^{-(s_0 t \lambda_{i_0}^2 + \dots + s_k t \lambda_{i_k}^2)} d^k s d\mu(t) \\ &= \sum_{i_0=i_k, i_1, \dots, i_k} \left( \prod_{j=1}^k ((1 - \varepsilon_j)(\lambda_{i_{j-1}} - \lambda_{i_j})\omega + \varepsilon_j \omega^2)_{i_{j-1} i_j} \right) g[\lambda_{i_0}^2, \dots, \lambda_{i_k}^2]. \end{aligned}$$

Glancing back at Proposition 9.27, we are finished if we establish a one-to-one relation between the order index sets  $I = \{0 = i_0 < i_1 < \dots < i_k = n\}$  such that  $i_{j-1} - i_j \leq 2$  for all  $1 \leq j \leq k$  and the multi-indices  $(\varepsilon_1, \dots, \varepsilon_k) \in \{0, 1\}^k$  such that  $\sum_{i=1}^k (1 + \varepsilon_i) = n$ . If  $I$  is such an index set, we define a multi-index

$$\varepsilon_j = \begin{cases} 0 & \text{if } \{i_j - 1, i_j\} \subset I, \\ 1 & \text{otherwise.} \end{cases}$$

Indeed,  $i_j = i_{j-1} + 1 + \varepsilon_j$ , so that

$$\sum_{i=1}^k (1 + \varepsilon_i) = i_0 + \sum_{i=1}^k (1 + \varepsilon_i) = i_k = n.$$

It is now clear that, *vice-versa*, if  $\varepsilon$  is as above, we define

$$I = \{0 = i_0 < i_1 < \dots < i_k = n\}$$

by  $i_j = i_{j-1} + 1 + \varepsilon_j$ , and starting with  $i_0 = 0$ .  $\square$

**Corollary 9.15** *If  $n \geq 0$  and  $\omega \in \Omega_D^1(\mathcal{A})$ , then*

$$S_b^{(n)}(0)(\omega, \dots, \omega) = (n-1)! \sum_{i_1, \dots, i_n} \omega_{i_1 i_2} \cdots \omega_{i_n i_1} f'[\lambda_{i_1}, \dots, \lambda_{i_n}].$$

Consequently,

$$S_b[\omega] = S_b[0] + \sum_{n=1}^{\infty} \frac{1}{n} \sum_{i_1, \dots, i_n} \omega_{i_1 i_2} \cdots \omega_{i_n i_1} f'[\lambda_{i_1}, \dots, \lambda_{i_n}].$$

An interesting consequence is the following.

**Corollary 9.16** *If  $n \geq 0$  and  $\omega \in \Omega_D^1(\mathcal{A})$  and if  $f'$  has compact support, then*

$$S_b^{(n)}(0)(\omega, \dots, \omega) = \frac{(n-1)!}{2\pi i} \operatorname{Tr} \oint f'(z) \omega(z-D)^{-1} \cdots \omega(z-D)^{-1},$$

where the contour integral encloses the intersection of the spectrum of  $D$  with  $\operatorname{supp} f'$ .

**Proof** This follows directly from Cauchy's formula for divided differences (see Note 16 on this chapter):

$$g[x_0, \dots, x_n] = \frac{1}{2\pi i} \oint \frac{g(z)}{(z-x_0) \cdots (z-x_n)} dz,$$

with the contour enclosing the points  $x_i$ .  $\square$

### 9.3.2 Cyclic Cocycles Underlying the Spectral Action

We now come to analyze in more details the structure of the terms in the Taylor expansion (9.3.1). We will write it as

$$S_b[\omega] - S_b[0] = \sum_n \frac{1}{n} \langle \underbrace{\omega, \dots, \omega}_n \rangle_f \quad (9.3.2)$$

where we have introduced brackets  $\langle \cdot \rangle_f$  as the following multilinear functionals  $\langle \cdot \rangle_f : (\Omega_D^1(\mathcal{A}))^{\times n} \rightarrow \mathbb{C}$ :

$$\langle \omega_1, \dots, \omega_n \rangle_f := \sum_{i_1, \dots, i_n} (\omega_1)_{i_1 i_2} \cdots (\omega_n)_{i_n i_1} f'[\lambda_{i_1}, \dots, \lambda_{i_n}] \quad (9.3.3)$$

For our algebraic results we only need two simple properties of the bracket  $\langle \cdot \rangle_f$ , stated in the following lemma. After proving this lemma, all analytical subtleties are taken care of, and we can focus on the algebra that ensues from these simple rules.

**Lemma 9.17** *For  $\omega_1, \dots, \omega_n \in \Omega^1(\mathcal{A})$  and  $a \in \mathcal{A}$  we have*

$$\langle \omega_1, \dots, \omega_n \rangle_f = \langle \omega_n, \omega_1, \dots, \omega_{n-1} \rangle_f, \quad (\text{I})$$

$$\begin{aligned} \langle \omega_1, \dots, a\omega_j, \dots, \omega_n \rangle_f &= \langle \omega_1, \dots, \omega_{j-1}a, \dots, \omega_n \rangle_f \\ &= \langle \omega_1, \dots, \omega_{j-1}, [D, a], \omega_j, \dots, \omega_n \rangle_f \end{aligned} \quad (\text{II})$$

where it is understood that for the edge case  $j = 1$  we need to substitute  $n$  for  $j - 1$  on the left-hand side.

**Proof** Property I follows immediately from definition (9.3.3) and it also reduces II to the edge case  $j = 1$ . For that case we compute

$$\begin{aligned} \langle a\omega_1, \dots, \omega_n \rangle_f &- \langle \omega_1, \dots, \omega_n a \rangle_f \\ &= \sum_{i_0, \dots, i_n} a_{i_0 i_1} (\omega_1)_{i_1 i_2} \cdots (\omega_n)_{i_n i_0} (f'[\lambda_{i_0}, \lambda_{i_2}, \dots, \lambda_{i_n}] - f'[\lambda_{i_1}, \lambda_{i_2}, \dots, \lambda_{i_n}]) \\ &= \sum_{i_0, \dots, i_n} a_{i_0 i_1} (\lambda_{i_0} - \lambda_{i_1}) (\omega_1)_{i_1 i_2} \cdots (\omega_n)_{i_n i_0} f'[\lambda_{i_0}, \lambda_{i_1}, \lambda_{i_2}, \dots, \lambda_{i_n}] \\ &= \langle [D, a], \omega_1, \omega_2, \dots, \omega_n \rangle_f \end{aligned}$$

where in the second equality we used the recursive definition of the divided difference (see Definition 9.25 below).  $\square$

### 9.3.2.1 Hochschild and Cyclic Cocycles

When the above brackets  $\langle \cdot \rangle_f$  are evaluated at one-forms  $a[D, b]$  associated to a spectral triple, the relations found in Lemma 9.17 can be translated nicely in terms of the coboundary operators  $b$  and  $B$  appearing in the definition of periodic cyclic cohomology (cf. Sect. 6.2). Let us write  $B = AB_0$  where  $A$  is the operator of full anti-symmetrization while  $B_0 : C^n(\mathcal{A}) \rightarrow C^{n+1}(\mathcal{A})$  is defined as

$$B_0\phi(a^0, a^1, \dots, a^n) := \phi(1, a^0, \dots, a^n)$$

We define the following  $n$ -cochain:

$$\phi_n(a_0, \dots, a_n) := \langle a_0[D, a_1], [D, a_2], \dots, [D, a_n] \rangle_f \quad (9.3.4)$$

We easily see that  $B_0\phi_n$  is invariant under cyclic permutations, so that  $B\phi_n = nB_0\phi_n$  for odd  $n$  and  $B\phi_n = 0$  for even  $n$ . Also,  $\phi_n(a_0, \dots, a_n) = 0$  when  $a_j = 1$  for some  $j \geq 1$ . We put  $\phi_0 := 0$ .

**Lemma 9.18** *We have  $b\phi_n = \phi_{n+1}$  for odd  $n$  and  $b\phi_n = 0$  for even  $n$ .*

**Proof** As  $b\phi_0 = 0$  by definition, and  $b^2 = 0$ , we need only check the case in which  $n$  is odd.

We find, by splitting up the sum, and shifting the second appearing sum by one, that

$$\begin{aligned} b\phi_n(a_0, \dots, a_{n+1}) &= \langle a_0a_1[D, a_1], [D, a_2], \dots, [D, a_{n+1}] \rangle_f - \langle a_0a_1[D, a_1], [D, a_2], \dots, [D, a_{n+1}] \rangle_f \\ &\quad + \sum_{j=2}^n (-1)^j \langle a_0[D, a_1], [D, a_2], \dots, a_j[D, a_{j+1}], \dots, [D, a_{n+1}] \rangle_f \\ &\quad - \sum_{j=2}^{n+1} (-1)^j \langle a_0[D, a_1], [D, a_2], \dots, [D, a_{j-1}]a_j, \dots, [D, a_{n+1}] \rangle_f \\ &\quad + \langle a_{n+1}a_0[D, a_1], [D, a_2], \dots, [D, a_n] \rangle_f \\ &= \sum_{j=2}^n (-1)^j \langle a_0[D, a_1], [D, a_2], \dots, [D, a_{n+1}] \rangle_f \\ &\quad - \langle a_0[D, a_1], [D, a_2], \dots, [D, a_n]a_{n+1} \rangle_f + \langle a_{n+1}a_0[D, a_1], \dots, [D, a_n] \rangle_f \\ &= \langle [D, a_{n+1}], a_0[D, a_1], [D, a_2], \dots, [D, a_n] \rangle_f \\ &= \phi_{n+1}(a_0, \dots, a_{n+1}), \end{aligned}$$

by I and II of Lemma 9.17. □

**Lemma 9.19** *Let  $n$  be even. We have  $bB_0\phi_n = 2\phi_n - B_0\phi_{n+1}$ .*

**Proof** Splitting the sum in two, and shifting the index of the second sum, we find

$$\begin{aligned}
& bB_0\phi_n(a_0, \dots, a_n) \\
&= \sum_{j=0}^{n-1} (-1)^j \langle [D, a_0], \dots, a_j [D, a_{j+1}], \dots, [D, a_n] \rangle_f \\
&\quad - \sum_{j=1}^n (-1)^j \langle [D, a_0], \dots, [D, a_{j-1}]a_j, \dots, [D, a_n] \rangle_f + \langle [D, a_n a_0], \dots, [D, a_{n-1}] \rangle_f \\
&= \langle a_0 [D, a_1], [D, a_2], \dots, [D, a_n] \rangle_f + \sum_{j=1}^{n-1} (-1)^j \langle [D, a_0], \dots, [D, a_n] \rangle_f \\
&\quad - \langle [D, a_0], \dots, [D, a_{n-2}], [D, a_{n-1}]a_n \rangle_f + \langle [D, a_n a_0], \dots, [D, a_{n-1}] \rangle_f \\
&= \phi_n(a_0, \dots, a_n) - \langle [D, a_0], \dots, [D, a_n] \rangle_f + \langle [D, a_n], [D, a_0], \dots, [D, a_{n-1}] \rangle_f \\
&\quad + \langle [D, a_n]a_0, [D, a_1], \dots, [D, a_{n-1}] \rangle_f \\
&= 2\phi_n(a_0, \dots, a_n) - B_0\phi_{n+1}(a_0, \dots, a_n),
\end{aligned}$$

by using both properties of the bracket  $\langle \cdot \rangle_f$  in the last step.  $\square$

Motivated by this we define

$$\psi_{2k-1} := \phi_{2k-1} - \frac{1}{2}B_0\phi_{2k}, \quad (9.3.5)$$

so that

$$B\psi_{2k+1} = 2(2k+1)b\psi_{2k-1}.$$

We can rephrase this property in terms of the  $(b, B)$ -complex as follows.

**Proposition 9.20** *Let  $\phi_n$  and  $\psi_{2k-1}$  be as defined above and set  $\tilde{\psi}_{2k-1} := (-1)^{k-1} \frac{(k-1)!}{(2k-1)!} \psi_{2k-1}$ .*

- (1) *The sequence  $(\phi_{2k})$  is a  $(b, B)$ -cocycle and each  $\phi_{2k}$  defines an even Hochschild cocycle:  $b\phi_{2k} = 0$ .*
- (2) *The sequence  $(\tilde{\psi}_{2k-1})$  is an odd  $(b, B)$ -cocycle.*

We use an noncommutative integral notation that is defined by linear extension of

$$\int_{\phi} a_0 \delta a_1 \cdots \delta a_n := \int_{\phi_n} a_0 \delta a_1 \cdots \delta a_n := \phi(a_0, a_1, \dots, a_n),$$

and similarly for  $\psi$ . The expression  $a_0 \delta a_1 \cdots \delta a_n$  is a so-called *universal differential n-form* in  $\Omega^n(\mathcal{A})$ , see Note 7 on Chap. 7 for more details.

### 9.3.3 Brackets and Noncommutative Integrals Over Universal Forms

In this section we will express the derivatives of the fluctuated spectral action (occurring in the Taylor series (9.3.1)) in terms of universal forms that are integrated along

$\phi$ . We thus make the jump from an expression in terms of  $\omega = \pi_D(A) \in \Omega_D^1(\mathcal{A})_{\text{sa}}$  to an expression in terms of  $A \in \Omega^1(\mathcal{A})$ . As  $\omega$  decomposes as a finite sum  $\omega = \sum a_j[D, b_j]$ , our task is to express  $\langle a_{j_1}[D, b_{j_1}], \dots, a_{j_n}[D, b_{j_n}] \rangle_f$  in terms of universal forms  $a_0 \delta a_1 \cdots \delta a_n$  integrated along  $\phi$ . This will turn out to be possible by just using II and the Leibniz rule  $[D, a_1 a_2] = a_1 [D, a_2] + [D, a_1] a_2$ . To find the exact expression we need to work in the algebra  $M_2(\Omega_D^\bullet(\mathcal{A})) = M_2(\mathbb{C}) \otimes \Omega_D^\bullet(\mathcal{A})$ .

**Proposition 9.21** *Let  $n \in \mathbb{N}$ . For  $a_1, \dots, a_n, b_1, \dots, b_n \in \mathcal{A}$ , denoting  $A_j := a_j \delta b_j$ , we have*

$$\langle a_1[D, b_1], \dots, a_n[D, b_n] \rangle_f = \int_\phi (A_1 \ 0) \prod_{j=2}^n \begin{pmatrix} A_j + \delta A_j & -A_j \\ \delta A_j & -A_j \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

**Proof** If we combine, for every  $n \in \mathbb{N}_0$ , the  $n$ -multilinear function  $\langle \cdot \rangle_f$  from (9.3.3), we obtain a linear function

$$\langle \cdot \rangle_f : T\Omega_D^1(\mathcal{A}) \rightarrow \mathbb{C}$$

on the tensor algebra  $T\Omega_D^1(\mathcal{A})$ . For any  $\mu, \nu \in T\Omega_D^1(\mathcal{A})$ , a straightforward calculation using the commutation rule II from Lemma 9.17 shows that

$$\langle \mu \otimes a_{j-1}[D, b_{j-1}] \otimes (a_j \ a_j b_j) \nu \rangle_f = \langle \mu \otimes (a_{j-1} \ a_{j-1} b_{j-1}) M_j \otimes \nu \rangle_f, \quad (9.3.6)$$

where  $M_j \in M_2(T\Omega^1(\mathcal{A}))$  is defined by

$$M_j := \begin{pmatrix} [D, b_{j-1} a_j] + [D, b_{j-1}] \otimes [D, a_j] & [D, b_{j-1} a_j b_j] + [D, b_{j-1}] \otimes [D, a_j b_j] \\ -[D, a_j] & -[D, a_j b_j] \end{pmatrix}. \quad (9.3.7)$$

Repeating (9.3.6), and subsequently using (9.3.4), it follows that

$$\begin{aligned} \langle a_1[D, b_1], \dots, a_n[D, b_n] \rangle_f &= \langle a_1[D, b_1] \otimes \dots \otimes a_{n-1}[D, b_{n-1}] \otimes (a_n \ a_n b_n) \begin{pmatrix} [D, b_n] \\ 0 \end{pmatrix} \rangle_f \\ &= \langle (a_1 \ a_1 b_1) \left( \prod_{j=2}^n M_j \right) \begin{pmatrix} [D, b_n] \\ 0 \end{pmatrix} \rangle_f \\ &= \int_\phi (a_1 \ a_1 b_1) \left( \prod_{j=2}^n N_j \right) \begin{pmatrix} \delta b_n \\ 0 \end{pmatrix}, \end{aligned}$$

where from (9.3.7) we obtain

$$\begin{aligned} N_j &= \begin{pmatrix} \delta(b_{j-1} a_j) + \delta b_{j-1} \delta a_j & \delta(b_{j-1} a_j b_j) + \delta b_{j-1} \delta(a_j b_j) \\ -\delta a_j & -\delta(a_j b_j) \end{pmatrix} \\ &= \begin{pmatrix} \delta b_{j-1} & b_{j-1} \\ 0 & -1 \end{pmatrix} \begin{pmatrix} a_j + \delta a_j & a_j b_j + \delta a_j b_j + a_j \delta b_j \\ \delta a_j & \delta a_j b_j + a_j \delta b_j \end{pmatrix}. \end{aligned}$$

By also writing  $\begin{pmatrix} \delta b_n \\ 0 \end{pmatrix} = \begin{pmatrix} \delta b_n & b_n \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ , we find that

$$\begin{aligned} & \langle a_1[D, b_1], \dots, a_n[D, b_n] \rangle_f \\ &= \int_{\phi} (a_1 \ a_1 b_1) \begin{pmatrix} \delta b_1 & b_1 \\ 0 & -1 \end{pmatrix} \left( \prod_{j=2}^n \begin{pmatrix} a_j + \delta a_j & a_j b_j + \delta a_j b_j + a_j \delta b_j \\ \delta a_j & \delta a_j b_j + a_j \delta b_j \end{pmatrix} \begin{pmatrix} \delta b_j & b_j \\ 0 & -1 \end{pmatrix} \right) \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ &= \int_{\phi} (A_1 \ 0) \left( \prod_{j=2}^n \begin{pmatrix} A_j + \delta A_j & -A_j \\ \delta A_j & -A_j \end{pmatrix} \right) \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \end{aligned}$$

which concludes the proof.  $\square$

**Corollary 9.22** *If  $n \in \mathbb{N}$ ,  $A \in \Omega^1(\mathcal{A})$  and  $\omega := \pi_D(A) \in \Omega_D^1(\mathcal{A})$ , then*

$$\langle \omega, \dots, \omega \rangle_f = \int_{\phi} (A \ 0) \begin{pmatrix} A + \delta A & -A \\ \delta A & -A \end{pmatrix}^{n-1} \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (9.3.8)$$

Using (9.3.8), we obtain in particular

$$\begin{aligned} \langle \omega \rangle_f &= \int_{\phi_1} A, \\ \langle \omega, \omega \rangle_f &= \int_{\phi_2} A^2 + \int_{\phi_3} A \delta A, \\ \langle \omega, \omega, \omega \rangle_f &= \int_{\phi_3} A^3 + \int_{\phi_4} A \delta A A + \int_{\phi_5} A \delta A \delta A, \\ \langle \omega, \omega, \omega, \omega \rangle_f &= \int_{\phi_4} A^4 + \int_{\phi_5} (A^3 \delta A + A \delta A A^2) + \int_{\phi_6} A \delta A \delta A A + \int_{\phi_7} A \delta A \delta A \delta A. \end{aligned}$$

With (9.3.2) this implies that

$$\begin{aligned} S_b[\omega] - S_b[0] &= \int_{\phi_1} A + \frac{1}{2} \int_{\phi_2} A^2 + \int_{\phi_3} \left( \frac{1}{2} A \delta A + \frac{1}{3} A^3 \right) \\ &\quad + \int_{\phi_4} \left( \frac{1}{3} A \delta A A + \frac{1}{4} A^4 \right) + \dots, \end{aligned}$$

where the dots indicate terms of form degree 5 and higher. Using  $\phi_{2k-1} = \psi_{2k-1} + \frac{1}{2} B_0 \phi_{2k}$ , this becomes

$$\begin{aligned} S_b[\omega] - S_b[0] &= \int_{\psi_1} A + \frac{1}{2} \int_{\phi_2} (\delta A + A^2) + \int_{\psi_3} \left( \frac{1}{2} A \delta A + \frac{1}{3} A^3 \right) \\ &\quad + \frac{1}{4} \int_{\phi_4} \left( \delta A \delta A + \frac{2}{3} (\delta A A^2 + A \delta A A + A^2 \delta A) + A^4 \right) + \dots. \end{aligned}$$

Notice that, if  $\phi_4$  would be tracial, we would be able to identify the terms  $\delta AA^2$ ,  $A\delta AA$  and  $A^2\delta A$ , and thus obtain the Yang–Mills form  $F^2 = (\delta A + A^2)^2$ , under the fourth integral. In the general case, however, cyclic permutations under  $\int_\phi$  produce correction terms, of which we will need to keep track. Even though the corresponding analysis is rather involved, it is only based on the properties I and II of the bracket (cf. Lemma 9.17); see Note 13 on this chapter. We then have the following

**Theorem 9.23** *The spectral action fluctuated by  $\omega = \pi_D(A) \in \Omega_D^1(\mathcal{A})_{sa}$  with corresponding  $A \in \Omega^1(\mathcal{A})$  and  $F = \delta A + A^2 \in \Omega^2(\mathcal{A})$  can be expanded as*

$$S_b[\omega] - S_b[0] \sim \sum_{k=1}^{\infty} \left( \int_{\psi_{2k-1}} \text{cs}_{2k-1}(A) + \frac{1}{2k} \int_{\phi_{2k}} F^k \right).$$

where the Chern–Simons forms of degree  $2k - 1$  are defined by

$$\text{cs}_{2k-1}(A) := \int_0^1 A(F_t)^{k-1} dt, \quad (9.3.9)$$

where  $F_t = t\delta A + t^2 A^2$  is the curvature two-form of the (connection) one-form  $A_t = tA$ .

**Proof** See Note 13 on this chapter. □

**Example 9.24** For the first three Chern–Simons forms one easily derives the following explicit expressions:

$$\begin{aligned} \text{cs}_1(A) &= A; & \text{cs}_3(A) &= \frac{1}{2} \left( A\delta A + \frac{2}{3} A^3 \right); \\ \text{cs}_5(A) &= \frac{1}{3} \left( A(\delta A)^2 + \frac{3}{4} A\delta AA^2 + \frac{3}{4} A^3\delta A + \frac{3}{5} A^5 \right). \end{aligned}$$

## 9.A Divided Differences

We recall the definition of and some basic results on divided differences.

**Definition 9.25** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  and let  $x_0, x_1, \dots, x_n$  be distinct points in  $\mathbb{R}$ . The divided difference of order  $n$  is defined by the recursive relations

$$\begin{aligned} f[x_0] &= f(x_0), \\ f[x_0, x_1, \dots, x_n] &= \frac{f[x_1, \dots, x_n] - f[x_0, x_1, \dots, x_{n-1}]}{x_n - x_0}. \end{aligned}$$

On coinciding points we extend this definition as the usual derivative:

$$f[x_0, \dots, x, \dots, x, \dots, x_n] := \lim_{u \rightarrow 0} f[x_0, \dots, x + u, \dots, x, \dots, x_n].$$

Finally, as a shorthand notation, for an index set  $I = \{i_1, \dots, i_n\}$  we write

$$f[x_I] = f[x_{i_1}, \dots, x_{i_n}].$$

Also note the following useful representation:

**Proposition 9.26** *For any  $x_0, \dots, x_n \in \mathbb{R}$ ,*

$$f[x_0, x_1, \dots, x_n] = \int_{\Delta_n} f^{(n)}(s_0 x_0 + s_1 x_1 + \dots + s_n x_n) d^n s.$$

**Proof** See Note 15 on this chapter. □

**Exercise 9.1** Prove Proposition 9.26 and show that it implies

$$\sum_{i=0}^n f[x_0, \dots, x_i, x_i, \dots, x_n] = f'[x_0, x_1, \dots, x_n].$$

**Proposition 9.27** *For any  $x_1, \dots, x_n \in \mathbb{R}$  for  $f(x) = g(x^2)$  we have,*

$$f[x_0, \dots, x_n] = \sum_I \left( \prod_{\{i-1, i\} \subset I} (x_i + x_{i+1}) \right) g[x_I^2],$$

where the sum is over all ordered index sets  $I = \{0 = i_0 < i_1 < \dots < i_k = n\}$  such that  $i_j - i_{j-1} \leq 2$  for all  $1 \leq j \leq k$  (i.e. there are no gaps in  $I$  of length greater than 1).

**Proof** This follows from the chain rule for divided differences (see Note 16 on this chapter): if  $f = g \circ \phi$ , then

$$f[x_0, \dots, x_n] = \sum_{k=1}^n \sum_{0=i_0 < i_1 < \dots < i_k=n} g[\phi(x_{i_0}), \dots, \phi(x_{i_k})] \prod_{j=0}^{k-1} \phi[x_{i_j}, \dots, x_{i_{j+1}}].$$

For  $\phi(x) = x^2$  we have  $\phi[x, y] = x + y$ ,  $\phi[x, y, z] = 1$  and all higher divided differences are zero. Thus, if  $i_{j+1} - i_j > 2$  then  $\phi[x_{i_j}, \dots, x_{i_{j+1}}] = 0$ . In the remaining cases one has

$$\phi[x_{i_j}, \dots, x_{i_{j+1}}] = \begin{cases} x_{i_j} + x_{i_{j+1}} & \text{if } i_{j+1} - i_j = 1 \\ 1 & \text{if } i_{j+1} - i_j = 2, \end{cases}$$

and in the above summation this selects precisely the index sets  $I$ . □

**Example 9.28** For the first few terms, we have

$$\begin{aligned} f[x_0, x_1] &= (x_0 + x_1)g[x_0^2, x_1^2], \\ f[x_0, x_1, x_2] &= (x_0 + x_1)(x_1 + x_2)g[x_0^2, x_1^2, x_2^2] + g[x_0^2, x_2^2], \\ f[x_0, x_1, x_2, x_3] &= (x_0 + x_1)(x_1 + x_2)(x_2 + x_3)g[x_0^2, x_1^2, x_2^2, x_3^2] \\ &\quad + (x_2 + x_3)g[x_0^2, x_2^2, x_3^2] + (x_0 + x_1)g[x_0^2, x_1^2, x_3^2]. \end{aligned}$$

## Notes

### Section 9.1 Spectral action functional

1. The spectral action principle was introduced by Chamseddine and Connes in [1, 2].
2. The spectral action has also been computed for spectral triples that are not the product of  $M$  with a finite space  $F$ , and which are further off the ‘commutative shore’. These include the noncommutative torus [3], the Moyal plane [4, 5], the quantum group  $SU_q(2)$  [6] and the Podleś sphere  $S_q^2$  [7]. We also refer to the book [8].
3. Note that we have put two restrictions on the fermions in the fermionic action  $S_f$  of Definition 9.3. The first is that we restrict ourselves to even vectors in  $\mathcal{H}^+$ , instead of considering all vectors in  $\mathcal{H}$ . The second restriction is that we do not consider the inner product  $\langle J\bar{\psi}', D_\omega\bar{\psi} \rangle$  for two independent vectors  $\psi$  and  $\psi'$ , but instead use the same vector  $\psi$  on both sides of the inner product. Each of these restrictions reduces the number of degrees of freedom in the fermionic action by a factor of 2, yielding a factor of 4 in total. It is precisely this approach that solves the problem of fermion doubling pointed out in [9] (see also the discussion in [10, Chap. 1, Sect. 16.3]). We shall discuss this in more detail in Chaps. 11 and 13, where we calculate the fermionic action for electrodynamics and the Standard Model, respectively.

### Section 9.2 Asymptotic expansion of the spectral action

4. For a complete treatment of the Laplace–Stieltjes transform, see [11].
5. Lemma 9.6 appeared as [10, Lemma 1.144].
6. Corollary 9.8 is [10, Theorem 1.145]. An analysis of the term  $\text{Tr } |D_\omega|^{-z}|_{z=0}$  therein, including a perturbative expansion in powers of  $\omega$  has been obtained in [12].

### Section 9.3 Perturbative expansion in the gauge field

7. Section 9.3 is based on [13, 14].
8. The notation  $\langle X_0, \dots, X_n \rangle_{t,n}$  should not be confused with the zeta functions  $\langle X_0, \dots, X_n \rangle_z$  introduced in Chap. 6. However, they are related through the formula

$$\langle X_0, \dots, X_n \rangle_{t,n} = \frac{(-1)^p}{2\pi i} \text{Tr} \int e^{-t\lambda} X_0 (\lambda - D^2)^{-1} X_1 \cdots A^n (\lambda - D^2)^{-1} d\lambda.$$

Multiplying this expression by  $t^{z-1}$  and integrating over  $t$  eventually yields  $\langle X_0, \dots, X_n \rangle_z$ . For details, we refer to [15, Appendix A].

9. For more details on Gâteaux derivatives, we refer to [16]. For instance, that the Gâteaux derivative of a linear map  $F$  between Fréchet spaces is a linear map  $F'(x)(\cdot)$  for any  $x \in X$  is shown in [16, Theorem 3.2.5].
10. The expansion in Eq. 9.3.1 is asymptotic in the sense that the partial sums  $\sum_{n=0}^N \frac{1}{n!} S_b^{(n)}(0) (\omega, \dots, \omega)$  can be estimated to differ from  $S_b[\omega]$  by  $\mathcal{O}(\|\omega\|^{N+1})$ . This is made precise in [13] and further improved upon in [14, 17].
11. Theorem 9.14 was proved in [13]. A similar result was obtained in finite dimensions in [18] and in a different setting in [17]. Corollary 9.16 was obtained at first order for bounded operators [19].
12. There is a close connection between the spectral action, the Krein spectral shift function [20, 21], as well as the spectral flow of Atiyah and Lusztig [22–24]. One way to see this is from Theorem 9.11, where we can control the asymptotic expansion of the spectral action using the remainder terms  $R_k$ . In [17] these terms are analyzed and related to a spectral shift formula [20, 21] (see also the book [25] and the review [26], and references therein). In fact, under the assumption that  $f$  has compact support, the first rest term  $S_b[\omega] - S_b^{(0)}(0)$  becomes

$$\mathrm{Tr} f(D + \omega) - \mathrm{Tr} f(D) = \int_{\mathbb{R}} f(x) d(\mathrm{Tr} E_{D+\omega}(x)) - \int_{\mathbb{R}} f(x) d(\mathrm{Tr} E_D(x)),$$

where  $E_{D+\omega}$  and  $E_D$  are the spectral projections of  $D + \omega$  and  $D$ , respectively. After a partial integration, we then obtain [17, Theorem 3.9]

$$\mathrm{Tr} f(D + \omega) - \mathrm{Tr} f(D) = \int_{\mathbb{R}} f'(x) \xi(x) dx, \quad (*)$$

where

$$\xi(x) = \mathrm{Tr} (E_{D+\omega}(x) - E_D(x))$$

is the so-called *spectral shift function*. Moreover, it turns out that the higher-order rest terms are related to higher-order spectral shift functions [27, 28].

Let us also briefly describe the intriguing connection between the spectral shift function and the local index formula of Chap. 6. In fact, [29] (using a result from [30, Appendix B]) relates the index of  $PuP$  which appears in the odd local index formula (Theorem 6.21) to the *spectral flow*  $\mathrm{sf}(\{D_t\})$  of the family  $D_t = (1-t)D + tuDu^* = D + tu[D, u^*]$  for  $0 \leq t \leq 1$ . Roughly speaking, the spectral flow of such a family of operators is given by the net number of eigenvalues of  $D_t$  that pass through 0 in the positive direction when  $t$  runs from 0 to 1. One then has

$$\mathrm{index} \ PuP = \mathrm{sf}(\{D_t\}_{t \in [0,1]}).$$

The connection between spectral flow and the spectral shift function was first hinted at in [31] and has been worked out in [32, 33]. Essentially, these latter papers build on the observation that the spectral flow from  $D_0 - x$  to  $D_1 - x$  for any real number  $x$  is equal to the spectral shift function  $\xi(x)$  defined above in terms of the spectral projections of  $D_0$  and  $D_1$ . Note that for a path connecting  $D$  and the unitarily equivalent operator  $uD u^*$  the spectral shift function is a constant. In fact, since  $D$  and  $uD u^*$  have identical spectrum, the left-hand side of  $(*)$  vanishes. Integration by parts on the right-hand side then ensures that  $\xi$  is constant (and in fact equal to the above index).

Eventually, a careful analysis of the spectral flow [34] (and [35] for the even case) allows one to prove the local index formula in the much more general setting of semi-finite spectral triples [29, 36–38].

Another encounter of spectral shift and spectral flow is in the computation of the index of the operator  $d/dt + A(t)$  with  $A(t)$  a suitable family of perturbations ( $t \in \mathbb{R}$ ). In fact, they were the operators studied by Atiyah, Patodi and Singer in [22–24]. The index of  $d/dt + A(t)$  can be expressed in terms of the spectral flow of  $A(t)$  under the assumptions that  $A(\pm\infty)$  is boundedly invertible, and that  $A(t)$  has discrete spectrum for all  $t \in \mathbb{R}$ . We refer to [39] for a careful historical account, and the extension of this result to relatively trace class perturbations  $A(t)$ .

13. For full details on the proof of Theorem 9.23, including its convergence aspects, we refer to the original work [14].
14. The Chern–Simons terms defined in Theorem 9.23 have been studied in [40] and also appear in the physics literature, see [41, pp. 391ff] and [42, ω bis 10]. Usually, they are evaluated using a *trace* on  $\Omega^\bullet$ , which is defined as a continuous degree zero map from  $\Omega^\bullet$  to a complex  $\omega$  such that  $\delta\tau = \tau\delta$  and  $\tau$  vanishes on graded commutators  $(\text{ad}X)(Y) = XY - (-1)^{\text{odd}(Y)} YX$ . Given a connection one-form  $A$  with curvature  $F = \delta A + A^2$  we may define the Chern character form as

$$\text{ch}_{2n} = \tau(F^n).$$

By the Bianchi identity  $(\delta + \text{ad}A)F = \delta F + [A, F] = 0$  we have:

$$\delta\tau(F^n) = \tau(\delta F^n) = \tau(-[A, F^n]) = 0,$$

and one may wonder whether we can write  $\text{ch}_{2n}$  as an exact form. This is where the Chern–Simons form enters, since one may derive the following transgression formula:

$$\delta(\tau(\text{cs}_{2n-1}(A))) = \frac{1}{n}\tau(F^n).$$

It follows directly by integrating the homotopy formula for a family of connection one-forms  $A_t = tA$ :

$$\frac{1}{n}\partial_t(\tau(F_t^n)) = \delta(\tau(\dot{A}_t F_t^{n-1})) \quad (*)$$

Let us for completeness prove Eq. (\*) for a family of connection one-forms  $A_t$  with curvature  $F_t = \delta A_t + A_t^2$ . Since

$$\partial_t(F_t) = \delta\dot{A}_t + \dot{A}_t A_t + A_t \dot{A}_t = (\delta + \text{ad}A_t)(\dot{A}_t),$$

we find indeed that

$$\begin{aligned} \frac{1}{n}\partial_t\tau(F_t^n) &= \tau((\delta + \text{ad}A_t)(\dot{A}_t)F_t^{n-1}) \\ &= \tau((\delta + \text{ad}A_t)(\dot{A}_t F_t^{n-1})) && \text{(Bianchi identity)} \\ &= \tau(\delta(\dot{A}_t F_t^{n-1})) && \text{(trace on commutator)} \\ &= \delta\tau(\dot{A}_t F_t^{n-1}). \end{aligned}$$

## Section 9.A Divided differences

15. Proposition 9.26 is due to Hermite [43].
16. The chain rule for divided differences is proved in [44]. For Cauchy's formula for divided differences, we refer to [45, Chap. I.1].

## References

1. Chamseddine, A.H., Connes, A.: Universal formula for noncommutative geometry actions: unifications of gravity and the standard model. *Phys. Rev. Lett.* **77**, 4868–4871 (1996)
2. Chamseddine, A.H., Connes, A.: The spectral action principle. *Commun. Math. Phys.* **186**, 731–750 (1997)
3. Essouabri, D., Iochum, B., Levy, C., Sitarz, A.: Spectral action on noncommutative torus. *J. Noncommut. Geom.* **2**, 53–123 (2008)
4. Gayral, V., Iochum, B.: The spectral action for Moyal planes. *J. Math. Phys.* **46**, 043503, 17 (2005)
5. Grosse, H., Wulkenhaar, R.: 8D-spectral triple on 4D-Moyal space and the vacuum of non-commutative gauge theory. *J. Geom. Phys.* **62**, 1583–1599 (2012)
6. Iochum, B., Levy, C., Sitarz, A.: Spectral action on  $SU_q(2)$ . *Commun. Math. Phys.* **289**, 107–155 (2009)
7. Eckstein, M., Iochum, B., Sitarz, A.: Heat trace and spectral action on the standard Podles sphere. [arXiv:1307.4188](https://arxiv.org/abs/1307.4188)
8. Eckstein, M., Iochum, B.: Spectral Action in Noncommutative Geometry. Mathematical Physics Studies. Springer (2018)
9. Lizzi, F., Mangano, G., Miele, G., Sparano, G.: Fermion Hilbert space and fermion doubling in the noncommutative geometry approach to gauge theories. *Phys. Rev. D* **55**, 6357–6366 (1997)
10. Connes, A., Marcolli, M.: Noncommutative Geometry, Quantum Fields and Motives. AMS, Providence (2008)
11. Widder, D.V.: The Laplace Transform. Princeton Mathematical Series, vol. 6. Princeton University Press, Princeton, N.J. (1941)
12. Connes, A., Chamseddine, A.H.: Inner fluctuations of the spectral action. *J. Geom. Phys.* **57**, 1–21 (2006)
13. van Suijlekom, W.D.: Perturbations and operator trace functions. *J. Funct. Anal.* **260**, 2483–2496 (2011)
14. van Nuland, T.D.H., van Suijlekom, W.D.: Cyclic cocycles in the spectral action. *J. Noncommut. Geom.* (online 22 December 2021) [arXiv:2104.09899](https://arxiv.org/abs/2104.09899)
15. Higson, N.: The residue index theorem of Connes and Moscovici. In: Surveys in Noncommutative Geometry. Clay Mathematics Proceedings, vol. 6, pp. 71–126. American Mathematical Society, Providence, RI (2006)
16. Hamilton, R.S.: The inverse function theorem of Nash and Moser. *Bull. Amer. Math. Soc.* **7** (1982)
17. Skripka, A.: Asymptotic expansions for trace functionals. *J. Funct. Anal.* **266**, 2845–2866 (2014)
18. Hansen, F.: Trace functions as Laplace transforms. *J. Math. Phys.* **47** 043504, 11 (2006)
19. Gilliam, D.S., Hohage, T., Ji, X., Ruymgaart, F.: The Fréchet derivative of an analytic function of a bounded operator with some applications. *Int. J. Math. Math. Sci.* (2009) Article ID 239025, 17
20. Lifshits, I.: On a problem in perturbation theory (Russian). *Uspehi Matem. Nauk.* **7**, 171–180 (1952)

21. Krein, M.: On a trace formula in perturbation theory. Matem. Sbornik **33**, 597–626 (1953)
22. Atiyah, M.F., Patodi, V.K., Singer, I.M.: Spectral asymmetry and Riemannian geometry. I. Math. Proc. Cambridge Philos. Soc. **77**, 43–69 (1975)
23. Atiyah, M.F., Patodi, V.K., Singer, I.M.: Spectral asymmetry and Riemannian geometry. II. Math. Proc. Cambridge Philos. Soc. **78**, 405–432 (1975)
24. Atiyah, M.F., Patodi, V.K., Singer, I.M.: Spectral asymmetry and Riemannian geometry. III. Math. Proc. Cambridge Philos. Soc. **79**, 71–99 (1976)
25. Yafaev, D.R.: Mathematical Scattering Theory. Translations of Mathematical Monographs, vol. 105. American Mathematical Society, Providence, RI (1992) General Theory, Translated from the Russian by J. R. Schulenberger
26. M. S. Birman and A. B. Pushnitski. Spectral shift function, amazing and multifaceted. *Integral Equations Operator Theory* 30 (1998), pp. 191–199. Dedicated to the memory of Mark Grigorievich Krein (1907–1989)
27. Koplienko, L.S.: The trace formula for perturbations of nonnuclear type. Sibirsk. Mat. Zh. **25**, 62–71 (1984)
28. Potapov, D., Skripka, A., Sukochev, F.: Spectral shift function of higher order. Invent. Math. **193**, 501–538 (2013)
29. Carey, A., Phillips, J.: Unbounded Fredholm modules and spectral flow. Canad. J. Math. **50**, 673–718 (1998)
30. Phillips, J., Raeburn, I.: An index theorem for Toeplitz operators with noncommutative symbol space. J. Funct. Anal. **120**, 239–263 (1994)
31. Müller, W.: Relative zeta functions, relative determinants and scattering theory. Commun. Math. Phys. **192**, 309–347 (1998)
32. Azamov, N.A., Carey, A.L., Sukochev, F.A.: The spectral shift function and spectral flow. Commun. Math. Phys. **276**, 51–91 (2007)
33. Azamov, N.A., Carey, A.L., Dodds, P.G., Sukochev, F.A.: Operator integrals, spectral shift, and spectral flow. Canad. J. Math. **61**, 241–263 (2009)
34. Carey, A.L., Phillips, J., Rennie, A., Sukochev, F.A.: The local index formula in semifinite von Neumann algebras. I. Spectral flow. Adv. Math. **202**, 451–516 (2006)
35. Carey, A.L., Phillips, J., Rennie, A., Sukochev, F.A.: The local index formula in semifinite von Neumann algebras. II. The even case. Adv. Math. **202**, 517–554 (2006)
36. Benameur, M.-T., Fack, T.: Type II non-commutative geometry. I. Dixmier trace in von Neumann algebras. Adv. Math. **199**, 29–87 (2006)
37. Carey, A.L., Phillips, J., Sukochev, F.A.: On unbounded  $p$ -summable Fredholm modules. Adv. Math. **151**, 140–163 (2000)
38. Carey, A., Phillips, J., Sukochev, F.: Spectral flow and Dixmier traces. Adv. Math. **173**, 68–113 (2003)
39. Gesztesy, F., Latushkin, Y., Makarov, K.A., Sukochev, F., Tomilov, Y.: The index formula and the spectral shift function for relatively trace class perturbations. Adv. Math. **227**, 319–420 (2011)
40. Quillen, D.: Chern-Simons forms and cyclic cohomology. In: The Interface of Mathematics and Particle Physics. Institute of Mathematics and its Applications Conference Series, vol. 24, pp. 117–134. Oxford (1988). Oxford University Press, New York (1990)
41. Choquet-Bruhat, Y., DeWitt-Morette, C., Dillard-Bleick, M.: Analysis, Manifolds and Physics. North-Holland Publishing Co., Amsterdam-New York-Oxford (1977)
42. Choquet-Bruhat, Y., DeWitt-Morette, C.: Analysis, Manifolds and Physics. Part II. North-Holland Publishing Co., Amsterdam (1989). 92 applications
43. Hermite, C.: Sur la formule d'interpolation de lagrange. J. Reine Angew. Math. **84**, 70–79 (1878)
44. Floater, M.S., Lyche, T.: Two chain rules for divided differences and Faà di Bruno's formula. Math. Comp. **76**, 867–877 (2007)
45. Donoghue, Jr., W.F.: Monotone Matrix Functions and Analytic Continuation. Springer, New York (1974). Die Grundlehren der mathematischen Wissenschaften, Band 207

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 10

## Almost-Commutative Manifolds and Gauge Theories



In this chapter we analyze the gauge theories corresponding (in the sense of Chap. 7) to a special class of noncommutative manifolds, to wit *almost-commutative*, or AC manifolds. We will see that this class leads to the usual gauge theories in physics. After identifying the gauge group, the gauge fields and the scalar fields, we compute the spectral action that yields the Lagrangian of physical interest.

### 10.1 Gauge Symmetries of AC Manifolds

We consider almost-commutative manifolds  $M \times F$  that are the products of a Riemannian spin manifold  $M$  with a finite noncommutative space  $F$ .

As such, these are reminiscent of the original Kaluza–Klein theories where one considers the product  $M \times \mathbb{S}^1$ . The crucial difference is that the space  $F$  is *finite* so that no extra dimensions appear, while it can have non-trivial (noncommutative) structure.

**Definition 10.1** Let  $M$  be a Riemannian spin manifold with canonical triple  $(C^\infty(M), L^2(S), D_M; J_M, \gamma_M)$ , and let  $(A_F, H_F, D_F; J_F, \gamma_F)$  be a finite real spectral triple. The *almost-commutative manifold*  $M \times F$  is given by the real spectral triple:

$$M \times F = (C^\infty(M, A_F), L^2(S \otimes (M \times H_F)), D_M \otimes 1 + \gamma_M \otimes D_F; J_M \otimes J_F, \gamma_M \otimes \gamma_F).$$

Recall the definition of the gauge group of a real spectral triple (cf. Definition 7.4). In the case of AC manifolds, it is given by

$$\mathfrak{G}(M \times F) := \{u Ju^{-1} : u \in C^\infty(M, \mathcal{U}(A_F))\},$$

with  $J = J_M \otimes J_F$ . Here we have identified  $\mathcal{U}(C^\infty(M, A_F)) = C^\infty(M, \mathcal{U}(A_F))$ . For the Lie algebra of the gauge group we have

$$\mathfrak{g}(M \times F) := \{X + JXJ^{-1} : X \in C^\infty(M, \mathfrak{u}(A_F))\}.$$

In the same way, we also obtain the groups  $\mathfrak{G}(M)$  and  $\mathfrak{G}(F)$ . For the canonical triple on the spin manifold  $M$ , we have seen in Example 8.2 that  $C^\infty(M)_{J_M} = C^\infty(M)$ , which means that the group  $\mathfrak{G}(M)$  is just the trivial group. For the finite space  $F$ , we obtain the *local* gauge group  $\mathfrak{G}(F)$ . Let us have a closer look at the structure of this local gauge group. We define two subsets of  $A_F$  by

$$\mathfrak{H}(F) := \mathcal{U}((A_F)_{J_F}), \quad (10.1.1a)$$

$$\mathfrak{h}(F) := \mathfrak{u}((A_F)_{J_F}). \quad (10.1.1b)$$

Note that the group  $\mathfrak{H}(F)$  is the counterpart for the finite space  $F$  of the group  $\mathcal{U}(\mathcal{A}_J)$  in Proposition 7.5, and  $\mathfrak{h}(F)$  is its Lie algebra.

**Proposition 10.2** *Let  $M$  be simply connected. Then the gauge group  $\mathfrak{G}(M \times F)$  of an almost-commutative manifold is given by  $C^\infty(M, \mathfrak{G}(F))$ , where  $\mathfrak{G}(F) = \mathcal{U}(A_F)/\mathfrak{H}(F)$  is the gauge group of the finite space. Consequently, the gauge Lie algebra  $\mathfrak{g}(M \times F)$  is given by  $C^\infty(M, \mathfrak{g}(F))$ , where  $\mathfrak{g}(F) = \mathfrak{u}(A_F)/\mathfrak{h}(F)$ .*

**Proof** This follows from Propositions 7.5 and 7.8, combined with the fact that for the algebra  $\mathcal{A} = C^\infty(M, A_F)$  we have  $\mathcal{U}(\mathcal{A}) \simeq C^\infty(M, \mathcal{U}(A_F))$ , while  $\mathcal{U}(\mathcal{A}_J) = C^\infty(M, \mathfrak{H}(F))$ . The quotient of the latter two groups is isomorphic to  $C^\infty(M, \mathfrak{G}(F))$  if the following homomorphism

$$C^\infty(M, \mathcal{U}(A_F)) \rightarrow C^\infty(M, \mathcal{U}(A_F)/\mathfrak{H}(F))$$

is surjective. This happens when  $M$  is simply connected, as in that case there exists a global lift from  $\mathcal{U}(A_F)/\mathfrak{H}(F)$  to  $\mathcal{U}(A_F)$  (see Note 4 of this chapter below).  $\square$

This is in concordance with the picture derived in Chap. 8, where the gauge group acts fiberwise on a  $C^*$ -bundle. Namely, in the case of an almost-commutative manifold we have a globally trivial  $C^*$ -bundle  $M \times A_F$  for which  $\mathcal{A}$  are the (smooth) sections. Since  $\mathfrak{G}(M \times F) \simeq C^\infty(M, \mathfrak{G}(F))$ , the gauge group is given by sections of the group bundle  $M \times \mathfrak{G}(F)$ , which then naturally acts fiberwise on the  $C^*$ -bundle  $M \times A_F$ .

Combined with the *outer* automorphisms on  $C^\infty(M)$ , we arrive at the full symmetry group of an almost-commutative manifold  $M \times F$  as a semi-direct product, where the ‘internal symmetries’ are given by the gauge group  $\mathfrak{G}(M \times F)$ . Furthermore, we also still have invariance under the group of diffeomorphisms  $\text{Diff}(M)$ , as in Example 7.2. There exists a group homomorphism  $\theta: \text{Diff}(M) \rightarrow \text{Aut}(\mathfrak{G}(M \times F))$  given by

$$\theta(\phi)U := U \circ \phi^{-1},$$

for  $\phi \in \text{Diff}(M)$  and  $U \in \mathfrak{G}(M \times F)$ . Hence, we can describe the *full symmetry group* by the semi-direct product

$$\mathfrak{G}(M \times F) \rtimes \text{Diff}(M).$$

### 10.1.1 Unimodularity

Suppose that  $A_F$  is a complex unital  $*$ -algebra, conform Definition 2.1. This algebra has a unit 1, and by complex linearity we see that  $\mathbb{C}1 \subset (A_F)_{J_F}$ . Restricting to unitary elements, we then find that  $U(1)$  is a subgroup of  $\mathfrak{H}(F)$ . Because  $\mathfrak{H}(F)$  is commutative,  $U(1)$  is then automatically a normal subgroup of  $\mathfrak{H}(F)$ .

If, on the other hand,  $A_F$  is a real algebra, we can only say that  $\mathbb{R}1 \subset (A_F)_{J_F}$ . Restricting to unitary (i.e. in this case orthogonal) elements, we then only obtain the insight that  $\{1, -1\}$  is a normal subgroup of  $\mathfrak{H}(F)$ .

**Proposition 10.3** *If  $A_F$  is a complex algebra, the gauge group is isomorphic to*

$$\mathfrak{G}(F) \cong S\mathcal{U}(A_F)/S\mathfrak{H}(F),$$

where

$$\begin{aligned} S\mathcal{U}(A_F) &:= \{g \in \mathcal{U}(A_F) \mid \det_{H_F} g = 1\}, \\ S\mathfrak{H}(F) &:= S\mathcal{U}(A_F) \cap \mathfrak{H}(F). \end{aligned}$$

In this case the gauge algebra is

$$\mathfrak{g}(F) \cong su(A_F)/s\mathfrak{h}(F),$$

with

$$\begin{aligned} su(A_F) &:= \{X \in \mathfrak{u}(A_F) \mid \text{Tr}_{H_F} X = 0\}, \\ s\mathfrak{h}(F) &:= su(A_F) \cap h_F. \end{aligned}$$

**Proof** Elements of the quotient  $\mathfrak{G}(F) = \mathcal{U}(A_F)/\mathfrak{H}(F)$  are given by the equivalence classes  $[u]$  for  $u \in \mathcal{U}(A_F)$ , subject to the equivalence relation  $[u] = [uh]$  for all  $h \in \mathfrak{H}(F)$ . Similarly, the quotient  $S\mathcal{U}(A_F)/S\mathfrak{H}(F)$  consists of classes  $[v]$  for  $v \in S\mathcal{U}(A_F)$ , with the equivalence relation  $[v] = [vg]$  for all  $g \in S\mathfrak{H}(F)$ . We first show that this quotient is well defined, i.e. that  $S\mathfrak{H}(F)$  is a normal subgroup of  $S\mathcal{U}(A_F)$ . For this we need to check that  $vgv^{-1} \in S\mathfrak{H}(F)$  for all  $v \in S\mathcal{U}(A_F)$  and  $g \in S\mathfrak{H}(F)$ . We already know that  $vgv^{-1} \in \mathfrak{H}(F)$ , because  $\mathfrak{H}(F)$  is a normal subgroup of  $\mathcal{U}(A_F)$ . We then also see that  $\det_{H_F}(vgv^{-1}) = \det_{H_F} g = 1$ , so  $vgv^{-1} \in S\mathfrak{H}(F)$ , and the quotient  $S\mathcal{U}(A_F)/S\mathfrak{H}(F)$  is indeed well defined.

As to for the claimed isomorphism, consider the map  $\varphi : \mathcal{U}(A_F) \rightarrow S\mathcal{U}(A_F)/S\mathfrak{H}(F)$  given by

$$\varphi(u) = [\lambda_u^{-1}u],$$

where  $\lambda_u \in U(1)$  is an element in  $U(1)$  such that  $\lambda_u^N = \det u$ , where  $N$  is the dimension of the finite-dimensional Hilbert space  $H_F$ .

Since  $U(1)$  is a subgroup of  $\mathcal{U}(A_F)$  (because we assume  $A_F$  to be a complex algebra), we see that indeed  $\lambda_u^{-1}u \in S\mathcal{U}(A_F)$ . Let us also check that  $\varphi$  does not depend on the choice of the  $N$ 'th root  $\lambda_u$  of  $\det u$  we take. Suppose  $\lambda'_u$  is such that  $\lambda'^N_u = \det u$ . We then must have  $\lambda_u^{-1}\lambda'_u \in \mu_N$ , where  $\mu_N$  is the multiplicative group of the  $N$ 'th roots of unity. Since  $U(1)$  is a subgroup of  $\mathfrak{H}(F)$ , we see that  $\mu_N$  is a subgroup of  $S\mathfrak{H}(F)$ , so  $[\lambda_u^{-1}u] = [\lambda'^{-1}_u u]$ , and hence the image of  $\varphi$  is indeed independent of the choice of  $\lambda_u$ .

Next, since  $S\mathcal{U}(A_F) \subset \mathcal{U}(A_F)$ , the homomorphism  $\varphi$  is clearly surjective. We determine its kernel:

$$\ker \varphi = \{u \in \mathcal{U}(A_F) : \lambda_u^{-1}u \in \mathfrak{H}(F)\} \simeq \{u \in \mathcal{U}(A_F) : u \in \mathfrak{H}(F)\} \equiv \mathfrak{H}(F),$$

since  $\lambda_u \in \mathfrak{H}(F)$ . □

The significance of Proposition 10.3 is that in the case of a complex algebra with a complex representation, equivalence classes of the quotient  $\mathfrak{G}(F) = \mathcal{U}(A_F)/\mathfrak{H}(F)$  can always be represented (though not uniquely) by elements of  $S\mathcal{U}(A_F)$ . In that sense, all elements  $g \in \mathfrak{G}(F)$  naturally satisfy the so-called **unimodularity condition**, i.e. they satisfy

$$\det_{H_F} g = 1.$$

In the case of an algebra with a real representation, this is not true and it is natural to impose the unimodularity condition for such representations by hand. We will see later in Chap. 13 how this works in the derivation of the Standard Model from noncommutative geometry.

**Example 10.4** Define the so-called **Yang–Mills finite spectral triple** (cf. Example 3.14)

$$F_{YM} = (M_N(\mathbb{C}), M_N(\mathbb{C}), D = 0; J_F = (\cdot)^*, \gamma_F = 1).$$

One easily checks that the commutative subalgebra  $(A_F)_{J_F}$  is given by  $\mathbb{C}\mathbb{I}_N$ . The group  $\mathfrak{H}(F)$  of unitary elements of this subalgebra is then equal to the group  $U(1)\mathbb{I}_N$ . Note that in this case  $\mathfrak{H}(F)$  is equal to the subgroup  $\mathcal{U}(Z(A_F))$  of  $U(N)$  that commutes with the algebra  $M_N(\mathbb{C})$ . We thus obtain that the gauge group is given by the quotient  $\mathfrak{G}(F_{YM}) = U(N)/U(1) =: PU(N)$ , which by Example 7.3 is equal to the group of inner automorphisms of  $M_N(\mathbb{C})$ . As in Proposition 10.3, this group can also be written as  $SU(N)/\mu_N$ , where the multiplicative group  $\mu_N$  of  $N$ 'th roots of unity is the center of  $SU(N)$ . The Lie algebra  $\mathfrak{g}(F_{YM})$  consists of the traceless anti-hermitian matrices, i.e. it is  $su(N)$ .

The almost-commutative manifold  $M \times F_{YM}$  will be referred to as the **Yang–Mills manifold**. By Proposition 10.2, in the simply connected case the global gauge group  $\mathfrak{G}(M \times F_{YM})$  is given by maps  $C^\infty(M, PU(N))$ , or, equivalently, by the space of smooth sections of the trivial group bundle  $M \times PU(N)$ .

**Exercise 10.1** In the context of the above example, check that indeed:

- (1) the commutative subalgebra  $M_N(\mathbb{C})_{J_F} \simeq \mathbb{C}\mathbb{I}_N$ ,
- (2)  $S\mathfrak{H}(F) = \mu_N$ , the multiplicative group of  $N$ 'th roots of unity.

Explain the difference with the case of  $M_N(\mathbb{R})$ .

## 10.2 Gauge Fields and Scalar Fields

Let us apply the discussion in Sect. 7.2 on Morita self-equivalences to the almost-commutative manifold  $M \times F$  and see what the corresponding gauge fields look like. For convenience, we restrict ourselves to simply connected manifolds  $M$  of dimension  $\dim M = 4$  and  $F$  of even KO-dimension so that  $\epsilon'_F = 1$  in Table 3.1; this is sufficient for the physical applications later on.

Thus, we determine  $\Omega_D^1(\mathcal{A})$  for almost-commutative manifolds, much as in Exercise 5.1. The Dirac operator  $D = D_M \otimes 1 + \gamma_M \otimes D_F$  consists of two terms, and hence we can also split the inner fluctuation  $\omega = a[D, b]$  into two terms. The first term is given by

$$a[D_M \otimes 1, b] = -i\gamma^\mu \otimes a\partial_\mu b =: \gamma^\mu \otimes A_\mu, \quad (10.2.1)$$

where  $A_\mu := -ia\partial_\mu b \in i\mathcal{A}$  must be hermitian.<sup>1</sup> The second term yields

$$a[\gamma_M \otimes D_F, b] = \gamma_M \otimes a[D_F, b] =: \gamma_M \otimes \phi, \quad (10.2.2)$$

for hermitian  $\phi := a[D_F, b]$ . Thus, the inner fluctuations of an even almost-commutative manifold  $M \times F$  take the form

$$\omega = \gamma^\mu \otimes A_\mu + \gamma_M \otimes \phi, \quad (10.2.3)$$

for certain hermitian operators  $A_\mu \in i\mathcal{A}$  and  $\phi \in \Gamma(\text{End}(V))$ , where  $V$  is the trivial vector bundle  $V = M \times H_F$ .

The ‘fluctuated’ Dirac operator is given by  $D_\omega = D + \omega + \epsilon' J \omega J^{-1}$  (cf. Sect. 7.2.2 above), for which we calculate

$$\gamma^\mu \otimes A_\mu + \epsilon' J \gamma^\mu \otimes A_\mu J^{-1} = \gamma^\mu \otimes (A_\mu - J_F A_\mu J_F^{-1}) =: \gamma^\mu \otimes B_\mu, \quad (10.2.4)$$

---

<sup>1</sup> Note that  $i\mathcal{A} = \mathcal{A}$  for complex algebras only.

which defines  $B_\mu \in \Gamma(\text{End}(V))$ , and where we have used that  $J_M \gamma^\mu J_M^{-1} = -\gamma^\mu$  in dimension 4. Note that if  $\nabla^E$  denotes the twisted connection on the tensor product bundle  $E := S \otimes V$ , i.e.

$$\nabla_\mu^E = \nabla_\mu^S \otimes 1 + i1 \otimes B_\mu,$$

we see that we can rewrite

$$D_M \otimes 1 + \gamma^\mu \otimes B_\mu = -i\gamma^\mu \nabla_\mu^E.$$

For the remainder of the fluctuated Dirac operator, we define  $\Phi \in \Gamma(\text{End}(E))$  by

$$\Phi := D_F + \phi + J_F \phi J_F^{-1}. \quad (10.2.5)$$

The fluctuated Dirac operator of a real even AC-manifold then takes the form

$$D_\omega = D_M \otimes 1 + \gamma^\mu \otimes B_\mu + \gamma_M \otimes \Phi = -i\gamma^\mu \nabla_\mu^E + \gamma_M \otimes \Phi. \quad (10.2.6)$$

In Sect. 10.1 we obtained the local gauge group  $\mathfrak{G}(F)$  with Lie algebra  $\mathfrak{g}(F)$ . For consistency we should now check that the gauge field  $A_\mu$  arising from the inner fluctuation indeed corresponds to this same gauge group.

The requirement that  $A_\mu$  is hermitian is equivalent to  $(iA_\mu)^* = -iA_\mu$ . Since  $A_\mu$  is of the form  $-ia\partial_\mu b$  for  $a, b \in \mathcal{A}$  (see (10.2.1)), we see that  $iA_\mu$  is an element of the algebra  $\mathcal{A}$  (also if  $\mathcal{A}$  is only a real algebra). Thus we have  $A_\mu(x) \in i\mathfrak{u}(A_F)$ .

The only way in which  $A_\mu$  appears in  $D_\omega$  is through the action of  $A_\mu - J_F A_\mu J_F^{-1}$ . If we take  $A'_\mu = A_\mu - a_\mu$  for some  $a_\mu \in i\mathfrak{h}(F) = i\mathfrak{u}((A_F)_{J_F})$  (which commutes with  $J_F$ ), we see that  $A'_\mu - J_F A'_\mu J_F^{-1} = A_\mu - J_F A_\mu J_F^{-1}$ . Therefore we may without any loss of generality assume that  $A_\mu(x)$  is an element of the quotient  $i\mathfrak{g}(F) = i(\mathfrak{u}(A_F)/\mathfrak{h}(F))$ . Since  $\mathfrak{g}(F)$  is the Lie algebra of the gauge group  $\mathfrak{G}(F)$ , we have therefore confirmed that

$$A_\mu \in C^\infty(M, i\mathfrak{g}(F)) \quad (10.2.7)$$

is indeed a gauge field for the local gauge group  $\mathfrak{G}(F)$ . For the field  $B_\mu$  found in (10.2.6), we can also write

$$B_\mu = \text{ad}(A_\mu) := A_\mu - J_F A_\mu J_F^{-1}.$$

So, we conclude that  $B_\mu$  is given by the adjoint action of a gauge field  $A_\mu$  for the gauge group  $\mathfrak{G}(F)$  with Lie algebra  $\mathfrak{g}(F)$ .

If the finite noncommutative space  $F$  has a grading  $\gamma_F$ , the field  $\phi$  satisfies  $\phi\gamma_F = -\gamma_F\phi$  and the field  $\Phi$  satisfies  $\Phi\gamma_F = -\gamma_F\Phi$  and  $\Phi J_F = J_F\Phi$ . These relations follow directly from the definitions of  $\phi$  and  $\Phi$  and the commutation relations for  $D_F$  according to Definition 3.1.

Using the cyclic property of the trace, it is easy to see that the traces of the fields  $B_\mu$ ,  $\phi$  and  $\Phi$  over the finite-dimensional Hilbert space  $H_F$  vanish identically: for  $B_\mu$  we find

$$\mathrm{Tr}_{H_F}(B_\mu) = \mathrm{Tr}_{H_F}(A_\mu - J_F A_\mu J_F^{-1}) = \mathrm{Tr}_{H_F}(A_\mu - A_\mu J_F^{-1} J_F) = 0,$$

whereas for the field  $\phi$  we find

$$\mathrm{Tr}_{H_F}(\phi) = \mathrm{Tr}_{H_F}(a[D_F, b]) = \mathrm{Tr}_{H_F}([b, a]D_F).$$

Since the grading commutes with the elements in the algebra and anti-commutes with the Dirac operator, it follows that this latter trace also vanishes. It then automatically follows that  $\Phi = D_F + \phi + J_F \phi J_F^{-1}$  is traceless too.

**Example 10.5** For the Yang–Mills manifold  $M \times F_{\text{YM}}$  of Example 10.4 the inner fluctuations take the form  $\omega = \gamma^\mu \otimes A_\mu$  for some traceless hermitian field  $A_\mu = A_\mu^* \in C^\infty(M, \text{is}u(N))$ . Since  $J_F A_\mu J_F^{-1} m = mA_\mu$  for  $m \in M_N(\mathbb{C})$ , we see that for the field  $B_\mu = A_\mu - J_F A_\mu J_F^{-1}$  we obtain the action

$$m \mapsto B_\mu m = A_\mu m - mA_\mu = [A_\mu, m] = (\mathrm{ad} A_\mu)m.$$

Thus  $A_\mu$  is a  $PU(N)$  gauge field which acts on the fermions in  $L^2(S) \otimes M_N(\mathbb{C})$  in the adjoint representation.

### 10.2.1 Gauge Transformations

Recall from Sect. 7.2 that an element  $U \in \mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$  acts on the inner fluctuations as a gauge transformation. In fact, the rule  $D_\omega \mapsto UD_\omega U^*$  with  $U = uJuJ^{-1}$  can be implemented by

$$u : \omega \mapsto \omega^u := u\omega u^* + u[D, u^*], \quad (10.2.8)$$

so that  $UD_\omega U^* = D_{\omega^u}$ . In physics, the resulting transformation on the inner fluctuation  $\omega \mapsto \omega^u$  will be interpreted as a gauge transformation of the gauge field.

Note that for an element  $U = uJuJ^{-1}$  in the gauge group  $\mathfrak{G}(M \times F)$ , there is an ambiguity in the corresponding transformation of  $\omega$ . Namely, for  $u \in \mathcal{U}(\mathcal{A})$  and  $h \in \mathcal{U}(\mathcal{A}_J)$ , we can also write  $U = uhJuuhJ^{-1}$ . Replacing  $u$  with  $uh$  using (5.2.1) we then obtain

$$\omega^{uh} = u\omega u^* + u[D, u^*] + h[D, h^*].$$

However, when considering the total inner fluctuation  $\omega^{uh} + J\omega^{uh}J^{-1}$ , the extra term  $h[D, h^*]$  cancels out:

$$h[D, h^*] + Jh[D, h^*]J^{-1} = h[D, h^*] + [D, h]h^* = [D, hh^*] = 0.$$

Hence the transformation of  $D_\omega = D + \omega + J\omega J^{-1}$  is well defined.

For an AC-manifold  $M \times F$ , by (10.2.3) we have  $\omega = \gamma^\mu \otimes A_\mu + \gamma_M \otimes \phi$  and  $D = -i\gamma^\mu \nabla_\mu^S \otimes 1 + \gamma_M \otimes D_F$ , and, using  $[\nabla_\mu^S, u^*] = \partial_\mu u^*$ , we thus obtain

$$\begin{aligned} A_\mu &\rightarrow uA_\mu u^* - iu\partial_\mu u^*, \\ \phi &\rightarrow u\phi u^* + u[D_F, u^*]. \end{aligned} \tag{10.2.9}$$

The first equation is precisely the gauge transformation for a gauge field  $A_\mu \in C^\infty(M, i\mathfrak{g}(F))$ , as desired. However, the transformation property of the field  $\phi$  is a bit surprising. In the Standard Model, the Higgs field is in the defining representation of the gauge group. The transformation for  $\phi$  derived above, on the other hand, is in the adjoint representation. From the framework of noncommutative geometry this is no surprise, since both bosonic fields  $A_\mu$  and  $\phi$  are obtained from the inner fluctuations of the Dirac operator, and are thereby expected to transform in a similar manner. Fortunately, for particular choices of the finite space  $F$ , the adjoint transformation property of  $\phi$  reduces to that of the defining representation. The key example of this will be discussed in Chap. 13, where we present the derivation of the Standard Model from an almost-commutative manifold.

## 10.3 The Heat Expansion of the Spectral Action

In the remainder of this chapter we shall derive an explicit formula for the bosonic Lagrangian of an almost-commutative manifold  $M \times F$  from the spectral action of Definition 9.1. We start by calculating a generalized Lichnerowicz formula for the square of the fluctuated Dirac operator. Subsequently, we show how we can use this formula to obtain an asymptotic expansion of the spectral action in the form of (9.2.1). We explicitly calculate the coefficients in this **heat kernel expansion**, allowing for a derivation of the general form of the Lagrangian for an almost-commutative manifold.

### 10.3.1 A Generalized Lichnerowicz Formula

Suppose we have a vector bundle  $E \rightarrow M$ . We say that a second-order differential operator  $H$  is a *generalized Laplacian* if it is of the form  $H = \Delta^E - F$ , where  $\Delta^E$  is a Laplacian in the sense of Definition 4.16 and  $F \in \Gamma(\text{End}(E))$ .

Our first task is to show that the fluctuated Dirac operator  $D_\omega$  on an almost-commutative manifold squares to a generalized Laplacian,  $D_\omega^2 = \Delta^E - F$ , and then determine  $F$ . Before we prove this, let us first have a closer look at some explicit

formulas for the fluctuated Dirac operator. Recall from (10.2.6) that we can write

$$D_\omega = -i\gamma^\mu \nabla_\mu^E + \gamma_M \otimes \Phi$$

for the connection  $\nabla_\mu^E = \nabla_\mu^S \otimes 1 + 1 \otimes (\partial_\mu + iB_\mu)$  on  $E = S \otimes V$ , and the scalar field  $\Phi \in \Gamma(\text{End}(E))$ . Let us evaluate the relations between the connection, its curvature and their adjoint actions. We define the operator  $D_\mu$  as the adjoint action of the connection  $\nabla_\mu^E$ , i.e.  $D_\mu = \text{ad}(\nabla_\mu^E)$ . In other words, we have

$$D_\mu \Phi = [\nabla_\mu^E, \Phi] = \partial_\mu \Phi + i[B_\mu, \Phi]. \quad (10.3.1)$$

We define the curvature  $F_{\mu\nu}$  of the gauge field  $B_\mu$  as usual by

$$F_{\mu\nu} := \partial_\mu B_\nu - \partial_\nu B_\mu + i[B_\mu, B_\nu]. \quad (10.3.2)$$

Recall the curvature of the connection  $\nabla^E$  from (4.2.3). Since in local coordinates we have  $[\partial_\mu, \partial_\nu] = 0$ , we find

$$\begin{aligned} \Omega_{\mu\nu}^E &= \nabla_\mu^E \nabla_\nu^E - \nabla_\nu^E \nabla_\mu^E \\ &= (\nabla_\mu^S \otimes 1 + i1 \otimes B_\mu)(\nabla_\nu^S \otimes 1 + i1 \otimes B_\nu) \\ &\quad - (\nabla_\nu^S \otimes 1 + i1 \otimes B_\nu)(\nabla_\mu^S \otimes 1 + i1 \otimes B_\mu) \\ &= \Omega_{\mu\nu}^S \otimes 1 + i1 \otimes \partial_\mu B_\nu - i1 \otimes \partial_\nu B_\mu - 1 \otimes [B_\mu, B_\nu]. \end{aligned}$$

Inserting (10.3.2), we obtain the formula

$$\Omega_{\mu\nu}^E = [\nabla_\mu^E, \nabla_\nu^E] = \Omega_{\mu\nu}^S \otimes 1 + i1 \otimes F_{\mu\nu}. \quad (10.3.3)$$

Next, let us have a look at the commutator  $[D_\mu, D_\nu]$ . Using the definition of  $D_\mu$  and the Jacobi identity, we obtain

$$\begin{aligned} [D_\mu, D_\nu]\Phi &= \text{ad}(\nabla_\mu^E)\text{ad}(\nabla_\nu^E)\Phi - \text{ad}(\nabla_\nu^E)\text{ad}(\nabla_\mu^E)\Phi \\ &= [\nabla_\mu^E, [\nabla_\nu^E, \Phi]] - [\nabla_\nu^E, [\nabla_\mu^E, \Phi]] \\ &= [[\nabla_\mu^E, \nabla_\nu^E], \Phi] = [\Omega_{\mu\nu}^E, \Phi] = \text{ad}(\Omega_{\mu\nu}^E)\Phi. \end{aligned}$$

Since  $\Omega_{\mu\nu}^S$  commutes with  $\Phi$ , we obtain the relation

$$[D_\mu, D_\nu] = i \text{ad}(F_{\mu\nu}).$$

Note that this relation simply reflects the fact that  $\text{ad} : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g})$  is a Lie algebra homomorphism.

In local coordinates, the Laplacian is given by

$$\Delta^E = -g^{\mu\nu} (\nabla_\mu^E \nabla_\nu^E - \Gamma_{\mu\nu}^\rho \nabla_\rho^E).$$

We can then calculate the explicit formula

$$\begin{aligned} \Delta^E &= -g^{\mu\nu} (\nabla_\mu^E \nabla_\nu^E - \Gamma_{\mu\nu}^\rho \nabla_\rho^E) \\ &= \Delta^S \otimes 1 - g^{\mu\nu} \left( i(\nabla_\mu^S \otimes 1)(1 \otimes B_\nu) + i(1 \otimes B_\mu)(\nabla_\nu^S \otimes 1) \right. \\ &\quad \left. - 1 \otimes B_\mu B_\nu - i\Gamma_{\mu\nu}^\rho \otimes B_\rho \right) \\ &= \Delta^S \otimes 1 - 2i(1 \otimes B^\mu)(\nabla_\mu^S \otimes 1) - ig^{\mu\nu}(1 \otimes \partial_\mu B_\nu) \\ &\quad + 1 \otimes B_\mu B^\mu + ig^{\mu\nu}\Gamma_{\mu\nu}^\rho \otimes B_\rho. \end{aligned} \quad (10.3.4)$$

We are now ready to prove that the fluctuated Dirac operator  $D_\omega$  of an almost-commutative manifold satisfies the following *generalized Lichnerowicz formula* or *Weitzenböck formula*. First, for the canonical Dirac operator  $D_M$  on a compact Riemannian spin manifold  $M$ , recall the Lichnerowicz formula of Theorem 4.21:

$$D_M^2 = \Delta^S + \frac{1}{4}s, \quad (10.3.5)$$

where  $\Delta^S$  is the Laplacian of the spin connection  $\nabla^S$ , and  $s$  is the scalar curvature of  $M$ .

**Proposition 10.6** *The square of the fluctuated Dirac operator on an almost-commutative manifold is a generalized Laplacian of the form*

$$D_\omega^2 = \Delta^E - F,$$

where the endomorphism  $F$  is given by

$$F = -\frac{1}{4}s \otimes 1 - 1 \otimes \Phi^2 + \frac{1}{2}i\gamma^\mu\gamma^\nu \otimes F_{\mu\nu} - i\gamma_M\gamma^\mu \otimes D_\mu\Phi, \quad (10.3.6)$$

in which  $D_\mu$  and  $F_{\mu\nu}$  are defined in (10.3.1) and (10.3.2), respectively.

**Proof** Rewriting the formula for  $D_\omega$ , we have

$$\begin{aligned} D_\omega^2 &= (D_M \otimes 1 + \gamma^\mu \otimes B_\mu + \gamma_M \otimes \Phi)^2 \\ &= D_M^2 \otimes 1 + \gamma^\mu\gamma^\nu \otimes B_\mu B_\nu + 1 \otimes \Phi^2 + (D_M\gamma^\mu \otimes 1)(1 \otimes B_\mu) \\ &\quad + (1 \otimes B_\mu)(\gamma^\mu D_M \otimes 1) + (D_M \otimes 1)(\gamma_M \otimes \Phi) + (\gamma_M \otimes \Phi)(D_M \otimes 1) \\ &\quad + (\gamma^\mu \otimes B_\mu)(\gamma_M \otimes \Phi) + (\gamma_M \otimes \Phi)(\gamma^\mu \otimes B_\mu). \end{aligned}$$

For the first term we use the Lichnerowicz formula of (10.3.5). We rewrite the second term into

$$\begin{aligned}\gamma^\mu \gamma^\nu \otimes B_\mu B_\nu &= \frac{1}{2} \gamma^\mu \gamma^\nu \otimes (B_\mu B_\nu + B_\nu B_\mu + [B_\mu, B_\nu]) \\ &= 1 \otimes B_\mu B^\mu + \frac{1}{2} \gamma^\mu \gamma^\nu \otimes [B_\mu, B_\nu],\end{aligned}$$

where we have used the Clifford relation (4.2.2) to obtain the second equality. For the fourth and fifth terms we use the local formula  $D_M = -i\gamma^\nu \nabla_\nu^S$  to obtain

$$\begin{aligned}(D_M \gamma^\mu \otimes 1)(1 \otimes B_\mu) + (1 \otimes B_\mu)(\gamma^\mu D_M \otimes 1) \\ = -(i\gamma^\nu \nabla_\nu^S \gamma^\mu \otimes 1)(1 \otimes B_\mu) - (1 \otimes B_\mu)(\gamma^\mu i\gamma^\nu \nabla_\nu^S \otimes 1).\end{aligned}$$

Using the identity  $[\nabla_\nu^S, c(\alpha)] = c(\nabla_\nu \alpha)$  for the spin connection, we find  $[\nabla_\nu^S \otimes 1, (\gamma^\mu \otimes 1)(1 \otimes B_\mu)] = c(\nabla_\nu(dx^\mu \otimes B_\mu))$ . We thus obtain

$$\begin{aligned}(D_M \gamma^\mu \otimes 1)(1 \otimes B_\mu) + (1 \otimes B_\mu)(\gamma^\mu D_M \otimes 1) \\ = -i(\gamma^\nu \otimes 1)c(\nabla_\nu(dx^\mu \otimes B_\mu)) \\ - i(\gamma^\nu \gamma^\mu \otimes 1)(1 \otimes B_\mu)(\nabla_\nu^S \otimes 1) - i(1 \otimes B_\mu)(\gamma^\mu \gamma^\nu \nabla_\nu^S \otimes 1) \\ = -i(\gamma^\nu \otimes 1)c(dx^\mu \otimes (\partial_\nu B_\mu) - \Gamma_{\mu\nu}^\rho dx^\mu \otimes B_\rho) - 2i(1 \otimes B^\nu)(\nabla_\nu^S \otimes 1) \\ = -i(\gamma^\nu \gamma^\mu \otimes 1)\left(1 \otimes \partial_\nu B_\mu - \Gamma_{\mu\nu}^\rho \otimes B_\rho\right) - 2i(1 \otimes B^\nu)(\nabla_\nu^S \otimes 1) \\ = -i(\gamma^\nu \gamma^\mu \otimes 1)(1 \otimes \partial_\nu B_\mu) + ig^{\mu\nu} \Gamma_{\mu\nu}^\rho \otimes B_\rho - 2i(1 \otimes B^\nu)(\nabla_\nu^S \otimes 1).\end{aligned}$$

The sixth and seventh terms are rewritten into

$$\begin{aligned}(D_M \otimes 1)(\gamma_M \otimes \Phi) + (\gamma_M \otimes \Phi)(D_M \otimes 1) &= -(\gamma_M \otimes 1)[D_M \otimes 1, 1 \otimes \Phi] \\ &= (\gamma_M \otimes 1)(i\gamma^\mu \otimes \partial_\mu \Phi) = i\gamma_M \gamma^\mu \otimes \partial_\mu \Phi.\end{aligned}$$

The eighth and ninth terms are rewritten as

$$(\gamma^\mu \otimes B_\mu)(\gamma_M \otimes \Phi) + (\gamma_M \otimes \Phi)(\gamma^\mu \otimes B_\mu) = -\gamma_M \gamma^\mu \otimes [B_\mu, \Phi].$$

Summing all these terms then yields the formula

$$\begin{aligned}D_\omega^2 &= \left(\Delta^S + \frac{1}{4}s\right) \otimes 1 + (1 \otimes B_\mu B^\mu) + \frac{1}{2} \gamma^\mu \gamma^\nu \otimes [B_\mu, B_\nu] \\ &\quad + 1 \otimes \Phi^2 - i(\gamma^\nu \gamma^\mu \otimes 1)(1 \otimes \partial_\nu B_\mu) + ig^{\mu\nu} \Gamma_{\mu\nu}^\rho \otimes B_\rho \\ &\quad - 2i(1 \otimes B^\nu)(\nabla_\nu^S \otimes 1) + i\gamma_M \gamma^\mu \otimes \partial_\mu \Phi - \gamma_M \gamma^\mu \otimes [B_\mu, \Phi].\end{aligned}$$

Inserting the formula for  $\Delta^E$  from (10.3.4), we obtain

$$\begin{aligned} D_\omega^2 &= \Delta^E + \frac{1}{4}s \otimes 1 + \frac{1}{2}\gamma^\mu\gamma^\nu \otimes [B_\mu, B_\nu] \\ &\quad + 1 \otimes \Phi^2 - i(\gamma^\nu\gamma^\mu \otimes 1)(1 \otimes \partial_\nu B_\mu) + ig^{\mu\nu}(1 \otimes \partial_\mu B_\nu) \\ &\quad + i\gamma_M\gamma^\mu \otimes \partial_\mu\Phi - \gamma_M\gamma^\mu \otimes [B_\mu, \Phi]. \end{aligned}$$

Using (10.3.2), we rewrite

$$\begin{aligned} &-i(\gamma^\nu\gamma^\mu \otimes 1)(1 \otimes \partial_\nu B_\mu) + ig^{\mu\nu}(1 \otimes \partial_\mu B_\nu) \\ &= -i(\gamma^\nu\gamma^\mu \otimes 1)(1 \otimes \partial_\nu B_\mu) + \frac{1}{2}i(\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu) \otimes (\partial_\mu B_\nu) \\ &= -\frac{1}{2}i\gamma^\mu\gamma^\nu \otimes (\partial_\mu B_\nu) + \frac{1}{2}i\gamma^\nu\gamma^\mu \otimes (\partial_\mu B_\nu) \\ &= -\frac{1}{2}i\gamma^\mu\gamma^\nu \otimes F_{\mu\nu} - \frac{1}{2}\gamma^\mu\gamma^\nu \otimes [B_\mu, B_\nu]. \end{aligned}$$

Using (10.3.1), we finally obtain

$$D_\omega^2 = \Delta^E + \frac{1}{4}s \otimes 1 + 1 \otimes \Phi^2 - \frac{1}{2}i\gamma^\mu\gamma^\nu \otimes F_{\mu\nu} + i\gamma_M\gamma^\mu \otimes D_\mu\Phi,$$

from which we can read off formula (10.3.6) for  $F$ .  $\square$

### 10.3.2 The Heat Expansion

Below, we present two important theorems (without proof) which we will need to calculate the spectral action of almost-commutative manifolds. The first of these theorems states that there exists a heat expansion for a generalized Laplacian. The second theorem gives explicit formulas for the first three non-zero coefficients of this expansion. Next, we will show how these theorems can be applied to obtain a perturbative expansion of the spectral action for an almost-commutative manifold, just as in Proposition 9.7.

**Theorem 10.7** *For a generalized Laplacian  $H = \Delta^E - F$  on  $E$  we have the following asymptotic expansion as  $t \rightarrow 0$ , known as the heat expansion:*

$$\text{Tr}(e^{-tH}) \sim \sum_{k \geq 0} t^{\frac{k-n}{2}} a_k(H), \quad (10.3.7)$$

where  $n$  is the dimension of the manifold, the trace is taken over the Hilbert space  $L^2(E)$  and the coefficients of the expansion are given by

$$a_k(H) := \int_M a_k(x, H) \sqrt{g} d^4x, \quad (10.3.8)$$

where  $\sqrt{g}d^4x$  denotes the Riemannian volume form. The coefficients  $a_k(x, H)$  are called the Seeley-DeWitt coefficients.

**Proof** See Note 6 of this chapter below.  $\square$

**Theorem 10.8** For a generalized Laplacian  $H = \Delta^E - F$  (as in Theorem 10.7), the Seeley-DeWitt coefficients are given by

$$\begin{aligned} a_0(x, H) &= (4\pi)^{-\frac{n}{2}} \text{Tr}(\text{id}), \\ a_2(x, H) &= (4\pi)^{-\frac{n}{2}} \text{Tr}\left(\frac{s}{6} + F\right), \\ a_4(x, H) &= (4\pi)^{-\frac{n}{2}} \frac{1}{360} \text{Tr}\left(-12\Delta s + 5s^2 - 2R_{\mu\nu}R^{\mu\nu} + 2R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}\right. \\ &\quad \left.+ 60sF + 180F^2 - 60\Delta F + 30\Omega_{\mu\nu}^E(\Omega^E)^{\mu\nu}\right), \end{aligned}$$

where this time the traces are taken over the fibre  $E_x$ . Here  $s$  is the scalar curvature of the Levi-Civita connection  $\nabla$ ,  $\Delta$  is the scalar Laplacian, and  $\Omega^E$  is the curvature of the connection  $\nabla^E$  corresponding to  $\Delta^E$ . All  $a_k(x, H)$  with odd  $k$  vanish.

**Proof** See Note 6 of this chapter below.  $\square$

We saw in Proposition 10.6 that the square of the fluctuated Dirac operator of an almost-commutative manifold is a generalized Laplacian. Applying Theorem 10.7 to  $D_\omega^2$  in dimension  $n = 4$  then yields the heat expansion:

$$\text{Tr}\left(e^{-tD_\omega^2}\right) \sim \sum_{k \geq 0} t^{\frac{k-4}{2}} a_k(D_\omega^2), \quad (10.3.9)$$

where the Seeley-DeWitt coefficients are given by Theorem 10.8. In the following proposition, we use this heat expansion for  $D_\omega^2$  to obtain an expansion of the spectral action.

**Proposition 10.9** For an almost-commutative manifold  $M \times F$  with  $M$  of dimension 4, the spectral action given by (9.1.1) can be expanded asymptotically (as  $\Lambda \rightarrow \infty$ ) as

$$\text{Tr}\left(f\left(\frac{D_\omega}{\Lambda}\right)\right) \sim a_4(D_\omega^2)f(0) + 2 \sum_{\substack{0 \leq k < 4 \\ k \text{ even}}} f_{4-k}\Lambda^{4-k} a_k(D_\omega^2) \frac{1}{\Gamma\left(\frac{4-k}{2}\right)} + \mathcal{O}(\Lambda^{-1}),$$

where  $f_j = \int_0^\infty f(v)v^{j-1}dv$  are the moments of the function  $f$ ,  $j > 0$ .

**Proof** Our proof is based on Proposition 9.7. Let  $g$  be the function  $g(u^2) = f(u)$ , so that its Laplace–Stieltjes transform

$$g(v) = \int_0^\infty e^{-sv} d\mu(s).$$

We can then formally write

$$g(tD_\omega^2) = \int_0^\infty e^{-stD_\omega^2} d\mu(s).$$

We now take the trace and use the heat expansion of  $D_\omega^2$  to obtain

$$\begin{aligned} \text{Tr}(g(tD_\omega^2)) &= \int_0^\infty \text{Tr}(e^{-stD_\omega^2}) d\mu(s) \sim \int_0^\infty \sum_{k \geq 0} (st)^{\frac{k-4}{2}} a_k(D_\omega^2) d\mu(s) \\ &= \sum_{k \geq 0} t^{\frac{k-4}{2}} a_k(D_\omega^2) \int_0^\infty s^{\frac{k-4}{2}} d\mu(s). \end{aligned} \quad (10.3.10)$$

The parameter  $t$  is considered to be a formal expansion parameter. From here on, we will drop the terms with  $k > 4$ . The term with  $k = 4$  equals

$$a_4(D_\omega^2) \int_0^\infty s^0 d\mu(s) = a_4(D_\omega^2) g(0).$$

We can rewrite the terms with  $k < 4$  using the definition of the  $\Gamma$ -function as the analytic continuation of

$$\Gamma(z) = \int_0^\infty r^{z-1} e^{-r} dr, \quad (10.3.11)$$

for  $z \in \mathbb{C}$  with  $\Re(z) > 0$ , and by inserting  $r = sv$ , we see that (for  $k < 4$ ) we have

$$\Gamma\left(\frac{4-k}{2}\right) = \int_0^\infty (sv)^{\frac{4-k}{2}-1} e^{-sv} d(sv) = s^{\frac{4-k}{2}} \int_0^\infty v^{\frac{4-k}{2}-1} e^{-sv} dv.$$

From this, we obtain an expression for  $s^{\frac{k-4}{2}}$ , which we insert into Eq. (10.3.10), and then we perform the integration over  $s$  to obtain

$$\begin{aligned} \text{Tr}(g(tD_\omega^2)) &\sim a_4(D_\omega^2) f(0) \\ &+ \sum_{0 \leq k < 4} t^{\frac{k-4}{2}} a_k(D_\omega^2) \frac{1}{\Gamma\left(\frac{4-k}{2}\right)} \int_0^\infty v^{\frac{4-k}{2}-1} g(v) dv + \mathcal{O}(\Lambda^{-1}). \end{aligned}$$

Now we choose the function  $g$  such that  $g(u^2) = f(u)$ . We rewrite the integration over  $v$  by substituting  $v = u^2$  and obtain

$$\int_0^\infty v^{\frac{4-k}{2}-1} g(v) dv = \int_0^\infty u^{4-k-2} g(u^2) d(u^2) = 2 \int_0^\infty u^{4-k-1} f(u) du,$$

which by definition equals  $2f_{4-k}$ . Upon writing  $t = \Lambda^{-2}$ , we have modulo  $\Lambda^{-1}$ ,

$$\begin{aligned} \mathrm{Tr} \left( f \left( \frac{D_\omega}{\Lambda} \right) \right) &= \mathrm{Tr} \left( g(\Lambda^{-2} D_\omega^2) \right) \\ &\sim a_4(D_\omega^2) f(0) + 2 \sum_{0 \leq k < 4} f_{4-k} \Lambda^{4-k} a_k(D_\omega^2) \frac{1}{\Gamma(\frac{4-k}{2})} + \mathcal{O}(\Lambda^{-1}). \end{aligned}$$

Using  $a_k(D_\omega^2) = 0$  for odd  $k$ , the claim follows.  $\square$

## 10.4 The Spectral Action on AC Manifolds

In the previous section we obtained a perturbative expansion of the spectral action for an almost-commutative manifold. We now explicitly calculate the coefficients in this expansion, first for the canonical triple (yielding the (Euclidean) Einstein–Hilbert action of General Relativity) for a four-dimensional Riemannian spin manifold  $M$  and then for a general almost-commutative manifold  $M \times F$ .

By Proposition 10.9 we have an asymptotic expansion as  $\Lambda \rightarrow \infty$ :

$$\mathrm{Tr} \left( f \left( \frac{D_\omega}{\Lambda} \right) \right) \sim 2f_4 \Lambda^4 a_0(D_\omega^2) + 2f_2 \Lambda^2 a_2(D_\omega^2) + f(0) a_4(D_\omega^2) + \mathcal{O}(\Lambda^{-1}). \quad (10.4.1)$$

**Proposition 10.10** *For the canonical triple  $(C^\infty(M), L^2(S), D_M)$ , the spectral action is given by:*

$$\mathrm{Tr} \left( f \left( \frac{D_M}{\Lambda} \right) \right) \sim \int_M \mathcal{L}_M(g_{\mu\nu}) \sqrt{g} d^4x + \mathcal{O}(\Lambda^{-1}), \quad (10.4.2)$$

where the Lagrangian is defined by

$$\mathcal{L}_M(g_{\mu\nu}) := \frac{f_4 \Lambda^4}{2\pi^2} - \frac{f_2 \Lambda^2}{24\pi^2} s + \frac{f(0)}{16\pi^2} \left( \frac{1}{30} \Delta s - \frac{1}{20} C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} + \frac{11}{360} R^* R^* \right).$$

Here the Weyl tensor  $C_{\mu\nu\rho\sigma}$  is given by the traceless part of the Riemann curvature tensor; so that

$$C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} = R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 2R_{\nu\sigma} R^{\nu\sigma} + \frac{1}{3} s^2, \quad (10.4.3)$$

and  $R^*$  is related to the Pontryagin class:

$$R^* R^* = s^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}. \quad (10.4.4)$$

**Proof** We have  $n = 4$ , and  $\text{Tr}(\text{id}) = \dim S_x = 4$  where  $S_x$  is the fiber of  $S$  at some  $x \in M$ . Inserting this into Theorem 10.8 gives

$$a_0(D_M^2) = \frac{1}{4\pi^2} \int_M \sqrt{g} d^4x.$$

From the Lichnerowicz formula (10.3.5) we see that  $F = -\frac{1}{4}s \text{id}$ , so

$$a_2(D_M^2) = -\frac{1}{48\pi^2} \int_M s \sqrt{g} d^4x.$$

Moreover,

$$5s^2\text{id} + 60sF + 180F^2 = \frac{5}{4}s^2\text{id}.$$

Inserting this into  $a_4(D_M^2)$  gives

$$\begin{aligned} a_4(D_M^2) &= \frac{1}{16\pi^2} \frac{1}{360} \int_M \text{Tr} \left( 3\Delta s \text{id} + \frac{5}{4}s^2\text{id} - 2R_{\mu\nu}R^{\mu\nu}\text{id} \right. \\ &\quad \left. + 2R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}\text{id} + 30\Omega_{\mu\nu}^S\Omega^{S\mu\nu} \right) \sqrt{g} d^4x. \end{aligned}$$

The curvature  $\Omega^S$  of the spin connection is defined as in (4.2.3), and its components are  $\Omega_{\mu\nu}^S = \Omega^S(\partial_\mu, \partial_\nu)$ . The spin curvature  $\Omega^S$  is related to the Riemannian curvature tensor by (see Note 8 of this chapter below),

$$\Omega_{\mu\nu}^S = \frac{1}{4}R_{\mu\nu\rho\sigma}\gamma^\rho\gamma^\sigma. \quad (10.4.5)$$

We use this as well as the trace identity

$$\text{Tr}(\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma) = 4(g^{\mu\nu}g^{\rho\sigma} - g^{\mu\rho}g^{\nu\sigma} + g^{\mu\sigma}g^{\nu\rho})$$

to calculate the last term of  $a_4(D_M^2)$ :

$$\begin{aligned} \text{Tr}(\Omega_{\mu\nu}^S\Omega^{S\mu\nu}) &= \frac{1}{16}R_{\mu\nu\rho\sigma}R^{\mu\nu}_{\lambda\kappa} \text{Tr}(\gamma^\rho\gamma^\sigma\gamma^\lambda\gamma^\kappa) \\ &= \frac{1}{4}R_{\mu\nu\rho\sigma}R^{\mu\nu}_{\lambda\kappa} (g^{\rho\sigma}g^{\lambda\kappa} - g^{\rho\lambda}g^{\sigma\kappa} + g^{\rho\kappa}g^{\sigma\lambda}) = -\frac{1}{2}R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, \end{aligned} \quad (10.4.6)$$

where the first term in the second line vanishes because of the antisymmetry of  $R_{\mu\nu\rho\sigma}$  in  $\rho$  and  $\sigma$ , and the other two terms contribute equally. We thus obtain

$$a_4(D_M^2) = \frac{1}{16\pi^2} \frac{1}{360} \int_M (12\Delta s + 5s^2 - 8R_{\mu\nu}R^{\mu\nu} - 7R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}) \sqrt{g}d^4x. \quad (10.4.7)$$

We rewrite this into a more convenient form, using (10.4.3) and (10.4.4), which together yield:

$$\begin{aligned} & -\frac{1}{20}C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma} + \frac{11}{360}R^*R^* \\ &= -\frac{1}{20}R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} + \frac{1}{10}R_{\nu\sigma}R^{\nu\sigma} - \frac{1}{60}s^2 \\ &+ \frac{11}{360}R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - \frac{44}{360}R_{\nu\sigma}R^{\nu\sigma} + \frac{11}{360}s^2 \\ &= \frac{1}{360}(-7R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 8R_{\nu\sigma}R^{\nu\sigma} + 5s^2). \end{aligned}$$

Therefore, we may rewrite (10.4.7) so as to obtain

$$a_4(D_M^2) = \frac{1}{16\pi^2} \int_M \left( \frac{1}{30}\Delta s - \frac{1}{20}C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma} + \frac{11}{360}R^*R^* \right) \sqrt{g}d^4x.$$

Inserting the obtained formulas for  $a_0(D_M^2)$ ,  $a_2(D_M^2)$  and  $a_4(D_M^2)$  into (10.4.1) proves the proposition.

**Remark 10.11** In general, an expression of the form

$$as^2 + bR_{\nu\sigma}R^{\nu\sigma} + cR_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma},$$

for certain constants  $a, b, c \in \mathbb{R}$ , can always be rewritten in the form  $\alpha s^2 + \beta C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma} + \gamma R^*R^*$ , for new constants  $\alpha, \beta, \gamma \in \mathbb{R}$ . One should note here that the term  $s^2$  is not present in the spectral action of the canonical triple as calculated in Proposition 10.10. The only higher-order gravitational term that arises is the conformal gravity term  $C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}$ .

Note that alternatively, using only (10.4.4), we could also have written

$$a_4(D_M^2) = \frac{1}{16\pi^2} \frac{1}{30} \int_M (\Delta s + s^2 - 3R_{\mu\nu}R^{\mu\nu} - \frac{7}{12}R^*R^*) \sqrt{g}d^4x.$$

The integral over  $\Delta s$  only yields a boundary term, so if the manifold  $M$  is compact without boundary, we can discard the term with  $\Delta s$ . Furthermore, for a 4-dimensional compact orientable manifold  $M$  without boundary, we have the formula

$$\int_M R^* R^* \sqrt{g} dx = 8\pi^2 \chi(M),$$

where  $\chi(M)$  is Euler characteristic. Hence the term with  $R^* R^*$  only yields a topological contribution to the action, which we will also disregard. From here on, we will therefore consider the Lagrangian

$$\mathcal{L}_M(g_{\mu\nu}) = \frac{f_4 \Lambda^4}{2\pi^2} - \frac{f_2 \Lambda^2}{24\pi^2} s - \frac{f(0)}{320\pi^2} C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma}, \quad (10.4.8)$$

or, which is the same,

$$\mathcal{L}_M(g_{\mu\nu}) = \frac{f_4 \Lambda^4}{2\pi^2} - \frac{f_2 \Lambda^2}{24\pi^2} s + \frac{f(0)}{480\pi^2} (s^2 - 3R_{\mu\nu}R^{\mu\nu}). \quad (10.4.9)$$

**Proposition 10.12** *The spectral action of the fluctuated Dirac operator of an almost-commutative manifold with  $\dim M = 4$  is given by*

$$\text{Tr} \left( f \left( \frac{D_\omega}{\Lambda} \right) \right) \sim \int_M \mathcal{L}(g_{\mu\nu}, B_\mu, \Phi) \sqrt{g} d^4x + \mathcal{O}(\Lambda^{-1}),$$

where

$$\mathcal{L}(g_{\mu\nu}, B_\mu, \Phi) := N \mathcal{L}_M(g_{\mu\nu}) + \mathcal{L}_B(B_\mu) + \mathcal{L}_\phi(g_{\mu\nu}, B_\mu, \Phi).$$

Here  $\mathcal{L}_M(g_{\mu\nu})$  is defined in Proposition 10.10,  $N$  is the dimension of the finite-dimensional Hilbert space  $H_F$ , and  $\mathcal{L}_B$  gives the kinetic term of the gauge field as

$$\mathcal{L}_B(B_\mu) := \frac{f(0)}{24\pi^2} \text{Tr}(F_{\mu\nu} F^{\mu\nu}),$$

and  $\mathcal{L}_\phi$  gives a scalar-field Lagrangian including its interactions plus a boundary term as

$$\begin{aligned} \mathcal{L}_\phi(g_{\mu\nu}, B_\mu, \Phi) &:= -\frac{2f_2 \Lambda^2}{4\pi^2} \text{Tr}(\Phi^2) + \frac{f(0)}{8\pi^2} \text{Tr}(\Phi^4) + \frac{f(0)}{24\pi^2} \Delta(\text{Tr}(\Phi^2)) \\ &\quad + \frac{f(0)}{48\pi^2} s \text{Tr}(\Phi^2) + \frac{f(0)}{8\pi^2} \text{Tr}((D_\mu \Phi)(D^\mu \Phi)). \end{aligned} \quad (10.4.10)$$

**Proof** The proof is very similar to Proposition 10.10, but we now use the formula for  $D_\omega^2$  given by Proposition 10.6. The trace over the Hilbert space  $H_F$  yields an overall factor  $N := \text{Tr}(1_{H_F})$ , so we have

$$a_0(D_\omega^2) = N a_0(D_M^2).$$

The square of the Dirac operator now contains three extra terms. The trace of  $\gamma_M \gamma^\mu$  vanishes, which follows from cyclicity of the trace and the fact that  $\gamma_M \gamma^\mu = -\gamma^\mu \gamma_M$ . Since  $\text{Tr}(\gamma^\mu \gamma^\nu) = 4g^{\mu\nu}$  and  $F_{\mu\nu}$  is anti-symmetric, the trace of  $\gamma^\mu \gamma^\nu F_{\mu\nu}$  also vanishes. Thus we find that

$$a_2(D_\omega^2) = Na_2(D_M^2) - \frac{1}{4\pi^2} \int_M \text{Tr}(\Phi^2) \sqrt{g} d^4x.$$

Furthermore we obtain several new terms from the formula for  $a_4(D_\omega^2)$ . First, we calculate

$$\frac{1}{360} \text{Tr}(60sF) = -\frac{1}{6}s(Ns + 4\text{Tr}(\Phi^2)).$$

The next contribution arises from the trace over  $F^2$ , which equals

$$\begin{aligned} F^2 &= \frac{1}{16}s^2 \otimes 1 + 1 \otimes \Phi^4 - \frac{1}{4}\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma \otimes F_{\mu\nu} F_{\rho\sigma} \\ &\quad + \gamma^\mu \gamma^\nu \otimes (D_\mu \Phi)(D_\nu \Phi) + \frac{1}{2}s \otimes \Phi^2 + \text{traceless terms}. \end{aligned}$$

Taking the trace then yields

$$\begin{aligned} \frac{1}{360} \text{Tr}(180F^2) &= \frac{N}{8}s^2 + 2\text{Tr}(\Phi^4) + \text{Tr}(F_{\mu\nu} F^{\mu\nu}) \\ &\quad + 2\text{Tr}((D_\mu \Phi)(D^\mu \Phi)) + s\text{Tr}(\Phi^2). \end{aligned}$$

Another contribution arises from  $-\Delta F$ . Again, we can simply ignore the traceless terms and obtain

$$\frac{1}{360} \text{Tr}(-60\Delta F) = \frac{1}{6}\Delta(Ns + 4\text{Tr}(\Phi^2)).$$

The final contribution comes from the term  $\Omega_{\mu\nu}^E \Omega^{E\mu\nu}$ , where the curvature  $\Omega^E$  is given by (10.3.3); we obtain

$$\Omega_{\mu\nu}^E \Omega^{E\mu\nu} = \Omega_{\mu\nu}^S \Omega^{S\mu\nu} \otimes 1 - 1 \otimes F_{\mu\nu} F^{\mu\nu} + 2i\Omega_{\mu\nu}^S \otimes F^{\mu\nu}.$$

Using (10.4.5), by the anti-symmetry of  $R_{\rho\sigma\mu\nu}$  we find

$$\text{Tr}(\Omega_{\mu\nu}^S) = \frac{1}{4}R_{\rho\sigma\mu\nu} \text{Tr}(\gamma^\rho \gamma^\sigma) = \frac{1}{4}R_{\rho\sigma\mu\nu} g^{\rho\sigma} = 0,$$

so the trace over the cross-terms in  $\Omega_{\mu\nu}^E \Omega^{E\mu\nu}$  vanishes. From (10.4.6) we then obtain

$$\frac{1}{360} \text{Tr}(30\Omega_{\mu\nu}^E \Omega^{E\mu\nu}) = \frac{1}{12} \left( -\frac{N}{2} R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4 \text{Tr}(F_{\mu\nu} F^{\mu\nu}) \right).$$

Gathering all terms, we obtain

$$\begin{aligned} a_4(x, D_\omega^2) &= \frac{1}{(4\pi)^2} \frac{1}{360} \left( -48N\Delta s + 20Ns^2 - 8NR_{\mu\nu}R^{\mu\nu} \right. \\ &\quad + 8NR_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 60s(Ns + 4\text{Tr}(\Phi^2)) \\ &\quad + 360 \left( \frac{N}{8}s^2 + 2\text{Tr}(\Phi^4) + \text{Tr}(F_{\mu\nu}F^{\mu\nu}) \right. \\ &\quad \left. \left. + 2\text{Tr}((D_\mu\Phi)(D^\mu\Phi)) + s\text{Tr}(\Phi^2) \right) \right. \\ &\quad + 60\Delta(Ns + 4\text{Tr}(\Phi^2)) \\ &\quad \left. - 30 \left( \frac{N}{2}R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} + 4\text{Tr}(F_{\mu\nu}F^{\mu\nu}) \right) \right) \\ &= \frac{1}{(4\pi)^2} \frac{1}{360} \left( 12N\Delta s + 5Ns^2 - 8NR_{\mu\nu}R^{\mu\nu} \right. \\ &\quad - 7NR_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} + 120s\text{Tr}(\Phi^2) \\ &\quad + 360 \left( 2\text{Tr}(\Phi^4) + 2\text{Tr}((D_\mu\Phi)(D^\mu\Phi)) \right) \\ &\quad \left. + 240\Delta(\text{Tr}(\Phi^2)) + 240\text{Tr}(F_{\mu\nu}F^{\mu\nu}) \right). \end{aligned}$$

Comparing the first line of the second equality to (10.4.7), we see that

$$\begin{aligned} a_4(x, D_\omega^2) &= Na_4(x, D_M^2) + \frac{1}{4\pi^2} \left( \frac{1}{12}s\text{Tr}(\Phi^2) + \frac{1}{2}\text{Tr}(\Phi^4) \right. \\ &\quad \left. + \frac{1}{2}\text{Tr}((D_\mu\Phi)(D^\mu\Phi)) + \frac{1}{6}\Delta(\text{Tr}(\Phi^2)) + \frac{1}{6}\text{Tr}(F_{\mu\nu}F^{\mu\nu}) \right). \end{aligned}$$

Inserting these Seeley-DeWitt coefficients into (10.4.1) proves the proposition.  $\square$

Note that the above Lagrangian is indeed gauge invariant. This is of course a consequence of the manifest gauge invariance of the spectral action, which follows from the invariance of the spectrum under unitary transformations.

**Example 10.13** Let us return to the Yang–Mills manifold  $M \times F_{\text{YM}}$  of Examples 10.4 and 10.5. We have already seen that the inner fluctuations are parametrized by a  $PU(N)$  gauge field  $A_\mu$ , which acts in the adjoint representation  $B_\mu = \text{ad } A_\mu$  on the fermions. There is no scalar field  $\phi$  and  $\Phi = D_F = 0$ . We can insert these fields into the result of Proposition 10.12. The dimension of the Hilbert space  $H_F = M_N(\mathbb{C})$  is  $N^2$ . We then find that the Lagrangian of the Yang–Mills manifold is given by

$$\mathcal{L}(g_{\mu\nu}, B_\mu) := N^2 \mathcal{L}_M(g_{\mu\nu}) + \frac{f(0)}{24\pi^2} \mathcal{L}_{YM}(B_\mu).$$

Here  $\mathcal{L}_{YM}$  is the Yang–Mills Lagrangian given by

$$\mathcal{L}_{YM}(B_\mu) := \text{Tr}(F_{\mu\nu} F^{\mu\nu}),$$

where  $F_{\mu\nu}$  denotes the curvature of  $B_\mu$ .

## Notes

### Section 10.1. Gauge Symmetries of AC Manifolds

1. Kaluza–Klein theories date back to [1, 2].
2. The name almost-commutative manifolds was coined in [3], suggesting that the noncommutativity is mild since it is simply given by the matrix product in  $A_F$ , pointwise on  $M$ . Almost-commutative manifolds essentially already appeared in [4], and somewhat later in the work of Connes and Lott [5]. Around the same time, a similar structure appeared in a series of papers by Dubois-Violette, Kerner and Madore [6–9], who studied the noncommutative differential geometry for the algebra of functions tensored with a matrix algebra, and its relevance to the description of gauge and scalar Higgs fields. Almost-commutative manifolds were later used by Chamseddine and Connes [10, 11], and by Chamseddine, Connes and Marcolli in [12] to geometrically describe Yang–Mills theories and the Standard Model of elementary particles, as we will see in the next chapters. We here base our treatment on [13].
3. We can regard  $C^\infty(M, A_F)$  as the space of smooth sections of a globally trivial  $*$ -algebra bundle  $M \times A_F$ . The natural question whether the above definition can be extended to the topologically non-trivial case is addressed in [14–16]. The special case of topologically non-trivial Yang–Mills theories is treated in [17] and in the next Chapter.
4. In the proof of Proposition 10.2 we have exploited a lift of group bundles, which exists if the manifold is simply connected. We refer to [16] for a careful discussion on this point.

### Section 10.3. The Heat Expansion of the Spectral Action

5. For more details on generalized Laplacians we refer to [18, Sect. 2.1].
6. Theorem 10.7 is proved by Gilkey in [19, Sect. 1.7]. Theorem 10.8 can be found as [19, Theorem 4.8.16]. For a more physicist-friendly approach, we refer to [20]. Note that the conventions used by Gilkey for the Riemannian curvature  $R$  are such that  $g^{\mu\rho} g^{\nu\sigma} R_{\mu\nu\rho\sigma}$  is negative for a sphere, in contrast to our own conventions. Therefore we have replaced  $s = -R$ .

### Section 10.4. The Spectral Action on AC Manifolds

7. The bosonic Lagrangian derived from the spectral action for AC manifolds was interpreted in [10] à la Wilson [21] as the bare Lagrangian at the cutoff scale  $\Lambda$ . A perturbative expansion of the full spectral action was obtained in [22–24], leading to unexpected and an intriguing behaviour for the propagation of particles at energies larger than the cutoff  $\Lambda$ . Alternatively, the interpretation of  $\Lambda$  as a regularization parameter has been worked out in [25–28], including the derivation of renormalizability conditions on the Krajewski diagrams.
8. The relation (10.4.5) is derived in [29, p. 395].
9. The derivation of Yang–Mills gauge theory from a noncommutative spin manifold as in Example 10.13 is due to Chamseddine and Connes in [10, 11].

## References

1. Kaluza, T.: Zum Unitätsproblem in der Physik . Sitzungsber. Preuss. Akad. Wiss. Berlin (Math.Phys.) **1921** 966–972 (1921)
2. Klein, O.: Quantentheorie und fünfdimensionale relativitätstheorie. Z. Phys. **37**, 895–906 (1926)
3. Iochum, B., Schucker, T., Stephan, C.: On a classification of irreducible almost commutative geometries. J. Math. Phys. **45**, 5003–5041 (2004)
4. Connes, A.: Essay on physics and noncommutative geometry. In: The Interface of Mathematics and Particle Physics (Oxford, 1988), Institute of Mathematics and its Applications Conference Series New Series. vol. 24, pp. 9–48. Oxford University Press, New York (1990)
5. Connes, A., Lott, J.: Particle models and noncommutative geometry. Nucl. Phys. Proc. Suppl. **18B**, 29–47 (1991)
6. Dubois-Violette, M., Kerner, R., Madore, J.: Classical bosons in a noncommutative geometry. Class. Quant. Grav. **6**, 1709 (1989)
7. Dubois-Violette, M., Kerner, R., Madore, J.: Gauge bosons in a noncommutative geometry. Phys. Lett. B **217**, 485–488 (1989)
8. Dubois-Violette, M., Madore, J., Kerner, R.: Noncommutative differential geometry and new models of gauge theory. J. Math. Phys. **31**, 323 (1990)
9. Dubois-Violette, M., Kerner, R., Madore, J.: Noncommutative differential geometry of matrix algebras. J. Math. Phys. **31**, 316 (1990)
10. Chamseddine, A.H., Connes, A.: Universal formula for noncommutative geometry actions: unifications of gravity and the standard model. Phys. Rev. Lett. **77**, 4868–4871 (1996)
11. Chamseddine, A.H., Connes, A.: The spectral action principle. Commun. Math. Phys. **186**, 731–750 (1997)
12. Chamseddine, A.H., Connes, A., Marcolli, M.: Gravity and the standard model with neutrino mixing. Adv. Theor. Math. Phys. **11**, 991–1089 (2007)
13. van den Dungen, K., van Suijlekom, W.D.: Particle physics from almost commutative space-times. Rev. Math. Phys. **24**, 1230004 (2012)
14. Čaćić, B.: A Reconstruction theorem for almost-commutative spectral triples. Lett. Math. Phys. **100**, 181–202 (2012)
15. Čaćić, B.: Real structures on almost-commutative spectral triples. Lett. Math. Phys. **103**, 793–816 (2013)
16. Boejink, J., van den Dungen, K.: On globally non-trivial almost-commutative manifolds. J. Math. Phys. **55**, 103508 (2014)
17. Boejink, J., van Suijlekom, W.D.: The noncommutative geometry of Yang-Mills fields. J. Geom. Phys. **61**, 1122–1134 (2011)
18. Berline, N., Getzler, E., Vergne, M.: Heat Kernels and Dirac Operators. Springer, Berlin (1992)
19. Gilkey, P.B.: Invariance Theory, the Heat Equation, and the Atiyah-Singer Index Theorem. Mathematics Lecture Series, vol. 11. Publish or Perish Inc., Wilmington (1984)
20. Vassilevich, D.V.: Heat kernel expansion: user's manual. Phys. Rept. **388**, 279–360 (2003)
21. Wilson, K.G.: Renormalization group methods. Adv. Math. **16**, 170–186 (1975)
22. Iochum, B., Levy, C., Vassilevich, D.: Spectral action beyond the weak-field approximation. Commun. Math. Phys. **316**, 595–613 (2012)
23. Iochum, B., Levy, C., Vassilevich, D.: Global and local aspects of spectral actions. J. Phys. A **45**, 374020 (2012)
24. Kurkov, M.A., Lizzi, F., Vassilevich, D.: High energy bosons do not propagate. Phys. Lett. B **731**, 311–315 (2014)
25. van Suijlekom, W.D.: Renormalization of the spectral action for the Yang-Mills system. JHEP **1103**, 146 (2011)
26. van Suijlekom, W.D.: Renormalization of the asymptotically expanded Yang-Mills spectral action. Commun. Math. Phys. **312**, 883–912 (2012)
27. van Suijlekom, W.D.: Renormalizability conditions for almost commutative manifolds. Ann. H. Poincaré **15**, 985–1011 (2014)

28. van Suijlekom, W.D.: Renormalizability conditions for almost-commutative geometries. *Phys. Lett. B* **711**, 434–438 (2012)
29. Gracia-Bondía, J.M., Várilly, J.C., Figueroa, H.: Elements of Noncommutative Geometry. Birkhäuser, Boston (2001)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 11

## The Noncommutative Geometry of Electrodynamics



In the previous chapters we have described the general framework for the description of gauge theories in terms of noncommutative manifolds. The present chapter serves two purposes. First, we describe abelian gauge theories within the framework of noncommutative geometry, which at first sight appears to be a *contradictio in terminis*. Second, in Sect. 11.2 we show how this example can be modified to provide a description of one of the simplest examples of a field theory in physics, namely electrodynamics. Because of its simplicity, it helps in gaining an understanding of the formulation of gauge theories in terms of almost-commutative manifolds, and as such it provides a first stepping stone towards the derivation of the Standard Model from noncommutative geometry in Chap. 13.

### 11.1 The Two-Point Space

In this section we discuss one of the simplest finite noncommutative spaces, namely the two-point space  $X = \{x, y\}$ . Recall from Chaps. 2 and 3 that such a space can be described by an even finite real spectral triple:

$$F_x := (C(X) = \mathbb{C}^2, H_F, D_F; J_F, \gamma_F). \quad (11.1.1)$$

As we require the action of  $C(X)$  on the finite-dimensional Hilbert space  $H_F$  to be faithful,  $H_F$  must at least be 2-dimensional. For now we restrict ourselves to the simplest case, taking  $H_F = \mathbb{C}^2$ . We use the  $\mathbb{Z}_2$ -grading  $\gamma_F$  to decompose  $H_F = H_F^+ \oplus H_F^- = \mathbb{C} \oplus \mathbb{C}$  into the two eigenspaces  $H_F^\pm = \{\psi \in H_F \mid \gamma_F \psi = \pm \psi\}$ . The action of  $C(X)$  on  $H_F$  respects this decomposition, whereas  $D_F$  interchanges the two subspaces  $H_F^\pm$ , say

$$D_F = \begin{pmatrix} 0 & t \\ \bar{t} & 0 \end{pmatrix},$$

for some  $t \in \mathbb{C}$ .

**Proposition 11.1** *The finite space  $F_x$  of (11.1.1) can only have a real structure  $J_F$  if  $D_F = 0$ . In that case, its KO-dimension is 0, 2 or 6.*

**Proof** The diagonal representation of the algebra  $\mathbb{C} \oplus \mathbb{C}$  on  $\mathbb{C} \oplus \mathbb{C}$  gives rise to one of the following two Krajewski diagrams (cf. Example 3.13):

$$\begin{array}{cc} \begin{array}{cc} \mathbf{1} & \mathbf{1} \\ \mathbf{1}^\circ & \circ \end{array} & \begin{array}{cc} \mathbf{1} & \mathbf{1} \\ \mathbf{1}^\circ & \circ \end{array} \\ \mathbf{1}^\circ \circ & \mathbf{1}^\circ \quad \circ \end{array}$$

As a Dirac operator  $D_F$  that fulfills the first-order condition 3.1.1 (for arbitrary  $J_F$ ) should connect nodes either vertically or horizontally, we find that  $D_F = 0$ .

The diagram on the left corresponds to KO-dimension 2 and 6, while the diagram on the right corresponds to KO-dimension 0 and 4. KO-dimension 4 is ruled out because of Lemma 3.8, combined with the fact that  $\dim H_F^\pm = 1$ , which does not allow for a  $J_F$  with  $J_F^2 = -1$ .  $\square$

### 11.1.1 The Product Space

Let  $M$  be a compact 4-dimensional Riemannian spin manifold. We now consider the almost-commutative manifold  $M \times F_x$  given by the product of  $M$  with the even finite space  $F_x$  corresponding to the two-point space (11.1.1). Thus we consider the almost-commutative manifold given by the data

$$M \times F_x := \left( C^\infty(M, \mathbb{C}^2), L^2(S) \otimes \mathbb{C}^2, D_M \otimes 1; J_M \otimes J_F, \gamma_M \otimes \gamma_F \right),$$

where we still need to make a choice for  $J_F$ . The algebra of this almost-commutative manifold is given by  $C^\infty(M, \mathbb{C}^2) \simeq C^\infty(M) \oplus C^\infty(M)$ . By Gelfand duality (Theorem 5.7) this algebra corresponds to the space

$$N := M \times X \simeq M \sqcup M,$$

which consists of the disjoint union of two copies of the space  $M$ , so we can write  $C^\infty(N) = C^\infty(M) \oplus C^\infty(M)$ . We can also decompose the total Hilbert space as  $\mathcal{H} = L^2(S) \oplus L^2(S)$ . For  $a, b \in C^\infty(M)$  and  $\psi, \phi \in L^2(S)$ , an element  $(a, b) \in C^\infty(N)$  then simply acts on  $(\psi, \phi) \in \mathcal{H}$  as  $(a, b)(\psi, \phi) = (a\psi, b\phi)$ .

**Remark 11.2** Let us consider Connes' distance formula (cf. Note 5 of this chapter below) on  $M \times F_x$ . First, as in (2.2.2), on the structure space  $X$  of  $A_F$  we may write a metric by:

$$d_{D_F}(x, y) = \sup \{ |a(x) - a(y)| : a \in A_F, \| [D_F, a] \| \leq 1 \}.$$

Note that now we only have two distinct points  $x$  and  $y$  in the space  $X$ , and we are going calculate the distance between these points. An element  $a \in \mathbb{C}^2 = C(X)$  is specified by two complex numbers  $a(x)$  and  $a(y)$ , so a small computation of the commutator with  $D_F$  gives

$$[D_F, a] = (a(y) - a(x)) \begin{pmatrix} 0 & t \\ -\bar{t} & 0 \end{pmatrix}.$$

The norm of this commutator is given by  $|a(y) - a(x)| |t|$ , so  $\| [D_F, a] \| \leq 1$  implies  $|a(y) - a(x)| \leq \frac{1}{|t|}$ . We therefore obtain that the distance between the two points  $x$  and  $y$  is given by

$$d_{D_F}(x, y) = \frac{1}{|t|}.$$

If there is a real structure  $J_F$ , we have  $t = 0$  by Proposition 11.1, so in that case the distance between the two points becomes infinite.

Let  $p$  be a point in  $M$ , and write  $(p, x)$  and  $(p, y)$  for the two corresponding points in  $N = M \times X$ . A function  $a \in C^\infty(N)$  is then determined by two functions  $a_x, a_y \in C^\infty(M)$ , given by  $a_x(p) := a(p, x)$  and  $a_y(p) := a(p, y)$ . Now the distance function on  $N$  is given by

$$d_{D_M \otimes 1}(n_1, n_2) = \sup \{ |a(n_1) - a(n_2)| : a \in \mathcal{A}, \| [D_M \otimes 1, a] \| \leq 1 \}.$$

If  $n_1$  and  $n_2$  are points in the same copy of  $M$ , for instance, if  $n_1 = (p, x)$  and  $n_2 = (q, x)$  for points  $p, q \in M$ , then their distance is determined by  $|a_x(p) - a_x(q)|$ , for functions  $a_x \in C^\infty(M)$  for which  $\| [D_M, a_x] \| \leq 1$ . Therefore, in this case we recover the geodesic distance on  $M$ , i.e.

$$d_{D_M \otimes 1}(n_1, n_2) = d_g(p, q).$$

However, if  $n_1$  and  $n_2$  lie in different copies of  $M$ , for instance if,  $n_1 = (p, x)$  and  $n_2 = (q, y)$ , then their distance is determined by  $|a_x(p) - a_y(q)|$  for two functions  $a_x, a_y \in C^\infty(M)$ , such that  $\| [D_M, a_x] \| \leq 1$  and  $\| [D_M, a_y] \| \leq 1$ . However, these requirements yield no restriction on  $|a_x(p) - a_y(q)|$ , so in this case the distance between  $n_1$  and  $n_2$  is infinite. We find that the space  $N$  is given by two disjoint copies of  $M$  that are separated by an infinite distance.

It should be noted that the only way in which the distance between the two copies of  $M$  could have been finite, is when the commutator  $[D_F, a]$  would be nonzero. This same commutator generates the scalar field  $\phi$  of (10.2.2), hence finiteness of the distance is related to the existence of scalar fields.

### 11.1.2 $U(1)$ Gauge Theory

We determine the gauge theory that corresponds to the almost-commutative manifold  $M \times F_x$ . The gauge group  $\mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$  from Definition 7.4 is given by the quotient  $\mathcal{U}(\mathcal{A})/\mathcal{U}(\mathcal{A}_J)$ , so if we wish to obtain a nontrivial gauge group, we need to choose  $J$  such that  $\mathcal{U}(\mathcal{A}_J) \neq \mathcal{U}(\mathcal{A})$ . Or, which in view of Example 8.2 is the same, we need to choose  $J_F$  so that  $\mathcal{U}((\mathcal{A}_F)_{J_F}) \neq \mathcal{U}(\mathcal{A}_F)$ . Looking at the form of  $J_F$  for the different (even) KO-dimensions (see the proof of Proposition 11.1), we conclude that we need KO-dimension 2 or 6. As we will see in the noncommutative description of the Standard Model in Chap. 13, the correct signature for the internal space is KO-dimension 6. Therefore, we choose to work in KO-dimension 6 as well. The almost-commutative manifold  $M \times F_x$  then has KO-dimension  $6 + 4 \bmod 8 = 2$ . This also means that we can use Definition 9.3 to calculate the fermionic action.

Summarizing, we will consider the finite space  $F_x$  given by the data

$$F_x := \left( \mathbb{C}^2, \mathbb{C}^2, D_F = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}; J_F = \begin{pmatrix} 0 & C \\ C & 0 \end{pmatrix}, \gamma_F = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right),$$

with  $C$  denoting complex conjugation, defining a real even finite space of KO-dimension 6. In the classification of irreducible geometries of Theorem 3.20, this space corresponds to the first case.

**Proposition 11.3** *The gauge group  $\mathfrak{G}(F)$  of the two-point space is given by  $U(1)$ .*

**Proof** First, note that  $\mathcal{U}(A_F) = U(1) \times U(1)$ . We now show that  $\mathcal{U}((A_F)_{J_F}) \equiv \mathcal{U}(A_F) \cap (A_F)_{J_F} \simeq U(1)$  so that the quotient  $\mathfrak{G}(F) \simeq U(1)$  as claimed. Indeed, for  $a \in \mathbb{C}^2$  to be in  $(A_F)_{J_F}$  it has to satisfy  $J_F a^* J_F = a$ . Since

$$J_F a^* J_F^{-1} = \begin{pmatrix} 0 & C \\ C & 0 \end{pmatrix} \begin{pmatrix} \bar{a}_1 & 0 \\ 0 & \bar{a}_2 \end{pmatrix} \begin{pmatrix} 0 & C \\ C & 0 \end{pmatrix} = \begin{pmatrix} a_2 & 0 \\ 0 & a_1 \end{pmatrix},$$

this is the case if and only if  $a_1 = a_2$ . Thus,  $(A_F)_{J_F} \simeq \mathbb{C}$ , whose unitary elements form the group  $U(1)$ , contained in  $\mathcal{U}(A_F)$  as the diagonal subgroup.  $\square$

In Proposition 10.12 we calculated the spectral action of an almost-commutative manifold. Before we can apply this to the two-point space, we need to find the exact form of the field  $B_\mu$ . Since we have  $(A_F)_{J_F} \simeq \mathbb{C}$ , we find  $\mathfrak{h}(F) = \mathfrak{u}((A_F)_{J_F}) \simeq i\mathbb{R}$ . From Proposition 10.3 and (10.2.7) we then see that the gauge field

$$A_\mu(x) \in i\mathfrak{g}_F = i(\mathfrak{u}(A_F)/(i\mathbb{R})) = i\mathfrak{su}(A_F) \simeq \mathbb{R}$$

becomes traceless.

Let us also explicitly derive this  $U(1)$  gauge field. An arbitrary hermitian field of the form  $A_\mu = -ia\partial_\mu b$  would be given by two  $U(1)$  gauge fields  $X_\mu^1, X_\mu^2 \in C^\infty(M, \mathbb{R})$ . However, because  $A_\mu$  only appears in the combination  $A_\mu - J_F A_\mu J_F^{-1}$ , we obtain

$$B_\mu = A_\mu - J_F A_\mu J_F^{-1} = \begin{pmatrix} X_\mu^1 & 0 \\ 0 & X_\mu^2 \end{pmatrix} - \begin{pmatrix} X_\mu^2 & 0 \\ 0 & X_\mu^1 \end{pmatrix} =: \begin{pmatrix} Y_\mu & 0 \\ 0 & -Y_\mu \end{pmatrix} = Y_\mu \otimes \gamma_F,$$

where we have defined the  $U(1)$  gauge field

$$Y_\mu := X_\mu^1 - X_\mu^2 \in C^\infty(M, \mathbb{R}) = C^\infty(M, i\mathfrak{u}(1)).$$

Thus, the fact that we only have the combination  $A_\mu - J_F A_\mu J_F^{-1}$  effectively identifies the  $U(1)$  gauge fields on the two copies of  $M$ , so that  $A_\mu$  is determined by only one  $U(1)$  gauge field. This ensures that we can take the quotient of the Lie algebra  $\mathfrak{u}(A_F)$  with  $\mathfrak{h}(F)$ . We can then write

$$A_\mu = \frac{1}{2} \begin{pmatrix} Y_\mu & 0 \\ 0 & -Y_\mu \end{pmatrix} = \frac{1}{2} Y_\mu \otimes \gamma_F,$$

which yields the same result:

$$B_\mu = A_\mu - J_F A_\mu J_F^{-1} = 2A_\mu = Y_\mu \otimes \gamma_F. \quad (11.1.2)$$

We summarize:

**Proposition 11.4** *The inner fluctuations of the almost-commutative manifold  $M \times F_X$  described above are parametrized by a  $U(1)$ -gauge field  $Y_\mu$  as*

$$D \mapsto D' = D + \gamma^\mu Y_\mu \otimes \gamma_F.$$

*The action of the gauge group  $\mathfrak{G}(M \times F_X) \simeq C^\infty(M, U(1))$  on  $D'$ , as in (10.2.8), is implemented by*

$$Y_\mu \mapsto Y_\mu - iu\partial_\mu u^*; \quad (u \in \mathfrak{G}(M \times F_X)).$$

## 11.2 Electrodynamics

Inspired by the previous section, which shows that one can use the framework of noncommutative geometry to describe a gauge theory with abelian gauge group  $U(1)$ , we proceed and try to describe the full theory of electrodynamics by an almost-commutative manifold. Our approach provides a unified description of gravity and electromagnetism, albeit at the classical level.

We have seen that the almost-commutative manifold  $M \times F_X$  describes a gauge theory with local gauge group  $U(1)$ , where the inner fluctuations of the Dirac operator provide the  $U(1)$  gauge field  $Y_\mu$ . There appear to be two problems if one wishes to use this model for a description of (classical) electrodynamics. First, by

Proposition 11.1, the finite Dirac operator  $D_F$  must vanish. However, we want our electrons to be massive, and for this purpose we need a finite Dirac operator that is non-zero.

Second, the Euclidean action for a free Dirac field is of the form

$$S = - \int i \bar{\psi} (\gamma^\mu \partial_\mu - m) \psi d^4x, \quad (11.2.1)$$

where the fields  $\psi$  and  $\bar{\psi}$  must be considered *independent variables*. Thus, we require that the fermionic action  $S_f$  should also yield two *independent* Dirac spinors. Let us write  $\{e, \bar{e}\}$  for the set of orthonormal basis vectors of  $H_F$ , where  $e$  is the basis element of  $H_F^+$  and  $\bar{e}$  of  $H_F^-$ . Note that on this basis, we have  $J_F e = \bar{e}$ ,  $J_F \bar{e} = e$ ,  $\gamma_F e = e$  and  $\gamma_F \bar{e} = -\bar{e}$ . The total Hilbert space  $\mathcal{H}$  is given by  $L^2(S) \otimes H_F$ . Since by means of  $\gamma_M$  we can also decompose  $L^2(S) = L^2(S)^+ \oplus L^2(S)^-$ , we obtain that the positive eigenspace  $\mathcal{H}^+$  of  $\gamma = \gamma_M \otimes \gamma_F$  is given by

$$\mathcal{H}^+ = L^2(S)^+ \otimes H_F^+ \oplus L^2(S)^- \otimes H_F^-.$$

Consequently, an arbitrary vector  $\xi \in \mathcal{H}^+$  can uniquely be written as

$$\xi = \psi_L \otimes e + \psi_R \otimes \bar{e},$$

for two Weyl spinors  $\psi_L \in L^2(S)^+$  and  $\psi_R \in L^2(S)^-$ . One should note here that  $\xi$  is completely determined by only one Dirac spinor  $\psi := \psi_L + \psi_R$ , instead of the required two independent spinors. Thus, the restrictions that are incorporated into the fermionic action of Definition 9.3 in fact constrain the finite space  $F_x$  too much.

### 11.2.1 The Finite Space

It turns out that both problems sketched above can be simply solved by doubling our finite-dimensional Hilbert space. Essentially, we introduce multiplicities in the Krajewski diagram that appeared in the proof of Proposition 11.1.

Thus, we start with the same algebra  $C^\infty(M, \mathbb{C}^2)$  that corresponds to the space  $N = M \times X \simeq M \sqcup M$ . The finite-dimensional Hilbert space will now be used to describe *four* particles, namely both the left-handed and the right-handed electrons and positrons. We choose the orthonormal basis  $\{e_R, e_L, \bar{e}_R, \bar{e}_L\}$  for  $H_F = \mathbb{C}^4$ , with respect to the standard inner product. The subscript  $L$  denotes left-handed particles, and the subscript  $R$  denotes right-handed particles, and we have  $\gamma_F e_L = e_L$  and  $\gamma_F e_R = -e_R$ .

We choose  $J_F$  such that it interchanges particles with their antiparticles, so  $J_F e_R = \bar{e}_R$  and  $J_F e_L = \bar{e}_L$ . We again choose the real structure such that it has KO-dimension 6, so we have  $J_F^2 = 1$  and  $J_F \gamma_F = -\gamma_F J_F$ . This last relation implies that the element  $\bar{e}_R$  is left-handed, whereas  $\bar{e}_L$  is right-handed.

The grading  $\gamma_F$  decomposes the Hilbert space  $H_F$  into  $H_F^+ \oplus H_F^-$ , where the bases of  $H_F^+$  and  $H_F^-$  are given by  $\{e_L, \overline{e_R}\}$  and  $\{e_R, \overline{e_L}\}$ , respectively. Alternatively, we can decompose the Hilbert space into  $H_e \oplus H_{\bar{e}}$ , where  $H_e$  contains the electrons  $\{e_R, e_L\}$ , and  $H_{\bar{e}}$  contains the positrons  $\{\overline{e_R}, \overline{e_L}\}$ .

The elements  $a \in A_F = \mathbb{C}^2$  now act as the following matrix with respect to the basis  $\{e_R, e_L, \overline{e_R}, \overline{e_L}\}$ :

$$a = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \rightarrow \begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_1 & 0 & 0 \\ 0 & 0 & a_2 & 0 \\ 0 & 0 & 0 & a_2 \end{pmatrix}. \quad (11.2.2)$$


Note that this action commutes with the grading, as it should. We can also easily check that  $[a, b^0] = 0$  for  $b^0 := J_F b^* J_F^{-1}$ , since both the left and the right action are given by diagonal matrices. For now, we still take  $D_F = 0$ , and hence the order one condition is trivially satisfied. We have therefore obtained the following result:

**Proposition 11.5** *The data*

$$\left( \mathbb{C}^2, \mathbb{C}^2, D_F = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}; J_F = \begin{pmatrix} 0 & C \\ C & 0 \end{pmatrix}, \gamma_F = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right)$$

define a real even spectral triple of  $KO$ -dimension 6.

This can be summarized by the following Krajewski diagram, with two nodes (of opposite grading) of multiplicity two:



### 11.2.2 A Non-trivial Finite Dirac Operator

Let us now consider the possibilities for adding a non-zero Dirac operator to the finite space  $F_{ED}$ . From the above Krajewski diagram, it can be easily seen that the only possible edges exist between the multiple vertices. That is, the only possible Dirac operator depends on one complex parameter and is given by

$$D_F = \begin{pmatrix} 0 & d & 0 & 0 \\ \bar{d} & 0 & 0 & 0 \\ 0 & 0 & 0 & \bar{d} \\ 0 & 0 & d & 0 \end{pmatrix}. \quad (11.2.3)$$

From here on, we will consider the finite space  $F_{ED}$  given by

$$F_{ED} := (\mathbb{C}^2, \mathbb{C}^4, D_F; J_F, \gamma_F).$$

### 11.2.3 The Almost-Commutative Manifold

Taking the product with the canonical triple, the almost-commutative manifold  $M \times F_{ED}$  (of KO-dimension 2) under consideration is given by the spectral triple

$$\begin{aligned} M \times F_{ED} := \\ (C^\infty(M, \mathbb{C}^2), L^2(S) \otimes \mathbb{C}^4, D_M \otimes 1 + \gamma_M \otimes D_F; J_M \otimes J_F, \gamma_M \otimes \gamma_F). \end{aligned} \quad (11.2.4)$$

As in Sect. 11.1, the algebra decomposes as

$$C^\infty(M, \mathbb{C}^2) = C^\infty(M) \oplus C^\infty(M),$$

and we now decompose the Hilbert space as

$$\mathcal{H} = (L^2(S) \otimes H_e) \oplus (L^2(S) \otimes H_{\bar{e}}).$$

The action of the algebra on  $\mathcal{H}$ , given by (11.2.2), is then such that one component of the algebra acts on the electron fields  $L^2(S) \otimes H_e$ , and the other component acts on the positron fields  $L^2(S) \otimes H_{\bar{e}}$ .

The derivation of the gauge group for  $F_{ED}$  is exactly the same as in Proposition 11.3, so again we have the finite gauge group  $\mathfrak{G}(F) \simeq U(1)$ . The field  $B_\mu := A_\mu - J_F A_\mu J_F^{-1}$  now takes the form

$$B_\mu = \begin{pmatrix} Y_\mu & 0 & 0 & 0 \\ 0 & Y_\mu & 0 & 0 \\ 0 & 0 & -Y_\mu & 0 \\ 0 & 0 & 0 & -Y_\mu \end{pmatrix} \quad \text{for } Y_\mu(x) \in \mathbb{R}. \quad (11.2.5)$$

Thus, we again obtain a single  $U(1)$  gauge field  $Y_\mu$ , carrying an action of the gauge group  $\mathfrak{G}(M \times F_{ED}) \simeq C^\infty(M, U(1))$  (as in Proposition 11.4).

As mentioned before, our space  $N$  consists of two copies of  $M$  and if  $D_F = 0$  the distance between these two copies is infinite (see Remark 11.2). This time we have introduced a non-zero Dirac operator, but it commutes with the algebra, i.e.  $[D_F, a] = 0$  for all  $a \in \mathcal{A}$ . Therefore, the distance between the two copies of  $M$  is still infinite.

To summarize, the  $U(1)$  gauge theory arises from the geometric space  $N = M \sqcup M$  as follows. On one copy of  $M$ , we have the vector bundle  $S \otimes (M \times H_e)$ , and on the other copy we have the vector bundle  $S \otimes (M \times H_{\bar{e}})$ . The gauge fields on each copy of  $M$  are identified with each other. The electrons  $e$  and positrons  $\bar{e}$  are then both

coupled to the same gauge field, and as such the gauge field provides an interaction between electrons and positrons. For comparison with Kaluza–Klein theories, note the different role that is played by the internal space.

### 11.2.4 The Spectral Action

We are now ready to explicitly calculate the Lagrangian that corresponds to the almost-commutative manifold  $M \times F_{ED}$ , and we will show that this yields the usual Lagrangian for electrodynamics (on a curved background manifold), as well as a purely gravitational Lagrangian. It consists of the spectral action  $S_b$  of Definition 9.1 and the fermionic action  $S_f$  of Definition 9.3, which we calculate separately (here and in the next section).

The spectral action for an almost-commutative manifold has been calculated in Proposition 10.12, and we only need to insert the fields  $B_\mu$  (given by (11.2.5)) and  $\Phi = D_F$ . We obtain the following result:

**Proposition 11.6** *The spectral action of the almost-commutative manifold  $M \times F_{ED}$  defined in (11.2.4) is given by*

$$\text{Tr} \left( f \left( \frac{D_\omega}{\Lambda} \right) \right) \sim \int_M \mathcal{L}(g_{\mu\nu}, Y_\mu) \sqrt{g} d^4x + O(\Lambda^{-1}),$$

with Lagrangian

$$\mathcal{L}(g_{\mu\nu}, Y_\mu) := 4\mathcal{L}_M(g_{\mu\nu}) + \mathcal{L}_Y(Y_\mu) + \mathcal{L}_\phi(g_{\mu\nu}, d).$$

Here  $\mathcal{L}_M(g_{\mu\nu})$  is defined in Proposition 10.10; the term  $\mathcal{L}_Y$  gives the kinetic term of the  $U(1)$  gauge field  $Y_\mu$  as

$$\mathcal{L}_Y(Y_\mu) := \frac{f(0)}{6\pi^2} Y_{\mu\nu} Y^{\mu\nu},$$

where the curvature  $Y_{\mu\nu}$  of the field  $Y_\mu$  is given by

$$Y_{\mu\nu} := \partial_\mu Y_\nu - \partial_\nu Y_\mu.$$

The scalar potential  $\mathcal{L}_\phi$  (ignoring the boundary term) gives two constant terms which add to the cosmological constant, plus an extra contribution to the Einstein–Hilbert action:

$$\mathcal{L}_\phi(g_{\mu\nu}) := -\frac{2f_2\Lambda^2}{\pi^2} |d|^2 + \frac{f(0)}{2\pi^2} |d|^4 + \frac{f(0)}{12\pi^2} s |d|^2,$$

where the constant  $d$  originates from (11.2.3).

**Proof** The trace over the Hilbert space  $\mathbb{C}^4$  yields an overall factor  $N = 4$ . The field  $B_\mu$  is given by (11.2.5), and we obtain  $\text{Tr}(F_{\mu\nu} F^{\mu\nu}) = 4Y_{\mu\nu} Y^{\mu\nu}$ . Inserting this into Proposition 10.12 provides the Lagrangian  $\mathcal{L}_Y$ . In addition, we have  $\Phi^2 = D_F^2 = |d|^2$ , and the scalar-field Lagrangian  $\mathcal{L}_\phi$  only yields extra numerical contributions to the cosmological constant and the Einstein–Hilbert action.  $\square$

### 11.2.5 The Fermionic Action

We have written the set of basis vectors of  $H_F$  as  $\{e_R, e_L, \overline{e_R}, \overline{e_L}\}$ , and the subspaces  $H_F^+$  and  $H_F^-$  are spanned by  $\{e_L, \overline{e_R}\}$  and  $\{e_R, \overline{e_L}\}$ , respectively. The total Hilbert space  $\mathcal{H}$  is given by  $L^2(S) \otimes H_F$ . Since we can also decompose

$$L^2(S) = L^2(S)^+ \oplus L^2(S)^-$$

by means of  $\gamma_M$ , we obtain for the +1-eigenspace of  $\gamma_M \otimes \gamma_F$ :

$$\mathcal{H}^+ = L^2(S)^+ \otimes H_F^+ \oplus L^2(S)^- \otimes H_F^-.$$

A spinor  $\psi \in L^2(S)$  can be decomposed as  $\psi = \psi_L + \psi_R$ . Each subspace  $H_F^\pm$  is now spanned by two basis vectors. A generic element of the tensor product of two spaces consists of sums of tensor products, so an arbitrary vector  $\xi \in \mathcal{H}^+$  can be uniquely written as

$$\xi = \chi_R \otimes e_R + \chi_L \otimes e_L + \psi_L \otimes \overline{e_R} + \psi_R \otimes \overline{e_L}, \quad (11.2.6)$$

for Weyl spinors  $\chi_L, \psi_L \in L^2(S)^+$  and  $\chi_R, \psi_R \in L^2(S)^-$ . Note that this vector  $\xi \in \mathcal{H}^+$  is now completely determined by two Dirac spinors  $\chi := \chi_L + \chi_R$  and  $\psi := \psi_L + \psi_R$ .

**Proposition 11.7** *The fermionic action of the almost-commutative manifold  $M \times F_{ED}$  defined in (11.2.4), is given by*

$$S_f = -i(J_M \tilde{\chi}, \gamma^\mu (\nabla_\mu^S - iY_\mu) \tilde{\psi}) + (J_M \tilde{\chi}_L, \bar{d} \tilde{\psi}_L) - (J_M \tilde{\chi}_R, d \tilde{\psi}_R).$$

**Proof** The fluctuated Dirac operator is given by

$$D_\omega = D_M \otimes 1 + \gamma^\mu \otimes B_\mu + \gamma_M \otimes D_F.$$

An arbitrary  $\xi \in \mathcal{H}^+$  has the form of (11.2.6), from which we obtain the following expressions:

$$\begin{aligned}
J\xi &= J_M \chi_R \otimes \overline{e_R} + J_M \chi_L \otimes \overline{e_L} + J_M \psi_L \otimes e_R + J_M \psi_R \otimes e_L, \\
(D_M \otimes 1)\xi &= D_M \chi_R \otimes e_R + D_M \chi_L \otimes e_L + D_M \psi_L \otimes \overline{e_R} + D_M \psi_R \otimes \overline{e_L}, \\
(\gamma^\mu \otimes B_\mu)\xi &= \gamma^\mu \chi_R \otimes Y_\mu e_R + \gamma^\mu \chi_L \otimes Y_\mu e_L - \gamma^\mu \psi_L \otimes Y_\mu \overline{e_R} - \gamma^\mu \psi_R \otimes Y_\mu \overline{e_L}, \\
(\gamma_M \otimes D_F)\xi &= \gamma_M \chi_L \otimes \overline{de_R} + \gamma_M \chi_R \otimes de_L + \gamma_M \psi_R \otimes d\overline{e_R} + \gamma_M \psi_L \otimes d\overline{e_L}.
\end{aligned}$$

We decompose the fermionic action into the three terms

$$\frac{1}{2}(J\tilde{\xi}, D_\omega \tilde{\xi}) = \frac{1}{2}(J\tilde{\xi}, (D_M \otimes 1)\tilde{\xi}) + \frac{1}{2}(J\tilde{\xi}, (\gamma^\mu \otimes B_\mu)\tilde{\xi}) + \frac{1}{2}(J\tilde{\xi}, (\gamma_M \otimes D_F)\tilde{\xi}),$$

and then continue to calculate each term separately. The first term is given by

$$\begin{aligned}
\frac{1}{2}(J\tilde{\xi}, (D_M \otimes 1)\tilde{\xi}) &= \frac{1}{2}(J_M \tilde{\chi}_R, D_M \tilde{\psi}_L) + \frac{1}{2}(J_M \tilde{\chi}_L, D_M \tilde{\psi}_R) \\
&\quad + \frac{1}{2}(J_M \tilde{\psi}_L, D_M \tilde{\chi}_R) + \frac{1}{2}(J_M \tilde{\psi}_R, D_M \tilde{\chi}_L).
\end{aligned}$$

Using the facts that  $D_M$  changes the chirality of a Weyl spinor, and that the subspaces  $L^2(S)^+$  and  $L^2(S)^-$  are orthogonal, we can rewrite this term as

$$\frac{1}{2}(J\tilde{\xi}, (D_M \otimes 1)\tilde{\xi}) = \frac{1}{2}(J_M \tilde{\chi}, D_M \tilde{\psi}) + \frac{1}{2}(J_M \tilde{\psi}, D_M \tilde{\chi}).$$

Using the symmetry of the form  $(J_M \tilde{\chi}, D_M \tilde{\psi})$ , we obtain

$$\frac{1}{2}(J\tilde{\xi}, (D_M \otimes 1)\tilde{\xi}) = (J_M \tilde{\chi}, D_M \tilde{\psi}) = -i(J_M \tilde{\chi}, \gamma^\mu \nabla_\mu^S \tilde{\psi}).$$

Note that the factor  $\frac{1}{2}$  has now disappeared from the result, which is the reason why this factor had to be included in the definition of the fermionic action. The second term is given by

$$\begin{aligned}
\frac{1}{2}(J\tilde{\xi}, (\gamma^\mu \otimes B_\mu)\tilde{\xi}) &= -\frac{1}{2}(J_M \tilde{\chi}_R, \gamma^\mu Y_\mu \tilde{\psi}_L) - \frac{1}{2}(J_M \tilde{\chi}_L, \gamma^\mu Y_\mu \tilde{\psi}_R) \\
&\quad + \frac{1}{2}(J_M \tilde{\psi}_L, \gamma^\mu Y_\mu \tilde{\chi}_R) + \frac{1}{2}(J_M \tilde{\psi}_R, \gamma^\mu Y_\mu \tilde{\chi}_L).
\end{aligned}$$

In a similar manner, we obtain

$$\frac{1}{2}(J\tilde{\xi}, (\gamma^\mu \otimes B_\mu)\tilde{\xi}) = -(J_M \tilde{\chi}, \gamma^\mu Y_\mu \tilde{\psi}),$$

where we have used the anti-symmetry of the form  $(J_M \tilde{\chi}, \gamma^\mu Y_\mu \tilde{\psi})$ . The third term is given by

$$\begin{aligned} \frac{1}{2}(J\tilde{\xi}, (\gamma_M \otimes D_F)\tilde{\xi}) &= \frac{1}{2}(J_M \tilde{\chi}_R, d\gamma_M \tilde{\psi}_R) + \frac{1}{2}(J_M \tilde{\chi}_L, \bar{d}\gamma_M \tilde{\psi}_L) \\ &\quad + \frac{1}{2}(J_M \tilde{\psi}_L, \bar{d}\gamma_M \tilde{\chi}_L) + \frac{1}{2}(J_M \tilde{\psi}_R, d\gamma_M \tilde{\chi}_R). \end{aligned}$$

The bilinear form  $(J_M \tilde{\chi}, \gamma_M \tilde{\psi})$  is again symmetric in the Grassmann variables  $\tilde{\chi}$  and  $\tilde{\psi}$ , but we now face the extra complication that two terms contain the parameter  $d$ , while the other two terms contain  $\bar{d}$ . Therefore we are left with two distinct terms:

$$\frac{1}{2}(J\tilde{\xi}, (\gamma_M \otimes D_F)\tilde{\xi}) = (J_M \tilde{\chi}_L, \bar{d}\tilde{\psi}_L) - (J_M \tilde{\chi}_R, d\tilde{\psi}_R). \quad \square$$

**Remark 11.8** It is interesting to note that the fermions acquire mass terms without being coupled to a scalar field. However, it seems that we obtain a complex mass parameter  $d$ , where we would desire a real parameter  $m$ . Simply requiring that our result should reproduce (11.2.1), we will therefore choose  $d := -im$ , so that

$$(J_M \tilde{\chi}_L, \bar{d}\tilde{\psi}_L) - (J_M \tilde{\chi}_R, d\tilde{\psi}_R) = i(J_M \tilde{\chi}, m\tilde{\psi}).$$

The results obtained in this section can now be summarized into the following theorem.

**Theorem 11.9** *The full Lagrangian of the almost-commutative manifold  $M \times F_{ED}$  as defined in Eq. (11.2.4), can be written as the sum of a purely gravitational Lagrangian,*

$$\mathcal{L}_{\text{grav}}(g_{\mu\nu}) = 4\mathcal{L}_M(g_{\mu\nu}) + \mathcal{L}_\phi(g_{\mu\nu}),$$

and a Lagrangian for electrodynamics,

$$\mathcal{L}_{ED} = -i \left\langle J_M \tilde{\chi}, (\gamma^\mu (\nabla_\mu^S - iY_\mu) - m)\tilde{\psi} \right\rangle + \frac{f(0)}{6\pi^2} Y_{\mu\nu} Y^{\mu\nu}.$$

**Proof** The spectral action  $S_b$  and the fermionic action  $S_f$  are given by Propositions 11.6 and 11.7. This immediately yields  $\mathcal{L}_{\text{grav}}$ . To obtain  $\mathcal{L}_{ED}$ , we need to rewrite the fermionic action  $S_f$  as the integral over a Lagrangian. The inner product  $(\cdot, \cdot)$  on the Hilbert space  $L^2(S)$  is given by

$$(\xi, \psi) = \int_M \langle \xi, \psi \rangle \sqrt{g} d^4x,$$

where the hermitian pairing  $\langle \cdot, \cdot \rangle$  is given by the pointwise inner product on the fibres. Choosing  $d = -im$  as in Remark 11.8, we can then rewrite the fermionic action into

$$S_f = - \int_M i \left\langle J_M \tilde{\chi}, (\gamma^\mu (\nabla_\mu^S - i Y_\mu) - m) \tilde{\psi} \right\rangle \sqrt{g} d^4x. \quad \square$$

### 11.2.6 Fermionic Degrees of Freedom

To conclude this chapter, let us make a final remark on the fermionic degrees of freedom in the Lagrangian derived above. We refer the reader to Appendix 11.3 for a short introduction to Grassmann variables and Grassmann integration.

As mentioned in Note 3 of this chapter below, the number of degrees of freedom of the fermion fields in the fermionic action is related to the restrictions that are incorporated into the definition of the fermionic action. These restrictions make sure that in this case we obtain two independent Dirac spinors in the fermionic action.

In fact, in quantum field theory one would consider the functional integral of  $e^S$  over the fields. We hence consider the case that  $\mathfrak{A}$  is the antisymmetric bilinear form on  $\mathcal{H}^+$  given by

$$\mathfrak{A}(\xi, \zeta) := (J\xi, D_\omega \zeta), \quad \text{for } \xi, \zeta \in \mathcal{H}^+,$$

and  $\mathfrak{A}'$  is the bilinear form on  $L^2(S)$  given by

$$\mathfrak{A}'(\chi, \psi) := -i \left( J_M \chi, (\gamma^\mu (\nabla_\mu^S - i Y_\mu) - m) \psi \right), \quad \text{for } \chi, \psi \in L^2(S).$$

We have shown in Proposition 11.7 that for  $\xi = \chi_L \otimes e_L + \chi_R \otimes e_R + \psi_L \otimes \overline{e_L} + \psi_R \otimes \overline{e_R}$ , where we can define two Dirac spinors by  $\chi := \chi_L + \chi_R$  and  $\psi := \psi_L + \psi_R$ , we obtain

$$\frac{1}{2} \mathfrak{A}(\xi, \xi) = \mathfrak{A}'(\chi, \psi).$$

Using the Grassmann integrals of (11.3.1) and (11.3.2), we then obtain for the bilinear forms  $\mathfrak{A}$  and  $\mathfrak{A}'$  the equality

$$\text{Pf}(\mathfrak{A}) = \int e^{\frac{1}{2} \mathfrak{A}(\tilde{\xi}, \tilde{\xi})} D[\tilde{\xi}] = \int e^{\mathfrak{A}'(\tilde{\chi}, \tilde{\psi})} D[\tilde{\psi}, \tilde{\chi}] = \det(\mathfrak{A}').$$

### 11.3 Grassmann Variables, Grassmann Integration and Pfaffians

We will give a short introduction to Grassmann variables, and use those to find the relation between the Pfaffian and the determinant of an antisymmetric matrix.

For a set of anti-commuting Grassmann variables  $\theta_i$ , we have  $\theta_i \theta_j = -\theta_j \theta_i$ , and in particular,  $\theta_i^2 = 0$ . On these Grassmann variables  $\theta_j$ , we define an integral by

$$\int 1 d\theta_j = 0, \quad \int \theta_j d\theta_j = 1.$$

If we have a Grassmann vector  $\theta$  consisting of  $N$  components, we define the integral over  $D[\theta]$  as the integral over  $d\theta_1 \cdots d\theta_N$ . Suppose we have two Grassmann vectors  $\eta$  and  $\theta$  of  $N$  components. We then define the integration element as  $D[\eta, \theta] = d\eta_1 d\theta_1 \cdots d\eta_N d\theta_N$ .

Consider the Grassmann integral over a function of the form  $e^{\theta^T \mathfrak{A} \eta}$  for Grassmann vectors  $\theta$  and  $\eta$  of  $N$  components. The  $N \times N$ -matrix  $\mathfrak{A}$  can be considered as a bilinear form on these Grassmann vectors. In the case where  $\theta$  and  $\eta$  are independent variables, we find

$$\int e^{\theta^T \mathfrak{A} \eta} D[\eta, \theta] = \det \mathfrak{A}, \quad (11.3.1)$$

where the determinant of  $\mathfrak{A}$  is given by the formula

$$\det(\mathfrak{A}) = \frac{1}{N!} \sum_{\sigma, \tau \in S_N} (-1)^{|\sigma| + |\tau|} \mathfrak{A}_{\sigma(1)\tau(1)} \cdots \mathfrak{A}_{\sigma(N)\tau(N)},$$

in which  $S_N$  denotes the set of all permutations of  $\{1, 2, \dots, N\}$ . Now let us assume that  $\mathfrak{A}$  is an antisymmetric  $N \times N$ -matrix  $\mathfrak{A}$  for  $N = 2l$ . If we then take  $\theta = \eta$ , we find

$$\int e^{\frac{1}{2} \eta^T \mathfrak{A} \eta} D[\eta] = \text{Pf}(\mathfrak{A}), \quad (11.3.2)$$

where the *Pfaffian* of  $\mathfrak{A}$  is given by

$$\text{Pf}(\mathfrak{A}) = \frac{(-1)^l}{2^l l!} \sum_{\sigma \in S_{2l}} (-1)^{|\sigma|} \mathfrak{A}_{\sigma(1)\sigma(2)} \cdots \mathfrak{A}_{\sigma(2l-1)\sigma(2l)}.$$

Finally, using these Grassmann integrals, one can show that the determinant of a  $2l \times 2l$  skew-symmetric matrix  $\mathfrak{A}$  is the square of the Pfaffian:

$$\det \mathfrak{A} = \text{Pf}(\mathfrak{A})^2.$$

So, by simply considering one instead of two independent Grassmann variables in the Grassmann integral of  $e^{\theta^T \mathfrak{A} \eta}$ , we are in effect taking the square root of a determinant.

## Notes

### Section 11.1. The Two-Point Space

1. The two-point space was first studied in [1, 2].
2. The need for KO-dimension 6 for the noncommutative description of the Standard Model has been observed independently by Barrett [3] and Connes [4].
3. In [5, Chap. 9] a proof is given for the claim that the inner fluctuation  $\omega + J\omega J^{-1}$  vanishes for commutative algebras. The proof is based on the assumption that the left and right action can be identified, i.e.  $a = a^0$ , for a commutative algebra. Though this holds in the case of the canonical triple describing a spin manifold, it need not be true for arbitrary commutative algebras. Indeed, the almost-commutative manifold  $M \times F_X$  provides a counter-example.  
What we can say about a commutative algebra, is that there exist no non-trivial inner automorphisms. Thus, it is an important insight that the gauge group  $\mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$  from Definition 7.4 is larger than the group of inner automorphisms, so that a commutative algebra may still lead to a non-trivial (necessarily abelian) gauge group.
4. It is shown in [6] that one can also obtain abelian gauge theories from a one-point space when one works with real algebras (cf. Sect. 3.3).

### Section 11.2. Electrodynamics

5. Earlier attempts at a unified description of gravity and electromagnetism originate from the work of Kaluza [7] and Klein [8] in the 1920s. In their approach, a new (compact) fifth dimension is added to the 4-dimensional spacetime  $M$ . The additional components in the 5-dimensional metric tensor are then identified with the electromagnetic gauge potential. Subsequently, it can be shown that the Einstein equations of the 5-dimensional spacetime can be reduced to the Einstein equations plus the Maxwell equations on 4-dimensional spacetime.
6. An interesting question that appears in the context of this Chapter is whether it is possible to describe the *abelian Higgs mechanism* (see e.g. [9, Sect. 8.3]) by an almost-commutative manifold. As already noticed, for  $M \times F_{ED}$  no scalar fields  $\Phi$  are generated since  $A_F$  commutes with  $D_F$ . In terms of the Krajewski diagram for  $M \times F_{ED}$ ,

$$\begin{array}{ccc} & 1 & 1 \\ & \circ & \circ \\ 1^\circ & & \circ \\ & \circ & \circ \end{array}$$

it follows that a component that runs counterdiagonally fails on the first-order condition (cf. Lemma 3.10). One is therefore tempted to look at the generalization of inner fluctuations to real spectral triples that do not necessarily satisfy the first-order condition, as was proposed in [10]. This generalization is crucial in the applications to Pati–Salam unification (see Chap. 15 below) but also in the present case one can show that non-zero off-diagonal components in (11.2.3) then generate a scalar field for which the spectral action yields a spontaneous breaking of the abelian gauge symmetry.

### Section 11.3. Grassmann Variables, Grassmann Integration and Pfaffians

7. For more details we refer the reader to [11].

## References

- Connes, A.: Essay on physics and noncommutative geometry. In: The interface of mathematics and particle physics (Oxford, 1988), Institute of Mathematics and its Applications Conference Series. New Series, vol. 24, pp. 9–48. Oxford University Press, New York (1990)
- Connes, A., Lott, J.: Particle models and noncommutative geometry. *Nucl. Phys. Proc. Suppl.* **18B**, 29–47 (1991)
- Barrett, J.W.: A Lorentzian version of the non-commutative geometry of the standard model of particle physics. *J. Math. Phys.* **48**, 012303 (2007)
- Connes, A.: On the foundations of noncommutative geometry. In: The Unity of Mathematics. Progress in Mathematics, vol. 244, pp. 173–204. Birkhäuser Boston, Boston (2006)
- Landi, G.: An Introduction to Noncommutative Spaces and their Geometry. Springer (1997)
- Bhowmick, J., D’Andrea, F., Das, B., Dabrowski, L.: Quantum gauge symmetries in Noncommutative Geometry, [arXiv:1112.3622](https://arxiv.org/abs/1112.3622)
- Kaluza, T.: Zum Unitätsproblem in der Physik. *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)*, vol. 1921, pp. 966–972 (1921)
- Klein, O.: Quantentheorie und fünfdimensionale relativitätstheorie. *Z. Phys.* **37**, 895–906 (1926)
- Kane, G.L.: Modern Elementary Particle Physics. Perseus (1993)
- Chamseddine, A.H., Connes, A., Van Suijlekom, W.D.: Inner fluctuations in noncommutative geometry without the first order condition. *J. Geom. Phys.* **73**, 222–234 (2013)
- Berezin, F.A.: The Method of Second Quantization. Translated from the Russian by Nobumichi Mugibayashi and Alan Jeffrey. Pure and Applied Physics, vol. 24. Academic Press, New York (1966)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book’s Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book’s Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 12

## The Noncommutative Geometry of Yang–Mills Fields



In this chapter we generalize the noncommutative description of Yang–Mills theory to topologically non-trivial gauge configurations.

### 12.1 Spectral Triple Obtained from an Algebra Bundle

Recall from Examples 10.4 and 10.5 that topologically trivial Yang–Mills gauge theory can be described by the almost-commutative manifold

$$M \times F_{YM} = (C^\infty(M) \otimes M_N(\mathbb{C}), L^2(S) \otimes M_N(\mathbb{C}), D_M \otimes 1; J_M \otimes (\cdot)^*, \gamma_M \otimes 1).$$

In fact, the tensor product of  $C^\infty(M)$  with the matrix algebra  $M_N(\mathbb{C})$  appearing here is equivalent to restricting the gauge theory to be defined on a *trivial* vector bundle. Indeed,  $C^\infty(M) \otimes M_N(\mathbb{C})$  is the algebra of smooth sections of the trivial algebra bundle  $M \times M_N(\mathbb{C})$  on  $M$ . For the topologically non-trivial case, this suggests considering an arbitrary  $*$ -algebra bundle with fiber  $M_N(\mathbb{C})$ . We work in a slightly more general setting more general  $*$ -algebras are allowed.

Thus, let  $\mathfrak{B}$  be some locally trivial  $*$ -algebra bundle whose fibers are copies of a fixed (finite-dimensional)  $*$ -algebra  $A$ . Furthermore, we require that for each  $x$  the fiber  $\mathfrak{B}_x$  is endowed with a faithful tracial state  $\tau_x$ , such that for each  $s \in \Gamma^\infty(\mathfrak{B})$  the function  $x \mapsto \tau_x s(x)$  is smooth. The corresponding Hilbert–Schmidt inner product in the fiber  $\mathfrak{B}_x$  that is induced by  $\tau_x$  is denoted by  $(\cdot, \cdot)_{\mathfrak{B}_x}$ . Consequently, the  $C^\infty(M)$ -valued form

$$\langle \cdot, \cdot \rangle_{\mathfrak{B}} : \Gamma^\infty(\mathfrak{B}) \times \Gamma^\infty(\mathfrak{B}) \rightarrow C^\infty(M); \quad \langle s, t \rangle_{\mathfrak{B}}(x) = (s(x), t(x))_{\mathfrak{B}_x}$$

is a hermitian structure on the  $C^\infty(M)$ -module  $\Gamma^\infty(\mathfrak{B})$ , satisfying the conditions of Proposition 7.14.

As in the previous chapters, we assume that  $M$  is a compact Riemannian spin manifold on which  $S \rightarrow M$  is a spinor bundle and  $D_M = -ic \circ \nabla^S$  is the Dirac operator. Combining the inner product on spinors with the above hermitian structure naturally induces the following inner product on  $\Gamma^\infty(\mathfrak{B} \otimes S)$ :

$$(\xi_1, \xi_2) := \int_M (\xi_1(x), \xi_2(x))_{\mathfrak{B}_x \otimes S_x}; \quad (\xi_1, \xi_2 \in \Gamma^\infty(\mathfrak{B} \otimes S)), \quad (12.1.1)$$

turning it into a pre-Hilbert space. Its completion with respect to the norm induced by this inner product consists of all square-integrable sections of  $\mathfrak{B} \otimes S$ , and is denoted by  $L^2(\mathfrak{B} \otimes S)$ .

**Remark 12.1** Note that we can identify  $\Gamma^\infty(\mathfrak{B}) \otimes_{C^\infty(M)} \Gamma^\infty(S)$  with  $\Gamma^\infty(\mathfrak{B} \otimes S)$  as  $C^\infty(M)$ -modules. In what follows, we will use this identification without further notice. The above inner product (12.1.1) can then be written as

$$(s_1 \otimes \psi_1, s_2 \otimes \psi_2) = (\psi_1, \langle s_1, s_2 \rangle_{\mathfrak{B}} \psi_2),$$

where  $\langle s_1, s_2 \rangle_{\mathfrak{B}} \in C^\infty(M)$  acts on  $\Gamma^\infty(S)$  by pointwise multiplication.

**Theorem 12.2** *In the above notation, let  $\nabla^{\mathfrak{B}}$  be a hermitian connection (with respect to the Hilbert–Schmidt inner product) on the  $*$ -algebra bundle  $\mathfrak{B}$  and let  $D_{\mathfrak{B}} = -i\gamma^\mu(\nabla_\mu^{\mathfrak{B}} \otimes 1 + 1 \otimes \nabla_\mu^S)$  be the twisted Dirac operator on  $\mathfrak{B} \otimes S$ . Then*

$$(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_{\mathfrak{B}})$$

is a spectral triple.

**Proof** First, it is obvious that fiberwise multiplication of  $a \in \Gamma^\infty(\mathfrak{B})$  on  $\Gamma^\infty(\mathfrak{B} \otimes S)$  extends to a bounded operator on  $L^2(\mathfrak{B} \otimes S)$ , since

$$\begin{aligned} \|as \otimes \psi\|^2 &= \int_M (\psi(x), (a(x)s(x), a(x)s(x))_{\mathfrak{B}_x} \psi(x))_{S_x} dx \\ &\leq \sup_{x \in M} \{\|a(x)\|_x^2\} \|s \otimes \psi\|^2. \end{aligned}$$

Here  $\|\cdot\|_x$  denotes the fiberwise operator  $C^*$ -norm. Since  $M$  is a compact manifold, the compactness of the resolvent follows from ellipticity of the twisted Dirac operator  $D_{\mathfrak{B}}$ . Moreover, the commutator  $[D_{\mathfrak{B}}, a]$  is bounded for  $a \in \Gamma^\infty(\mathfrak{B})$  since  $D_{\mathfrak{B}}$  is a first-order differential operator. More precisely, in local coordinates one computes

$$[D_{\mathfrak{B}}, a](s \otimes \psi) = -i(\partial_\mu a + [\omega_\mu^{\mathfrak{B}}, a])s \otimes \gamma^\mu \psi,$$

where  $\nabla_\mu^{\mathfrak{B}} = \partial_\mu + \omega_\mu^{\mathfrak{B}}$ . This operator is bounded on  $L^2(\mathfrak{B} \otimes S)$ , provided  $a$  is differentiable and  $\omega_\mu^{\mathfrak{B}}$  is smooth.  $\square$

Next, we would like to extend our construction to arrive at a real spectral triple. For this, we introduce an anti-linear operator on  $L^2(\mathfrak{B} \otimes S)$  of the form

$$J(s \otimes \psi) = s^* \otimes J_M \psi,$$

with  $J_M$  charge conjugation on  $M$  as in Definition 4.13. For this operator to be a real structure on our spectral triple  $(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_{\mathfrak{B}})$ , we need some extra conditions on the connection  $\nabla^{\mathfrak{B}}$  on  $\mathfrak{B}$ .

**Definition 12.3** Let  $\mathfrak{B}$  be a  $*$ -algebra bundle over a manifold  $M$ . A  $*$ -algebra connection  $\nabla$  on  $\mathfrak{B}$  is a connection on  $\mathfrak{B}$  that satisfies

$$\nabla(st) = s\nabla t + (\nabla s)t, \quad (\nabla s)^* = \nabla s^*; \quad (s, t \in \Gamma^\infty(\mathfrak{B})).$$

If  $\mathfrak{B}$  is a hermitian  $*$ -algebra bundle and  $\nabla$  is also a hermitian connection, then  $\nabla$  is called a *hermitian*  $*$ -algebra connection.

**Lemma 12.4** Every locally trivial hermitian  $*$ -algebra bundle  $\mathfrak{B}$  defined over a compact space  $M$  admits a hermitian  $*$ -algebra connection.

**Proof** Let  $\{U_i\}$  be a finite open covering of  $M$  such that  $\mathfrak{B}$  is trivialized over  $U_i$  for each  $i$ . Then on each  $U_i$  there exists a hermitian  $*$ -algebra connection  $\nabla_i$ , for instance the trivial connection  $d$  on  $U_i$ . Now, let  $\{f_i\}$  be a partition of unity subordinate to the open covering  $\{U_i\}$  (note that all  $f_i$  are real-valued). Then the linear map  $\nabla$  defined by

$$(\nabla s)(x) = \sum_i f_i(x)(\nabla_i s)(x); \quad (x \in M)$$

is a hermitian  $*$ -algebra connection on  $\Gamma^\infty(\mathfrak{B})$ . □

**Remark 12.5** The fact that locally, i.e. on some trivializing neighborhood, the exterior derivative  $d$  is a hermitian  $*$ -algebra connection shows that on such a local chart every hermitian  $*$ -algebra connection is of the form

$$d + \omega^{\mathfrak{B}},$$

where  $\omega^{\mathfrak{B}}$  is a real connection one-form with values in the real Lie algebra of  $*$ -derivations of the fiber that are anti-hermitian with respect to the inner product on the fiber. For instance, when the fiber is the  $*$ -algebra  $M_N(\mathbb{C})$  endowed with the Hilbert–Schmidt inner product, this Lie algebra is precisely  $\text{ad}(u(N)) \cong su(N)$ .

**Theorem 12.6** In addition to the conditions of Theorem 12.2, suppose that  $\nabla^{\mathfrak{B}}$  is a hermitian  $*$ -algebra connection and set  $\gamma = 1 \otimes \gamma_M$  as a self-adjoint operator on  $L^2(\mathfrak{B} \otimes S)$ . Then

$$(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_{\mathfrak{B}}; J, \gamma)$$

is a real and even spectral triple whose KO-dimension is equal to the dimension of  $M$ .

**Proof** First of all, we check that  $J$  is anti-unitary:

$$\begin{aligned} (J(s \otimes \psi), J(t \otimes \eta)) &= (J_M \psi, \langle s^*, t^* \rangle J_M \eta) = (J_M \psi, J_M \overline{\langle s^*, t^* \rangle \eta}) \\ &= (\overline{\langle s^*, t^* \rangle \eta}, \psi) = (\langle s, t \rangle \eta, \psi) = (t \otimes \eta, s \otimes \psi), \end{aligned}$$

where we used in the second step that  $J_M f = \bar{f} J_M$  for every  $f \in C^\infty(M)$ , in the third step that  $J_M$  is anti-unitary, and in the fourth step that  $\langle s, t \rangle = \langle t^*, s^* \rangle$  (by definition of the hermitian structure as a fiberwise trace). Moreover, if  $J_M^2 = \epsilon$  it follows that  $J^2 = \epsilon$ .

We next establish  $DJ = \epsilon' JD$  by a local calculation:

$$\begin{aligned} (JD - \epsilon' DJ)(s \otimes \psi) &= J(\nabla_\mu^\mathfrak{B} s \otimes (-i\gamma^\mu \psi) + s \otimes D_M \psi) - \epsilon' D_\mathfrak{B}(s^* \otimes J_M \psi) \\ &= (\nabla_\mu^\mathfrak{B} s)^* \otimes i J_M \gamma^\mu \psi + s^* \otimes J_M D_M \psi \\ &\quad - \epsilon' \nabla_\mu^\mathfrak{B} s^* \otimes (-i\gamma^\mu J_M \psi) - \epsilon' s^* \otimes D_M J_M \psi \\ &= i((\nabla_\mu^\mathfrak{B} s)^* - \nabla_\mu^\mathfrak{B} s^*) \otimes J_M \gamma^\mu \psi = 0, \end{aligned}$$

since  $J_M \gamma^\mu = -\epsilon' \gamma^\mu J_M$ , and the last step follows from the definition of a  $*$ -algebra connection, i.e.  $(\nabla s)^* = \nabla s^*$  for all  $s \in \Gamma^\infty(\mathfrak{B})$ .

The commutant property follows easily:

$$\begin{aligned} [a, b^0](s \otimes \psi) &= a J b^* J^{-1}(s \otimes \psi) - J b^* J^{-1} a(s \otimes \psi) \\ &= a J(b^* s^* \otimes J_M^* \psi) - J b^*(s^* a^* \otimes J_M^* \psi) \\ &= asb \otimes \psi - asb \otimes \psi = 0, \end{aligned}$$

where  $a, b \in \Gamma^\infty(\mathfrak{B})$  and  $s \otimes \psi \in \Gamma^\infty(\mathfrak{B}) \otimes_{C^\infty(M)} \Gamma^\infty(S)$ . Since  $[a, b^0] = 0$  on  $\Gamma^\infty(\mathfrak{B}) \otimes_{C^\infty(M)} \Gamma^\infty(S) \cong \Gamma^\infty(\mathfrak{B} \otimes S)$ , it is zero on the entire Hilbert space  $L^2(\mathfrak{B} \otimes S)$ . It remains to check the order one condition for the Dirac operator. First note that

$$[[D, a], b^0](s \otimes \psi) = -i\gamma^\mu([[\nabla_\mu, a], b^0](s \otimes \psi)); \quad (a, b, s \in \Gamma^\infty(\mathfrak{B})).$$

This is zero because  $[[\nabla, a], b^0](s \otimes \psi)$  is zero:

$$\begin{aligned} ([\nabla_\mu, a]sb) \otimes \psi - J b^* J^{-1}([\nabla_\mu, a]s \otimes \psi) \\ &= \nabla_\mu(asb) \otimes \psi - a\nabla_\mu(sb) \otimes \psi - \nabla_\mu(as)b \otimes \psi + a(\nabla_\mu s)b \otimes \psi \\ &= ((\nabla_\mu a)sb + a(\nabla_\mu s)b + as(\nabla_\mu b) - a(\nabla_\mu s)b \\ &\quad - as(\nabla_\mu b) - (\nabla_\mu a)sb - a(\nabla_\mu s)b + a(\nabla_\mu s)b) \otimes \psi, \\ &= 0 \end{aligned}$$

using the defining property for  $\nabla^\mathfrak{B}$  to be a  $*$ -algebra connection. Thus,  $J$  fulfills all of the necessary conditions for a real structure on the spectral triple

$(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_{\mathfrak{B}})$ . The conditions on  $\gamma$  to be a grading operator for this spectral triple are easily checked too.  $\square$

## 12.2 Yang–Mills Theory as a Noncommutative Manifold

The real spectral triple  $(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_{\mathfrak{B}}; J, \gamma)$  that we obtained in Theorem 12.6 will turn out to be the correct triple to describe a topologically non-trivial  $PU(N)$ -gauge theory on the spin manifold  $M$  if the fibers of  $\mathfrak{B}$  are taken to be isomorphic to the  $*$ -algebra  $M_N(\mathbb{C})$ . Moreover, this triple not only describes a non-trivial  $PU(N)$ -gauge theory: every  $PU(N)$ -gauge theory on  $M$  is described by such a triple. In this section we prove these claims by first showing how a principal  $PU(N)$ -bundle can be constructed from this spectral triple. As in the topologically trivial case (cf. Remark 10.13) the spectral action applied to this triple will give the Einstein–Yang–Mills action, but now the gauge potential can be interpreted as a connection one-form on the  $PU(N)$ -bundle  $P$ . In fact, the original algebra bundle  $\mathfrak{B}$  will turn out to be an associated bundle of the principal bundle  $P$ . From now on, then, the fibers of  $\mathfrak{B}$  are assumed to be  $M_N(\mathbb{C})$ .

### 12.2.1 From Algebra Bundles to Principal Bundles

In order to construct a principal  $PU(N)$ -bundle  $P$  out of  $\mathfrak{B}$ , first of all note that since all  $*$ -automorphisms of  $M_N(\mathbb{C})$  are obtained by conjugation with a unitary element  $u \in M_N(\mathbb{C})$  (see Example 7.3), the transition functions of the bundle  $\Gamma^\infty(\mathfrak{B})$  take their values in

$$\text{Ad } U(N) \cong U(N)/Z(U(N)) \cong PU(N).$$

Thus the bundle  $\mathfrak{B}$  provides us with an open covering  $\{U_i\}$  of  $M$  as well as transition functions  $\{g_{ij}\}$  with values in  $PU(N)$ . Using the reconstruction theorem for principal bundles, we can then construct a principal  $PU(N)$ -bundle. By construction, the bundle  $\mathfrak{B}$  is an associated bundle to  $P$ .

Furthermore, for the real spectral triple

$$(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_{\mathfrak{B}}; J, \gamma)$$

of Theorem 12.6, the hermitian connection  $\nabla^{\mathfrak{B}}$  on the bundle  $\mathfrak{B}$  can locally be written as  $\nabla^{\mathfrak{B}} = d + \omega^{\mathfrak{B}}$ , where  $\omega^{\mathfrak{B}}$  is a  $su(N)$ -valued one-form, (cf. Remark 12.5). Moreover, the transformation rule for  $\omega^{\mathfrak{B}}$  is  $\omega_i^{\mathfrak{B}} = g_{ij}^{-1} dg_{ij} + g_{ij}^{-1} \omega_j^{\mathfrak{B}} g_{ij}$ , with  $g_{ij}$  the  $PU(N)$ -valued transition function of  $\mathfrak{B}$ . Comparing this expression with the usual transformation property of a connection one-form, one concludes that the hermitian  $*$ -algebra connection  $\nabla^{\mathfrak{B}}$  on  $\mathfrak{B}$  induces a connection one-form on the principal bundle  $P$  constructed in the previous paragraph.

Conversely, given a  $PU(N)$ -gauge theory  $(P, \omega^P)$  on some compact Riemannian spin manifold, we can construct the locally trivial hermitian  $*$ -algebra bundle  $\mathfrak{B} := P \times_{PU(N)} M_N(\mathbb{C})$ , where  $PU(N)$  acts on  $M_N(\mathbb{C})$  in the usual way. Moreover, the connection  $\omega^P$  on  $P$  induces a hermitian  $*$ -algebra connection on  $\mathfrak{B}$ . Following the steps described in the previous paragraph, it is not difficult to see that the principal bundle and connection obtained from the ensuing spectral triple,

$$(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), -i\gamma^\mu (\nabla_\mu^\mathfrak{B} \otimes 1 + 1 \otimes \nabla_\mu^S); J, \gamma),$$

coincide with  $(P, \omega^P)$ .

**Proposition 12.7** *Let  $(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_\mathfrak{B}; J, \gamma)$  be as before with  $M$  simply connected and  $\mathfrak{B}$  a locally trivial  $*$ -algebra bundle with fiber  $M_N(\mathbb{C})$  and a faithful smoothly-varying tracial state. Then:*

- (1) *there exists a principal  $PU(N)$ -bundle  $P$  such that  $\mathfrak{B}$  is an associated bundle of  $P$ , as well as a connection one-form  $\omega^P$  on  $P$  corresponding to  $\nabla^\mathfrak{B}$ ;*
- (2) *the gauge group  $\mathfrak{G}(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S); J)$  of this spectral triple (as in Definition 7.4) is isomorphic to the space of smooth sections of the associated group bundle  $\text{Ad } P := P \times_{PU(N)} PU(N)$ .*

Every  $PU(N)$ -gauge theory  $(P, \omega^P)$  on  $M$  is determined by such a spectral triple.

**Proof** The only statement left to prove is (2). If  $\mathfrak{B} = P \times_{PU(N)} M_N(\mathbb{C})$ , then  $\mathcal{U}(\Gamma^\infty(\mathfrak{B})) = \Gamma^\infty(P \times_{PU(N)} U(N))$ . As a consequence,

$$\begin{aligned} \mathfrak{G}(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S); J) &\simeq \{uJu^{-1} : u \in \Gamma^\infty(P \times_{PU(N)} U(N))\} \\ &\simeq \Gamma^\infty(P \times_{PU(N)} PU(N)), \end{aligned}$$

where we argue as in the proof of Proposition 10.2 (see also Note 4 on Chap. 10).  $\square$

### 12.2.2 Inner Fluctuations and Spectral Action

In this section, we calculate the spectral action for the real spectral triple of Theorem 12.6 in the case that  $\dim M = 4$ . We show that the spectral action applied to the spectral triple  $(\Gamma^\infty(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_\mathfrak{B}; J, \gamma)$  produces the Einstein–Yang–Mills action for a connection one-form on the  $PU(N)$ -bundle  $P$ . If  $\mathfrak{B}$  is a trivial algebra bundle, this reduces to Example 10.13. In fact, most of these local computations can be adopted in this case as well, since locally the bundle  $\mathfrak{B}$  is trivial. Nevertheless, for completeness we include the computations in the case at hand.

First of all, in Remark 12.5 we noticed that locally, i.e. on some local trivialization  $U$ , the connection  $\nabla^\mathfrak{B}$  is expressed as  $d + \omega^\mathfrak{B}$ , where  $\omega^\mathfrak{B}$  is an  $su(N)$ -valued one-form that acts in the adjoint representation on  $\Gamma^\infty(\mathfrak{B})$ . Therefore,  $\omega^\mathfrak{B}$  already induces

a connection one-form on  $P$ . To get the full gauge potential we need to take the fluctuations of the Dirac operator into account as well.

Recall from Sect. 7.2 that inner fluctuations of the Dirac operator are given by a perturbation term of the form

$$\omega = \sum_j a_j [D, b_j]; \quad (a_j, b_j \in \Gamma(\mathfrak{B})), \quad (12.2.1)$$

with the additional condition that  $\sum_j a_j [D, b_j]$  is a self-adjoint operator. Explicitly, we have

$$\omega = \sum_j -i\gamma^\mu \circ (a_j [\nabla_\mu, b_j] \otimes 1).$$

Locally, on some trivializing neighborhood  $U$ , the expression in (12.2.1) can be written as

$$\omega = \gamma^\mu A_\mu,$$

where  $A_\mu$  are the components of the one-form  $\sum_j a_j [\nabla, b_j]$  with values in  $\Gamma^\infty(\mathfrak{B})$ . Since  $\omega$  is self-adjoint, the one-form  $A_\mu$  can be considered a real one-form taking values in the hermitian elements of  $\Gamma^\infty(\mathfrak{B})$ .

Similarly, the expression  $\omega + J\omega J^{-1}$  is locally written as

$$\gamma^\mu A_\mu - \gamma^\mu J A_\mu J^{-1},$$

since in 4 dimensions  $\gamma^\mu$  anti-commutes with  $J$ . Writing out the second term gives:

$$(\gamma^\mu J A_\mu J^{-1})(s \otimes \psi) = s A_\mu \otimes \gamma^\mu \psi; \quad (s \otimes \psi \in \Gamma^\infty(\mathfrak{B} \otimes S)),$$

so that on this local patch,  $\omega + J\omega J^{-1}$  can be written as

$$\gamma^\mu \text{ad } A_\mu.$$

Consequently,  $\omega + J\omega J^{-1}$  eliminates the  $iu(1)$ -part of  $\omega$ , so that  $\omega$  effectively satisfies the *unimodularity condition*

$$\text{Tr } \omega = 0.$$

Thus,  $i \text{ad } A_\mu$  is a one-form on  $M$  with values in  $\Gamma^\infty(\text{ad } P)$  where  $\text{ad } P = P \times_{PU(N)} su(N)$ .

The expression for  $D + \omega + J\omega J^{-1}$  on a local chart  $U$  is then given by

$$D_\omega = -i\gamma^\mu (\nabla_\mu^\mathfrak{B} \otimes 1 + 1 \otimes \nabla_\mu^S + i \text{ad } A_\mu \otimes 1),$$

where the connection  $\nabla^{\mathfrak{B}}$  can be expressed on  $U$  as  $d + \omega^{\mathfrak{B}}$  for some unique  $su(N)$ -valued one-form  $\omega^{\mathfrak{B}}$  on  $U$ . Thus, on  $U$  the fluctuated Dirac operator can be rewritten as

$$D_{\omega} = -i\gamma^{\mu}(1 \otimes \nabla_{\mu}^S + (\partial_{\mu} + \omega_{\mu}^{\mathfrak{B}} + i \operatorname{ad} A_{\mu}) \otimes 1).$$

We interpret  $(\omega_{\mu}^{\mathfrak{B}} + i \operatorname{ad} A_{\mu})$  as the full gauge potential on  $U$ , acting in the adjoint representation on the spinors. The natural action of an element  $g$  in the group  $\mathfrak{G}(\Gamma^{\infty}(\mathfrak{B}), L^2(\mathfrak{B} \otimes S); J) \simeq \Gamma^{\infty}(\operatorname{Ad} P)$  by conjugation on  $D_{\omega}$  then induces the familiar gauge transformation:

$$\omega_{\mu}^{\mathfrak{B}} + i \operatorname{ad} A_{\mu} \mapsto (g^{-1}\omega_{\mu}^{\mathfrak{B}} g + g^{-1}(dg)) + g^{-1}(i \operatorname{ad} A_{\mu})g,$$

where the first two terms on the right-hand side are the transformation of  $\omega^{\mathfrak{B}}$  under a change of local trivialization, and the last term is the transformation of  $i \operatorname{ad} A_{\mu}$ . Therefore, since  $\mathfrak{B}$  is an associated bundle of  $P$ , it follows that  $\omega_{\mu}^{\mathfrak{B}} + i \operatorname{ad} A_{\mu}$  induces a  $su(N)$ -valued connection one-form on the principal  $PU(N)$ -bundle  $P$  that acts on  $\Gamma^{\infty}(\mathfrak{B})$  in the adjoint representation.

Let us summarize what we have obtained so far.

**Proposition 12.8** *Let  $(\Gamma^{\infty}(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_{\mathfrak{B}}; J, \gamma)$  and let  $P$  be as before, so that  $P \times_{PU(N)} M_N(\mathbb{C}) \simeq \mathfrak{B}$ . Then, the inner fluctuations of  $D_{\mathfrak{B}}$  are parametrized by sections of  $\Gamma^{\infty}(T^*M \otimes \operatorname{ad} P)$  where  $\operatorname{ad} P = P \times_{PU(N)} su(N)$ . Moreover, the action of  $\mathfrak{G}(\Gamma^{\infty}(\mathfrak{B}), L^2(\mathfrak{B} \otimes S); J)$  on the inner fluctuations of  $D_{\mathfrak{B}}$  by conjugation coincides with the adjoint action of  $\Gamma^{\infty}(\operatorname{Ad} P)$  on  $\Gamma^{\infty}(\operatorname{ad} P)$ .*

Let us now proceed to compute the spectral action for these inner fluctuations. We apply the results of Sect. 10.3, using the following result.

**Lemma 12.9** *For the spectral triple  $(\Gamma^{\infty}(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_{\mathfrak{B}}; J, \gamma)$ , the square of the fluctuated Dirac operator is a generalized Laplacian of the form  $\Delta^E - F$ , with  $E = \mathfrak{B} \otimes S$  (notation as in Theorem 10.7), and we have the following local expressions for the corresponding curvature  $\Omega_{\mu\nu}^E$  and the bundle endomorphism  $F$ :*

$$\begin{aligned} F &= -\frac{1}{4}s \otimes \mathbb{I}_{N^2} + \frac{1}{2}i\gamma^{\mu}\gamma^{\nu} \otimes F_{\mu\nu}; \\ \Omega_{\mu\nu}^E &= \Omega_{\mu\nu}^S \otimes \mathbb{I}_{N^2} + i\mathbb{I}_4 \otimes F_{\mu\nu}, \end{aligned}$$

where  $F_{\mu\nu}$  is the curvature of the connection  $\nabla_{\mu}^{\mathfrak{B}} + i \operatorname{ad} A_{\mu}$ .

As before, this result allows us to compute the bosonic spectral action for the fluctuated Dirac operator  $D_{\omega}$ , essentially reducing the computation in terms of a local trivialization to the trivial case (cf. Example 10.13), with the following result.

**Theorem 12.10** *For the spectral triple  $(\Gamma^{\infty}(\mathfrak{B}), L^2(\mathfrak{B} \otimes S), D_{\mathfrak{B}}; J, \gamma)$ , the spectral action yields the Yang–Mills action for  $\nabla^{\mathfrak{B}} + i \operatorname{ad} A_{\mu}$  minimally coupled to gravity:*

$$\mathrm{Tr}(f(D_\omega/\Lambda)) \sim \frac{f(0)}{24\pi^2} \int_M \mathrm{Tr} F_{\mu\nu} F^{\mu\nu} \sqrt{g} dx + N^2 \int_M \mathcal{L}_M(g_{\mu\nu}) \sqrt{g} dx,$$

asymptotically as  $\Lambda \rightarrow \infty$  and up to terms  $\propto \Lambda^{-2}$ . The Lagrangian  $\mathcal{L}_M(g^{\mu\nu})$  is given by (10.4.8).

### 12.2.3 Topological Spectral Action

A natural invariant in this topologically non-trivial context is the topological spectral action, given in Eq. (9.1.2). With Proposition 9.5 we find that, in general,

$$S_{\mathrm{top}}[\omega] = f(0) \mathrm{index} D_\omega.$$

Hence, in the setting of Theorem 12.10, using the Atiyah–Singer index theorem (cf. Note 18 on Chap. 6), we find an extra contribution of the form

$$S_{\mathrm{top}}[\omega] = \frac{f(0)}{(2\pi i)^{n/2}} \int_M \hat{A}(M) \mathrm{ch}(\mathfrak{B}),$$

in terms of the  $\hat{A}$ -form of  $M$  and the Chern character of the algebra bundle  $\mathfrak{B}$ .

## Notes

- For an exposition of Yang–Mills theory in terms of principal bundle and connections, we refer to [1, Sect. 2,3] and [2].
- This chapter extends the noncommutative description of Yang–Mills gauge theory of [3, 4] to the topologically non-trivial case; it is based on [5]. For a more general treatment of topologically non-trivial almost-commutative geometries we refer to [6–8].

### Section 12.1 Spectral triple obtained from an algebra bundle

- Our approach to locally trivial  $*$ -algebra bundles gains in substance with the Serre–Swan Theorem, establishing a duality between vector bundles over a topological space  $X$  and finite projective modules over  $C(X)$  [9, 10]. A smooth version was obtained in [11] (see also [12, Proposition 4.2.1] or [13, Sect. 2.3]). The fiberwise inner product gives rise to the hermitian structure found in Proposition 7.14. A version of the Serre–Swan Theorem for  $*$ -algebra bundles has been obtained in [5].

### Section 12.2 Yang–Mills theory as a noncommutative manifold

- A special case of Proposition 12.7 occurs when  $\mathfrak{B}$  is an endomorphism bundle. It follows from a result by Dixmier and Douady in [14] (cf. [15]) that a bundle  $\mathfrak{B}$  with continuously varying

trace is an endomorphism bundle if and only if the Dixmier–Douady class  $\delta(\Gamma(\mathfrak{B})) \in H^3(\mathbb{Z})$  of the  $C^*$ -algebra of continuous sections  $\Gamma(\mathfrak{B})$  of this bundle is equal to zero. Because the Dixmier–Douady class of the bundle  $\mathfrak{B}$  vanishes one can lift the  $PU(N)$ -valued transition functions  $g_{ij}$  to  $U(N)$ -valued functions  $\mu_{ij}$  such that  $g_{ij} = \text{Ad } \mu_{ij}$ , and  $\mu_{ij}\mu_{jk} = \mu_{ik}$  (see for instance [15], Theorem 4.85). One may therefore construct a principal  $U(N)$ -bundle instead of a  $PU(N)$ -bundle, to which  $\mathfrak{B}$  is associated if and only if  $\mathfrak{B}$  is an endomorphism bundle.

## References

- Atiyah, M.F., Bott, R.: The Yang-Mills equations over Riemann surfaces. *Philos. Trans. Roy. Soc. London Ser. A* **308**, 523–615 (1983)
- Bleecker, D.: Gauge Theory and Variational Principles. Addison-Wesley, Reading (1981)
- Chamseddine, A.H., Connes, A.: Universal formula for noncommutative geometry actions: unifications of gravity and the standard model. *Phys. Rev. Lett.* **77**, 4868–4871 (1996)
- Chamseddine, A.H., Connes, A.: The spectral action principle. *Commun. Math. Phys.* **186**, 731–750 (1997)
- Boeijink, J., van Suijlekom, W.D.: The noncommutative geometry of Yang-Mills fields. *J. Geom. Phys.* **61**, 1122–1134 (2011)
- Ćaćić, B.: A reconstruction theorem for almost-commutative spectral triples. *Lett. Math. Phys.* **100**, 181–202 (2012)
- Ćaćić, B.: Real structures on almost-commutative spectral triples. *Lett. Math. Phys.* **103**, 793–816 (2013)
- Boeijink, J., van den Dungen, K.: On globally non-trivial almost-commutative manifolds. *J. Math. Phys.* **55**, 103508 (2014)
- Serre, J.-P.: Modules projectifs et espaces fibrés à fibre vectorielle. In: Dubreil, S.P., Dubreil-Jacotin, M.-L., Pisot, C. (eds.) 1957/58, Fasc. 2, Exposé 23, page 18. Secrétariat mathématique, Paris (1958)
- Swan, R.G.: Vector bundles and projective modules. *Trans. Am. Math. Soc.* **105**, 264–277 (1962)
- Connes, A.: Non-commutative geometry and physics. In: Gravitation et Quantifications, pp. 805–950. Les Houches (1992). North-Holland, Amsterdam, 1995
- Landi, G.: An Introduction to Noncommutative Spaces and Their Geometry. Springer (1997)
- Gracia-Bondía, J.M., Várilly, J.C., Figueroa, H.: Elements of Noncommutative Geometry. Birkhäuser, Boston (2001)
- Dixmier, J., Douady, A.: Champs continus d’espaces hilbertiens et de  $C^*$ -algèbres. *Bull. Soc. Math. France* **91**, 227–284 (1963)
- Raeburn, I., Williams, D.P.: Morita equivalence and continuous-trace  $C^*$ -algebras. Mathematical Surveys and Monographs, vol. 60. American Mathematical Society, Providence, RI (1998)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book’s Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book’s Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 13

## The Noncommutative Geometry of the Standard Model



One of the major applications of noncommutative geometry to physics has been the derivation of the Standard Model of particle physics from a suitable almost-commutative manifold. In this Chapter we present this derivation, using the results of Chap. 10.

### 13.1 The Finite Space

Our starting point is the classification of irreducible finite geometries of KO-dimension 6 from Sect. 3.4, based on the matrix algebra  $M_N(\mathbb{C}) \oplus M_N(\mathbb{C})$  for  $N \geq 1$ . We have already seen in Chap. 11 that  $N = 1$  is the finite geometry corresponding to electrodynamics. We now proceed and aim for the full Standard Model of particle physics. Let us make the following two additional requirements on the irreducible finite geometry  $(A, H_F, D_F; J_F, \gamma_F)$ :

- (1) The finite-dimensional Hilbert space  $H_F$  carries a symplectic structure  $I^2 = -1$ ;
- (2) the grading  $\gamma_F$  induces a non-trivial grading on  $A$ , by mapping


$$a \mapsto \gamma_F a \gamma_F,$$

and selects an even subalgebra  $A^{\text{ev}} \subset A$  consisting of elements that commute with  $\gamma_F$ .

We have already seen in Sect. 3.4 that the first demand sets  $A = M_k(\mathbb{H}) \oplus M_{2k}(\mathbb{C})$ , represented on the Hilbert space  $\mathbb{C}^{2(2k)^2}$ . The second requirement sets  $k \geq 2$ ; we will take the simplest  $k = 2$  so that  $H_F = \mathbb{C}^{32}$ . Indeed, this allows for a  $\gamma_F$  such that

$$A^{\text{ev}} = \mathbb{H}_R \oplus \mathbb{H}_L \oplus M_4(\mathbb{C}),$$

where  $\mathbb{H}_R$  and  $\mathbb{H}_L$  are two copies (referred to as *right* and *left*) of the quaternions; they are the diagonal of  $M_2(\mathbb{H}) \subset A$ . The Hilbert space can then be decomposed



**Fig. 13.1** The Krajewski diagram for the finite real spectral triple  $(A^{\text{ev}} = \mathbb{H}_R \oplus \mathbb{H}_L \oplus M_4(\mathbb{C}), H_F, D_F; J_F, \gamma_F)$ . The dashed line corresponds to an ‘off-diagonal’ component of the Dirac operator, thus failing on the first-order condition. The labels + and – represent the value of the grading  $\gamma_F$  on the corresponding summands of  $H_F$

according to the defining representations of  $A^{\text{ev}}$ ,

$$H_F = (\mathbb{C}_R^2 \oplus \mathbb{C}_L^2) \otimes \mathbb{C}^{4^o} \oplus \mathbb{C}^4 \otimes (\mathbb{C}_R^{2^o} \oplus \mathbb{C}_L^{2^o}). \quad (13.1.1)$$

According to this direct sum decomposition, we write

$$D_F = \begin{pmatrix} S & T^* \\ T & \bar{S} \end{pmatrix} \quad (13.1.2)$$

where

$$\begin{aligned} S : (\mathbb{C}_R^2 \oplus \mathbb{C}_L^2) \otimes \mathbb{C}^{4^o} &\rightarrow (\mathbb{C}_R^2 \oplus \mathbb{C}_L^2) \otimes \mathbb{C}^{4^o}, \\ T : (\mathbb{C}_R^2 \oplus \mathbb{C}_L^2) \otimes \mathbb{C}^{4^o} &\rightarrow \mathbb{C}^4 \otimes (\mathbb{C}_R^{2^o} \oplus \mathbb{C}_L^{2^o}). \end{aligned}$$

This gives rise to the Krajewski diagram of Fig. 13.1. We now make an additional assumption,


- (3) The off-diagonal components  $T$  and  $T^*$  of the Dirac operator in (13.1.2) are non-zero.

In Fig. 13.1 such an off-diagonal component corresponds to the dashed line. As this line runs neither vertically, horizontally, or between the same vertex, it follows from Lemma 3.10 that the corresponding component of  $D_F$  breaks the first-order condition.

**Proposition 13.1** *Up to \*-automorphisms of  $A^{\text{ev}}$ , there is a unique \*-subalgebra  $A_F \subset A^{\text{ev}}$  of maximal dimension that allows  $T \neq 0$  in (13.1.2). It is given by*

$$A_F = \left\{ \left( q_\lambda, q, \begin{pmatrix} q & 0 \\ 0 & m \end{pmatrix} \right) : \lambda \in \mathbb{C}, q \in \mathbb{H}_L, m \in M_3(\mathbb{C}) \right\} \subset \mathbb{H}_R \oplus \mathbb{H}_L \oplus M_4(\mathbb{C}),$$

**Fig. 13.2** The Krajewski diagram of the space  $F_{SM}$  describing the Standard Model



where  $\lambda \mapsto q_\lambda$  is the embedding of  $\mathbb{C} \hookrightarrow \mathbb{H}$ , with

$$q_\lambda = \begin{pmatrix} \lambda & 0 \\ 0 & \bar{\lambda} \end{pmatrix}.$$

Consequently,  $A_F \simeq \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ .

**Proof** We give a diagrammatic proof. From Fig. 13.1, we see that in order to fulfill the first-order condition, we should bring the dashed line to run horizontally or vertically, or to begin and start at the same node on the diagonal. We do so by considering the Krajewski diagrams for subalgebras  $A_F \subset A^{\text{ev}}$  which are induced by Fig. 13.1. If  $T$  is of rank 1, the only possibility is to bring the dashed line to the diagonal. In other words, the subalgebra we are looking for should have a component that is embedded diagonally in  $\mathbb{H}_R$  and  $M_4(\mathbb{C})$ . Such a component can only be  $\mathbb{C}$ , and the resulting subalgebra is embedded as

$$\begin{aligned} \mathbb{C} \oplus M_3(\mathbb{C}) &\rightarrow \mathbb{H}_R \oplus M_4(\mathbb{C}); \\ (\lambda, m) &\mapsto \left( \begin{pmatrix} \lambda & 0 \\ 0 & \bar{\lambda} \end{pmatrix}, \begin{pmatrix} \lambda & 0 \\ 0 & m \end{pmatrix} \right). \end{aligned}$$

This breaks the Krajewski diagram to the diagram of Fig. 13.2, where the dashed line now connects the two vertices labeled by  $(\mathbf{1}, \mathbf{1}^\circ)$ . The other edges of Fig. 13.1 are now torn apart to the resulting edges in Fig. 13.2.

If  $T$  has rank greater than 1, then a similar argument shows that one obtains a subalgebra of smaller dimension than  $A_F$ .  $\square$

In order to connect to the physics of the Standard Model, let us introduce an orthonormal basis for  $H_F$  that can be recognized as the fermionic particle content of the Standard Model, and subsequently write the representation of  $A_F$  in terms of this basis. Starting with the Krajewski diagram of Fig. 13.2, we let the first three nodes in the top row be represented by basis vectors  $\{\nu_R, e_R, (\nu_L, e_L)\}$  of the so-called **lepton space**  $H_l$ , while the three nodes in the bottom row represent the basis vectors  $\{u_R, d_R, (u_L, d_L)\}$  of the **quark space**  $H_q$ . Their reflections with respect to the diagonal represent are the **anti-lepton space**  $H_{\bar{l}}$  and the **anti-quark space**  $H_{\bar{q}}$ ,

spanned by  $\{\overline{\nu_R}, \overline{e_R}, (\overline{\nu_L}, \overline{e_L})\}$  and  $\{\overline{u_R}, \overline{d_R}, (\overline{u_L}, \overline{d_L})\}$ , respectively. The three colors of the quarks are given by a tensor factor  $\mathbb{C}^3$  and when we take into account *three generations* of fermions and anti-fermions by tripling the above finite-dimensional Hilbert space we obtain

$$H_F := (H_l \oplus H_{\bar{l}} \oplus H_q \oplus H_{\bar{q}})^{\oplus 3}.$$

Note that  $H_l = \mathbb{C}^4$ ,  $H_q = \mathbb{C}^4 \otimes \mathbb{C}^3$ ,  $H_{\bar{l}} = \mathbb{C}^4$ , and  $H_{\bar{q}} = \mathbb{C}^4 \otimes \mathbb{C}^3$ . An element  $a = (\lambda, q, m) \in A_F$  acts on the space of leptons  $H_l$  as  $q_\lambda \oplus q$ , and acts on the space of quarks  $H_q$  as  $(q_\lambda \oplus q) \otimes \mathbb{I}_3$ . That is,

$$\begin{aligned} a = (\lambda, q, m) &\xrightarrow{H_l} \begin{pmatrix} \lambda & 0 & 0 & 0 \\ 0 & \bar{\lambda} & 0 & 0 \\ 0 & 0 & \alpha & \beta \\ 0 & 0 & -\bar{\beta} & \bar{\alpha} \end{pmatrix}, \\ a = (\lambda, q, m) &\xrightarrow{H_q} \begin{pmatrix} \lambda & 0 & 0 & 0 \\ 0 & \bar{\lambda} & 0 & 0 \\ 0 & 0 & \alpha & \beta \\ 0 & 0 & -\bar{\beta} & \bar{\alpha} \end{pmatrix} \otimes \mathbb{I}_3. \end{aligned}$$

For the action of  $a$  on an anti-lepton  $\bar{l} \in H_{\bar{l}}$  we have  $a\bar{l} = \lambda\mathbb{I}_4\bar{l}$ , and on an anti-quark  $\bar{q} \in H_{\bar{q}}$  we have  $a\bar{q} = (\mathbb{I}_4 \otimes m)\bar{q}$ .

The  $\mathbb{Z}_2$ -grading  $\gamma_F$  is such that left-handed particles have eigenvalue +1 and right-handed particles have eigenvalue -1. The anti-linear operator  $J_F$  interchanges particles with their anti-particles, so  $J_F f = \bar{f}$  and  $J_F \bar{f} = f$ , with  $f$  a lepton or quark.

Finally, we write the Dirac operator of (13.1.2) in terms of the decomposition of  $H_F$  in particle  $(H_l^{\oplus 3} \oplus H_q^{\oplus 3})$  and anti-particles  $(H_{\bar{l}}^{\oplus 3} \oplus H_{\bar{q}}^{\oplus 3})$ . The operator  $S$  will be chosen to be

$$\begin{aligned} S_l := S|_{H_l^{\oplus 3}} &= \begin{pmatrix} 0 & 0 & Y_\nu^* & 0 \\ 0 & 0 & 0 & Y_e^* \\ Y_\nu & 0 & 0 & 0 \\ 0 & Y_e & 0 & 0 \end{pmatrix}, \\ S_q \otimes \mathbb{I}_3 := S|_{H_q^{\oplus 3}} &= \begin{pmatrix} 0 & 0 & Y_u^* & 0 \\ 0 & 0 & 0 & Y_d^* \\ Y_u & 0 & 0 & 0 \\ 0 & Y_d & 0 & 0 \end{pmatrix} \otimes \mathbb{I}_3, \end{aligned}$$

where  $Y_\nu$ ,  $Y_e$ ,  $Y_u$  and  $Y_d$  are  $3 \times 3$  **Yukawa mass matrices** acting on the three generations, and  $\mathbb{I}_3$  acting on the three colors of the quarks. The symmetric operator  $T$  only acts on the right-handed (anti)neutrinos, so it is given by  $T\nu_R = Y_R\overline{\nu_R}$ , for a certain  $3 \times 3$  symmetric **Majorana mass matrix**  $Y_R$ , and  $Tf = 0$  for all other

fermions  $f \neq \nu_R$ . Note that  $\nu_R$  here stands for a vector with 3 components for the number of generations.

Let us summarize what we have obtained so far.

**Proposition 13.2** *The data*

$$F_{SM} := (A_F, H_F, D_F; J_F, \gamma_F)$$

as given above define a finite real even spectral triple of KO-dimension 6.

## 13.2 The Gauge Theory

### 13.2.1 The Gauge Group

We shall now describe the gauge theory corresponding to the almost-commutative manifold  $M \times F_{SM}$ . In order to determine the gauge group  $\mathfrak{G}(F_{SM})$  of Definition 7.4, let us start by examining the subalgebra  $(A_F)_{J_F}$  of the algebra  $A_F$  of Proposition 13.1, as defined in Sect. 8.1. For an element  $a = (\lambda, q, m) \in \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ , the relation  $aJ_F = J_F a^*$  now yields  $\lambda = \bar{\lambda} = \alpha = \bar{\alpha}$  and  $\beta = 0$ , as well as  $m = \lambda \mathbb{I}_3$ . So,  $a \in (A_F)_{J_F}$  if and only if  $a = (x, x, x)$  for  $x \in \mathbb{R}$ . Hence we find

$$(A_F)_{J_F} \simeq \mathbb{R}.$$

Next, let us consider the Lie algebra  $\mathfrak{h}(F) = \mathfrak{u}((A_F)_{J_F})$  of (10.1.1b). Since  $\mathfrak{u}(A_F)$  consists of the anti-hermitian elements of  $A_F$ , we obtain that the  $\mathfrak{h}(F) = \mathfrak{u}((A_F)_{J_F})$  is given by the trivial subalgebra  $\{0\}$ .

**Proposition 13.3** *The local gauge group  $\mathfrak{G}(F_{SM})$  of the finite space  $F_{SM}$  is given by*

$$\mathfrak{G}(F_{SM}) \simeq (U(1) \times SU(2) \times U(3)) / \{1, -1\},$$

where  $\{1, -1\}$  is the diagonal normal subgroup in  $U(1) \times SU(2) \times U(3)$ .

**Proof** The unitary elements of the algebra form the group  $\mathcal{U}(A_F) \simeq U(1) \times \mathcal{U}(\mathbb{H}) \times U(3)$ . Now, a quaternion  $q = q_0 \mathbb{I} + i q_1 \sigma_1 + i q_2 \sigma_2 + i q_3 \sigma_3$  is unitary if and only if  $|q|^2 = q_0^2 + q_1^2 + q_2^2 + q_3^2 = 1$ . Using the embedding of  $\mathbb{H}$  in  $M_2(\mathbb{C})$ , we find  $|q|^2 = \det(q) = 1$ , and this yields the isomorphism  $\mathcal{U}(\mathbb{H}) \simeq SU(2)$ . Hence, the unitary group  $\mathcal{U}(A_F)$  is given by  $U(1) \times SU(2) \times U(3)$ . By Proposition 10.2, the gauge group is given by the quotient of the unitary group with the subgroup  $\mathfrak{H}(F) = \mathfrak{u}((A_F)_{J_F})$ , which is the diagonal normal subgroup

$$\{\pm(1, \mathbb{I}_2, \mathbb{I}_3)\} \subset U(1) \times SU(2) \times U(3).$$

□

The gauge group that we obtain here is not the gauge group of the Standard Model, because (even ignoring the quotient with the finite group  $\{1, -1\}$ ) we have a factor  $U(3)$  instead of  $SU(3)$ . As mentioned in Proposition 10.3, the unimodularity condition is only satisfied for complex algebras, but in our case, the algebra  $\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$  is only a real algebra. Therefore, the unimodularity condition is not automatically satisfied. Instead, we shall *require* that the unimodularity condition is satisfied, so for  $u = (\lambda, q, m) \in U(1) \times SU(2) \times U(3)$  we impose

$$\det|_{H_F}(u) = 1 \implies (\lambda \det m)^{12} = 1.$$

For  $u \in U(1) \times SU(2) \times U(3)$ , we denote the corresponding element in  $\mathfrak{G}(F_{SM})$  by  $U = uJuJ^{-1}$ . We shall then consider the subgroup

$$S\mathfrak{G}(F_{SM}) = \left\{ U = uJuJ^{-1} \in \mathfrak{G}(F_{SM}) \mid u = (\lambda, q, m), (\lambda \det m)^{12} = 1 \right\}.$$

The effect of the unimodularity condition is that the determinant of  $m \in U(3)$  is identified (modulo the multiplicative group  $\mu_{12}$  of 12'th roots of unity) with  $\bar{\lambda}$ . In other words, imposing the unimodularity condition provides us, modulo some finite abelian group, with the gauge group  $U(1) \times SU(2) \times SU(3)$ . This agrees with the Standard Model, as even the group  $U(1) \times SU(2) \times SU(3)$  is actually not the true gauge group of the Standard Model. Indeed, it contains a finite abelian subgroup (isomorphic to)  $\mu_6$  which acts trivially on all bosonic and fermionic particles in the Standard Model. The group  $\mu_6$  is embedded in  $U(1) \times SU(2) \times SU(3)$  by  $\lambda \mapsto (\lambda, \lambda^3, \lambda^2)$ . The true gauge group of the Standard Model is therefore given by

$$\mathfrak{G}_{SM} := U(1) \times SU(2) \times SU(3)/\mu_6.$$

**Proposition 13.4** *The unimodular gauge group  $S\mathfrak{G}(F_{SM})$  is isomorphic to*

$$S\mathfrak{G}(F_{SM}) \cong \mathfrak{G}_{SM} \rtimes \mu_{12}.$$

**Proof** Proposition 13.3 shows that  $S\mathfrak{G}(F_{SM}) \cong S\mathcal{U}(A_F)/\mu_2$ , so we determine  $S\mathcal{U}(A_F)$ . We do so in two steps:

$$S\mathcal{U}(A_F) \cong G \times SU(2) \times SU(3)/\mu_3, \tag{I}$$

where  $G = \{(\lambda, \mu) \in U(1) \times U(1) : (\lambda\mu^3)^{12} = 1\}$ , containing  $\mu_3$  as the subgroup  $\{e\} \times \mu_3$ , and

$$G \cong \mu_{12} \times U(1). \tag{II}$$

For (I), consider the map

$$(\lambda, \mu, q, m) \in G \times SU(2) \times SU(3) \mapsto (\lambda, q, \mu m) \in SU(A_F).$$

We claim that this map is surjective and has kernel  $\mu_3$ . If  $(\lambda, q, m) \in SU(A_F)$ , then there exists  $\mu \in U(1)$  such that  $\mu^3 = \det m \in U(1)$ . Since  $(\lambda\mu^3)^{12} = (\lambda \det m)^{12} = 1$ , the element  $(\lambda, \mu, q, m)$  lies in the pre-image of  $(\lambda, q, m)$ . The kernel of the above map consists of pairs  $(\lambda, \mu, q, m) \in G \times SU(2) \times SU(3)$  such that  $\lambda = 1$ ,  $q = 1$  and  $m = \mu^{-1}\mathbb{I}_3$ . Since  $m \in SU(3)$ , this  $\mu$  satisfies  $\mu^3 = 1$ . So we have established (I).

For (II) we show that the following sequence is split-exact:

$$1 \rightarrow U(1) \rightarrow G \rightarrow \mu_{12} \rightarrow 1,$$

where the group homomorphisms are given by  $\lambda \in U(1) \mapsto (\lambda^3, \lambda^{-1}) \in G$  and  $(\lambda, \mu) \in G \rightarrow \lambda\mu^3 \in \mu_a$ . Exactness can be easily checked, and the splitting map is given by  $\lambda \in \mu_{12} \rightarrow (\lambda, 1) \in G$ . In this abelian case, the corresponding action of  $\mu_{12}$  on  $U(1)$  is trivial so that the resulting semi-direct product is

$$G \simeq U(1) \rtimes \mu_{12} \simeq U(1) \times \mu_{12}.$$

□

A similar argument shows that the gauge algebra of Definition 7.4 is

$$\mathfrak{g}(F_{SM}) \simeq u(1) \oplus su(2) \oplus u(3),$$

and the restriction to traceless matrices gives the gauge algebra of the Standard Model:

$$sg(F_{SM}) \simeq u(1) \oplus su(2) \oplus su(3).$$

### 13.2.2 The Gauge and Scalar Fields

As we have seen in more generality in (10.2.7), the gauge field corresponding to  $F_{SM}$  takes values in  $\mathfrak{g}(F_{SM})$ . We here confirm this result and derive the precise form of the gauge field  $A_\mu$  of (10.2.1), and also of the scalar field  $\phi$  of (10.2.2).

Take two elements  $a = (\lambda, q, m)$  and  $b = (\lambda', q', m')$  of the algebra  $\mathcal{A} = C^\infty(\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}))$ . According to the representation of  $A_F$  on  $H_F$ , the inner fluctuations  $A_\mu = -ia\partial_\mu b$  decompose as

$$\Lambda_\mu := -i\lambda\partial_\mu\lambda'$$

on  $\nu_R$ ,

$$\Lambda'_\mu := -i\bar{\lambda}\partial_\mu\bar{\lambda}'$$

on  $e_R$ ,

$$Q_\mu := -iq\partial_\mu q'$$

on  $(\nu_l, e_L)$ , and

$$V'_\mu := -im\partial_\mu m'$$

acting on  $H_{\bar{q}}$ ; on all other components of  $H_F$  the gauge field  $A_\mu$  acts as zero. Imposing the hermiticity  $\Lambda_\mu = \Lambda_\mu^*$  implies  $\Lambda_\mu \in \mathbb{R}$ , and also automatically yields  $\Lambda'_\mu = -\Lambda_\mu$ . Furthermore,  $Q_\mu = Q_\mu^*$  implies that  $Q_\mu$  is a real-linear combination of the Pauli matrices, which span  $i\mathfrak{su}(2)$ . Finally, the condition that  $V'_\mu$  be hermitian yields  $V'_\mu \in i\mathfrak{u}(3)$ , so  $V'_\mu$  is a  $U(3)$  gauge field. As mentioned above, we need to impose the unimodularity condition to obtain an  $SU(3)$  gauge field. Hence, we require that the trace of the gauge field  $A_\mu$  over  $H_F$  vanishes, and we obtain

$$\text{Tr}|_{H_{\bar{l}}}(\Lambda_\mu \mathbb{I}_4) + \text{Tr}|_{H_{\bar{q}}}(\mathbb{I}_4 \otimes V'_\mu) = 0 \implies \text{Tr}(V'_\mu) = -\Lambda_\mu.$$

Therefore, we can define a traceless  $SU(3)$  gauge field  $V_\mu$  by  $\bar{V}_\mu := -V'_\mu - \frac{1}{3}\Lambda_\mu$ . The gauge field  $A_\mu$  is given by

$$\begin{aligned} A_\mu|_{H_l} &= \begin{pmatrix} \Lambda_\mu & 0 \\ 0 & -\Lambda_\mu \\ & Q_\mu \end{pmatrix}, & A_\mu|_{H_q} &= \begin{pmatrix} \Lambda_\mu & 0 \\ 0 & -\Lambda_\mu \\ & Q_\mu \end{pmatrix} \otimes \mathbb{I}_3, \\ A_\mu|_{H_{\bar{l}}} &= \Lambda_\mu \mathbb{I}_4, & A_\mu|_{H_{\bar{q}}} &= -\mathbb{I}_4 \otimes \left( \bar{V}_\mu + \frac{1}{3}\Lambda_\mu \right), \end{aligned}$$

for some  $U(1)$  gauge field  $\Lambda_\mu$ , an  $SU(2)$  gauge field  $Q_\mu$  and an  $SU(3)$  gauge field  $V_\mu$ . The action of the field  $B_\mu = A_\mu - J_F A_\mu J_F^{-1}$  on the fermions is then given by

$$\begin{aligned} B_\mu|_{H_l} &= \begin{pmatrix} 0 & 0 \\ 0 & -2\Lambda_\mu \\ & Q_\mu - \Lambda_\mu \mathbb{I}_2 \end{pmatrix}, \\ B_\mu|_{H_q} &= \begin{pmatrix} \frac{4}{3}\Lambda_\mu \mathbb{I}_3 + V_\mu & 0 \\ 0 & -\frac{2}{3}\Lambda_\mu \mathbb{I}_3 + V_\mu \\ & (Q_\mu + \frac{1}{3}\Lambda_\mu \mathbb{I}_2) \otimes \mathbb{I}_3 + \mathbb{I}_2 \otimes V_\mu \end{pmatrix}. \end{aligned} \tag{13.2.1}$$

Note that the coefficients in front of  $\Lambda_\mu$  in the above formulas are precisely the well-known hypercharges of the corresponding particles, as given by the following table:

Particle	$\nu_R$	$e_R$	$\nu_L$	$e_L$	$u_R$	$d_R$	$u_L$	$d_L$
Hypercharge	0	-2	-1	-1	$\frac{4}{3}$	$-\frac{2}{3}$	$\frac{1}{3}$	$\frac{1}{3}$

Next, let us turn to the scalar field  $\phi$ , which is given by

$$\phi|_{H_l} = \begin{pmatrix} 0 & Y^* \\ Y & 0 \end{pmatrix}, \quad \phi|_{H_q} = \begin{pmatrix} 0 & X^* \\ X & 0 \end{pmatrix} \otimes \mathbb{I}_3, \quad \phi|_{H_{\bar{l}}} = 0, \quad \phi|_{H_{\bar{q}}} = 0, \quad (13.2.2)$$

where we now have, for complex fields  $\phi_1, \phi_2$ ,

$$Y = \begin{pmatrix} Y_\nu \phi_1 - Y_e \bar{\phi}_2 \\ Y_\nu \phi_2 + Y_e \bar{\phi}_1 \end{pmatrix}, \quad X = \begin{pmatrix} Y_u \phi_1 - Y_d \bar{\phi}_2 \\ Y_u \phi_2 + Y_d \bar{\phi}_1 \end{pmatrix}.$$

The scalar field  $\Phi$  is then given by

$$\Phi = D_F + \begin{pmatrix} \phi & 0 \\ 0 & 0 \end{pmatrix} + J_F \begin{pmatrix} \phi & 0 \\ 0 & 0 \end{pmatrix} J_F^* = \begin{pmatrix} S + \phi & T^* \\ T & (S + \phi) \end{pmatrix}. \quad (13.2.3)$$

**Proposition 13.5** *The action of the gauge group  $S\mathfrak{G}(M \times F_{SM})$  on the fluctuated Dirac operator*

$$D_\omega = D_M \otimes \mathbb{I} + \gamma^\mu \otimes B_\mu + \gamma_M \otimes \Phi$$

is implemented by

$$\begin{aligned} \Lambda_\mu &\mapsto \Lambda_\mu - i\lambda\partial_\mu\bar{\lambda}, \quad Q_\mu \mapsto qQ_\mu q^* - iq\partial_\mu q^*, \quad \bar{V}_\mu \mapsto m\bar{V}_\mu m^* - im\partial_\mu m^*, \\ \begin{pmatrix} \phi_1 + 1 \\ \phi_2 \end{pmatrix} &\mapsto \bar{\lambda}q \begin{pmatrix} \phi_1 + 1 \\ \phi_2 \end{pmatrix}, \end{aligned}$$

for  $\lambda \in C^\infty(M, U(1))$ ,  $q \in C^\infty(M, SU(2))$  and  $m \in C^\infty(M, SU(3))$ .

**Proof** We simply insert the formulas for the fields obtained in (13.2.1) into the transformations given by (10.2.9). Let us write

$$u = (\lambda, q, m) \in C^\infty(M, U(1) \times SU(2) \times SU(3)).$$

The term  $u\omega u^*$  replaces  $Q_\mu$  by  $qQ_\mu q^*$ , and  $\bar{V}_\mu$  by  $m\bar{V}_\mu m^*$ , respectively. We also see that the term  $-iu\partial_\mu u^*$  is given by  $-i\lambda\partial_\mu\bar{\lambda}$  on  $\nu_R, u_R$  and  $H_{\bar{l}}$ , by the expression  $-i\bar{\lambda}\partial_\mu\lambda = i\lambda\partial_\mu\bar{\lambda}$  on  $e_R$  and  $d_R$ , by  $-iq\partial_\mu q^*$  on  $(\nu_L, e_L)$  and  $(u_L, d_L)$ , and, finally, by  $-im\partial_\mu m^*$  on  $H_{\bar{q}}$ . We thus obtain the desired transformation rules for  $\Lambda_\mu$ ,  $Q_\mu$ , and  $\bar{V}_\mu$ .

For the transformation of  $\phi$ , we separately calculate  $u\phi u^*$  and  $u[D_F, u^*]$ . Since  $\phi = 0$  on  $H_{\bar{l}}$  and  $H_{\bar{q}}$ , we may restrict our calculation of  $u\phi u^*$  to  $H_l$  and  $H_q$ . On  $H_l$  we find

$$u\phi u^* = \begin{pmatrix} q_\lambda & 0 \\ 0 & q \end{pmatrix} \begin{pmatrix} 0 & Y^* \\ Y & 0 \end{pmatrix} \begin{pmatrix} q_\lambda^* & 0 \\ 0 & q^* \end{pmatrix} = \begin{pmatrix} 0 & q_\lambda Y^* q^* \\ q Y q_\lambda^* & 0 \end{pmatrix},$$

which is still hermitian. We then calculate

$$\begin{aligned} q Y q_\lambda^* &= \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \begin{pmatrix} Y_\nu \phi_1 - Y_e \bar{\phi}_2 \\ Y_\nu \phi_2 + Y_e \bar{\phi}_1 \end{pmatrix} \begin{pmatrix} \bar{\lambda} & 0 \\ 0 & \lambda \end{pmatrix} \\ &= \begin{pmatrix} \bar{\lambda} Y_\nu (\alpha \phi_1 + \beta \phi_2) & \lambda Y_e (\beta \bar{\phi}_1 - \alpha \bar{\phi}_2) \\ \bar{\lambda} Y_\nu (-\bar{\beta} \phi_1 + \bar{\alpha} \phi_2) & \lambda Y_e (\bar{\alpha} \bar{\phi}_1 + \bar{\beta} \bar{\phi}_2) \end{pmatrix}. \end{aligned}$$

A similar computation on  $H_q$  gives the same transformation for the  $\phi_1$  and  $\phi_2$ .

Next, let us calculate the second term  $u[D_F, u^*]$ . The operator  $T$  in  $D_F$  only acts on  $\nu_R$ , and therefore commutes with the algebra. Upon restricting to  $H_{\bar{l}}$  and  $\mathcal{H}_{\bar{q}}$ , the operator  $\bar{S}$  commutes with the algebra. Hence, once again we may restrict our calculation to  $H_l$  and  $H_q$ . The term  $u[S, u^*]$  is  $u S u^* - S$  and we compute

$$u S u^* = \begin{pmatrix} 0 & q_\lambda Y_0^* q^* \\ q Y_0 q_\lambda^* & 0 \end{pmatrix},$$

where  $Y_0 = \begin{pmatrix} Y_\nu & 0 \\ 0 & Y_e \end{pmatrix}$  on  $H_l$  and  $Y_0 = \begin{pmatrix} Y_u & 0 \\ 0 & Y_d \end{pmatrix}$  on  $H_q$ . We find that on  $H_l$ ,

$$q Y_0 q_\lambda^* = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \begin{pmatrix} Y_\nu & 0 \\ 0 & Y_e \end{pmatrix} \begin{pmatrix} \bar{\lambda} & 0 \\ 0 & \lambda \end{pmatrix} = \begin{pmatrix} \bar{\lambda} Y_\nu \alpha & \lambda Y_e \beta \\ -\bar{\lambda} Y_\nu \bar{\beta} & \lambda Y_e \bar{\alpha} \end{pmatrix},$$

and a similar expression holds on  $H_q$  after replacing  $Y_\nu$  and  $Y_e$  by  $Y_u$  and  $Y_d$ , respectively.

Combining the two contributions to the transformation, we find that the transformation  $u\phi u^* + u[S, u^*]$  maps

$$Y = \begin{pmatrix} Y_\nu \phi_1 - Y_e \bar{\phi}_2 \\ Y_\nu \phi_2 + Y_e \bar{\phi}_1 \end{pmatrix} \mapsto Y' = \begin{pmatrix} Y_\nu \phi'_1 - Y_e \bar{\phi}'_2 \\ Y_\nu \phi'_2 + Y_e \bar{\phi}'_1 \end{pmatrix},$$

where we defined

$$\phi'_1 := \bar{\lambda}(\alpha \phi_1 + \beta \phi_2 + \alpha) - 1, \quad \phi'_2 := \bar{\lambda}(-\bar{\beta} \phi_1 + \bar{\alpha} \phi_2 - \bar{\beta}).$$

Rewriting this in terms of  $q$  completes the proof.  $\square$

Summarizing, the gauge fields derived from  $F_{SM}$  take values in the Lie algebra  $u(1) \oplus su(2) \oplus su(3)$  and transform according to the usual Standard Model gauge transformations. The scalar field  $\phi$  transforms as the Standard Model Higgs field in the defining representation of  $SU(2)$ , with hypercharge  $-1$ .

### 13.3 The Spectral Action

In this section we calculate the spectral action for the almost-commutative manifold  $M \times F_{SM}$  and derive the bosonic part of the Lagrangian of the Standard Model. The general form of this Lagrangian has already been calculated for almost-commutative manifolds in Sect. 10.12, so we only need to insert the expressions (13.2.1) and (13.2.3) for the fields  $\Phi$  and  $B_\mu$ . We start with a few lemmas that capture the rather tedious calculations that are needed to obtain the traces of  $F_{\mu\nu}F^{\mu\nu}$ ,  $\Phi^2$ ,  $\Phi^4$  and  $(D_\mu\Phi)(D^\mu\Phi)$ .

We denote the curvatures of the  $U(1)$ ,  $SU(2)$  and  $SU(3)$  gauge fields by

$$\begin{aligned}\Lambda_{\mu\nu} &:= \partial_\mu\Lambda_\nu - \partial_\nu\Lambda_\mu, \\ Q_{\mu\nu} &:= \partial_\mu Q_\nu - \partial_\nu Q_\mu + i[Q_\mu, Q_\nu], \\ V_{\mu\nu} &:= \partial_\mu V_\nu - \partial_\nu V_\mu + i[V_\mu, V_\nu].\end{aligned}\tag{13.3.1}$$

**Lemma 13.6** *The trace of the square of the curvature of  $B_\mu$  is given by*

$$\mathrm{Tr}_{H_F}(F_{\mu\nu}F^{\mu\nu}) = 24\left(\frac{10}{3}\Lambda_{\mu\nu}\Lambda^{\mu\nu} + \mathrm{Tr}(Q_{\mu\nu}Q^{\mu\nu}) + \mathrm{Tr}(V_{\mu\nu}V^{\mu\nu})\right).$$

**Proof** Let us first consider the trace over the lepton sector. Using (13.2.1), we find that the curvature  $F_{\mu\nu}$  of  $B_\mu$  can be written as

$$\begin{aligned}F_{\mu\nu}\Big|_{H_l} &= \begin{pmatrix} 0 & 0 \\ 0 & -2\Lambda_{\mu\nu} \\ & Q_{\mu\nu} - \Lambda_{\mu\nu}\mathbb{I}_2 \end{pmatrix}, \\ F_{\mu\nu}\Big|_{H_{\bar{l}}} &= \begin{pmatrix} 0 & 0 \\ 0 & 2\Lambda_{\mu\nu} \\ & \Lambda_{\mu\nu}\mathbb{I}_2 - (\bar{Q})_{\mu\nu} \end{pmatrix},\end{aligned}$$

where  $(\bar{Q})_{\mu\nu}$  is the curvature of  $\bar{Q}_\mu$ . The square of the curvature therefore becomes

$$\begin{aligned}F_{\mu\nu}F^{\mu\nu}\Big|_{H_l} &= \begin{pmatrix} 0 & 0 \\ 0 & 4\Lambda_{\mu\nu}\Lambda^{\mu\nu} \\ & Q_{\mu\nu}Q^{\mu\nu} + \Lambda_{\mu\nu}\Lambda^{\mu\nu}\mathbb{I}_2 - 2\Lambda_{\mu\nu}Q^{\mu\nu} \end{pmatrix}, \\ F_{\mu\nu}F^{\mu\nu}\Big|_{H_{\bar{l}}} &= \begin{pmatrix} 0 & 0 \\ 0 & 4\Lambda_{\mu\nu}\Lambda^{\mu\nu} \\ & (\bar{Q})_{\mu\nu}(\bar{Q})^{\mu\nu} + \Lambda_{\mu\nu}\Lambda^{\mu\nu}\mathbb{I}_2 - 2\Lambda_{\mu\nu}(\bar{Q})^{\mu\nu} \end{pmatrix}.\end{aligned}$$

Since  $Q_{\mu\nu}$  is traceless, the cross-term  $-2\Lambda_{\mu\nu}Q^{\mu\nu}$  drops out after taking the trace. Note that since  $Q_\mu$  is hermitian we have  $\bar{Q}_\mu = Q_\mu^T$ , and this also holds for  $\bar{Q}_{\mu\nu}$ . This implies that

$$\mathrm{Tr}(\overline{(Q_{\mu\nu})(Q^{\mu\nu})}) = \mathrm{Tr}((Q_{\mu\nu})^T(Q^{\mu\nu})^T) = \mathrm{Tr}(Q_{\mu\nu}Q^{\mu\nu}).$$

Thus, with three generations we obtain

$$\mathrm{Tr}_{H_l \oplus H_{\bar{l}}} (F_{\mu\nu} F^{\mu\nu}) = 36 \Lambda_{\mu\nu} \Lambda^{\mu\nu} + 6 \mathrm{Tr}(Q_{\mu\nu} Q^{\mu\nu}).$$

For the quark sector, on  $H_q$ , we obtain the curvature

$$F_{\mu\nu}|_{H_q} = \begin{pmatrix} \frac{4}{3} \Lambda_{\mu\nu} \mathbb{I}_3 + V_{\mu\nu} & 0 \\ 0 & -\frac{2}{3} \Lambda_{\mu\nu} \mathbb{I}_3 + V_{\mu\nu} \\ & (Q_{\mu\nu} + \frac{1}{3} \Lambda_{\mu\nu} \mathbb{I}_2) \otimes \mathbb{I}_3 + \mathbb{I}_2 \otimes V_{\mu\nu} \end{pmatrix},$$

where we have defined the curvature of the  $SU(3)$  gauge field by

$$V_{\mu\nu} := \partial_\mu V_\nu - \partial_\nu V_\mu + i[V_\mu, V_\nu].$$

A similar expression can be derived on  $H_{\bar{q}}$ .

If we calculate the trace of the square of the curvature  $F_{\mu\nu}$ , the cross-terms again vanish, so we obtain

$$\begin{aligned} \mathrm{Tr}|_{H_q} (F_{\mu\nu} F^{\mu\nu}) &= \left( \frac{16}{3} + \frac{4}{3} + \frac{1}{3} + \frac{1}{3} \right) \Lambda_{\mu\nu} \Lambda^{\mu\nu} \\ &\quad + 3 \mathrm{Tr}(Q_{\mu\nu} Q^{\mu\nu}) + 4 \mathrm{Tr}(V_{\mu\nu} V^{\mu\nu}). \end{aligned}$$

We multiply this by a factor of 2 to include the trace over the anti-quarks, and by a factor of 3 for the number of generations. Adding the result to the trace over the lepton sector, we finally obtain

$$\mathrm{Tr}(F_{\mu\nu} F^{\mu\nu}) = 80 \Lambda_{\mu\nu} \Lambda^{\mu\nu} + 24 \mathrm{Tr}(Q_{\mu\nu} Q^{\mu\nu}) + 24 \mathrm{Tr}(V_{\mu\nu} V^{\mu\nu}). \quad \square$$

**Lemma 13.7** *The traces of  $\Phi^2$  and  $\Phi^4$  are given by*

$$\begin{aligned} \mathrm{Tr}(\Phi^2) &= 4a|H|^2 + 2c, \\ \mathrm{Tr}(\Phi^4) &= 4b|H|^4 + 8e|H|^2 + 2d, \end{aligned}$$

where  $H$  denotes the complex doublet  $(\phi_1 + 1, \phi_2)$  and

$$\begin{aligned} a &= \mathrm{Tr}(Y_\nu^* Y_\nu + Y_e^* Y_e + 3Y_u^* Y_u + 3Y_d^* Y_d), \\ b &= \mathrm{Tr}((Y_\nu^* Y_\nu)^2 + (Y_e^* Y_e)^2 + 3(Y_u^* Y_u)^2 + 3(Y_d^* Y_d)^2), \\ c &= \mathrm{Tr}(Y_R^* Y_R), \\ d &= \mathrm{Tr}((Y_R^* Y_R)^2), \\ e &= \mathrm{Tr}(Y_R^* Y_R Y_\nu^* Y_\nu). \end{aligned} \tag{13.3.2}$$

**Proof** The field  $\Phi$  is given by (13.2.3), and its square equals

$$\Phi^2 = \begin{pmatrix} (S + \phi)^2 + T^*T & (S + \phi)T^* + T^*\overline{(S + \phi)} \\ T(S + \phi) + \overline{(S + \phi)}T & \overline{(S + \phi)}^2 + TT^* \end{pmatrix}.$$

The square of the off-diagonal part yields  $T^*T = TT^* = |Y_R|^2$  on  $\nu_R$  and  $\overline{\nu_R}$ , and zero on  $l \neq \nu_R, \overline{\nu_R}$ . On the lepton sector of the Hilbert space, the component  $S + \phi$  is given by

$$S + \phi|_{H_l} = \begin{pmatrix} 0 & Y^* + Y_0^* \\ Y + Y_0 & 0 \end{pmatrix}.$$

We then calculate

$$\mathfrak{X} := (Y + Y_0)^*(Y + Y_0) = |H|^2 \begin{pmatrix} |Y_\nu|^2 & 0 \\ 0 & |Y_e|^2 \end{pmatrix},$$

where we defined the complex doublet  $H := (\phi_1 + 1, \phi_2)$ . Similarly, we define  $\mathfrak{X}' := (Y + Y_0)(Y + Y_0)^*$ , and note that  $\text{Tr}(\mathfrak{X}) = \text{Tr}(\mathfrak{X}')$  by the cyclic property of the trace. Since  $\mathfrak{X} = \mathfrak{X}^*$  and  $\text{Tr}(\mathfrak{X}) = \text{Tr}(\mathfrak{X}^T)$ , we also have  $\text{Tr}(\overline{\mathfrak{X}}) = \text{Tr}(\mathfrak{X})$ . Thus, on the lepton sector we obtain

$$\begin{aligned} \text{Tr}_{H_l \oplus H_{\bar{l}}}(\Phi^2) &= \text{Tr}(\mathfrak{X} + \mathfrak{X}' + \overline{\mathfrak{X}} + \overline{\mathfrak{X}'}) + 2|Y_R|^2 \\ &= 4\text{Tr}(\mathfrak{X}) + 2|Y_R|^2 = 4(|Y_\nu|^2 + |Y_e|^2)|H|^2 + 2|Y_R|^2. \end{aligned}$$

On the quark sector we similarly find

$$\text{Tr}_{H_q \oplus H_{\bar{q}}}(\Phi^2) = 4 \cdot 3(|Y_\nu|^2 + |Y_e|^2)|H|^2,$$

leading to the stated formula for  $\text{Tr}(\Phi^2)$ .

In order to find the trace of  $\Phi^4$ , we calculate

$$(\mathfrak{X} + T^*T)^2 = |H|^4 \begin{pmatrix} |Y_\nu|^4 & 0 \\ 0 & |Y_e|^4 \end{pmatrix} + 2|H|^2 \begin{pmatrix} |Y_R|^2|Y_\nu|^2 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} |Y_R|^4 & 0 \\ 0 & 0 \end{pmatrix}.$$

We hence obtain

$$\begin{aligned} \text{Tr}_{H_l \oplus H_{\bar{l}}}(\Phi^4) &= \text{Tr}(4\mathfrak{X}^2 + 4\mathfrak{X}T^*T + 2(T^*T)^2) + 4|H|^2|Y_R|^2|Y_\nu|^2 \\ &= 4|H|^4(|Y_\nu|^4 + |Y_e|^4) + 8|H|^2|Y_R|^2|Y_\nu|^2 + 2|Y_R|^4. \end{aligned}$$

On the quark sector, we obtain a similar result with  $Y_\nu$  replaced by  $Y_u$  and  $Y_e$  by  $Y_d$ , leaving out the  $Y_R$ , and including a factor of 3 for the trace in colour space.  $\square$

**Lemma 13.8** *The trace of  $(D_\mu \Phi)(D^\mu \Phi)$  is given by*

$$\mathrm{Tr} \left( (D_\mu \Phi) (D^\mu \Phi) \right) = 4a |D_\mu H|^2,$$

where  $H$  denotes the complex doublet  $(\phi_1 + 1, \phi_2)$ , and the covariant derivative  $D_\mu$  on  $H$  is defined as

$$D_\mu H = \partial_\mu H + i Q_\mu^a \sigma^a H - i \Lambda_\mu H.$$

**Proof** We need to calculate the commutator  $[B_\mu, \Phi]$ . We note that  $B_\mu$  commutes with the off-diagonal part of  $D_F$ . It is therefore sufficient to calculate the commutator  $[B_\mu, S + \phi]$  on  $H_l$ . We shall write  $Q_\mu = Q_\mu^1 \sigma^1 + Q_\mu^2 \sigma^2 + Q_\mu^3 \sigma^3$  as a linear combination of Pauli matrices with real coefficients  $Q_\mu^a$ . By direct calculation on the lepton sector, we then obtain

$$[B_\mu, S + \phi]|_{H_l} = \begin{pmatrix} 0 & 0 & -\bar{Y}_\nu \chi_1 & -\bar{Y}_\nu \bar{\chi}_2 \\ 0 & 0 & -\bar{Y}_e \chi_2 & \bar{Y}_e \bar{\chi}_1 \\ Y_\nu \chi_1 & Y_e \bar{\chi}_2 & 0 & 0 \\ Y_\nu \chi_2 & -Y_e \bar{\chi}_1 & 0 & 0 \end{pmatrix},$$

where we defined the new doublet  $\chi = (\chi_1, \chi_2)$  by

$$\begin{aligned} \chi_1 &:= (\phi_1 + 1)(Q_\mu^3 - \Lambda_\mu) + \phi_2(Q_\mu^1 - iQ_\mu^2), \\ \chi_2 &:= (\phi_1 + 1)(Q_\mu^1 + iQ_\mu^2) + \phi_2(-Q_\mu^3 - \Lambda_\mu). \end{aligned}$$

We then obtain

$$\begin{aligned} D_\mu(S + \phi)|_{H_l} &= \partial_\mu \phi + i[B_\mu, S + \phi] \\ &= \begin{pmatrix} 0 & 0 & \bar{Y}_\nu(\partial_\mu \bar{\phi}_1 - i\bar{\chi}_1) & \bar{Y}_\nu(\partial_\mu \bar{\phi}_2 - i\bar{\chi}_2) \\ 0 & 0 & -\bar{Y}_e(\partial_\mu \phi_2 + i\chi_2) & \bar{Y}_e(\partial_\mu \phi_1 + i\chi_1) \\ Y_\nu(\partial_\mu \phi_1 + i\chi_1) - Y_e(\partial_\mu \bar{\phi}_2 - i\bar{\chi}_2) & 0 & 0 & 0 \\ Y_\nu(\partial_\mu \phi_2 + i\chi_2) & Y_e(\partial_\mu \phi_1 - i\chi_1) & 0 & 0 \end{pmatrix}. \end{aligned}$$

As  $\phi$  commutes with the gauge field  $V_\mu$ , the corresponding formula for  $D_\mu(S + \phi)$  on the quark sector is identical (after having tensored with  $\mathbb{I}_3$  in colour space).

Since we want to calculate the trace of the square of  $D_\mu \Phi$ , it is sufficient to determine only the terms on the diagonal of  $(D_\mu \Phi)(D^\mu \Phi)$ . We find

$$\mathrm{Tr}_{H_l \oplus H_q} \left( (D_\mu(S + \phi))(D^\mu(S + \phi)) \right) = 2a \left( |\partial_\mu \phi_1 + i\chi_1|^2 + |\partial_\mu \phi_2 + i\chi_2|^2 \right),$$

where we have used

$$a = \mathrm{Tr} (Y_\nu^* Y_\nu + Y_e^* Y_e + 3Y_u^* Y_u + 3Y_d^* Y_d)$$

as in (13.3.2). The column vector  $H$  is given by the complex doublet  $(\phi_1 + 1, \phi_2)$ . We then note that  $\partial_\mu \phi + i\chi$  is equal to the covariant derivative  $D_\mu H$ , so that

$$\mathrm{Tr}_{H_l \oplus H_q} \left( (D_\mu(S + \phi))(D^\mu(S + \phi)) \right) = 2a|D_\mu H|^2.$$

The trace over  $H_{\bar{l}} \oplus H_{\bar{q}}$  yields exactly the same contribution, so we need to multiply this by 2, which gives the desired result.  $\square$

**Proposition 13.9** *The spectral action of the almost-commutative manifold  $M \times F_{SM}$  is given by*

$$\mathrm{Tr} \left( f \left( \frac{D_\omega}{\Lambda} \right) \right) \sim \int_M \mathcal{L}(g_{\mu\nu}, \Lambda_\mu, Q_\mu, V_\mu, H) \sqrt{g} d^4x + O(\Lambda^{-1}),$$

for the Lagrangian

$$\mathcal{L}(g_{\mu\nu}, \Lambda_\mu, Q_\mu, V_\mu, H) := 96\mathcal{L}_M(g_{\mu\nu}) + \mathcal{L}_A(\Lambda_\mu, Q_\mu, V_\mu) + \mathcal{L}_H(g_{\mu\nu}, \Lambda_\mu, Q_\mu, H),$$

where  $\mathcal{L}_M(g_{\mu\nu})$  is defined in Proposition 10.10,  $\mathcal{L}_A$  gives the kinetic terms of the gauge fields as

$$\mathcal{L}_A(\Lambda_\mu, Q_\mu, V_\mu) := \frac{f(0)}{\pi^2} \left( \frac{10}{3} \Lambda_{\mu\nu} \Lambda^{\mu\nu} + \mathrm{Tr}(Q_{\mu\nu} Q^{\mu\nu}) + \mathrm{Tr}(V_{\mu\nu} V^{\mu\nu}) \right),$$

and the Higgs potential  $\mathcal{L}_H$  (ignoring the boundary term) equals

$$\begin{aligned} \mathcal{L}_H(g_{\mu\nu}, \Lambda_\mu, Q_\mu, H) := & \frac{bf(0)}{2\pi^2} |H|^4 + \frac{-2af_2\Lambda^2 + ef(0)}{\pi^2} |H|^2 \\ & - \frac{cf_2\Lambda^2}{\pi^2} + \frac{df(0)}{4\pi^2} + \frac{af(0)}{12\pi^2} s |H|^2 + \frac{cf(0)}{24\pi^2} s + \frac{af(0)}{2\pi^2} |D_\mu H|^2. \end{aligned}$$

**Proof** We use the general form of the spectral action of an almost-commutative manifold as calculated in Proposition 10.12, and combine it with the previous Lemmas. The gravitational Lagrangian  $\mathcal{L}_M$  obtains a factor 96 from the trace over  $H_F$ . From Lemma 13.6 we immediately find the term  $\mathcal{L}_A$ . Combining the formulas of  $\mathrm{Tr}(\Phi^2)$  and  $\mathrm{Tr}(\Phi^4)$  obtained in Lemma 13.7, we find the Higgs potential

$$\begin{aligned} & - \frac{f_2\Lambda^2}{2\pi^2} \mathrm{Tr}(\Phi^2) + \frac{f(0)}{8\pi^2} \mathrm{Tr}(\Phi^4) \\ & = \frac{bf(0)}{2\pi^2} |H|^4 + \frac{-2af_2\Lambda^2 + ef(0)}{\pi^2} |H|^2 - \frac{cf_2\Lambda^2}{\pi^2} + \frac{df(0)}{4\pi^2}. \end{aligned}$$

The coupling of the Higgs field to the scalar curvature  $s$  is given by

$$\frac{f(0)}{48\pi^2} s \mathrm{Tr}(\Phi^2) = \frac{af(0)}{12\pi^2} s |H|^2 + \frac{cf(0)}{24\pi^2} s,$$

where the second term yields a contribution to the Einstein-Hilbert term  $-\frac{f_2 \Lambda^2}{3\pi^2} s$  of  $\mathcal{L}_M$ . Finally, the kinetic term of the Higgs field including minimal coupling to the gauge fields is obtained from Lemma 13.8 as

$$\frac{f(0)}{8\pi^2} \text{Tr}((D_\mu \Phi)(D^\mu \Phi)) = \frac{af(0)}{2\pi^2} |D_\mu H|^2. \quad \square$$

### 13.3.1 Coupling Constants and Unification

In Proposition 13.9 we calculated the bosonic Lagrangian from the spectral action. We now rescale the Higgs and gauge fields  $\Lambda_\mu, Q_\mu, V_\mu$  in such a way that their kinetic terms are properly normalized.

We start with the Higgs field, and require that its kinetic term is normalized as usual, i.e.,

$$\int_M \frac{1}{2} |D_\mu H|^2 \sqrt{g} d^4x.$$

This normalization is evidently achieved by rescaling the Higgs field as

$$H \mapsto \sqrt{\frac{\pi^2}{af(0)}} H. \quad (13.3.3)$$

Next, write the non-abelian gauge fields as  $Q_\mu = Q_\mu^a \sigma^a$  and  $V_\mu = V_\mu^i \lambda^i$ , for the Gell-Mann matrices  $\lambda^i$  and real coefficients  $V_\mu^i$ . We introduce coupling constants  $g_1, g_2$  and  $g_3$  into the model by rescaling the gauge fields as

$$\Lambda_\mu = \frac{1}{2} g_1 Y_\mu, \quad Q_\mu^a = \frac{1}{2} g_2 W_\mu^a, \quad V_\mu^i = \frac{1}{2} g_3 G_\mu^i.$$

Using the relations  $\text{Tr}(\sigma^a \sigma^b) = 2\delta^{ab}$  and  $\text{Tr}(\lambda^i \lambda^j) = 2\delta^{ij}$ , we now find that the Lagrangian  $\mathcal{L}_A$  of Proposition 13.9 can be written as

$$\mathcal{L}_A(Y_\mu, W_\mu, G_\mu) = \frac{f(0)}{2\pi^2} \left( \frac{5}{3} g_1^2 Y_{\mu\nu} Y^{\mu\nu} + g_2^2 W_{\mu\nu} W^{\mu\nu} + g_3^2 G_{\mu\nu} G^{\mu\nu} \right).$$

It is natural to require that these kinetic terms are properly normalized, and this imposes the relations

$$\frac{f(0)}{2\pi^2} g_3^2 = \frac{f(0)}{2\pi^2} g_2^2 = \frac{5f(0)}{6\pi^2} g_1^2 = \frac{1}{4}. \quad (13.3.4)$$

The coupling constants are then related by

$$g_3^2 = g_2^2 = \frac{5}{3}g_1^2, \quad (13.3.5)$$

which is precisely the relation between the coupling constants at unification, common to grand unified theories (GUT). We shall further discuss this in Sect. 14.2.

In terms of the rescaled fields, we obtain the following result:

**Theorem 13.10** *The spectral action (ignoring topological and boundary terms) of the almost-commutative manifold  $M \times F_{SM}$  is given by*

$$\begin{aligned} S_B = & \int_M \left( \frac{48f_4\Lambda^4}{\pi^2} - \frac{cf_2\Lambda^2}{\pi^2} + \frac{df(0)}{4\pi^2} + \left( \frac{cf(0)}{24\pi^2} - \frac{4f_2\Lambda^2}{\pi^2} \right)s - \frac{3f(0)}{10\pi^2}(C_{\mu\nu\rho\sigma})^2 \right. \\ & + \frac{1}{4}Y_{\mu\nu}Y^{\mu\nu} + \frac{1}{4}W_{\mu\nu}^a W^{\mu\nu,a} + \frac{1}{4}G_{\mu\nu}^i G^{\mu\nu,i} + \frac{b\pi^2}{2a^2 f(0)}|H|^4 \\ & \left. - \frac{2af_2\Lambda^2 - ef(0)}{af(0)}|H|^2 + \frac{1}{12}s|H|^2 + \frac{1}{2}|D_\mu H|^2 \right) \sqrt{g} d^4x, \end{aligned}$$

where the covariant derivative  $D_\mu H$  is given by

$$D_\mu H = \partial_\mu H + \frac{1}{2}ig_2 W_\mu^a \sigma^a H - \frac{1}{2}ig_1 Y_\mu H. \quad (13.3.6)$$

### 13.3.2 The Higgs Mechanism


Writing down a gauge theory with massive gauge bosons, one encounters the notorious difficulty that the mass terms of these gauge bosons are not gauge invariant. The Higgs field plays a central role in obtaining these mass terms within a gauge theory. The celebrated Higgs mechanism provides a *spontaneous breaking* of the gauge symmetry and thus generates mass terms. In this section we describe how the Higgs mechanism breaks the  $U(1) \times SU(2)$  symmetry and introduces mass terms for some of the gauge bosons of the Standard Model.

In Theorem 13.10 we obtained the Higgs Lagrangian  $\mathcal{L}_H$ . If we drop all the terms that are independent of the Higgs field  $H$ , and also ignore the coupling of the Higgs to the gravitational field, we obtain the Lagrangian

$$\mathcal{L}(g_{\mu\nu}, Y_\mu, W_\mu^a, H) := \frac{b\pi^2}{2a^2 f(0)}|H|^4 - \frac{2af_2\Lambda^2 - ef(0)}{af(0)}|H|^2 + \frac{1}{2}|D_\mu H|^2. \quad (13.3.7)$$

We wish to find the value of  $H$  for which this Lagrangian obtains its minimum value.

Hence, we consider the Higgs potential



**Fig. 13.3** The potential  $\mathcal{L}_{\text{pot}}(H)$  of (13.3.8) with  $2af_2\Lambda^2 > ef(0)$

$$\mathcal{L}_{\text{pot}}(H) := \frac{b\pi^2}{2a^2 f(0)} |H|^4 - \frac{2af_2\Lambda^2 - ef(0)}{af(0)} |H|^2. \quad (13.3.8)$$

If  $2af_2\Lambda^2 < ef(0)$ , the minimum of this potential is obtained at  $H = 0$ , and in this case there will be no symmetry breaking. Indeed, the minimum  $H = 0$  is symmetric under the full symmetry group  $U(1) \times SU(2)$ .

We now assume that  $2af_2\Lambda^2 > ef(0)$ , so that the potential has the form depicted in Fig. 13.3. The minimum of the Higgs potential is then reached if the field  $H$  satisfies

$$|H|^2 = \frac{2a^2 f_2\Lambda^2 - af(0)}{b\pi^2}, \quad (13.3.9)$$

and none such minimum is invariant any more under  $U(1) \times SU(2)$ . The fields that satisfy this relation are called the *vacuum states* of the Higgs field. We choose a vacuum state  $(v, 0)$ , where the *vacuum expectation value*  $v$  is a real parameter such that  $v^2$  is given by the right-hand side of (13.3.9). From the transformation rule of Proposition 13.5, we see that the vacuum state  $(v, 0)$  is still invariant under a subgroup of  $U(1) \times SU(2)$ . This subgroup is isomorphic to  $U(1)$  and is given by

$$\left\{ \left( \lambda, q_\lambda = \begin{pmatrix} \lambda & 0 \\ 0 & \bar{\lambda} \end{pmatrix} \right) : \lambda \in U(1) \right\} \subset U(1) \times SU(2).$$

Let us simplify the expression for the Higgs potential. First, we note that the potential only depends on the absolute value  $|H|$ . A transformation of the doublet  $H$  by an element  $(\lambda, q) \in U(1) \times SU(2)$  is written as  $H \mapsto uH$  with  $u = \bar{\lambda}q$  a unitary matrix. Since a unitary transformation preserves absolute values, we see that  $\mathcal{L}_{\text{pot}}(uH) = \mathcal{L}_{\text{pot}}(H)$  for any  $u \in U(1) \times SU(2)$ . We can use this *gauge freedom* to transform the Higgs field into a simpler form. Consider elements of  $SU(2)$  of the form

$$\begin{pmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix}$$

such that  $|\alpha|^2 + |\beta|^2 = 1$ . The doublet  $H$  can in general be written as  $(h_1, h_2)$ , for some  $h_1, h_2 \in \mathbb{C}$ . We then see that we may write

$$\begin{pmatrix} h_1 \\ h_2 \end{pmatrix} = \begin{pmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix} \begin{pmatrix} |H| \\ 0 \end{pmatrix}, \quad \alpha = \frac{h_1}{|H|}, \quad \beta = \frac{h_2}{|H|},$$

which means that we may always use the gauge freedom to write the doublet  $H$  in terms of one real parameter. Let us define a new real-valued field  $h$  by setting  $h(x) := |H(x)| - v$ . We then obtain

$$H = u(x) \begin{pmatrix} v + h(x) \\ 0 \end{pmatrix}, \quad u(x) := \begin{pmatrix} \alpha(x) & -\bar{\beta}(x) \\ \beta(x) & \bar{\alpha}(x) \end{pmatrix}. \quad (13.3.10)$$

Inserting this transformed Higgs field into the Higgs potential, we obtain the following expression in terms of the real parameter  $v$  and the real field  $h(x)$ :

$$\begin{aligned} \mathcal{L}_{\text{pot}}(h) &= \frac{bf(0)}{2\pi^2} (v + h)^4 - \frac{2af_2\Lambda^2 - ef(0)}{\pi^2} (v + h)^2 \\ &= \frac{b\pi^2}{2a^2f(0)} (h^4 + 4vh^3 + 6v^2h^2 + 4v^3h + v^4) \\ &\quad - \frac{2af_2\Lambda^2 - ef(0)}{af(0)} (h^2 + 2vh + v^2). \end{aligned}$$

Using (13.3.9), the value of  $v^2$  is given by

$$v^2 = \frac{2a^2f_2\Lambda^2 - af(0)}{b\pi^2}.$$

We then see that in  $\mathcal{L}_{\text{pot}}$  the terms linear in  $h$  cancel out. This is of course no surprise, since the change of variables  $|H(x)| \mapsto v + h(x)$  means that at  $h(x) = 0$  we are at the minimum of the potential, where the first order derivative of the potential with respect to  $h$  must vanish. We thus obtain the simplified expression

$$\mathcal{L}_{\text{pot}}(h) = \frac{b\pi^2}{2a^2 f(0)} \left( h^4 + 4vh^3 + 4v^2h^2 - v^4 \right). \quad (13.3.11)$$

We now observe that the field  $h(x)$  has acquired a mass term and has two self-interactions given by  $h^3$  and  $h^4$ . We also have another contribution to the cosmological constant, given by  $-v^4$ .

### 13.3.2.1 Massive Gauge Bosons

Next, let us consider what this procedure entails for the remainder of the Higgs Lagrangian  $\mathcal{L}_H$ . We first consider the kinetic term of  $H$ , including its minimal coupling to the gauge fields, given by

$$\mathcal{L}_{\text{min}}(Y_\mu, W_\mu^a, H) := \frac{1}{2} |D_\mu H|^2.$$

The transformation of (13.3.10) is a gauge transformation, and to make sure that  $\mathcal{L}_{\text{min}}$  is invariant under this transformation, we also need to transform the gauge fields. The field  $Y_\mu$  is unaffected by the local  $SU(2)$ -transformation  $u(x)$ . The transformation of  $W_\mu = W_\mu^a \sigma^a$  is obtained from Proposition 13.5 and is given by

$$W_\mu \rightarrow u W_\mu u^* - \frac{2i}{g_2} u \partial_\mu u^*.$$

One then easily checks that we obtain the transformation  $D_\mu H \mapsto u D_\mu H$ , so that  $|D_\mu H|^2$  is invariant under such transformations. So we can just insert the doublet  $(v + h, 0)$  into (13.3.6) and obtain

$$\begin{aligned} D_\mu H &= \partial_\mu \begin{pmatrix} v + h \\ 0 \end{pmatrix} + \frac{1}{2} i g_2 W_\mu^a \sigma^a \begin{pmatrix} v + h \\ 0 \end{pmatrix} - \frac{1}{2} i g_1 Y_\mu \begin{pmatrix} v + h \\ 0 \end{pmatrix} \\ &= \partial_\mu \begin{pmatrix} h \\ 0 \end{pmatrix} + \frac{1}{2} i g_2 W_\mu^1 \begin{pmatrix} 0 \\ v + h \end{pmatrix} + \frac{1}{2} i g_2 W_\mu^2 \begin{pmatrix} 0 \\ i(v + h) \end{pmatrix} \\ &\quad + \frac{1}{2} i g_2 W_\mu^3 \begin{pmatrix} v + h \\ 0 \end{pmatrix} - \frac{1}{2} i g_1 Y_\mu \begin{pmatrix} v + h \\ 0 \end{pmatrix}. \end{aligned}$$

We can then calculate its square as

$$\begin{aligned} |D_\mu H|^2 &= (D^\mu H)^*(D_\mu H) \\ &= (\partial^\mu h)(\partial_\mu h) + \frac{1}{4} g_2^2 (v + h)^2 (W^{\mu,1} W_\mu^1 + W^{\mu,2} W_\mu^2 + W^{\mu,3} W_\mu^3) \\ &\quad + \frac{1}{4} g_1^2 (v + h)^2 B'^\mu Y_\mu - \frac{1}{2} g_1 g_2 (v + h)^2 B'^\mu W_\mu^3. \end{aligned}$$

Note that the last term yields a mixing of the gauge fields  $Y_\mu$  and  $W_\mu^3$ , parametrized by the electroweak mixing angle  $\theta_w$  defined by

$$c_w := \cos \theta_w = \frac{g_2}{\sqrt{g_1^2 + g_2^2}}, \quad s_w := \sin \theta_w = \frac{g_1}{\sqrt{g_1^2 + g_2^2}}.$$

Note that the relation  $g_2^2 = 3g_1^2$  for the coupling constants implies that we obtain the values  $\cos^2 \theta_w = \frac{1}{4}$  and  $\sin^2 \theta_w = \frac{3}{4}$  at the electroweak unification scale  $\Lambda_{EW}$ . Let us now define new gauge fields by

$$\begin{aligned} W_\mu &:= \frac{1}{\sqrt{2}}(W_\mu^1 + iW_\mu^2), & W_\mu^* &:= \frac{1}{\sqrt{2}}(W_\mu^1 - iW_\mu^2), \\ Z_\mu &:= c_w W_\mu^3 - s_w Y_\mu, & A'_\mu &:= s_w W_\mu^3 + c_w Y_\mu, \end{aligned} \quad (13.3.12)$$

where we have added a prime to  $A_\mu$  to distinguish the (photon) field from the general form of the inner fluctuations in Eq. (10.2.1). We now show that the new fields  $Z_\mu$  and  $A'_\mu$  become mass eigenstates. The fields  $W_\mu^1$  and  $W_\mu^2$  were already mass eigenstates, but the fields  $W_\mu$  and  $W_\mu^*$  are chosen so that they obtain a definite charge. We can write

$$\begin{aligned} W_\mu^1 &= \frac{1}{\sqrt{2}}(W_\mu + W_\mu^*), & W_\mu^2 &= \frac{-i}{\sqrt{2}}(W_\mu - W_\mu^*), \\ W_\mu^3 &= s_w A'_\mu + c_w Z_\mu, & Y_\mu &= c_w A'_\mu - s_w Z_\mu, \end{aligned}$$

and inserting this into the expression for  $|D_\mu H|^2$  yields

$$\frac{1}{2}|D_\mu H|^2 = \frac{1}{2}(\partial^\mu h)(\partial_\mu h) + \frac{1}{4}g_2^2(v+h)^2 W^{\mu*}W_\mu + \frac{1}{8}\frac{g_2^2}{c_w^2}(v+h)^2 Z^\mu Z_\mu. \quad (13.3.13)$$

Thus, we see that the fields  $W_\mu$ ,  $W_\mu^*$  and  $Z_\mu$  acquire a mass term (where  $Z_\mu$  has a larger mass than  $W_\mu$ ,  $W_\mu^*$ ) and that the fields  $A'_\mu$  are massless. The (tree-level) masses of the  $W$ -boson and  $Z$ -boson are evidently given by

$$M_W = \frac{1}{2}vg_2, \quad M_Z = \frac{1}{2}v\frac{g_2}{c_w}. \quad (13.3.14)$$

## 13.4 The Fermionic Action

In order to obtain the full Lagrangian for the Standard Model, we also need to calculate the fermionic action  $S_f$  of Definition 9.3. First, let us have a closer look at the fermionic particle fields and their interactions.

By an abuse of notation, let us write  $\nu^\lambda, \bar{\nu}^\lambda, e^\lambda, \bar{e}^\lambda, u^{\lambda c}, \bar{u}^{\lambda c}, d^{\lambda c}, \bar{d}^{\lambda c}$  for a set of independent Dirac spinors. We then write a generic Grassmann vector  $\tilde{\xi} \in \mathcal{H}_{\text{cl}}^+$  as follows:

$$\begin{aligned}\tilde{\xi} = & \nu_L^\lambda \otimes \nu_L^\lambda + \nu_R^\lambda \otimes \nu_R^\lambda + \bar{\nu}_R^\lambda \otimes \overline{\nu_L^\lambda} + \bar{\nu}_L^\lambda \otimes \overline{\nu_R^\lambda} \\ & + e_L^\lambda \otimes e_L^\lambda + e_R^\lambda \otimes e_R^\lambda + \bar{e}_R^\lambda \otimes \overline{e_L^\lambda} + \bar{e}_L^\lambda \otimes \overline{e_R^\lambda} \\ & + u_L^{\lambda c} \otimes u_L^{\lambda c} + u_R^{\lambda c} \otimes u_R^{\lambda c} + \bar{u}_R^{\lambda c} \otimes \overline{u_L^{\lambda c}} + \bar{u}_L^{\lambda c} \otimes \overline{u_R^{\lambda c}} \\ & + d_L^{\lambda c} \otimes d_L^{\lambda c} + d_R^{\lambda c} \otimes d_R^{\lambda c} + \bar{d}_R^{\lambda c} \otimes \overline{d_L^{\lambda c}} + \bar{d}_L^{\lambda c} \otimes \overline{d_R^{\lambda c}},\end{aligned}$$

where in each tensor product it should be clear that the first component is a Weyl spinor, and the second component is a basis element of  $H_F$ . Here  $\lambda = 1, 2, 3$  labels the generation of the fermions, and  $c = r, g, b$  labels the color index of the quarks.

Let us have a closer look at the gauge fields of the electroweak sector. For the physical gauge fields of (13.3.12) we can write

$$\begin{aligned}Q_\mu^1 + i Q_\mu^2 &= \frac{1}{\sqrt{2}} g_2 W_\mu, & Q_\mu^1 - i Q_\mu^2 &= \frac{1}{\sqrt{2}} g_2 W_\mu^*, \\ Q_\mu^3 - \Lambda_\mu &= \frac{g_2}{2c_w} Z_\mu, & \Lambda_\mu &= \frac{1}{2} s_w g_2 A'_\mu - \frac{1}{2} \frac{s_w^2 g_2}{c_w} Z_\mu, \\ -Q_\mu^3 - \Lambda_\mu &= -s_w g_2 A'_\mu + \frac{g_2}{2c_w} (1 - 2c_w^2) Z_\mu, \\ Q_\mu^3 + \frac{1}{3} \Lambda_\mu &= \frac{2}{3} s_w g_2 A'_\mu - \frac{g_2}{6c_w} (1 - 4c_w^2) Z_\mu, \\ -Q_\mu^3 + \frac{1}{3} \Lambda_\mu &= -\frac{1}{3} s_w g_2 A'_\mu - \frac{g_2}{6c_w} (1 + 2c_w^2) Z_\mu.\end{aligned}\tag{13.4.1}$$

Here we have rescaled the Higgs field in (13.3.3), so we can write  $H = \frac{\sqrt{af(0)}}{\pi}(\phi_1 + 1, \phi_2)$ . We parametrize the Higgs field as

$$H = (v + h + i\phi^0, i\sqrt{2}\phi^-),$$

where  $\phi^0$  is real and  $\phi^-$  is complex. We write  $\phi^+$  for the complex conjugate of  $\phi^-$ . Thus, we can write

$$(\phi_1 + 1, \phi_2) = \frac{\pi}{\sqrt{af(0)}}(v + h + i\phi^0, i\sqrt{2}\phi^-).\tag{13.4.2}$$

As in Remark 11.8, we will need to impose a further restriction on the mass matrices in  $D_F$ , in order to obtain physical mass terms in the fermionic action. From here on, we will require that the matrices  $Y_x$  are anti-hermitian, for  $x = \nu, e, u, d$ . We then define the hermitian mass matrices  $m_x$  by writing

$$Y_x =: -i \frac{\sqrt{af(0)}}{\pi v} m_x. \quad (13.4.3)$$

Similarly, we also take  $Y_R$  to be anti-hermitian, and we introduce a hermitian (and symmetric) Majorana mass matrix  $m_R$  by writing

$$Y_R = -i m_R. \quad (13.4.4)$$

**Theorem 13.11** *The fermionic action of the almost-commutative manifold  $M \times F_{SM}$  is given by*

$$S_F = \int_M (\mathcal{L}_{kin} + \mathcal{L}_{gf} + \mathcal{L}_{Hf} + \mathcal{L}_R) \sqrt{g} d^4x,$$

where, suppressing all generation and color indices, the kinetic terms of the fermions are given by

$$\begin{aligned} \mathcal{L}_{kin} := & -i \langle J_M \bar{\nu}, \gamma^\mu \nabla_\mu^S \nu \rangle - i \langle J_M \bar{e}, \gamma^\mu \nabla_\mu^S e \rangle \\ & - i \langle J_M \bar{u}, \gamma^\mu \nabla_\mu^S u \rangle - i \langle J_M \bar{d}, \gamma^\mu \nabla_\mu^S d \rangle, \end{aligned}$$

the minimal coupling of the gauge fields to the fermions is given by

$$\begin{aligned} \mathcal{L}_{gf} := & s_w g_2 A'_\mu \left( - \langle J_M \bar{e}, \gamma^\mu e \rangle + \frac{2}{3} \langle J_M \bar{u}, \gamma^\mu u \rangle - \frac{1}{3} \langle J_M \bar{d}, \gamma^\mu d \rangle \right) \\ & + \frac{g_2}{4c_w} Z_\mu \left( \langle J_M \bar{\nu}, \gamma^\mu (1 + \gamma_M) \nu \rangle + \langle J_M \bar{e}, \gamma^\mu (4s_w^2 - 1 - \gamma_M) e \rangle \right. \\ & \quad + \langle J_M \bar{u}, \gamma^\mu (-\frac{8}{3}s_w^2 + 1 + \gamma_M) u \rangle \\ & \quad \left. + \langle J_M \bar{d}, \gamma^\mu (\frac{4}{3}s_w^2 - 1 - \gamma_M) d \rangle \right) \\ & + \frac{g_2}{2\sqrt{2}} W_\mu \left( \langle J_M \bar{e}, \gamma^\mu (1 + \gamma_M) \nu \rangle + \langle J_M \bar{d}, \gamma^\mu (1 + \gamma_M) u \rangle \right) \\ & + \frac{g_2}{2\sqrt{2}} W_\mu^* \left( \langle J_M \bar{\nu}, \gamma^\mu (1 + \gamma_M) e \rangle + \langle J_M \bar{u}, \gamma^\mu (1 + \gamma_M) d \rangle \right) \\ & + \frac{g_3}{2} G_\mu^i \left( \langle J_M \bar{u}, \gamma^\mu \lambda_i u \rangle + \langle J_M \bar{d}, \gamma^\mu \lambda_i d \rangle \right), \end{aligned}$$

the Yukawa couplings of the Higgs field to the fermions are given by

$$\begin{aligned} \mathcal{L}_{Hf} := & i \left( 1 + \frac{h}{v} \right) \left( \langle J_M \bar{\nu}, m_\nu \nu \rangle + \langle J_M \bar{e}, m_e e \rangle \right. \\ & \quad \left. + \langle J_M \bar{u}, m_u u \rangle + \langle J_M \bar{d}, m_d d \rangle \right) \\ & + \frac{\phi^0}{v} \left( \langle J_M \bar{\nu}, \gamma_M m_\nu \nu \rangle - \langle J_M \bar{e}, \gamma_M m_e e \rangle \right. \\ & \quad \left. + \langle J_M \bar{u}, \gamma_M m_u u \rangle - \langle J_M \bar{d}, \gamma_M m_d d \rangle \right) \end{aligned}$$

$$\begin{aligned}
& + \frac{\phi^-}{\sqrt{2}v} \left( \langle J_M \bar{e}, m_e(1 + \gamma_M)\nu \rangle - \langle J_M \bar{e}, m_\nu(1 - \gamma_M)\nu \rangle \right) \\
& + \frac{\phi^+}{\sqrt{2}v} \left( \langle J_M \bar{\nu}, m_\nu(1 + \gamma_M)e \rangle - \langle J_M \bar{\nu}, m_e(1 - \gamma_M)e \rangle \right) \\
& + \frac{\phi^-}{\sqrt{2}v} \left( \langle J_M \bar{d}, m_d(1 + \gamma_M)u \rangle - \langle J_M \bar{d}, m_u(1 - \gamma_M)u \rangle \right) \\
& + \frac{\phi^+}{\sqrt{2}v} \left( \langle J_M \bar{u}, m_u(1 + \gamma_M)d \rangle - \langle J_M \bar{u}, m_d(1 - \gamma_M)d \rangle \right),
\end{aligned}$$

and, finally, the Majorana masses of the right-handed neutrinos (and left-handed anti-neutrinos) are given by

$$\mathcal{L}_R := i \langle J_M \nu_R, m_R \nu_R \rangle + i \langle J_M \bar{\nu}_L, m_R \bar{\nu}_L \rangle.$$

**Proof** The proof is similar to Proposition 11.7, though the calculations are now a little more complicated. From Definition 9.3 we know that the fermionic action is given by  $S_F = \frac{1}{2} \langle J \tilde{\xi}, D_\omega \tilde{\xi} \rangle$ , where the fluctuated Dirac operator is given by

$$D_\omega = D_M \otimes 1 + \gamma^\mu \otimes B_\mu + \gamma_M \otimes \Phi.$$

We rewrite the inner product on  $\mathcal{H}$  as  $(\xi, \psi) = \int_M \langle \xi, \psi \rangle \sqrt{g} d^4x$ . As in Proposition 11.7, the expressions for  $J \tilde{\xi} = (J_M \otimes J_F) \tilde{\xi}$  and  $(D_M \otimes 1) \tilde{\xi}$  are obtained straightforwardly. Using the symmetry of the form  $(J_M \tilde{\chi}, D_M \tilde{\psi})$ , and then we obtain the kinetic terms as

$$\begin{aligned}
\frac{1}{2} \langle J \tilde{\xi}, (D_M \otimes 1) \tilde{\xi} \rangle &= \langle J_M \bar{\nu}^\lambda, D_M \nu^\lambda \rangle + \langle J_M \bar{e}^\lambda, D_M e^\lambda \rangle \\
&\quad + \langle J_M \bar{u}^{\lambda c}, D_M u^{\lambda c} \rangle + \langle J_M \bar{d}^{\lambda c}, D_M d^{\lambda c} \rangle.
\end{aligned}$$

The other two terms in the fluctuated Dirac operator yield more complicated expressions. For the calculation of  $(\gamma^\mu \otimes B_\mu) \tilde{\xi}$ , we use (13.2.1) for the gauge field  $B_\mu$ , and insert the expressions of (13.4.1). As in Proposition 11.7, we then use the antisymmetry of the form  $(J_M \tilde{\chi}, \gamma^\mu \tilde{\psi})$ . For the coupling of the fermions to the gauge fields, a direct calculation then yields

$$\begin{aligned}
\frac{1}{2} \langle J \tilde{\xi}, (\gamma^\mu \otimes B_\mu) \tilde{\xi} \rangle &= \\
s_w g_2 A'_\mu & \left( - \langle J_M \bar{e}^\lambda, \gamma^\mu e^\lambda \rangle + \frac{2}{3} \langle J_M \bar{u}^{\lambda c}, \gamma^\mu u^{\lambda c} \rangle - \frac{1}{3} \langle J_M \bar{d}^{\lambda c}, \gamma^\mu d^{\lambda c} \rangle \right) \\
+ \frac{g_2}{4c_w} Z_\mu & \left( \langle J_M \bar{\nu}^\lambda, \gamma^\mu (1 + \gamma_M) \nu^\lambda \rangle + \langle J_M \bar{e}^\lambda, \gamma^\mu (4s_w^2 - 1 - \gamma_M) e^\lambda \rangle \right. \\
&\quad + \langle J_M \bar{u}^{\lambda c}, \gamma^\mu (-\frac{8}{3}s_w^2 + 1 + \gamma_M) u^{\lambda c} \rangle \\
&\quad \left. + \langle J_M \bar{d}^{\lambda c}, \gamma^\mu (\frac{4}{3}s_w^2 - 1 - \gamma_M) d^{\lambda c} \rangle \right)
\end{aligned}$$

$$\begin{aligned}
& + \frac{g_2}{2\sqrt{2}} W_\mu \left( \langle J_M \bar{e}^\lambda, \gamma^\mu (1 + \gamma_M) \nu^\lambda \rangle + \langle J_M \bar{d}^{\lambda c}, \gamma^\mu (1 + \gamma_M) u^{\lambda c} \rangle \right) \\
& + \frac{g_2}{2\sqrt{2}} W_\mu^* \left( \langle J_M \bar{\nu}^\lambda, \gamma^\mu (1 + \gamma_M) e^\lambda \rangle + \langle J_M \bar{u}^{\lambda c}, \gamma^\mu (1 + \gamma_M) d^{\lambda c} \rangle \right) \\
& + \frac{g_3}{2} G_\mu^i \lambda_i^{dc} \left( \langle J_M \bar{u}^{\lambda d}, \gamma^\mu u^{\lambda c} \rangle + \langle J_M \bar{d}^{\lambda d}, \gamma^\mu d^{\lambda c} \rangle \right),
\end{aligned}$$

where in the weak interactions the projection operator  $\frac{1}{2}(1 + \gamma_M)$  is used to select only the left-handed spinors.

Next, we need to calculate  $\frac{1}{2}(J_M \tilde{\chi}, (\gamma_M \otimes \Phi) \tilde{\xi})$ . The Higgs field is given by  $\Phi = D_F + \phi + J_F \phi J_F^*$ , where  $\phi$  is given by (13.2.2). Let us first focus on the four terms involving only the Yukawa couplings for the neutrinos. Using the symmetry of the form  $(J_M \tilde{\chi}, \gamma_M \tilde{\psi})$ , we obtain

$$\begin{aligned}
& \frac{1}{2} \langle J_M \bar{\nu}_R^\kappa, \gamma_M Y_\nu^{\kappa\lambda} (\phi_1 + 1) \nu_R^\lambda \rangle + \frac{1}{2} \langle J_M \nu_R^\kappa, \gamma_M Y_\nu^{\lambda\kappa} (\phi_1 + 1) \bar{\nu}_R^\lambda \rangle \\
& + \frac{1}{2} \langle J_M \bar{\nu}_L^\kappa, \gamma_M \bar{Y}_\nu^{\lambda\kappa} (\bar{\phi}_1 + 1) \nu_L^\lambda \rangle + \frac{1}{2} \langle J_M \nu_L^\kappa, \gamma_M \bar{Y}_\nu^{\lambda\kappa} (\bar{\phi}_1 + 1) \bar{\nu}_L^\lambda \rangle \\
& = \langle J_M \bar{\nu}_R^\kappa, \gamma_M Y_\nu^{\kappa\lambda} (\phi_1 + 1) \nu_R^\lambda \rangle + \langle J_M \bar{\nu}_L^\kappa, \gamma_M \bar{Y}_\nu^{\lambda\kappa} (\bar{\phi}_1 + 1) \nu_L^\lambda \rangle.
\end{aligned}$$

Using (13.4.2) and (13.4.3), and dropping the generation labels, we can now rewrite

$$\begin{aligned}
& \langle J_M \bar{\nu}_R, \gamma_M Y_\nu (\phi_1 + 1) \nu_R \rangle + \langle J_M \bar{\nu}_L, \gamma_M \bar{Y}_\nu (\bar{\phi}_1 + 1) \nu_L \rangle \\
& = i \left( 1 + \frac{h}{v} \right) \langle J_M \bar{\nu}, m_\nu \nu \rangle - \frac{\phi^0}{v} \langle J_M \bar{\nu}, \gamma_M m_\nu \nu \rangle.
\end{aligned}$$

For  $e, u, d$  we obtain similar terms, the only difference being that for  $e$  and  $d$  the sign for  $\phi^0$  is changed. We also find terms that mix neutrino's and electrons; by the symmetry of the form  $(J_M \tilde{\chi}, \gamma_M \tilde{\psi})$ , these are given by the four terms

$$\begin{aligned}
& \frac{\sqrt{2}}{v} \left( \phi^- \langle J_M \bar{e}_L, m_e \nu_L \rangle + \phi^+ \langle J_M \bar{\nu}_L, m_\nu e_L \rangle \right. \\
& \quad \left. - \phi^- \langle J_M \bar{e}_R, m_\nu \nu_R \rangle - \phi^+ \langle J_M \bar{\nu}_R, m_e e_R \rangle \right).
\end{aligned}$$

There are four similar terms with  $\nu$  and  $e$  replaced by  $u$  and  $d$ , respectively. We can use the projection operators  $\frac{1}{2}(1 \pm \gamma_M)$  to select left- or right-handed spinors. Lastly, the off-diagonal part  $T$  in the finite Dirac operator  $D_F$  yields the Majorana mass terms for the right-handed neutrinos (and left-handed anti-neutrinos). Using (13.4.4), these Majorana mass terms are given by

$$\langle J_M \nu_R, \gamma_M Y_R \nu_R \rangle + \langle J_M \bar{\nu}_L, \gamma_M \bar{Y}_R \bar{\nu}_L \rangle = i \langle J_M \nu_R, m_R \nu_R \rangle + i \langle J_M \bar{\nu}_L, m_R \bar{\nu}_L \rangle.$$

Thus, we find that the mass terms of the fermions and their couplings to the Higgs field are given by

$$\begin{aligned}
& \frac{1}{2} \langle J \tilde{\xi}, (\gamma_M \otimes \Phi) \tilde{\xi} \rangle = \\
& i \left( 1 + \frac{h}{v} \right) \left( \langle J_M \bar{\nu}, m_\nu \nu \rangle + \langle J_M \bar{e}, m_e e \rangle + \langle J_M \bar{u}, m_u u \rangle + \langle J_M \bar{d}, m_d d \rangle \right) \\
& + \frac{\phi^0}{v} \left( \langle J_M \bar{\nu}, \gamma_M m_\nu \nu \rangle - \langle J_M \bar{e}, \gamma_M m_e e \rangle + \langle J_M \bar{u}, \gamma_M m_u u \rangle - \langle J_M \bar{d}, \gamma_M m_d d \rangle \right) \\
& + \frac{\phi^-}{\sqrt{2}v} \left( \langle J_M \bar{e}, m_e (1 + \gamma_M) \nu \rangle - \langle J_M \bar{e}, m_\nu (1 - \gamma_M) \nu \rangle \right) \\
& + \frac{\phi^+}{\sqrt{2}v} \left( \langle J_M \bar{\nu}, m_\nu (1 + \gamma_M) e \rangle - \langle J_M \bar{\nu}, m_e (1 - \gamma_M) e \rangle \right) \\
& + \frac{\phi^-}{\sqrt{2}v} \left( \langle J_M \bar{d}, m_d (1 + \gamma_M) u \rangle - \langle J_M \bar{d}, m_u (1 - \gamma_M) u \rangle \right) \\
& + \frac{\phi^+}{\sqrt{2}v} \left( \langle J_M \bar{u}, m_u (1 + \gamma_M) d \rangle - \langle J_M \bar{u}, m_d (1 - \gamma_M) d \rangle \right) \\
& + i \langle J_M \nu_R, m_R \nu_R \rangle + i \langle J_M \bar{\nu}_L, m_R \bar{\nu}_L \rangle,
\end{aligned}$$

where we have suppressed all indices.  $\square$

In Theorems 13.10 and 13.11 we have calculated the action functional of Definitions 9.1 and 9.3 for the almost-commutative manifold  $M \times F_{SM}$  defined in this Chapter. To summarize, we have geometrically derived:

- (1) The full particle contents of the Standard Model, to wit,
  - the  $W$ ,  $Z$  bosons, photons, and gluons, corresponding to the  $U(1) \times SU(2) \times SU(3)$  Standard Model gauge group.
  - the Higgs boson.
  - three generations of left and right-handed leptons and quarks.
- (2) The dynamics and all interactions of the Standard Model, including
  - self-interactions of the gauge bosons, and coupling to fermions
  - masses for the fermions, including masses for the neutrinos, and coupling to the Higgs field
  - Higgs spontaneous symmetry breaking mechanism, giving masses to the  $W$  and  $Z$  boson, and also to the Higgs boson itself.
- (3) Minimal coupling to gravity.

In addition to the usual Standard Model, there are relations between the coupling constants in the Lagrangian of Theorem 13.10. In the next Chapter, we will analyze this in more detail and derive physical predictions from these relations.

## Notes

- For an exposition of the Standard Model of particle physics, we refer to [1, 2].

### Section 13.1. The Finite Space

- The first description of the finite space  $F_{SM}$  yielding the Standard Model (without right-handed neutrinos though) was given by Connes in [3], based on [4, 5] (see also the review [6]). As already mentioned in the Notes to Chap. 9, the spectral action principle was formulated in [7, 8] where it was also applied to the Standard Model. Extensive computations on this model can be found in [9].

In [10] the noncommutative geometric formulation of the Standard Model got in good shape, mainly because of the choice for the finite space to be of KO-dimension 6 [11, 12]. This solved the problem of fermion doubling pointed out in [13] (see also the discussion in [14, Chap. 1, Sect. 16.3]), and at the same time allowed for the introduction of Majorana masses for right-handed neutrinos, along with the seesaw mechanism. Here, we follow [15].

The derivation of the Standard Model algebra  $A_F$  from the list of finite irreducible geometries of Sect. 3.4 was first obtained in [16]. This includes Proposition 13.1 of which we here give an alternative, diagrammatic proof.

The moduli space of Dirac operators  $D_F$  of the form (13.1.2) was analyzed in [10, Sect. 2.7] (cf. [14, Sect. 1.13.5]) and in [17].

### Section 13.2. The Gauge Theory

- The condition of unimodularity was imposed in the context of the Standard Model in [10, Sect. 2.5] (see also [14, Chap. 1, Sect. 13.3]). The derivation of the hypercharges from the unimodularity condition is closely related to the equivalence between unimodularity in the almost-commutative Standard Model and anomaly cancellation for the usual Standard Model [18].
- Proposition 13.4 agrees with [10, Proposition 2.16] (see also [14, Proposition 1.185]). For the derivation of the Standard Model gauge group  $\mathfrak{G}_{SM}$ , we refer to [19].

### Section 13.3. The Spectral Action

- The coefficients  $a, b, c, d$  and  $e$  in Lemma 13.7 agree with those appearing in [10] (see also [14, Chap. 1, Sect. 15.2]).
- The Higgs mechanism is attributed to Englert, Brout and Higgs [20, 21].
- The form of the Higgs field in (13.3.10) that is obtained after a suitable change of basis is called *unitary gauge* and was introduced by Weinberg in [22, 23] (see also [24, Chap. 21]).

## References

- Cottingham, W.N., Greenwood, D.A.: An Introduction to the Standard Model of Particle Physics, 2nd edn. Cambridge University Press, Cambridge (2007)
- Kane, G.L.: Modern Elementary Particle Physics. Perseus (1993)
- Connes, A.: Gravity coupled with matter and the foundation of non-commutative geometry. Commun. Math. Phys. **182**, 155–176 (1996)
- Connes, A.: Essay on physics and noncommutative geometry. In: The Interface of Mathematics and Particle Physics (Oxford, 1988), volume 24 of Institute of Mathematics and its Applications Conference Series, pp. 9–48. Oxford University Press, New York (1990)

5. Connes, A., Lott, J.: Particle models and noncommutative geometry. *Nucl. Phys. Proc. Suppl.* **18B**, 29–47 (1991)
6. Martín, C.P., Gracia-Bondía, J.M., Várilly, J.C.: The standard model as a noncommutative geometry: the low-energy regime. *Phys. Rep.* **294**, 363–406 (1998)
7. Chamseddine, A.H., Connes, A.: Universal formula for noncommutative geometry actions: unifications of gravity and the Standard Model. *Phys. Rev. Lett.* **77**, 4868–4871 (1996)
8. Chamseddine, A.H., Connes, A.: The spectral action principle. *Commun. Math. Phys.* **186**, 731–750 (1997)
9. Scheck, F., Upmeier, H., Werner, W. (eds.): *Noncommutative Geometry and the Standard Model of Elementary Particle Physics*, volume 596 of *Lecture Notes in Physics*. Springer, Berlin (2002). Papers from the conference held in Hesselberg, 14–19 Mar. 1999
10. Chamseddine, A.H., Connes, A., Marcolli, M.: Gravity and the standard model with neutrino mixing. *Adv. Theor. Math. Phys.* **11**, 991–1089 (2007)
11. Barrett, J.W.: A Lorentzian version of the non-commutative geometry of the standard model of particle physics. *J. Math. Phys.* **48**, 012303 (2007)
12. Connes, A.: Noncommutative geometry and the standard model with neutrino mixing. *JHEP* **0611**, 081 (2006)
13. Lizzi, F., Mangano, G., Miele, G., Sparano, G.: Fermion Hilbert space and fermion doubling in the noncommutative geometry approach to gauge theories. *Phys. Rev. D* **55**, 6357–6366 (1997)
14. Connes, A., Marcolli, M.: *Noncommutative Geometry. Quantum Fields and Motives*, AMS Providence (2008)
15. van den Dungen, K., van Suijlekom, W.D.: Particle physics from almost commutative space-times. *Rev. Math. Phys.* **24**, 1230004 (2012)
16. Chamseddine, A.H., Connes, A.: Why the standard model. *J. Geom. Phys.* **58**, 38–47 (2008)
17. Čaćić, B.: Moduli spaces of Dirac operators for finite spectral triples. *Quantum Groups and Noncommutative Spaces: Perspectives on Quantum Geometry* (Marcolli, M., Parashar, D. (eds.)), Vieweg Verlag (2010)
18. Alvarez, E., Gracia-Bondía, J.M., Martin, C.: Anomaly cancellation and the gauge group of the standard model in NCG. *Phys. Lett. B* **364**, 33–40 (1995)
19. Baez, J., Huerta, J.: The algebra of grand unified theories. *Bull. Am. Math. Soc. (N.S.)* **47**, 483–552 (2010)
20. Englert, F., Brout, R.: Broken symmetry and the mass of gauge vector mesons. *Phys. Rev. Lett.* **13**, 321–323 (1964)
21. Higgs, P.W.: Broken symmetries and the masses of gauge bosons. *Phys. Rev. Lett.* **13**, 508–509 (1964)
22. Weinberg, S.: Physical processes in a convergent theory of the weak and electromagnetic interactions. *Phys. Rev. Lett.* **27**, 1688–1691 (1971)
23. Weinberg, S.: General theory of broken local symmetries. *Phys. Rev. D* **7**, 1068–1082 (1973)
24. Weinberg, S.: *The Quantum Theory of Fields*, vol. II. Cambridge University Press (1996)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 14

## Phenomenology of the Noncommutative Standard Model



In Theorems 13.10 and 13.11, we have derived the full Lagrangian for the Standard Model from the almost-commutative manifold  $M \times F_{SM}$ . The coefficients in this Lagrangian are given in terms of:

- the value  $f(0)$  and the moments  $f_2$  and  $f_4$  of the function  $f$  in the spectral action;
- the cut-off scale  $\Lambda$  in the spectral action;
- the vacuum expectation value  $v$  of the Higgs field;
- the coefficients  $a, b, c, d, e$  of (13.3.2) that are determined by the mass matrices in the finite Dirac operator  $D_F$ .

One can find several relations among these coefficients in the Lagrangian, which we shall derive in the following section. Inspired by the relation  $g_3^2 = g_2^2 = \frac{5}{3}g_1^2$  obtained from (13.3.4), we will assume that these relations hold at the unification scale. Subsequently, we use the renormalization group equations to obtain predictions for the Standard Model at ‘lower’ (i.e. particle accelerator) energies.

### 14.1 Mass Relations

#### 14.1.1 Fermion Masses

Recall from (13.4.3) that we defined the mass matrices  $m_x$  of the fermions by rewriting the matrices  $Y_x$  in the finite Dirac operator  $D_F$ . Inserting the formula (13.4.3) for  $Y_x$  into the expression for  $a$  given by (13.3.2), we obtain

$$a = \frac{af(0)}{\pi^2 v^2} \text{Tr} (m_v^* m_v + m_e^* m_e + 3m_u^* m_u + 3m_d^* m_d),$$

which yields

$$\mathrm{Tr} (m_v^* m_v + m_e^* m_e + 3m_u^* m_u + 3m_d^* m_d) = \frac{\pi^2 v^2}{f(0)}.$$

From (13.3.14) we know that the mass of the  $W$ -boson is given by  $M_W = \frac{1}{2}v g_2$ . Using the normalization (13.3.4), expressing  $g_2$  in terms of  $f(0)$ , we can then write

$$f(0) = \frac{\pi^2 v^2}{8M_W^2}. \quad (14.1.1)$$

Inserting this into the expression above, we obtain a relation between the fermion mass matrices  $m_x$  and the  $W$ -boson mass  $M_W$ , *viz.*

$$\mathrm{Tr} (m_v^* m_v + m_e^* m_e + 3m_u^* m_u + 3m_d^* m_d) = 2g_2^2 v^2 = 8M_W^2. \quad (14.1.2)$$

If we assume that the mass of the top quark is much larger than all other fermion masses, we may neglect the other fermion masses. In that case, the above relation would yield the constraint

$$m_{\text{top}} \lesssim \sqrt{\frac{8}{3}} M_W. \quad (14.1.3)$$

### 14.1.2 The Higgs Mass

We obtain a mass  $m_h$  for the Higgs boson  $h$  by writing the term proportional to  $h^2$  in (13.3.11) in the form

$$\frac{b\pi^2}{2a^2 f(0)} 4v^2 h^2 = \frac{1}{2} m_h^2 h^2.$$

Thus, the Higgs mass is given by

$$m_h = \frac{2\pi\sqrt{b}v}{a\sqrt{f(0)}}. \quad (14.1.4)$$

Inserting (14.1.1) into this expression for the Higgs mass, we see that  $M_W$  and  $m_h$  are related by

$$m_h^2 = 32 \frac{b}{a^2} M_W^2.$$

Next, we introduce the quartic Higgs coupling constant  $\lambda$  by writing

$$\frac{b\pi^2}{2a^2 f(0)} h^4 =: \frac{1}{24} \lambda h^4.$$

From (13.3.4) we then find

$$\lambda = 24 \frac{b}{a^2} g_2^2, \quad (14.1.5)$$

so that the (tree-level) Higgs mass can be expressed in terms of the mass  $M_W$  of the  $W$ -boson, the coupling constant  $g_2$  and the quartic Higgs coupling  $\lambda$  as

$$m_h^2 = \frac{4\lambda M_W^2}{3g_2^2}. \quad (14.1.6)$$

### 14.1.3 The Seesaw Mechanism

Let us consider the mass terms for the neutrinos. The matrix  $D_F$  described in Sect. 13.1 provides the Dirac masses as well as the Majorana masses of the fermions. After a rescaling as in (13.4.3), the mass matrix restricted to the subspace of  $H_F$  with basis  $\{\nu_L, \nu_R, \overline{\nu}_L, \overline{\nu}_R\}$  is given by

$$\begin{pmatrix} 0 & m_\nu^* & m_R^* & 0 \\ m_\nu & 0 & 0 & 0 \\ m_R & 0 & 0 & \overline{m}_\nu^* \\ 0 & 0 & \overline{m}_\nu & 0 \end{pmatrix}.$$

Suppose we consider only one generation, so that  $m_\nu$  and  $m_R$  are just scalars. The eigenvalues of the above mass matrix are then given by

$$\pm \frac{1}{2} m_R \pm \frac{1}{2} \sqrt{m_R^2 + 4m_\nu^2}.$$

If we assume that  $m_\nu \ll m_R$ , then these eigenvalues are approximated by  $\pm m_R$  and  $\pm \frac{m_\nu^2}{m_R}$ . This means that there is a heavy neutrino, for which the Dirac mass  $m_\nu$  may be neglected, so that its mass is given by the Majorana mass  $m_R$ . However, there is also a light neutrino, for which the Dirac and Majorana terms conspire to yield a mass  $\frac{m_\nu^2}{m_R}$ , which is in fact much smaller than the Dirac mass  $m_\nu$ . This is called the *seesaw mechanism*. Thus, even though the observed masses for these neutrinos may be very small, they might still have large Dirac masses (or Yukawa couplings).

From (14.1.2) we obtained a relation between the masses of the top quark and the  $W$ -boson by neglecting all other fermion masses. However, because of the seesaw mechanism it might be that one of the neutrinos has a Dirac mass of the same order

of magnitude as the top quark. In that case, it would not be justified to neglect all other fermion masses, but instead we need to correct for such massive neutrinos.

Let us introduce a new parameter  $\rho$  (typically taken to be of order 1) for the ratio between the Dirac mass  $m_\nu$  for the tau-neutrino and the mass  $m_{\text{top}}$  of the top quark at unification scale, so we write  $m_\nu = \rho m_{\text{top}}$ . Instead of (14.1.3), we then obtain the restriction

$$m_{\text{top}} \lesssim \sqrt{\frac{8}{3 + \rho^2}} M_W. \quad (14.1.7)$$

## 14.2 Renormalization Group Flow

In this section we evaluate the renormalization group equations (RGEs) for the Standard Model from ordinary energies up to the unification scale. For the validity of these RGEs we need to assume the existence of a ‘big desert’ up to the grand unification scale. This means that one assumes that:

- there exist no new particles (besides the known Standard Model particles) with a mass below the unification scale;
- perturbative quantum field theory remains valid throughout the big desert.

Furthermore, we also ignore any gravitational contributions to the renormalization group flow.

### 14.2.1 Coupling Constants

In (13.3.1) we introduced the coupling constants for the gauge fields, and we obtained the relation  $g_3^2 = g_2^2 = \frac{5}{3}g_1^2$ . This is precisely the relation between the coupling constants at (grand) unification, common to grand unified theories (GUT). Thus, it would be natural to assume that our model is defined at the scale  $\Lambda_{\text{GUT}}$ . However, it turns out that there is no scale at which the relation  $g_3^2 = g_2^2 = \frac{5}{3}g_1^2$  holds exactly, as we show below.

The renormalization group  $\beta$ -functions of the (minimal) standard model read

$$\frac{dg_i}{dt} = -\frac{1}{16\pi^2} b_i g_i^3; \quad (b_1, b_2, b_3) = \left(-\frac{41}{6}, \frac{19}{6}, 7\right),$$

where  $t = \log \mu$ . At first order, these equations are uncoupled from all other parameters of the Standard Model, and the solutions for the running coupling constants  $g_i(\mu)$  at the energy scale  $\mu$  are easily seen to satisfy

$$g_i(\mu)^{-2} = g_i(M_Z)^{-2} + \frac{b_i}{8\pi^2} \log \frac{\mu}{M_Z}, \quad (14.2.1)$$

where  $M_Z$  is the experimental mass of the Z-boson:

$$M_Z = 91.1876 \pm 0.0021 \text{ GeV.}$$

For later convenience, we also recall that the experimental mass of the W-boson is

$$M_W = 80.399 \pm 0.023 \text{ GeV.} \quad (14.2.2)$$


The experimental values of the coupling constants at the energy scale  $M_Z$  are known too, and are given by

$$g_1(M_Z) = 0.3575 \pm 0.0001, \quad (14.2.3)$$

$$g_2(M_Z) = 0.6519 \pm 0.0002, \quad (14.2.4)$$

$$g_3(M_Z) = 1.220 \pm 0.004. \quad (14.2.5)$$

Using these experimental values, we obtain the running of the coupling constants in Fig. 14.1. As can be seen in this figure, the running coupling constants do not meet at any single point, and hence they do not determine a unique unification scale  $\Lambda_{GUT}$ . In other words, the relation  $g_3^2 = g_2^2 = \frac{5}{3}g_1^2$  cannot hold exactly at any energy scale, unless we drop the big desert hypothesis. Nevertheless, in the remainder of this section we assume that this relation holds at least approximately and we will come back to this point in the next section. We consider the range for  $\Lambda_{GUT}$  determined



**Fig. 14.1** The running of the gauge coupling constants

by the triangle of the running coupling constants in Fig. 14.1. The scale  $\Lambda_{12}$  at the intersection of  $\sqrt{\frac{5}{3}}g_1$  and  $g_2$  determines the lowest value for  $\Lambda_{GUT}$ , given by

$$\Lambda_{12} = M_Z \exp \left( \frac{8\pi^2(\frac{3}{5}g_1(M_Z)^{-2} - g_2(M_Z)^{-2})}{b_2 - \frac{3}{5}b_1} \right) = 1.03 \times 10^{13} \text{ GeV}. \quad (14.2.6)$$

The highest value  $\Lambda_{23}$  is given by the solution of  $g_2 = g_3$ , which yields

$$\Lambda_{23} = M_Z \exp \left( \frac{8\pi^2(g_3(M_Z)^{-2} - g_2(M_Z)^{-2})}{b_2 - b_3} \right) = 9.92 \times 10^{16} \text{ GeV}. \quad (14.2.7)$$

We assume that the Lagrangian we have derived from the almost-commutative manifold  $M \times F_{SM}$  is valid at some scale  $\Lambda_{GUT}$ , which we take to be between  $\Lambda_{12}$  and  $\Lambda_{23}$ . All relations obtained in Fig. 14.1 are assumed to hold approximately at this scale, and all predictions that will follow from these relations are therefore also only approximate.

### 14.2.2 Renormalization Group Equations

The running of the neutrino masses has been studied in a general setting for non-degenerate seesaw scales. In what follows we consider the case where only the tau-neutrino has a large Dirac mass  $m_\nu$ , which cannot be neglected with respect to the mass of the top-quark. In the remainder of this section we calculate the running of the Yukawa couplings for the top-quark and the tau-neutrino, as well as the running of the quartic Higgs coupling. Let us write  $y_{top}$  and  $y_\nu$  for the Yukawa couplings of the top quark and the tau-neutrino, defined by

$$m_{top} = \frac{1}{2}\sqrt{2}y_{top}v, \quad m_\nu = \frac{1}{2}\sqrt{2}y_\nu v, \quad (14.2.8)$$

where  $v$  is the vacuum expectation value of the Higgs field.

Let  $m_R$  be the Majorana mass for the right-handed tau-neutrino. By the Appelquist–Carazzone decoupling theorem (cf. Note 5 on Sect. 14.2.4) we can distinguish two energy domains:  $\mu > m_R$  and  $\mu < m_R$ . We again neglect all fermion masses except for the top quark and the tau neutrino. For high energies  $\mu > m_R$ , the renormalization group equations are given by

$$\begin{aligned}\frac{dy_{\text{top}}}{dt} &= \frac{1}{16\pi^2} \left( \frac{9}{2}y_{\text{top}}^2 + y_v^2 - \frac{17}{12}g_1^2 - \frac{9}{4}g_2^2 - 8g_3^2 \right) y_{\text{top}}, \\ \frac{dy_v}{dt} &= \frac{1}{16\pi^2} \left( 3y_{\text{top}}^2 + \frac{5}{2}y_v^2 - \frac{3}{4}g_1^2 - \frac{9}{4}g_2^2 \right) y_v, \\ \frac{d\lambda}{dt} &= \frac{1}{16\pi^2} \left( 4\lambda^2 - (3g_1^2 + 9g_2^2)\lambda + \frac{9}{4}(g_1^4 + 2g_1^2g_2^2 + 3g_2^4) \right. \\ &\quad \left. + 4(3y_{\text{top}}^2 + y_v^2)\lambda - 12(3y_{\text{top}}^4 + y_v^4) \right).\end{aligned}\quad (14.2.9)$$

Below the threshold  $\mu = m_R$ , the Yukawa coupling of the tau-neutrino drops out of the RG equations and is replaced by an effective coupling

$$\kappa = 2 \frac{y_v^2}{m_R},$$

which provides an effective mass  $m_l = \frac{1}{4}\kappa v^2$  for the light tau-neutrino. The renormalization group equations of  $y_{\text{top}}$  and  $\lambda$  for  $\mu < m_R$  are then given by

$$\begin{aligned}\frac{dy_{\text{top}}}{dt} &= \frac{1}{16\pi^2} \left( \frac{9}{2}y_{\text{top}}^2 - \frac{17}{12}g_1^2 - \frac{9}{4}g_2^2 - 8g_3^2 \right) y_{\text{top}}, \\ \frac{d\lambda}{dt} &= \frac{1}{16\pi^2} \left( 4\lambda^2 - (3g_1^2 + 9g_2^2)\lambda + \frac{9}{4}(g_1^4 + 2g_1^2g_2^2 + 3g_2^4) \right. \\ &\quad \left. + 12y_{\text{top}}^2\lambda - 36y_{\text{top}}^4 \right).\end{aligned}\quad (14.2.10)$$

Finally, the equation for  $y_v$  is replaced by an equation for the effective coupling  $\kappa$  given by

$$\frac{d\kappa}{dt} = \frac{1}{16\pi^2} \left( 6y_{\text{top}}^2 - 3g_2^2 + \frac{\lambda}{6} \right) \kappa.\quad (14.2.11)$$

### 14.2.3 Running Masses

The numerical solutions to the coupled differential equations of (14.2.9) (14.2.10) and (14.2.11) for  $y_{\text{top}}$ ,  $y_v$  and  $\lambda$  depend on the choice of three input parameters:

- the scale  $\Lambda_{GUT}$  at which our model is defined;
- the ratio  $\rho$  between the masses  $m_v$  and  $m_{\text{top}}$ ;
- the Majorana mass  $m_R$  that produces the threshold in the renormalization group flow.

The scale  $\Lambda_{GUT}$  is taken to be either  $\Lambda_{12} = 1.03 \times 10^{13}$  GeV or  $\Lambda_{23} = 9.92 \times 10^{16}$  GeV, as given by (14.2.6) and (14.2.7), respectively. We now deter-

mine the numerical solution to (14.2.9), (14.2.10) and (14.2.11) for a range of values for  $\rho$  and  $m_R$ . First, we need to start with the initial conditions of the running parameters at the scale  $\Lambda_{GUT}$ . Inserting the top-quark mass  $m_{\text{top}} = \frac{1}{2}\sqrt{2}y_{\text{top}}v$ , the tau-neutrino mass  $m_\nu = \rho m_{\text{top}}$ , and the  $W$ -boson mass  $M_W = \frac{1}{2}g_2 v$  into (14.1.7), we obtain the constraints

$$y_{\text{top}}(\Lambda_{GUT}) \lesssim \frac{2}{\sqrt{3 + \rho^2}} g_2(\Lambda_{GUT}), \quad y_\nu(\Lambda_{GUT}) \lesssim \frac{2\rho}{\sqrt{3 + \rho^2}} g_2(\Lambda_{GUT}),$$

where (14.2.1) yields the values  $g_2(\Lambda_{12}) = 0.5444$  and  $g_2(\Lambda_{23}) = 0.5170$ .

Furthermore, from (14.1.5) we obtain an expression for the quartic coupling  $\lambda$  at  $\Lambda_{GUT}$ . Approximating the coefficients  $a$  and  $b$  from (13.3.2) by  $a \approx (3 + \rho^2)m_{\text{top}}^2$  and  $b \approx (3 + \rho^4)m_{\text{top}}^4$ , we obtain the boundary condition

$$\lambda(\Lambda_{GUT}) \approx 24 \frac{3 + \rho^4}{(3 + \rho^2)^2} g_2(\Lambda_{GUT})^2.$$

Using these boundary conditions, we can now numerically solve the RG equations of (14.2.9) from  $\Lambda_{GUT}$  down to  $m_R$ , which provides us with values for  $y_{\text{top}}(m_R)$ ,  $y_\nu(m_R)$  and  $\lambda(m_R)$ . At this point, the Yukawa coupling  $y_\nu$  is replaced by the effective coupling  $\kappa$  with boundary condition

$$\kappa(m_R) = 2 \frac{y_\nu(m_R)^2}{m_R}.$$

Next, we numerically solve the RG equations of (14.2.10) and (14.2.11) down to  $M_Z$  to obtain the values for  $y_{\text{top}}$ ,  $\kappa$  and  $\lambda$  at ‘low’ energy scales.

The running mass of the top quark at these energies is given by (14.2.8). We find the running Higgs mass by inserting  $\lambda$  into (14.1.6). We shall evaluate these running masses at their own energy scale. For instance, our predicted mass for the Higgs boson is the solution for  $\mu$  of the equation  $\mu = \sqrt{\lambda(\mu)/3}v$ , in which we ignore the running of the vacuum expectation value  $v$ .

The effective mass of the light neutrino is determined by the effective coupling  $\kappa$ , and we choose to evaluate this mass at scale  $M_Z$ . Thus, we calculate the masses by

$$\begin{aligned} m_{\text{top}}(m_{\text{top}}) &= \frac{1}{2}\sqrt{2}y_{\text{top}}(m_{\text{top}})v, \\ m_l(M_Z) &= \frac{1}{4}\kappa(M_Z)v^2, \\ m_h(m_h) &= \sqrt{\frac{\lambda(m_h)}{3}}v, \end{aligned}$$

where, from the  $W$ -boson mass (14.2.2) we can insert the value  $v = 246.66 \pm 0.15$ . The results of this procedure for  $m_{\text{top}}$ ,  $m_l$  and  $m_h$  are given in Table 14.1. In this table,

**Table 14.1** Numerical results for the masses  $m_{\text{top}}$  of the top-quark,  $m_l$  of the light tau-neutrino, and  $m_h$  of the Higgs boson, as a function of  $\Lambda_{\text{GUT}}$ ,  $\rho$ , and  $m_R$

$\Lambda_{\text{GUT}}$ ( $10^{13}$ GeV)	1.03	1.03	1.03	1.03	1.03	1.03	1.03
$\rho$	0	0.90	0.90	1.00	1.00	1.10	1.10
$m_R$ ( $10^{13}$ GeV)	–	0.25	1.03	0.30	1.03	0.35	1.03
$m_{\text{top}}$ (GeV)	183.2	173.9	174.1	171.9	172.1	169.9	170.1
$m_l$ (eV)	0	2.084	0.5037	2.076	0.6030	2.080	0.7058
$m_h$ (GeV)	188.3	175.5	175.7	173.4	173.7	171.5	171.8
$\Lambda_{\text{GUT}}$ ( $10^{16}$ GeV)	9.92	9.92	9.92	9.92	9.92		
$\rho$	0	1.10	1.10	1.20	1.20		
$m_R$ ( $10^{13}$ GeV)	–	0.30	2.0	0.35	9900		
$m_{\text{top}}$ (GeV)	186.0	173.9	174.2	171.9	173.5		
$m_l$ (eV)	0	1.939	0.2917	1.897	$6.889 \times 10^{-5}$		
$m_h$ (GeV)	188.1	171.3	171.6	169.1	171.2		
$\Lambda_{\text{GUT}}$ ( $10^{16}$ GeV)	9.92	9.92	9.92	9.92			
$\rho$	1.30	1.30	1.35	1.35			
$m_R$ ( $10^{13}$ GeV)	0.40	9900	100	9900			
$m_{\text{top}}$ (GeV)	169.9	171.6	169.8	170.6			
$m_l$ (eV)	1.866	$7.818 \times 10^{-5}$	$8.056 \times 10^{-3}$	$8.286 \times 10^{-5}$			
$m_h$ (GeV)	167.1	169.3	167.4	168.4			

we have chosen the range of values for  $\rho$  and  $m_R$  such that the mass of the top-quark and the light tau-neutrino are in agreement with their experimental values


$$m_{\text{top}} = 172.0 \pm 0.9 \pm 1.3 \text{ GeV}, \quad m_l \leq 2 \text{ eV}.$$

For comparison, we have also included the simple case where we ignore the Yukawa coupling of the tau-neutrino (by setting  $\rho = 0$ ), in which case there is no threshold at the Majorana mass scale either. As an example, we have plotted the running of  $\lambda$ ,  $y_{\text{top}}$ ,  $y_\nu$  and  $\kappa$  for the values of  $\Lambda_{\text{GUT}} = \Lambda_{23} = 9.92 \times 10^{16}$  GeV,  $\rho = 1.2$ , and  $m_R = 3 \times 10^{12}$  GeV in Figs. 14.2, 14.3, 14.4 and 14.5.


For the allowed range of values for  $\rho$  and  $m_R$  that yield plausible results for  $m_{\text{top}}$  and  $m_l$ , we see that the mass  $m_h$  of the Higgs boson takes its value within the range

$$167 \text{ GeV} \leq m_h \leq 176 \text{ GeV}.$$


The errors in this prediction, which result from the initial conditions (other than  $m_{\text{top}}$  and  $m_l$ ) taken from experiment, as well as from ignoring higher-loop corrections to the RGEs, are smaller than this range of possible values for the Higgs mass, and therefore we may ignore these errors.




**Fig. 14.2** The running of the quartic Higgs coupling  $\lambda$  for  $\Lambda_{GUT} = 9.92 \times 10^{16}$  GeV,  $\rho = 1.2$ , and  $m_R = 3 \times 10^{12}$  GeV



**Fig. 14.3** The running of the top-quark Yukawa coupling  $y_{\text{top}}$  for  $\Lambda_{GUT} = 9.92 \times 10^{16}$  GeV,  $\rho = 1.2$ , and  $m_R = 3 \times 10^{12}$  GeV



**Fig. 14.4** The running of the tau-neutrino Yukawa coupling  $y_\nu$  for  $\Lambda_{GUT} = 9.92 \times 10^{16}$  GeV,  $\rho = 1.2$ , and  $m_R = 3 \times 10^{12}$  GeV




**Fig. 14.5** The running of the effective coupling  $\kappa$  for  $\Lambda_{GUT} = 9.92 \times 10^{16}$  GeV,  $\rho = 1.2$ , and  $m_R = 3 \times 10^{12}$  GeV

#### 14.2.4 Higgs Mass: Comparison to Experimental Results

Since the discovery of the Higgs boson at the ATLAS and CMS experiments at the Large Hadron Collider at CERN in 2012 we also know with increasing accuracy that

**Fig. 14.6** Observed and expected exclusion limits for a Standard Model Higgs boson at the 95-percent confidence level for the combined CDF and DZero analyses. (Fermilab)



the experimental value for the Higgs mass is around  $m_h \simeq 125.5$  GeV. Strictly speaking, this is a falsification of the noncommutative Standard Model since it evidently lies outside of the above predicted range. In fact, ironically enough, the above range was among the first to be excluded by Fermilab's D0 experiment (see Fig. 14.6).

As usual, in the derivation of the model and the renormalization group equation several assumptions were made and simplifications were applied, so that it is important to look back at them. And indeed, lifting the curtain slightly for what is to come, in the next Chapter we will see that the reduction in Proposition 13.1 from the irreducible finite geometry of KO-dimension 6 based on  $M_2(\mathbb{H}) \oplus M_4(\mathbb{C})$  to the Standard Model based on  $\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$  may not be necessary nor desired. As we will see, the irreducible geometry describes a Pati–Salam gauge model that goes Beyond the Standard Model and allows for a Higgs mass that is compatible with the observed value. Moreover, this will solve the incompatibility between the grand unification of the gauge couplings suggested by the spectral model in Eq.(13.3.5) and the existence of a GUT-triangle in Fig. 14.2.1.

## Notes

- In the first part of this Chapter, we mainly follow [1, Sect. 5] (see also [2, Chap. 1, Sect. 17]). In Sect. 14.2 we have also incorporated the running of the neutrino masses as in [3] (see also [4]).

### Section 14.1. Mass Relations

- Further details on the see-saw mechanism can be found in e.g. [5].

### Section 14.2. Renormalization Group Flow

- The renormalization group  $\beta$ -functions of the (minimal) standard model are taken from [6–9]. We simplify the expressions by ignoring the 2-loop contributions, and instead consider only the 1-loop approximation. The renormalization group  $\beta$ -functions are [6, Eq. (B.2)] or [9, Eq. (A.1)].

### Section 14.2.1. Coupling Constants

4. The experimental masses of the  $Z$  and  $W$ -boson and the top quark, as well as the experimental values of the coupling constants at the energy scale  $M_Z$  are found in [10].
5. In arriving at (14.2.9) we have followed the approach of [3] where two energy domains are considered:  $\mu > m_R$  and  $\mu < m_R$ . The Appelquist–Carazzone decoupling theorem is found in [11]. For the renormalization group equations, we refer to [7, Eq. (B.4)], [12, Eq. (14) and (15)] and [8, Eq. (B.3)].
6. The discovery of the Higgs boson at the ATLAS and CMS experiments is published in [13, 14].
7. In this chapter and Chap. 15 we exploit renormalization group techniques to run couplings and masses down from the GUT-scale to ordinary energies. The renormalization group equations were derived in a perturbative approach to quantum field theory, which was supposed to be valid at all scales. Moreover, we have adopted the one-loop beta-functions, something which can definitely be improved. Even though this might lead to more accurate predictions, it is not expected to resolve the incompatibility between the predicted range for  $m_h$  and the experimentally measured value. In our analysis we have discarded all possible gravitational effects on the running of the coupling constants. It might very well be that gravitational correction terms alter the predicted values to a more realistic value. A possible approach to incorporate gravitational effects in the running of the coupling constants is discussed in [15].

## References

1. Chamseddine, A.H., Connes, A., Marcolli, M.: Gravity and the Standard Model with neutrino mixing. *Adv. Theor. Math. Phys.* **11**, 991–1089 (2007)
2. Connes, A., Marcolli, M.: Noncommutative Geometry, Quantum Fields and Motives. AMS, Providence (2008)
3. Jureit, J., Krajewski, T., Schücker, T., Stephan, C.A.: Seesaw and noncommutative geometry. *Phys. Lett. B* **654**, 127–132 (2007)
4. van den Dungen, K., van Suijlekom, W.D.: Particle physics from almost commutative space-times. *Rev. Math. Phys.* **24**, 1230004 (2012)
5. Mohapatra, R., Pal, P.: Massive Neutrinos in Physics and Astrophysics, 2nd ed. World Scientific Lecture Notes in Physics, vol. 60, pp. 1–397 (1998)
6. Machacek, M., Vaughn, M.: Two-loop renormalization group equations in a general quantum field theory: (i). Wave function renormalization. *Nucl. Phys. B* **222**, 83–103 (1983)
7. Machacek, M., Vaughn, M.: Two-loop renormalization group equations in a general quantum field theory (ii). Yukawa couplings. *Nucl. Phys. B* **236**, 221–232 (1984)
8. Machacek, M., Vaughn, M.: Two-loop renormalization group equations in a general quantum field theory?: (iii). Scalar quartic couplings. *Nucl. Phys. B* **249**, 70–92 (1985)
9. Ford, C., Jones, D.R.T., Stephenson, P.W., Einhorn, M.B.: The effective potential and the renormalisation group. *Nucl. Phys. B* **395**, 17–34 (1993)
10. Nakamura, K., et al.: Review of particle physics. *J. Phys. G: Nucl. Part. Phys.* **37**, 075021 (2010)
11. Appelquist, T., Carazzone, J.: Infrared singularities and massive fields. *Phys. Rev. D* **11**, 2856–2861 (1975)
12. Antusch, S., Kersten, J., Lindner, M., Ratz, M.: Neutrino mass matrix running for non-degenerate see-saw scales. *Phys. Lett. B* **538**, 87–95 (2002)
13. Aad, G., et al.: Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett. B* **716**, 1–29 (2012)

14. Chatrchyan, S., et al.: Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett. B* **716**, 30–61 (2012)
15. Estrada, C., Marcolli, M.: Asymptotic safety, hypergeometric functions, and the Higgs mass in spectral action models. *Int. J. Geom. Meth. Mod. Phys.* **10**, 1350036 (2013)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 15

## Beyond the Standard Model: Pati–Salam Unification



One of the pressing questions at this point is whether noncommutative geometry may point to new physics beyond the Standard Model. The success of the spectral construction of the Standard Model, predicting its particle content, including gauge fields, Higgs fields as well as a singlet whose vev gives Majorana mass to the right handed neutrino, is a strong signal that we are on the right track. However, the mismatch between the predicted range of the Higgs mass and the experimentally observed value suggests that we should reconsider the path we took.

### 15.1 The Finite Noncommutative Space of the Pati–Salam Model

Recall from Sect. 13.1 that the route that led to the above conclusion starts with the simplest irreducible geometry that allows for a symplectic constraint (condition 1 on Page 1). It was based on the algebra

$$A := M_2(\mathbb{H}) \oplus M_4(\mathbb{C}). \quad (15.1.1)$$

The existence of the chirality operator  $\gamma$  that commutes with the algebra breaks the quaternionic matrices  $M_2(\mathbb{H})$  to the diagonal subalgebra and leads us to consider the finite algebra

$$A_{PS} := \mathbb{H}_R \oplus \mathbb{H}_L \oplus M_4(\mathbb{C}). \quad (15.1.2)$$

In view of this structure, a convenient tensorial representation of our Hilbert space vectors  $\Psi$  in  $H_F$  (cf. Eq. (13.1.1)) is given by

$$\Psi = \begin{pmatrix} \psi_A \\ \psi_{A'} \end{pmatrix}, \quad \psi_{A'} = \psi_A^c \quad (15.1.3)$$

where  $\psi_A^c$  is the conjugate vector to  $\psi_A$ . Thus all primed indices  $A'$  correspond to the Hilbert space of conjugate vectors. It is acted on by both the left algebra  $M_2(\mathbb{H})$  and the right algebra  $M_4(\mathbb{C})$ . Therefore the index  $A$  can take 16 values and is represented by

$$A = \alpha I \quad (15.1.4)$$

where the index  $\alpha$  is acted on by quaternionic matrices and the index  $I$  by  $M_4(\mathbb{C})$  matrices. Moreover, when the grading breaks  $M_2(\mathbb{H})$  into  $\mathbb{H}_R \oplus \mathbb{H}_L$  the index  $\alpha$  is decomposed to  $\alpha = \dot{a}, a$  where  $\dot{a} = \dot{1}, \dot{2}$  (dotted index) is acted on by the first quaternionic algebra  $\mathbb{H}_R$  and  $a = 1, 2$  is acted on by the second quaternionic algebra  $\mathbb{H}_L$ . When  $M_4(\mathbb{C})$  breaks into  $\mathbb{C} \oplus M_3(\mathbb{C})$  (due to symmetry breaking or through the use of the order one condition as in Proposition 13.1) the index  $I$  is decomposed into  $I = 1, i$  and thus distinguishing leptons and quarks, where the 1 is acted on by the  $\mathbb{C}$  and the  $i$  by  $M_3(\mathbb{C})$ . Therefore the various components of the spinor  $\psi_A$  are

$$\begin{aligned} \psi_{\alpha I} &= \begin{pmatrix} v_R & u_{iR} & v_L & u_{iL} \\ e_R & d_{iR} & e_L & d_{iL} \end{pmatrix}, \quad i = 1, 2, 3 \\ &= (\psi_{\dot{a}1}, \psi_{\dot{a}i}, \psi_{a1}, \psi_{ai}), \quad a = 1, 2, \quad \dot{a} = \dot{1}, \dot{2} \end{aligned} \quad (15.1.5)$$

The (finite) Dirac operator can be written in matrix form

$$D_F = \begin{pmatrix} D_A^B & D_{A'}^{B'} \\ D_{A'}^B & D_{A'}^{B'} \end{pmatrix}, \quad (15.1.6)$$

and must satisfy the properties

$$\gamma_F D_F = -D_F \gamma_F \quad J_F D_F = D_F J_F \quad (15.1.7)$$

where  $J_F^2 = 1$ . A matrix realization of  $\gamma_F$  and  $J_F$  are given by

$$\gamma_F = \begin{pmatrix} G_F & 0 \\ 0 & -\bar{G}_F \end{pmatrix}, \quad G_F = \begin{pmatrix} 1_2 & 0 \\ 0 & -1_2 \end{pmatrix}, \quad J_F = \begin{pmatrix} 0_4 & 1_4 \\ 1_4 & 0_4 \end{pmatrix} \circ cc \quad (15.1.8)$$

where cc stands for complex conjugation.

**Proposition 15.1** (1) *The data*

$$F_{PS} := (A_{PS}, H_F, D_F; J_F, \gamma_F)$$

as defined above is a finite real even spectral triple of KO-dimension 6 that fulfills the first-order condition on a subalgebra  $A_{SM} = \mathbb{C} \oplus \mathbb{H}_L \oplus M_3(\mathbb{C}) \subset A_{PS}$ .

- (2) The unimodular gauge group  $S\mathfrak{G}(F_{PS})$  is isomorphic to the Pati–Salam gauge group  $SU(2)_R \times SU(2)_L \times SU(4)$ .

**Proof** (1) follows from Proposition 13.1 while (2) is a straightforward computation, using that  $\mathcal{U}(\mathbb{H}) \cong SU(2)$  (see the proof of Proposition 13.3).

## 15.2 The Gauge and Scalar Field Contents

Since we are dealing with a real spectral triple that does not fulfill the first-order condition, we have to apply the general framework of inner fluctuations developed in Sect. 7.3. More precisely, the initial operator for the almost-commutative manifold  $M \times F_{PS}$  is given by

$$D = D_M \otimes 1 + \gamma_M \otimes D_F$$

for which the inner fluctuations are given by

$$D_\omega = D + \omega_{(1)} + J\omega_{(1)}J^{-1} + \omega_{(2)}$$

$$\omega_{(1)} = \sum a [D, b]; \quad \omega_{(2)} = \sum a [J\omega_{(1)}J^{-1}, b].$$

The computation is rather involved due to the second-order term, see Note 3 on Sect. 15.4. One finds that the different components of the operator  $D_\omega$  are given by

$$(D_\omega)_{\dot{a}I}^{\dot{b}J} = \gamma^\mu \left( \nabla_\mu^S \delta_{\dot{a}}^{\dot{b}} \delta_I^J - \frac{i}{2} g_R W_{\mu R}^\alpha (\sigma^\alpha)_a^{\dot{b}} \delta_I^J - \delta_{\dot{a}}^{\dot{b}} \left( \frac{i}{2} g V_\mu^m (\lambda^m)_I^J + \frac{i}{2} g V_{\mu I} \delta_J^J \right) \right)$$

$$(D_\omega)_{aI}^{bJ} = \gamma^\mu \left( \nabla_\mu^S \delta_a^b \delta_I^J - \frac{i}{2} g_L W_{\mu L}^\alpha (\sigma^\alpha)_a^b \delta_I^J - \delta_a^b \left( \frac{i}{2} g V_\mu^m (\lambda^m)_I^J + \frac{i}{2} g V_{\mu I} \delta_J^J \right) \right)$$

where the fifteen  $4 \times 4$  matrices  $(\lambda^m)_I^J$  are traceless and generate the group  $SU(4)$  and  $W_{\mu R}^\alpha$ ,  $W_{\mu L}^\alpha$ ,  $V_\mu^m$  are the gauge fields of  $SU(2)_R$ ,  $SU(2)_L$ , and  $SU(4)$ .

In addition we have

$$(D_\omega)_{\dot{a}I}^{bJ} = \gamma_M ((Y_v \phi_{\dot{a}}^b + Y_e \tilde{\phi}_{\dot{a}}^b) \Sigma_I^J + (Y_u \phi_{\dot{a}}^b + Y_d \tilde{\phi}_{\dot{a}}^b) (\delta_I^J - \Sigma_I^J)) \equiv \gamma_M \Sigma_{\dot{a}I}^{bJ}$$

$$(D_\omega)_{\dot{a}I}^{\dot{b}J'} = \gamma_M Y_R^* \Delta_{\dot{a}J} \Delta_{bI} \equiv \gamma_M H_{\dot{a}I\dot{b}J}$$

where the Higgs field  $\phi_{\dot{a}}^b$  is in the  $(2_R, \bar{2}_L, 1)$  of the product gauge group  $SU(2)_R \times SU(2)_L \times SU(4)$ , and  $\Delta_{\dot{a}J}$  is in the  $(2_R, 1_L, 4)$  representation while  $\Sigma_I^J$  is in the  $(1_R, 1_L, 1 + 15)$  representation. The field  $\tilde{\phi}_{\dot{a}}^b$  is not an independent field and is given by

$$\tilde{\phi}_{\dot{a}}^b = \sigma_2 \bar{\phi}_{\dot{a}}^b \sigma_2.$$

**Table 15.1** Pati–Salam scalar particle content and their representations for a first-order Dirac operator. The field  $\Sigma_J^I$  in the last row is decoupled if there is quark-lepton coupling unification

Particle	$SU(2)_R$	$SU(2)_L$	$SU(4)$
$\phi_{\dot{a}}^b$	2	2	1
$\Delta_{\dot{a}I}$	2	1	4
$\Sigma_J^I$	1	1	15

Note that the field  $\Sigma_J^I$  decouples (and set to  $\delta_I^1 \delta_1^J$ ) in the special case when there is lepton and quark unification of the couplings

$$Y_v = Y_u, \quad Y_e = Y_d.$$

This is summarized in Table 15.1.

**Proposition 15.2** *The spectral action (ignoring topological and boundary terms) of the almost-commutative manifold  $M \times F_{PS}$  is given by*

$$\begin{aligned} S_B = & \int_M \frac{48f_4}{\pi^2} \Lambda^4 - \frac{2f_2 \Lambda^2}{\pi^2} \left( R + \frac{1}{4} \left( H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{a}I} + 2\Sigma_{\dot{a}I}^{cK} \Sigma_{cK}^{\dot{a}I} \right) \right) \\ & + \frac{f(0)}{2\pi^2} \left[ \frac{1}{30} (-18C_{\mu\nu\rho\sigma}^2 + 11R^*R^*) + g_L^2 (W_{\mu\nu L}^\alpha)^2 + g_R^2 (W_{\mu\nu R}^\alpha)^2 + g^2 (V_{\mu\nu}^m)^2 \right. \\ & + D_\mu \Sigma_{\dot{a}I}^{\dot{c}K} D^\mu \Sigma_{\dot{c}K}^{\dot{a}I} + \frac{1}{2} D_\mu H_{\dot{a}I\dot{b}J} D^\mu H^{\dot{a}I\dot{b}J} + \frac{1}{12} R \left( H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{a}I} + 2\Sigma_{\dot{a}I}^{cK} \Sigma_{cK}^{\dot{a}I} \right) \\ & \left. + \frac{1}{2} \left| H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{b}J} \right|^2 + 2H_{\dot{a}I\dot{c}K} \Sigma_{\dot{b}J}^{\dot{c}K} H^{\dot{a}I\dot{d}L} \Sigma_{\dot{d}L}^{\dot{b}J} + \Sigma_{\dot{a}I}^{\dot{c}K} \Sigma_{\dot{c}K}^{\dot{b}J} \Sigma_{\dot{b}J}^{\dot{d}L} \Sigma_{\dot{d}L}^{\dot{a}I} \right] \sqrt{g} d^4x. \end{aligned}$$

**Proof** We proceed using the same notation and formulas as in Sect. 10.4. The first Seeley-de Witt coefficient is

$$\begin{aligned} a_0 &= \frac{1}{16\pi^2} \text{Tr}(1) \\ &= \frac{1}{16\pi^2} (4)(32)(3) \int \sqrt{g} d^4x \\ &= \frac{24}{\pi^2} \int_M \sqrt{g} d^4x \end{aligned}$$

where the numerical factors come, respectively, from the traces on the Clifford algebra, the dimensions of the Hilbert space and number of generations. The second coefficient is

$$a_2 = \frac{1}{16\pi^2} \int_M \text{Tr} \left( E + \frac{1}{6} R \right) \sqrt{g} d^4x$$

where  $E$  is a  $384 \times 384$  matrix over Hilbert space of three generations of spinors, whose components are derived and listed in the appendix. Taking the various traces we get

$$\begin{aligned} a_2 &= \frac{1}{16\pi^2} \int_M \left( (R(-96 + 64) - 8 \left( H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{a}I} + 2\Sigma_{\dot{a}I}^{cK} \Sigma_{cK}^{\dot{a}I} \right)) \sqrt{g} d^4x \right. \\ &\quad \left. = -\frac{2}{\pi^2} \int_M \left( R + \frac{1}{4} \left( H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{a}I} + 2\Sigma_{\dot{a}I}^{cK} \Sigma_{cK}^{\dot{a}I} \right) \right) \sqrt{g} d^4x. \right) \end{aligned}$$

It should be understood in the above formula and in what follows, that whenever the matrices  $Y_v$ ,  $Y_u$ ,  $Y_e$ ,  $Y_d$  and  $Y_R$  appear in an action, one must take the trace over generation space. The mass terms can be expressed in terms of the fundamental Higgs field to give

$$H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{a}I} = |Y_R|^2 \left( \Delta_{\dot{a}K} \bar{\Delta}^{\dot{a}K} \right)^2$$

and

$$\begin{aligned} 2\Sigma_{\dot{a}I}^{cK} \Sigma_{cK}^{\dot{a}I} &= 2 \left( ((Y_v - Y_u) \phi_{\dot{a}}^c + (Y_e - Y_d) \tilde{\phi}_{\dot{a}}^c) \Sigma_I^K + (Y_u \phi_{\dot{a}}^c + Y_d \tilde{\phi}_{\dot{a}}^c) \delta_I^K \right) \\ &\quad \left( ((Y_{*v} - Y_u^*) \phi_{\dot{a}}^{\dot{a}} + (Y_{*e} - Y_{*d}) \tilde{\phi}_{\dot{a}}^{\dot{a}}) \Sigma_K^I + (Y_u^* \phi_{\dot{a}}^{\dot{a}} + Y_{*d} \tilde{\phi}_{\dot{a}}^{\dot{a}}) \delta_K^I \right). \end{aligned}$$

The next coefficient is

$$a_4 = \frac{1}{16\pi^2} \int_M \text{Tr} \left( \frac{1}{360} \left( 5R^2 - 2R_{\mu\nu}^2 + 2R_{\mu\nu\rho\sigma}^2 \right) 1 + \frac{1}{2} \left( E^2 + \frac{1}{3} RE + \frac{1}{6} \Omega_{\mu\nu}^2 \right) \right) \sqrt{g} d^4x$$

where  $\Omega_{\mu\nu}$  is the  $384 \times 384$  curvature matrix of the connection  $\omega_\mu$ . Using the expressions for the matrices  $E$  and  $\Omega_{\mu\nu}$  derived in the appendix, and taking the traces, we get

$$\begin{aligned} a_4 &= \frac{1}{2\pi^2} \int_M \left[ -\frac{3}{5} C_{\mu\nu\rho\sigma}^2 + \frac{11}{30} R^* R^* + g_L^2 \left( W_{\mu\nu L}^\alpha \right)^2 + g_R^2 \left( W_{\mu\nu R}^\alpha \right)^2 + g^2 \left( V_{\mu\nu}^m \right)^2 \right. \\ &\quad + \nabla_\mu \Sigma_{\dot{a}I}^{\dot{c}K} \nabla^\mu \Sigma_{\dot{c}K}^{\dot{a}I} + \frac{1}{2} \nabla_\mu H_{\dot{a}I\dot{b}J} \nabla^\mu H^{\dot{a}I\dot{b}J} + \frac{1}{12} R \left( H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{a}I} + 2\Sigma_{\dot{a}I}^{cK} \Sigma_{cK}^{\dot{a}I} \right) \\ &\quad \left. + \frac{1}{2} \left| H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{b}J} \right|^2 + 2H_{\dot{a}I\dot{c}K} \Sigma_{bJ}^{\dot{c}K} H^{\dot{a}I\dot{d}L} \Sigma_{dL}^{bJ} + \Sigma_{\dot{a}I}^{\dot{c}K} \Sigma_{cK}^{\dot{b}J} \Sigma_{bJ}^{\dot{d}L} \Sigma_{dL}^{\dot{a}I} \right] \sqrt{g} d^4x \end{aligned}$$

where  $C_{\mu\nu\rho\sigma}$  is the Weyl tensor. The stated result now follows from Eq.(10.4.1).  $\square$

The physical content of this action is a cosmological constant term, the Einstein Hilbert term  $R$ , a Weyl tensor square term  $C_{\mu\nu\rho\sigma}^2$ , kinetic terms for the  $SU(2)_R \times SU(2)_L \times SU(4)$  gauge fields, kinetic terms for the composite Higgs fields  $H_{\dot{a}I\dot{b}J}$  and  $\Sigma_{bJ}^{cK}$  as well as mass terms and quartic terms for the Higgs fields. We also notice that this action gives the gauge coupling unification

$$g_R = g_L = g. \quad (15.2.1)$$

Having determined the full Dirac operators, including fluctuations, we can write all the fermionic interactions including the ones with the gauge vectors and Higgs scalars. We write the fermionic action using our tensorial notation:

$$\begin{aligned} & \psi_A^* D_A^B \psi_B + \psi_{A'}^* D_{A'}^B \psi_B + \psi_A^* D_A^{B'} \psi_{B'} + \psi_{A'}^* D_{A'}^{B'} \psi_{B'} \\ &= \psi_A^* D_A^B \psi_B + \psi_A C D^{AB} \psi_B + \text{h.c} \end{aligned}$$

where  $C$  is the charge conjugation matrix, and we have denoted  $D_{A'}^B = D^{AB}$ . We then find

$$\begin{aligned} & \int_M \left[ \psi_{\dot{a}I}^* \gamma^\mu \left( \nabla_\mu^S \delta_{\dot{a}}^{\dot{b}} \delta_I^J - \frac{i}{2} g_R W_{\mu R}^\alpha (\sigma^\alpha)_{\dot{a}}^{\dot{b}} \delta_I^J - \delta_{\dot{a}}^{\dot{b}} \left( \frac{i}{2} g V_\mu^m (\lambda^m)_I^J + \frac{i}{2} g V_\mu \delta_I^J \right) \right) \psi_{\dot{b}J} \right. \\ &+ \psi_{\dot{a}I}^* \gamma^\mu \left( \nabla_\mu^S \delta_a^b \delta_I^J - \frac{i}{2} g_L W_{\mu L}^\alpha (\sigma^\alpha)_a^b \delta_I^J - \delta_a^b \left( \frac{i}{2} g V_\mu^m (\lambda^m)_I^J + \frac{i}{2} g V_\mu \delta_I^J \right) \right) \psi_{bJ} \\ &+ \psi_{\dot{a}I}^* \gamma_5 \left( (Y_v \phi_a^b + Y_e \tilde{\phi}_a^b) \Sigma_I^J + (Y_u \phi_a^b + Y_d \tilde{\phi}_a^b) (\delta_I^J - \Sigma_I^J) \right) \psi_{bJ} \\ &+ \psi_{\dot{a}I}^* \gamma_5 \left( (Y_v^* \phi_a^b + Y_e^* \tilde{\phi}_a^b) \Sigma_I^J + (Y_u^* \phi_a^b + Y_d^* \tilde{\phi}_a^b) (\delta_I^J - \Sigma_I^J) \right) \psi_{bJ} \\ &+ \left. C \psi_{\dot{a}I} \gamma_5 Y_R \bar{\Delta}^{\dot{a}J} \bar{\Delta}^{\dot{b}I} \psi_{\dot{b}J} + \text{h.c} \right] \sqrt{g} d^4 x \end{aligned}$$

### 15.3 Truncation to the Standard Model

In this section we show that the above grand unified Pati-Salam type model can break to the  $U(1) \times SU(2) \times SU(3)$  symmetry of the SM.

First of all, the scalar field  $\phi_a^b = (2_R, 2_L, 1)$  must be truncated to the Higgs doublet  $H$  by writing

$$\phi_a^b = \delta_a^i \epsilon^{bc} H_c.$$

The other scalar field  $\Delta_{\dot{a}I} = (2_R, 1, 4)$  is truncated to a real singlet scalar field

$$\Delta_{\dot{a}I} = \delta_{\dot{a}}^i \delta_I^j \sqrt{\sigma}.$$

These then imply the relations

$$\begin{aligned} \Sigma_{\dot{a}I}^{bJ} &= \left( \delta_{\dot{a}}^i Y_v \epsilon^{bc} H_c + \delta_{\dot{a}}^i \bar{H}^b Y_e \right) \delta_I^i \delta_J^l + \left( \delta_{\dot{a}}^i Y_u \epsilon^{bc} H_c + \delta_{\dot{a}}^i Y_d \bar{H}^b \right) \delta_I^i \delta_J^l \delta_i^j \\ H_{\dot{a}I} \delta_{\dot{b}J} &= \delta_{\dot{a}}^i \delta_{\dot{b}}^j Y_R \delta_I^i \delta_J^l \sigma \\ g_R W_{\mu R}^3 &= g_1 B_\mu, \quad W_{\mu R}^\pm = 0 \\ \sqrt{\frac{3}{2}} g V_\mu^{15} &= -g_1 B_\mu \quad (V_\mu)_1^i = 0 \end{aligned}$$

where  $V_\mu^{15}$  is the  $SU(4)$  gauge field corresponding to the generator

$$\lambda^{15} = \frac{1}{\sqrt{6}} \text{diag}(3, -1, -1, -1)$$

which could be identified with the  $B - L$  generator. In particular the components  $(D_A)_{11}^{11}$  and  $(D_A)_{21}^{21}$  of the Dirac operator simplify to

$$\begin{aligned} (D_A)_{11}^{11} &= \gamma^\mu \left( \nabla_\mu^S - \frac{i}{2} g_R W_{\mu R}^\alpha (\sigma^\alpha)_1^1 - \left( \frac{i}{2} g V_\mu^m (\lambda^m)_1^1 \right) \right) \\ &= \gamma^\mu \left( \nabla_\mu^S - \frac{i}{2} g_R W_{\mu R}^3 - \left( \frac{i}{2} g V_\mu^{15} \sqrt{\frac{3}{2}} \right) \right) \\ &= \gamma^\mu \nabla_\mu^S \\ (D_A)_{21}^{21} &= \gamma^\mu \left( \nabla_\mu^S - \frac{i}{2} g_R W_{\mu R}^\alpha (\sigma^\alpha)_2^2 - \left( \frac{i}{2} g V_\mu^m (\lambda^m)_2^2 \right) \right) \\ &= \gamma^\mu \left( \nabla_\mu^S + \frac{i}{2} g_R W_{\mu R}^3 - \left( \frac{i}{2} g V_\mu^{15} \sqrt{\frac{3}{2}} \right) \right) \\ &= \gamma^\mu (\nabla_\mu^S + i g_1 B_\mu) \end{aligned}$$

which are identified with the Dirac operators acting on the right-handed neutrino and right-handed electron. Similar substitutions give the action of the Dirac operators on the remaining fermions and give the expected results. We now compute the various terms in the spectral action. First for the mass terms we have

$$\begin{aligned} \frac{1}{4} H_{aI\dot{b}J} H^{\dot{b}J\dot{a}I} &= \frac{1}{4} \left( \delta_a^1 \delta_{\dot{b}}^1 Y_R \delta_I^1 \delta_{\dot{I}}^J \sigma \right) \left( \delta_1^{\dot{a}} \delta_1^{\dot{b}} \delta_I^J \delta_{\dot{I}}^J Y_R^* \sigma \right) \\ &= \frac{1}{4} \text{tr} |Y_R|^2 \sigma^2 = \frac{1}{4} c \sigma^2 \\ \frac{1}{2} \Sigma_{aI}^{cK} \Sigma_{cK}^{\dot{a}I} &= \frac{1}{2} \left| \left( \delta_a^1 Y_v \epsilon^{bc} H_c + \delta_a^2 \bar{H}^b Y_e \right) \delta_I^1 \delta_{\dot{I}}^J + \left( \delta_a^1 Y_u \epsilon^{bc} H_c + \delta_a^2 Y_d \bar{H}^b \right) \delta_I^J \delta_{\dot{J}}^I \delta_{\dot{I}}^J \right|^2 \\ &= \frac{1}{2} a \bar{H} H \end{aligned}$$

where

$$\begin{aligned} a &= \text{tr} (Y_v^* Y_v + Y_e^* Y_e + 3 (Y_u^* Y_u + Y_d^* Y_d)) \\ c &= \text{tr} (Y_R^* Y_R) \end{aligned}$$

Next for the  $a_4$  term, starting with the gauge kinetic energies we have

$$g_L^2 (W_{\mu\nu L}^\alpha)^2 + g_R^2 (W_{\mu\nu R}^\alpha)^2 + g^2 (V_{\mu\nu}^m)^2 \rightarrow g_L^2 (W_{\mu\nu L}^\alpha)^2 + \frac{5}{3} g_1^2 B_{\mu\nu}^2 + g_3^2 (V_{\mu\nu}^m)^2$$

where  $m = 1, \dots, 8$  for  $V_{\mu\nu}^m$  restricted to the  $SU(3)$  gauge group. Next for the scalar kinetic and quartic terms we have

$$\begin{aligned}\nabla_\mu \Sigma_{aI}^{\dot{c}K} \nabla^\mu \Sigma_{\dot{c}K}^{aI} &\rightarrow a \nabla_\mu \bar{H} \nabla^\mu H \\ \frac{1}{2} \nabla_\mu H_{\dot{a}I\dot{b}J} \nabla^\mu H^{\dot{a}I\dot{b}J} &\rightarrow \frac{1}{2} c \partial_\mu \sigma \partial^\mu \sigma \\ \frac{1}{12} R \left( H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{a}I} + 2 \Sigma_{aI}^{\dot{c}K} \Sigma_{\dot{c}K}^{aI} \right) &\rightarrow \frac{1}{12} R (2a \bar{H} H + c \sigma^2) \\ \frac{1}{2} \left| H_{\dot{a}I\dot{c}K} H^{\dot{c}K\dot{b}J} \right|^2 &\rightarrow \frac{1}{2} d \sigma^4 \\ 2 H_{\dot{a}I\dot{c}K} \Sigma_{bJ}^{\dot{c}K} H^{\dot{a}I\dot{d}L} \Sigma_{dL}^{bJ} &\rightarrow 2e \bar{H} H \sigma^2 \\ \Sigma_{aI}^{\dot{c}K} \Sigma_{cK}^{bJ} \Sigma_{bJ}^{dL} \Sigma_{dL}^{aI} &\rightarrow b (\bar{H} H)^2\end{aligned}$$

Collecting all terms we end up with the bosonic action for the Standard Model:

$$\begin{aligned}S_B = \int_M \left( \frac{48f_4\Lambda^4}{\pi^2} - \frac{2f_2}{\pi^2} \Lambda^2 \left( R + \frac{1}{2}a \bar{H} H + \frac{1}{4}c \sigma^2 \right) \right. \\ \left. + \frac{f(0)}{2\pi^2} \left[ \frac{1}{30} (-18C_{\mu\nu\rho\sigma}^2 + 11R^*R^*) + \frac{5}{3}g_1^2 B_{\mu\nu}^2 + g_2^2 (W_{\mu\nu}^\alpha)^2 + g_3^2 (V_{\mu\nu}^m)^2 \right. \right. \\ \left. \left. + \frac{1}{6}aR \bar{H} H + b(\bar{H} H)^2 + a|\nabla_\mu H_a|^2 + 2e \bar{H} H \sigma^2 \right. \right. \\ \left. \left. + \frac{1}{2}d \sigma^4 + \frac{1}{12}cR \sigma^2 + \frac{1}{2}c(\partial_\mu \sigma)^2 \right] \right) \sqrt{g} d^4x\end{aligned}\quad (15.3.1)$$

where

$$\begin{aligned}b &= \text{tr} \left( (Y_v^* Y_v)^2 + (Y_e^* Y_e)^2 + 3 \left( (Y_u^* Y_u)^2 + (Y_d^* Y_d)^2 \right) \right) \\ d &= \text{tr} \left( (Y_R^* Y_R)^2 \right) \\ e &= \text{tr} \left( Y_v^* Y_v Y_R^* Y_R \right).\end{aligned}$$

This action completely agrees with the Standard Model Lagrangian obtained in Theorem 13.10, under the replacement  $Y_R \rightarrow Y_R \sigma$ .

## 15.4 Phenomenology of the Noncommutative Pati–Salam Model

An important test of the above Pati–Salam model is to check whether the gauge coupling unification (15.2.1) when run using RG equations would give values consistent with the values in Eq.(14.2.3) of the gauge couplings for electromagnetic,

weak and strong interactions at the scale of the  $Z$ -boson mass. Moreover, in view of the observations at the end of Chap. 14 it is important to make sure that the model is compatible with the relatively low observed mass of the Higgs boson.

### 15.4.1 Grand Unification of the Gauge Couplings

We have already computed the inner perturbations of the finite Dirac operator for the Pati–Salam model in Sect. 15.2. Recall in particular the scalar content from Table 15.1. We will assume that there is lepton quark unification, so that the  $\Sigma_I^J$  is decoupled.

The boundary conditions between the couplings are taken at the intermediate mass scale  $\mu = m_R$  to be the usual

$$\frac{1}{g_1^2} = \frac{2}{3} \frac{1}{g^2} + \frac{1}{g_R^2}, \quad \frac{1}{g_2^2} = \frac{1}{g_L^2}, \quad \frac{1}{g_3^2} = \frac{1}{g^2}, \quad (15.4.1)$$

in terms of the Standard Model gauge couplings  $g_1, g_2, g_3$ . At the mass scale  $m_R$  the Pati–Salam symmetry is broken to that of the Standard Model, and we take it to be the same scale that is present in the see-saw mechanism. It should thus be of the order  $10^{11} - 10^{13}$  GeV.

Before turning to the computation of the  $\beta$ -functions of the Pati–Salam gauge couplings for the composite model, let us discuss the scalar sector that remains after spontaneous symmetry breaking to the Standard Model gauge group. A quick analysis leads to the scalar fields listed in Table 15.2. Note that this includes the SM Higgs and a real scalar singlet.

The presence of the above scalar fields of course also have an influence on the running of the Standard Model gauge couplings (at one loop). We compute that instead of the usual  $\beta$ -functions  $(b_1, b_2, b_3) = (-\frac{41}{6}, \frac{19}{6}, 7)$  (cf. Sect. 14.2.1) we have

$$(b_1, b_2, b_3) = \left( -\frac{64}{9}, 3, \frac{41}{6} \right).$$

**Table 15.2** Scalar particle content induced by the Pati–Salam model with SM-representations

	$U(1)_Y$	$SU(2)_L$	$SU(3)$
$\begin{pmatrix} \phi_1^0 \\ \phi_1^+ \end{pmatrix} = \begin{pmatrix} \phi_1^1 \\ \phi_1^2 \end{pmatrix}$	1	2	1
$\begin{pmatrix} \phi_2^- \\ \phi_2^0 \end{pmatrix} = \begin{pmatrix} \phi_2^1 \\ \phi_2^2 \end{pmatrix}$	-1	2	1
$\sigma$	0	1	1
$\eta$	$-\frac{2}{3}$	1	3

One observes that this difference is relatively small (less than 5%). In fact, the scalar fields that appear in addition to the SM Higgs have a negligible effect in our study of the running of the gauge couplings below.

Next, we compute the  $\beta$ -functions for the Pati–Salam couplings  $g_R, g_L, g$  in the presence of the above composite particle content (cf. Table 15.1):

$$(b_R, b_L, b) = \left( \frac{7}{3}, 3, \frac{31}{3} \right). \quad (15.4.2)$$

The solutions of the RG-equations are easily found to be


$$g_R(\mu)^{-2} = g_R(m_R)^{-2} + \frac{1}{8\pi^2} \frac{7}{3} \log \frac{\mu}{m_R}, \quad (15.4.3)$$

$$g_L(\mu)^{-2} = g_L(m_R)^{-2} + \frac{1}{8\pi^2} 3 \log \frac{\mu}{m_R}, \quad (15.4.4)$$

$$g(\mu)^{-2} = g(m_R)^{-2} + \frac{1}{8\pi^2} \frac{31}{3} \log \frac{\mu}{m_R}, \quad (15.4.5)$$

We impose the boundary conditions (15.4.1) at the mass scale  $\mu = m_R$ .

Our approach for finding a unification scale is as follows. We search for an energy scale where the couplings  $g_R, g_L$  and  $g$  are equal by varying the scale  $m_R$  at which the boundary conditions (15.4.1) are imposed. With the running of the Pati–Salam couplings governed by the coefficients (15.4.2) there is a unique value of  $m_R$  for which the three lines meet. The unification scale is  $\Lambda \approx 2.5 \times 10^{15}$  GeV and the value found for the intermediate scale is  $m_R = 4.25 \times 10^{13}$  GeV (Fig. 15.1).



**Fig. 15.1** Running of coupling constants for the spectral Pati–Salam model with composite Higgs fields:  $g_1, g_2, g_3$  for  $\mu < m_R$  and  $g_R, g_L, g$  for  $\mu > m_R$  with unification scale  $\Lambda \approx 2.5 \times 10^{15}$  GeV for  $m_R = 4.25 \times 10^{13}$  GeV

If the scalar field  $\Sigma_I^J$  is not decoupled—in other words, if there is no lepton-quark coupling unification—then there is an additional scalar ( $1_R, 1_L, 15$ ) irreducible representation contributing to the  $\beta$ -function, giving a slightly different  $(b_R, b_L, b) = (\frac{7}{3}, 3, 9)$ . This in turn gives a unification scale  $\Lambda \approx 6.3 \times 10^{15}$  GeV for  $m_R = 4.1 \times 10^{13}$  GeV.

### 15.4.2 Running of the Higgs Mass

Let us analyze the additional terms in the spectral action displayed in Eq.(15.3.1), focusing on the scalar part:

$$\begin{aligned} \mathcal{L}'_H(g_{\mu\nu}, \Lambda_\mu, Q_\mu, H, \sigma) := & \frac{bf(0)}{2\pi^2}|H|^4 - \frac{2af_2\Lambda^2}{\pi^2}|H|^2 + \frac{ef(0)}{\pi^2}\sigma^2|H|^2 \\ & - \frac{cf_2\Lambda^2}{\pi^2}\sigma^2 + \frac{df(0)}{4\pi^2}\sigma^4 + \frac{af(0)}{2\pi^2}|D_\mu H|^2 + \frac{1}{4\pi^2}f(0)c(\partial_\mu\sigma)^2, \end{aligned}$$

where we ignored the coupling to the scalar curvature.

As in Chap. 14, we exploit the approximation that  $m_{top}$ ,  $m_v$  and  $m_R$  are the dominant mass terms. Moreover, as before we write  $m_v = \rho m_{top}$ . That is, the expressions for  $a, b, c, d$  and  $e$  in (13.3.2) now become

$$\begin{aligned} a &\approx m_{top}^2(\rho^2 + 3), \\ b &\approx m_{top}^4(\rho^4 + 3), \\ c &\approx m_R^2, \\ d &\approx m_R^4, \\ e &\approx \rho^2 m_R^2 m_{top}^2. \end{aligned}$$

In a unitary gauge, where  $H = \begin{pmatrix} h \\ 0 \end{pmatrix}$ , we arrive at the following potential:

$$\mathcal{L}_{pot}(h, \sigma) = \frac{1}{24}\lambda_h h^4 + \frac{1}{2}\lambda_{h\sigma} h^2\sigma^2 + \frac{1}{4}\lambda_\sigma \sigma^4 - \frac{4g_2^2}{\pi^2}f_2\Lambda^2(h^2 + \sigma^2),$$

where we have defined coupling constants

$$\lambda_h = 24 \frac{\rho^4 + 3}{(\rho^2 + 3)^2} g_2^2, \quad \lambda_{h\sigma} = \frac{8\rho^2}{\rho^2 + 3} g_2^2, \quad \lambda_\sigma = 8g_2^2. \quad (15.4.6)$$

This potential can be minimized, and if we replace  $h$  by  $v + h$  and  $\sigma$  by  $w + \sigma$ , respectively, expanding around a minimum for the terms quadratic in the fields, we obtain:

$$\begin{aligned}\mathcal{L}_{pot}(v + h, w + \sigma)|_{\text{quadratic}} &= \frac{1}{6}v^2\lambda_h v^2 + 2vw\lambda_{h\sigma}\sigma h + w^2\lambda_\sigma\sigma^2 \\ &= \frac{1}{2}(h\ \sigma) M^2 \begin{pmatrix} h \\ \sigma \end{pmatrix},\end{aligned}$$

where we have defined the mass matrix  $M$  by

$$M^2 = 2 \begin{pmatrix} \frac{1}{6}\lambda_h v^2 & \lambda_{h\sigma} vw \\ \lambda_{h\sigma} vw & \lambda_\sigma w^2 \end{pmatrix}.$$

This mass matrix can be easily diagonalized, and if we make the natural assumption that  $w$  is of the order of  $m_R$ , while  $v$  is of the order of  $M_W$ , so that  $v \ll w$ , we find that the two eigenvalues are


$$\begin{aligned}m_+^2 &\sim 2\lambda_\sigma w^2 + 2\frac{\lambda_{h\sigma}^2}{\lambda_\sigma}v^2, \\ m_-^2 &\sim 2\lambda_h v^2 \left( \frac{1}{6} - \frac{\lambda_{h\sigma}^2}{\lambda_h \lambda_\sigma} \right).\end{aligned}$$

We can now determine the value of these two masses by running the scalar coupling constants  $\lambda_h$ ,  $\lambda_{h\sigma}$  and  $\lambda_\sigma$  down to ordinary energy scalar. The renormalization group equations for these couplings are given by

$$\begin{aligned}\frac{d\lambda_h}{dt} &= \frac{1}{16\pi^2} \left( 4\lambda_h^2 + 12\lambda_{h\sigma}^2 - (3g_1^2 + 9g_2^2)\lambda_h + \frac{9}{4}(g_1^4 + 2g_1^2g_2^2 + 3g_2^4) \right. \\ &\quad \left. + 4(3y_{\text{top}}^2 + y_v^2)\lambda_h - 12(3y_{\text{top}}^4 + y_v^4) \right), \\ \frac{d\lambda_{h\sigma}}{dt} &= \frac{1}{16\pi^2} \left( 8\lambda_{h\sigma}^2 + 6\lambda_{h\sigma}\lambda_\sigma + 2\lambda_{h\sigma}\lambda_h \right. \\ &\quad \left. - \frac{3}{2}(g_1^2 + 3g_2^2)\lambda_{h\sigma} + 2(3y_{\text{top}}^2 + y_v^2)\lambda_{h\sigma} \right), \\ \frac{d\lambda_\sigma}{dt} &= \frac{1}{16\pi^2} \left( 8\lambda_{h\sigma}^2 + 18\lambda_\sigma^2 \right).\end{aligned}$$

As before, at lower energy the coupling  $y_v$  drops out of the RG equations and is replaced by an effective coupling.

At one-loop, the other couplings obey the renormalization group equations of the Standard Model, that is, they satisfy (14.2.9) and (14.2.10). As before, we can solve these differential equations, with boundary conditions at  $\Lambda_{GUT}$  given for the scalar couplings by (15.4.6). The result varies with the chosen value for  $\Lambda_{GUT}$  and the parameter  $\rho$ . The mass of  $\sigma$  is essentially given by the largest eigenvalue  $m_+$  which is of the order  $10^{12}$  GeV for all values of  $\Lambda_{GUT}$  and the parameter  $\rho$ . The allowed mass range for the Higgs, i.e. for  $m_-$ , is depicted in Fig. 15.2. The expected value



**Fig. 15.2** A contour plot of the Higgs mass  $m_h$  as a function of  $\rho^2$  and  $t = \log(\Lambda_{GUT}/M_Z)$ . The red line corresponds to  $m_h = 125.5$  GeV

$m_h = 125.5$  GeV is therefore compatible with the above noncommutative model, while the scalar field  $\sigma$  stabilizes the Higgs vacuum at higher energies. Furthermore, this calculation implies that there is a relation (given by the red line in the Figure) between the ratio  $m_\nu/m_{top}$  and the unification scale  $\Lambda_{GUT}$ .

We conclude that with noncommutative geometry we can proceed beyond the Standard Model, enlarging the field content of the Standard Model by a real scalar field with a mass of the order of  $10^{12}$  GeV. At the time of writing of the second edition of this book (Summer 2021), this is completely compatible with experiment and also guarantees stability of the Higgs vacuum at higher energy scales. Of course, the final word is to experiment in the years to come. What we can say at this point is that noncommutative geometry provides a fascinating dialogue between abstract mathematics and concrete measurements in experimental high-energy physics.

## Notes

### Section 15.1. The Finite Noncommutative Space of the Pati–Salam Model

1. The Pati–Salam model was introduced in [1]. The particle content that we find is very similar to the one considered by Marshak and Mohapatra [2].
2. Coincidentally the algebra  $M_2(\mathbb{H}) \oplus M_4(\mathbb{C})$  comes out as a solution of the two-sided Heisenberg quantization relation between the Dirac operator  $D$  and the two maps from the four spin-manifold and the two four spheres  $S^4 \times S^4$  [3, 4]. This removes the arbitrary symplectic constraint and replaces it with a relation that quantize the four-volume in terms of two quanta of geometry.

### Section 15.2. The Gauge and Scalar Field Contents

3. We refer to [5, Appendix A] for all details on the derivation of the inner fluctuations for the Pati–Salam model; see also [6].
4. The important point to notice in the derivation of the inner fluctuations for the Pati–Salam model is the novel phenomena of the appearance of composite Higgs field as is apparent in the above formulas where the Higgs field  $\Sigma_{aI}^{bJ}$  is formed out of the products of the fields  $\phi_a^b$  and  $\Sigma_I^J$  while the Higgs field  $H_{aIbJ}$  is made from the product of  $\Delta_{aI}\Delta_{bJ}$ . This composite structure is a result of the quadratic dependence of the gauge fields  $\omega_{(2)}$  on those appearing in  $\omega_{(1)}$ . The importance of this point should not be underestimated. The reason is that the main disadvantage of grand unified theories is the need for complicated Higgs representations with arbitrary potentials. In the noncommutative geometric setting, this problem is now solved by having minimal representations of the Higgs fields allowing for (quadratic) products of these representations. We also note that a very close model to the one deduced here is the one considered by Marshak and Mohapatra where the  $U(1)$  of the left-right model is identified with the  $B - L$  symmetry. They proposed the same Higgs fields  $(2_R, 2_L, 1)$ ,  $(2_R, 1, 4)$  and  $(1, 1, 15)$  we have, but also in addition the field  $(1, 2_L, 4)$ . However, they assumed that this Higgs fields does not get a vev, and thus does not effect the symmetry breaking. Although the broken generators of the  $SU(4)$  gauge fields can mediate lepto-quark interactions leading to proton decay, it was shown that in all such types of models with partial unification, the proton is stable. In addition this type of model arises in the first phase of breaking of  $SO(10)$  to  $SU(2)_R \times SU(2)_L \times SU(4)$  and these have been extensively studied [7]. The recent work in [8] considers noncommutative grand unification based on the  $k = 8$  algebra  $M_4(\mathbb{H}) \oplus M_8(\mathbb{C})$  keeping the first order condition.
5. The obstruction to allow for lower  $m_h$  (see Sect. 15.4.2) in the spectral Standard Model was overcome in [9] simply by taking into account the scalar field  $\sigma$  which was already present in the full model that was computed previously in [10].

### Section 15.4. Phenomenology of the Noncommutative Pati–Salam Model

6. This section is based on [5, 11]. For a recent overview, we also refer to [6].
7. The boundary condition (15.4.1) can be found in [12, Eq. (5.8.3)].
8. The field  $\sigma$  played a key role in [10] in lowering the Higgs mass prediction to a realistic value [9]. A qualitative study of the form of the scalar potential that we have done for the present Pati–Salam composite model indicates that this result continues to hold here. However, being interested mainly in the running of the gauge couplings, we leave a full study of the potential and its physical implications for future work.
9. Note that in our analysis we have disregarded the non-renormalizable, order eight terms that appear in the expansion of the spectral action for the composite model [5, Sect. 8], so let us argue

why they can be ignored. In fact, since we consider only the running of the gauge couplings at the one loop level, we can safely ignore these non-renormalizable terms. Moreover, their contribution to the running of other (scalar) couplings will be suppressed by negative powers of  $m_R$ , at least at the one loop level.

10. In view of the assumptions made in our analysis, we trust the values for  $m_r$  only as indicative of the corresponding orders of magnitudes. Other possible Pati–Salam models (for different initial  $D_F$ ) were considered in [11].
11. The renormalization group equations for the couplings  $\lambda_h$ ,  $\lambda_{h\sigma}$ ,  $\lambda_\sigma$  have been derived in [13].
12. For stability bounds on the Higgs mass, we refer to [14].
13. The small correction to the space  $M \times F_{SM}$  was realized in [9] (and already tacitly present in [10]) and the results of Sect. 15.4.2 confirm their conclusions.
14. Other noncommutative geometric models that go beyond the Standard Model include [15–19], adopting a slightly different approach to almost-commutative manifolds as we do (cf. Note 3 on Sect. 7.2). The intersection between supersymmetry and almost-commutative manifolds is analyzed in [20–22].

## References

1. Pati, J.C., Salam, A.: Lepton number as the fourth color. *Phys. Rev. D* **10**, 275–289 (1974)
2. Mohapatra, R.N., Marshak, R.E.: Local B-L Symmetry of electroweak interactions, majorana neutrinos and neutron oscillations. *Phys. Rev. Lett.* **44**, 1316–1319 (1980). [Erratum: *Phys. Rev. Lett.* **44**, 1643 (1980)]
3. Chamseddine, A.H., Connes, A., Mukhanov, V.: Geometry and the quantum: basics. *JHEP* **1412**, 098 (2014)
4. Chamseddine, A.H., Connes, A., Mukhanov, V.: Quanta of geometry: noncommutative aspects. *Phys. Rev. Lett.* **114**, 091302 (2015)
5. Chamseddine, A.H., Connes, A., van Suijlekom, W.D.: Beyond the spectral standard model: emergence of Pati-Salam unification. *JHEP* **1311**, 132 (2013)
6. Chamseddine, A., van Suijlekom, W.D.: A survey of spectral models of gravity coupled to matter. In: *Advances in Noncommutative Geometry—on the Occasion of Alain Connes' 70th Birthday*, pp. 1–51. Springer, Cham (2019)
7. Babu, K.S., Mohapatra, R.N.: Coupling unification, GUT-Scale baryogenesis and neutron-antineutron oscillation in SO(10). *Phys. Lett. B* **715**, 328–334 (2012)
8. Devastato, A., Lizzi, F., Martinetti, P.: Grand symmetry, spectral action, and the Higgs mass. *JHEP* **1401**, 042 (2014)
9. Chamseddine, A.H., Connes, A.: Resilience of the spectral standard model. *JHEP* **1209**, 104 (2012)
10. Chamseddine, A.H., Connes, A.: Noncommutative geometry as a framework for unification of all fundamental interactions including gravity. Part I. *Fortsch. Phys.* **58**, 553–600 (2010)
11. Chamseddine, A.H., Connes, A., van Suijlekom, W.D.: Grand unification in the spectral Pati–Salam model. *JHEP* **11**, 011 (2015)
12. Mohapatra, R.: Unification and Supersymmetry. *The Frontiers of Quark-Lepton Physics*, 3rd edn. Springer, New York (2003)
13. Gonderinger, M., Li, Y., Patel, H., Ramsey-Musolf, M.J.: Vacuum stability, perturbativity, and scalar singlet dark matter. *JHEP* **1001**, 053 (2010)
14. Sher, M.: Electroweak Higgs potentials and vacuum stability. *Phys. Rept.* **179**, 273–418 (1989)
15. Stephan, C.A.: Almost-commutative geometries beyond the Standard Model. *J. Phys. A* **39**, 9657 (2006)
16. Stephan, C.A.: Almost-commutative geometries beyond the Standard Model. II. New colours. *J. Phys. A* **40**, 9941 (2007)

17. Stephan, C.A.: New scalar fields in noncommutative geometry. *Phys. Rev. D* **79**, 065013 (2009)
18. Stephan, C.A.: Beyond the Standard Model: A Noncommutative Approach, 0905.0997
19. Stephan, C.A.: A dark sector extension of the almost-commutative standard model. *Int. J. Mod. Phys. A* **29**, 1450005 (2014)
20. van den Broek, T., van Suijlekom, W.D.: Supersymmetric QCD and noncommutative geometry. *Commun. Math. Phys.* **303**, 149–173 (2011)
21. van den Broek, T., van Suijlekom, W.D.: Supersymmetric QCD from noncommutative geometry. *Phys. Lett. B* **699**, 119–122 (2011)
22. Beenakker, W., van den Broek, T., van Suijlekom, W. D.: Supersymmetry and Noncommutative Geometry. SpringerBrief in Mathematical Physics 9, Springer (2016)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Chapter 16

## Towards a Quantum Theory



In the final Chapter of this book we present an overview of recent and ongoing work, taking the first steps towards a quantum theory for noncommutative geometry. Indeed, in the applications to particle physics phenomenology one applies the usual, physicist' textbook renormalization group methods to the spectral action, in order to arrive at couplings and mass parameters at lower energy. And even though the appearance of such experimentally testable results from a geometrical framework valid at high-energies is very intriguing, this step remains a weak point of the noncommutative approach to particle physics. In other words, it means that in the passage to the quantum theory one loses the elegant spectral and unifying picture that one started with.

We indicate two paths that could lead to a quantum theory for noncommutative manifolds. Working in the general context of spectral triples, the first approach quantizes the fermionic content by applying the procedure of second quantization to  $(\mathcal{A}, \mathcal{H}, D)$ , while the second computes one-loop corrections to the perturbative expansion of the spectral action, given in its general form as in Sect. 9.3.

### 16.1 Second Quantization of Spectral Triples

We use the operator algebra formalism of  $C^*$ -dynamical systems and KMS condition to pass from the first-quantized or “one-particle” level of spectral triples  $(\mathcal{A}, \mathcal{H}, D)$  to the second-quantized level. The Hilbert space  $\mathcal{H}$  is used to construct the complexified Clifford algebra  $C := \text{Cliff}_{\mathbb{C}}(\mathcal{H}_{\mathbb{R}})$  of its underlying real Hilbert space  $\mathcal{H}_{\mathbb{R}}$  when considered as a Euclidean space. The operator  $D$  is used as the generator of a one-parameter group  $\sigma_t \in \text{Aut}(C)$  of automorphisms of the Clifford algebra. The algebra  $\mathcal{A}$  manifests itself through the inner fluctuations (cf. Eq. (7.2.5)) deforming the operator  $D$  to  $D'$ . These inner fluctuations continue to make sense at the second-quantized level and give rise to deformations  $\sigma'_t \in \text{Aut}(C)$  of the above one-parameter group

of automorphisms. We concentrate here on the meaning of the spectral action, and thus take the  $C^*$ -dynamical system  $(\mathcal{C}, \sigma_t)$  as our starting point, keeping in mind that the results will automatically apply to the deformations  $(\mathcal{C}, \sigma'_t)$ .

### 16.1.1 KMS and a Dynamical System

We first briefly recall the KMS condition for  $C^*$ -dynamical systems, that is, a  $C^*$ -algebra  $\mathcal{C}$  together with a one-parameter group of automorphisms  $\sigma_t \in \text{Aut}(\mathcal{C})$ ,  $t \in \mathbb{R}$ .

**Definition 16.1** Let  $(\mathcal{C}, \sigma_t)$  be a  $C^*$ -dynamical system. For a given  $0 < \beta < \infty$ , a state  $\varphi$  on the unital  $C^*$ -algebra  $\mathcal{C}$  satisfies the KMS condition at inverse temperature  $\beta$  if for all  $a, b \in \mathcal{C}$ , there exists a function  $F_{a,b}(z)$  which is holomorphic on the strip

$$I_\beta = \{z \in \mathbb{C} \mid 0 < \Im(z) < \beta\}, \quad (16.1.1)$$

continuous on the boundary  $\partial I_\beta$  and bounded, with the property that for all  $t \in \mathbb{R}$

$$F_{a,b}(t) = \varphi(a\sigma_t(b)) \quad \text{and} \quad F_{a,b}(t + i\beta) = \varphi(\sigma_t(b)a). \quad (16.1.2)$$

In short the KMS condition at inverse temperature  $\beta$  means that one has the formal equality

$$\varphi(a\sigma_t(b))|_{t=i\beta} = \varphi(ba), \quad \forall a, b \in \mathcal{C}. \quad (16.1.3)$$

What matters in the context of the present paper is the existence and uniqueness of KMS states on a matrix algebra  $M_n(\mathbb{C})$  and we give the short proof for convenience. A one-parameter group of automorphisms  $\sigma_t \in \text{Aut}(M_n(\mathbb{C}))$  is always associated to a self-adjoint  $H = H^* \in M_n(\mathbb{C})$  by

$$\sigma_t(A) = e^{itH} A e^{-itH}, \quad \forall t \in \mathbb{R}, \quad A \in M_n(\mathbb{C}).$$

Given a state  $\psi$  on the matrix algebra  $M_n(\mathbb{C})$  there exists a unique density matrix  $\rho \geq 0$  such that

$$\psi(T) = \text{Tr}(\rho T), \quad \forall T \in M_n(\mathbb{C}).$$

By uniqueness of the trace it follows that

$$\psi(AB) = \psi(BA), \quad \forall A, B \Rightarrow \rho = \frac{1}{n} \text{id}.$$

A state which is KMS $_\beta$  for  $\sigma_t \in \text{Aut}(M_n(\mathbb{C}))$  is invariant. In fact

$$\psi(e^{-\beta H} B e^{\beta H}) = \psi(B), \quad \forall B \Rightarrow e^{\beta H} \rho e^{-\beta H} = \rho \Rightarrow \rho e^{\beta H} = e^{\beta H} \rho.$$

It follows using (16.1.3) that if  $\psi$  is KMS $_{\beta}$  then, with  $B' = e^{-\beta H}B$

$$\psi(BA) = \psi(Ae^{-\beta H}Be^{\beta H}) \Rightarrow \text{Tr}(\rho e^{\beta H}B'A) = \text{Tr}(\rho AB'e^{\beta H})$$

so that  $\rho e^{\beta H} = e^{\beta H}\rho$  defines a trace and hence is a scalar multiple of id. This shows that  $\rho = Ze^{-\beta H}$  for  $Z = 1/\text{Tr}(e^{-\beta H})$  and gives the uniqueness of the KMS $_{\beta}$  state. The same formula gives the existence. For completeness, we include a proof of the following result on KMS-states on Clifford algebras

**Proposition 16.2** *Let  $\mathcal{H}$  be a complex Hilbert space,  $D$  a self-adjoint operator in  $\mathcal{H}$  with compact resolvent. Let  $C := \text{Cliff}_{\mathbb{C}}(\mathcal{H}_{\mathbb{R}})$  be the complexified Clifford algebra of the underlying real Hilbert space  $\mathcal{H}_{\mathbb{R}}$  and  $\sigma_t \in \text{Aut}(C)$  be the one-parameter group of automorphisms associated to  $\exp(itD) \in \text{Aut}(\mathcal{H}_{\mathbb{R}})$ . Then for any  $\beta > 0$  there exists a unique KMS $_{\beta}$  state  $\psi_{\beta}$  on the  $C^*$ -dynamical system  $(C, \sigma_t)$ .*

**Proof** One applies the existence and uniqueness of KMS states on a matrix algebra to the subalgebra of  $C = \text{Cliff}_{\mathbb{C}}(\mathcal{H}_{\mathbb{R}})$  associated to the subspace corresponding to a finite dimensional spectral projection of  $D$ . This is enough to prove the uniqueness of the KMS $_{\beta}$  state. The existence also follows since the existence part for matrix algebras gives a coherent system of states which define a state on the inductive limit of the  $C^*$ -algebras.  $\square$

**Proposition 16.3** *Let  $\mathcal{H}, D, C := \text{Cliff}_{\mathbb{C}}(\mathcal{H}_{\mathbb{R}})$ ,  $\sigma_t \in \text{Aut}(C)$  and  $\psi_{\beta}$  be as in Proposition 16.2. Then if the operator  $\exp(-\beta|D|)$  is of trace class, the state  $\psi_{\beta}$  is of type I and the associated irreducible representation is given by the fermionic second quantization associated to the complex structure  $I := i \operatorname{sign} D$  on  $\mathcal{H}_{\mathbb{R}}$ .*

The proof of this proposition will be given in Sect. 16.1.2, Proposition 16.5 after recalling some terminology.

### 16.1.2 Fermionic Second Quantization

In this section we recall the procedure of (fermionic) second quantization.

Consider the real Euclidean vector space  $V := \mathcal{H}_{\mathbb{R}}$  that underlies the complex Hilbert space  $\mathcal{H}$ . Our first goal is to find irreducible representations of the Clifford algebra associated to the real Euclidean vector space  $\mathcal{H}_{\mathbb{R}}$  underlying  $\mathcal{H}$  and for this it turns out that a crucial role is played by complex structures. We let  $I$  be an orthogonal complex structure on  $V$ , which is not necessarily the one coming from  $\mathcal{H}$ . Then we may regard  $V$  as a complex vector space when we define  $i$  to act as  $I$ . The resulting complex Hilbert space will be denoted by  $V_I$ .

A representation of the complexified Clifford algebra  $\text{Cliff}_{\mathbb{C}}(V)$  is given on the Fock space  $\bigwedge V_I$  that is built on  $V_I$  by the usual formula

$$\begin{aligned} \gamma_I : \text{Cliff}_{\mathbb{C}}(V) &\rightarrow \mathcal{L}(\bigwedge V_I) \\ v &\mapsto a_I^*(v) + a_I(v); \quad (v \in V). \end{aligned}$$

Here the *creation operators*  $a_I^*(v)$  depend  $\mathbb{C}$ -linearly on  $v \in V_I$  and are given by exterior multiplication by  $v$  while the *annihilation operator*  $a_I(v)$  is its adjoint. We choose a unit vector  $\Omega_I \in \bigwedge^0 V_I$  and call it the *vacuum vector*. It is annihilated by  $a_I(v)$  for all  $v \in V$ . The following is well-known.

**Lemma 16.4** *The above representation  $\gamma_I$  of the complexified Clifford algebra  $\text{Cliff}_{\mathbb{C}}(V)$  on Fock space  $\bigwedge V_I$  is irreducible.*

**Proof** We may assume that  $V$  is a inductive limit of finite-dimensional Hilbert spaces, and, accordingly, that  $\text{Cliff}_{\mathbb{C}}(V)$  is the  $C^*$ -algebraic inductive limit of finite-dimensional Clifford algebras. Without loss of generality we may thus assume that  $\dim V < \infty$  so that  $\text{Cliff}_{\mathbb{C}}(V)$  are simple matrix algebras. We invoke Schur's Lemma to conclude that  $\gamma_I$  is irreducible if and only if every operator  $T : V \rightarrow V$  commuting with all  $\gamma_I(v)$  ( $v \in V^{\mathbb{C}}$ ) is a scalar.

For any  $v$ , one has, using the  $\mathbb{C}$ -linearity of  $a_I^*(v)$  and  $\mathbb{C}$ -anti-linearity of  $a_I(v)$

$$a_I^*(v) = \frac{1}{2} (\gamma_I(v) - i\gamma_I(Iv)), \quad a_I(v) = \frac{1}{2} (\gamma_I(v) + i\gamma_I(Iv)).$$

Hence any  $T$  that commutes with  $\gamma_I(v)$  for all  $v$  commutes with  $a_I^*(v)$  and  $a_I(v)$ . From this it follows that

$$a_I(v)(T\Omega_I) = T(a_I(v)\Omega_I) = 0$$

so that  $T\Omega_I \in \Lambda^0(V_I)$ . In other words,  $T\Omega_I = t\Omega_I$  for some  $t \in \mathbb{C}$ . Moreover,

$$T(v_1 \wedge \cdots \wedge v_k) = T(a_I^*(v_1) \cdots a_I^*(v_k)\Omega) = (a_I^*(v_1) \cdots a_I^*(v_k))(T\Omega) = t(v_1 \wedge \cdots \wedge v_k),$$

so that  $T = t \cdot \text{id}$ .  $\square$

Any orthogonal operator  $T : V \rightarrow V$  induces an automorphism of  $\text{Cliff}_{\mathbb{C}}(V)$  by sending  $\gamma_I(v) \rightarrow \gamma_I(Tv)$ . In some cases this automorphism can be lifted to the Fock space  $\bigwedge V_I$ , for instance, if  $U$  is a unitary operator on  $V_I$ . Then, if  $\bigwedge U$  is the unitary operator in the Fock space such that on simple tensors

$$\bigwedge U(v_1 \wedge \cdots \wedge v_n) := U(v_1) \wedge \cdots \wedge U(v_n),$$

one has the covariance

$$\bigwedge U \circ a_I^*(v) \circ \bigwedge U^* = a_I^*(Uv).$$

We thus get the equality

$$(\bigwedge U)\gamma_I(v)(\bigwedge U^*) = \gamma_I(Uv).$$

Suppose now that we are given a (complex) Hilbert space  $\mathcal{H}$  and a self-adjoint operator  $D$  in  $\mathcal{H}$  with compact resolvent. Again, let  $V = \mathcal{H}_{\mathbb{R}}$  denote the underlying real vector space. Suppose that we take the natural complex structure on  $V$  so that  $V_I = \mathcal{H}$ . Then the above construction gives us an irreducible representation  $\gamma$  of the canonical anti-commutation relations (CAR) algebra on  $\bigwedge \mathcal{H}$  but, from a physical point of view this representation is not the right one to consider. In fact, one needs to choose another irreducible representation, corresponding to a different complex structure on  $\mathcal{H}$ . Let us describe it in some detail.

If  $E_{\pm}$  are the spectral projections of  $D$  corresponding to the positive and negative eigenspaces of  $D$  in  $\mathcal{H}$ , let us define the following complex structure:

$$I = i(E_+ - E_-).$$

In other words,  $I = iF$  where  $F = D|D|^{-1}$  is the sign<sup>1</sup> of  $D$ . In view of the previous section, this gives rise to another irreducible representation  $\gamma_I$  of  $\text{Cliff}_{\mathbb{C}}(V)$  in Fock space, where the key difference with respect to the original Fock space representation  $\gamma$  is that  $i$  acts as  $-i$  on  $E_-(\mathcal{H})$ . In other words, the operator  $D$  can be considered to act as  $|D|$  which now only has positive eigenvalues. More precisely,

**Proposition 16.5** (i) *The one-parameter group  $\sigma_t \in \text{Aut}(C)$  is implemented in the (physical) Fock representation by the one-parameter unitary group  $W(t) = \bigwedge \exp(it|D|)$ , i.e. one has*

$$\gamma_I(\sigma_t(A)) = \bigwedge (e^{it|D|}) \gamma_I(A) \bigwedge (e^{-it|D|}), \quad \forall A \in \text{Cliff}_{\mathbb{C}}(V). \quad (16.1.4)$$

(ii) *If  $\exp(-\beta|D|)$  is of trace class the state  $\psi_{\beta}$  is of type I and is given by*

$$\psi_{\beta}(A) = \frac{1}{Z} \text{Tr} \left( \bigwedge \exp(-\beta|D|) \gamma_I(A) \right), \quad \forall A \in \text{Cliff}_{\mathbb{C}}(V) \quad (16.1.5)$$

where the normalization factor  $Z$  is finite.

**Proof** The associated KMS $_{\beta}$  state is obtained after normalization from the density matrix  $W(i\beta)$  obtained by analytic continuation. Thus it coincides here with the operator

$$\rho = \bigwedge \exp(-\beta|D|).$$

Since  $T = \exp(-\beta|D|)$  is positive and of trace class we get that  $\rho = \bigwedge T$  is also positive and of trace class (with trace given by the determinant of  $1 + T$ ). Thus  $Z < \infty$ . The state  $\psi_{\beta}$  is implemented by a density matrix in an irreducible representation and is thus of type I.  $\square$

---

<sup>1</sup> We take the convention that the sign of 0 is 1.

### 16.1.3 von Neumann Information Theoretic Entropy

We start by briefly recalling von Neumann's notion of entropy. Consider a density matrix  $\rho$  on a Hilbert space  $\mathcal{H}$ , *i.e.* a positive trace-class operator with normalized trace. It induces a state  $\phi$  on any  $C^*$ -subalgebra of  $\mathcal{L}(\mathcal{H})$  by setting  $\phi(\cdot) = \text{Tr}(\rho \cdot)$ . The *entropy* of this state  $\phi$  is then defined to be

$$S(\phi) := -\text{Tr}(\rho \log \rho).$$

For composite systems  $\phi_1 \otimes \phi_2$  on  $\mathcal{H}_1 \otimes \mathcal{H}_2$  one finds the following important additivity property for entropy

$$S(\phi_1 \otimes \phi_2) = S(\phi_1) + S(\phi_2).$$

Let us start with a basic example of entropy that will play a crucial role in what follows.

**Lemma 16.6** *Let  $x > 0$ , the entropy of the partition of the unit interval in two intervals with ratio of size  $x$ , is given by*

$$\mathcal{E}(x) := \log(x+1) - \frac{x \log(x)}{x+1}.$$

**Proof** The sizes of the intervals are  $\frac{1}{x+1}$  and  $\frac{x}{x+1}$ . One has

$$-\frac{\log(\frac{1}{x+1})}{x+1} - \frac{x \log(\frac{x}{x+1})}{x+1} = \frac{x+1}{x+1} \log(x+1) - \frac{x \log(x)}{x+1} = \mathcal{E}(x).$$

□

**Corollary 16.7** *One has  $\mathcal{E}(x) = \mathcal{E}(1/x)$  for any  $x > 0$ .*

**Proof** The obtained partitions are isomorphic. □

The following result gives an expression for the entropy of density matrices that arise as a second-quantized operator.

**Lemma 16.8** *Let  $T \in \mathcal{L}^1(\mathcal{H})^+$  be a positive trace class operator and  $\phi$  the state associated to  $\bigwedge T$  then*

$$S(\phi) = \text{Tr}(\mathcal{E}(T)). \quad (16.1.6)$$

**Proof** Let first  $T = T_1 \oplus T_2$  be an orthogonal decomposition. Let us show that the associated states fulfill  $\phi = \phi_1 \otimes \phi_2$ . One has with  $\rho_I = \bigwedge T_I$  the equality  $\rho = \rho_1 \otimes \rho_2$  by the compatibility of the wedge functor with direct sums. Then

$$\text{Tr}(\rho_1 \otimes \rho_2) = \text{Tr}(\rho_1) \text{Tr}(\rho_2) \Rightarrow \phi = \phi_1 \otimes \phi_2.$$

Next the entropy functional fulfills

$$S(\phi_1 \otimes \phi_2) = S(\phi_1) + S(\phi_2).$$

This shows that the functional  $S(\phi)$  is additive for direct sum decompositions. It also applies to infinite sums and one can thus consider only the one dimensional case. In this case the state associated to the operator  $T$  of multiplication by  $x$  corresponds to  $\bigwedge T$  whose spectrum is  $\{1, x\}$  and hence has entropy given by the function  $\mathcal{E}(x)$  of Lemma 16.6.  $\square$

**Theorem 16.9** *Let  $\mathcal{H}, D, C := \text{Cliff}_{\mathbb{C}}(\mathcal{H}_{\mathbb{R}})$ ,  $\sigma_t \in \text{Aut}(C)$  and  $\psi_{\beta}$  be as in Proposition 16.2. Then if the operator  $\exp(-\beta|D|)$  is of trace class, the state  $\psi_{\beta}$  is of type I and its von Neumann entropy is equal to the spectral action  $\text{Tr}(h(\beta D))$  for the spectral function  $h(x) := \mathcal{E}(e^{-x})$ .*

**Proof** The first statement follows from Proposition 16.5. The statement about the entropy follows from Lemma 16.8 together with the fact that  $x \mapsto \mathcal{E}(e^{-x})$  is an even function (cf. Corollary 16.7).  $\square$

## 16.2 One-Loop Corrections to the Spectral Action

We start with the expansion of the spectral action derived in Eq. 9.3.2:

$$S_b[\omega] - S_b[0] = \sum_n \frac{1}{n} \langle \underbrace{\omega, \dots, \omega}_n \rangle_f$$

We thus work under the same assumptions as those stated in Sect. 9.3.

The bracket will be represented as the following Feynman diagram:

$$\langle \omega_1, \dots, \omega_n \rangle = \begin{array}{c} \text{Diagram: A circular loop with } n \text{ external wavy lines labeled } \omega_1, \dots, \omega_n. \\ \text{The lines are labeled clockwise starting from the bottom-left.} \end{array} \quad (16.2.1)$$

The loop diagram nicely reflects the cyclicity of the bracket:  $\langle \omega_1, \dots, \omega_n \rangle_f = \langle \omega_n, \omega_1, \dots, \omega_{n-1} \rangle_f$ . The second crucial property is that

$$\langle a\omega_1, \dots, \omega_n \rangle_f - \langle \omega_1, \dots, \omega_n a \rangle_f = \langle [D, a], \omega_1, \dots, \omega_n \rangle_f \quad (16.2.2)$$

In fact, this identity boils down to a *Ward identity*, represented diagrammatically as


$$\begin{array}{c} \text{---} \\ | \\ a \end{array} - \begin{array}{c} \text{---} \\ | \\ a \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ [D, a] \end{array} \quad (16.2.3)$$

In order to analyze the quantum theory corresponding to the above classical action functional  $S_b[\omega]$  we adopt the background field method. We take the background fields to be the usual gauge fields of the form  $\omega = \sum_j a_j[D, b_j] \in \Omega_D^1(\mathcal{A})$  but allow the path integral to integrate over all finite-size hermitian complex-valued matrices  $Q$ . We consider the dimension, say  $N$ , of these matrices as a regularizing cutoff of our model, which should eventually be sent to  $\infty$ .

For such matrices  $Q = (Q_{kl})$ , the brackets can be conveniently expressed in terms of divided differences of  $f'$ . Indeed, as in Eq.(9.3.3) we have:

$$\begin{aligned} \frac{1}{2}\langle Q, Q \rangle &= \frac{1}{2} \sum_{k,l} Q_{kl} Q_{lk} f'[\lambda_k, \lambda_l] \\ \frac{1}{3}\langle Q, Q, Q \rangle &= \frac{1}{3} \sum_{k,l,m} Q_{kl} Q_{lm} Q_{mk} f'[\lambda_k, \lambda_l, \lambda_m] \end{aligned}$$

*et cetera*, where  $\lambda_k$  are the eigenvalues of  $D$ .



(a) An example of a positive function:  $f(x) = (1 + ax^2)\Phi(bx)$  with  $\Phi$  a bump function and  $a = 1/900, b = 1/100$ .

(b) The divided difference  $f'[\lambda_k, \lambda_l]$  for this function  $f$ .

**Fig. 16.1** The inverse gauge propagator  $f'[\lambda_k, \lambda_l]$  for the  $N = 61$  smallest eigenvalues of the Dirac operator on the circle (i.e.  $\lambda_k, \lambda_l = -30, -29, \dots, 30$ )

We now make the assumption that the first divided difference of  $f'$  is strictly positive on the  $N$  relevant eigenvalues of  $D$  (see Fig. 16.1 for an example of such an  $f$ ). We may then perform the Gaussian integration to get for the propagator:

$$\overline{Q_{kl} Q_{mn}} = \frac{\int Q_{kl} Q_{mn} e^{-\frac{1}{2}\langle Q, Q \rangle_f} dQ}{\int e^{-\frac{1}{2}\langle Q, Q \rangle_f} dQ} = \delta_{kn} \delta_{lm} G_{kl}$$

in terms of  $G_{kl} := \frac{1}{f'[\lambda_k, \lambda_l]}$ . Notice that the inverse propagator is bounded, which is in stark contrast to the usual unbounded nature of inverse propagators in ordinary local quantum field theory.

In any case, we are now in a position to consider higher-loop contributions to the spectral action, and, in particular, all one-particle irreducible  $n$ -point Feynman graphs. Their (possibly divergent) amplitudes form the starting point of the renormalization process of the spectral action.

### 16.2.1 Ward Identity for the Gauge Propagator

In addition to the Ward identity (16.2.3) for the fermion propagator, we claim that we also have the following Ward identity for the gauge propagator:

$$(16.2.4)$$

where every fermion loop adds a minus sign. Indeed, the left-hand side is

$$\begin{aligned} & \overline{Q_{ik} Q_{lm} a_{mn}} - a_{im} \overline{Q_{mk} Q_{ln}} \\ &= G_{ik} \delta_{im} \delta_{kl} a_{mn} - G_{ln} \delta_{mn} \delta_{kl} a_{im} \\ &= (G_{ik} - G_{nk}) \delta_{kl} a_{in} \end{aligned}$$

while for the right-hand side we use the defining property of the divided differences to find:

**Table 16.1** The two-point graphs at one-loop

$$\begin{aligned}
 & -\overline{Q_{ik}Q_{rp}}a_{pq}(\lambda_p - \lambda_q)\overline{Q_{qr}Q_{ln}}f'[\lambda_p, \lambda_q, \lambda_r] \\
 & = -G_{ik}\delta_{ip}\delta_{kr}G_{qr}\delta_{qn}\delta_{rl}a_{pq}(\lambda_p - \lambda_q)f'[\lambda_p, \lambda_q, \lambda_r] \\
 & = G_{ik}G_{nk}(f'[\lambda_k, \lambda_n] - f'[\lambda_i, \lambda_k])\delta_{kl}a_{in}.
 \end{aligned}$$

The two expressions coincide because of the very fact that the free propagator is the inverse of the divided difference.

### 16.2.2 Two-Point Functions at One-Loop

The two-point graphs at one-loop are given in Table 16.1. The external fields  $\omega_1, \omega_2$  should be assigned to the external legs in all different cyclical manners.


The amplitude for the first graph is given by

$$\begin{aligned}
 \omega_1 \text{---} \text{loop} \text{---} \omega_2 &= \sum_{\substack{i,j,k \\ l,m,n}} (\omega_1)_{ij} \overbrace{Q_{jk}Q_{ki}(\omega_2)_{lm}Q_{mn}Q_{nl}}^{\text{loop}} \times f'[\lambda_i, \lambda_j, \lambda_k]f'[\lambda_l, \lambda_m, \lambda_n] \\
 &= \sum_{i,k} (\omega_1)_{ii}(\omega_2)_{kk}G_{ik}^2f'[\lambda_i, \lambda_i, \lambda_k]f'[\lambda_i, \lambda_k, \lambda_k].
 \end{aligned} \tag{16.2.5}$$

In particular, there is no running loop index in this expression and so this diagram remains finite even when the size  $N$  of the matrices is sent to  $\infty$ . We conclude that the amplitude of this graph is not relevant for renormalization purposes.

We then turn to the second graph in Table 16.1, and compute

$$\begin{aligned}
 \omega_1 \text{---} \text{loop} \text{---} \omega_2 &= \sum_{\substack{i,j,k \\ l,m,n}} (\omega_1)_{ij} \overbrace{Q_{jk}Q_{ki}(\omega_2)_{lm}Q_{mn}Q_{nl}}^{\text{loop}} f'[\lambda_i, \lambda_j, \lambda_k]f'[\lambda_l, \lambda_m, \lambda_n] \\
 &= \sum_{i,j,k} (\omega_1)_{ij}(\omega_2)_{ji}G_{ik}G_{kj}f'[\lambda_i, \lambda_j, \lambda_k]^2.
 \end{aligned} \tag{16.2.6}$$



**Fig. 16.2** The behaviour of the summands (indexed by  $\lambda_k$  running from  $-30$  to  $30$ ) for the vertex contribution in (16.2.6) and (16.2.7) for the Dirac operator on the circle and function  $f$  as in Fig. 16.2a

We find that this amplitude has a potential divergence in the limit that  $N \rightarrow \infty$  (see Fig. 16.2 for the behaviour of the summands). As such it should be subtracted from the effective action in order to render the theory finite after removal of the regulator.

For the final diagram with two external lines we compute its amplitude to be:

$$\begin{aligned}
 \text{Diagram with two external lines} &= \sum_{i,j,k,l} (\omega_1)_{ij} Q_{jk} \overline{Q_{kl}} (\omega_2)_{li} f'[\lambda_i, \lambda_j, \lambda_k, \lambda_l] \\
 &= \sum_{i,j,k} (\omega_1)_{ij} (\omega_2)_{ji} G_{jk} f''[\lambda_i, \lambda_j, \lambda_k].
 \end{aligned} \tag{16.2.7}$$

Again, this graph amplitude is potentially divergent in the limit  $N \rightarrow \infty$  and should thus be subtracted. The same applies to the same graph but with  $\omega_1$  and  $\omega_2$  exchanged.

### 16.2.3 One-Loop Counterterms to the Spectral Action

The computations of the graph amplitudes in the previous section show that the second two graphs in Table 16.1 are the relevant ones to consider as counterterms for the spectral action. However, since the spectral action is in particular a gauge theory, it is crucial that such counterterms are of the same form as the terms appearing in the spectral action.

As may be expected, a crucial role will be played by so-called *quantum Ward identities*. They form the analogue of (16.2.3) for the divergent component of the 1PI  $n$ -point functions at one loop. Let us denote by  $\langle\!\langle \omega_1, \dots, \omega_n \rangle\!\rangle^{1L}$  all one-loop  $n$ -point graphs whose amplitudes involve a sum over a loop index. The skeletons for such graphs are depicted in Table 16.2, for which all external lines are written outside the graph diagram, and labelled in cyclical order. Indeed, if an external line would be in the interior of the diagram, it is surrounded by the loop in the diagram, and will thus prevent the loop index from running (as in Eq. 16.2.5).

The quantum Ward identities are now given by

$$\begin{aligned} \langle\!\langle \omega_1, \dots, a\omega_j, \dots, \omega_n \rangle\!\rangle^{1L} - \langle\!\langle \omega_1, \dots, \omega_{j-1}a, \dots, \omega_n \rangle\!\rangle^{1L} \\ = \langle\!\langle \omega_1, \dots, \omega_{j-1}, [D, a], \omega_j, \dots, \omega_n \rangle\!\rangle^{1L}. \end{aligned}$$

It is this identity, in combination with cyclicity of the bracket  $\langle\!\langle \omega_1, \dots, \omega_n \rangle\!\rangle = \langle\!\langle \omega_n, \omega_1, \dots, \omega_{n-1} \rangle\!\rangle$ , which allows us to follow line-by-line the derivation of the Chern–Simons and Yang–Mills terms in Theorem 9.23.

**Theorem 16.10** *The divergent part of the one-loop quantum effective spectral action can be expanded as*

$$\sum_n \frac{1}{n} \langle\!\langle \omega, \dots, \omega \rangle\!\rangle_\infty^{1L} = \sum_{k=1}^{\infty} \left( \int_{\tilde{\psi}_{2k-1}} \text{cs}_{2k-1}(A) + \frac{1}{2k} \int_{\tilde{\phi}_{2k}} F^k \right).$$


Here  $\tilde{\phi}$  and  $\tilde{\psi}$  are the analogues of  $\phi$  and  $\psi$  as defined in (9.3.4) and (9.3.5) but now using the double bracket.

**Proof** All divergent one-loop diagrams have skeletons as depicted in Table 16.2, with the external lines labelled cyclically from 1 to  $n$ . The decoration of the external legs of our graphs with the external fields  $\omega_1, \dots, \omega_n$  then proceeds according to this labelling 1, ...,  $n$  and, upon summing over all such decorated graphs  $G$ , we get

$$\langle\!\langle \omega_1, \dots, \omega_n \rangle\!\rangle^{1L} = \sum_G G_{\omega_1, \dots, \omega_n}.$$

The left-hand side of the quantum Ward identity essentially comes down to connecting external edges to the graphs  $G$ . We will write  $G_i$  for the graph  $G$  with an insertion of an external gauge edge at a point  $i$  in between  $n$  and 1: this insertion

**Table 16.2** Skeletons for divergent one-loop  $n$ -point functions with increasing number of vertices. The fermion loops that define the vertices are all oriented as clockwise



point  $i$  can be either an outer fermion line in  $G$  (as in (16.2.3)) or, if 1 and  $n$  are not attached to the same vertex in  $G$ , a gauge propagator (as in 16.2.4). We then find

$$\langle\langle a\omega_1, \dots, \omega_n \rangle\rangle^{1L} - \langle\langle \omega_1, \dots, \omega_n a \rangle\rangle^{1L} = \sum_{G,i} (G_i)_{[D,a],\omega_1,\dots,\omega_n},$$

where the decoration  $[D, a]$  is attached to the external gauge edge inserted at the point  $i$  of  $G_i$ .

It is clear that the sum over  $G$  and  $i$  yield all decorated  $n + 1$ -point graphs, and, moreover, that any  $n + 1$ -point graph with labels  $[D, a], \omega_1, \dots, \omega_n$  is obtained in a unique manner from an insertion of an external edge in an  $n$ -point graph, as described above. We are thus left with  $\langle\langle [D, a], \omega_1, \dots, \omega_n \rangle\rangle^{1L}$ , as desired.  $\square$

We conclude that the passage to the one-loop renormalized spectral action can be realized by a transformation in the space of noncommutative integrals, sending  $\phi \mapsto \phi + \tilde{\phi}$  and  $\psi \mapsto \psi + \tilde{\psi}$ , thus rendering the theory (one-loop) renormalizable as a gauge theory. Of course, a general “power-counting” procedure and diagrammatics beyond the one-loop order is of great importance, but at the moment of writing still waiting to be developed.

## Notes

### Section 16.1. Second Quantization of Spectral Triples

1. Section 16.1 is based on [1].
2. A proof of Proposition 16.2 can be found in [2, Proposition 5.2.23].
3. We refer to [3, 4] and [5, Sect. 5.3 and 6.1] for excellent expositions on fermionic second quantization.
4. The lift of the unitary operator  $U$  to the Fock space is a special case of the *Shale–Stinespring Theorem* [6]. It states that the automorphism on  $\text{Cliff}_{\mathbb{C}}(V)$  defined by an orthogonal operator  $T : V \rightarrow V$  is implementable by a unitary operator on Fock space  $\bigwedge V_I$  if and only if  $T + ITI$  is Hilbert–Schmidt.
5. The discussion of the representation  $\gamma_I$  for the different complex structure  $I$  on  $\mathcal{H}$  derives from the work of Dirac who realized in [7] that in order to avoid unwanted negative energy solutions to his Dirac equation, one has to fill up (what is now called) the *Dirac sea*.
6. There is an intriguing relation between the function  $h$  that appears in Theorem 16.9 and the Riemann zeta function. All details can be found in [1]. See also the more general treatment in [8], including the chemical potential for both the bosonic and fermionic case.

### Section 16.2. One-loop Corrections to the Spectral Action

7. Section 16.2 is based on [9].
8. The boundeness of the inverse propagator is another manifestation of the regularizing properties of the spectral action, in line with [10–13]. It is an interesting problem to analyze the form of the propagator for more general  $f$ , including a possible gauge fixing, for instance along the lines of [14, 15] or by means of orthogonal polynomials as in [16].
9. Note that due to the cyclic symmetry of the vertices, an equivalent representation of the Feynman diagrams may be given by ribbon graphs as in [17], identifying

We however stick with the original fermion cycle vertices, as they are especially convenient to capture the Ward identities (16.2.3) and (16.2.4).

10. The type of one-loop graphs derived from the spectral action are familiar in the context of matrix models. In fact, it is interesting to confront this to the proof of renormalizability for noncommutative scalar field theories [17]. One of the main differences is that they consider so-called non-local matrix models [18] with a quartic vertex, while instead we have a local matrix model but with vertices of arbitrary valence.

## References

1. Chamseddine, A.H., Connes, A., van Suijlekom, W.D.: Entropy and the spectral action. *Commun. Math. Phys.* **373**, 457–471 (2020)
2. Bratteli, O., Robinson, D.W.: Operator Algebras and Quantum-Statistical Mechanics. II, 2nd ed. Springer, New York (1997)
3. Araki, H.: Bogoliubov automorphisms and Fock representations of canonical anticommutation relations. In: *Operator Algebras and Mathematical Physics* (Iowa City, Iowa, 1985), volume 62

- of Contemporary Mathematics, pp. 23–141. American Mathematical Society, Providence, RI (1987)
- 4. Carey, A.L., Ruijsenaars, S.N.M.: On fermion gauge groups, current algebras and Kac-Moody algebras. *Acta Appl. Math.* **10**, 1–86 (1987)
  - 5. Gracia-Bondía, J.M., Várilly, J.C., Figueroa, H.: Elements of Noncommutative Geometry. Birkhäuser, Boston (2001)
  - 6. Shale, D., Stinespring, W.F.: Spinor representations of infinite orthogonal groups. *J. Math. Mech.* **14**, 315–322 (1965)
  - 7. Dirac, P.A.M.: Quantum theory of emission and absorption of radiation. *Proc. Roy. Soc. Lond. A* **114**, 243 (1927)
  - 8. Dong, R., Khalkhali, M., van Suijlekom, W.D.: Second quantization and the spectral action. *J. Geom. Phys.* **167**, 104285 (2021)
  - 9. van Nuland, T.D.H., van Suijlekom, W.D.: One-loop corrections to the spectral action. *JHEP* **078** (2022)
  - 10. van Suijlekom, W.D.: Renormalization of the spectral action for the Yang-Mills system. *JHEP* **1103**, 146 (2011)
  - 11. Iochum, B., Levy, C., Vassilevich, D.: Spectral action beyond the weak-field approximation. *Commun. Math. Phys.* **316**, 595–613 (2012)
  - 12. Kurkov, M.A., Lizzi, F., Vassilevich, D.: High energy bosons do not propagate. *Phys. Lett. B* **731**, 311–315 (2014)
  - 13. Alkofer, N., Saueressig, F., Zanusso, O.: Spectral dimensions from the spectral action. *Phys. Rev. D* **91**, 025025 (2015)
  - 14. Iseppi, R.A., van Suijlekom, W.D.: Noncommutative geometry and the BV formalism: application to a matrix model. *J. Geom. Phys.* **120**, 129–141 (2017)
  - 15. Iseppi, R.A.: The BV formalism: theory and application to a matrix model. *Rev. Math. Phys.* **31**, 1950035, 24 (2019)
  - 16. Bessis, D., Itzykson, C., Zuber, J.B.: Quantum field theory techniques in graphical enumeration. *Adv. Appl. Math.* **1**, 109–157 (1980)
  - 17. Grosse, H., Wulkenhaar, R.: Renormalization of  $\phi^4$  theory on noncommutative  $\mathbb{R}^4$  in the matrix base. *Commun. Math. Phys.* **256**, 305–374 (2005)
  - 18. Grosse, H., Wulkenhaar, R.: Power-counting theorem for non-local matrix models and renormalisation. *Commun. Math. Phys.* **254**, 91–127 (2005)

**Open Access** This book is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



# Appendix

## Alphabetic Index, and Notation Index

$(\cdot, \cdot)$	Inner product (with values in $\mathbb{C}$ )
$\langle \cdot, \cdot \rangle$	Hermitian structure/algebra valued inner product
$\langle \cdot, \cdot \rangle_f$	Multi-linear functionals on differential forms
$\langle \cdot, \cdot \rangle_z$	Zeta functions defining the improper cocycles
$\langle\langle \cdot, \cdot \rangle\rangle$	Loop corrections to spectral action
$\ \cdot\ _{\text{Lip}}$	Lipschitz (semi)-norm
$\ \cdot\ _s$	Sobolev norm
$\ \cdot\ $	Operator/ $C^*$ -norm
$\mathcal{A}$	$*$ -algebra
$\mathcal{A}_\theta$	Smooth noncommutative torus algebra
$A$	$C^*$ -algebra
$A_\theta$	Noncommutative torus $C^*$ -algebra
$\mathcal{A}^\circ$ or $A^\circ$	Opposite algebra
$\widehat{A}$	Structure space of $A$
$(A, H, D)$	Finite spectral triple
$(A, H, D; J, \gamma)$	Finite real spectral triple
$(\mathcal{A}, \mathcal{H}, D)$	Spectral triple
$(\mathcal{A}, \mathcal{H}, D; J, \gamma)$	Real spectral triple
$\mathcal{A}_J$	Commutative subalgebra of $\mathcal{A}$
$\mathcal{A}^N$	Direct sum of $N$ copies of $\mathcal{A}$
$\widehat{A}(R)$	$\widehat{A}$ -form of Riemannian curvature
$\alpha$	Algebra automorphism
$\text{Aut}(\mathcal{A})$	Group of algebra automorphisms
$A_\mu$	Gauge field
$\text{ad } A_\mu$	Gauge field in adjoint representation
$\text{Ad}$	Adjoint action
$a_k$	Seeley-DeWitt coefficient
$\mathcal{B}(\mathcal{H})$	$C^*$ -algebra of bounded operators on Hilbert space $\mathcal{H}$

$\mathfrak{B}$	$C^*$ -bundle
$b, B$	Boundary operators on cochains
$B_\mu$	Gauge field in adjoint representation
$F_{\mu\nu}$	Field strength (curvature) of $B_\mu$
$C(X)$	$C^*$ -algebra of continuous functions on compact topological space $X$
$C^\infty(M)$	$*$ -algebra of smooth functions on compact smooth manifold $M$
$C^\infty(M_\theta)$	$*$ -algebra of the toric noncommutative manifold $M_\theta$
$\mathbb{C}^{n^\circ}$	Defining representation of $M_n(\mathbb{C})^\circ$
$\text{Cl}(V, Q)$	Clifford algebra
$\chi$	$\mathbb{Z}_2$ -grading on Clifford algebra
$\text{Cl}^0(V, Q)$	Even part of Clifford algebra
$\text{Cl}^1(V, Q)$	Odd part of Clifford algebra
$\text{Cl}_n^\pm$	Clifford algebra $\text{Cl}(\mathbb{R}^n, \pm Q_n)$
$\mathbb{C}\text{Cl}_n$	Clifford algebra $\text{Cl}(\mathbb{C}^n, Q_n)$
$\mathbb{C}\text{Cl}_n^0$	Even part of Clifford algebra $\text{Cl}_n$
$\mathbb{C}\text{Cl}_n^1$	Ddd part of Clifford algebra $\text{Cl}_n$
$(\text{Cl}_n^\pm)^0$	Even part of Clifford algebra $\text{Cl}_n^\pm$
$(\text{Cl}_n^\pm)^1$	Odd part of Clifford algebra $\text{Cl}_n^\pm$
$\text{Cl}^\pm(TM)$	Clifford algebra bundle
$\text{Cliff}^\pm(M)$	Space of sections of Clifford algebra bundle
$\mathbb{C}\text{Cl}(TM)$	Complexified Clifford algebra bundle
$\text{Cliff}(M)$	Space of sections of complexified Clifford algebra bundle
$\text{Cliff}_{\mathbb{C}}(V)$	Complexified Clifford algebra of vector space $V$
$c$	Clifford multiplication
$C^n(\mathcal{A})$	$n$ -cochains on algebra $\mathcal{A}$
$C^{\text{ev}}(\mathcal{A})$	Even cocains
$C^{\text{odd}}(\mathcal{A})$	Odd cochains
$c_{p,k}$	Combinatorial coefficients in residue cocycle
$c(k_1, \dots, k_j)$	Combinatorial coefficients in residue cocycle
$\text{ch}(E)$	Chern character of vector bundle $E$
$C_{\mu\nu\rho\sigma}$	Weyl curvature
$\chi(M)$	Euler characteristic
$D$	Self-adjoint operator
$D_F$	Finite Dirac operator
$d_{ij}$	Metric on finite discrete space
$D_{ij}$	Components of Dirac operator in finite spectral triple
$D_{ij,kl}$	Components of Dirac operator in finite real spectral triple
$D_e$	Component of finite Dirac operator labeling the edge $e$
$D_M$	Dirac operator on Riemannian spin manifold
$D_{\mathcal{A}_\theta}$	Dirac operator on the noncommutative torus
$\Delta^E$	Laplacian on vector bundle
$\Delta^S$	Laplacian on spinor bundle

$d_g$	Riemannian distance function
$d(x, y)$	Distance function
$d$	Derivation given by $[D, \cdot]$
$\delta$	Derivation given by $[ D , \cdot]$
$D_p$	$p D p$
$D_u$	$P u P$ with $P = \frac{1}{2}(1 + \text{Sign } D)$
$\text{Der}(\mathcal{A})$	Lie algebra of algebra derivations
$D_\omega$	Operator $D$ in the presence of inner fluctuation $\omega$
$d_R$	Right-handed down quark
$d_L$	Left-handed down quark
$\mathcal{E}$ or $E$	Algebra module
$\mathcal{E}^\circ$ or $E^\circ$	Conjugate module
$E_A$	Right $A$ -module
$_A E$	Left $A$ -module
${}_A E_B$	$A - B$ -bimodule
$E \otimes_A F$	Balanced tensor product
$\text{End}_{\mathcal{A}}(\mathcal{E})$	Algebra of module endomorphisms
$\text{End}(S)$	Endomorphism bundle of $S$
$e_R$	Right-handed electron
$e_L$	Left-handed electron
$\mathbb{F}$	Field ( $\mathbb{R}, \mathbb{C}$ or $\mathbb{H}$ )
$F_B$	Right $B$ -module
$F_B \circ {}_B E$	Kasparov product
$\phi_n$	$(b, B)$ -cocycle
$\langle \phi, p \rangle$	Pairing between even $(b, B)$ -cocycle and projection
$\langle \phi, u \rangle$	Pairing between odd $(b, B)$ -cocycle and unitary
$f[x_0, \dots, x_n]$	Divided difference of $f$ of order $n$
$f_k$	Moments of function $f$
$F_X$	Two-point space
$F$	Finite noncommutative space/finite real spectral triple
$F_{ED}$	Finite real spectral triple for electrodynamics
$F_{YM}$	Finite real spectral triple for Yang–Mills theory
$F_{SM}$	Finite real spectral triple for the Standard Model
$\phi, \Phi$	Scalar field
$\Gamma = (\Gamma^{(0)}, \Gamma^{(1)})$	Graph
$(\Gamma, \Lambda)$	$\Lambda$ -decorated graph/Krajewski diagram
$\Gamma(E)$	Space of continuous sections of a vector bundle $E$
$\Gamma^\infty(E)$	Space of smooth sections of a vector bundle $E$
$\Gamma_{\mu\nu}^\kappa$	Christoffel symbols
$\tilde{\Gamma}_{\mu a}^b$	Christoffel symbols in orthonormal basis
$\gamma_{n+1}$	Chirality operator
$\gamma_\mu$	Dirac gamma matrices, generators of $\text{Cliff}^+(M)$

$\gamma_a$	Generators of $\text{Cliff}^+(M)$ in orthonormal basis
$\gamma_M$	$\mathbb{Z}_2$ -grading on spinors
$\gamma$	$\mathbb{Z}_2$ -grading
$g$	Riemannian metric
$\mathfrak{G}(\mathcal{A}, \mathcal{H}; J)$	Gauge group
$\mathfrak{G}(M \times F)$	Gauge group of almost-commutative manifold
$\mathfrak{G}(F)$	Gauge group of finite noncommutative space
$\mathfrak{g}(\mathcal{A}, \mathcal{H}; J)$	Gauge Lie algebra
$\mathfrak{g}(M \times F)$	Gauge Lie algebra of almost-commutative manifold
$\mathfrak{g}(F)$	Gauge Lie algebra of finite noncommutative space
$G_\mu$	$SU(3)$ Standard Model gauge field
$G_{\mu\nu}$	Field strength (curvature) of $G_\mu$
$g_1$	$U(1)$ coupling constant
$g_2$	$SU(2)$ coupling constant
$g_3$	$SU(3)$ coupling constant
$\mathbb{H}$	Quaternions
$\mathcal{H}(H)$	Hilbert space (finite-dimensional Hilbert space)
$\mathcal{H}^+$	Positive eigenspace of grading $\gamma$
$\mathcal{H}^s$	Sobolev space
$\mathcal{H}_{cl}$	Set of Grassmann variables in $\mathcal{H}$
$H_l$	Hilbert space of leptons
$H_q$	Hilbert space of quarks
$HH^n(\mathcal{A})$	Hochschild cohomology
$HCP^{\text{ev}}(\mathcal{A})$	Even cyclic cohomology
$HCP^{\text{odd}}(\mathcal{A})$	Odd cyclic cohomology
$\text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{F})$	Space of module endomorphisms
$\mathfrak{H}(F)$	Group of unitary elements in commutative subalgebra
$\mathfrak{h}(F)$	Lie algebra of skew-hermitian elements in commutative subalgebra
$H$	Higgs field
$h$	Higgs field (unitary gauge)
$\mathbb{I}_N$	$N \times N$ identity matrix
$\text{Inn}(\mathcal{A})$	Group of inner automorphisms
$J$	Anti-unitary operator/real structure
$j : \Gamma \rightarrow \Gamma$	Involutive graph automorphism
$J_n^\pm$	Anti-linear map
$J_M$	Charge conjugation
$\mathsf{KK}_f(A, B)$	Set of Kasparov modules for $(A, B)$
$\kappa$	Effective Yukawa coupling for tau-neutrino
$\mathcal{L}$	Lagrangian
$\Lambda_\mu$	$U(1)$ Standard Model gauge field

$\Lambda_{\mu\nu}$	Field strength (curvature) of $\Lambda_\mu$
$\lambda$	Higgs quartic coupling
$\Lambda_{12}$	Electroweak unification scale
$\Lambda_{23}$	Weak-strong unification scale
$\Lambda_{GUT}$	Grand unification scale
$M_n(\mathbb{F})$	$*$ -algebra of $n \times n$ matrices with entries in $\mathbb{F}$
$M_n(A)$	$*$ -algebra of $n \times n$ matrices with entries in $A$
$M$	Manifold
$M \times F$	Almost-commutative manifold
$\mu_N$	Group of $N$ 'th roots of unity
$M_W$	Mass of $W$ -boson
$M_Z$	Mass of $Z$ -boson
$m_{top}$	Mass of top quark
$m_h$	Higgs mass
$m_\nu$	Neutrino mass
$m_R$	Majorana mass matrix
$m_l$	Effective mass of tau-neutrino
<b>n</b>	Defining representation $\mathbb{C}^n$ of $M_n(\mathbb{C})$
$\nabla$	Connection on module/vector bundle
$\nabla^{\mathfrak{B}}$	$*$ -algebra connection
$\nabla^S$	Spin connection
$\nu_R$	Right-handed neutrino
$\nu_L$	Left-handed neutrino
$\omega$	Connection one-form/inner fluctuation
$\omega^\#$	Vector field corresponding to one-form $\omega$
$\Omega^E$	Curvature of connection on bundle $E$
$\Omega_{dR}^k(M)$	De Rham differential k-forms
$\Omega_{dR}^k(M, A)$	De Rham differential k-forms with values in algebra $A$
$\Omega_k(M)$	De Rham $k$ -currents on $M$
$\Omega_D^1(\mathcal{A})$	Connes' differential one-forms associated to a spectral triple
$op^r$	Space of operators of analytic order $\leq r$
$Out(\mathcal{A})$	Group of outer automorphisms
$\pi$	Algebra representation
$\pi(A)'$	Commutant of $\pi(A)$
$\Psi_p(a^0, \dots, a^p)$	Improper $(b, B)$ -cocycle
$\Psi(\mathcal{A})$	Pseudodifferential operators
$\Psi^k(\mathcal{A})$	Pseudodifferential operators of order $k$
$p$	Projection
$PU(N)$	Projective unitary group
$q$	Quaternion
$q_\lambda$	Embedding of $\mathbb{C}$ in $\mathbb{H}$

$Q$	Quadratic form
$Q_n$	Standard quadratic form on $\mathbb{R}^n$ or $\mathbb{C}^n$
$Q_\mu$	$SU(2)$ Standard Model gauge field
$Q_{\mu\nu}$	Field strength (curvature) of $Q_\mu$
$R(X, Y)$	Riemannian curvature tensor
$R_{\mu\nu\kappa\lambda}$	Riemannian curvature tensor
$R_{\nu\lambda}$	Ricci tensor
$R^* R^*$	Pontryagin class
$\text{res}_{s=0} \Psi_p$	Residue cocycle
$\rho$	Ratio between mass of tau-neutrino and top quark
$s$	Scalar curvature
$S$	Spinor bundle
$Sd$	Dimension spectrum
$S_b[\omega]$	Spectral action
$S_b^{(n)}$	$n$ 'th Gâteaux derivative of spectral action
$S_{top}[\omega]$	Topological spectral action
$S_f[\omega, \psi]$	Fermionic action
$S\mathcal{U}(\mathcal{A}_F)$	Group of elements in $\mathcal{U}(\mathcal{A}_F)$ with determinant 1
$S\mathfrak{H}(F)$	Group of elements in $\mathfrak{H}(F)$ with determinant 1
$su(\mathcal{A}_F)$	Lie algebra of traceless elements in $\mathfrak{u}(\mathcal{A}_F)$
$s\mathfrak{h}(F)$	Lie algebra of traceless elements in $\mathfrak{h}(F)$
$S_N$	Group of permutations on $N$ elements
$SO(N)$	Group of special orthogonal transformations of $\mathbb{R}^N$
$SU(N)$	Group of special unitary transformations of $\mathbb{C}^N$
$su(N)$	Lie algebra of traceless skew-hermitian transformations of $\mathbb{C}^N$
$\sigma$	Real scalar field
$\sigma_t$	One-parameter group of isomorphism of $C^*$ -algebra
$TM$	Tangent bundle
$T^*M$	Cotangent bundle
$T^{(k)}$	$k$ 'th iteration of derivation $T \mapsto [D^2, T]$
$u$	Unitary algebra element
$U$	Unitary operator
$\mathcal{U}(\mathcal{A})$	Group of unitary elements in $*$ -algebra $\mathcal{A}$
$\mathfrak{u}(\mathcal{A})$	Lie algebra of skew-hermitian algebra elements
$U(N)$	Group of unitary transformations of $\mathbb{C}^N$
$u(N)$	Lie algebra of skew-hermitian transformations of $\mathbb{C}^N$
$u_R$	Right-handed up quark
$u_L$	Left-handed up quark
$V_\mu$	$SU(3)$ Standard Model gauge field
$V_{\mu\nu}$	Field strength (curvature) of $V_\mu$
$v$	Higgs vacuum expectation value

$W_\mu$	$SU(2)$ Standard Model gauge field
$W_{\mu\nu}$	Field strength (curvature) of $W_\mu$
$Y_\nu, Y_e, Y_u, Y_d$	Yukawa mass matrices
$Y_R$	Majorana mass matrix
$Y_\mu$	$U(1)$ Standard Model gauge field
$Y_{\mu\nu}$	Field strength (curvature) of $Y_\mu$
$y_\nu$	Yukawa coupling of tau-neutrino
$y_{top}$	Yukawa coupling of top quark
$Z(\mathcal{A})$	Center of $\mathcal{A}$
$\zeta(z)$	Riemann zeta function
$\zeta_E(z)$	Epstein zeta function
$\zeta_b(z)$	Zeta function.

# Index

## A

- Abelian gauge theory, 4  
Algebra, 1–4, 9–17, 21, 23, 25, 26, 28, 31–33, 35, 36, 43–47, 49–52, 56–58, 60, 62, 66, 71, 73, 74, 76, 77, 80–86, 95–97, 102, 103, 106, 114, 115, 121–129, 132, 136–143, 145–152, 154, 155, 166, 169, 180–185, 187, 199, 204, 207, 208, 210, 217, 219–224, 227–229, 231, 233–235, 238, 255, 271, 272, 284, 287–292  
Algebra bundle, 4, 57, 151, 154, 199, 219–221, 223, 224, 227  
Algebra connection, 221–224  
Algebra of module homomorphisms, 125  
Almost-commutative manifold, 79, 179, 180, 183, 186, 188, 190, 191, 193, 196, 199, 203, 204, 206, 207, 210–212, 214, 217, 219, 229, 233, 239, 243, 245, 251, 254, 257, 262, 273, 274, 285  
Analytic order, 102–105, 115  
Anti-unitary operator, 31, 33, 37–39, 47, 58, 62, 81  
Asymptotic expansion, 4, 104–106, 109, 159, 160, 173, 174, 186, 190, 193  
Atiyah–Bott formula, 110, 113  
Atiyah–Singer index theorem, 227  
Automorphism, 40, 41, 44, 82, 83, 85, 121–124, 139, 140, 143, 145, 146, 148, 150, 152, 153, 180, 182, 217, 223, 230, 287–290, 300

## B

- Background field, 2, 294  
 $(b, B)$ -coboundaries, 97

© The Editor(s) (if applicable) and The Author(s) 2025

W. D. van Suijlekom, *Noncommutative Geometry and Particle Physics*,

Mathematical Physics Studies, <https://doi.org/10.1007/978-3-031-59120-4>

$(b, B)$ -cocycle, 97, 99, 100, 105, 106, 108–110, 168

$b$  (differential), 95, 96

$B$  (differential), 95, 96

Big desert, 260

Big desert, 260

Bimodule, 14, 16, 31, 33, 34, 46, 128

## C

Canonical triple, 77, 85, 89, 91, 101, 115, 179, 180, 193, 195, 210, 217

Center, 53, 122, 124, 145, 146, 152, 154, 182

Chern character, 115, 116, 175, 227

Chirality operator, 52, 54, 55, 77, 271

Christoffel symbols, 60–62, 66

Classification of finite real spectral triples, 3, 31, 35, 40, 47

Classification of finite spectral triples, 3, 25, 35, 45

Classification of irreducible geometries, 3, 45, 206

Clifford algebra, 32, 49–53, 55–58, 62, 71, 274, 287, 289, 290

Clifford algebra bundle, 56, 57

Clifford multiplication, 57, 61, 63

Cochain, 95, 96, 98–100, 108, 112, 167

Cohomology, 95, 97

Cohomology groups, 95–97

Commutant, 12, 33, 44, 46, 54

Commutant property, 32, 76, 125, 134, 145, 222

Commutative subalgebra, 124, 145, 146, 151, 154, 182

Compatible connection, 35

Complex algebra, 43, 181–183, 234  
 Complex involutive algebra, 145  
 Conjugate module, 34, 128  
 Connection, 2, 24, 25, 28, 34, 35, 59–62,  
     130–133, 150, 154, 171, 174, 184, 187,  
     191, 221, 224, 226, 227, 275  
 Connection one-form, 175, 221, 223–226  
 Coupling constant, 3, 4, 244, 245, 249, 254,  
     259–261, 269, 280–282  
 $C^*$ -algebra, 4, 13, 28, 73–75, 80, 84–86, 122,  
     143, 228  
 $C^*$ -bundle, 145–148, 150, 152–155, 180  
 $C^*$ -dynamical system, 287–289  
 $C(X)$ -algebra, 146, 150, 154  
 Cyclic cocycles, 91, 99, 166, 167  
 Cyclic cohomology, 95, 97, 99, 112, 114,  
     116  
 Cyclic vector, 33

**D**

Defining representation, 12, 17, 20, 53, 123,  
     186, 230, 238  
 De Rham current, 114  
 De Rham differential form, 79  
 Derivation, 1–4, 22, 24, 25, 54, 79, 81, 82,  
     99, 103, 124, 125, 130, 138, 148, 182,  
     186, 199, 203, 210, 229, 255, 268, 284,  
     298  
 Determinant, 216, 217, 234, 291  
 Differential one-form, 22, 23, 79, 123, 133  
 Dimension spectrum, 102, 110, 115, 159,  
     160  
 Dirac operator, 1–3, 22, 23, 26, 34, 35, 37,  
     39, 43, 49, 55, 59, 63, 64, 68, 69, 71,  
     75–78, 81–84, 110, 115, 134, 146, 155,  
     183–188, 191, 196, 197, 204, 207–210,  
     212, 220, 222, 225, 226, 230, 232, 237,  
     252, 253, 255, 257, 272, 274, 276, 277,  
     279, 284, 294, 297  
 Discrete metric, 19  
 Distance function, 56, 75, 85, 205  
 Divided difference, 22, 164–166, 171, 172,  
     176, 294–296  
 Dixmier-Douady class, 228

**E**

Effective action, 297  
 Electrodynamics, 4, 173, 203, 207, 211, 214,  
     229  
 Entropy, 292, 293  
 Epstein zeta function, 93, 114  
 Euler characteristic, 196

Even cochain, 98  
 Even cyclic cohomology, 112

**F**

Fermion doubling, 173, 255  
 Fermionic action, 2, 4, 157, 158, 173, 206,  
     208, 211–215, 249–252, 276  
 Finitely generated projective module, 126  
 Finite noncommutative space, 9, 31, 79, 121,  
     179, 184, 203, 284  
 Finite projective module, 227  
 Finite real noncommutative space, 31  
 Finite real spectral triple, 3, 31–35, 38, 40,  
     41, 43, 45–47, 179, 203, 230  
 Finite spectral triple, 3, 21–25, 27, 28, 31,  
     33, 35, 43, 45, 47, 174  
 Finite summability, 111, 115  
 First-order condition, 133, 134, 204, 217,  
     230, 231, 272, 273  
 Fredholm operator, 90, 92, 110

**G**

Gâteaux derivative, 162, 163, 174  
 Gauge bosons of the Standard Model, 245  
 Gauge field, 1–4, 121, 125, 132, 146, 150,  
     151, 157–159, 161, 179, 183–187, 196,  
     198, 206, 207, 210, 211, 260, 271, 273,  
     275, 277, 284, 294  
 Gauge fields on almost-commutative mani-  
     folds, 183  
 Gauge fields of the Standard Model, 4, 238,  
     239, 260, 277, 284, 306, 308, 309  
 Gauge group, 224  
 Gauge group of the Standard Model, 234  
 Gauge group on almost-commutative mani-  
     folds, 179, 180, 183, 186, 188, 190,  
     191, 199, 203, 206, 233, 306  
 Gauge invariant, 2, 4, 157, 158, 198, 245  
 Gauge Lie algebra, 124, 180  
 Gauge theory, 1–4, 79, 121, 145, 146, 151–  
     153, 199, 206, 207, 210, 219, 223, 224,  
     227, 233, 245, 298, 299  
 Gauge transformation, 2, 4, 132, 134, 137,  
     150, 185, 186, 226, 238, 248  
 Gelfand duality, 73–75, 84  
 Generalized Laplacian, 186, 188, 190, 191,  
     199, 226  
 Generations, 232, 233, 240, 250, 251, 253,  
     254, 259, 274, 275  
 Grand unified theories, 260, 284  
 Graph, 25–28, 35, 40, 41, 69, 71, 293, 295–  
     300

- Grassmann connection, 130  
 Grassmann integral, 215–217  
 Grassmann variables, 158, 214–217
- H**  
 Heat kernel expansion, 159, 160, 186  
 Hermitian  $*$ -algebra connection, 221, 223, 224  
 Hermitian structure, 129, 219, 220, 222, 227
- Higgs field, 1, 2, 4, 186, 199, 238, 243–247, 250, 251, 253–255, 257, 262, 271, 273, 275, 280, 284  
 Higgs mass, 3, 4, 258, 259, 264, 265, 268, 269, 271, 281, 282, 284, 285  
 Higgs mechanism, 217, 245, 255  
 Higgs potential, 243, 245–247  
 Hilbert bimodule, 15–18, 23, 28, 33, 34  
 Hilbert–Schmidt inner product, 32, 219–221
- Hilbert transform, 89  
 Hochschild coboundary, 95  
 Hochschild cocycle, 99, 167, 168  
 Hochschild cohomology, 95, 96, 167  
 Hypercharge, 236–238, 255
- I**  
 Idempotent, 126–128  
 Improper cocycle, 111  
 Inner automorphism, 121–123, 140, 143  
 Inner derivation, 125  
 Inner fluctuation, 121  
 Inner unitary equivalence, 119, 141  
 Involutive algebra, 9, 43, 136  
 Irreducible geometry, 3, 45, 47, 206, 255, 268, 271  
 Irreducible representation, 12, 13, 17, 26, 33, 35, 36, 45, 46, 53, 54, 74, 123, 281, 289, 291
- K**  
 Kaluza–Klein theory, 179, 199, 211  
 Kasparov product, 15, 16, 155  
 KMS, 287–289, 291, 293, 299  
 KO-dimension, 31, 32, 35, 38–41, 45, 47, 55, 58, 59, 76–78, 81, 82, 158, 183, 204, 206, 208–210, 217, 229, 233, 255, 268, 272  
 Krajewski diagram, 3, 35, 38, 40, 41, 43, 45–47, 199, 204, 208, 209, 217, 230, 231
- L**  
 $\Lambda$ -decorated graph, 35  
 Laplace–Stieltjes transform, 159, 160, 173, 192  
 Laplacian, 1, 59, 60, 63, 64, 186, 188, 190, 191, 199, 226  
 Leibniz rule, 24, 34, 59, 64, 108, 130, 132, 143, 169  
 Leptons of the Standard Model, 231, 254  
 Levi–Civita connection, 60, 61, 77, 191  
 Lichnerowicz formula, 63, 186, 188, 189, 194  
 Local index formula, 3, 89, 92, 94, 110, 113–115, 174  
 Localization, 3, 145, 148, 150, 151
- M**  
 Majorana mass, 252, 253, 255, 259, 262, 263, 265, 271  
 Majorana mass matrix, 232, 251  
 Mass matrix, 232, 251, 259, 282  
 Mass relations, 257  
 Matrix algebra, 2, 3, 9, 11, 13–18, 21, 23, 26–28, 32, 33, 44, 45, 47, 53, 73, 74, 92, 121, 141–143, 199, 219, 229, 288–290  
 McKean–Singer formula, 111, 115  
 Metric, 3, 18–21, 24, 57, 58, 60, 83, 204, 217
- Module, 14, 16, 17, 24–26, 28, 31, 34, 125–130, 133  
 Module homomorphism, 125  
 Morita equivalence, 3, 14, 18, 23, 25, 28, 33, 121, 125, 126, 128, 130, 133, 143  
 Morita self-equivalence, 125, 132, 183  
 Multiplicity, 26, 27, 33, 36, 39, 41, 46, 208, 209
- N**  
 Noncommutative Riemannian spin manifolds, 3, 59, 64, 73, 76, 84  
 Noncommutative torus, 73, 80–82, 85, 99, 100, 114, 115, 173
- O**  
 Odd cochain, 96, 98, 99  
 Odd cyclic cohomology, 97  
 One-loop correction, 287, 300  
 Opposite algebra, 32, 137  
 Order one condition, 32, 39, 40, 76, 209, 222, 272

Orientation axiom, 47  
 Outer automorphism, 121

## Q

Quarks of the Standard Model, 231, 254

## P

Pati–Salam model, 284  
 Periodic cyclic cohomology, 97, 99, 100, 167  
 Perturbative expansion, 159, 161, 173, 190, 193, 199, 287  
 Pfaffian, 216  
 Pontryagin class, 194  
 Powers–Rieffel projection, 114, 115  
 Product spectral triple, 28, 78  
 Projection, 89, 92, 93, 97, 99–101, 110, 113–115, 127, 128, 130, 151, 174, 253, 289, 291  
 Pseudodifferential operator, 103, 104

## Q

Quaternions, 43, 229, 233

## R

Real algebra, 43, 46, 53, 184, 217, 234  
 Real involutive algebra, 43  
 Real spectral triple, 3, 31–35, 38, 40, 41, 43, 45–47, 55, 59, 76–78, 80–82, 84, 85, 121, 123, 125, 131–133, 136, 137, 139, 140, 145, 151, 152, 154, 179, 203, 217, 221, 223, 224, 230, 273  
 Real structure, 3, 41, 44, 46, 76–78, 123, 133, 146, 221, 222  
 Real subalgebra, 43, 44, 53, 54  
 Reconstruction theorem, 21, 28, 85, 146, 223  
 Regular spectral triple, 102–104, 110, 115, 159  
 Renormalization group equation, 4, 257, 260, 262, 263, 268, 269, 282, 285  
 Renormalization group flow, 260, 263  
 Representation, 2, 11–22, 26, 31–33, 35, 36, 39, 44–47, 53, 54, 74, 79, 81, 83–85, 123, 133, 152, 172, 182, 185, 186, 198, 204, 224, 226, 230, 231, 235, 238, 271, 273, 274, 279, 281, 284, 289–291, 300

Residue cocycle, 109

Ricci tensor, 60

Riemannian curvature tensor, 60, 194  
 Riemannian metric, 55, 56  
 Riemannian spin geometry, 3, 55, 73, 75  
 Riemannian spin manifold, 1, 2, 22, 49, 55, 59, 63, 64, 71, 73, 75, 77, 79, 84, 85, 115, 146, 151, 179, 188, 193, 204, 220, 224  
 Riemann zeta function, 90, 300

## S

Scalar curvature, 63, 64, 188, 191, 243, 281  
 Scalar field, 2, 4, 179, 183, 187, 198, 205, 214, 217, 235, 237, 238, 273, 276, 279–281, 283, 284, 300  
 Scalar fields on almost-commutative manifolds, 183  
 Scalar fields of the Standard Model, 276  
 Second quantization, 287, 289, 299  
 Seeley–De Witt coefficients, 191, 198  
 Seesaw mechanism, 255, 259  
 Self-adjoint, 1, 2, 128, 131, 133, 139, 140, 288, 289, 291  
 Separating vector, 33, 45, 46  
 Serre–Swan Theorem, 227  
 Simple dimension spectrum, 102, 115, 159  
 Sobolev space, 102, 115  
 Spectral action, 2–4, 157–160, 162, 164, 166, 168, 171, 173, 174, 179, 186, 190, 191, 193, 195, 196, 198, 199, 206, 211, 214, 217, 223, 224, 226, 227, 239, 243–245, 255, 257, 274, 277, 281, 284, 287, 288, 293, 295, 298–300  
 Spectral action of the Standard Model, 3, 173, 239, 255, 257, 274, 277, 281, 284  
 Spectral action on almost-commutative manifolds, 190  
 Spectral flow, 174, 175  
 Spectral shift, 174, 175  
 Spectral triple, 1–3, 21–25, 27, 28, 31–35, 38, 40–43, 45–47, 49, 58, 73, 75–82, 84, 85, 89, 91, 94, 101–104, 110, 111, 115, 116, 121, 123–125, 130–133, 136, 137, 139, 140, 143, 145, 146, 151–154, 157–159, 167, 173, 174, 179, 182, 203, 209, 210, 217, 219–224, 226, 230, 233, 272, 273, 287  
 Spin connection, 59, 62, 63, 188, 189, 194  
 Spin manifold, 1–3, 22, 49, 55, 58, 59, 62–64, 71, 73, 75–77, 79, 83–85, 101, 115, 116, 146, 151, 179, 180, 188, 193, 199, 204, 217, 220, 223, 224  
 Spin<sup>c</sup> manifold, 58, 61, 62, 68, 69, 71, 85

Spinor, 2, 57, 58, 61, 62, 65, 66, 77, 81, 83, 92, 93, 146, 208, 212, 213, 215, 220, 226, 250, 253, 272, 275  
Spinor bundle, 2, 61, 62, 65, 77, 146, 220  
Spin structure, 49, 58, 59, 63, 85  
Spin<sup>c</sup> structure, 57  
Spontaneous symmetry breaking, 1, 3, 4, 254, 279  
Square-integrable spinor, 58  
Standard Model of particle physics, 1–4, 229, 255  
Strong interaction, 279  
Structure space, 12, 13, 17, 18, 21, 26, 28, 35, 74, 122, 204  
Symmetric operator, 19–22, 24, 66, 67, 82, 232

**T**

Taylor expansion, 162, 164, 166  
Taylor's formula, 162  
Tensor product (of modules), 14  
Tensor product (of spectral triples), 28  
Topological spectral action, 4, 157–159, 227  
  
Toric noncommutative manifolds, 3, 73, 113, 145, 151, 155  
Two-point space, 203, 204, 206, 217

**U**

Unification, 244, 245, 249  
Unimodularity, 181, 182, 225, 234, 236, 255  
  
Unitary equivalence, 12, 23, 25, 27, 33, 35, 41, 47, 123, 132, 143  
Unitary gauge, 255, 281  
Unitary operator, 2, 23, 33, 37, 79, 83, 290, 300

**V**

Vacuum expectation value, 246, 257, 262, 264  
Vacuum state, 246

**W**

Ward identities, 294, 295, 298, 300  
Weak interaction, 253  
Weitzenböck formula, 188  
Weyl tensor, 193, 275  
Winding number, 89, 91, 94

**Y**

Yang–Mills action, 223, 224, 226  
Yang–Mills field, 219  
Yang–Mills manifold, 183, 185, 198  
Yang–Mills theory, 199, 219, 223, 227  
Yukawa coupling, 251, 253, 259, 262–267  
Yukawa mass matrix, 232