Canadian Syncope Risk Score

adults (≥16 y) with syncope · casualty · 30-day serious outcome

CSRS −3 to 11

CRITERIA

Item	Points
Clinical evaluation	
Predisposition to vasovagal symptoms (warm crowded place, prolonged standing, fear, emotion, pain)	**−1**
History of heart disease (CAD, AF/flutter, CCF, valvular disease)	**+1**
Any SBP <90 or >180 mmHg in casualty	**+2**
Investigations	
Troponin >99th percentile	**+2**
Abnormal QRS axis (<−30° or >100°)	**+1**
QRS duration >130 ms	**+1**
Corrected QT >480 ms	**+2**
Diagnosis in casualty	
Vasovagal syncope	**−2**
Cardiac syncope	**+2**
Total	**−3 to 11**

30-DAY SERIOUS ADVERSE OUTCOME

- **≤−2** Very low (≈0.2%)
- **−1 to 0** Low (≈0.7%)
- **1–3** Medium (≈8%)
- **4–5** High (≈19%)
- **≥6** Very high (≈51%)

USE

- Very low / low: consider discharge with follow-up
- Medium–very high: monitor, admit / specialist input
- Serious outcome = death, arrhythmia, MI, structural heart disease, PE, haemorrhage, procedural intervention

DO NOT USE IF

- LOC >5 min, change in mental status, obvious seizure, head trauma causing LOC
- Serious diagnosis already found in casualty
- Not in the SA STGs

CSRS Thiruganasambandamoorthy V et al. Canadian Syncope Risk Score. CMAJ 2016;188:E289-98

Not in the SA STGs — all content from the international sources named above. ANDH · csrs · updated 2026-10-14