GRACE Score — ACS In-hospital & 6-month Mortality

adults with confirmed ACS (STEMI / NSTE-ACS) · ward / casualty

GRACE (in-hospital)

POINTS (IN-HOSPITAL NOMOGRAM)

Age (y)	Pts	HR (/min)	Pts	SBP (mmHg)	Pts	Creatinine µmol/L	Pts
<30	0	<50	0	<80	58	<35	1
30–39	8	50–69	3	80–99	53	35–70	4
40–49	25	70–89	9	100–119	43	71–105	7
50–59	41	90–109	15	120–139	34	106–140	10
60–69	58	110–149	24	140–159	24	141–176	13
70–79	75	150–199	38	160–199	10	177–353	21
80–89	91	≥200	46	≥200	0	≥354	28
≥90	100						

OTHER VARIABLES

Finding	Pts
Killip I — no heart failure	**0**
Killip II — crackles, S3, ↑JVP	**+20**
Killip III — pulmonary oedema	**+39**
Killip IV — cardiogenic shock	**+59**
Cardiac arrest at admission	**+39**
ST-segment deviation	**+28**
Elevated cardiac enzymes / troponin	**+14**

RISK CATEGORIES (TOTAL POINTS)

	Low	Intermediate	High
NSTE-ACS			
In-hospital death	≤108 (<1%)	109–140 (1–3%)	>140 (>3%)
6-month death	≤88 (<3%)	89–118 (3–8%)	>118 (>8%)
STEMI			
In-hospital death	≤125 (<2%)	126–154 (2–5%)	>154 (>5%)
6-month death	≤99 (<4.5%)	100–127 (4.5–11%)	>127 (>11%)

CONVERSION

Creatinine: **`µmol/L = mg/dL × 88.4`** · original bands in mg/dL (<0.40 … ≥4.0)

SA STG

UA/NSTEMI: TIMI and GRACE are well-validated risk scores; high-risk patients (incl. positive troponin) → discuss with cardiology for angiography ✚

USE

- NSTE-ACS: GRACE >140 → early invasive strategy (<24 h) — discuss with cardiology
- Score at presentation; Killip class from examination
- Prefer the online GRACE 2.0 calculator if available (continuous variables)

HOSP STG & EML Hospital Level Adults, 6th ed. 2024 (updated Mar 2026) — pp 3.12 · **GRACE** Granger CB et al. GRACE in-hospital mortality. Arch Intern Med 2003;163:2345-53