SOFA Score & qSOFA — Sepsis (Sepsis-3)

adults with suspected infection · casualty / ward / ICU

SOFA 0–24

SOFA (TABLE 23.6) ✚

System	0	1	2	3	4
PaO_2/FiO_2 (mmHg)	≥400	<400	<300	<200 + resp. support	<100 + resp. support
Platelets $\times 10^9$/L	≥150	<150	<100	<50	<20
Bilirubin µmol/L	<20	20–32	33–101	102–204	>204
Cardiovascular	MAP ≥70	MAP <70	Dopamine <5 or dobutamine (any)	Dopamine 5.1–15 or adrenaline ≤0.1	Dopamine >15 or adrenaline >0.1
GCS	15	13–14	10–12 ‡	6–9	<6
Creatinine µmol/L (or UO)	<110	110–170	171–299	300–440 or UO <500 mL/d	>440 or UO <200 mL/d

DEFINITIONS

SOFA ✚: `sum of 6 systems (0–4 each) = 0–24`

Sepsis ✚: `infection + acute ↑SOFA ≥2 (baseline 0 if unknown)`

Septic shock ✚: `vasopressors for MAP ≥65 AND lactate >2 mmol/L despite fluids`

qSOFA: `RR ≥22 + altered mentation + SBP ≤100 (1 each, 0–3)`

MAP: `(SBP + 2 × DBP) ÷ 3`

QSOFA (BEDSIDE SCREEN)

≥2 High risk of poor outcome — full SOFA, lactate, cultures, escalate

0–1 Does not exclude sepsis — keep reassessing

SA STG — FIRST HOUR

Balanced crystalloid 30 mL/kg over 3 h; reassess for fluid overload ✚

Target MAP >65 mmHg; lactate clearance >20% over 2 h ✚

Adrenaline 0.01–1.0 mcg/kg/min if hypotensive after fluids ✚

NOTES

- Catecholamine doses in mcg/kg/min for ≥1 h
- ‡ STG Table 23.6 prints GCS 10–11 for 2 points; Sepsis-3 uses 10–12 ‡ ✚
- STG table unit labels incomplete — values as Sepsis-3 (SI) ‡ ✚

HOSP STG & EML Hospital Level Adults, 6th ed. 2024 (updated Mar 2026) — pp 23.34, 23.35 · **SEPSIS3** Singer M et al. Sepsis-3. JAMA 2016;315:801-10; Seymour CW et al. JAMA 2016;315:762-74

✚ = SA STG / national guideline (pages listed above). Unmarked content: international source named above. ‡ = STG misprint or sources differ (see app). ANDH · sofa · updated 2026-10-14