Upper GI Bleed — Glasgow-Blatchford & Rockall

adults with haematemesis / melaena · casualty

GBS 0–23 · Rockall 0–7

GLASGOW-BLATCHFORD SCORE (PRE-ENDOSCOPY)

Admission finding	Points
Blood urea (mmol/L)	
6.5–7.9	+2
8.0–9.9	+3
10.0–24.9	+4
≥25	+6
Haemoglobin (g/dL) — men | women	
Men 12.0–12.9 | women 10.0–11.9	+1
Men 10.0–11.9	+3
Men or women <10.0	+6
Systolic BP (mmHg)	
100–109	+1
90–99	+2
<90	+3
Other markers	
Pulse ≥100/min	+1
Presentation with melaena	+1
Presentation with syncope	+2
Hepatic disease (history or clinical/lab evidence)	+2
Cardiac failure	+2
Total	**0–23**

GBS INTERPRETATION

- **0** Low risk — consider early discharge with outpatient endoscopy
- **1–5** Admit; risk of intervention rises with score
- **≥6** >50% need intervention (transfusion / endoscopy / surgery)

PRE-ENDOSCOPY (CLINICAL) ROCKALL

Variable	Pts
Age	
<60 y	0
60–79 y	+1
≥80 y	+2
Shock	
None (HR <100, SBP ≥100)	0
Tachycardia: HR ≥100, SBP ≥100	+1
Hypotension: SBP <100	+2
Comorbidity	
None major	0
Cardiac failure, IHD, any major comorbidity	+2
Renal failure, liver failure, disseminated malignancy	+3
Total	**0–7**

ROCKALL (PRE-ENDOSCOPY)

- **0** Low risk of death
- **≥1** Not low risk — full Rockall after endoscopy adds diagnosis & stigmata (max 11)

SA STG

- Variceal bleeding: endoscopic variceal ligation and/or immediate referral for advanced management ✚
- Alarm symptoms (haematemesis or melaena, anaemia, weight loss, dysphagia, chest pain, >60 y new dyspepsia) → endoscopy ✚

HOSP STG & EML Hospital Level Adults, 6th ed. 2024 (updated Mar 2026) — pp 1.11, 1.3 · **GBS** Blatchford O et al. Lancet 2000;356:1318-21 · **ROCKALL** Rockall TA et al. Gut 1996;38:316-21 · **NICEUGIB** NICE CG141 Acute upper GI bleeding in over 16s (2012, updated 2016)

✚ = SA STG / national guideline (pages listed above). Unmarked content: international source named above. ‡ = STG misprint or sources differ (see app). ANDH · ugib · updated 2026-10-14