

Table 1: Diversity estimates by normalized Shannon’s entropy, considering each of 26 nucleotide positions containing amino acid substitutions conserved at greater than 75 percent in the FNV strains analyzed, relative to FVV.

Nucleotide Position	FVV	FNV-IP	FNV-Yale	FNV-FC	FNV-NT
357	0.278	0.005	0.004	0.010	0.066
386	0.173	0.025	0.014	0.028	0.097
554	0.062	0.054	0.018	0.009	0.080
854	0.182	0.014	0.004	0.014	0.038
992	0.148	0.007	0.001	0.010	0.131
1134	0.057	0.025	0.009	0.011	0.094
1432	0.133	0.003	0.005	0.021	0.123
1572	0.038	0.147	0.140	0.147	0.184
1718	0.091	0.003	0.007	0.013	0.068
1965	0.039	0.021	0.020	0.020	0.024
2344	0.121	0.020	0.007	0.006	0.085
3752	0.153	0.029	0.018	0.009	0.005
3812	0.126	0.065	0.057	0.057	0.061
4358	0.097	0.028	0.010	0.009	0.003
4607	0.076	0.036	0.022	0.234	0.053
5145	0.180	0.033	0.023	0.017	0.014
6527	0.169	0.008	0.004	0.035	0.006
6843	0.133	0.005	0.004	0.033	0.016
6942	0.106	0.031	0.020	0.099	0.166
7171	0.049	0.010	0.021	0.017	0.020
7178	0.128	0.011	0.020	0.126	0.033
7380	0.253	0.041	0.038	0.158	0.144
7641	0.168	0.013	0.012	0.113	0.279
8409	0.140	0.007	0.014	0.180	0.035
8640	0.054	0.024	0.018	0.128	0.038
9227	0.211	0.001	0.006	0.367	0.012