

Table 1: Amino acid variants concordantly observed in both SA14-14-2/P1 and SA14-14-2/P2, relative to the newly sequenced SA14 strain.

			Nucleotide		Amino Acid		Variant Percent	
Position	Gene	Codon	Cons ^a	Var ^b	Cons	Var	SA14-14-2/P1	SA14-14-2/P2
948	M	66	C	U	L	F	1.026	0.8972
1116	E	47	A	G	N	D	22.87	36.18
1453		159	U	A	V	D	10.59	11.38
1512		179	A	G	K	E	36.48	42.28
1911		312	A	G	K	E	1.538	0.9815
2646	NS1	57	G	A	V	I	0.9662	0.386
3652		392	U	G	V	G ^c	0.9572	1.365
3669		398	C	U	L	F	6.045	8.313
3695		406	G	C	L	F	3.907	4.841
5311	NS3	235	U	C	V	A ^c	1.693	1.707
5781		392	G	A	D	N	0.6344	1.04
5835		410	G	A	D	N	11.18	8.654
7167	NS4B	86	G	A	V	M	2.13	3.166
7657		249	A	G	D	G	20.12	15.99
7774	NS5	24	U	C	I	T	1.901	1.767
8791		372	C	U	A	V	1.077	2.365
9292		539	G	C	G	A	0.736	1.795
10069		798	C	U	T	I	0.8492	0.8072
10092		806	G	C	G	R	1.096	0.7716
10329		906	G	A	V	I	2.132	1.709

^a Consensus nucleotide or residue in the newly computed consensus sequence.

^b Variant nucleotide or residue in the alignment, which is observed below the consensus frequency.

^c Variant residue is a revertant to wild-type SA14 genotype.