

# **The Knuth-Yao Quadrangle Inequality Speedup is a Consequence of Total Monotonicity**

Wolfgang W. Bein    (University of Nevada)

Mordecai J. Golin    (Hong Kong UST)

Lawrence L. Larmore    (University of Nevada)

Yan Zhang    (Hong Kong UST)

# Motivation

- Nothing new: material here goes back 20-30 years.
- There are two classic cookbooks
  - Dynamic Programming Speedups in the literature:  
Knuth-Yao technique & SMAWK algorithm.
- They “feel” similar. Are they related?
- Knuth-Yao predates online algorithms.
  - Can the KY speedup be maintained online?
- Answers to the two questions turned out to be related.
- Note: major confusion arises in the analysis because certain essential terms, e.g., quadrangle-inequality, monotone and online-algorithm have been used in very different ways in the two techniques’ literature.

# Outline

- Background

- Kunth-Yao (KY) Quadrangle Inequality (QI) Speedup
- SMAWK Algorithm for finding  
Row Minima of Totally Monotone (TM) Matrices

- The  $D^d$  Decomposition

A transformation from QI to TM such that  
SMAWK solves KY problem as quickly as KY.

- The  $L^m$  and  $R^m$  Decompositions

Another transformation from QI to TM that  
(1) implies KY speedup and (2) enables online solution.

- Extensions

Applying the technique to known generalizations of KY.

# Outline

## ● Background

- Kunth-Yao (KY) Quadrangle Inequality (QI) Speedup
- SMAWK Algorithm for finding  
Row Minima of Totally Monotone (TM) Matrices

## ● The $D^d$ Decomposition

A transformation from QI to TM such that  
SMAWK solves KY problem as quickly as KY.

## ● The $L^m$ and $R^m$ Decompositions

Another transformation from QI to TM that  
(1) implies KY speedup and (2) enables online solution.

## ● Extensions

Applying the technique to known generalizations of KY.

# Background

# Background

- Kunth-Yao Quadrangle Inequality Speedup

# Background

- Kunth-Yao Quadrangle Inequality Speedup
  - D. E. Knuth (1971) and F. F. Yao (1980,1982)

# Background

- Kunth-Yao Quadrangle Inequality Speedup
  - D. E. Knuth (1971) and F. F. Yao (1980,1982)
  - $\Theta(n)$  speedup:  $O(n^3)$  down to  $O(n^2)$



# Background

- Kunth-Yao Quadrangle Inequality Speedup
  - D. E. Knuth (1971) and F. F. Yao (1980,1982)
  - $\Theta(n)$  speedup:  $O(n^3)$  down to  $O(n^2)$
- SMAWK Algorithm for finding  
Row Minima of Totally Monotone Matrices

# Background

- Kunth-Yao Quadrangle Inequality Speedup
  - D. E. Knuth (1971) and F. F. Yao (1980,1982)
  - $\Theta(n)$  speedup:  $O(n^3)$  down to  $O(n^2)$
- SMAWK Algorithm for finding  
Row Minima of Totally Monotone Matrices
  - A. Aggarwal, M. M. Klawe, S. Moran, P. Shor, R. Wilber (1986)

# Background

- Kunth-Yao Quadrangle Inequality Speedup
  - D. E. Knuth (1971) and F. F. Yao (1980,1982)
  - $\Theta(n)$  speedup:  $O(n^3)$  down to  $O(n^2)$
- SMAWK Algorithm for finding Row Minima of Totally Monotone Matrices
  - A. Aggarwal, M. M. Klawe, S. Moran, P. Shor, R. Wilber (1986)
  - $\Theta(n)$  speedup:  $O(n^2)$  down to  $O(n)$

# Background

- Kunth-Yao Quadrangle Inequality Speedup
  - D. E. Knuth (1971) and F. F. Yao (1980,1982)
  - $\Theta(n)$  speedup:  $O(n^3)$  down to  $O(n^2)$
- SMAWK Algorithm for finding Row Minima of Totally Monotone Matrices
  - A. Aggarwal, M. M. Klawe, S. Moran, P. Shor, R. Wilber (1986)
  - $\Theta(n)$  speedup:  $O(n^2)$  down to  $O(n)$
- Both techniques are often used to speed up DPs.

# Background

- Kunth-Yao Quadrangle Inequality Speedup
  - D. E. Knuth (1971) and F. F. Yao (1980,1982)
  - $\Theta(n)$  speedup:  $O(n^3)$  down to  $O(n^2)$
- SMAWK Algorithm for finding Row Minima of Totally Monotone Matrices
  - A. Aggarwal, M. M. Klawe, S. Moran, P. Shor, R. Wilber (1986)
  - $\Theta(n)$  speedup:  $O(n^2)$  down to  $O(n)$
- Both techniques are often used to speed up DPs.
- How are the two techniques related?

# Quadrangle Inequality

# Quadrangle Inequality

- Original Motivation

Computing **Optimal Binary Search Trees (Optimal BST)**

[Gilbert and Moore (1959)]

# Quadrangle Inequality

- Original Motivation  
Computing **Optimal Binary Search Trees (Optimal BST)**  
[Gilbert and Moore (1959)]
- Optimal BST
  - Construct a search tree for  $n$  keys



# Quadrangle Inequality

- Original Motivation  
Computing **Optimal Binary Search Trees (Optimal BST)**  
[Gilbert and Moore (1959)]
- Optimal BST
  - Construct a search tree for  $n$  keys
  - $n$  internal nodes corresponds to successful search  
 $p_l$ , ( $l = 1 \dots n$ ) is the weight that **search-key = Key <sub>$l$</sub>**

# Quadrangle Inequality

- Original Motivation

Computing **Optimal Binary Search Trees (Optimal BST)**

[Gilbert and Moore (1959)]

- Optimal BST

- Construct a search tree for  $n$  keys

- $n$  internal nodes corresponds to successful search

$p_l, (l = 1 \dots n)$  is the weight that **search-key = Key <sub>$l$</sub>**

- $n + 1$  external nodes corresponds to unsuccessful search

$q_l, (l = 0 \dots n)$  is the weight that **Key <sub>$l$</sub>  < search-key < Key <sub>$l+1$</sub>**

# Quadrangle Inequality

## ● Original Motivation

Computing **Optimal Binary Search Trees (Optimal BST)**

[Gilbert and Moore (1959)]

## ● Optimal BST

- Construct a search tree for  $n$  keys

- $n$  internal nodes corresponds to successful search

$p_l, (l = 1 \dots n)$  is the weight that **search-key = Key <sub>$l$</sub>**

- $n + 1$  external nodes corresponds to unsuccessful search

$q_l, (l = 0 \dots n)$  is the weight that **Key <sub>$l$</sub>  < search-key < Key <sub>$l+1$</sub>**

- Minimize the number of comparisons

$$\sum_{1 \leq l \leq n} p_l \cdot (1 + \underbrace{d(p_l)}_{\text{depth}}) + \sum_{0 \leq l \leq n} q_l \cdot \underbrace{d(q_l)}_{\text{depth}}$$

# Optimal BST

• Minimize  $\sum_{1 \leq l \leq n} p_l \cdot (1 + d(p_l)) + \sum_{0 \leq l \leq n} q_l \cdot d(q_l)$

# Optimal BST

- Minimize  $\sum_{1 \leq l \leq n} p_l \cdot (1 + d(p_l)) + \sum_{0 \leq l \leq n} q_l \cdot d(q_l)$
- An example

# Optimal BST

• Minimize  $\sum_{1 \leq l \leq n} p_l \cdot (1 + d(p_l)) + \sum_{0 \leq l \leq n} q_l \cdot d(q_l)$

• An example


$$n = 2 \quad p = (19, 12), \quad q = (36, 20, 11)$$

# Optimal BST

• Minimize  $\sum_{1 \leq l \leq n} p_l \cdot (1 + d(p_l)) + \sum_{0 \leq l \leq n} q_l \cdot d(q_l)$

• An example

$n = 2$       $p = (19, 12), \quad q = (36, 20, 11)$




# Optimal BST

• Minimize  $\sum_{1 \leq l \leq n} p_l \cdot (1 + d(p_l)) + \sum_{0 \leq l \leq n} q_l \cdot d(q_l)$

• An example

$n = 2$       $p = (19, 12),$       $q = (36, 20, 11)$






# Optimal BST

● Minimize  $\sum_{1 \leq l \leq n} p_l \cdot (1 + d(p_l)) + \sum_{0 \leq l \leq n} q_l \cdot d(q_l)$

● An example

$n = 2$       $p = (19, 12),$       $q = (36, 20, 11)$





Cost = 141

# Optimal BST

• Minimize  $\sum_{1 \leq l \leq n} p_l \cdot (1 + d(p_l)) + \sum_{0 \leq l \leq n} q_l \cdot d(q_l)$

• An example

$n = 2$       $p = (19, 12)$ ,      $q = (36, 20, 11)$





Cost = 141

# Optimal BST

• Minimize  $\sum_{1 \leq l \leq n} p_l \cdot (1 + d(p_l)) + \sum_{0 \leq l \leq n} q_l \cdot d(q_l)$

• An example

$n = 2$      $p = (19, 12)$ ,     $q = (36, 20, 11)$




Cost = 141

# Optimal BST


• Minimize  $\sum_{1 \leq l \leq n} p_l \cdot (1 + d(p_l)) + \sum_{0 \leq l \leq n} q_l \cdot d(q_l)$

• An example

$n = 2$      $p = (19, 12)$ ,     $q = (36, 20, 11)$



Cost = 141



Cost = 173

# Optimal BST

- Solution: Dynamic Programming (DP)


# Optimal BST

- Solution: Dynamic Programming (DP)
  - $B_{i,j}$  the optimal BST for the subproblem  $\text{Key}_{i+1}, \dots, \text{Key}_j$

# Optimal BST

- Solution: Dynamic Programming (DP)
  - $B_{i,j}$  the optimal BST for the subproblem  $\text{Key}_{i+1}, \dots, \text{Key}_j$
  - DP recurrence

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$



# Optimal BST

- DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$



# Optimal BST

## ● DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

	0	1	2	3	4	5	6
0	0						
1		0					
2			0				
3				0			
4					0		
5						0	
6							0

$B_{i,j}$  depends  
on the entries  
to the left and  
below.

# Optimal BST

- DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- An example

$$n = 6 \quad p = (88, 21, 19, 12, 14, 18) \quad q = (53, 89, 36, 20, 11, 19, 15)$$

# Optimal BST

- DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- An example

$n = 6$      $p = (88, 21, 19, 12, 14, 18)$      $q = (53, 89, 36, 20, 11, 19, 15)$

	0	1	2	3	4	5	6
0	0						
1		0					
2			0				
3				0			
4					0		
5						0	
6							0

# Optimal BST

## ● DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

## ● An example

$n = 6$      $p = (88, 21, 19, 12, 14, 18)$      $q = (53, 89, 36, 20, 11, 19, 15)$

	0	1	2	3	4	5	6
0	0	230					
1		0	146				
2			0	75			
3				0	43		
4					0	44	
5						0	52
6							0

# Optimal BST

## ● DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

## ● An example

$n = 6$      $p = (88, 21, 19, 12, 14, 18)$      $q = (53, 89, 36, 20, 11, 19, 15)$

	0	1	2	3	4	5	6
0	0	230	433				
1		0	146	260			
2			0	75	141		
3				0	43	119	
4					0	44	121
5						0	52
6							0

# Optimal BST

## ● DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

## ● An example

$n = 6$       $p = (88, 21, 19, 12, 14, 18)$       $q = (53, 89, 36, 20, 11, 19, 15)$

	0	1	2	3	4	5	6
0	0	230	433	586			
1		0	146	260	349		
2			0	75	141	250	
3				0	43	119	204
4					0	44	121
5						0	52
6							0

# Optimal BST

## ● DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

## ● An example

$n = 6$      $p = (88, 21, 19, 12, 14, 18)$      $q = (53, 89, 36, 20, 11, 19, 15)$

	0	1	2	3	4	5	6
0	0	230	433	586	698		
1		0	146	260	349	491	
2			0	75	141	250	357
3				0	43	119	204
4					0	44	121
5						0	52
6							0

# Optimal BST

## ● DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

## ● An example

$n = 6$      $p = (88, 21, 19, 12, 14, 18)$      $q = (53, 89, 36, 20, 11, 19, 15)$

	0	1	2	3	4	5	6
0	0	230	433	586	698	862	
1		0	146	260	349	491	624
2			0	75	141	250	357
3				0	43	119	204
4					0	44	121
5						0	52
6							0



# Optimal BST

- DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- An example

$n = 6$       $p = (88, 21, 19, 12, 14, 18)$       $q = (53, 89, 36, 20, 11, 19, 15)$

	0	1	2	3	4	5	6
0	0	230	433	586	698	862	1002
1		0	146	260	349	491	624
2			0	75	141	250	357
3				0	43	119	204
4					0	44	121
5						0	52
6							0

# Optimal BST

- DP: Straightforward Calculation

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- An example

$n = 6$       $p = (88, 21, 19, 12, 14, 18)$       $q = (53, 89, 36, 20, 11, 19, 15)$

	0	1	2	3	4	5	6
0	0	230	433	586	698	862	1002
1		0	146	260	349	491	624
2			0	75	141	250	357
3				0	43	119	204
4					0	44	121
5						0	52
6							0

# Optimal BST

● Naive:  $O(n^3) = \sum_{i=1}^n \sum_{j=i}^n \Theta(j - i)$

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

# Optimal BST

- Naive:  $O(n^3) = \sum_{i=1}^n \sum_{j=i}^n \Theta(j - i)$

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- Speedup:  $O(n^2)$  [Knuth (1971)]

# Optimal BST

- Naive:  $O(n^3) = \sum_{i=1}^n \sum_{j=i}^n \Theta(j - i)$

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- Speedup:  $O(n^2)$  [Knuth (1971)]

- $K_B(i, j)$  the largest index  $t$  that achieves the minimum.

# Optimal BST

- Naive:  $O(n^3) = \sum_{i=1}^n \sum_{j=i}^n \Theta(j - i)$

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- Speedup:  $O(n^2)$  [Knuth (1971)]

- $K_B(i, j)$  the largest index  $t$  that achieves the minimum.

- Theorem in [Knuth (1971)]

$$K_B(i, j) \leq K_B(i, j + 1) \leq K_B(i + 1, j + 1)$$

# Optimal BST

● Naive:  $O(n^3) = \sum_{i=1}^n \sum_{j=i}^n \Theta(j - i)$

$$B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

● Speedup:  $O(n^2)$  [Knuth (1971)]

●  $K_B(i, j)$  the largest index  $t$  that achieves the minimum.

● Theorem in [Knuth (1971)]

$$K_B(i, j) \leq K_B(i, j + 1) \leq K_B(i + 1, j + 1)$$

	$i$	$i + 1$
$j$	$K_B(i, j)$	$K_B(i, j + 1)$
$j + 1$		$K_B(i + 1, j + 1)$

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$



# Optimal BST

- Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

- The index table

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0					
1			1				
2				2			
3					3		
4						4	
5							5
6							

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0				
1			1				
2				2			
3					3		
4						4	
5							5
6							

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0				
1			1	1			
2				2			
3					3		
4						4	
5							5
6							

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0				
1			1	1			
2				2	2		
3					3		
4						4	
5							5
6							

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0				
1			1	1			
2				2	2		
3					3	4	
4						4	
5							5
6							

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0				
1			1	1			
2				2	2		
3					3	4	
4						4	5
5							5
6							

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0				
1			1	1			
2				2	2		
3					3	4	
4						4	5
5							5
6							



# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0	0			
1			1	1	1		
2				2	2	2	
3					3	4	4
4						4	5
5							5
6							

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0	0	0		
1			1	1	1	1	
2				2	2	2	4
3					3	4	4
4						4	5
5							5
6							

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0	0	0	1	
1			1	1	1	1	2
2				2	2	2	4
3					3	4	4
4						4	5
5							5
6							

# Optimal BST

● Speedup:  $B_{i,j} = \sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

$$K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$$

● The index table

	0	1	2	3	4	5	6
0		0	0	0	0	1	1
1			1	1	1	1	2
2				2	2	2	4
3					3	4	4
4						4	5
5							5
6							

# Optimal BST

● Speedup:

●  $K_B(i, j) \leq K_B(i, j + 1) \leq K_B(i + 1, j + 1)$

# Optimal BST

## ● Speedup:

●  $K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$

● Each diagonal  $j - i = d$

$$\begin{aligned} O(n) &= \sum_{i=1}^{n-d} (K_B(i+1, i+d) - K_B(i, i+d-1)) \\ &= K_B(n-d+1, n) - K_B(1, d) \end{aligned}$$

# Optimal BST

## Speedup:

•  $K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$

• Each diagonal  $j - i = d$

$$\begin{aligned} O(n) &= \sum_{i=1}^{n-d} (K_B(i+1, i+d) - K_B(i, i+d-1)) \\ &= K_B(n-d+1, n) - K_B(1, d) \end{aligned}$$

•  $O(n^2)$  total work over all  $n$  diagonals.

# Quadrangle Inequality



# Quadrangle Inequality

- Definition [Yao (1980, 1982)]

# Quadrangle Inequality

● Definition [Yao (1980, 1982)]

● Function  $f(i, j)$ ,  $(0 \leq i \leq j \leq n)$

satisfies a **Quadrangle Inequality (QI)**, if  $\forall i \leq i' \leq j \leq j'$

$$f(i, j) + f(i', j') \leq f(i', j) + f(i, j')$$


# Quadrangle Inequality

● Definition [Yao (1980, 1982)]

● Function  $f(i, j)$ ,  $(0 \leq i \leq j \leq n)$

satisfies a **Quadrangle Inequality (QI)**, if  $\forall i \leq i' \leq j \leq j'$

$$f(i, j) + f(i', j') \leq f(i', j) + f(i, j')$$



# Quadrangle Inequality

- Definition [Yao (1980, 1982)]

- Function  $f(i, j)$ ,  $(0 \leq i \leq j \leq n)$

satisfies a **Quadrangle Inequality (QI)**, if  $\forall i \leq i' \leq j \leq j'$

$$f(i, j) + f(i', j') \leq f(i', j) + f(i, j')$$

- Function  $f(i, j)$ ,  $(0 \leq i \leq j \leq n)$

is **Monotone over the integer lattice (MIL)**, if  $\forall [i, j] \subseteq [i', j']$

$$f(i, j) \leq f(i', j')$$

# Speedup using Quadrangle Inequality

$$B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

# Speedup using Quadrangle Inequality

$$B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- Lemmas from [Yao (1980)]

# Speedup using Quadrangle Inequality

$$B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- Lemmas from [Yao (1980)]
  - (A) If  $w(i,j)$  satisfies **QI** and is **MIL**,  
 $\Rightarrow B_{i,j}$  satisfies QI.

# Speedup using Quadrangle Inequality

$$B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

● Lemmas from [Yao (1980)]

● (A) If  $w(i,j)$  satisfies **QI** and is **MIL**,  
 $\Rightarrow B_{i,j}$  satisfies **QI**.

● (B) If  $B_{i,j}$  satisfies **QI**,  
 $\Rightarrow K_B(i,j) \leq K_B(i,j+1) \leq K_B(i+1,j+1)$



# Speedup using Quadrangle Inequality

$$B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

● Lemmas from [Yao (1980)]

● (A) If  $w(i,j)$  satisfies **QI** and is **MIL**,  
 $\Rightarrow B_{i,j}$  satisfies **QI**.

● (B) If  $B_{i,j}$  satisfies **QI**,  
 $\Rightarrow K_B(i,j) \leq K_B(i,j+1) \leq K_B(i+1,j+1)$

● In optimal BST problem,

$$B_{i,j} = \underbrace{\sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l}_{w(i,j)} + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

# Speedup using Quadrangle Inequality

$$B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

● Lemmas from [Yao (1980)]

● (A) If  $w(i,j)$  satisfies **QI** and is **MIL**,  
 $\Rightarrow B_{i,j}$  satisfies **QI**.

● (B) If  $B_{i,j}$  satisfies **QI**,  
 $\Rightarrow K_B(i,j) \leq K_B(i,j+1) \leq K_B(i+1,j+1)$

● In optimal BST problem,

$$B_{i,j} = \underbrace{\sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l}_{w(i,j)} + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

● Optimal BST  $w(i,j)$  satisfies **QI** as equality and is **MIL**.

# Speedup using Quadrangle Inequality

$$B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

## ● Lemmas from [Yao (1980)]

● (A) If  $w(i,j)$  satisfies **QI** and is **MIL**,  
 $\Rightarrow B_{i,j}$  satisfies **QI**.

● (B) If  $B_{i,j}$  satisfies **QI**,  
 $\Rightarrow K_B(i,j) \leq K_B(i,j+1) \leq K_B(i+1,j+1)$

## ● In optimal BST problem,

$$B_{i,j} = \underbrace{\sum_{l=i+1}^j p_l + \sum_{l=i}^j q_l}_{w(i,j)} + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

● Optimal BST  $w(i,j)$  satisfies **QI** as equality and is **MIL**.

●  $\Rightarrow$  exactly Knuth's result.

# Online Problem

# Online Problem

- Definition: Two-sided online problem

# Online Problem

- Definition: Two-sided online problem
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$

# Online Problem

- Definition: Two-sided online problem
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
  - Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .

# Online Problem

- Definition: Two-sided online problem
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
  - Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .
- An example



# Online Problem


## ● Definition: Two-sided online problem

- Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
- Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .

## ● An example

$$p = ( \quad 19, 12, 14 \quad ) \quad q = ( \quad 36, 20, 11, 19 \quad )$$

	1	2	3	4	5	6
1						
2		0	75	141	250	
3			0	43	119	
4				0	44	
5					0	
6						



# Online Problem


## Definition: Two-sided online problem

- Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
- Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .

## An example

$p = (19, 12, 14, 18)$     $q = (36, 20, 11, 19, 15)$

	1	2	3	4	5	6
1						
2		0	75	141	250	357
3			0	43	119	204
4				0	44	121
5					0	52
6						0



# Online Problem


## Definition: Two-sided online problem

- Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
- Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .

## An example

$$p = (21, 19, 12, 14, 18) \quad q = (89, 36, 20, 11, 19, 15)$$

	1	2	3	4	5	6
1	0	146	260	349	491	624
2		0	75	141	250	357
3			0	43	119	204
4				0	44	121
5					0	52
6						0



# Outline

## ● Background

- Kunth-Yao (KY) Quadrangle Inequality (QI) Speedup
- SMAWK Algorithm for finding  
Row Minima of Totally Monotone (TM) Matrices

## ● The $D^d$ Decomposition

A transformation from QI to TM such that  
SMAWK solves KY problem as quickly as KY.

## ● The $L^m$ and $R^m$ Decompositions

Another transformation from QI to TM that  
(1) implies KY speedup and (2) enables online solution.

## ● Extensions

Applying the technique to known generalizations of KY.

# Totally Monotone Matrices

- Definition

# Totally Monotone Matrices

● Definition      $M$  is an  $m \times n$  matrix

# Totally Monotone Matrices

- Definition  $M$  is an  $m \times n$  matrix
- $RM_M(i)$  is **index** of rightmost minimum item of row  $i$  of  $M$ .

# Totally Monotone Matrices

- Definition  $M$  is an  $m \times n$  matrix
  - $RM_M(i)$  is **index** of rightmost minimum item of row  $i$  of  $M$ .
  - $M$  is **Monotone** if  $\forall i \leq i', \quad RM_M(i) \leq RM_M(i')$ .



# Totally Monotone Matrices

● Definition  $M$  is an  $m \times n$  matrix

●  $RM_M(i)$  is **index** of rightmost minimum item of row  $i$  of  $M$ .

●  $M$  is **Monotone** if  $\forall i \leq i', \quad RM_M(i) \leq RM_M(i')$ .

7	2	4	3	8	9
5	1	5	1	6	5
7	1	2	0	3	1
9	4	5	1	3	2
8	4	5	3	4	3
9	6	7	5	6	5

$$RM_M(1) = 2$$

$$RM_M(2) = 4$$

$$RM_M(3) = 4$$

$$RM_M(4) = 4$$

$$RM_M(5) = 6$$

$$RM_M(6) = 6$$

# Totally Monotone Matrices

## ● Definition (Cond.)

● A  $2 \times 2$  **Monotone** matrix

2	4
4	5

2	3
5	3

7	1
2	2

# Totally Monotone Matrices

## ● Definition (Cond.)

● A  $2 \times 2$  **Monotone** matrix

2	4
4	5

2	3
5	3

7	1
2	2

● An  $m \times n$  matrix  $M$  is **Totally Monotone** (TM)  
if every  $2 \times 2$  submatrix is **Monotone**.

(submatrix: not necessarily contiguous in the original matrix)

# Totally Monotone Matrices

## ● Definition (Cond.)

- A  $2 \times 2$  **Monotone** matrix

2	4
4	5

2	3
5	3

7	1
2	2

- An  $m \times n$  matrix  $M$  is **Totally Monotone** (TM)  
if every  $2 \times 2$  submatrix is **Monotone**.  
(submatrix: not necessarily contiguous in the original matrix)

## ● Property

$M$  is **Totally Monotone**  $\Rightarrow M$  is **Monotone**

# Totally Monotone Matrices

## ● Definition (Cond.)

- A  $2 \times 2$  **Monotone** matrix

2	4
4	5

2	3
5	3

7	1
2	2

- An  $m \times n$  matrix  $M$  is **Totally Monotone** (TM)  
if every  $2 \times 2$  submatrix is **Monotone**.

(submatrix: not necessarily contiguous in the original matrix)

- Property

$M$  is **Totally Monotone**  $\Rightarrow M$  is **Monotone**

$M$  is **Totally Monotone**  $\nLeftarrow M$  is **Monotone**

# SMAWK Algorithm

# SMAWK Algorithm

## ● Motivation

Find all  $m$  row minima of an **implicitly** given  $m \times n$  matrix  $M$

# SMAWK Algorithm

- Motivation

Find all  $m$  row minima of an **implicitly** given  $m \times n$  matrix  $M$

- Naive Algorithm:  $O(mn)$



# SMAWK Algorithm

- Motivation  
Find all  $m$  row minima of an **implicitly** given  $m \times n$  matrix  $M$
- Naive Algorithm:  $O(mn)$
- **SMAWK** Algorithm  
[Aggarwal, Klawe, Moran, Shor, Wilber (1986)]

# SMAWK Algorithm

- Motivation
  - Find all  $m$  row minima of an **implicitly** given  $m \times n$  matrix  $M$
- Naive Algorithm:  $O(mn)$
- SMAWK Algorithm
  - [Aggarwal, Klawe, Moran, Shor, Wilber (1986)]
  - If  $M$  is **Totally Monotone**,
    - all  $m$  row minima can be found in  $O(m + n)$  time.

# SMAWK Algorithm

- Motivation

Find all  $m$  row minima of an **implicitly** given  $m \times n$  matrix  $M$

- Naive Algorithm:  $O(mn)$

- SMAWK Algorithm

[Aggarwal, Klawe, Moran, Shor, Wilber (1986)]

- If  $M$  is **Totally Monotone**,  
all  $m$  row minima can be found in  $O(m + n)$  time.

- Usually  $m = \Theta(n)$   
 $\Theta(n)$  speedup:  $O(n^2)$  down to  $O(n)$ .

# SMAWK Algorithm

- Motivation
  - Find all  $m$  row minima of an **implicitly** given  $m \times n$  matrix  $M$
- Naive Algorithm:  $O(mn)$
- SMAWK Algorithm
  - [Aggarwal, Klawe, Moran, Shor, Wilber (1986)]
  - If  $M$  is **Totally Monotone**,
    - all  $m$  row minima can be found in  $O(m + n)$  time.
  - Usually  $m = \Theta(n)$ 
    - $\Theta(n)$  speedup:  $O(n^2)$  down to  $O(n)$ .
- SMAWK was culmination of decade(s) of work on similar problems; speedups using convexity and concavity.

# SMAWK Algorithm

- Motivation
  - Find all  $m$  row minima of an **implicitly** given  $m \times n$  matrix  $M$
- Naive Algorithm:  $O(mn)$
- SMAWK Algorithm
  - [Aggarwal, Klawe, Moran, Shor, Wilber (1986)]
  - If  $M$  is **Totally Monotone**,
    - all  $m$  row minima can be found in  $O(m + n)$  time.
  - Usually  $m = \Theta(n)$ 
    - $\Theta(n)$  speedup:  $O(n^2)$  down to  $O(n)$ .
- SMAWK was culmination of decade(s) of work on similar problems; speedups using convexity and concavity.
- Has been used to speed up many DP problems, e.g., computational geometry, bioinformatics,  $k$ -center on a line, etc.

# The Monge Property

# The Monge Property

- Motivation

TM property is often established via Monge property.

# The Monge Property

- Motivation

TM property is often established via Monge property.

- Definition

An  $m \times n$  matrix  $M$  is **Monge** if  $\forall i \leq i'$  and  $\forall j \leq j'$

$$M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$



# The Monge Property

## Quadrangle Inequality

Function  $f(i, j)$

$$\forall i \leq i' \leq j \leq j'$$

$$f(i, j) + f(i', j') \leq f(i', j) + f(i, j')$$

## Monge

Matrix  $M$

$$\forall i \leq i' \text{ and } \forall j \leq j'$$

$$M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

# The Monge Property

## Quadrangle Inequality

Function  $f(i, j)$

$$\forall i \leq i' \leq j \leq j'$$

$$f(i, j) + f(i', j') \leq f(i', j) + f(i, j')$$

## Monge

Matrix  $M$

$$\forall i \leq i' \text{ and } \forall j \leq j'$$

$$M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

## ● QI vs. Monge

# The Monge Property

## Quadrangle Inequality

Function  $f(i, j)$

$$\forall i \leq i' \leq j \leq j'$$

$$f(i, j) + f(i', j') \leq f(i', j) + f(i, j')$$

## Monge

Matrix  $M$

$$\forall i \leq i' \text{ and } \forall j \leq j'$$

$$M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

## ● QI vs. Monge

- Different names for same type of inequality.

# The Monge Property

## Quadrangle Inequality

Function  $f(i, j)$

$$\forall i \leq i' \leq j \leq j'$$

$$f(i, j) + f(i', j') \leq f(i', j) + f(i, j')$$

## Monge

Matrix  $M$

$$\forall i \leq i' \text{ and } \forall j \leq j'$$

$$M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

## ● QI vs. Monge

- Different names for same type of inequality.
- Used differently in literature.

# The Monge Property

## Quadrangle Inequality

Function  $f(i, j)$

$$\forall i \leq i' \leq j \leq j'$$

$$f(i, j) + f(i', j') \leq f(i', j) + f(i, j')$$

## Monge

Matrix  $M$

$$\forall i \leq i' \text{ and } \forall j \leq j'$$

$$M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

## ● QI vs. Monge

- Different names for same type of inequality.
- Used differently in literature.
  - QI:  $f(i, j)$  is function to be calculated.
  - Monge:  $M_{i,j}$  implicitly given.

# The Monge Property

## Quadrangle Inequality

Function  $f(i, j)$

$$\forall i \leq i' \leq j \leq j'$$

$$f(i, j) + f(i', j') \leq f(i', j) + f(i, j')$$

## Monge

Matrix  $M$

$$\forall i \leq i' \text{ and } \forall j \leq j'$$

$$M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

## ● QI vs. Monge

- Different names for same type of inequality.
- Used differently in literature.
- QI:  $f(i, j)$  is function to be calculated.  
Need all  $f(i, j)$  entries.
- Monge:  $M_{i,j}$  implicitly given.  
Only need the row minima, but not other entries.

# Monge Property

$$\forall i \leq i' \quad \forall j \leq j' \quad M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

## ● Theorems

# Monge Property

$$\forall i \leq i' \quad \forall j \leq j' \quad M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

## ● Theorems

- $M$  is Monge  $\Rightarrow M$  is Totally Monotone
- $M$  is Monge  $\nLeftarrow M$  is Totally Monotone



# Monge Property

$$\forall i \leq i' \quad \forall j \leq j' \quad M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

## ● Theorems

- $M$  is **Monge**  $\Rightarrow$   $M$  is **Totally Monotone**  
 $M$  is **Monge**  $\nLeftarrow$   $M$  is **Totally Monotone**
- If  $\forall i$  and  $\forall j$ ,  $M_{i,j} + M_{i+1,j+1} \leq M_{i+1,j} + M_{i,j+1}$ ,  
then  $M$  is Monge.

# Monge Property

$$\forall i \leq i' \quad \forall j \leq j' \quad M_{i,j} + M_{i',j'} \leq M_{i',j} + M_{i,j'}$$

## ● Theorems

- $M$  is **Monge**  $\Rightarrow M$  is **Totally Monotone**  
 $M$  is **Monge**  $\nLeftarrow M$  is **Totally Monotone**
- If  $\forall i$  and  $\forall j$ ,  $M_{i,j} + M_{i+1,j+1} \leq M_{i+1,j} + M_{i,j+1}$ ,  
then  $M$  is Monge.
- $\Rightarrow$  Only need to prove Monge property for **adjacent** rows and columns.

# Monge Property

- General Scheme

# Monge Property

- General Scheme

1. Prove **Monge** Property for adjacent rows and columns

# Monge Property

## ● General Scheme

1. Prove **Monge** Property for adjacent rows and columns
2. (Automatically implies) **Monge** Property

# Monge Property

## ● General Scheme

1. Prove **Monge** Property for adjacent rows and columns
2. (Automatically implies) **Monge** Property
3. (Automatically implies) **Totally Monotone** Property

# Monge Property

## ● General Scheme

1. Prove **Monge** Property for adjacent rows and columns
2. (Automatically implies) **Monge** Property
3. (Automatically implies) **Totally Monotone** Property
4. Use **SMAWK** algorithm to find row minima

# Monge Property

## ● General Scheme

1. Prove **Monge** Property for adjacent rows and columns
2. (Automatically implies) **Monge** Property
3. (Automatically implies) **Totally Monotone** Property
4. Use **SMAWK** algorithm to find row minima
5. Usually  $\Theta(n)$  speedup



# Relationship?

Quadrangle Inequality

Totally Monotone (Monge)

# Relationship?

## Quadrangle Inequality

A matrix to be calculated

## Totally Monotone (Monge)

A matrix given implicitly

# Relationship?

## Quadrangle Inequality

A matrix to be calculated

Need all  $O(n^2)$  entries

## Totally Monotone (Monge)

A matrix given implicitly

Need only  $O(n)$  row minima

# Relationship?

## Quadrangle Inequality

A matrix to be calculated

Need all  $O(n^2)$  entries

$O(n^3)$  to  $O(n^2)$  speedup

## Totally Monotone (Monge)

A matrix given implicitly

Need only  $O(n)$  row minima

$O(n^2)$  to  $O(n)$  speedup

# Relationship?

## Quadrangle Inequality

A matrix to be calculated

Need all  $O(n^2)$  entries

$O(n^3)$  to  $O(n^2)$  speedup

## Totally Monotone (Monge)

A matrix given implicitly

Need only  $O(n)$  row minima

$O(n^2)$  to  $O(n)$  speedup

● This talk

# Relationship?

## Quadrangle Inequality

A matrix to be calculated

Need all  $O(n^2)$  entries

$O(n^3)$  to  $O(n^2)$  speedup

## Totally Monotone (Monge)

A matrix given implicitly

Need only  $O(n)$  row minima

$O(n^2)$  to  $O(n)$  speedup

## ● This talk

- QI instance is decomposed into  $\Theta(n)$  TM instances

# Relationship?

## Quadrangle Inequality

A matrix to be calculated

Need all  $O(n^2)$  entries

$O(n^3)$  to  $O(n^2)$  speedup

## Totally Monotone (Monge)

A matrix given implicitly

Need only  $O(n)$  row minima

$O(n^2)$  to  $O(n)$  speedup

## ● This talk

- QI instance is decomposed into  $\Theta(n)$  TM instances
- Each TM instance requires  $O(n)$  time

# Relationship?

## Quadrangle Inequality

A matrix to be calculated

Need all  $O(n^2)$  entries

$O(n^3)$  to  $O(n^2)$  speedup

## Totally Monotone (Monge)

A matrix given implicitly

Need only  $O(n)$  row minima

$O(n^2)$  to  $O(n)$  speedup

## ● This talk

- QI instance is decomposed into  $\Theta(n)$  TM instances
- Each TM instance requires  $O(n)$  time
- $\Rightarrow$  QI instance requires  $O(n^2)$  time in total



# Decompositions

QI instance  $\longrightarrow \Theta(n)$  TM instances

# Decompositions

QI instance  $\longrightarrow \Theta(n)$  TM instances

- $D^d$  decomposition
- $L^m$  and  $R^m$  decompositions

# Decompositions

QI instance  $\longrightarrow \Theta(n)$  TM instances

- $D^d$  decomposition
  - Each diagonal  $\longrightarrow$  TM instance
- $L^m$  and  $R^m$  decompositions

# Decompositions

QI instance  $\longrightarrow \Theta(n)$  TM instances

- $D^d$  decomposition
  - Each diagonal  $\longrightarrow$  TM instance
- $L^m$  and  $R^m$  decompositions
  - $L^m$ : Each row  $\longrightarrow$  TM instance
  - $R^m$ : Each column  $\longrightarrow$  TM instance

# Decompositions

QI instance  $\longrightarrow \Theta(n)$  TM instances

- $D^d$  decomposition

- Each diagonal  $\longrightarrow$  TM instance
- Permits solving QI problem directly using SMAWK.  
Same time bound as KY but different technique.

- $L^m$  and  $R^m$  decompositions

- $L^m$ : Each row  $\longrightarrow$  TM instance
- $R^m$ : Each column  $\longrightarrow$  TM instance

# Decompositions

QI instance  $\longrightarrow \Theta(n)$  TM instances

- $D^d$  decomposition

- Each diagonal  $\longrightarrow$  TM instance
- Permits solving QI problem directly using SMAWK.  
Same time bound as KY but different technique.

- $L^m$  and  $R^m$  decompositions

- $L^m$ : Each row  $\longrightarrow$  TM instance
- $R^m$ : Each column  $\longrightarrow$  TM instance
- Immediately implies the [original](#) KY speedup

# Decompositions

QI instance  $\longrightarrow \Theta(n)$  TM instances

- $D^d$  decomposition

- Each diagonal  $\longrightarrow$  TM instance
- Permits solving QI problem directly using SMAWK.  
Same time bound as KY but different technique.

- $L^m$  and  $R^m$  decompositions

- $L^m$ : Each row  $\longrightarrow$  TM instance
- $R^m$ : Each column  $\longrightarrow$  TM instance
- Immediately implies the **original** KY speedup
- Permits using algorithm of [Larmore, Schieber (1990)], to get **“online”** KY speedup.

# $D^d$ Decomposition



# $D^d$ Decomposition


- Each **diagonal**  $d$  in original QI matrix corresponds to a new Monge Matrix  $D^d$

# $D^d$ Decomposition

- Each **diagonal**  $d$  in original QI matrix corresponds to a new Monge Matrix  $D^d$
- Entries on **diagonal**  $d$  are row minima of corresponding rows in  $D^d$ .


# $D^d$ Decomposition

- Each **diagonal**  $d$  in original QI matrix corresponds to a new Monge Matrix  $D^d$
- Entries on **diagonal**  $d$  are row minima of corresponding rows in  $D^d$ .




# $D^d$ Decomposition

- Each **diagonal**  $d$  in original QI matrix corresponds to a new Monge Matrix  $D^d$
- Entries on **diagonal**  $d$  are row minima of corresponding rows in  $D^d$ .




# $D^d$ Decomposition

- Each **diagonal**  $d$  in original QI matrix corresponds to a new Monge Matrix  $D^d$
- Entries on **diagonal**  $d$  are row minima of corresponding rows in  $D^d$ .




# $D^d$ Decomposition

- Each **diagonal**  $d$  in original QI matrix corresponds to a new Monge Matrix  $D^d$
- Entries on **diagonal**  $d$  are row minima of corresponding rows in  $D^d$ .



# $D^d$ Decomposition

- Each **diagonal**  $d$  in original QI matrix corresponds to a new Monge Matrix  $D^d$
- Entries on **diagonal**  $d$  are row minima of corresponding rows in  $D^d$ .



# $L^m$ and $R^m$ Decompositions ( $R^m$ shown)



# $L^m$ and $R^m$ Decompositions ( $R^m$ shown)


- Each **column (row)**  $m$  in original QI matrix corresponds to a new Monge Matrix  $R^m$  ( $L^m$ )

# $L^m$ and $R^m$ Decompositions ( $R^m$ shown)

- Each **column (row)**  $m$  in original QI matrix corresponds to a new Monge Matrix  $R^m$  ( $L^m$ )
- Entries on **column (row)**  $m$  are row minima of corresponding rows in  $R^m$  ( $L^m$ ).


# $L^m$ and $R^m$ Decompositions ( $R^m$ shown)

- Each **column (row)**  $m$  in original QI matrix corresponds to a new Monge Matrix  $R^m$  ( $L^m$ )
- Entries on **column (row)**  $m$  are row minima of corresponding rows in  $R^m$  ( $L^m$ ).




# $L^m$ and $R^m$ Decompositions ( $R^m$ shown)

- Each **column (row)**  $m$  in original QI matrix corresponds to a new Monge Matrix  $R^m$  ( $L^m$ )
- Entries on **column (row)**  $m$  are row minima of corresponding rows in  $R^m$  ( $L^m$ ).




# $L^m$ and $R^m$ Decompositions ( $R^m$ shown)

- Each **column (row)**  $m$  in original QI matrix corresponds to a new Monge Matrix  $R^m$  ( $L^m$ )
- Entries on **column (row)**  $m$  are row minima of corresponding rows in  $R^m$  ( $L^m$ ).




# $L^m$ and $R^m$ Decompositions ( $R^m$ shown)

- Each **column (row)**  $m$  in original QI matrix corresponds to a new Monge Matrix  $R^m$  ( $L^m$ )
- Entries on **column (row)**  $m$  are row minima of corresponding rows in  $R^m$  ( $L^m$ ).



# $L^m$ and $R^m$ Decompositions ( $R^m$ shown)

- Each **column (row)**  $m$  in original QI matrix corresponds to a new Monge Matrix  $R^m$  ( $L^m$ )
- Entries on **column (row)**  $m$  are row minima of corresponding rows in  $R^m$  ( $L^m$ ).



# Outline

## ● Background

- Kunth-Yao (KY) Quadrangle Inequality (QI) Speedup
- SMAWK Algorithm for finding  
Row Minima of Totally Monotone (TM) Matrices

## ● The $D^d$ Decomposition

A transformation from QI to TM such that  
SMAWK solves KY problem as quickly as KY.

## ● The $L^m$ and $R^m$ Decompositions

Another transformation from QI to TM that  
(1) implies KY speedup and (2) enables online solution.

## ● Extensions

Applying the technique to known generalizations of KY.



# $D^d$ Decomposition

- Definition

# $D^d$ Decomposition

- Definition

- General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

# $D^d$ Decomposition

## ● Definition

### ● General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

### ● For diagonal $d$ , ( $1 \leq d < n$ )

$$B_{i,i+d} = w(i, i+d) + \min_{i < j \leq i+d} \{B_{i,j-1} + B_{j,i+d}\}$$

# $D^d$ Decomposition

## ● Definition

### ● General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

### ● For diagonal $d$ , ( $1 \leq d < n$ )

$$B_{i,i+d} = w(i, i+d) + \min_{i < j \leq i+d} \{B_{i,j-1} + B_{j,i+d}\}$$

### ● Define $(n - d + 1) \times (n + 1)$ matrix $D^d$

$$D_{i,j}^d = \begin{cases} w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} & \text{if } 0 \leq i < j \leq i+d \leq n \\ \infty & \text{otherwise} \end{cases}$$

# $D^d$ Decomposition

## Definition

### General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

### For diagonal $d$ , ( $1 \leq d < n$ )

$$B_{i,i+d} = w(i, i+d) + \min_{i < j \leq i+d} \{B_{i,j-1} + B_{j,i+d}\}$$

### Define $(n - d + 1) \times (n + 1)$ matrix $D^d$

$$D_{i,j}^d = \begin{cases} w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} & \text{if } 0 \leq i < j \leq i+d \leq n \\ \infty & \text{otherwise} \end{cases}$$

### Then,

$$B_{i,i+d} = \min_{i < j \leq i+d} D_{i,j}^d = \min_{0 \leq j \leq n} D_{i,j}^d$$

# $D^d$ Decomposition

## Definition

### General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

### For diagonal $d$ , ( $1 \leq d < n$ )

$$B_{i,i+d} = w(i, i+d) + \min_{i < j \leq i+d} \{B_{i,j-1} + B_{j,i+d}\}$$

### Define $(n - d + 1) \times (n + 1)$ matrix $D^d$

$$D_{i,j}^d = \begin{cases} w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} & \text{if } 0 \leq i < j \leq i+d \leq n \\ \infty & \text{otherwise} \end{cases}$$

### Then,

$$B_{i,i+d} = \min_{i < j \leq i+d} D_{i,j}^d = \min_{0 \leq j \leq n} D_{i,j}^d$$

## Lemma

### $D^d$ is Monge, for each $1 \leq d < n$ .

# $D^d$ Decomposition


$$D_{i,j}^d = \begin{cases} w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} & \text{if } 0 \leq i < j \leq i+d \leq n \\ \infty & \text{otherwise} \end{cases}$$

● Shape of  $D^d$

# $D^d$ Decomposition

$$D_{i,j}^d = \begin{cases} w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} & \text{if } 0 \leq i < j \leq i+d \leq n \\ \infty & \text{otherwise} \end{cases}$$

● Shape of  $D^d$





$D^d$  is Monge

# $D^d$ is Monge

Definition  $D_{i,j}^d = w(i, i + d) + \{B_{i,j-1} + B_{j,i+d}\}$

---

# $D^d$ is Monge

Definition  $D_{i,j}^d = w(i, i + d) + \{B_{i,j-1} + B_{j,i+d}\}$

---

Goal

$$D_{i,j}^d + D_{i+1,j+1}^d \leq D_{i+1,j}^d + D_{i,j+1}^d$$

# $D^d$ is Monge

Definition  $D_{i,j}^d = w(i, i + d) + \{B_{i,j-1} + B_{j,i+d}\}$

---

By definition

$$D_{i,j}^d + D_{i+1,j+1}^d = \{w(i, i + d) + w(i + 1, i + d + 1)\} + \\ \{B_{i,j-1} + B_{i+1,j}\} + \{B_{j,i+d} + B_{j+1,i+d+1}\}$$

$$D_{i+1,j}^d + D_{i,j+1}^d = \{w(i + 1, i + d + 1) + w(i, i + d)\} + \\ \{B_{i+1,j-1} + B_{i,j}\} + \{B_{j,i+d+1} + B_{j+1,i+d}\}$$

Goal

$$D_{i,j}^d + D_{i+1,j+1}^d \leq D_{i+1,j}^d + D_{i,j+1}^d$$

# $D^d$ is Monge

Definition  $D_{i,j}^d = w(i, i + d) + \{B_{i,j-1} + B_{j,i+d}\}$

---

By definition

$$D_{i,j}^d + D_{i+1,j+1}^d = \{w(i, i + d) + w(i + 1, i + d + 1)\} + \\ \{B_{i,j-1} + B_{i+1,j}\} + \{B_{j,i+d} + B_{j+1,i+d+1}\}$$

$$D_{i+1,j}^d + D_{i,j+1}^d = \{w(i + 1, i + d + 1) + w(i, i + d)\} + \\ \{B_{i+1,j-1} + B_{i,j}\} + \{B_{j,i+d+1} + B_{j+1,i+d}\}$$

Since  $B$  satisfies QI,

$$B_{i,j-1} + B_{i+1,j} \leq B_{i+1,j-1} + B_{i,j} \\ B_{j,i+d} + B_{j+1,i+d+1} \leq B_{j,i+d+1} + B_{j+1,i+d}$$

Goal

$$D_{i,j}^d + D_{i+1,j+1}^d \leq D_{i+1,j}^d + D_{i,j+1}^d$$

# $D^d$ is Monge

Definition  $D_{i,j}^d = w(i, i + d) + \{B_{i,j-1} + B_{j,i+d}\}$

---

By definition

$$D_{i,j}^d + D_{i+1,j+1}^d = \{w(i, i + d) + w(i + 1, i + d + 1)\} + \\ \{B_{i,j-1} + B_{i+1,j}\} + \{B_{j,i+d} + B_{j+1,i+d+1}\}$$

$$D_{i+1,j}^d + D_{i,j+1}^d = \{w(i + 1, i + d + 1) + w(i, i + d)\} + \\ \{B_{i+1,j-1} + B_{i,j}\} + \{B_{j,i+d+1} + B_{j+1,i+d}\}$$

Since  $B$  satisfies QI,

$$B_{i,j-1} + B_{i+1,j} \leq B_{i+1,j-1} + B_{i,j} \\ B_{j,i+d} + B_{j+1,i+d+1} \leq B_{j,i+d+1} + B_{j+1,i+d}$$

So

$$D_{i,j}^d + D_{i+1,j+1}^d \leq D_{i+1,j}^d + D_{i,j+1}^d$$

# **SMAWK replaces KY**

# SMAWK replaces KY

● We know

$$● D_{i,j}^d = \begin{cases} w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} & \text{if } 0 \leq i < j \leq i+d \leq n \\ \infty & \text{otherwise} \end{cases}$$

$$● B_{i,i+d} = \min_{0 \leq j \leq n} D_{i,j}^d = \text{minimum of row } i \text{ of } D^d$$

●  $D^d$  is Monge, for each  $1 \leq d < n$ .



# SMAWK replaces KY

- We know

- $$D_{i,j}^d = \begin{cases} w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} & \text{if } 0 \leq i < j \leq i+d \leq n \\ \infty & \text{otherwise} \end{cases}$$

- $B_{i,i+d} = \min_{0 \leq j \leq n} D_{i,j}^d = \text{minimum of row } i \text{ of } D^d$

- $D^d$  is Monge, for each  $1 \leq d < n$ .

- For fixed  $d$ , SMAWK can be used to find all the  $B_{i,i+d}$  (row minima of  $D^d$ ) in  $O(n)$  time.

# SMAWK replaces KY

- We know

- $$D_{i,j}^d = \begin{cases} w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} & \text{if } 0 \leq i < j \leq i+d \leq n \\ \infty & \text{otherwise} \end{cases}$$

- $B_{i,i+d} = \min_{0 \leq j \leq n} D_{i,j}^d = \text{minimum of row } i \text{ of } D^d$

- $D^d$  is Monge, for each  $1 \leq d < n$ .

- For fixed  $d$ , SMAWK can be used to find all the  $B_{i,i+d}$  (row minima of  $D^d$ ) in  $O(n)$  time.

- $\Rightarrow O(n^2)$  time for all  $D^d$ .

# SMAWK replaces KY

- We know

- $$D_{i,j}^d = \begin{cases} w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} & \text{if } 0 \leq i < j \leq i+d \leq n \\ \infty & \text{otherwise} \end{cases}$$

- $B_{i,i+d} = \min_{0 \leq j \leq n} D_{i,j}^d = \text{minimum of row } i \text{ of } D^d$

- $D^d$  is Monge, for each  $1 \leq d < n$ .

- For fixed  $d$ , SMAWK can be used to find all the  $B_{i,i+d}$  (row minima of  $D^d$ ) in  $O(n)$  time.

- $\Rightarrow O(n^2)$  time for all  $D^d$ .

- Note: Must run SMAWK on  $D^d$  in the order  $d = 1, 2, 3, \dots$

Entries in  $D^d$  depend upon row minima of  $D^{d'}$  where  $d' < d$ .

# Outline

## ● Background

- Kunth-Yao (KY) Quadrangle Inequality (QI) Speedup

- SMAWK Algorithm for finding  
Row Minima of Totally Monotone (TM) Matrices

## ● The $D^d$ Decomposition

A transformation from QI to TM such that  
SMAWK solves KY problem as quickly as KY.

## ● The $L^m$ and $R^m$ Decompositions

Another transformation from QI to TM that  
(1) implies KY speedup and (2) enables online solution.

## ● Extensions


Applying the technique to known generalizations of KY.

# $R^m$ Decomposition

- $R^m$  decomposition


# $R^m$ Decomposition

●  $R^m$  decomposition




# $R^m$ Decomposition

●  $R^m$  decomposition



# $R^m$ Decomposition


●  $R^m$  decomposition






# $R^m$ Decomposition

●  $R^m$  decomposition



# $R^m$ Decomposition

●  $R^m$  decomposition



# $R^m$ Decomposition

- Definition

# $R^m$ Decomposition

- Definition

- General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

# $R^m$ Decomposition

## ● Definition

### ● General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

### ● For column $m$ , ( $1 \leq m \leq n$ )

$$B_{i,m} = w(i, m) + \min_{i < j \leq m} \{B_{i,j-1} + B_{j,m}\}$$

# $R^m$ Decomposition

## ● Definition

### ● General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

### ● For column $m$ , ( $1 \leq m \leq n$ )

$$B_{i,m} = w(i, m) + \min_{i < j \leq m} \{B_{i,j-1} + B_{j,m}\}$$

### ● Define $(m+1) \times (m+1)$ matrix $R^m$

$$R_{i,j}^m = \begin{cases} w(i, m) + \{B_{i,j-1} + B_{j,m}\} & \text{if } 0 \leq i < j \leq m \\ \infty & \text{otherwise} \end{cases}$$

# $R^m$ Decomposition

## Definition

### General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

### For column $m$ , ( $1 \leq m \leq n$ )

$$B_{i,m} = w(i, m) + \min_{i < j \leq m} \{B_{i,j-1} + B_{j,m}\}$$

### Define $(m+1) \times (m+1)$ matrix $R^m$

$$R_{i,j}^m = \begin{cases} w(i, m) + \{B_{i,j-1} + B_{j,m}\} & \text{if } 0 \leq i < j \leq m \\ \infty & \text{otherwise} \end{cases}$$

### Then

$$B_{i,m} = \min_{i < j \leq m} R_{i,j}^m = \min_{0 < j \leq m} R_{i,j}^m$$

# $R^m$ Decomposition

## Definition

### General recurrence

$$B_{i,j} = w(i, j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

### For column $m$ , ( $1 \leq m \leq n$ )

$$B_{i,m} = w(i, m) + \min_{i < j \leq m} \{B_{i,j-1} + B_{j,m}\}$$

### Define $(m+1) \times (m+1)$ matrix $R^m$

$$R_{i,j}^m = \begin{cases} w(i, m) + \{B_{i,j-1} + B_{j,m}\} & \text{if } 0 \leq i < j \leq m \\ \infty & \text{otherwise} \end{cases}$$

### Then

$$B_{i,m} = \min_{i < j \leq m} R_{i,j}^m = \min_{0 < j \leq m} R_{i,j}^m$$

## Lemma

### $R^m$ is Monge, for each $1 \leq m \leq n$ .



# $R^m$ Decomposition


$$R_{i,j}^m = \begin{cases} w(i, m) + \{B_{i,j-1} + B_{j,m}\} & \text{if } 0 \leq i < j \leq m \\ \infty & \text{otherwise} \end{cases}$$

● Shape of  $R^m$

# $R^m$ Decomposition

$$R_{i,j}^m = \begin{cases} w(i, m) + \{B_{i,j-1} + B_{j,m}\} & \text{if } 0 \leq i < j \leq m \\ \infty & \text{otherwise} \end{cases}$$

● Shape of  $R^m$



# $R^m$ is Monge

# $R^m$ is Monge

Definition  $R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\}$

---

# $R^m$ is Monge

Definition  $R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\}$

---

Goal

$$R_{i,j}^m + R_{i+1,j+1}^m \leq R_{i+1,j}^m + R_{i,j+1}^m$$

# $R^m$ is Monge

Definition  $R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\}$

---

By definition

$$R_{i,j}^m + R_{i+1,j+1}^m = \{w(i, m) + w(i+1, m)\} + \\ \{B_{i,j-1} + B_{i+1,j}\} + \{B_{j,m} + B_{j+1,m}\}$$

$$R_{i+1,j}^m + R_{i,j+1}^m = \{w(i+1, m) + w(i, m)\} + \\ \{B_{i+1,j-1} + B_{i,j}\} + \{B_{j,m} + B_{j+1,m}\}$$

Goal

$$R_{i,j}^m + R_{i+1,j+1}^m \leq R_{i+1,j}^m + R_{i,j+1}^m$$

# $R^m$ is Monge

Definition  $R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\}$

---

By definition

$$R_{i,j}^m + R_{i+1,j+1}^m = \{w(i, m) + w(i+1, m)\} + \\ \{B_{i,j-1} + B_{i+1,j}\} + \{B_{j,m} + B_{j+1,m}\}$$

$$R_{i+1,j}^m + R_{i,j+1}^m = \{w(i+1, m) + w(i, m)\} + \\ \{B_{i+1,j-1} + B_{i,j}\} + \{B_{j,m} + B_{j+1,m}\}$$

Since  $B$  satisfies QI,

$$B_{i,j-1} + B_{i+1,j} \leq B_{i+1,j-1} + B_{i,j}$$

Goal

$$R_{i,j}^m + R_{i+1,j+1}^m \leq R_{i+1,j}^m + R_{i,j+1}^m$$

# $R^m$ is Monge

Definition  $R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\}$

---

By definition

$$R_{i,j}^m + R_{i+1,j+1}^m = \{w(i, m) + w(i+1, m)\} + \\ \{B_{i,j-1} + B_{i+1,j}\} + \{B_{j,m} + B_{j+1,m}\}$$

$$R_{i+1,j}^m + R_{i,j+1}^m = \{w(i+1, m) + w(i, m)\} + \\ \{B_{i+1,j-1} + B_{i,j}\} + \{B_{j,m} + B_{j+1,m}\}$$

Since  $B$  satisfies QI,

$$B_{i,j-1} + B_{i+1,j} \leq B_{i+1,j-1} + B_{i,j}$$

So

$$R_{i,j}^m + R_{i+1,j+1}^m \leq R_{i+1,j}^m + R_{i,j+1}^m$$



# Outline

## ● Background

- Kunth-Yao (KY) Quadrangle Inequality (QI) Speedup
- SMAWK Algorithm for finding  
Row Minima of Totally Monotone (TM) Matrices

## ● The $D^d$ Decomposition

A transformation from QI to TM such that  
SMAWK solves KY problem as quickly as KY.

## ● The $L^m$ and $R^m$ Decompositions

Another transformation from QI to TM that  
(1) implies KY speedup and (2) enables online solution.

## ● Extensions

Applying the technique to known generalizations of KY.

# $L^m$ and $R^m$ Imply Original KY Result

- KY Speedup

- $K_B(i, j) \leq K_B(i, j + 1) \leq K_B(i + 1, j + 1)$

# $L^m$ and $R^m$ Imply Original KY Result

- KY Speedup

- $K_B(i, j) \leq K_B(i, j + 1) \leq K_B(i + 1, j + 1)$

- $R^m \longrightarrow K_B(i, j + 1) \leq K_B(i + 1, j + 1)$

# $L^m$ and $R^m$ Imply Original KY Result

## ● KY Speedup

●  $K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$

●  $R^m \longrightarrow K_B(i, j+1) \leq K_B(i+1, j+1)$

## ● Recall

$\text{RM}_{R^m}(i)$  is **index** of rightmost minimum of row  $i$  of  $R^m$ .

1	1	2	2	2	2
1	1	1	1	2	2
1	1	1	1	2	2
1	1	1	1	2	2
1	1	1	1	1	1
1	1	1	1	1	1

$$\text{RM}_M(1) = 2$$

$$\text{RM}_M(2) = 4$$

$$\text{RM}_M(3) = 4$$

$$\text{RM}_M(4) = 4$$

$$\text{RM}_M(5) = 6$$

$$\text{RM}_M(6) = 6$$

# $L^m$ and $R^m$ Imply Original KY Result

## ● KY Speedup

●  $K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$

●  $R^m \longrightarrow K_B(i, j+1) \leq K_B(i+1, j+1)$

## ● Recall

$\text{RM}_{R^m}(i)$  is **index** of rightmost minimum of row  $i$  of  $R^m$ .

## ● From the definition

$$B_{i,m} = \min_{i < j \leq m} R_{i,j}^m \longrightarrow K_B(i, m) = \text{RM}_{R^m}(i)$$

# $L^m$ and $R^m$ Imply Original KY Result

## ● KY Speedup

●  $K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$

●  $R^m \longrightarrow K_B(i, j+1) \leq K_B(i+1, j+1)$

## ● Recall

$\text{RM}_{R^m}(i)$  is **index** of rightmost minimum of row  $i$  of  $R^m$ .

## ● From the definition

$$B_{i,m} = \min_{i < j \leq m} R_{i,j}^m \longrightarrow K_B(i, m) = \text{RM}_{R^m}(i)$$

## ● So

$$R^m \text{ is } \text{TM} \longrightarrow \text{RM}_{R^m}(i) \leq \text{RM}_{R^m}(i+1)$$

$$\longrightarrow K_B(i, m) \leq K_B(i+1, m)$$

# $L^m$ and $R^m$ Imply Original KY Result

## ● KY Speedup

●  $K_B(i, j) \leq K_B(i, j+1) \leq K_B(i+1, j+1)$

●  $R^m \longrightarrow K_B(i, j+1) \leq K_B(i+1, j+1)$

## ● Recall

$\text{RM}_{R^m}(i)$  is **index** of rightmost minimum of row  $i$  of  $R^m$ .

## ● From the definition

$$B_{i,m} = \min_{i < j \leq m} R_{i,j}^m \longrightarrow K_B(i, m) = \text{RM}_{R^m}(i)$$

## ● So

$$R^m \text{ is } \text{TM} \longrightarrow \text{RM}_{R^m}(i) \leq \text{RM}_{R^m}(i+1)$$

$$\longrightarrow K_B(i, m) \leq K_B(i+1, m)$$

●  $L^m \longrightarrow K_B(i, j) \leq K_B(i, j+1)$

## ● Similar

# Outline

## ● Background

- Kunth-Yao (KY) Quadrangle Inequality (QI) Speedup

- SMAWK Algorithm for finding  
Row Minima of Totally Monotone (TM) Matrices

## ● The $D^d$ Decomposition

A transformation from QI to TM such that  
SMAWK solves KY problem as quickly as KY.

## ● The $L^m$ and $R^m$ Decompositions

Another transformation from QI to TM that  
(1) implies KY speedup and (2) enables online solution.

## ● Extensions

Applying the technique to known generalizations of KY.



# LARSCH Algorithm

# LARSCH Algorithm

●  $D^d$  decomposition

# LARSCH Algorithm

●  $D^d$  decomposition

●  $D_{i,j}^d = w(i, i + d) + \{B_{i,j-1} + B_{j,i+d}\} \quad (0 \leq i < j \leq i + d \leq n)$

# LARSCH Algorithm

- $D^d$  decomposition

- $D_{i,j}^d = w(i, i + d) + \{B_{i,j-1} + B_{j,i+d}\} \quad (0 \leq i < j \leq i + d \leq n)$

- SMAWK algorithm

# LARSCH Algorithm

- $D^d$  decomposition

- $D_{i,j}^d = w(i, i + d) + \{B_{i,j-1} + B_{j,i+d}\} \quad (0 \leq i < j \leq i + d \leq n)$

- SMAWK algorithm

- $L^m$  and  $R^m$  decomposition

# LARSCH Algorithm

- $D^d$  decomposition

- $D_{i,j}^d = w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} \quad (0 \leq i < j \leq i+d \leq n)$

- SMAWK algorithm

- $L^m$  and  $R^m$  decomposition

- $R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\} \quad (0 \leq i < j \leq m)$

# LARSCH Algorithm

## ● $D^d$ decomposition

●  $D_{i,j}^d = w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} \quad (0 \leq i < j \leq i+d \leq n)$

● SMAWK algorithm

## ● $L^m$ and $R^m$ decomposition

●  $R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\} \quad (0 \leq i < j \leq m)$

● Can **not** use SMAWK algorithm:

$B_{j,m} = \min_t R_{j,t}^m$  is **row-minima** of row  $j$  of  $R^m$   
and is therefore **not** known.

# LARSCH Algorithm

## ● $D^d$ decomposition

●  $D_{i,j}^d = w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} \quad (0 \leq i < j \leq i+d \leq n)$

● SMAWK algorithm

## ● $L^m$ and $R^m$ decomposition

●  $R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\} \quad (0 \leq i < j \leq m)$

● Can **not** use SMAWK algorithm:

$B_{j,m} = \min_t R_{j,t}^m$  is **row-minima of row  $j$  of  $R^m$**   
and is therefore **not** known.

● LARSCH algorithm [Larmore, Schieber (1990)]

permits calculating row minima of TM matrices in  $O(n)$  time,  
even with this dependency.



# LARSCH Algorithm

## • $D^d$ decomposition

•  $D_{i,j}^d = w(i, i+d) + \{B_{i,j-1} + B_{j,i+d}\} \quad (0 \leq i < j \leq i+d \leq n)$

• SMAWK algorithm

## • $L^m$ and $R^m$ decomposition

•  $R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\} \quad (0 \leq i < j \leq m)$

• Can **not** use SMAWK algorithm:

$B_{j,m} = \min_t R_{j,t}^m$  is **row-minima of row  $j$  of  $R^m$**   
and is therefore **not** known.

• LARSCH algorithm [Larmore, Schieber (1990)]

permits calculating row minima of TM matrices in  $O(n)$  time,  
even with this dependency.

•  $O(n)$  time for each column  $\Rightarrow O(n^2)$  in total.

# LARSCH Algorithm

Finding row minima in totally monotone matrices **with limited dependency**.  
This is also known as **online TM problem**.


# LARSCH Algorithm

Finding row minima in totally monotone matrices **with limited dependency**.  
This is also known as **online TM problem**.

Entries of column  $j$  can depend on the row minima of rows  $i$  where  $M_{i,j} = \infty$ .

Green: the column  $j$ .

Red: rows that column  $j$  can depend on.




# LARSCH Algorithm

Finding row minima in totally monotone matrices **with limited dependency**.  
This is also known as **online TM problem**.

Entries of column  $j$  can depend on the row minima of rows  $i$  where  $M_{i,j} = \infty$ .

Green: the column  $j$ .

Red: rows that column  $j$  can depend on.




# LARSCH Algorithm

Finding row minima in totally monotone matrices **with limited dependency**.  
This is also known as **online TM problem**.

Entries of column  $j$  can depend on the row minima of rows  $i$  where  $M_{i,j} = \infty$ .

Green: the column  $j$ .

Red: rows that column  $j$  can depend on.






# LARSCH Algorithm

Finding row minima in totally monotone matrices **with limited dependency**.  
This is also known as **online TM problem**.

Entries of column  $j$  can depend on the row minima of rows  $i$  where  $M_{i,j} = \infty$ .

Green: the column  $j$ .

Red: rows that column  $j$  can depend on.



$$R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\} \quad (0 \leq i < j \leq m)$$


# LARSCH Algorithm

Finding row minima in totally monotone matrices **with limited dependency**.  
This is also known as **online TM problem**.

Entries of column  $j$  can depend on the row minima of rows  $i$  where  $M_{i,j} = \infty$ .

Green: the column  $j$ .

Red: rows that column  $j$  can depend on.



$$R_{i,j}^m = w(i, m) + \{B_{i,j-1} + B_{j,m}\} \quad (0 \leq i < j \leq m)$$

$R^m$  satisfies the condition of LARSCH.



# Note

- Aggarwal and Park (FOCS '88) developed a 3-D monotone matrix representation of the KY problem and then showed how to use an algorithm due to Wilber (for online computation of maxima of certain concave sequences) to calculate “tube-maxima” of their matrices.
- Careful decomposition of their work yields a decomposition similar to  $L^m$  and an  $O(n)$  algorithm for calculating its row-minima. This provides an alternative derivation of the previous result (with a symmetry argument extending it to  $R^m$ ).

# Online Algorithm

# Online Algorithm

- Recall: Two-sided online

# Online Algorithm

- Recall: Two-sided online
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$

# Online Algorithm

- Recall: Two-sided online
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
  - Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .

# Online Algorithm

- Recall: Two-sided online
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
  - Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .

	1	2	3	4	5	6
1	0	146	260	349	491	624
2		0	75	141	250	357
3			0	43	119	204
4				0	44	121
5					0	52
6						0

# Online Algorithm

- Recall: Two-sided online
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
  - Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .
- Online algorithm: using LARSCH

# Online Algorithm

- Recall: Two-sided online
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
  - Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .
- Online algorithm: using LARSCH
  - Add  $\text{Key}_{r+1}$



# Online Algorithm

- Recall: Two-sided online
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
  - Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .
- Online algorithm: using LARSCH
  - Add  $\text{Key}_{r+1}$
  - Construct  $R^{r+1}$

# Online Algorithm

- Recall: Two-sided online
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
  - Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .
- Online algorithm: using LARSCH
  - Add  $\text{Key}_{r+1}$
  - Construct  $R^{r+1}$
  - Solve by LARSCH

# Online Algorithm

- Recall: Two-sided online
  - Current step: Optimal BST for  $\text{Key}_{l+1}, \dots, \text{Key}_r$
  - Next step: Add either  $\text{Key}_l$  or  $\text{Key}_{r+1}$ .
- Online algorithm: using LARSCH
  - Add  $\text{Key}_{r+1}$
  - Construct  $R^{r+1}$
  - Solve by LARSCH
  - $O(n)$  time worst case

# Outline

## ● Background

- Kunth-Yao (KY) Quadrangle Inequality (QI) Speedup
- SMAWK Algorithm for finding  
Row Minima of Totally Monotone (TM) Matrices

## ● The $D^d$ Decomposition

A transformation from QI to TM such that  
SMAWK solves KY problem as quickly as KY.

## ● The $L^m$ and $R^m$ Decompositions

Another transformation from QI to TM that  
(1) implies KY speedup and (2) enables online solution.

## ● Extensions

Applying the technique to known generalizations of KY.

# Extensions

- Some known extensions

# Extensions

- Some known extensions
  - [Michelle L. Wachs (1989)]

# Extensions

- Some known extensions
  - [Michelle L. Wachs (1989)]
  - [Al Borchers, Prosenjit Gupta (1994)]

# Recurrence



# Recurrence

## ● Original Knuth-Yao

$$B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

# Recurrence

- Original Knuth-Yao

$$B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$$

- Borchers and Gupta

$$B_{i,j} = \min_{i < t \leq j} \{w(i,t,j) + aB_{i,t-1} + bB_{t,j}\}$$

# Generalization of QI

# Generalization of QI

● Original Knuth-Yao

●  $B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

# Generalization of QI

## Original Knuth-Yao

•  $B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

•  $w(i,j)$  satisfies QI, if  $\forall i \leq i' \leq j \leq j'$

$$w(i,j) + w(i',j') \leq w(i',j) + w(i,j')$$

# Generalization of QI

## ● Original Knuth-Yao

●  $B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

●  $w(i,j)$  satisfies QI, if  $\forall i \leq i' \leq j \leq j'$

$$w(i,j) + w(i',j') \leq w(i',j) + w(i,j')$$

## ● Borchers and Gupta

●  $B_{i,j} = \min_{i < t \leq j} \{w(i,t,j) + aB_{i,t-1} + bB_{t,j}\}$

# Generalization of QI

## Original Knuth-Yao

- $B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

- $w(i,j)$  satisfies QI, if  $\forall i \leq i' \leq j \leq j'$

$$w(i,j) + w(i',j') \leq w(i',j) + w(i,j')$$

## Borchers and Gupta

- $B_{i,j} = \min_{i < t \leq j} \{w(i,t,j) + aB_{i,t-1} + bB_{t,j}\}$

- $w(i,t,j)$  satisfies QI, if  $\forall i \leq i' < t \leq t' \leq j'$  and  $t \leq j \leq j'$

$$w(i,t,j) + w(i',t',j') \leq w(i',t,j) + w(i,t',j')$$

and  $\forall i < t \leq t' \leq j \leq j'$  and  $i \leq i' < t'$

$$w(i',t',j') + w(i,t,j) \leq w(i',t',j) + w(i,t,j')$$

# Generalization of QI

## Original Knuth-Yao

- $B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

- $w(i,j)$  satisfies QI, if  $\forall i \leq i' \leq j \leq j'$

$$w(i,j) + w(i',j') \leq w(i',j) + w(i,j')$$

## Borchers and Gupta

- $B_{i,j} = \min_{i < t \leq j} \{w(i,t,j) + aB_{i,t-1} + bB_{t,j}\}$

- $w(i,t,j)$  satisfies QI, if  $\forall i \leq i' < t \leq t' \leq j'$  and  $t \leq j \leq j'$

$$w(i,t,j) + w(i',t',j') \leq w(i',t,j) + w(i,t',j')$$

and  $\forall i < t \leq t' \leq j \leq j'$  and  $i \leq i' < t'$

$$w(i',t',j') + w(i,t,j) \leq w(i',t',j) + w(i,t,j')$$

- If the value of  $w(i,t,j)$  is independent of  $t$ , the Borchers and Gupta definition becomes the original Knuth-Yao definition.



# Generalization of MIL

# Generalization of MIL

● Original Knuth-Yao

●  $B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

# Generalization of MIL

## ● Original Knuth-Yao

●  $B_{i,j} = w(i,j) + \min_{i < t \leq j} \{ B_{i,t-1} + B_{t,j} \}$

●  $w(i,j)$  is Monotone on the integer lattice (MIL),  
if  $\forall [i,j] \subseteq [i',j'], w(i,j) \leq w(i',j')$ .

# Generalization of MIL

## ● Original Knuth-Yao

- $B_{i,j} = w(i,j) + \min_{i < t \leq j} \{B_{i,t-1} + B_{t,j}\}$

- $w(i,j)$  is Monotone on the integer lattice (MIL),  
if  $\forall [i,j] \subseteq [i',j'], w(i,j) \leq w(i',j')$ .

## ● Borchers and Gupta

- $B_{i,j} = \min_{i < t \leq j} \{w(i,t,j) + aB_{i,t-1} + bB_{t,j}\}$

# Generalization of MIL

## ● Original Knuth-Yao

●  $B_{i,j} = w(i,j) + \min_{i < t \leq j} \{ B_{i,t-1} + B_{t,j} \}$

●  $w(i,j)$  is Monotone on the integer lattice (MIL),  
if  $\forall [i,j] \subseteq [i',j']$ ,  $w(i,j) \leq w(i',j')$ .

## ● Borchers and Gupta

●  $B_{i,j} = \min_{i < t \leq j} \{ w(i,t,j) + aB_{i,t-1} + bB_{t,j} \}$

●  $w(i,t,j)$  is Monotone on the integer lattice (MIL),  
if  $\forall [i,j] \subseteq [i',j']$  and  $i < t \leq j$ ,  $w(i,t,j) \leq w(i',t,j')$ .

# Applications

## Applications

# Applications

- [Borchers, Gupta (1994)]  
Rectilinear Steiner Minimal Arborescence (RSMA) of a slide

# Applications

- [Borchers, Gupta (1994)]  
Rectilinear Steiner Minimal Arborescence (RSMA) of a slide
- Slide: a set of points  $(x_i, y_i)$  such that,  
if  $i < j$ , then  $x_i < x_j$  and  $y_i > y_j$ .



# Applications


- [Borchers, Gupta (1994)]  
Rectilinear Steiner Minimal Arborescence (RSMA) of a slide
- Slide: a set of points  $(x_i, y_i)$  such that,  
if  $i < j$ , then  $x_i < x_j$  and  $y_i > y_j$ .
- RSMA: a directed tree where each edge  
either goes up or to the right.

# Applications

- [Borchers, Gupta (1994)]

Rectilinear Steiner Minimal Arborescence (RSMA) of a **slide**

- **Slide**: a set of points  $(x_i, y_i)$  such that, if  $i < j$ , then  $x_i < x_j$  and  $y_i > y_j$ .
- **RSMA**: a directed tree where each edge either goes up or to the right.




# Applications

## ● [Borchers, Gupta (1994)]

Rectilinear Steiner Minimal Arborescence (RSMA) of a slide

- Slide: a set of points  $(x_i, y_i)$  such that, if  $i < j$ , then  $x_i < x_j$  and  $y_i > y_j$ .
- RSMA: a directed tree where each edge either goes up or to the right.




- $$B_{i,j} = \min_{i < t \leq j} \left\{ \underbrace{(x_t - x_i + y_{t-1} - y_j)}_{w(i,t,j)} + B_{i,t-1} + B_{t,j} \right\}$$

# Applications

## ● [Borchers, Gupta (1994)]

Rectilinear Steiner Minimal Arborescence (RSMA) of a slide

- Slide: a set of points  $(x_i, y_i)$  such that, if  $i < j$ , then  $x_i < x_j$  and  $y_i > y_j$ .
- RSMA: a directed tree where each edge either goes up or to the right.



- $B_{i,j} = \min_{i < t \leq j} \left\{ \underbrace{(x_t - x_i + y_{t-1} - y_j)}_{w(i,t,j)} + B_{i,t-1} + B_{t,j} \right\}$
- $w(i, t, j)$  satisfies generalized QI and MIL.

# Outline

- Background

- Kunth-Yao (KY) Quadrangle Inequality (QI) Speedup

- SMAWK Algorithm for finding  
Row Minima of Totally Monotone (TM) Matrices

- The  $D^d$  Decomposition

A transformation from QI to TM such that  
SMAWK solves KY problem as quickly as KY.

- The  $L^m$  and  $R^m$  Decompositions

Another transformation from QI to TM that  
(1) implies KY speedup and (2) enables online solution.

- Extensions

Applying the technique to known generalizations of KY.

# Questions?