

Steinberg modules and Donkin pairs.

Wilberd van der Kallen

November 27, 2017

1 Summary

We prove that in characteristic $p > 0$ a module with good filtration for a group of type E_6 restricts to a module with good filtration for a group of type F_4 . Thus we confirm a conjecture of Brundan for one more case. Our method relies on the canonical Frobenius splittings of Mathieu. Next we settle the remaining cases, in characteristic not 2, with a computer-aided variation on the old method of Donkin.

2 Preliminaries

Our base field k is algebraically closed of characteristic p . Let G be a connected semisimple group and H a connected semisimple subgroup. (Good filtrations with more general groups are treated in [3].) We refer to [5] and [16] for unexplained terminology and notation.

Now choose a Borel subgroup B in G and a maximal torus T in B so that, if B^- is the opposite Borel subgroup, then $B \cap H$ and $B^- \cap H$ are a Borel subgroups in H and $T \cap H$ is a maximal torus in H .

We follow the convention that the roots of B are positive. If $\lambda \in X(T)$ is dominant, then $\text{ind}_B^G(-\lambda)$ is the dual Weyl module $\nabla_G(\lambda^*)$ with highest weight $\lambda^* = -w_0\lambda$ and lowest weight $-\lambda$. Its dual is the Weyl module $\Delta_G(\lambda)$. In a good filtration of a G -module the layers are of the form $\nabla_G(\mu)$.

Definition 2.1 We say that (G, H) is a Donkin pair if for any G -module M with good filtration, the H -module $\text{res}_H^G M$ has good filtration.

Let $\mathcal{U}(U)$ denote the hyperalgebra of the unipotent radical U of B . We recall the presentation of Weyl modules.

Lemma 2.2 *Let λ be dominant and let $v_{-\lambda^*}$ be a nonzero weight vector of lowest weight $-\lambda^*$ in $\Delta_G(\lambda)$. Then $v_{-\lambda^*}$ generates $\Delta_G(\lambda)$ as a $\mathcal{U}(U)$ -module, and the annihilator of $v_{-\lambda^*}$ equals the left ideal of $\mathcal{U}(U)$ generated by the $X_\alpha^{(n)}$ with α simple and $n > (\lambda^*, \alpha^\vee)$.*

Proof Note that $\mathcal{U}(U)$ is a graded algebra graded by height. Therefore the left ideal in the lemma is the intersection of all ideals I of finite codimension that contain it and that lie inside the annihilator. But by the proof of [12, Proposition Fondamentale] such ideals I are equal to the annihilator. \square

Let X be a smooth projective B -variety with canonical bundle ω . (Generalizations to other varieties will be left to the reader.) There is by [10, §2] a natural map $\epsilon : H^0(X, \omega^{1-p}) \rightarrow k$ so that $\phi \in H^0(X, \omega^{1-p})$ determines a Frobenius splitting if and only if $\epsilon(\phi) = 1$. Let St_G be the Steinberg module of the simply connected cover \tilde{G} of G . For simplicity of notation we further assume that St_G is actually a G -module. Its B -socle is the highest weight space $k_{(p-1)\rho}$.

Recall that a Frobenius splitting of X is called *canonical* if the corresponding ϕ is T -invariant and lies in the image of a B -module map $\text{St}_G \otimes k_{(p-1)\rho} \rightarrow H^0(X, \omega^{1-p})$. (Compare lemma 2.2 and [16, Definition 4.3.5].) If the group G needs to be emphasized, we will speak of a G -canonical splitting. Now suppose X is actually a G -variety.

Lemma 2.3 *X has a canonical splitting if and only if there is a G -module map $\psi : \text{St}_G \otimes \text{St}_G \rightarrow H^0(X, \omega^{1-p})$ so that $\epsilon\psi \neq 0$.*

Proof There is, up to scalar multiple, only one possibility for a map $\text{St}_G \otimes \text{St}_G \rightarrow k$. If $\epsilon\psi \neq 0$ then the subspace of T -invariants in $\text{St}_G \otimes k_{(p-1)\rho}$ maps isomorphically to k . Conversely, a map from $\text{St}_G \otimes k_{(p-1)\rho}$ to a G -module M can be extended to $\text{St}_G \otimes \text{St}_G$ because the G -module generated by the image of $k_{(p-1)\rho}$ in $M \otimes \text{St}_G^*$ is St_G . \square

We have the following fundamental result of Mathieu [8].

Theorem 2.4 [9, 6.2] *Assume X has a canonical splitting and \mathcal{L} is a G -linearized line bundle on X . Then $H^0(X, \mathcal{L})$ has a good filtration.*

3 Pairings

Now apply this to $X = G/B$. Of course the $\nabla_G(\mu)$ are of the form $H^0(X, \mathcal{L})$, see [5, I 5.12]. It follows that (G, H) is a Donkin pair if X has an H -canonical splitting. We also have a surjection $\mathrm{St}_G \otimes \mathrm{St}_G \rightarrow H^0(X, \omega^{1-p})$, by [5, II 14.20]. The composite with $H^0(X, \omega^{1-p}) \rightarrow k$ may be identified as in [6], [11] with the natural pairing on the self dual representation St_G . Thus we get

Theorem 3.1 (Pairing criterion) *Assume there is an H -module map*

$$\mathrm{St}_H^* \otimes \mathrm{St}_H \rightarrow \mathrm{St}_G^* \otimes \mathrm{St}_G$$

whose composite with the evaluation map $\mathrm{St}_G^ \otimes \mathrm{St}_G \rightarrow k$ is nonzero. Then (G, H) is a Donkin pair.*

Remark 3.2 Despite the notation, the \tilde{H} -module St_H need not be an H -module. Even if St_H is not an H -module, $\mathrm{St}_H^* \otimes \mathrm{St}_H$ is one. It may be better to replace $H \subset G$ with the homomorphism $\tilde{H} \rightarrow \tilde{G}$. Thus an operation like res_H^G would really mean restriction along $\tilde{H} \rightarrow \tilde{G}$.

Remark 3.3 The pairing criterion is satisfied if and only if G/B has an H -canonical splitting. Indeed suppose we are given a map $\mathrm{St}_H^* \otimes \mathrm{St}_H = \mathrm{St}_H \otimes \mathrm{St}_H \rightarrow H^0(G/B, \omega^{1-p})$ as in Lemma 2.3. We have to factor it through the surjection $\pi : \mathrm{St}_G^* \otimes \mathrm{St}_G \rightarrow H^0(G/B, \omega^{1-p})$. But the kernel K of π has good filtration by [8] (or by the proof in [5, II 4.16]), so $\mathrm{Ext}_H^1(\mathrm{St}_H^* \otimes \mathrm{St}_H, \mathrm{res}_H^G K)$ vanishes by theorem 2.4 and the main properties of good filtrations ([8, Theorem 1], [5, II 4.13]).

Now we illustrate the criterion with some old examples of Donkin pairs.

Example 3.4 Let G still be semisimple and connected. It is easy to see from the formulas in the proof of [6, 3.2] that the pairing criterion applies to the diagonal G inside a product $G \times \cdots \times G$.

Example 3.5 Let H be the commutator subgroup of a Levi subgroup of a parabolic in the semisimple connected group G . Then, after passing to simply connected covers if necessary, St_H is a direct summand of $\mathrm{res}_H^G \mathrm{St}_G$, so again the pairing criterion applies.

Lemma 3.6 *Let (G, H) satisfy the pairing criterion and let X be a smooth projective G -variety. If X has a G -canonical splitting, then it has an H -canonical one.*

Proof Use lemma 2.3. □

The following lemma was pointed out to me by Jesper Funch Thomsen.

Lemma 3.7 *Let X, Y be smooth projective G -varieties with canonical splitting. Then $X \times Y$ has a G -canonical splitting.*

Proof Use example 3.4. □

Remark 3.8 For the users of our book, let us now point out how to get theorem 2.4. We have $G \times^B X = G/B \times X$ by remark [16, 1.2.2], so [16, lemma 4.4.2] applies with $Y = X$ in the notations of that lemma.

Remark 3.9 In lemma 3.7 one cannot replace G with B . Here is an example. Take $G = SL_3$ in characteristic 2 and let Z be the Demazure resolution of a Schubert divisor. Then $H^0(Z, \omega_Z^{-1})$ is a nine dimensional B -module. There is a fundamental representation V so that $H^0(Z, \omega_Z^{-1})$ is isomorphic to a codimension one submodule of the degree three part of the ring of regular functions on V . Using this, one checks with computer assisted computations that $Z, Z \times Z, Z \times Z \times Z$ have B -canonical splittings, while $Z \times Z \times Z \times Z$ does not have one.

Our next aim is to treat the following example.

Example 3.10 For G we take the simply connected group of type E_6 . From the symmetry of its Dynkin diagram we have a graph automorphism which is an involution. For H we take the group of fixed points of the involution. It is connected ([15, 8.2]) of type F_4 . It has been conjectured by Brundan [2, 4.4] that (G, H) is a Donkin pair.

More generally, with our usual notations we have.

Theorem 3.11 *Assume there are dominant weights $\sigma_1, \sigma_2, \sigma_3$, so that*

1. *The highest weight $(p-1)\rho_G$ of St_G equals $\sigma_1 + \sigma_2 + \sigma_3$.*
2. *$\sigma_1 + \sigma_2$ and $\sigma_2 + \sigma_3$ both restrict to the highest weight $(p-1)\rho_H$ of St_H .*
3. *The natural map $\nabla_G(\sigma_1) \rightarrow \nabla_H(\text{res}_{B \cap H}^B \sigma_1)$ is surjective.*

Then (G, H) is a Donkin pair. In fact it satisfies the pairing criterion.

Remark 3.12 If (G, H) is a Donkin pair and λ is dominant, then one knows that $\nabla_G(\lambda) \rightarrow \nabla_H(\text{res}_{B \cap H}^B \lambda) = \text{ind}_{H \cap B^-}^H(\text{res}_{H \cap B^-}^{B^-} \lambda)$, induced by the projection of $\nabla_G(\lambda)$ onto its highest weight space, is surjective. (Exercise. Use a good filtration as in the proof of [5, II 4.16].)

Remark 3.13 Our theorem 3.11 also applies to the Levi subgroup case of example 3.5 (take $\sigma_1 = 0$). One hopes to find a more general method to attack at least all graph automorphisms. Theorem 3.11 applies if the graph automorphism is an involution and different simple roots in an orbit are perpendicular to each other. But for the graph automorphism of a group of type A_{2n} in characteristic $p > 2$ there are no $\sigma_1, \sigma_2, \sigma_3$ as in the theorem. The coefficient of $\text{res}_{B \cap H}^B \rho_G$ with respect to the fundamental weight that corresponds to the short root is four, which is too high.

Proof of Theorem 3.11.

We will often write the restriction of a weight to $T \cap H$ with the same symbol as the weight. We will repeatedly use basic properties of Weyl modules and their duals. See [5, II 14.20] for surjectivity of cup product between dual Weyl modules and [5, II 2.13] for Weyl modules as universal highest weight modules. We first need a number of nonzero maps of H -modules. They are natural up to nonzero scalars that do not interest us.

The first map is the map

$$\epsilon_H : \nabla_H(2(p-1)\rho_H) \rightarrow k$$

which detects Frobenius splittings on $H/(H \cap B)$. Together with the surjection

$$\nabla_H(\sigma_2) \otimes \nabla_H((p-1)\rho_G) \rightarrow \nabla_H(2(p-1)\rho_H)$$

it gives a nonzero map $\nabla_H(\sigma_2) \otimes \nabla_H((p-1)\rho_G) \rightarrow k$ and hence a nonzero

$$\eta_1 : \nabla_H(\sigma_2) \rightarrow \nabla_H((p-1)\rho_G)^*.$$

The map $\nabla_G(\sigma_2 + \sigma_3) \rightarrow \text{St}_H$ is nonzero, hence surjective. The map $\nabla_G(\sigma_1) \rightarrow \nabla_H(\sigma_1)$ is surjective by assumption. In the commutative diagram

$$\begin{array}{ccc} \nabla_G(\sigma_2 + \sigma_3) \otimes \nabla_G(\sigma_1) & \longrightarrow & \text{St}_G \\ \downarrow & & \downarrow \\ \text{St}_H \otimes \nabla_H(\sigma_1) & \longrightarrow & \nabla_H((p-1)\rho_G) \end{array}$$

the horizontal maps are also surjective. So the map

$$\eta_2 : \nabla_H((p-1)\rho_G)^* \rightarrow \text{St}_G^*$$

is injective. We obtain a nonzero

$$\eta_2 \eta_1 : \nabla_H(\sigma_2) \rightarrow \text{St}_G^*.$$

The nonzero $\text{St}_H \rightarrow \nabla_G(\sigma_2 + \sigma_3)$ combines with the map

$$\nabla_G(\sigma_1) \otimes \nabla_G(\sigma_2 + \sigma_3) \rightarrow \text{St}_G$$

to yield

$$\nabla_G(\sigma_1) \otimes \text{St}_H \rightarrow \text{St}_G$$

and combining this with $\eta_2 \eta_1$ we get

$$\eta_3 : \nabla_H(\sigma_2) \otimes \nabla_G(\sigma_1) \otimes \text{St}_H \rightarrow \text{St}_G^* \otimes \text{St}_G.$$

We claim that its image is detected by the evaluation map

$$\eta_4 : \text{St}_G^* \otimes \text{St}_G \rightarrow k.$$

This is because η_3 factors through $\nabla_H((p-1)\rho_G)^* \otimes \text{St}_G$, on which the restriction of η_4 factors through $\nabla_H((p-1)\rho_G)^* \otimes \nabla_H((p-1)\rho_G)$, the map η_1 is nonzero, the image of $\nabla_G(\sigma_1) \otimes \text{St}_H \rightarrow \text{St}_G$ maps onto $\nabla_H((p-1)\rho_G)$.

From the nontrivial $\eta_4\eta_3$ we get a nontrivial

$$\eta_5 : \nabla_H(\sigma_2) \otimes \nabla_G(\sigma_1) \rightarrow \text{St}_H^*.$$

Then η_5 must be split surjective. Choose a left inverse

$$\eta_6 : \text{St}_H^* \rightarrow \nabla_H(\sigma_2) \otimes \nabla_G(\sigma_1)$$

of η_5 . It leads to

$$\eta_7 : \text{St}_H^* \otimes \text{St}_H \rightarrow \nabla_H(\sigma_2) \otimes \nabla_G(\sigma_1) \otimes \text{St}_H$$

and the map we use in the pairing criterion is $\eta_3\eta_7$. Indeed the map $\text{St}_H^* \rightarrow \text{St}_H^*$ defined by $\eta_4\eta_3\eta_7$ equals $\eta_5\eta_6$, hence is nonzero. \square

4 The E_6 - F_4 pair.

We turn to the E_6 - F_4 pair of example 3.10. First observe that for $p > 13$ one could simply follow the method of [2] to prove that the pair is a Donkin pair. Indeed the restriction to F_4 of a fundamental representation then has its dominant weights in the bottom alcove. Looking a little closer and applying the linkage principle one can treat $p \geq 11$ in the same manner.

But for $p = 5$ one has $\varpi_4 \uparrow \varpi_1 + \varpi_4$ and for $p = 7$ one has $\varpi_1 \uparrow \varpi_1 + \varpi_4$. This makes that one has more trouble to see that the restriction of $\nabla_G(\varpi_4)$ has a good filtration with respective layers $\nabla_H(\varpi_1)$, $\nabla_H(\varpi_3)$, $\nabla_H(\varpi_3)$, $\nabla_H(\varpi_1 + \varpi_4)$, $\nabla_H(\varpi_2)$. For $p = 2$ or $p = 3$ it is even worse.

So let us apply theorem 3.11 instead. We take $\sigma_1 = (p-1)(\varpi_1 + \varpi_3)$, $\sigma_2 = (p-1)(\varpi_2 + \varpi_4)$, $\sigma_3 = (p-1)(\varpi_5 + \varpi_6)$ in the notations of Bourbaki for E_6 [1, Planches]. Then $\text{res}_{B \cap H}^B \varpi_i$ equals ϖ_4 , ϖ_1 , ϖ_3 , ϖ_2 , ϖ_3 , ϖ_4 for $i = 1, \dots, 6$ respectively.

First let let $p = 2$. Then $\nabla_H(\text{res}_{B \cap H}^B \sigma_1) = \nabla_H(\varpi_3 + \varpi_4)$ is irreducible. Indeed its dominant weights come in two parts. The weights 0 , ϖ_4 , ϖ_1 , ϖ_3 , $2\varpi_4$, $\varpi_1 + \varpi_4$, ϖ_2 lie in one orbit, and the highest weight lies in a different orbit under the affine Weyl group. To be more specific, $\varpi_1 - \rho_H \uparrow \varpi_3 + \varpi_4$, but $\varpi_4 - \rho_H \uparrow 0 \uparrow \varpi_4 \uparrow \varpi_1 \uparrow \varpi_3 \uparrow 2\varpi_4 \uparrow \varpi_1 + \varpi_4 \uparrow \varpi_2$. So $\nabla_G(\sigma_1) \rightarrow \nabla_H(\text{res}_{B \cap H}^B \sigma_1)$ is surjective.

Remains the case $p > 2$. To see that $\nabla_G(\lambda) \rightarrow \nabla_H(\text{res}_{B \cap H}^B \lambda)$ is surjective for $\lambda = \sigma_1$, it suffices to do this for $\lambda = \varpi_1$ and $\lambda = \varpi_3$. For $p > 3$ one could now use that $\nabla_H(\text{res}_{B \cap H}^B \lambda)$ is irreducible for both $\lambda = \varpi_1$ and $\lambda = \varpi_3$,

because each of the dominant weights of $\nabla_H(\text{res}_{B \cap H}^B \lambda)$ is in a different orbit under the affine Weyl group.

But we need an argument that works for $p \geq 3$. Now $\nabla_G(\varpi_1)$ is a miniscule representation of dimension 27, and $\nabla_H(\varpi_4) = \nabla_H(\text{res}_{B \cap H}^B \varpi_1)$ has dimension 26. There are 24 short roots and they have multiplicity one in $\nabla_H(\varpi_4)$. So the map from $M := \nabla_G(\varpi_1)$ to $\nabla_H(\varpi_4)$ hits at least 24 dimensions and its kernel consists of H -invariants. Indeed there are three weights of $\nabla_G(\varpi_1)$ that restrict to zero. In Bourbaki notation they are $\zeta_1 = 1/6(\epsilon_8 - \epsilon_7 - \epsilon_6) + 1/2(-\epsilon_1 + \epsilon_2 + \epsilon_3 - \epsilon_4 - \epsilon_5)$, $\zeta_2 = 1/6(\epsilon_8 - \epsilon_7 - \epsilon_6) + 1/2(\epsilon_1 - \epsilon_2 - \epsilon_3 + \epsilon_4 - \epsilon_5)$, $\zeta_3 = -1/3(\epsilon_8 - \epsilon_7 - \epsilon_6) + \epsilon_5$. Put $\zeta_4 = 1/6(\epsilon_8 - \epsilon_7 - \epsilon_6) + 1/2(\epsilon_1 - \epsilon_2 - \epsilon_3 - \epsilon_4 + \epsilon_5)$, $\zeta_5 = 1/6(\epsilon_8 - \epsilon_7 - \epsilon_6) + 1/2(-\epsilon_1 + \epsilon_2 - \epsilon_3 + \epsilon_4 - \epsilon_5)$. Then X_{α_1} induces an isomorphism $M_{\zeta_3} \rightarrow M_{\zeta_4}$ and it annihilates $M_{\zeta_1} + M_{\zeta_2}$. Similarly X_{α_6} induces an isomorphism $M_{\zeta_2} \rightarrow M_{\zeta_4}$ and annihilates $M_{\zeta_1} + M_{\zeta_3}$. The same space is annihilated by X_{α_3} , which induces an isomorphism $M_{\zeta_2} \rightarrow M_{\zeta_5}$. Finally X_{α_5} induces an isomorphism $M_{\zeta_1} \rightarrow M_{\zeta_5}$ and annihilates $M_{\zeta_2} + M_{\zeta_3}$.

It follows that in $M_{\zeta_1} + M_{\zeta_2} + M_{\zeta_3}$ there is just a one dimensional subspace of vectors annihilated by both $X_{\alpha_1} + X_{\alpha_6}$ and $X_{\alpha_3} + X_{\alpha_5}$. (These two operators come from the Lie algebra of H .) We conclude that $\text{res}_H^G M$ has a good filtration and that $M \rightarrow \nabla_H(\varpi_4)$ is surjective. As $p > 2$, we then also have that $M \wedge M$ and $\text{res}_H^G(M \wedge M)$ have a good filtration. It then follows from the character that $M \wedge M = \nabla_G(\varpi_3)$. (We use the program LiE [17].) So $\text{res}_H^G \nabla_G(\varpi_3)$ has a good filtration and therefore maps onto $\nabla_G(\text{res}_{B \cap H}^B \varpi_3)$.

Summing up, we have shown

Theorem 4.1 *The E_6 - F_4 pair is a Donkin pair. In fact it satisfies the pairing criterion.*

Remark 4.2 When Steve Donkin received this proof, he proceeded to show that one could also prove E_6 - F_4 to be a Donkin pair with the ‘ancient methods’ of his book [3]. Of course he had to treat more representations than we do. We will use his method in the last section to treat the remaining cases of Brundan’s conjecture in characteristic $p > 2$, where we have no alternative yet.

5 Induction and canonical splitting

We finish the discussion of canonical splittings with an analogue of proposition [9, 5.5]. It makes a principle from [8] more explicit. The result was explained to us by O. Mathieu at a reception of the mayor of Aarhus in August 1998. It shows once more that canonical splittings combine well with Demazure desingularisation of Schubert varieties.

Proposition 5.1 *Let X be a projective B -variety with canonical splitting. Let P be a minimal parabolic. Then $P \times^B X$ has a canonical splitting.*

Corollary 5.2 *The same conclusion holds for any parabolic subgroup.*

Proof If P is not minimal, take a Demazure resolution $Z = P_1 \times^B P_2 \times^B \dots \times^B P_r / B$ of P/B and apply the proposition to get a canonical splitting on $P_1 \times^B P_2 \times^B \dots \times^B P_r \times^B X$. Then push the splitting forward ([10, Prop. 4]) to $P \times^B X$. \square

Proof of Proposition We use notations as in [16, Ch. 4, A.4]. Let ζ be the highest weight of St and s the simple reflection corresponding with P . One checks as in [16, A.4.6] that

$$\mathcal{E}nd_F(P \times^B X) = (P \times^B \mathcal{E}nd_F(X)) \otimes \pi^* \mathcal{L}(s\zeta - \zeta),$$

where $\pi : P \times^B X \rightarrow P/B$. We are given a map $\phi : k_\zeta \otimes \text{St} \rightarrow \text{End}_F(X)$. The required map $\psi : k_\zeta \otimes \text{St} \rightarrow \text{End}_F(P \times^B X)$ may be constructed by composing maps

$$\begin{aligned} k_\zeta \otimes \text{St} &\cong \\ k_{-s\zeta} \otimes \text{ind}_B^P(k_{\zeta+s\zeta} \otimes \text{St}) &\rightarrow \\ k_{-s\zeta} \otimes \text{ind}_B^P(k_{s\zeta} \otimes \text{End}_F(X)) &\cong \\ k_{-s\zeta} \otimes H^0(P \times^B X, P \times^B (\mathcal{E}nd_F(X)[s\zeta])) &\cong \\ \text{End}_F(P \times^B X, B \times^B X) &\rightarrow \\ \text{End}_F(P \times^B X) & \end{aligned}$$

Here $k_{-s\zeta}$ is identified with the weight space of weight $-s\zeta$ of

$$H^0(P \times^B X, \pi^* \mathcal{L}(-\zeta)).$$

An element of that weight space has divisor $(p-1)B \times^B X = (p-1)X$.

To see that the image of ψ is not in the kernel of

$$\epsilon_{P \times^B X} : \text{End}_F(P \times^B X) \rightarrow k,$$

it suffices to show that the diagram

$$\begin{array}{ccccc} k_\zeta \otimes \text{St} & \rightarrow & \text{End}_F(P \times^B X, B \times^B X) & \rightarrow & \text{End}_F(P \times^B X) \\ \parallel & & \downarrow & & \downarrow \\ k_\zeta \otimes \text{St} & \xrightarrow{\phi} & \text{End}_F(X) & \rightarrow & k \end{array}$$

commutes. Now

$$\begin{array}{ccc} k_{-s\zeta} \otimes \text{ind}_B^P(k_{\zeta+s\zeta} \otimes \text{St}) & \longrightarrow & k_\zeta \otimes \text{St} \\ \downarrow & & \downarrow \\ k_{-s\zeta} \otimes \text{ind}_B^P(k_{s\zeta} \otimes \text{End}_F(X)) & \longrightarrow & \text{End}_F(X) \end{array}$$

commutes and by restricting to the trivial fibration $BsB \times^B X \rightarrow BsB/B$ one shows through the following lemma that the bottom map in this last diagram agrees with the map that factors through $\text{End}_F(P \times^B X, B \times^B X)$. \square

Lemma 5.3 *Let A be a commutative k -algebra. Then*

$$\text{End}_F(A[t]) = \text{End}_F(A) \otimes \text{End}_F(k[t]) = \text{End}_F(A) \otimes k[t]$$

and the map $\text{End}_F(A[t], (t)) \rightarrow \text{End}_F(A)$ is induced by the map

$$t^{p-1}k[t] = t^{p-1} * \text{End}_F(k[t]) = \text{End}_F(k[t], (t)) \rightarrow \text{End}_F(k) = k$$

which sends $t^{p-1}f(t)$ to $f(0)$.

Proof Straightforward, provided one keeps in mind how $\text{End}_F(R)$ is an R -module ([16, 4.3.3]). Compare also [16, A.4.5]. \square

6 More Donkin pairs

In this section we do not use the pairing criterion. Instead we return to the methods of Donkin's book [3], combined with computer calculations of characters, of linkage, and of the Jantzen sum formula.

Let G , H be as before, with G simply connected. In fact H will be the commutator subgroup of the group of fixed points of an involution of G which leaves invariant the maximal torus T and the Borel subgroup B . We refer to [14] for the classification of the possibilities, assuming $p > 2$. (Of course involutions of the simply connected G are lifted [15, 9.16] from the involutions of the corresponding adjoint group, which are treated in [14].)

Remark 6.1 Let H be the fixed point group of an involution that leaves T and B invariant in the simply connected semisimple G . Then H is connected reductive by [15, 8.2]. Now an H -module has good filtration in the sense of [3] if and only if its restriction to the commutator subgroup of H has good filtration. That is why we look only at semisimple subgroups H .

Let \mathcal{M} denote the set of finite dimensional G -modules M with good filtration for which $\text{res}_H^G M$ has good filtration. Let \mathcal{S} denote the set of dominant weights λ of G so that $\nabla_G(\lambda) \in \mathcal{M}$. As always we try to show that all dominant weights of G are in \mathcal{S} . For this purpose we recall some useful lemmas.

Lemma 6.2 1. If $M_1 \oplus M_2 \in \mathcal{M}$, then $M_1 \in \mathcal{M}$.

2. If $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ is exact, and $M' \in \mathcal{M}$, then $M \in \mathcal{M}$ if and only if $M'' \in \mathcal{M}$.

3. If $M_1, M_2 \in \mathcal{M}$, then $M_1 \otimes M_2 \in \mathcal{M}$.

If M is a G -module with good filtration, write $\text{supp}_\nabla(M)$ for the set of dominant weights λ so that $\nabla_G(\lambda)$ occurs as a layer in a good filtration of M . We order the dominant weights of G by the partial order in which $\mu \leq \lambda$ if and only if $\lambda - \mu$ is in the closed cone spanned by the positive roots. In particular, if λ is dominant, then $0 \leq \lambda$, and all dominant weights μ of $\nabla_G(\lambda)$ satisfy $\mu \leq \lambda$. We say that a filtration of M is a good filtration *adapted* to the partial order if there are λ_i so that the i -th layer is a direct sum of copies of $\nabla_G(\lambda_i)$, and $i \leq j$ if $\lambda_i \leq \lambda_j$. (So we still call it a good filtration, even though $\nabla_G(\lambda_i)$ may have multiplicity in the i -th layer.) If M has a good filtration, then it also has one adapted to the partial order, by the proof of [5, II 4.16].

Lemma 6.3 Let $M \in \mathcal{M}$ and $\lambda \in \text{supp}_\nabla(M)$. Assume for every weight μ in $\text{supp}_\nabla(M)$, distinct from λ , that one of the following holds

1. $\mu < \lambda$ and $\mu \in \mathcal{S}$.
2. μ and λ are in different orbits under the affine Weyl group.

Then $\lambda \in \mathcal{S}$.

Proof We may replace M by an indecomposable direct summand M_1 with $\lambda \in \text{supp}_\nabla(M_1)$. The linkage principle tells that we thus get rid of the second possibility in the lemma. Then in a good filtration adapted to the partial order, the module $\nabla_G(\lambda)$ occurs only as a summand of the top layer, which is in \mathcal{M} by lemma 6.2. \square

Lemma 6.4 *Let λ be a dominant weight of G . If λ is in the bottom alcove, or if the Jantzen sum formula yields zero, then $\nabla_G(\lambda)$ is irreducible.*

Proof See [5, II Cor. 5.6 and 8.21] \square

6.5 The pairs E_8 , D_8 and E_8 , E_7A_1

Say G is of type E_8 in characteristic $p > 2$ and H is the fixed point group of an involution. There are two cases, up to conjugacy. One may have H of type D_8 or one may have H of type E_7A_1 .

In either case we wish to show that G, H is a Donkin pair. In other words, we want that all dominant weights are in \mathcal{S} . We will argue by induction along the partial order. Thus when trying to prove that $\lambda \in \mathcal{S}$, we shall always assume that $\mu \in \mathcal{S}$ for $\mu < \lambda$. Of course the zero weight is in \mathcal{S} , so say λ is nonzero. If λ is not a fundamental weight, write $\lambda = \lambda_1 + \lambda_2$ where λ_i are nonzero dominant weights. As $\lambda_i < \lambda$, we may apply lemma 6.3 with $M = \nabla_G(\lambda_1) \otimes \nabla_G(\lambda_2)$ to conclude that $\lambda \in \mathcal{S}$.

Remain the fundamental weights. Observe that $\varpi_8 < \varpi_1 < \varpi_7 < \varpi_2 < \varpi_6 < \varpi_3 < \varpi_5 < \varpi_4$. But we will not discuss them in this exact order.

To see that $\varpi_8 \in \mathcal{S}$ we compute the character of $\text{res}_H^G \nabla_G(\varpi_8)$ with the program LiE [17], decompose this character in terms of Weyl characters, and use lemma 6.4 to see that $\text{res}_H^G \nabla_G(\varpi_8)$ has a composition series whose factors are irreducible (dual) Weyl modules. Here we use a Java applet of Lauritzen for the Jantzen sum formula.

To see that $\varpi_1 \in \mathcal{S}$ we may argue the same way if H is of type D_8 . If H is of type E_7A_1 , let K be the subgroup of type E_7 in H , and F the

subgroup of type A_1 . Then G, K is a Donkin pair (Levi subgroup case), so we may consider a good filtration adapted to the partial order (on weights for K) of $\text{res}_K^G \nabla_G(\varpi_1)$. It is a filtration by H -modules. It suffices to show that its layers have good filtration as H -modules. Let N be such a layer. Its character is the character of some $\nabla_H(\lambda_1, \lambda_2)$ where λ_1 is a dominant weight for K and λ_2 is one for F . (From now on we do not mention the computer calculations that are needed to support such statements.) Moreover, $\nabla_F(\lambda_2)$ is irreducible, so that the $K \cap B$ -socle of N is an irreducible F -module. It follows that the natural map $N \rightarrow \nabla_H(\lambda_1, \lambda_2)$ is an isomorphism, and thus $\varpi_1 \in \mathcal{S}$.

To see that $\varpi_2 \in \mathcal{S}$, we apply lemma 6.3 with $M = \nabla_G(\varpi_1) \otimes \nabla_G(\varpi_8)$. (Note $\varpi_1, \varpi_8 < \varpi_2$, so that indeed $M \in \mathcal{M}$ by the inductive assumption.) One may find the necessary statement about linkage in Donkin's book. (This is no accident, as we follow him in our choices.) We also checked the non-linkage with a straightforward Mathematica program.

To see that $\varpi_7 \in \mathcal{S}$, we similarly use $M = \nabla_G(\varpi_8) \wedge \nabla_G(\varpi_8)$. (Recall $p > 2$.) To get $\varpi_3 \in \mathcal{S}$, use $M = \nabla_G(\varpi_1) \wedge \nabla_G(\varpi_1)$. To get $\varpi_4 \in \mathcal{S}$, use $M = \nabla_G(\varpi_2) \wedge \nabla_G(\varpi_2)$. To get $\varpi_5 \in \mathcal{S}$, use $M = \nabla_G(\varpi_1) \otimes \nabla_G(\varpi_2)$ if $p = 3$, and $M = \wedge^4 \nabla_G(\varpi_8)$ if $p > 3$. To get $\varpi_6 \in \mathcal{S}$, use $M = \nabla_G(\varpi_1) \otimes \nabla_G(\varpi_1)$ if $p = 3$, and $M = \wedge^3 \nabla_G(\varpi_8)$ if $p > 3$.

6.6 The pair E_6, A_5A_1

Let G be the simply connected group of type E_6 in characteristic $p > 2$ and let H be the fixed point group of such an inner involution that H is of type A_5A_1 and the involution commutes with the graph automorphism. We wish to show again this is a Donkin pair. We argue as in the E_8, E_7A_1 case.

If $\lambda = \varpi_1$ or ϖ_2 , then we argue with socles as we did to show $\varpi_1 \in \mathcal{S}$ for the E_8, E_7A_1 pair.

To treat ϖ_3 we use $M = \nabla_G(\varpi_1) \wedge \nabla_G(\varpi_1)$. To get $\varpi_4 \in \mathcal{S}$, use $M = \nabla_G(\varpi_2) \wedge \nabla_G(\varpi_2)$. The remaining two fundamental weights are in \mathcal{S} by symmetry.

6.7 The pair E_6, C_4

Let G be the simply connected group of type E_6 in characteristic $p > 2$ and let H be the fixed point group of such an outer involution that H is of type C_4 and the involution commutes with the graph automorphism.

The module $\nabla_G(\varpi_2)$ is the Lie algebra of the adjoint form of G . Its restriction $\text{res}_H^G \nabla_G(\varpi_2)$ has a six dimensional weight space for the weight zero, just like $\nabla_G(\varpi_2)$ itself. We claim that it contains no nonzero invariant. Indeed we may choose the involution so that $X_{\alpha_3} + X_{\alpha_5}$, $X_{\alpha_1} + X_{\alpha_6}$, $X_{\alpha_3+\alpha_4} + X_{\alpha_5+\alpha_4}$, X_{α_2} are in the Lie algebra of H , where we have put $X_{\alpha_3+\alpha_4} = [X_{\alpha_3}, X_{\alpha_4}]$ and $X_{\alpha_5+\alpha_4} = [X_{\alpha_4}, X_{\alpha_5}]$. The only element in the weight zero weight space of $\nabla_G(\varpi_2)$ that is annihilated by all these elements is the zero vector. Now $\text{res}_H^G \nabla_G(\varpi_2)$ contains an irreducible $\nabla_H(2\varpi_1)$ and the quotient by that submodule is either irreducible, or $p = 3$ and there are two composition factors, one of which is one dimensional. (This also uses the Jantzen sum formula.) As there is no invariant in $\text{res}_H^G \nabla_G(\varpi_2)$ and there is no extension between $\nabla_H(2\varpi_1)$ and the other composition factors ([5, II 4.13, 4.14]), we get $\varpi_2 \in \mathcal{S}$.

As $\text{res}_H^G \nabla_G(\varpi_1)$ is irreducible, we also have $\varpi_1 \in \mathcal{S}$. The rest goes as for the pair E_6 , A_5A_1 .

6.8 The pairs E_7 , A_7 and E_7 , D_6A_1

Say G is simply connected of type E_7 in characteristic $p > 2$ and H is the fixed point group of an involution. There are two cases, up to conjugacy. One may have H of type A_7 or one may have H of type D_6A_1 .

We argue as before. If H is of type D_6A_1 we show that $\varpi_1, \varpi_2, \varpi_7 \in \mathcal{S}$ by the argument with socles used to show $\varpi_1 \in \mathcal{S}$ for the E_8 , E_7A_1 pair. If H is of type A_7 we have $\varpi_1, \varpi_7 \in \mathcal{S}$ for the same reason, involving the sum formula, as why $\varpi_8 \in \mathcal{S}$ for the E_8 , D_8 pair. If $p = 7$ we see in the same manner that $\varpi_2 \in \mathcal{S}$. If $p \neq 7$ use $M = \nabla_G(\varpi_1) \otimes \nabla_G(\varpi_7)$ to get $\varpi_2 \in \mathcal{S}$. To get $\varpi_3 \in \mathcal{S}$ use $M = \nabla_G(\varpi_1) \wedge \nabla_G(\varpi_1)$. To get $\varpi_4 \in \mathcal{S}$ use $M = \nabla_G(\varpi_2) \wedge \nabla_G(\varpi_2)$. To get $\varpi_5 \in \mathcal{S}$ use $M = \wedge^3 \nabla_G(\varpi_7)$ if $p \neq 3$ and $M = \nabla_G(\varpi_1) \wedge \nabla_G(\varpi_2)$ otherwise. To get $\varpi_6 \in \mathcal{S}$ use $M = \nabla_G(\varpi_7) \wedge \nabla_G(\varpi_7)$.

6.9 The pairs F_4 , B_4 and F_4 , C_3A_1

Say G is of type F_4 in characteristic $p > 2$ and H is the fixed point group of an involution. There are two cases, up to conjugacy. One may have H of type B_4 or one may have H of type C_3A_1 .

We argue as before. If H is of type C_3A_1 we show that $\varpi_1, \varpi_4 \in \mathcal{S}$ by the argument with socles used to show $\varpi_1 \in \mathcal{S}$ for the E_8 , E_7A_1 pair. If H is of type B_4 we have $\varpi_1, \varpi_4 \in \mathcal{S}$ for the same reason, involving the

sum formula, as why $\varpi_8 \in \mathcal{S}$ for the E_8 , D_8 pair. To get $\varpi_3 \in \mathcal{S}$ use $M = \nabla_G(\varpi_4) \wedge \nabla_G(\varpi_4)$. To get $\varpi_2 \in \mathcal{S}$ use $M = \nabla_G(\varpi_1) \wedge \nabla_G(\varpi_1)$.

Theorem 6.10 (Brundan’s Conjecture [2]) *Let G be semisimple simply connected. If either*

- (i) *H is the centralizer of a graph automorphism of G ; or*
- (ii) *H is the centralizer of an involution of G and the characteristic is at least three,*

then G, H is a Donkin pair.

Proof

We have either a Levi subgroup case, first settled in [3] (see also remarks 3.5, 6.1), or a case treated in [2], or a case treated above, up to conjugacy. \square

Remark 6.11 Of course we would much prefer a case-free proof, based on the pairing criterion say.

References

- [1] N. Bourbaki, Groupes et Algèbres de Lie, Ch. 4, 5 et 6, Paris: Hermann 1968.
- [2] J. Brundan, Dense orbits and double cosets, in: Carter, R. W. (ed.) et al., Algebraic groups and their representations (Cambridge, 1997), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci. 517, (1998), 259–274.
- [3] S. Donkin, Rational Representations of Algebraic Groups: tensor products and filtrations, Lecture Notes in Mathematics 1140, Berlin: Springer 1985.
- [4] E. Friedlander, A canonical filtration for certain rational modules, Math. Z. 188 (1985), 433–438.
- [5] J.-C. Jantzen, Representations of Algebraic Groups, Pure and Applied Mathematics v. 131, Boston: Academic Press 1987.

- [6] N. Lauritzen, J. F. Thomsen, Frobenius splitting and hyperplane sections of flag manifolds, *Inv. math.* 128 (1997), 437–442.
- [7] O. Mathieu, Filtrations of B -modules, *Duke Math. Journal* 59 (1989), 421–442.
- [8] O. Mathieu, Filtrations of G -modules, *Ann. Sci. École Norm. Sup.* 23 (1990), 625–644.
- [9] O. Mathieu, Tilting modules and their applications, preprint.
- [10] V.B. Mehta and A. Ramanathan, Frobenius splitting and cohomology vanishing for Schubert varieties, *Annals of Math.* 122 (1985), 27–40.
- [11] V.B. Mehta and T.N. Venkataramana, A note on Steinberg modules and Frobenius splitting, *Inv. math.* 123 (1996), 467–469.
- [12] P. Polo, Variétés de Schubert et excellentes filtrations, *Astérisque* 173–174 (1989), 281–311.
- [13] S. Ramanan and A. Ramanathan, Projective normality of flag varieties and Schubert varieties, *Inv. math.* 79 (1985), 217–224. *Math.* 122 (1985), 27–40.
- [14] T.A. Springer, The classification of involutions of simple algebraic groups, *J. Fac. Sci., Univ. Tokyo, Sect. I A* 34 (1987), 655–670.
- [15] R. Steinberg, Endomorphisms of linear algebraic groups, *Memoirs of the Amer. Math. Soc.* 80 (1968).
- [16] W. van der Kallen, Lectures on Frobenius splittings and B -modules. Notes by S. P. Inamdar, Tata Institute of Fundamental Research, Bombay, and Springer-Verlag, Berlin, 1993.
- [17] M.A.A. van Leeuwen, A.M. Cohen, B. Lissers, LiE, A package for Lie group computations, *Computer Algebra Netherlands*, Amsterdam, 1992.