

HIGHER ORBITAL INTEGRALS, SHALIKA GERMS, AND THE HOCHSCHILD HOMOLOGY OF HECKE ALGEBRAS

VICTOR NISTOR¹

Version: 3.0A; Revised: 8/16/2000; Run: September 8, 2017

ABSTRACT. We give a detailed calculation of the Hochschild and cyclic homology of the algebra $\mathcal{C}_c^\infty(G)$ of locally constant, compactly supported functions on a reductive p -adic group G . We use these calculations to extend to arbitrary elements the definition the higher orbital integrals introduced by Blanc and Brylinski for regular semisimple elements. Then we extend to higher orbital integrals some results of Shalika. We also investigate the effect of the “induction morphism” on Hochschild homology.

CONTENTS

Introduction	1
1. Homology of Hecke algebras	3
2. Higher orbital integrals and their Shalika germs	10
3. The cohomology of the unipotent variety	15
4. Induction and the unipotent variety	17
5. Examples	23
References	26

INTRODUCTION

Orbital integrals play an important role in the harmonic analysis of a reductive p -adic group G ; they are, for instance, one of the main ingredients in the Arthur–Selberg trace formula. Orbital integrals on unimodular groups are a particular case of invariant distribution, which have been used in [2] to prove the irreducibility of certain induced representations of GL_n over a p -adic field.

By definition, an invariant distribution on a unimodular group G gives rise to a trace (i.e., a Hochschild cocycle of degree zero) on $\mathcal{C}_c^\infty(G)$, the Hecke algebra of compactly supported, locally constant, complex valued functions on G . It is interesting then to try to analyse as completely as possible the Hochschild homology and cohomology groups of the algebra $\mathcal{C}_c^\infty(G)$ (denoted $\mathrm{HH}_*(\mathcal{C}_c^\infty(G))$ and $\mathrm{HH}^*(\mathcal{C}_c^\infty(G))$, respectively).

In this paper, G will be the set of \mathbb{F} -rational points of a linear algebraic group \mathbb{G} defined over a finite extension \mathbb{F} of the field \mathbb{Q}_p of p -adic numbers, p being a fixed prime number. The group \mathbb{G} does not have to be reductive, although this is

¹partially supported by a NSF Young Investigator Award DMS-9457859 and a Sloan Research Fellowship. Manuscripts are available from <http://www.math.psu.edu/nistor/>.

certainly the most interesting case. When we shall assume \mathbb{G} (or G , by abuse of language) to be reductive, we shall state this explicitly. For us, the most important topology to consider on G will be the locally compact topology induced from an embedding of $G \subset GL_n(\mathbb{F})$. Nevertheless, the Zariski topology on G will also play a role in our study.

To state the main result of this paper, we need to introduce first the concept of standard subgroup. For any set $A \subset G$, we shall denote

$$C(A) := \{g \in G, ga = ag, \forall a \in A\}$$

and $Z(A) := A \cap C(A)$. This latter notation will be used only when A is a subgroup of G . A commutative subgroup S of G is called *standard* if $S = Z(C(s))$ for some semi-simple element $s \in G$. Our results will be stated in terms of standard subgroups of G . We shall denote by H_u the set of unipotent elements of a subgroup H . Sometimes, the set $C(S)_u$ is also denoted by \mathcal{U}_S , in order to avoid having to many paranthesis in our formulae. One of the main results of this paper (Theorem 1) identifies the groups $\mathrm{HH}_*(\mathcal{C}_c^\infty(G))$ in terms of the following data: the set Σ of (conjugacy classes of) standard subgroups S of G , the subset $S^{\mathrm{reg}} \subset S$ of S -regular elements, the action of the Weyl group $W(S)$ of S on $\mathcal{C}_c^\infty(S)$, and the continuous cohomology of the $C(S)$ -module

$$\mathcal{C}_c^\infty(\mathcal{U}_S)_\delta := \mathcal{C}_c^\infty(C(S)_u) \otimes \Delta_{C(S)},$$

where $\Delta_{C(S)}$ denotes the modular function of the group $C(S)$. More precisely, if G is a p -adic group defined over a field of characteristic zero, as before, then Theorem 1 states the existence of an isomorphism

$$(1) \quad \mathrm{HH}_q(\mathcal{C}_c^\infty(G)) \simeq \bigoplus_{S \in \Sigma} \mathcal{C}_c^\infty(S^{\mathrm{reg}})^{W(S)} \otimes \mathrm{H}_q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta),$$

which can be made natural by using a generalization of the Shalika germs.

It is important to relate this result with the periodic cyclic homology groups of $\mathcal{C}_c^\infty(G)$. For the Hecke algebra $\mathcal{C}_c^\infty(G)$, the periodic cyclic homology is related to Hochschild homology by

$$(2) \quad \mathrm{HP}_q(\mathcal{C}_c^\infty(G)) \simeq \bigoplus_{k \in \mathbb{Z}} \mathrm{HH}_{q+2k}(\mathcal{C}_c^\infty(G))_{\mathrm{comp}},$$

that is, $\mathrm{HP}_*(\mathcal{C}_c^\infty(G))$ is the localization of Hochschild homology to the G -invariant subset of compact elements of G . This relation is implicit in [10]. Consequently, the results of this paper complement the results on the cyclic homology of p -adic groups in [10, 19]. It is interesting to remark that $\mathrm{HP}_*(\mathcal{C}_c^\infty(G))$ can also be described in terms of the admissible spectrum of G , see [12], and hence our results have significance for the representation theory of p -adic groups. See also [16] for similar results on (the groups of real points of) algebraic groups defined over \mathbb{R} . These periodic cyclic cohomology groups are isomorphic to $K_*(C_r^*(G))$, by combining results from [1], [13], and [10].

Assume for the moment that G is reductive. Then, in order to better understand the role played by the groups $\mathrm{HH}_*(\mathcal{C}_c^\infty(G))$ and $\mathrm{H}_*(G, \mathcal{C}_c^\infty(G_u))$ in the representation theory of G , we relate $\mathrm{H}_*(G, \mathcal{C}_c^\infty(G_u))$ to the analogous cohomology groups, $\mathrm{H}_*(P, \mathcal{C}_c^\infty(P_u)_\delta)$ and $\mathrm{H}_*(M, \mathcal{C}_c^\infty(M_u)_\delta)$, associated to parabolic subgroups P of G and to their Levi components M . In particular, we define morphisms between these Hochschild homology groups that are analogous to the induction and inflation morphisms that play such a prominent role in the representation theory of p -adic groups. These morphisms are induced by morphisms of algebras.

In [4], Blanc and Brylinski have introduced higher orbital integrals associated to regular semisimple elements by proving first that

$$(3) \quad \mathrm{HH}_q(\mathcal{C}_c^\infty(G)) \simeq \mathrm{H}_q(G, \mathcal{C}_c^\infty(G)_\delta),$$

a result which they dubbed “the MacLane isomorphism.” (Actually, they did not have to twist with the modular function, because they worked only with unimodular groups G , see Lemma 1 for the slightly more general version needed in this paper), after that we rely more on filtrations of the G -module $\mathcal{C}_c^\infty(G)$, rather than on localization. This allows us to define higher orbital integrals at arbitrary elements. Then, we study the properties of these orbital integrals and we obtain in particular a proof of the existence of abstract Shalika germs for the higher orbital integrals. Actually, the existence of Shalika germs turns out to be a consequence of some general homological properties of the ring of invariant, locally constant functions on the group G . We also use the techniques developed in [16] in the framework of real algebraic groups. It would be interesting to relate the results of this paper to those of [11].

This is the revised version of an preprint that was first circulated in February 1999. I would like to thank Paul Baum, David Kazhdan, Robert Langlands, George Lusztig, Roger Plymen, and Peter Schneider for useful comments and discussions.

1. HOMOLOGY OF HECKE ALGEBRAS

In this section we shall use several general results on Hochschild homology of algebras, on algebraic groups, and on the continuous cohomology of totally disconnected groups. Good references are [5, 6, 15], for the general theory, and [12] for questions related to Hochschild homology.

If G is a group and $A \subset G$ is a subset, we denote by $C(A)$ the *centralizer* of A , that is, the set of elements of G that commute with every element of A , and by $N(A)$ the *normalizer* of A , that is, the set of elements $g \in G$ such that $gAg^{-1} = A$. By $Z = Z(G) = C(G)$ we denote the center of G .

If X is a totally disconnected, locally compact, non-discrete space X , we denote by $\mathcal{C}_c^\infty(X)$ the space of compactly supported, locally constant, complex valued functions on X . Recall that, if $U \subset X$ is an open subset of X as above, then restriction defines an isomorphism

$$(4) \quad \mathcal{C}_c^\infty(X)/\mathcal{C}_c^\infty(U) \simeq \mathcal{C}_c^\infty(X \setminus U).$$

Let \mathbb{G} be a linear algebraic group defined over a totally disconnected locally compact field \mathbb{F} . Thus \mathbb{F} is a finite algebraic extension of \mathbb{Q}_p , the field of p -adic numbers. The set $\mathbb{G}(\mathbb{F})$ of \mathbb{F} -rational points of \mathbb{G} is called a p -adic group and will be denoted simply by G . It is known [5] that $G = \mathbb{G}(\mathbb{F})$ identifies with a closed subgroup of $GL_n(\mathbb{F})$, and hence it has a natural locally compact topology that makes it a totally disconnected space. We fix a Haar measure dg on G .

Consider now the space $\mathcal{C}_c^\infty(G)$ of compactly supported, locally constant functions on G . Fix a Haar measure dh on G . Then the convolution product, denoted $*$, is defined by

$$f_1 * f_2(g) = \int_G f_1(h) f_2(h^{-1}g) dh,$$

makes $\mathcal{C}_c^\infty(G)$ an algebra, the *Hecke algebra* of G . It is important in representation theory to determine the (Ad_G-) invariant linear functionals on $\mathcal{C}_c^\infty(G)$. If G is

unimodular, the space of invariant linear functionals on $\mathcal{C}_c^\infty(G)$ coincides with the space of traces on $\mathcal{C}_c^\infty(G)$; moreover, since the space of traces of $\mathcal{C}_c^\infty(G)$ identifies with $\mathrm{HH}^0(\mathcal{C}_c^\infty(G))$, the first Hochschild cohomology group of $\mathcal{C}_c^\infty(G)$, it is reasonable to ask what are the groups $\mathrm{HH}^q(\mathcal{C}_c^\infty(G))$, the Hochschild homology groups of $\mathcal{C}_c^\infty(G)$, in general. Since $\mathrm{HH}^q(\mathcal{C}_c^\infty(G))$ is the algebraic dual of $\mathrm{HH}_q(\mathcal{C}_c^\infty(G))$, it is enough to concentrate on Hochschild homology. The computation of the groups $\mathrm{HH}_q(\mathcal{C}_c^\infty(G))$ is the main purpose of this paper.

We first recall the definition of $\mathrm{HH}_*(\mathcal{C}_c^\infty(G))$. Let

$$\mathcal{C}_c^\infty(G^{q+1}) = \mathcal{C}_c^\infty(G) \otimes \mathcal{C}_c^\infty(G) \otimes \dots \otimes \mathcal{C}_c^\infty(G),$$

$(q+1)$ -times, be the usual (algebraic) tensor product of vector spaces. The Hochschild differential $b : \mathcal{C}_c^\infty(G^{q+2}) \rightarrow \mathcal{C}_c^\infty(G^{q+1})$ is given by

$$(5) \quad (bf)(g_0, g_1, \dots, g_q) = \sum_{j=0}^q (-1)^j \int_G f(g_0, \dots, g_{j-1}, \gamma, \gamma^{-1}g_j, g_{j+1}, \dots, g_q) d\gamma \\ + (-1)^{q+1} \int_G f(\gamma^{-1}g_0, g_1, \dots, g_q, \gamma) d\gamma.$$

By definition, the q -th Hochschild homology group of $\mathcal{C}_c^\infty(G)$, denoted $\mathrm{HH}_q(\mathcal{C}_c^\infty(G))$, is the q -th homology group of the complex $(\mathcal{C}_c^\infty(G^{q+1}), b)$. Since $\mathcal{C}_c^\infty(G)$ is an inductive limit of unital algebras, this definition coincides with the usual definition of Hochschild homology for non-unital algebras (using algebras with adjoint unit).

The group G acts by conjugation on $\mathcal{C}_c^\infty(G)$, and we denote by $\mathcal{C}_c^\infty(G)_{\mathrm{ad}}$ the G -module defined by this action. Also, let Δ_G denote the modular function of G , which we recall, is defined by the relation

$$\Delta_G(h) \int_G f(gh) dg = \int_G f(g) dg.$$

We shall be especially interested in the G -module $\mathcal{C}_c^\infty(G)_\delta$ obtained from $\mathcal{C}_c^\infty(G)_{\mathrm{ad}}$ by twisting it with the modular function. More precisely, let $\mathcal{C}_c^\infty(G)_\delta = \mathcal{C}_c^\infty(G)$ as vector spaces, and let the action of G on functions be given by the formula

$$(6) \quad (\gamma \cdot f)(g) = \Delta_G(\gamma) f(\gamma^{-1}g\gamma), \quad f \in \mathcal{C}_c^\infty(G)_\delta.$$

The reason for this twisting is that, for G non-unimodular, the traces of $\mathcal{C}_c^\infty(G)$ are the G -invariant functionals on $\mathcal{C}_c^\infty(G)_\delta$, not on $\mathcal{C}_c^\infty(G)$ (this is an immediate consequence of Lemma 1). More generally, our approach to the Hochschild homology of $\mathcal{C}_c^\infty(G)$ is based on Lemma 1.

Before stating and proving Lemma 1, we need to introduce some notation. First, if M is an arbitrary G -module, we denote by $M \otimes \Delta_G$ the tensor product of the G -modules M and \mathbb{C} , where the action on \mathbb{C} is given by the multiplication with the modular function of G . (We sometimes write this as $\mathcal{C}_c^\infty(G)_\delta = \mathcal{C}_c^\infty(G) \otimes \Delta_G$.)

If M is a right G -module and M' is a left G -module, then $M \otimes_G M'$ is defined as the quotient of $M \otimes M'$ by the submodule generated by $mg \otimes m' - m \otimes gm'$. For example, if $H \subset G$ is a closed subgroup and if X is a left H -space, then we have an isomorphism of G -spaces

$$(7) \quad \mathcal{C}_c^\infty(G) \otimes_H (\mathcal{C}_c^\infty(X) \otimes \Delta_H) \simeq \mathcal{C}_c^\infty(G \times_H X),$$

where $G \times X$ is the quotient $(G \times X)/H$ for the action $h(g, x) = (gh^{-1}, hx)$. This isomorphism is obtained by observing that the natural map

$$t_X : \mathcal{C}_c^\infty(G) \otimes \mathcal{C}_c^\infty(X) = \mathcal{C}_c^\infty(G \times X) \longrightarrow \mathcal{C}_c^\infty(G \times_H X),$$

$$t_X(f)(\overline{(g, x)}) = \int_H f(gh, h^{-1}x)dh,$$

passes to the quotient to give the desired isomorphism. Sometimes it will be convenient to regard a left G -module as a right G -module by replacing g with g^{-1} .

Also, recall that a G -module M is *smooth* if, and only if, the stabilizer of each element of M is open in G . Then one can define the *continuous homology* groups of G with coefficients in the smooth module M , denoted $H_k(G, M)$, using tensor products as follows. Let $\mathcal{B}_q(G) = \mathcal{C}_c^\infty(G^{q+1})$, $q = 0, 1, \dots$, be the Bar complex of the group G , with differential

$$(df)(g_0, g_1, \dots, g_q) = \sum_{j=0}^{q+1} (-1)^j \int_G f(g_0, \dots, g_{j-1}, \gamma, g_j, \dots, g_q) d\gamma.$$

Then the complex (\mathcal{B}_q, d) gives a resolution of \mathbb{C} with projective $\mathcal{C}_c^\infty(G)$ -modules, and the complex

$$(8) \quad \mathcal{B}_q(G) \otimes_G M$$

computes $H_q(G, M)$. See [3, 6].

We shall need the following extension of a result from [4]:

Lemma 1. *Let $\mathcal{C}_c^\infty(G)_\delta = \mathcal{C}_c^\infty(G) \otimes \Delta_G$ be the G -module obtained by twisting the adjoint action of G on $\mathcal{C}_c^\infty(G)$ by the modular function. Then we have a natural isomorphism*

$$(9) \quad \mathrm{HH}_q(\mathcal{C}_c^\infty(G)) \simeq H_q(G, \mathcal{C}_c^\infty(G)_\delta).$$

Proof. Consider the complex (8), which computes the continuous cohomology of $M = \mathcal{C}_c^\infty(G)_\delta$, and let $h_G : \mathcal{B}_q(G) \otimes \mathcal{C}_c^\infty(G)_\delta \simeq \mathcal{C}_c^\infty(G) \otimes \mathcal{C}_c^\infty(G^{q+1}) = \mathcal{C}_c^\infty(G^{q+2}) \rightarrow \mathcal{C}_c^\infty(G)^{\otimes q+1}$ be the map

$$h_G(f)(g_0, g_1, \dots, g_q) = \int_G f(g^{-1}hg, g^{-1}g_0, g^{-1}g_0g_1, \dots, g^{-1}g_0g_1 \dots g_q) dg,$$

$$h = g_0g_1 \dots g_q.$$

As in equation (7), the map h_G descends to the quotient to induce an isomorphism

$$(10) \quad \tilde{h}_G : \mathcal{B}_q(G) \otimes_G \mathcal{C}_c^\infty(G)_\delta \simeq \mathcal{C}_c^\infty(G^{q+2}) \otimes_G \mathbb{C} \simeq \mathcal{C}_c^\infty(G)^{\otimes q+1}$$

of complexes, that is $\tilde{h}_G \circ (d \otimes_G 1) = b \circ \tilde{h}_G$, which establishes the isomorphism $H_q(G, \mathcal{C}_c^\infty(G)_\delta) \simeq \mathrm{HH}_q(\mathcal{C}_c^\infty(G))$, as desired. \square

To better justify the twisting of the module $\mathcal{C}_c^\infty(G)$ by the modular function in the above lemma, note that the trivial representation of G gives rise to an obvious morphism $\pi_0 : \mathcal{C}_c^\infty(G) \rightarrow \mathbb{C}$, by $\pi_0(f) = \int_G f(g)dg$, which hence defines a trace on $\mathcal{C}_c^\infty(G)$. However π_0 is not G -invariant for the usual action of G , but is invariant if we twist the adjoint action of G by the modular function, as indicated.

We proceed now to a detailed study of the G -module $\mathcal{C}_c^\infty(G)_\delta$. First we define a natural Ad_G -invariant stratification of G , called the *standard stratification* of G .

Let \mathfrak{g} be the Lie algebra of G in the sense of linear algebraic groups. Denote by $a_i(g)$ the coefficients of the polynomial $\det(t + 1 - \text{Ad}_g)$,

$$\det(t + 1 - \text{Ad}_g) = \sum_{i=0}^m a_i(g) t^i \in \mathbb{F}[t].$$

Let a_r be the first non-zero coefficient a_i , and define

$$V_k = \{a_r = a_{r+1} = \dots = a_{r+k-1} = 0\}.$$

Thus $V_0 = G$, by convention, and $G \setminus V_1 = G'$, the set of regular elements of G if G is reductive. Also, $V_{m+1} = \emptyset$ because $a_m = 1$. We observe that the functions $a_i(g)$ are G -invariant polynomial functions on G , and that they depend only on the semisimple part of g .

The description of the Hochschild homology of $\mathcal{C}_c^\infty(G)$ that we shall obtain is formulated in terms of certain commutative subgroups of G , called *standard*, that we now define.

Definition 1. A commutative subgroup $S \subset G$ is called *standard* if, and only if, there exists a semisimple element $s_0 \in G$ such that S is the center of $C(s_0)$, the centralizer of s_0 in G .

If s_0 and S are as in the definition above, then S is the set of \mathbb{F} -rational points of a subgroup \mathbb{S} of \mathbb{G} and $C(S) = C(s_0)$. The element s_0 in the above definition is not unique in general, and, for a standard subgroup $S \subset G$, we denote by $S^{\text{reg}} \subset S$ the set of semisimple elements $s \in S$ such that $C(S) = C(s)$. An element $s \in S^{\text{reg}}$ will be called *S -regular*. This set is not empty by the definition of a standard subgroup. Note that S^{reg} depends on G also, and not only on S .

Standard subgroups exist. Indeed, if $\gamma \in G$ is semisimple, then $S(\gamma) := Z(C(\gamma))$, the center of the centralizer of γ , is a standard subgroup of G . In particular, every semisimple element of G belongs to S^{reg} , for some standard subgroup S of G . For all standard subgroups S , the set S^{reg} is an open subset of S in the Zariski topology.

For any p -adic group H , we denote by H_u the set of unipotent elements of H , and call it the *unipotent variety* of H . In the particular case of $H = C(S)$, where $S \subset G$ is a standard subgroup, we also denote $C(S)_u = \mathcal{U}_S$.

In order to proceed further, recall that the Jordan decomposition of an element $g \in G$ is $g = g_s g_u$, where g_s is semisimple, g_u is unipotent, and $g_s g_u = g_u g_s$. This decomposition is unique [5]. If $g = g_s g_u$ is the Jordan decomposition of $g \in G$ and if $g_s \in S^{\text{reg}}$, then $g_u \in \mathcal{U}_S$, by definition, and hence $g \in S^{\text{reg}} \mathcal{U}_S$.

Consider now a standard subgroup $S \subset G$ and let

$$F_S = \text{Ad}_G(S^{\text{reg}}), \text{ and} \\ F_S^u = \text{Ad}_G(S^{\text{reg}} \mathcal{U}_S)$$

be the set of semisimple elements of G conjugated to an element of S^{reg} and, respectively, the set of elements $g \in G$ conjugated to an element of $S^{\text{reg}} \mathcal{U}_S$ (i.e., such that the semisimple part of g is in F_S). Also, let $N(S)$ be the normalizer of S and $W(S) = N(S)/C(S)$. Since $N(S)$ leaves S^{reg} invariant and is actually the normalizer of this set, it follows that the quotient $W(S)$ can be identified with a set of automorphisms of S_{ss} , the subgroup of semisimple elements of S . Since $N(S)$ is the set of \mathbb{F} -rational points of an algebraic group, the rigidity of tori ([5], page 117) shows that $W(S)$ is finite.

The natural map $(g, s) \rightarrow gsg^{-1}$ descends to a map

$$\phi_S : (G \times S^{\text{reg}})/N(S) \ni (g, s) \rightarrow gsg^{-1} \in F_S.$$

Similarly, we obtain a map

$$\phi_S^u : (G \times S^{\text{reg}}\mathcal{U}_S)/N(S) \ni (g, su) \rightarrow gsug^{-1} \in F_S^u.$$

In the following proposition we consider the locally compact (and Hausdorff) topology of G , and *not* the Zariski topology. Denote by G_{ss} the set of semisimple elements of G .

Proposition 1. *Let S be a standard subgroup of G . Using the above notation, we have:*

(i) *The set F_S is an analytic submanifold of G , and the maps ϕ_S and ϕ_S^u are homeomorphisms.*

(ii) *For each k , the set $V_k \setminus V_{k+1}$ is a disjoint union of sets of the form F_S^u , and each $F_S^u \subset V_k \setminus V_{k+1}$ is an open subset of V_k . Moreover, the set $G_{ss} \cap (V_k \setminus V_{k+1})$ is a disjoint union of sets of the form F_S , and each $F_S \subset V_k \setminus V_{k+1}$ is an open subset of $G_{ss} \cap V_k$.*

Proof. (i) First we check that ϕ_S and ϕ_S^u are injective. Indeed, assume that $g_1s_1g_1^{-1} = g_2s_2g_2^{-1}$, for some $s_1, s_2 \in S^{\text{reg}}$. Then, if $g = g_2^{-1}g_1$, we have

$$gC(s_1)g^{-1} = C(s_2) \Rightarrow gC(S)g^{-1} = C(S) \Rightarrow gSg^{-1} = S,$$

and hence $g \in N(S)$. Consequently, we have $(g_1, s_1) = (g_2g, g^{-1}s_2g) = g^{-1}(g_2, s_2)$, with $g \in N(S)$, as desired. The same argument shows that if F_S and $F_{S'}$ have a point in common, then the standard subgroups S and S' are conjugated in G .

The injectivity of ϕ_S^u follows from the injectivity of ϕ_S , indeed, if $g_1(s_1u_1)g_1^{-1} = g_2(s_2u_2)g_2^{-1}$, let $g = g_2^{-1}g_1$ as above, and conclude that $gs_1g^{-1} = s_2$, by the uniqueness of the Jordan decomposition. As above, this implies that $g \in N(S)$.

Since the differential $d\phi_S$ is a linear isomorphism onto its image (i.e. it is injective) and ϕ_S is injective, it follows that ϕ_S is a local homeomorphism onto its image (for the locally compact topologies), and that its image is an analytic submanifold ([21], p. 38, Theorem (2.3)). The set $G_{ss} \cap (V_k \setminus V_{k+1})$ is an algebraic variety on which G acts with orbits of the same dimension, and hence ϕ_S is proper. This proves that ϕ_S is a homeomorphism. Using an inverse for ϕ_S , we obtain that ϕ_S^u is also a homeomorphism.

To prove now (ii), consider a standard subgroup $S \subset G$, and let d be the dimension of $C(S)$. Then $a_0 = a_1 = \dots = a_{d-1} = 0$ on S , and $S \cap \{a_d \neq 0\}$ contains S^{reg} as an open component. It follows that, if $s \in G_{ss} \cap (V_k \setminus V_{k+1}) \cap S^{\text{reg}}$, then $F_S \subset V_k \setminus V_{k+1}$. This shows that $G_{ss} \cap (V_k \setminus V_{k+1})$ is a union of sets of the form F_S . This must then be a disjoint union because the sets F_S are either equal or disjoint, as proved above.

Now, if $g \in V_k \setminus V_{k+1}$ has semisimple part s , then $s \in F_S \subset G_s \cap (V_k \setminus V_{k+1})$, for some standard subgroup S , and hence $g \in F_S^u \subset V_k \setminus V_{k+1}$. The sets F_S^u are open in the induced topology because the map $V_k \setminus V_{k+1} \rightarrow G_s \cap (V_k \setminus V_{k+1})$ is continuous. \square

See also [24].

Let $R^\infty := R^\infty(G)$ be the ring of locally constant Ad_G -invariant functions on G with the pointwise product, which we regard as a subset of the set of endomorphisms

of the G -module $\mathcal{C}_c^\infty(G)_\delta = \mathcal{C}_c^\infty(G) \otimes \Delta_G$. For each $k \geq 1$, denote by $I_k \subset R^\infty$ the ideal generated by functions $f : G \rightarrow \mathbb{C}$ of the form

$$f = \phi(a_r, a_{r+1}, \dots, a_{r+k-1}),$$

where ϕ is a locally constant function $\phi : \mathbb{F}^k \rightarrow \mathbb{C}$ such that $\phi(0, 0, \dots, 0) = 0$. (Recall that each of the polynomials a_0, \dots, a_{r-1} is the 0 polynomial.) By convention, we set $I_0 = (0)$; also, it follows that $I_{m+1} = R^\infty$.

Fix now k , and let $p_n \in I_k$ be the function $\phi_n(a_r, a_{r+1}, \dots, a_{r+k-1})$, where $\phi_n : \mathbb{F}^k \rightarrow \mathbb{C}$ is 1 on the set

$$\{\xi = (\xi_0, \dots, \xi_{k-1}) \in \mathbb{F}^k, \max |\xi_i| \geq q^{-n}\}$$

and vanishes outside this set. (Here q is the number of elements of the residual field of \mathbb{F} , and the non-archimedean norm “ $|\cdot|$ ” is normalized such that its range is $\{0\} \cup \{q^n, n \in \mathbb{Z}\}$.) Then $p_n = p_n^2 = p_n p_{n+1}$ and $I_k = \cup p_n R^\infty$. For further reference, we state as a lemma a basic property of the constructions we have introduced.

Lemma 2. *If M is a R^∞ -module, then $I_k M = \cup p_n M$.*

As a consequence of this above lemma, we obtain the following result.

Corollary 1. *Consider the G -module*

$$M_k = (I_{k+1}/I_k) \otimes_{R^\infty} \mathcal{C}_c^\infty(G)_\delta \simeq I_{k+1} \mathcal{C}_c^\infty(G)_\delta / I_k \mathcal{C}_c^\infty(G)_\delta.$$

Then, for each $q \geq 0$, we have an isomorphism

$$H_q(G, \mathcal{C}_c^\infty(G)_\delta) \simeq \bigoplus_{k=0}^m H_q(G, M_k)$$

of vector spaces.

Proof. There exists a (not natural) isomorphism

$$H_q(G, \mathcal{C}_c^\infty(G)_\delta) \simeq \bigoplus_{k=0}^m I_{k+1} H_q(G, \mathcal{C}_c^\infty(G)_\delta) / I_k H_q(G, \mathcal{C}_c^\infty(G)_\delta),$$

of vector spaces.

By the above lemma, the inclusion of $I_k \mathcal{C}_c^\infty(G)_\delta \rightarrow \mathcal{C}_c^\infty(G)_\delta$ of G -modules induces natural isomorphisms

$$\begin{aligned} H_q(G, I_k \mathcal{C}_c^\infty(G)_\delta) &\simeq H_q(G, \lim_{\rightarrow} p_n \mathcal{C}_c^\infty(G)_\delta) \\ &\simeq \lim_{\rightarrow} p_n H_q(G, \mathcal{C}_c^\infty(G)_\delta) \simeq I_k H_q(G, \mathcal{C}_c^\infty(G)_\delta), \end{aligned}$$

because the functor H_q is compatible with inductive limits and with direct sums.

The naturality of these isomorphisms and the Five Lemma show that

$$H_q(G, I_{k+1} \mathcal{C}_c^\infty(G)_\delta / I_k \mathcal{C}_c^\infty(G)_\delta) \simeq I_{k+1} H_q(G, \mathcal{C}_c^\infty(G)_\delta) / I_k H_q(G, \mathcal{C}_c^\infty(G)_\delta).$$

This is enough to complete the proof. \square

We now study the homology of the subquotients $M_k = I_{k+1} \mathcal{C}_c^\infty(G)_\delta / I_k \mathcal{C}_c^\infty(G)_\delta$ by identifying them with induced modules. Let Σ_k be a set of representative of conjugacy classes of standard subgroups S such that $F_S \subset V_k \setminus V_{k+1}$ (or, equivalently, $F_S^u \subset V_k \setminus V_{k+1}$).

Lemma 3. *Using the above notation, we have $I_k \mathcal{C}_c^\infty(G) = \mathcal{C}_c^\infty(G \setminus V_k)$ and*

$$I_{k+1} \mathcal{C}_c^\infty(G)_\delta / I_k \mathcal{C}_c^\infty(G)_\delta \simeq \bigoplus_{S \in \Sigma_k} \mathcal{C}_c^\infty(F_S^u).$$

Proof. It follows from the definition of I_k that, if $f \in I_k \mathcal{C}_c^\infty(G)_\delta$, then f vanishes in a neighborhood of V_k . Conversely, if f is in $\mathcal{C}_c^\infty(G \setminus V_k)$, then we can find some polynomial a_i , with $i \leq r + k - 1$, such that $|a_i|$ is bounded from below on the support of f by, say, q^{-n} , then $p_n f = f$. The second isomorphism follows from the first isomorphism using (4) and Lemma 2. \square

If $H \subset G$ is a closed subgroup and M is a smooth (left) H -module (that is, the stabilizer of each $m \in M$ is an open subgroup of M), we denote

$$(11) \quad \text{ind}_H^G(M) = \mathcal{C}_c^\infty(G) \otimes_H M = \mathcal{C}_c^\infty(G) \otimes M / \{fh \otimes m - f \otimes hm, h \in H\}.$$

Where the right H -module structure on $\mathcal{C}_c^\infty(G)$ is $(fh)(g) = f(gh^{-1})$. Then Shapiro's lemma, see [7], states that

$$(12) \quad H_k(G, \text{ind}_H^G(M) \otimes \Delta_G) \simeq H_k(H, M).$$

(A proof of Shapiro's Lemma in our setting is contained in the proof of Theorem 4.)

The basic examples of induced modules are obtained from H -spaces. If X is an H -space (we agree that H acts on X from the left), then

$$\mathcal{C}_c^\infty((G \times X)/H) \simeq \text{ind}_H^G(\mathcal{C}_c^\infty(X) \otimes \Delta_H) \simeq \text{ind}_H^G(\mathcal{C}_c^\infty(X)_\delta)$$

as G -modules, where H acts on $G \times X$ by $h(g, x) = (gh^{-1}, hx)$. For example, Proposition 1 identifies $\mathcal{C}_c^\infty(F_S^u)$ with an induced module:

$$\mathcal{C}_c^\infty(F_S^u) \simeq \text{ind}_{N(S)}^G(\mathcal{C}_c^\infty(S^{\text{reg}} \mathcal{U}_S) \otimes \Delta_{N(S)}) = \text{ind}_{N(S)}^G(\mathcal{C}_c^\infty(S^{\text{reg}} \mathcal{U}_S)_\delta).$$

Shapiro's lemma is an easy consequence of the Serre–Hochschild spectral sequence, see [7], which states the following. Let M be a smooth G -module and $H \subset G$ be a normal subgroup. Then the action of G on $H_q(H, M)$ descends to an action of G/H , and there exists a spectral sequence with $E_{p,q}^2 = H_p(G/H, H_q(H, M))$, convergent to $H_{p+q}(G, M)$.

Let $M_k = I_{k+1} \mathcal{C}_c^\infty(G)_\delta / I_k \mathcal{C}_c^\infty(G)_\delta$, as before.

Proposition 2. *Using the above notation, we have a natural*

$$H_q(G, M_k) \simeq \bigoplus_{\langle S \rangle \in \Sigma_k} \mathcal{C}_c^\infty(S^{\text{reg}})^{W(S)} \otimes H_q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta)$$

isomorphism of R^∞ -modules

Proof. Let S be a standard subgroup of G . Recall first that $W(S) = N(S)/C(S)$ is a finite group that acts freely on S^{reg} , which gives a $N(S)$ -equivariant isomorphism

$$\mathcal{C}_c^\infty(S^{\text{reg}} \mathcal{U}_S)_\delta = \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta \otimes \mathcal{C}_c^\infty(S^{\text{reg}}).$$

Let M be a smooth $N(S)$ -module. The Hochschild–Serre spectral sequence applied to the module M and the normal subgroup $C(S) \subset N(S)$ gives natural isomorphisms

$$H_q(N(S), M) \simeq H_0(W(S), H_q(C(S), M)) \simeq H_q(C(S), M)^{W(S)}.$$

Combining these two isomorphisms, we obtain

$$\begin{aligned} \mathrm{H}_k(G, \mathcal{C}_c^\infty(F_S^u)_\delta) &\simeq \mathrm{H}_k(G, \mathrm{ind}_{N(S)}^G(\mathcal{C}_c^\infty(S^{\mathrm{reg}}\mathcal{U}_S)_\delta)_\delta) \simeq \mathrm{H}_k(N(S), \mathcal{C}_c^\infty(S^{\mathrm{reg}}\mathcal{U}_S)_\delta) \\ &\simeq (\mathrm{H}_k(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta) \otimes \mathcal{C}_c^\infty(S^{\mathrm{reg}}))^{W(S)} \simeq \mathcal{C}_c^\infty(S^{\mathrm{reg}})^{W(S)} \otimes \mathrm{H}_k(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta). \end{aligned}$$

The result then follows from Lemma 3, which implies directly that

$$M_k \simeq \oplus_{S \in \Sigma_k} \mathcal{C}_c^\infty(F_S^u)_\delta.$$

The proof is now complete. \square

Combining this proposition with Corollary 1, we obtain the main result of this section. Recall that a p -adic group $G = \mathbb{G}(\mathbb{F})$ is the set of \mathbb{F} -rational points of a linear algebraic group \mathbb{G} defined over a non-archimedean, non-discrete, locally compact field \mathbb{F} of characteristic zero. Also, recall that \mathcal{U}_S is the set of unipotent elements commuting with the standard subgroup S , and that the action of $C(S)$ on $\mathcal{C}_c^\infty(\mathcal{U}_S)$ is twisted by the modular function of $C(S)$, yielding the module $\mathcal{C}_c^\infty(\mathcal{U}_S)_\delta = \mathcal{C}_c^\infty(\mathcal{U}_S) \otimes \Delta_{C(S)}$.

Theorem 1. *Let G be a p -adic group. Let Σ be a set of representative of conjugacy classes of standard subgroups of $S \subset G$ and $W(S) = N(S)/C(S)$, then we have an isomorphism*

$$\mathrm{HH}_q(\mathcal{C}_c^\infty(G)) \simeq \bigoplus_{S \in \Sigma} \mathcal{C}_c^\infty(S^{\mathrm{reg}})^{W(S)} \otimes \mathrm{H}_q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta).$$

Remark. The isomorphism of the above theorem is not natural. A more natural description of $\mathrm{HH}_q(\mathcal{C}_c^\infty(G))$ will be obtained in one of the following sections by considering higher orbital integrals and their Shalika germs

2. HIGHER ORBITAL INTEGRALS AND THEIR SHALIKA GERMS

Proposition 2 of the previous section allows us to determine the structure of the localized cohomology groups $\mathrm{HH}_*(\mathcal{C}_c^\infty(G))_{\mathfrak{m}}$, where \mathfrak{m} is a maximal ideal of $R^\infty(G)$. This will lead to an extension of the *higher orbital integrals* introduced by Blanc and Brylinski in [4], and to a generalization of some results of Shalika [22] to higher orbital integrals. In this way, we shall also obtain a more natural description of $\mathrm{HH}_q(\mathcal{C}_c^\infty(G))$.

First recall the following result.

Proposition 3. *Let G be a reductive p -adic group over a field of characteristic 0, $S \subset G$ be a standard subgroup, and $\gamma \in S^{\mathrm{reg}}$ (that is, γ is a semisimple element such that $C(S) = C(\gamma)$). Then there exists a $N(S)$ -invariant closed and open neighborhood of γ in $C(S)$ such that*

$$G \times U \in (g, h) \rightarrow ghg^{-1} \in G$$

defines a homeomorphism of $(G \times U)/N(S)$ onto a G -invariant, closed and open subset $V \subset G$ containing γ .

Proof. The result follows from Luna's Lemma. For p -adic groups, Luna's Lemma is proved in [17], page 109, Properties "C" and "D." \square

From this proposition we obtain the following consequences for the ring $R^\infty(G)$.

Corollary 2. *Let U and V be as in Proposition 3 above.*

(i) *The ring $R^\infty(G)$ decomposes as the direct sum $\mathcal{C}^\infty(V)^G \oplus \mathcal{C}^\infty(V^c)^G$, and $\mathcal{C}^\infty(V)^G \simeq \mathcal{C}^\infty(U)^{N(S)} \subset \mathcal{C}^\infty(C(S))^{N(S)} = R^\infty(C(S))^{W(S)}$. (Here V^c is the complement of V in G .)*

(ii) *For any two semisimple elements $\gamma, \gamma' \in G$, if $\phi(\gamma) = \phi(\gamma')$ for all $\phi \in R^\infty(G)$, then γ and γ' are conjugated in G .*

(iii) *Let $\mathfrak{m} \in R^\infty(G)$ be the maximal ideal consisting of functions that vanish at a semisimple element $\gamma \in G$. Then \mathfrak{p} is generated by an increasing sequence of projections, and $M_{\mathfrak{m}} \simeq M/\mathfrak{m}M$, for any $R^\infty(G)$ -module M .*

Proof. (i) is an immediate consequence of Proposition 3. (ii) follows from [17], Proposition 2.5.

To prove (iii), observe that the maximal ideal \mathfrak{m} is generated by a sequence of projections p_n , that is, $\mathfrak{m} = \cup p_n R^\infty(G)$, with $p_n^2 = p_n$.

We know from [17], Proposition 2.5, that $R^\infty(G)$ is isomorphic to C^∞ , for some locally compact, totally disconnected topological space X . Moreover, if M is a $C^\infty(X)$ -module and \mathfrak{m} is the maximal ideal of functions vanishing at x_0 , for some fixed point $x_0 \in X$, then $\mathcal{C}^\infty(X)_{\mathfrak{m}} \simeq \mathcal{C}^\infty(X)/\mathfrak{m}\mathcal{C}^\infty(X)$, and hence

$$M_{\mathfrak{m}} = M \otimes_{\mathcal{C}^\infty(X)} \mathcal{C}^\infty(X)_{\mathfrak{m}} \simeq M \otimes_{\mathcal{C}^\infty(X)} \mathcal{C}^\infty(X)/\mathfrak{m}\mathcal{C}^\infty(X) \simeq M/\mathfrak{m}M.$$

Since X is metrizable, we can choose a basis V_n of compact open neighborhoods of x_0 in X . Then, if we let p_n to be the characteristic function of V_n^c , then p_n are projections generating \mathfrak{m} . By choosing V_n to be decreasing, we obtain a decreasing sequence p_n . \square

We now consider for each maximal ideal $\mathfrak{m} \subset R^\infty = R^\infty(G)$ the localization $\mathrm{HH}_q(\mathcal{C}_c^\infty(G))_{\mathfrak{m}}$.

Proposition 4. *Let \mathfrak{m} be a maximal ideal of $R^\infty(G)$. If \mathfrak{m} consists of the functions that vanish at the semisimple element $\gamma \in G$ and $S \subset G$ is a standard subgroup such that $\gamma \in S^{\mathrm{reg}}$, then*

$$\mathrm{HH}_q(\mathcal{C}_c^\infty(G))_{\mathfrak{m}} \simeq H_q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta).$$

For all other maximal ideals $\mathfrak{m} \subset R^\infty(G)$, we have $\mathrm{HH}_q(\mathcal{C}_c^\infty(G))_{\mathfrak{m}} = 0$.

Note that $H_q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)) \simeq H_q(C(\gamma), \mathcal{C}_c^\infty(C(\gamma)_u))$.

Proof. The vanishing of $\mathrm{HH}_*(\mathcal{C}_c^\infty(G))_{\mathfrak{m}}$ in the second half of the proposition follows because $\mathcal{C}_c^\infty(G)_{\mathfrak{m}} = 0$ in that case.

Assume now that \mathfrak{m} consists of functions vanishing at γ , a semisimple element of G . The localization functor $V \rightarrow V_{\mathfrak{m}}$ is exact by standard homological algebra. The sequence of ideals $(I_k)_{\mathfrak{m}}$ is an increasing sequence satisfying $(I_0)_{\mathfrak{m}} \simeq 0$ and $(I_{m+1})_{\mathfrak{m}} \simeq \mathbb{C}$. Choose k such that $(I_k)_{\mathfrak{m}} \simeq 0$ and $(I_{k+1})_{\mathfrak{m}} \simeq \mathbb{C}$. (This happens if, and only if, $\gamma \in V_k \setminus V_{k+1}$.) It follows that

$$H_q(G, \mathcal{C}_c^\infty(G)_\delta)_{\mathfrak{m}} \simeq H_q(G, I_{k+1}\mathcal{C}_c^\infty(G)_\delta / I_k\mathcal{C}_c^\infty(G)_\delta)_{\mathfrak{m}}.$$

Since all the isomorphisms of Proposition 2 are compatible with this the localization functor, we obtain that

$$\begin{aligned} \mathrm{HH}_q(\mathcal{C}_c^\infty(G))_{\mathfrak{m}} &\simeq H_q(G, \mathcal{C}_c^\infty(G)_\delta)_{\mathfrak{m}} \\ &\simeq \bigoplus_{\langle S \rangle \in \Sigma_k} H_q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta) \otimes (\mathcal{C}_c^\infty(S^{\mathrm{reg}})^{W(S)} / \mathfrak{m}\mathcal{C}_c^\infty(S^{\mathrm{reg}})^{W(S)}). \end{aligned}$$

The only quotient $\mathcal{C}_c^\infty(S^{\text{reg}})^{W(S)}/\mathfrak{m}\mathcal{C}_c^\infty(S^{\text{reg}})^{W(S)}$ that does not vanish is the one containing (a conjugate of) γ , and then it is isomorphic to \mathbb{C} . This completes the proof. \square

An alternative proof can be obtained by writing

$$H_q(G, \mathcal{C}_c^\infty(G)_\delta)_\mathfrak{m} \simeq H_q(G, \mathcal{C}_c^\infty(G)_{\delta\mathfrak{m}}) \simeq H_q(G, \mathcal{C}_c^\infty(G)_\delta/\mathfrak{m}\mathcal{C}_c^\infty(G)_\delta),$$

and then observing that $\mathcal{C}_c^\infty(G)_\delta/\mathfrak{m}\mathcal{C}_c^\infty(G)_\delta \simeq \mathcal{C}_c^\infty(\mathcal{U}_S)$, by Corollary 2 (iii). However the above proof is more convenient when dealing with orbital integrals. See also [16], first circulated in 1990 as a preprint of the Mathematical Institute of the Romanian Academy (INCREST) Nr. 18, March 1990, and where the localization techniques were first introduced.

We now extend the definition of higher orbital integrals introduced by Blanc and Brylinski to cover non-regular semisimple elements also. Fix a standard subgroup $S \subset G$, and let k be such that $S^{\text{reg}} \subset V_k \setminus V_{k+1}$. As in the above proof, Proposition 2 gives a natural $R^\infty(G)$ -linear, degree preserving, surjective morphism

$$H_*(G, I_{k+1}\mathcal{C}_c^\infty(G)_\delta/I_k\mathcal{C}_c^\infty(G)_\delta) \longrightarrow \mathcal{C}_c^\infty(S^{\text{reg}})^{W(S)} \otimes H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta),$$

and hence a linear map

(13)

$$I_{k+1} \text{HH}_*(\mathcal{C}_c^\infty(G)) = H_*(G, I_{k+1}\mathcal{C}_c^\infty(G)_\delta) \longrightarrow \mathcal{C}_c^\infty(S^{\text{reg}})^{W(S)} \otimes H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta).$$

Fix $c \in H^q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta)$ and $\gamma \in S^{\text{reg}}$, and let

$$\mathcal{O}_{\gamma,c} = \mathcal{O}_{\gamma,c}^S : I_{k+1} \text{HH}_q(\mathcal{C}_c^\infty(G)) \longrightarrow \mathbb{C}$$

be the evaluation of the map at γ and c in (13). We obtain, in particular, that for any $f \in I_{k+1} \text{HH}_q(\mathcal{C}_c^\infty(G))$, the function $\gamma \rightarrow \mathcal{O}_{\gamma,c}(f)$ is a locally constant, compactly supported function on S^{reg} . The function $\mathcal{O}_{\gamma,c}$ can then be extended to the whole group $\text{HH}_q(\mathcal{C}_c^\infty(G))$ using a simple observation. For any $\gamma \in S^{\text{reg}}$ there exists a locally constant function $\phi \in I_{k+1}$ such that $\phi(\gamma) = 1$. Then let

$$\mathcal{O}_{\gamma,c}(f) := \mathcal{O}_{\gamma,c}(\phi f),$$

which is independent of ϕ . It follows from definition of $\mathcal{O}_{\gamma,c}$ that, for any $f \in \text{HH}_q(\mathcal{C}_c^\infty(G))$, the function $\gamma \rightarrow \mathcal{O}_{\gamma,c}(f)$, obtained as above, is a locally constant function on S^{reg} , but not necessarily compactly supported. We thus obtain the following result.

Proposition 5. *Let $S \subset G$ be a standard subgroup. Then there exists a degree-preserving, $R^\infty(G)$ -linear map*

$$\mathcal{O}^S : \text{HH}_*(\mathcal{C}_c^\infty(G)) \longrightarrow \mathcal{C}^\infty(S^{\text{reg}})^{W(S)} \otimes H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta),$$

which is an isomorphism when localized at each maximal ideal $\mathfrak{m} \subset R^\infty(G)$ consisting of functions vanishing at an element $\gamma \in S^{\text{reg}}$.

We call the maps \mathcal{O}^S and $\mathcal{O}_{\gamma,c} = \mathcal{O}_{\gamma,c}^S$ “higher orbital integrals” because they generalize the usual notion of orbital integral. (If c is a cocycle of dimension q , we call $\mathcal{O}_{\gamma,c}$ a q -higher orbital integral.) Indeed, assume that G and $C(S)$ are unimodular, following thorough the identifications in the previous section, we obtain,

for $c_0 = 1 \in H^0(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta)$ the evaluation at the identity element $e \in G$, and $f \in \mathcal{C}_c^\infty(G) = \mathrm{HH}_0(\mathcal{C}_c^\infty(G))$, that

$$\mathcal{O}_{\gamma, c_0}(f) = \mathcal{O}_{\gamma, 1}(f) = \int_{G/C(S)} f(g\gamma g^{-1}) d\bar{g},$$

where $d\bar{g}$ is the induced measure on $G/C(S)$.

If $\gamma \in G$ is a semisimple element and S is a standard subgroup of G such that $C(\gamma) = C(S)$, (i.e., $\gamma \in S^{\mathrm{reg}}$), then restriction at γ defines a map

$$\mathcal{O}_\gamma = \mathcal{O}_\gamma^S : \mathrm{HH}_*(\mathcal{C}_c^\infty(G)) \rightarrow H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta)$$

such that $c(\mathcal{O}_\gamma(f)) = \mathcal{O}_{\gamma, c}(f)$, for all $c \in H^q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta)$.

A word on notation, whenever we write $\mathcal{O}_{\gamma, c}^S$ or \mathcal{O}_γ^S , we assume that $\gamma \in S^{\mathrm{reg}}$, which actually determines S . This means that we can omit S from notation. However, if we want to write that $\mathcal{O}_{\gamma, c} = \mathcal{O}_{\gamma, c}^S$ is obtained by specializing

$$\mathcal{O}^S : \mathrm{HH}_*(\mathcal{C}_c^\infty(G)) \rightarrow \mathcal{C}^\infty(S^{\mathrm{reg}})^{W(S)} \otimes H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta)$$

to a point $\gamma \in S^{\mathrm{reg}}$ and then by evaluating at c , that is that

$$\mathcal{O}_{\gamma, c}^S(f) = \langle \mathcal{O}^S(f)(\gamma), c \rangle,$$

then it is obviously more convenient to include S in the notation.

Let $\gamma \in G$ be a semisimple element. We want now to investigate the behavior of the orbital integrals $\mathcal{O}_{g, c}$ with g in a small neighborhood of γ . Fix a standard subgroup $S \subset G$ such that γ is in the closure of $Ad_G(S^{\mathrm{reg}})$, but is not in $Ad_G(S^{\mathrm{reg}})$, and a class $c \in H^q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta)$. More precisely, we want to study the germ of the function $g \rightarrow \mathcal{O}_{g, c}(f)$ at an element γ , where $f \in \mathrm{HH}_q(\mathcal{C}_c^\infty(G))$ is arbitrary. The germ of a function h at γ will be denoted h_γ .

The following theorem extends one of the basic properties of Shalika germs from usual orbital integrals to higher orbital integrals.

Theorem 2. *Let $S \in G$ be a standard subgroup $\gamma \in S$ an element in the closure of S^{reg} , but $\gamma \notin S^{\mathrm{reg}}$. Then there exists a degree preserving linear map*

$$\sigma_\gamma^S : H_*(C(\gamma), \mathcal{C}_c^\infty(C(\gamma)_u)_\delta) \rightarrow \mathcal{C}^\infty(S^{\mathrm{reg}})_\gamma^{W(S)} \otimes H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta),$$

such that

$$\mathcal{O}^S(f)_\gamma = \sigma_\gamma^S(\mathcal{O}_\gamma(f)),$$

for all $f \in \mathrm{HH}_*(\mathcal{C}_c^\infty(G))$.

Note that, in the notation for the maps σ_γ^S , the standard subgroup S is no longer determined by γ .

Proof. By the definition of the localization of a module, the map

$$\mathcal{O}^S : \mathrm{HH}_*(\mathcal{C}_c^\infty(G)) \rightarrow \mathcal{C}^\infty(S^{\mathrm{reg}})_\gamma^{W(S)} \otimes H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta)$$

factors through a map

$$F : \mathrm{HH}_*(\mathcal{C}_c^\infty(G))_\gamma \rightarrow \mathcal{C}^\infty(S^{\mathrm{reg}})_\gamma^{W(S)} \otimes H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta).$$

Since $\mathcal{O}_\gamma : \mathrm{HH}_*(\mathcal{C}_c^\infty(G))_\gamma \rightarrow H_*(C(\gamma), \mathcal{C}_c^\infty(C(\gamma)_u)_\delta)$ is an isomorphism by Proposition 4, we may define

$$\sigma_\gamma^S = F \circ \mathcal{O}_\gamma^{-1},$$

and all desired properties for σ_γ^S will be satisfied. \square

Let $\gamma \in S \setminus S^{\text{reg}}$ be such that γ is in the closure of S^{reg} , as above, and also let $c \in H^q(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta)$. Then a consequence of the above theorem, Theorem 2, is that the germ at γ of the higher orbital integrals $\mathcal{O}_{g,c}^S$ depends only on \mathcal{O}_γ . More precisely, if $g \in S^{\text{reg}}$, $f \in \text{HH}_q(\mathcal{C}_c^\infty(G))$, and we regard $\mathcal{O}_{g,c}^S(f)$ as a function of g , then its germ at γ , denoted $\mathcal{O}_{g,c}^S(f)_\gamma$, is given by

$$(14) \quad \mathcal{O}_{g,c}^S(f)_\gamma = \langle \sigma_\gamma^S(\mathcal{O}_\gamma(f)), c \rangle.$$

This observation allows us to relate Theorem 2 with results of Shalika [22] and Vigneras [24]. So assume now that G is reductive and let $\xi_i \in H_0(C(\gamma), \mathcal{C}_c^\infty(C(\gamma)_u)_\delta)$ be the basis dual to the basis of $H^0(C(\gamma), \mathcal{C}_c^\infty(C(\gamma)_u)_\delta)$ given by the orbital integrals over the orbits of γu , for u nilpotent in $C(\gamma)$. If we let $F_i = \sigma_\gamma(\xi_i)$, then we recover the usual definition of Shalika germs. Due to this fact, we shall call the maps σ_γ^S introduced in Theorem 2 the *higher* Shalika germs.

We can now characterize the range of the higher orbital integrals. Combining all higher orbital integrals for $S \subset G$ ranging through a set Σ of representatives of standard subgroups of G , we obtain a map

$$\mathcal{O} : \text{HH}_*(\mathcal{C}_c^\infty(G)) \rightarrow \bigoplus_{S \in \Sigma} \mathcal{C}^\infty(S^{\text{reg}})^{W(S)} \otimes H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)).$$

Theorem 3. *Let Σ be a set of representatives of standard subgroups of G and σ_γ^S be the maps introduced in Theorem 2 for $\gamma \in \overline{S^{\text{reg}}} \setminus S^{\text{reg}}$. Also, let*

$$\mathcal{F} \subset \bigoplus_{S \in \Sigma} \mathcal{C}^\infty(S^{\text{reg}}) \otimes H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S)_\delta),$$

be the space of sections ξ satisfying $\xi_\gamma = \sigma_\gamma^S(\xi(\gamma))$. Then \mathcal{O} establishes a $R^\infty(G)$ -linear isomorphism

$$\mathcal{O} : \text{HH}_*(\mathcal{C}_c^\infty(G)) \longrightarrow \mathcal{F}.$$

Proof. Note first that the map \mathcal{O} is well defined, that is, that its range is contained in \mathcal{F} , by Theorem 2.

To prove that \mathcal{O} is an isomorphism, filter both $\text{HH}_*(\mathcal{C}_c^\infty(G))$ and \mathcal{F} by the subgroups $I_k \text{HH}_*(\mathcal{C}_c^\infty(G))$ and, respectively, by $I_k \mathcal{F}$, using the ideals I_k introduced in Section 1. Since \mathcal{O} is $R^\infty(G)$ -linear, it preserves this filtration and induces maps

$$I_{k+1} \text{HH}_*(\mathcal{C}_c^\infty(G)) / I_k \text{HH}_*(\mathcal{C}_c^\infty(G)) \rightarrow I_{k+1} \mathcal{F} / I_k \mathcal{F}.$$

These maps are, by construction, exactly the isomorphisms of Proposition 2. Standard homological algebra then implies that \mathcal{O} itself is an isomorphism, as desired. \square

A consequence of the above result the following “density” corollary.

Corollary 3. *Let $a \in \text{HH}_q(\mathcal{C}_c^\infty(G))$. If all q -higher orbital integrals of a vanish, then $a = 0$.*

We shall also need certain specific cocycles below. Let τ_0 be the trace $\tau_0(f) = f(e)$ on $\mathcal{C}_c^\infty(G)$, G unimodular, obtained by evaluating f at the identity e of G . Let G_0 be the kernel of all unramified characters of G . Then $G/G_0 \simeq \mathbb{Z}^r$, where r is the rank of a split component of G . Let $p_j : G \rightarrow \mathbb{Z}$ be the morphisms obtained by considering the j th component of \mathbb{Z}^r . Then

$$\delta_j(f)(g) = p_j(g)f(g)$$

defines a derivation of $\mathcal{C}_c^\infty(G)$. Moreover, we can identify $H^*(G)$ with $\Lambda^*\mathbb{C}^r$, the exterior algebra with generators $\delta_1, \dots, \delta_r$. Fix $c \in H^*(G)$. We can assume that $c = \delta_1 \wedge \dots \wedge \delta_q$, and then we define the map $D_c : \mathcal{C}_c^\infty(G)^{\otimes q+1} \rightarrow \mathcal{C}_c^\infty(G)$ by the formula

$$(15) \quad D_c(f_0, \dots, f_q) = (q!)^{-1} \sum_{\sigma \in S_q} \epsilon(\sigma) f_0 \delta_{\sigma(1)}(f_1) \delta_{\sigma(2)}(f_2) \dots \delta_{\sigma(q)}(f_q)$$

Then $\tau_c = \tau_0 \circ D_c(f_0, \dots, f_q)$ defines a Hochschild q cocycle on $\mathcal{C}_c^\infty(G)$, and

$$(16) \quad \tau_c = \mathcal{O}_{e,c},$$

if we naturally identify c with an element of the cohomology group $H^*(G, \mathcal{C}_c^\infty(G_u))$.

3. THE COHOMOLOGY OF THE UNIPOTENT VARIETY

It follows from the main result of the first section, Theorem 1, that, in order to obtain a more precise description of the Hochschild homology of $\mathcal{C}_c^\infty(G)$, we need to understand the continuous cohomology of the H -module $\mathcal{C}_c^\infty(H_u)_\delta$, where H ranges through the set of centralizers of standard subgroups of G and H_u is the variety of unipotent elements in H . (We call the variety H_u the *unipotent variety* of H .) Since the cohomology groups $H_k(H, \mathcal{C}_c^\infty(H_u)_\delta)$ depend only on H , it is enough to consider the case $H = G$. In this section we gather some results on the groups $H_q(G, \mathcal{C}_c^\infty(G_u)_\delta)$.

We first need to recall the computation of the groups $H_*(G) = H_*(G, \mathbb{C})$, [6]. More generally, we also need to compute $H_*(G, \mathbb{C}_\chi)$, where $\chi : G \rightarrow \mathbb{C}^*$ is a character of G and $\mathbb{C}_\chi = \mathbb{C}$ as a vector space, but with G -action given by the character χ .

Assume first that $G = S$ is a commutative p -adic group, and let S_0 be the union of all compact-open subgroups of S . Then S_0 is a subgroup of S and S/S_0 is a free abelian subgroup, whose rank we denote by $\text{rk}(S)$. For this group we then have

$$H_k(S) \simeq H_k(S/S_0, \mathbb{C}) \simeq \Lambda^k \mathbb{C}^{\text{rk}(S)}.$$

For an arbitrary p -adic group G , we may identify the cohomology groups $H_q(G)$ with those of a commutative p -adic group. Indeed, if G^0 is the connected component of G (in the sense of algebraic groups) then G/G^0 is finite, and hence $H_q(G) \simeq H_q(G^0)$, by the Hochschild–Serre spectral sequence. This tells us that we may assume G to be connected as an algebraic group. Choose then a Levi decomposition $G = MN$, where N is the unipotent radical of G , M is a reductive subgroup, uniquely determined up to conjugation, and the product MN is a semidirect product. Since $H_q(N) = 0$ for $q > 0$, it follows that $H_q(G) \simeq H_q(M)$. Let $M_1 \subset M$ be the commutator subgroup of M , which is also a p -adic group, see [5]. The cohomology groups $H_q(M_1)$ were computed in [4], Proposition 6.1, page 316, and [6] and they also vanish for $q > 0$ (because the fundamental domain of the building of G is a simplex). All in all, we obtain that

$$H_q(G) \simeq H_q(M) \simeq H_q(M^{\text{ab}}),$$

where $M^{\text{ab}} = M/M_1$ is the abelianization of M .

We summarize the above discussion in the following well known statement.

Lemma 4. *Let G be a p -adic group, not necessarily reductive, and let r be the rank of a split component of the reductive quotient of G . Then*

$$H_q(G) = H_q(G, \mathbb{C}) \simeq \Lambda^q \mathbb{C}^r,$$

and $H_q(G, \mathbb{C}_\chi) = 0$, if χ is a nontrivial character of G .

We continue with a few elementary remarks on $H_k(G, \mathcal{C}_c^\infty(G_u)_\delta)$.

Remark 1. If $G_1 \rightarrow G$ is a surjective morphism with finite kernel F , then there exists a natural homeomorphism $G_{1u} \simeq G_u$ of the unipotent varieties of the two groups. Since the kernel F acts trivially on G_{1u} , using the Hochschild-Serre spectral sequence we obtain an isomorphism

$$(17) \quad H_k(G_1, \mathcal{C}_c^\infty(G_{1u})_\delta) \simeq H_k(G_1, \mathcal{C}_c^\infty(G_u)) \simeq H_k(G, \mathcal{C}_c^\infty(G_u)_\delta).$$

Remark 2. If $G \subset G_1$ is a normal p -adic subgroup with $F \simeq G_1/G$ finite, then we again have a natural homeomorphism $G_{1u} = G_u$. This gives

$$(18) \quad H_k(G_1, \mathcal{C}_c^\infty(G_{1u})_\delta) \simeq H_k(G_1, \mathcal{C}_c^\infty(G_u)_\delta) \simeq H_k(G, \mathcal{C}_c^\infty(G_u)_\delta)^F,$$

using once again the Hochschild-Serre spectral sequence. In particular, if the characteristic morphism $F \rightarrow \text{Aut}(G)/\text{Inn}(G)$ is trivial, then we get a natural isomorphism $H_k(G, \mathcal{C}_c^\infty(G_u)_\delta) \simeq H_k(G_1, \mathcal{C}_c^\infty(G_{1u})_\delta)$.

Remark 3. If $G = G' \times G''$, then $G_u = G'_u \times G''_u$ naturally, and hence $\mathcal{C}_c^\infty(G_u) \simeq \mathcal{C}_c^\infty(G'_u) \otimes \mathcal{C}_c^\infty(G''_u)$. This gives

$$(19) \quad H_k(G, \mathcal{C}_c^\infty(G_u)_\delta) \simeq \bigoplus_{i+j=k} H_i(G', \mathcal{C}_c^\infty(G'_u)) \otimes H_j(G'', \mathcal{C}_c^\infty(G''_u)).$$

Remark 4. If Z is a commutative p -adic group of split rank r , then

$$H_k(Z, \mathcal{C}_c^\infty(Z_u)_\delta) \simeq \mathcal{C}_c^\infty(Z_u) \otimes \Lambda^k \mathbb{C}^r.$$

Remark 5. The above isomorphisms reduce the computation of $H_k(G, \mathcal{C}_c^\infty(G_u)_\delta)$ for G reductive, to the computation of the cohomology groups corresponding to its semisimple quotient $H := G/Z(G)$:

$$(20) \quad H_k(G, \mathcal{C}_c^\infty(G_u)_\delta) = H_k(G, \mathcal{C}_c^\infty(G_u)) \simeq \bigoplus_{i+j=k} H_i(H, \mathcal{C}_c^\infty(H_u)_\delta) \otimes \Lambda^j \mathbb{C}^r,$$

where r is the rank of a split component of G .

Let τ_0 be the trace obtained by evaluating at the identity. Using τ_0 , we obtain an injection $H^j(G) \ni c \rightarrow \tau_0 \otimes c \in H^j(G, \mathcal{C}_c^\infty(G_u))$.

In order to obtain more precise results on $H_*(G, \mathcal{C}_c^\infty(G_u)_\delta)$, we need to take a closer look at the structure of $\mathcal{C}_c^\infty(G_u)$ as a G -module. For a G -space X , we denote by $\langle X \rangle$ the quotient space X/G with the induced topology, which may be non-Hausdorff. Thus $\langle G_u \rangle$ is the set of unipotent conjugacy classes of G .

Assume now that $\langle G_u \rangle$ is a finite set. (This happens for example if G is reductive, because the ground field \mathbb{F} has characteristic zero.) Then the space G_u can be written as an increasing union of open G -invariant sets $U_l \subset G_u$, $U_{-1} = \emptyset$, such that each difference set $U_l \setminus U_{l-1}$ is a disjoint union of open and closed G -orbits,

$$(21) \quad U_l \setminus U_{l-1} = \cup X_{l,j}.$$

A filtration U_l with these properties will be called “nice.” There may be several nice filtrations of G_u .

A nice filtration of G_u , as above, gives rise, by standard arguments, to a spectral sequence converging to $H_k(G, \mathcal{C}_c^\infty(G_u)_\delta)$, as follows. First, let $\langle g \rangle \in \langle G_u \rangle$ be the

orbit through an element $g \in G_u$. Also, let $C(g)$ denote the centralizer of $g \in G_u$ and r_g denote the rank of a split component of $C(g)$ if $C(g)$ is unimodular, $r_g = 0$ otherwise. This definition of r_g is justified by $H_k(C(g), \Delta_{C(g)}) \simeq \Lambda^k \mathbb{C}^{r_g}$.

Proposition 6. *Let G be a p -adic group with finitely many unipotent orbits (i.e., $\langle G_u \rangle$ is finite). Then, for any nice filtration (U_l) of G_u by open G -invariant subsets, there exists a natural spectral sequence with*

$$E_{p,q}^2 = \bigoplus_{\langle u \rangle \in \langle U_p \setminus U_{p-1} \rangle} \Lambda^{p+q} \mathbb{C}^{r_u},$$

convergent to $H_{p+q}(G, \mathcal{C}_c^\infty(G_u)_\delta)$.

Proof. The argument is standard and goes as follows. Recall first that any filtration $0 = F_0 \subset F_1 \subset \dots \subset F_N = \mathcal{C}_c^\infty(G_u)_\delta$ by G -submodules gives rise to a spectral sequence with $E_{p,q}^1 = H_{p+q}(G, F_p/F_{p-1})$, convergent to $H_{p+q}(G, \mathcal{C}_c^\infty(G_u)_\delta)$.

Now, associated to the open sets U_l of a nice filtration, there exists an increasing filtration $F_l = \mathcal{C}_c^\infty(U_l)_\delta \subset \mathcal{C}_c^\infty(G_u)_\delta$ by G -submodules such that

$$\mathcal{C}_c^\infty(U_l)_\delta / \mathcal{C}_c^\infty(U_{l-1})_\delta \simeq \bigoplus_j \mathcal{C}_c^\infty(X_{l,j})_\delta,$$

where each $X_{l,j}$ is the orbit of a unipotent element, and $U_l \setminus U_{l-1}$ has the topology given by the disjoint union of the orbits $X_{l,j}$. Fix l and j , and let u be a unipotent element in $X_{l,j}$ (so that then $X_{l,j}$ is the orbit through u), which implies that $\mathcal{C}_c^\infty(X_{l,j})_\delta \simeq \text{ind}_{C(u)}^G(\Delta_{C(u)})$. Finally, from Shapiro's lemma we obtain that

$$H_k(G, \mathcal{C}_c^\infty(X_{l,j})_\delta) \simeq H_k(C(u), \Delta_{C(u)}) \simeq \Lambda^k \mathbb{C}^{r_u},$$

and this completes the proof. \square

We expect this spectral sequence to converge for G reductive. This is the case, for example for $G = GL_n(\mathbb{F})$ and for $SL_n(\mathbb{F})$. See Section 5. The convergence of the spectral sequence implies, in particular, the convergence of the orbital integrals of unipotent elements in reductive groups (which is a well known fact due to Deligne and Rao [18]). In general, the convergence of the spectral sequence of the above proposition can be interpreted to represent the convergence of “higher orbital integrals.”

4. INDUCTION AND THE UNIPOTENT VARIETY

We assume from now on that G is reductive, and we fix a parabolic subgroup $P \subset G$, $P \neq G$, and a Levi subgroup $M \subset P$, so that $P = MN$, where N is the unipotent radical of P , and the product is a semidirect product. In this section, we relate $H_*(G, \mathcal{C}_c^\infty(G_u)_\delta)$ to the groups, $H_*(P, \mathcal{C}_c^\infty(P_u)_\delta)$ and $H_*(M, \mathcal{C}_c^\infty(M_u)_\delta)$. We have considered non-unimodular subgroups in the previous sections in order to be able to handle subgroups like P .

Let K be a “good” maximal compact subgroup of G (see [9], Theorem 5), so that $G = KP$. This decomposition shows that the map

$$K \times P \ni (k, p) \rightarrow kp k^{-1} \in G$$

is proper, and hence the map $G \times_P P := (G \times P)/P \ni (g, p) \rightarrow gp g^{-1} \in G$ is also proper. This gives a map

$$\mathcal{C}_c^\infty(G)_\delta = \mathcal{C}_c^\infty(G) \rightarrow \mathcal{C}_c^\infty(G \times_P P) \simeq \text{ind}_P^G(\mathcal{C}_c^\infty(P) \otimes \Delta_P) = \text{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta)$$

of G -modules. This map of G -modules and the standard identification of Hochschild homology with continuous cohomology, equation (9), then give a morphism

$$(22) \quad \text{ind}_P^G : \text{HH}_*(\mathcal{C}_c^\infty(G)) \longrightarrow \text{HH}_*(\mathcal{C}_c^\infty(P)),$$

defined as the composition of the following sequence of morphisms

$$\begin{aligned} \text{HH}_*(\mathcal{C}_c^\infty(G)) &\simeq \text{H}_*(G, \mathcal{C}_c^\infty(G)_\delta) \longrightarrow \text{H}_*(G, \text{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta) \otimes \Delta_G) \\ &\simeq \text{H}_*(P, \mathcal{C}_c^\infty(P)_\delta) \simeq \text{HH}_*(\mathcal{C}_c^\infty(P)) \end{aligned}$$

of Hochschild homology groups. The main result of this section states that ind_P^G is induced by a morphism of algebras, which we now proceed to define.

Let dk be the normalized Haar measure on the maximal compact subgroup K , normalized such that K has volume 1. The composition of kernels

$$T_1 T_2(k_1, k_2) = \int_{G/P} T_1(k_1, k) T_2(k, k_2) dk$$

defines on $\mathcal{C}^\infty(K \times K)$ an algebra structure. Let

$$(23) \quad \phi_P^G : \mathcal{C}_c^\infty(G) \longrightarrow \mathcal{C}^\infty(K \times K) \otimes \mathcal{C}_c^\infty(P),$$

be defined by $\phi_P^G(f)(k_1, k_2, p) = f(k_1 p k_2^{-1})$.

Recall [9] that the push-forward of the product $dpdk$ of Haar measure on $P \times K$, via the multiplication map $P \times K \ni (p, k) \rightarrow pk \in G$, is a left invariant measure on G , and hence a multiple λdg of the Haar measure dg on G . Suppose that the measure dk of K is the restriction of dg to K , and has total mass 1. Then the Haar measures on G and P will be called *compatible* if $\lambda = 1$. We shall need the following result of Harish Chandra (implicitly stated in [23]):

Lemma 5. *Suppose the Haar measures on G and P are compatible. Then the linear map ϕ_P^G , defined above in equation (23), is a morphism of algebras.*

Proof. The product on $\mathcal{C}^\infty(K \times K) \otimes \mathcal{C}_c^\infty(P) = \mathcal{C}_c^\infty(K \times K \times P)$ is given by the formula

$$(h_1 h_2)(k_1, k_2, p) = \int_K \int_P h_1(k_1, k, q) h_2(k, k_2, q^{-1}p) dq dk.$$

Let $*$ denote the multiplication (i.e., convolution product) on $\mathcal{C}_c^\infty(G)$. Thus, we need to prove that

$$f_1 * f_2(k_1 p k_2^{-1}) = \int_K \int_P f_1(k_1 q k^{-1}) f_2(k q^{-1} p k_2^{-1}) dq dk,$$

for all $f_1, f_2 \in \mathcal{C}_c^\infty(G)$. Consider the map $P \times K \ni (q, k) \rightarrow g := qk^{-1} \in G$, and let $d\mu$ be the push-forward of the measure $dqdk$. Then the right-hand side of the above formula becomes

$$\int_K \int_P f_1(k_1 q k^{-1}) f_2(k q^{-1} p k_2^{-1}) dq dk = \int_G f_1(k_1 g) f_2(g^{-1} p k_2^{-1}) d\mu(g).$$

We know that $d\mu = dg$, by assumptions (see the discussion before the statement of this lemma), and then

$$\int_G f_1(k_1 g) f_2(g^{-1} p k_2^{-1}) d\mu(g) = \int_G f_1(g) f_2(g^{-1} k_1 p k_2^{-1}) d\mu(g) = f_1 * f_2(k_1 p k_2^{-1}),$$

by the invariance of the Haar measure. The lemma is proved. \square

The trace $\mathcal{C}^\infty(K \times K) \rightarrow \mathbb{C}$ induces an isomorphism

$$\tau : \mathrm{HH}_*(\mathcal{C}^\infty(K \times K) \otimes \mathcal{C}_c^\infty(P)) \simeq \mathrm{HH}_*(\mathcal{C}_c^\infty(P)).$$

Explicitly, this isomorphism is given at the level of chains by

$$\begin{aligned} \tau(f_0 \otimes f_1 \otimes \dots \otimes f_q)(p_0, p_1, \dots, p_q) \\ := \int_{K^{q+1}} f_0(k_0, k_1, p_0) f_1(k_1, k_2, p_1) \dots f_q(k_q, k_0, p_q) dk_0 \dots dk_q. \end{aligned}$$

This isomorphism combines with ϕ_P^G to give a morphism

$$(24) \quad (\phi_P^G)_* : \mathrm{HH}_*(\mathcal{C}_c^\infty(G)) \longrightarrow \mathrm{HH}_*(\mathcal{C}_c^\infty(P)).$$

Theorem 4. *Let P be a parabolic subgroup of a reductive p -adic group G . Consider the morphisms $(\phi_P^G)_*$ and $\mathrm{ind}_P^G : \mathrm{HH}_*(\mathcal{C}_c^\infty(G)) \longrightarrow \mathrm{HH}_*(\mathcal{C}_c^\infty(P))$, defined above (Equations (24) and (22)). Then $(\phi_P^G)_* = \mathrm{ind}_P^G$.*

Proof. Let M_1 and M_2 be two left G -modules. We can regard M_1 as a right module, and then the tensor product $M_1 \otimes_G M_2$ is the quotient of $M_1 \otimes M_2$ by the group generated by the elements $gm_1 \otimes gm_2 - m_1 \otimes m_2$, as before. Alternatively, we can think of $M_1 \otimes_G M_2$ as $(M_1 \otimes M_2) \otimes_G \mathbb{C}$. This justifies the notation $f \otimes_G 1$ for a morphism $M_1 \otimes_G M_2 \rightarrow M'_1 \otimes_G M'_2$ induced by a morphism

$$f = f_1 \otimes f_2 : M_1 \otimes M_2 \rightarrow M'_1 \otimes M'_2.$$

We shall prove the theorem by an explicit computation. To this end, we shall use the results and notation (h_G and $\tilde{h}_G = h_G \otimes_G 1$) of Lemma 1.

By direct computation, we see that the morphism

$$\tau \circ \phi_P^G : \mathcal{C}_c^\infty(G)^{\otimes q+1} \rightarrow \mathcal{C}_c^\infty(P)^{\otimes q+1}$$

between Hochschild complexes, is given by the formula

$$(25) \quad \tau \circ \phi_P^G(f)(p_0, p_1, \dots, p_q) = \int_{K^{q+1}} f(k_0 p_0 k_1^{-1}, k_1 p_1 k_2^{-1}, \dots, k_q p_q k_0^{-1}) dk_0 dk_1 \dots dk_q.$$

We now want to realize the map $\mathrm{ind}_P^G : \mathrm{HH}_*(\mathcal{C}_c^\infty(G)) \longrightarrow \mathrm{HH}_*(\mathcal{C}_c^\infty(P))$, at the level of complexes. In the process, it will be convenient to identify the smooth G -module $\mathcal{C}_c^\infty((G \times P)/P) \simeq \mathrm{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta)$ with a subspace of the space of functions on $G \times P$, using the projection $G \times P \rightarrow (G \times P)/P$.

Consider the G -morphism

$$l : \mathcal{B}_q(G) \otimes \mathcal{C}_c^\infty(G) \rightarrow \mathcal{B}_q(G) \otimes \mathrm{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta)$$

induced by the morphism

$$\mathcal{C}_c^\infty(G) \rightarrow \mathrm{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta) \subset \mathcal{C}^\infty(G \times P).$$

Explicitly,

$$l(f)(g_0, g_1, \dots, g_q, g, p) = f(g_0, g_1, \dots, g_q, gpg^{-1}).$$

Then the resulting morphism

$$l \otimes_G 1 : \mathrm{H}_q(G, \mathcal{C}_c^\infty(G)_\delta) = \mathrm{H}_q(G, \mathcal{C}_c^\infty(G)_\delta) \rightarrow \mathrm{H}_q(G, \mathrm{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta))$$

is the morphism $\mathrm{H}_q(G, \mathcal{C}_c^\infty(G)_\delta) \rightarrow \mathrm{H}_q(G, \mathrm{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta))$ on homology corresponding to the G -morphism $\mathcal{C}_c^\infty(G) \rightarrow \mathrm{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta)$.

The G -morphism

$$(26) \quad r : \mathcal{B}_q(G) \otimes \operatorname{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta) \rightarrow \operatorname{ind}_P^G(\mathcal{B}_q(P) \otimes \mathcal{C}_c^\infty(P)_\delta) \\ = \mathcal{C}_c^\infty(G) \otimes_P (\mathcal{B}_q(P) \otimes \mathcal{C}_c^\infty(P)_\delta) \subset \mathcal{C}_c^\infty(G \times P^{q+2}),$$

given by the formula

$$r(f)(g, p_0, p_1, \dots, p_q, p) = \int_{K^{q+1}} f(gp_0k_1^{-1}, gp_1k_2^{-1}, \dots, gp_nk_0^{-1}, g, p) dk,$$

($dk = dk_0 \dots dk_q$) is well defined and induces an isomorphism in homology, because the only nonzero homology groups are in dimension 0, and they are both isomorphic to $\operatorname{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta)$. We have an isomorphism

$$\chi : \operatorname{ind}_P^G(\mathcal{B}_q(P) \otimes \mathcal{C}_c^\infty(P)_\delta) \otimes_G \mathbb{C} \rightarrow (\mathcal{B}_q(P) \otimes \mathcal{C}_c^\infty(P)_\delta) \otimes_P \mathbb{C},$$

of complexes. This shows that the homology of the second complex in (26) is isomorphic to $H_q(P, \mathcal{C}_c^\infty(P)_\delta)$, and that the map induced on homology, that is

$$\chi(r \otimes_G 1) : H_q(G, \operatorname{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta)) \rightarrow H_q(P, \mathcal{C}_c^\infty(P)_\delta),$$

is the Shapiro isomorphism.

Recall now that the isomorphism $H_q(G, \mathcal{C}_c^\infty(G)_\delta) \simeq \operatorname{HH}_q(\mathcal{C}_c^\infty(G))$ is induced by the morphism of complexes \tilde{h}_q defined in Lemma 1, Equation (9). From the definition of the morphism $\operatorname{ind}_P^G : \operatorname{HH}_*(\mathcal{C}_c^\infty(G)) \rightarrow \operatorname{HH}_*(\mathcal{C}_c^\infty(P))$ and the above discussion, we obtain the equality of the morphisms $H_q(G, \mathcal{C}_c^\infty(G)_\delta) \rightarrow H_q(P, \mathcal{C}_c^\infty(P)_\delta)$ induced by $\chi(rl \otimes_G 1)$ and $\tilde{h}_P^{-1} \circ \operatorname{ind}_P^G \circ \tilde{h}_G$. Thus, in order to complete the proof, it would be enough to check that $\tilde{h}_P \circ \chi(rl \otimes_G 1) = \tau \circ \phi_P^G \circ \tilde{h}_G$ at the level of complexes. Let

$$t : \mathcal{B}_q(G) \otimes \operatorname{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta) \rightarrow \mathcal{B}_q(G) \otimes_G \operatorname{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta)$$

be the projection. Since h_G is surjective, it is also enough to check that $\tilde{h}_P \chi(r \otimes_G 1)tl = \tau \phi_P^G h_G$.

Let

$$r'(f)(p_0, p_1, \dots, p_q, p) = \int_{K \times K^{q+1}} f(k'p_0k_1^{-1}, k'p_1k_2^{-1}, \dots, k'p_nk_0^{-1}, k', p) dk' dk,$$

where $dk = dk_0 \dots dk_q$, as before. Then r' induces a morphism

$$r' : \mathcal{B}_q(G) \otimes \operatorname{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta) \rightarrow \mathcal{B}_q(P) \otimes \mathcal{C}_c^\infty(P)_\delta$$

of complexes satisfying $h_P \circ r' = \tilde{h}_P \chi(r \otimes_G 1)t$. Directly from the definitions we obtain then that $h_P \circ r' \circ l = \tau \circ \phi_P^G \circ h_G$. This completes the proof. \square

For simplicity, we have stated and proved the above result only for G reductive, however, it extends to general G and P such that G/P is compact, by including the modular function of G , where appropriate.

In order to better understand the effect of the morphism

$$\operatorname{ind}_P^G = (\phi_P^G)_* : \operatorname{HH}_*(\mathcal{C}_c^\infty(G)) \longrightarrow \operatorname{HH}_*(\mathcal{C}_c^\infty(P)),$$

it is sometimes useful to look at its action on the geometric fibers of the group $\operatorname{HH}_*(\mathcal{C}_c^\infty(G))$. This is especially useful because the action on the geometric fibers also recovers the classical results on the characters of induced representations.

First we observe that restriction defines a morphism $\rho_P^G : R^\infty(G) \rightarrow R^\infty(P)$. In case the group G is reductive and M is a Levi component of the parabolic subgroup P , we also have $R^\infty(P) \simeq R^\infty(M)$.

Lemma 6. *Let P be a parabolic subgroup of a reductive p -adic group G , and let $\rho_P^G : R^\infty(G) \rightarrow R^\infty(P)$ be the morphism induced by restriction, used to define a $R^\infty(G)$ -module structure on $\mathrm{HH}_*(\mathcal{C}_c^\infty(P))$. Then*

$$\mathrm{ind}_P^G : \mathrm{HH}_*(\mathcal{C}_c^\infty(G)) \longrightarrow \mathrm{HH}_*(\mathcal{C}_c^\infty(P))$$

is $R^\infty(G)$ -linear, in the sense that $\mathrm{ind}_P^G(f\xi) = \rho_P^G(f)\mathrm{ind}_P^G(\xi)$, for all $f \in R^\infty(G)$ and all $\xi \in \mathrm{HH}_(\mathcal{C}_c^\infty(G))$.*

Proof. The result of the lemma follows from the fact that the map

$$\mathcal{C}_c^\infty(G) \rightarrow \mathrm{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta)$$

is $R^\infty(G)$ -linear and the isomorphism of Shapiro's Lemma,

$$\mathrm{H}_q(G, \mathrm{ind}_P^G(\mathcal{C}_c^\infty(P)_\delta)) \simeq \mathrm{H}_q(P, \mathcal{C}_c^\infty(P)_\delta),$$

is natural.

Alternatively, one can use the explicit formula of equation (25). \square

If $\mathfrak{m} = \mathfrak{m}_\gamma \subset R^\infty(G)$ is the maximal ideal of functions vanishing at a semisimple element $\gamma \in G$, then its image $(\rho_P^G)_*(\mathfrak{m}) := \rho_P^G(\mathfrak{m})R^\infty(P) \subset R^\infty(P) = R^\infty(M)$ is the ideal of functions vanishing at all $g \in M$ that are conjugated to γ in G . If γ is elliptic, then $\mathfrak{m} = R^\infty(P)$. If $\gamma \in M$, then \mathfrak{m} need not, in general, be maximal. Nevertheless, we obtain a morphism

$$(\rho_P^G)_\gamma : \mathbb{C} \simeq R^\infty(G)_\gamma = R^\infty(G)/\mathfrak{m} \rightarrow R^\infty(M)/(\rho_P^G)_*(\mathfrak{m}) \simeq \mathbb{C}^{\#\langle\gamma\rangle},$$

where $\#\langle\gamma\rangle = l$ is the set of conjugacy classes in M that consist of elements that are conjugated to γ in the bigger group G . Let $\gamma_1, \gamma_2, \dots, \gamma_l \in M$ be representatives of the conjugacy classes of element in M that are conjugated to γ in G .

We are ready now to study the morphisms

$$\begin{aligned} (\mathrm{ind}_P^G)_\gamma : \mathrm{HH}_q(\mathcal{C}_c^\infty(G))_\gamma &= \mathrm{HH}_q(\mathcal{C}_c^\infty(G))/\mathfrak{m} \mathrm{HH}_q(\mathcal{C}_c^\infty(G)) \\ &\longrightarrow \mathrm{HH}_q(\mathcal{C}_c^\infty(P))/(\rho_P^G)_*(\mathfrak{m}) \simeq \bigoplus_{j=1}^l \mathrm{HH}_q(\mathcal{C}_c^\infty(P))_{\gamma_j}. \end{aligned}$$

Let $C_P(\gamma_j)$ be the centralizer of γ_j in P and $C_G(\gamma_j) \simeq C_G(\gamma)$ be the centralizer of γ_j in G . Then $C_P(\gamma_j)_u$ identifies with a subspace of $C_G(\gamma)_u$, which gives rise to a continuous proper map $C_G(\gamma) \times_{C_P(\gamma)} C_P(\gamma)_u \rightarrow C_G(\gamma)_u$, and hence to a morphism

$$\mathcal{C}_c^\infty(C_G(\gamma)_u) \rightarrow \mathrm{ind}_{C_P(\gamma_j)}^{C_G(\gamma)}(\mathcal{C}_c^\infty(C_P(\gamma_j)_u)_\delta)$$

of $C_G(\gamma)$ -modules. Passing to cohomology, we obtain using Shapiro's Lemma a morphism

$$\iota_{\gamma_j}^\gamma : \mathrm{H}_q(C_G(\gamma), \mathcal{C}_c^\infty(C_G(\gamma)_u)) \longrightarrow \mathrm{H}_q(C_P(\gamma), \mathcal{C}_c^\infty(C_P(\gamma_j)_u)_\delta).$$

Recall that Proposition 4 gives isomorphisms

$$\mathrm{HH}_q(\mathcal{C}_c^\infty(G))_\gamma \simeq \mathrm{H}_q(C_G(\gamma), \mathcal{C}_c^\infty(C_G(\gamma)_u))$$

and

$$\mathrm{HH}_q(\mathcal{C}_c^\infty(P))_{\gamma_j} \simeq \mathrm{H}_q(C_P(\gamma_j), \mathcal{C}_c^\infty(C_P(\gamma_j)_u)_\delta).$$

Proposition 7. *Let $\gamma \in G$ be a semisimple element and $M \subset P$ be as above. If the conjugacy class of γ does not intersect M , then $\mathrm{HH}_*(\mathcal{C}_c^\infty(P))_\gamma = 0$, and hence $(\mathrm{ind}_P^G)_\gamma = 0$. Otherwise, using the notation above, we have*

$$(\mathrm{ind}_P^G)_\gamma = \oplus_{j=1}^l \iota_{\gamma_j}^\gamma : \mathrm{HH}_*(\mathcal{C}_c^\infty(G))_\gamma \rightarrow \mathrm{HH}_*(\mathcal{C}_c^\infty(P))_\gamma.$$

Proof. This follows from definitions if we observe that in the sequence of maps

$$\begin{aligned} G \times_P P \times_{C_P(\gamma_i)} (\gamma_i C_P(\gamma_i)_u) &\simeq G \times_{C_G(\gamma_i)} C_G(\gamma_i) \times_{C_P(\gamma_i)} (\gamma_i C_P(\gamma_i)_u) \\ &\rightarrow G \times_{C_G(\gamma_i)} (\gamma_i C_G(\gamma_i)_u) \end{aligned}$$

the second map is induced by $C_G(\gamma_i) \times_{C_P(\gamma_i)} C_P(\gamma_i)_u \rightarrow C_G(\gamma_i)_u$ and their composition induces on homology the direct summand $\iota_{\gamma_j}^\gamma$ of the map $(\mathrm{ind}_P^G)_\gamma$. \square

Another morphism that is likely to play an important role is the “inflation,” which we now define. Let $N \subset P$ be the unipotent radical of an algebraic p -adic group, and let $M = P/N$ be its reductive quotient. Then integration over N defines an algebra morphism

$$\psi_M^P : \mathcal{C}_c^\infty(P) \rightarrow \mathcal{C}_c^\infty(M), \quad \psi_M^P(f)(m) = \int_N f(mn) dn.$$

Integration over N also defines a G -morphism $\mathcal{C}_c^\infty(P)_\delta \rightarrow \mathcal{C}_c^\infty(M)$, and, since N is a union of compact subgroups, we finally obtain morphisms

$$\mathrm{H}_k(P, \mathcal{C}_c^\infty(P)_\delta) \rightarrow \mathrm{H}_k(P, \mathcal{C}_c^\infty(M)) \simeq \mathrm{H}_k(M, \mathcal{C}_c^\infty(M)),$$

whose composition we denote inf_M^P .

Theorem 5. *If M is a Levi component of a p -adic group P over a field of characteristic zero, as above. Then we have*

$$(\psi_M^P)_* = \mathrm{inf}_M^P : \mathrm{HH}_*(\mathcal{C}_c^\infty(P)) \rightarrow \mathrm{HH}_*(\mathcal{C}_c^\infty(M)).$$

Proof. Integration over N defines a morphism

$$f : \mathcal{B}(P) \otimes \mathcal{C}_c^\infty(P)_\delta \rightarrow \mathcal{B}(M) \otimes \mathcal{C}_c^\infty(M),$$

which commutes with the action of P . Then $f \otimes_P 1$ coincides with the morphism of complexes induced by ψ_M^P .

Consider now the maps h_G defined in the proof of Lemma 1. Then $\psi_M^P \circ h_P = h_M \circ f$, and hence $\psi_M^P \circ \tilde{h}_P = \tilde{h}_M \circ (f \otimes_P 1)$, from which the result follows. \square

We now want to proceed by analogy and establish the explicit form of the action of inf_M^P on the geometric fibers. Fix $\gamma \in M$. Again, integration over the nilpotent radical of $C_P(\gamma)$, the centralizer of γ in P , induces a morphism

$$\mathcal{C}_c^\infty(C_P(\gamma)_u)_\delta = \mathcal{C}_c^\infty(C_P(\gamma)_u) \otimes \Delta_{C_P(\gamma)} \rightarrow \mathcal{C}_c^\infty(C_M(\gamma)_u),$$

of P -modules. Let

$$\begin{aligned} j_\gamma : \mathrm{HH}_*(\mathcal{C}_c^\infty(P))_\gamma &= \mathrm{H}_*(C_P(\gamma), \mathcal{C}_c^\infty(C_P(\gamma)_u)_\delta) \\ &\longrightarrow \mathrm{H}_*(C_M(\gamma), \mathcal{C}_c^\infty(C_M(\gamma)_u)) = \mathrm{HH}_*(\mathcal{C}_c^\infty(M))_\gamma \end{aligned}$$

be the induced morphism.

Proposition 8. *Let γ be a semisimple element of a Levi component M of the group P . Let $d(\gamma)$ be the determinant of $\text{Ad}_\gamma^{-1} - 1$ acting on $\text{Lie}(N)/\ker(\text{Ad}_\gamma - 1)$. Then, using localization at the maximal ideal defined by γ in $R^\infty(G) = R^\infty(P)$ and the above notation,*

$$(\inf_M^P)_\gamma = |d(\gamma)|^{-1} j_\gamma : \text{HH}_*(\mathcal{C}_c^\infty(P))_\gamma \rightarrow \text{HH}_*(\mathcal{C}_c^\infty(M))_\gamma.$$

Proof. Fix $\gamma \in G$, not necessarily semisimple and let N_γ be the subgroup of elements of N commuting with γ . We choose a complement V_γ of $\text{Lie}(N_\gamma)$ in $\text{Lie}(N)$ and we use the exponential map to identify V_γ with a subset of N . Then the Jacobian of the map

$$V_\gamma \times N_\gamma \ni (n, n') \rightarrow \gamma^{-1} n \gamma n^{-1} n' \in N = V_\gamma N_\gamma$$

is $d(\gamma)$, and from this the result follows. \square

This result is compatible with the results of van Dijk on characters of induced representations, see [8].

5. EXAMPLES

Our results can be used to obtain some very explicit results in certain particular cases.

Example 1. Let Z be a commutative p -adic group of split rank r (so that $H_q(Z) \simeq \Lambda^q \mathbb{C}^r$, for all $q \geq 0$). Then

$$(27) \quad \text{HH}_q(\mathcal{C}_c^\infty(Z)) \simeq \mathcal{C}_c^\infty(Z) \otimes \Lambda^q \mathbb{C}^r.$$

Example 2. Let P be the (parabolic) subgroup of upper triangular matrices in $SL_2(\mathbb{F})$, and $A \subset P$ be the subgroup of diagonal matrices. Then inflation defines a morphism

$$\inf_A^P : \text{HH}_*(\mathcal{C}_c^\infty(P)) \rightarrow \text{HH}_*(\mathcal{C}_c^\infty(A)) = \mathcal{C}_c^\infty(A) \otimes \Lambda^* \mathbb{C}$$

whose range is $\mathcal{C}_c^\infty(A) \oplus \mathcal{C}_c^\infty(A \setminus \{\pm I\})$, with I the identity matrix of $SL_2(\mathbb{F})$. (We see this by localizing at each γ .) To describe the kernel of \inf_A^P , let

$$(28) \quad u_b = \begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix}.$$

Then, if we choose b to range through Σ_u , a set of representative of $\mathbb{F}^*/\mathbb{F}^{*2}$, the set of elements u_b forms a set of representatives of the set of nontrivial conjugacy classes of unipotent elements of P . Recall that \mathbb{F} has characteristic zero, so Σ_u is a discrete set. Let $\mathcal{O}_{b,+}$ be the orbital integral associated to u_b , and let $\mathcal{O}_{b,-}$ be the orbital integral associated to $-u_b$, then

$$F_\pm = \oplus_b \mathcal{O}_{b,\pm} : \mathcal{C}_c^\infty(G) \rightarrow \mathbb{C}^{\Sigma_u}$$

identify the kernel of \inf_A^P as follows. The map $F_+ \oplus F_- : \ker(\inf_A^P) \rightarrow \mathbb{C}^{\pm \Sigma_u}$ is injective, and the range of each of F_\pm is the set of elements with zero sum.

All in all, let us consider the map $\Phi = \inf_A^P \oplus F_+ \oplus F_-$,

$$\Phi : \text{HH}_*(\mathcal{C}_c^\infty(P)) \rightarrow (\mathcal{C}_c^\infty(A) \oplus \mathbb{C}^{\pm \Sigma_u})_{(0)} \oplus (\mathcal{C}_c^\infty(A \setminus \{\pm I\}))_{(1)},$$

where the lower index (i) represents the degree. Then Φ is surjective in degree 1, and, in degree 0, it consists of $(f, \lambda_{b,\epsilon})$, $f \in \mathcal{C}_c^\infty(A)$, $\lambda \in \mathbb{C}$, for $\epsilon \in \{\pm 1\}$ and $b \in \Sigma_u \simeq \mathbb{F}^*/\mathbb{F}^{*2}$, such that $\sum_b \lambda_{b,\epsilon} = f(\epsilon I)$, for $\epsilon = \pm 1$.

Note that evaluation at $\pm I$ does not define a trace on $\mathcal{C}_c^\infty(P)$. Actually, in the spectral sequence of Proposition 6, the obstruction to extend the evaluation at I to a trace is responsible for “killing” the 1 cohomology supported at I (which explains our claim on the range of \inf_A^P above).

Example 3. Consider now the group $G = SL_2(\mathbb{F})$, where \mathbb{F} is a p -adic field of characteristic zero such that the characteristic of the residual field is not 2, for simplicity. Let F_q be the residual field of \mathbb{F} (thus $q = p^n$, for some $n \in \mathbb{N}$ and some prime p , denotes the number of elements of F_q). We choose ϵ in the valuation ring of \mathbb{F} , whose image in F_q is not a square. Also, let τ be a generator of the (unique) maximal ideal of the valuation ring of \mathbb{F} . We shall use the notation of [20] and thus let θ range through the set $\{\epsilon, \tau, \epsilon\tau\}$ and let $T_\theta, T_\theta^\#$ be the elliptic tori defined there. (Recall that $T_\theta = \{[a_{ij}], a_{11} = a_{22}, a_{21} = \theta a_{12}\}$ and $T_\theta^\# = \{[a_{ij}], a_{11} = a_{22}, a_{21} = \theta^\# a_{12}\}$, where $\theta^\# = \theta a^2$, for some $a \in \mathbb{F}^\times$ not in the image of the norm map $N : \mathbb{F}[\theta]^\times \rightarrow \mathbb{F}^\times$.) We distinguish two cases, first the case where -1 is a square and then the case where it is not a square. If -1 is a square, then the Weyl group of each of the tori $T = T_\theta, T_\theta^\#$ has order 2. Otherwise $W(T) = \{1\}$, for each $T = T_\theta$ or $T = T_\theta^\#$, but T_θ and $T_\theta^\#$ are conjugate for each fixed θ .

Let $X = \cup_\theta T_\theta / S_2 \cup \cup_\theta T_\theta^\# / S_2$, if -1 is a square, and $X = \cup_\theta T_\theta$ otherwise, with the induced topology. Then $X \setminus \{\pm 1\}$ identifies with the set of elliptic conjugacy classes of $SL_2(\mathbb{F})$.

Denote by $A \subset SL_2(\mathbb{F})$ the set of diagonal matrices in $SL_2(\mathbb{F})$. Let $W(A) = S_2$ act on $\mathcal{C}_c^\infty(A) \otimes \Lambda^* \mathbb{C}$ by conjugation on $\mathcal{C}_c^\infty(A)$ and act by the nontrivial character on \mathbb{C} . Then we have the following. Recall that there are 10 conjugacy classes of unipotent elements in $SL_2(\mathbb{F})$, if $p \neq 2$.

Proposition 9. *The composition*

$$\phi := \inf_A^P \circ \text{ind}_P^G : \text{HH}_*(\mathcal{C}_c^\infty(SL_2(\mathbb{F}))) \rightarrow \text{HH}_*(A) = \mathcal{C}_c^\infty(A) \otimes \Lambda^* \mathbb{C}$$

has range consisting of $W(A)$ invariant elements, and the kernel of ϕ is isomorphic to $\mathcal{C}_c^\infty(X \setminus \{\pm 1\}) \oplus \mathbb{C}^{10}$, via orbital integrals with respect to elliptic and unipotent elements.

Proof. First of all, it is clear that the composition $\phi = \inf_A^P \circ \text{ind}_P^G$ is invariant with respect to the Weyl group $W(A)$, and hence its range consists of $W(A)$ -invariant elements.

The localization of ϕ at a regular, diagonal conjugacy class γ is onto by Proposition 4. Next, we know that every orbital integral extends to $\mathcal{C}_c^\infty(SL_2(\mathbb{F}))$, and this implies directly that the spectral sequence of Proposition 6 collapses at the E^2 term. This proves that the localization of ϕ at $\gamma = 1$ is also onto, and hence ϕ is onto. The rest of the proposition follows also from Proposition 6 by localization. \square

This example is also discussed in [4], but from a different perspective.

Example 4. We end this section with a description of the ingredients entering in the formula (1) for the of the Hochschild homology of $\mathcal{C}_c^\infty(G)$, if $G = GL_n(\mathbb{F})$. Let $\gamma \in G$ be a semisimple element. The minimal polynomial Q_γ of γ decomposes as $Q_\gamma = p_1 p_2 \dots p_r$ into irreducible polynomials with coefficients in \mathbb{F} . (We assume, for simplicity, that each polynomial p_j is a monic polynomial.) Also, let $P_\gamma = p_1^{l_1} p_2^{l_2} \dots p_r^{l_r}$ be the characteristic polynomial of γ . Then the algebra generated by

γ is

$$\mathbb{F}[\gamma] \simeq \mathbb{K}_1 \oplus \dots \oplus \mathbb{K}_r,$$

where $\mathbb{K}_i = \mathbb{F}[t]/(p_i(t))$ are not necessarily distinct fields. The commutant $\{\gamma\}'$ of γ in $M_n(\mathbb{F})$ is the commutant of this algebra, and hence

$$\{\gamma\}' \simeq M_{l_1}(\mathbb{K}_1) \oplus M_{l_2}(\mathbb{K}_2) \oplus \dots \oplus M_{l_r}(\mathbb{K}_r),$$

$$C(\gamma) \simeq \prod_{i=1}^r GL_{l_i}(\mathbb{K}_i), \quad S := Z(C(\gamma)) \simeq \prod_{i=1}^r \mathbb{K}_i^*,$$

and

$$S^{\text{reg}} = \{(x_i) \in S, x_i \text{ generates } \mathbb{K}_i \text{ and the minimal polynomials of } x_i \text{ are distinct}\}.$$

By the Skolem–Noether theorem, the Weyl group $W(S) = N(S)/C(S)$ coincides with the group of algebra automorphisms of $\{\gamma\}'$. This group has as quotient a group isomorphic to the subgroup $\Pi \subset N(S)$ which permutes the algebras $M_{l_i}(\mathbb{K}_i)$. Then $\Pi \cong S_{m_1} \times \dots \times S_{m_t}$, that is, Π is a product of symmetric groups. We denote the kernel of this morphism by $W_0(S)$. It is isomorphic to $\prod_{i=1}^r \text{Aut}_{\mathbb{F}}(\mathbb{K}_i)$ (again by the Skolem–Noether theorem). The group $W(S)$ is then the semidirect product of $W_0(S)$ by Π . We hence obtain exact sequences

$$1 \longrightarrow N_0(S) \longrightarrow N(S) \longrightarrow \Pi \longrightarrow 1$$

and

$$1 \longrightarrow C(S) \longrightarrow N_0(S) \longrightarrow W_0(S) \longrightarrow 1.$$

According to (1), the only other ingredients necessary to compute $\text{HH}_*(\mathcal{C}_c^\infty(G))$ are the groups $H_*(C(S), \mathcal{C}_c^\infty(\mathcal{U}_S))$.

Now, the unipotent variety of $C(S)$ is the product of the unipotent varieties of $GL_{l_i}(\mathbb{K}_i)$, $i = 1, r$, and the subgroup $C(S)$ preserves this product decomposition. We see then that in order to prove that the spectral sequence of Proposition 6 collapses (for any choice of open subsets U_i), it is enough to check this for the spectral sequence converging to the cohomology of $\mathcal{C}_c^\infty(GL_n(\mathbb{K})_u)$, for an arbitrary characteristic zero p -adic field \mathbb{K} .

Fix a unipotent element $\gamma \in GL_n(\mathbb{K})$. Define then $V_0 = 0$, $V_l = \ker(\gamma - 1)^l \subset \mathbb{K}^n$, if $l > 0$. Also, choose W_l such that $V_l = V_{l-1} \oplus W_l$, and define

$$P = \{\gamma \in GL_n(\mathbb{K}), \gamma V_l \subset V_l\}, \text{ and}$$

$$M = \{\gamma \in GL_n(\mathbb{K}), \gamma W_l = W_l\}.$$

Then P is a parabolic subgroup with unipotent radical

$$N = \{\gamma \in GL_n(\mathbb{K}), (\gamma - 1)V_l \subset V_{l-1}\},$$

and M is a Levi component of P . It is easy to check, from definition, that the P -orbit of u in N is dense. The centralizer of u is then contained in P and has split rank \leq the split rank of P . Fix a maximal split torus A in the centralizer of u . We can assume that this split torus is contained in M . From the definition and by direct inspection, the map $H_*(A) \rightarrow H_*(M)$ is injective, and hence the map

$$(29) \quad H^*(M) \longrightarrow H^*(A) = H^*(C(u))$$

is surjective.

Fix now a cohomology class $c_0 \in H^q(C(\gamma)) \simeq H^q(A)$ and choose a cohomology class $c \in H^q(M)$ that maps to c_0 under the above restriction map. Also, let τ be

the trace on $\tau_0(f) = f(e)$ on $\mathcal{C}_c^\infty(M)$ (obtained by evaluation at the identity e). Then the formula

$$(30) \quad \phi_0(f_0, \dots, f_q) = \tau_0(D_c(f_0, \dots, f_q))$$

defines a Hochschild cyclic cocycle on $\mathcal{C}_c^\infty(M)$. Consequently,

$$\phi = \phi_0 \circ \inf_A^P \circ \circ \circ \text{ind}_P^G$$

defines a Hochschild cocycle on $\mathcal{C}_c^\infty(G)$. For any filtration U_i of G_u by open, invariant open sets, such that each $U_l \setminus U_{l-1}$ consists of a single orbit. Suppose that the orbit $U_l \setminus U_{l-1}$ is the orbit of $\gamma \in GL_n(\mathbb{F})$ considered above. Then the cocycle ϕ will vanish on $\mathcal{C}_c^\infty(U_l)$ and represent the cohomology class

$$c \in H^q(C(\gamma)) \simeq H^q(G, \mathcal{C}_c^\infty(U_l \setminus U_{l-1})).$$

From this it follows that the spectral sequence of Proposition 6 degenerates at E^2 .

It is very likely that the above argument extends to arbitrary reductive G by choosing M and P as in [18].

REFERENCES

- [1] P. Baum, N. Higson, and R. Plymen, *Equivariant homology for SL_s of a p -adic field*, in Index theory and operator algebras (Boulder, CO, 1991), 1–18, Contemp. Math. **148**, Amer. Math. Soc., Providence, R.I., 1993.
- [2] J. N. Bernstein, *P -invariant distributions on $GL(N)$ and the classification of unitary representations of $GL(N)$* , in Lie Group Representations II, (Herb and all. ed.), Lecture Notes in Mathematics, Springer Verlag, **1041**, (1984), 51–102.
- [3] P. Blanc, *(Co)homologie différentiable et changement de groupes*, Astérisque **124-125** (1985), 13–30.
- [4] P. Blanc and J. L. Brylinski, *Cyclic Homology and the Selberg Principle*, J. Funct. Anal. **109** (1992), 289–330.
- [5] A. Borel, *Linear Algebraic Groups*, second edition, Springer Verlag, Graduate Texts in Mathematics, vol 126.
- [6] A. Borel and N. Wallach, *Continuous Cohomology, Discrete Subgroups, and Representations of Reductive Groups*, Ann. of Math. Studies, Princeton University Press, no. 94, 1980.
- [7] W. Casselman and D. Wigner, *Continuous cohomology and a conjecture of Serre's*, Invent. Math. **25** (1974), 199–211.
- [8] G. van Dijk, *Computation of Certain Induced Characters of p -Adic Groups*, Math. Ann., **199** (1972), 229–240.
- [9] Harish-Chandra, *Harmonic Analysis on Reductive p -adic Groups*, Notes by G. van Dijk, Springer Lecture Notes in Mathematics, **162**, 1970.
- [10] N. Higson and V. Nistor, *Cyclic cohomology of groups acting on buildings*, J. Funct. Anal. **141** (1996), 466–495.
- [11] D. Kazhdan, *Cuspidal geometry of p -adic groups*, J. Analyse Math. **47** (1986), 1–36.
- [12] D. Kazhdan, V. Nistor, and P. Schneider, *Hochschild and cyclic homology of finite type algebras*, to appear in Selecta.
- [13] V. Lafforgue, *Une démonstration de la Conjecture de Baum-Connes pour les groupes réductifs sur un corps p -adique et pour certains groupes discrets possédant la propriété (T)*, C. R. Acad. Sci. Paris Sér. I Math **327** (1998), 439–444.
- [14] R. Langlands, *Base change for GL_2* , Institute for Advanced Study, 1975.
- [15] S. Mac Lane, *Homology*, Springer-Verlag, Berlin-Heidelberg-New York, 1995.
- [16] V. Nistor, *Cyclic cohomology of crossed products by algebraic groups*, Invent. Math. **112** (1993), 615–638.
- [17] C. Rader and S. Rallis, *Spherical characters on p -adic symmetric spaces*, Am. J. Math. **118** (1996), 91–178.
- [18] R. Ranga Rao, *Orbital integrals in reductive groups*, Annals of Math., **96** (1972), 505–510.
- [19] P. Schneider, *The cyclic homology of p -adic reductive groups*, J. Reine Angew. Math. **475** (1996), 39–54.

- [20] P. J. Sally, Jr and J. A. Shalika, *The Fourier transform of orbital integrals on SL_2 over a p -adic field*, in Lie Group Representations II, (Herb and all. ed.), Lecture Notes in Mathematics, Springer Verlag, **1041** (1984), 303–340.
- [21] W. H. Schikhof, *Differentiation in non-archimedean valued fields*, Indagationes Math. **32** (1970), 35–44.
- [22] J. A. Shalika, *A theorem on semisimple p -adic groups*, Annals of Math. **95** (1972), 226–242.
- [23] A. J. Silberger, *Introduction to Harmonic Analysis on reductive p -adic groups*, Mathematical Notes, Princeton University Press, 1979.
- [24] M.-F. Vigneras, *Caractérisation des intégrales orbitales sur un group reductif p -adique*, J. Fac. Sci. Univ. Tokyo Sect IA Math. **28** (1981), 945–961.

PENNSYLVANIA STATE UNIVERSITY, MATH. DEPT., UNIVERSITY PARK, PA 16802, USA
E-mail address: `nistor@math.psu.edu`