

QUIVERS OF TYPE A , FLAG VARIETIES AND REPRESENTATION THEORY

OLIVIER SCHIFFMANN

Introduction. In this survey, we describe and relate various occurrences of quivers of type A (both finite and affine) and their canonical bases in combinatorics, in algebraic geometry and in representation theory. The ubiquity of these quivers makes them especially important to study : they are pervasive in very classical topics (such as the theory of symmetric functions) as well as in some of the most recent and exciting areas of representation theory (such as representation theory of quantum affine algebras, affine Hecke algebras or Cherednik algebras).

. There is a vast litterature on the subject and we have been forced to make a choice in the selection of the results presented here. We believe that one of the main reasons for the omnipresence of quivers of type A is the fact that they are related (in more than one way) to classical and fundamental objects in geometric representation theory of type A such as (partial) flag varieties and nilpotent orbits. This is the point we tried to emphasize in this survey. For that reason, many interesting and important results are only alluded to or sketched here and we apologize to all those whose work we did not mention through a lack of space and competence. Also, we do not give complete proofs of all results presented here and refer to the original papers whenever possible. Rather, we have tried to convey the fundamental ideas in a nontechnical way as much as possible.

. This survey is organized as follows. In Section 1 we describe some basic geometric properties of quivers of type A and their representation spaces. Section 2 is a brief reminder on Ringel-Hall algebras and their canonical bases, and in Sections 3 and 4 we describe the Ringel-Hall algebras of the categories of representations of quivers of type $A_1^{(1)}$ and type $A_n, A_n^{(1)}$ respectively. This makes a link with combinatorics (symmetric functions, [Mac]) and with quantum group theory ([Ri1], [S1]). In Section 5 we recall some classical geometric constructions in type A such as (partial, affine) flag varieties, nilpotent orbits, Hecke algebras and various convolution algebras. In Section 6, we give a first link between quivers of type A and (partial, affine) Schubert varieties; one important consequence is that the coefficients of the canonical basis of the Hall algebras of quivers of type A are given by Kazhdan-Lusztig polynomials ([Lu1], [Z1]). Section 7 gives the geometric realization of Schur-Weyl duality and its affine and quantum cousins. This gives a conceptual explanation for some of the results in Section 4. We totally change perspective in Section 8 and, following [LLT], [A] and [VV1], we present a surprising and beautiful application of the theory of canonical bases and quivers of type A to character formulae for quantum affine algebras and affine Hecke algebras (at roots of unity). Finally, we gather a few more interesting occurrences of quivers of type A in relation to geometry (Nakajima quiver varieties, parabolic structures on curves and surfaces) in the last Section.

1. QUIVERS OF TYPE A

1.1. We will denote by Δ_n the (equioriented) quiver of type A_n , with vertex set $I_n = \{1, \dots, n\}$ and with edge set $\{(i, i+1) \mid i = 1, \dots, n-1\}$. Similarly, we let $\Delta_n^{(1)}$ stand for the (equioriented) affine quiver of type $A_{n-1}^{(1)}$, with vertex set $\mathbb{Z}/n\mathbb{Z}$ and edge set $\{(i, i+1) \mid i \in \mathbb{Z}/n\mathbb{Z}\}$ (note that this also makes sense for $n = 1$). Let k be any field. Recall that a *representation* of Δ_n (resp. of $\Delta_n^{(1)}$) over k consists of a pair $(\mathbf{V}, \underline{\phi})$ where $\mathbf{V} = \bigoplus_i V_i$ is an I_n -graded (resp. $\mathbb{Z}/n\mathbb{Z}$ -graded) k -vector space and $\underline{\phi} = (\phi_i) \in \bigoplus_i \text{Hom}(V_i, V_{i+1})$. We let $\text{Mod}_k(\Delta_n)$ be the abelian category of representations of Δ_n over k . In the case of $\Delta_n^{(1)}$ we will impose an additional *nilpotency* condition; namely, we denote by $\text{Mod}_k(\Delta_n^{(1)})$ the abelian category of representations $(\mathbf{V}, \underline{\phi})$ of $\Delta_n^{(1)}$ such that $\phi_N \cdots \phi_2 \phi_1 = 0$ for $N \gg 0$. This restriction may seem artificial from the strict point of view of quivers, but it will appear naturally later in the context of flag varieties.

1.2. Let Δ be either Δ_n or $\Delta_n^{(1)}$. The simple objects in $\text{Mod}_k(\Delta)$ are naturally in bijection with the vertices of Δ , and we will denote by S_i the simple module with $\mathbf{V} = V_i = k$. Indecomposable objects of $\text{Mod}_k(\Delta_n)$ are parametrized by pairs $(i; l)$ with $i \in I_n$ and $1 \leq l \leq n+1-i$: to $(i; l)$ corresponds the unique indecomposable object $S_{i,l}$ with top S_i and length l . Similarly, indecomposable objects of $\text{Mod}_k(\Delta_n^{(1)})$ are parametrized by pairs $(i; l)$ with $i \in \mathbb{Z}/n\mathbb{Z}$ and $l \geq 1$, and we use the same notation $S_{i,l}$ for the unique indecomposable with top S_i and length l . In particular, in both cases, the set of isomorphism classes of objects in $\text{Mod}_k(\Delta)$ is in bijection with a set of *n-multipartitions* (or *n-multisegments*)

$$\mathbf{m} = (\lambda^{(i)})_i = \sum_i \sum_t (i; \lambda_t^{(i)})$$

where $\lambda_t^{(i)} \leq n+1-i$ if $\Delta = \Delta_n$.

1.3. Let Δ again stand for either Δ_n or $\Delta_n^{(1)}$. Let us fix a dimension vector $\mathbf{d} = (d_i)_i \in \mathbb{N}^I$ with $I = I_n$ or $I = \mathbb{Z}/n\mathbb{Z}$ according to Δ . The group $G_{\mathbf{d}} = \prod_i GL_k(d_i)$ acts naturally by conjugation on the space

$$E(\mathbf{d}) = \bigoplus_{i=1}^{n-1} \text{Hom}(k^{d_i}, k^{d_{i+1}})$$

if $\Delta = \Delta_n$ and on the space

$$E(\mathbf{d}) = E_{\text{nilp}}(\mathbf{d}) = \{(\phi_i) \in \bigoplus_{i \in \mathbb{Z}/n\mathbb{Z}} \text{Hom}(k^{d_i}, k^{d_{i+1}}) \mid \phi_N \cdots \phi_2 \phi_1 = 0 \text{ for } N \gg 0\}$$

if $\Delta = \Delta_n^{(1)}$. Moreover, the set of $G_{\mathbf{d}}$ -orbits is (by definition) in bijection with the set of isomorphism classes of objects in $\text{Mod}_k(\Delta)$ of dimension \mathbf{d} . We will denote by $\mathcal{O}_{\mathbf{m}}$ the $G_{\mathbf{d}}$ -orbit (in the appropriate space) corresponding to a multipartition \mathbf{m} .

Since there are only finitely many multipartitions of a given dimension \mathbf{d} the group $G_{\mathbf{d}}$ acts on $E(\mathbf{d})$ with only finitely many orbits. It follows that there is a (unique) dense orbit (the “generic representation of dimension \mathbf{d} ”); it is also easy to see that there is a unique closed orbit (the semisimple representation of dimension \mathbf{d} , in which all maps ϕ_i are zero). Observe that when $\Delta = \Delta_1^{(1)}$ we recover the classical situation of the group $GL(d)$ acting on the nilpotent cone $\mathcal{N} = \{x \in gl_d \mid x^N = 0 \text{ for } N \gg 0\}$.

1.4. We use the same notation as in **1.3**. Let \mathbf{m} be a multisegment of dimension \mathbf{d} . The Zariski closure $\overline{\mathcal{O}_{\mathbf{m}}}$ is a union of $G_{\mathbf{d}}$ -orbits. This induces a partial order \leq on the set of orbits, which can be explicitly described. More important for us will be the geometric properties of the singular variety $\overline{\mathcal{O}_{\mathbf{m}}}$. The first result in this direction is the following.

Theorem 1.1 ([ADK],[LM]). *The variety $\overline{\mathcal{O}_{\mathbf{m}}}$ is normal.*

This theorem has been generalized to quivers of type A and D with arbitrary orientations by Bobinski-Zwara ([BZ1], [BZ2]), and is conjectured to hold for arbitrary quivers (in the case of the nilpotent cone this is a well-known result of Kostant). In the same spirit, Abeasis, Del Fra and Kraft constructed in [ADK] an explicit desingularization of $\overline{\mathcal{O}_{\mathbf{m}}}$ as follows :

. We first introduce some notation. Let $\underline{\mathbf{d}} = (\mathbf{d}^{(1)}, \dots, \mathbf{d}^{(r)})$ be a sequence of dimension vectors for Δ and let us set $\mathbf{d} = \sum_t \mathbf{d}^{(t)}$. We denote by $\mathcal{F}_{\underline{\mathbf{d}}}$ the variety of flags

$$0 = L_0 \subset L_1 \subset \dots \subset L_r = \bigoplus_i k^{\mathbf{d}_i}$$

of I -graded vector spaces such that $\dim L_t/L_{t-1} = \mathbf{d}^{(t)}$. This is a smooth projective variety. Now consider

$$X_{\underline{\mathbf{d}}} = \{((L_t), x) \in \mathcal{F}_{\underline{\mathbf{d}}} \times E(\mathbf{d}) \mid x(L_i) \subset L_{i-1}\}.$$

This is a vector bundle over $\mathcal{F}_{\underline{\mathbf{d}}}$ of rank $\sum_i \sum_{s < t} \mathbf{d}_i^{(s)} \mathbf{d}_{i-1}^{(t)}$. In particular, it is a smooth (quasi-projective) variety. On the other hand, the second projection $\pi : X_{\underline{\mathbf{d}}} \rightarrow E(\mathbf{d})$ is a projective morphism.

Proposition 1.1. *Let \mathbf{m} be a multipartition and let $x \in \mathcal{O}_{\mathbf{m}}$. Set*

$$(1.1) \quad \mathbf{d}^{(t)} = \dim \text{Ker}(x^t) / \text{Ker}(x^{t-1}).$$

Then $\pi(X_{\underline{\mathbf{d}}}) = \overline{\mathcal{O}_{\mathbf{m}}}$ and $\pi : X_{\underline{\mathbf{d}}} \rightarrow \overline{\mathcal{O}_{\mathbf{m}}}$ is a resolution of singularities.

Proof. Since π is projective and $G_{\mathbf{d}}$ -equivariant, its image is a closed $G_{\mathbf{d}}$ -stable subvariety. Moreover, since $X_{\underline{\mathbf{d}}}$ is irreducible, so is $\pi(X_{\underline{\mathbf{d}}})$. Therefore $\pi(X_{\underline{\mathbf{d}}}) = \overline{\mathcal{O}}$ for some orbit \mathcal{O} . Now let $x \in \pi(X_{\underline{\mathbf{d}}})$. By definition, the points of the fiber of π over x are the flags $(L_t) \in \mathcal{F}_{\underline{\mathbf{d}}}$ which are compatible with x . It is easy to see that such a flag is unique if and only if $x \in \mathcal{O}_{\mathbf{m}}$. On the other hand, if (L_t) is a flag compatible with x and if $(L_t) \in \pi^{-1}(y)$ with $y \in \mathcal{O}_{\mathbf{m}}$ then $x + uy \in \mathcal{O}_{\mathbf{m}}$ for u in a dense subset of \mathbb{C}^* . Thus $x \in \overline{\mathcal{O}_{\mathbf{m}}}$. The Proposition follows. \diamond

. Observe that when $\Delta = \Delta_1^{(1)}$ and $\mathcal{O}_{\mathbf{m}}$ is the regular nilpotent orbit then $X_{\underline{\mathbf{d}}} = T^*\mathfrak{B}$ where \mathfrak{B} is the flag variety of GL_d and the above resolution is just the Springer resolution of the nilpotent cone. This construction of desingularizations of orbit closures has been generalized to arbitrary finite type quivers by Reineke ([Re]).

Corollary 1.1. *The dimension of the orbit $\mathcal{O}_{\mathbf{m}}$ is*

$$\dim \mathcal{O}_{\mathbf{m}} = \sum_i \sum_{s < t} \mathbf{d}_i^{(s)} (\mathbf{d}_i^{(t)} + \mathbf{d}_{i-1}^{(t)}).$$

\diamond

2. HALL ALGEBRAS AND CANONICAL BASES

In this section we let Δ stand for either Δ_n or $\Delta_n^{(1)}$.

2.1. We briefly recall the definition of Ringel-Hall algebras. Let $k = \mathbb{F}_q$ be a finite field with q elements and let $\mathcal{C} = \text{Mod}_k(\Delta)$. It is known that \mathcal{C} satisfies the following conditions

- i) $\dim \text{Ext}^i(M, N) < \infty$ for any $M, N \in \mathcal{C}$, $i = 0, 1$,
- ii) $\text{Ext}^i(M, N) = 0$ for any $i > 1$ and $M, N \in \mathcal{C}$.

Introduce two bilinear forms on $K_0(\mathcal{C}) \simeq \mathbb{Z}^I$ by the formulas

$$\langle M, N \rangle = \dim \text{Hom}(M, N) - \dim \text{Ext}^1(M, N),$$

$$(M, N) = \langle M, N \rangle + \langle N, M \rangle.$$

Let us consider a \mathbb{C} -vector space $\mathbf{U}_{\mathcal{C}}$ (or simply \mathbf{U}) with a basis $\{[M]\}$ indexed by the isomorphism classes of objects in \mathcal{C} . Let us choose $v \in \mathbb{C}$ such that $v^{-2} = q$. Finally let us denote by $\mathbf{d}(M) \in \mathbb{N}^I$ the dimension of a representation M .

. We define an algebra structure on \mathbf{U} by setting

$$(2.1) \quad [M] \cdot [N] = v^{\mathbf{d}(M) \cdot \mathbf{d}(N) - \langle N, M \rangle} \sum_{[P]} F_{M,N}^P [P]$$

where

$$\mathbf{d}_1 \cdot \mathbf{d}_2 = \dim (G_{\mathbf{d}_1 + \mathbf{d}_2} / G_{\mathbf{d}_1} \times G_{\mathbf{d}_2}) = 2 \sum_i \mathbf{d}_{1,i} \mathbf{d}_{2,i},$$

$$F_{M,N}^P = \#\{Q \subset P \mid Q \simeq M, P/Q \simeq N\}.$$

In order to define a bialgebra structure, we need to slightly enlarge \mathbf{U} . Let $\tilde{\mathbf{U}}$ be the algebra generated by \mathbf{U} and elements K_M , $M \in K_0(\mathcal{C})$ subject to the relations $K_M K_N = K_{M+N}$ and $K_M[P] = v^{-(M,P)}[P]K_M$. We define a bialgebra structure on $\tilde{\mathbf{U}}$ by putting

$$\Delta(K_M) = K_M \otimes K_M,$$

$$(2.2) \quad \Delta([P]) = \sum_{[M],[N]} v^{-\mathbf{d}(M) \cdot \mathbf{d}(N) - \langle N, M \rangle} \frac{|\text{Aut}(M)| \cdot |\text{Aut}(N)|}{|\text{Aut}(P)|} F_{M,N}^P [M] K_N \otimes [N].$$

Finally, define a scalar product on $\tilde{\mathbf{U}}$ by the formula

$$\langle [M] K_I, [N] K_J \rangle_G = v^{-(I,J) - \mathbf{d}(M) \cdot \mathbf{d}(N)} \delta_{M,N} \frac{(1 - v^2)^{|\mathbf{d}(M)|}}{|\text{Aut}(M)|}$$

where we set $|\mathbf{d}| = \sum_i \mathbf{d}_i$.

Theorem 2.1 ([Gre]). *($\tilde{\mathbf{U}}, \cdot, \Delta$) is a bialgebra and $\langle \cdot, \cdot \rangle_G$ is a nondegenerate Hopf pairing, i.e*

$$\langle xy, z \rangle_G = \langle x \otimes y, \Delta(z) \rangle_G.$$

Remarks. i) The algebra \mathbf{U} is clearly $K_0(\mathcal{C}) = \mathbb{Z}^I$ -graded. The same holds for $\tilde{\mathbf{U}}$ where the elements K_N are put in degree 0.

ii) Our conventions differ slightly from the usual ones regarding Ringel-Hall algebras. The standard basis $(\{M\})$ of \mathbf{U} is related to ours by the change of variables $[M] = v^{|\mathbf{d}(M)| - \frac{1}{2}\mathbf{d}(M) \cdot \mathbf{d}(M)} \{M\}$, and we renormalized the scalar product $\langle \cdot, \cdot \rangle_G$ by the factor $(1 - v^2)^{|\mathbf{d}(M)|}$. This will make the relation to symmetric functions more transparent.

2.2. Recall that the set of isomorphism classes of objects in $Mod_k(\Delta)$ is in bijection with a set of multipartitions.

Proposition 2.1 ([Mac], [Ri2]). *Let $\mathbf{m}, \mathbf{n}, \mathbf{l}$ be arbitrary multipartitions. There exists a polynomial $P_{\mathbf{m}, \mathbf{n}}^{\mathbf{l}} \in \mathbb{Z}[t]$ such that for any finite field k with q elements we have $F_{\mathbf{m}, \mathbf{n}}^{\mathbf{l}} = P_{\mathbf{m}, \mathbf{n}}^{\mathbf{l}}(q)$.*

The nice consequence of the above Proposition is that we now may (and we now will) consider the Hall algebras $\mathbf{U}_{\mathcal{C}}$ and $\tilde{\mathbf{U}}_{\mathcal{C}}$ for a *generic* q , i.e as $\mathbb{C}[v, v^{-1}]$ -algebras. When $\Delta = \Delta_1^{(1)}$ then the isomorphism classes of objects are labelled by usual partitions and the above polynomials are the *Hall polynomials* (see [Mac]).

2.3. Let us denote by $\mathbb{C}_{G_{\mathbf{d}}}[E(\mathbf{d})]$ the set of $G_{\mathbf{d}}$ -invariant functions on $E(\mathbf{d})$. Observe that we have a natural identification of vector spaces

$$\mathbf{U} \simeq \bigoplus_{\mathbf{d}} \mathbb{C}_{G_{\mathbf{d}}}[E(\mathbf{d})], \quad [\mathbf{m}] \mapsto \mathbf{1}_{\mathcal{O}_{\mathbf{m}}}.$$

In this context, Lusztig gave in [Lu0] a geometric construction of the generic Hall algebra \mathbf{U} , in which one works over the algebraically closed field $\overline{\mathbb{F}}_q$ and where $G_{\mathbf{d}}$ -invariant functions are replaced by (complexes of) $G_{\mathbf{d}}$ -equivariant constructible sheaves (see Saito's paper in this volume). Observe that in our case there are only finitely many $G_{\mathbf{d}}$ -orbits and these induce an algebraic stratification of $E(\mathbf{d})$. In particular, for every orbit $\mathcal{O}_{\mathbf{m}} \subset E(\mathbf{d})$, let $IC_{\mathcal{O}_{\mathbf{m}}}$ be the simple $G_{\mathbf{d}}$ -equivariant perverse sheaf associated to the constant sheaf $\mathbb{C}_{\mathcal{O}_{\mathbf{m}}}$. We set

$$\begin{aligned} \mathbf{f}_{\mathbf{m}} &= v^{\dim \mathcal{O}_{\mathbf{m}}} \mathbf{1}_{\mathcal{O}_{\mathbf{m}}}, \\ \mathbf{b}_{\mathbf{m}} &= \sum_{i, \mathbf{m}'} v^{-i + \dim \mathcal{O}_{\mathbf{m}} - \dim \mathcal{O}_{\mathbf{m}'}} \dim \mathcal{H}_{\mathcal{O}_{\mathbf{m}'}}^i(IC_{\mathcal{O}_{\mathbf{m}}}) \mathbf{f}_{\mathbf{m}'}, \end{aligned}$$

where $\mathcal{H}_{\mathcal{O}_{\mathbf{m}'}}^i(IC_{\mathcal{O}_{\mathbf{m}}})$ is the stalk over any point of $\mathcal{O}_{\mathbf{m}'}$ of the i th cohomology sheaf of $IC_{\mathcal{O}_{\mathbf{m}}}$. The bases $\{\mathbf{f}_{\mathbf{m}}\}_{\mathbf{m}}$ and $\{\mathbf{b}_{\mathbf{m}}\}_{\mathbf{m}}$ are called the *PBW basis* and the *canonical basis* respectively. Observe that Green's scalar product restricts to a nondegenerate form on \mathbf{U} . It will be important for us to consider the *dual PBW basis* $\{\mathbf{f}_{\mathbf{m}}^*\}$ and the *dual canonical basis* $\{\mathbf{b}_{\mathbf{m}}^*\}$ with respect to \langle, \rangle_G . Note that $\mathbf{f}_{\mathbf{m}}^*$ and $\mathbf{f}_{\mathbf{m}}$ only differ by a scalar in $\mathbb{C}[v, v^{-1}]$. For $\mathbf{d} \in \mathbb{N}^I$ we simply denote by $\mathbf{f}_{\mathbf{d}} (= \mathbf{b}_{\mathbf{d}})$ the basis element associated to the unique closed (=semisimple) orbit.

2.4. Finally, we note the following well-known recursive procedure to compute the canonical basis $\{\mathbf{b}_{\mathbf{m}}\}$ of \mathbf{U} . The Verdier duality functor on the (derived) category of constructible $G_{\mathbf{d}}$ -sheaves on $E(\mathbf{d})$ descends to a semilinear involution $u \mapsto \bar{u}$ on \mathbf{U} . This involution can be computed :

Proposition 2.2. *The involution $u \mapsto \bar{u}$ is the unique ring homomorphism $\mathbf{U} \rightarrow \mathbf{U}$ such that $\bar{v} = v^{-1}$ and $\bar{\mathbf{f}}_{\mathbf{d}} = \mathbf{f}_{\mathbf{d}}$ for all $\mathbf{d} \in \mathbb{N}^I$.*

Proof. It is standard that the involution $u \mapsto \bar{u}$ has the above properties. Let $\mathbf{U}' \subset \mathbf{U}$ be the subalgebra generated $\mathbf{f}_{\mathbf{d}}$ for $\mathbf{d} \in \mathbb{N}^I$. It is enough to prove that $\mathbf{U}' = \mathbf{U}$. We argue by induction on the (partial) degeneration order on the set of orbits. Let \mathbf{m} be a multipartition and assume that $\mathbf{f}_{\mathbf{m}'} \in \mathbf{U}'$ for all $\mathbf{m}' \neq \mathbf{m}$ such that $\mathcal{O}_{\mathbf{m}'} \subset \overline{\mathcal{O}_{\mathbf{m}}}$. Pick $x \in \mathcal{O}_{\mathbf{m}}$ and define dimension vectors $\mathbf{d}^{(t)}$, $t = 1, \dots, r$ as in (1.1). Then from Proposition 1.1. we have

$$\mathbf{f}^{\mathbf{d}_r} \dots \mathbf{f}^{\mathbf{d}_1} \in v^{\mathbb{Z}} \mathbf{f}_{\mathbf{m}} \oplus \bigoplus_{\mathbf{m}' < \mathbf{m}} \mathbb{C}[v, v^{-1}] \mathbf{f}_{\mathbf{m}'}$$

and hence $\mathbf{f}_{\mathbf{m}} \in \mathbf{U}'$. ◇

The properties of simple perverse sheaves now translate as follows.

Proposition 2.3. *The canonical basis element \mathbf{b}_m is characterized by the two conditions*

- i) $\overline{\mathbf{b}_m} = \mathbf{b}_m$,
- ii) $\mathbf{b}_m \in \mathbf{f}_m \oplus \bigoplus_{m' < m} v\mathbb{C}[v]\mathbf{f}_{m'}$.

◇

3. HALL ALGEBRAS AND SYMMETRIC FUNCTIONS

3.1. In this section we assume that $\Delta = \Delta_1^{(1)}$ and set $\mathbf{U}_1^{(1)} = \mathbf{U}_{Mod_k(\Delta)}$. In that case, the set of isomorphism classes of objects in $Mod_k(\Delta)$ is in natural bijection with the set Π of all partitions, and we will denote by the same letter λ the isoclass corresponding to a partition λ .

3.2. Let $\Gamma = \mathbb{C}[x_1, x_2, \dots]^{\infty}$ be Macdonald's ring of symmetric polynomials in infinitely many variables. This ring is equipped with various bases. Following classical notations, we denote by p_λ the power-sum symmetric function, by e_λ the elementary symmetric function, and by s_λ the Schur function.

Theorem 3.1 ([Hal]). *The assignment $\mathbf{f}_{(1r)} \mapsto e_r$ extends to an isomorphism of algebras $\phi : \mathbf{U}_1^{(1)} \xrightarrow{\sim} \Gamma \otimes_{\mathbb{C}} \mathbb{C}[v, v^{-1}]$. Moreover, $\phi(\mathbf{f}_\lambda) = P_\lambda(v^2)$ where P_λ is the Hall-Littlewood polynomial.*

Now let us consider the scalar product $\langle \cdot, \cdot \rangle_G$. From the above theorem one can deduce by direct computation :

Proposition 3.1. *The scalar product $\langle \cdot, \cdot \rangle = \phi^* \langle \cdot, \cdot \rangle_G$ coincides with the Hall-Littlewood scalar product :*

$$\langle p_\lambda, p_\mu \rangle = \delta_{\lambda, \mu} z_\lambda \frac{1}{\prod_i (1 - v^{2\lambda_i})},$$

where $z_{(1^{m_1} 2^{m_2} \dots)} = \prod_i m_i! i^{m_i}$.

Observe that the form $\langle \cdot, \cdot \rangle$ on $K_0(\mathcal{C})$ vanishes here. It follows that $\mathbf{U}_1^{(1)}$ can itself be equipped with a coproduct by the formula

$$\Delta_0([P]) = \sum_{[M], [N]} v^{-\mathbf{d}(M) \cdot \mathbf{d}(N)} \frac{|\text{Aut}(M)| \cdot |\text{Aut}(N)|}{|\text{Aut}(P)|} F_{M,N}^P[M] \otimes [N].$$

From Green's theorem (Theorem 2.1) and the above proposition we see that $\phi : \mathbf{U}_1^{(1)} \xrightarrow{\sim} \Gamma \otimes_{\mathbb{C}} \mathbb{C}[v, v^{-1}]$ is an isomorphism of bialgebras, where $\Gamma \otimes_{\mathbb{C}} \mathbb{C}[v, v^{-1}]$ is equipped with Zelevinsky's coproduct (see [Z2]).

Finally, the canonical basis also admits a nice interpretation in terms of symmetric functions :

Proposition 3.2 ([Lu1]). *We have $\phi(\mathbf{b}_\lambda) = s_\lambda$. Hence $\phi(\mathbf{b}_\lambda^*) = S_\lambda(v)$ where $S_\lambda(v)$ is the dual Schur function.*

We will give a geometric interpretation of the above theorem in Section 7.

4. HALL ALGEBRAS AND QUANTUM GROUPS

4.1. Let $A = (a_{ij})_{i=1,\dots,r}$ be a symmetric generalized Cartan matrix. Associated to A is a complex Kac-Moody algebra \mathfrak{g} with a Cartan decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$. The quantized enveloping algebra $\mathbf{U}_v(\mathfrak{g})$ of \mathfrak{g} is the $\mathbb{C}(v)$ -algebra generated by elements $E_i, F_i, K_i^{\pm 1}$ for $i = 1, \dots, r$ satisfying the relations

$$K_i K_i^{-1} = K_i^{-1} K_i = 1, \quad K_i K_j = K_j K_i,$$

$$K_i E_j = v^{a_{ij}} E_j K_i, \quad K_i F_j = v^{-a_{ij}} F_j K_i, \quad [E_i, F_j] = \delta_{ij} \frac{K_i - K_i^{-1}}{v - v^{-1}},$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k E_i^{(k)} E_j E_i^{(1-a_{ij}-k)} = \sum_{k=0}^{1-a_{ij}} (-1)^k F_i^{(k)} F_j F_i^{(1-a_{ij}-k)} = 0 \quad \text{if } i \neq j,$$

where as usual

$$[k] = \frac{v^k - v^{-k}}{v - v^{-1}}, \quad [k]! = [1] \cdots [k], \quad E_i^{(k)} = \frac{E_i^k}{[k]!}, \quad F_i^{(k)} = \frac{F_i^k}{[k]!}.$$

The Lusztig integral form $\mathbf{U}_A(\mathfrak{g})$ is the $\mathbb{C}[v, v^{-1}]$ -subalgebra generated by $E_i^{(k)}, F_j^{(k)}$ and $K_i^{\pm 1}$ for $i = 1, \dots, r$ and $k \geq 1$. The above algebras are equipped with a Hopf algebra structure with a coproduct

$$\Delta(K_i) = K_i \otimes K_i, \quad \Delta(E_i) = E_i \otimes K_i + 1 \otimes E_i, \quad \Delta(F_i) = F_i \otimes 1 + K_i^{-1} \otimes F_i.$$

We will denote by $\mathbf{U}_A^-(\mathfrak{g}) \subset \mathbf{U}_A(\mathfrak{g})$ the subalgebra generated by $F_i^{(k)}$ for $i = 1, \dots, r$ and $k \geq 1$.

. To an arbitrary quiver Q with vertex set I and edge set Ω is associated a Cartan matrix A as above : we put $r = |I|$ and let a_{ij} be the number of (unoriented) edges between i and j . In particular, Δ_n corresponds to the Lie algebra \mathfrak{sl}_{n+1} while $\Delta_n^{(1)}$ corresponds to the affine Lie algebra $\widehat{\mathfrak{sl}}_n$ (for $n > 1$).

4.2. Let us first assume that $\Delta = \Delta_n$ and set $\mathbf{U}_n = \mathbf{U}_{\text{Mod}_k(\Delta_n)}$. In that case we have the following theorem of Ringel :

Theorem 4.1 ([Ri1]). *The assignment $F_i^{(k)} \mapsto [S_i^{\oplus k}]$ for $i = 1, \dots, n$ and $k \geq 1$ extends to an isomorphism of algebras $\phi : \mathbf{U}_A^-(\mathfrak{sl}_{n+1}) \xrightarrow{\sim} \mathbf{U}_n$.*

4.3. For the rest of this section we assume that $\Delta = \Delta_n^{(1)}$ and set $\mathbf{U}_n^{(1)} = \mathbf{U}_{\text{Mod}_k(\Delta_n^{(1)})}$.

The map ϕ defined above gives rise to an embedding $\mathbf{U}_A^-(\widehat{\mathfrak{sl}}_n) \rightarrow \mathbf{U}_n^{(1)}$ but it is no longer surjective (this can easily be checked by comparing graded dimensions). Let us denote by $\mathbf{C}_n \subset \mathbf{U}_n^{(1)}$ the image of ϕ . We first give a description of \mathbf{C}_n .

. Recall that the set of isomorphism classes of objects in \mathcal{C} is indexed by (ordered) n -tuples of partitions $\mathbf{m} = (\lambda^{(1)}, \dots, \lambda^{(n)})$. Call such a multipartition *aperiodic* if, for every $k > 0$ there exists a least one i such that k is not a part of $\lambda^{(i)}$. Let us denote by \mathcal{M} the set of all multipartitions and by \mathcal{M}^{ap} the subset of aperiodic multipartitions.

Theorem 4.2 ([Lu4]). *We have*

$$\mathbf{C}_n = \bigoplus_{\mathbf{m} \in \mathcal{M}^{ap}} \mathbb{C}[v, v^{-1}] \mathbf{b}_{\mathbf{m}}.$$

4.4. Let us call a multipartition $\mathbf{m} = (\lambda^{(1)}, \dots, \lambda^{(n)})$ *completely periodic* if we have $\lambda^{(1)} = \dots = \lambda^{(n)}$, and let $\mathcal{M}^{per} \subset \mathcal{M}$ be the set of all such multipartitions. Consider the space

$$\mathbf{K} = \bigoplus_{\mathbf{m} \in \mathcal{M}^{per}} \mathbb{C}[v, v^{-1}] \mathbf{b}_{\mathbf{m}}^*.$$

Observe that \mathbf{K} has nonzero terms only in periodic degrees $\mathbf{d} = (d, \dots, d)$ (and hence it is in fact \mathbb{Z} -graded).

Theorem 4.3 ([S1]). *The following holds :*

- i) \mathbf{K} is a central subalgebra of $\mathbf{U}_n^{(1)}$,
- ii) The multiplication map induces an isomorphism of algebras

$$\mathbf{K} \otimes_{\mathbb{C}[v, v^{-1}]} \mathbf{C}_n \rightarrow \mathbf{U}_n^{(1)},$$

- iii) The algebra \mathbf{K} is stable under the involution $u \mapsto \bar{u}$ and, as a \mathbb{Z} -graded algebra, $\mathbf{K} \simeq \mathbb{C}[v, v^{-1}][z_1, z_2, \dots]$ where $\deg(z_l) = l$ for $l \geq 1$.

Observe that if $n = 1$ then \mathbf{C}_n is not defined and $\mathbf{U}_1^{(1)} = \mathbf{K}$. Part iii) of the above theorem can then be deduced from Theorem 3.1 and the well-known fact that $\Gamma \simeq \mathbb{C}[z_1, z_2, \dots]$ as a \mathbb{Z} -graded algebra. However, we will see that, as in the case $n = 1$, there is *for every* n a geometrically defined *canonical* identification $\mathbf{K} \simeq \Gamma \otimes \mathbb{C}[v, v^{-1}]$.

Remarks. i) Theorem 4.3 can be reinterpreted as follows : $\mathbf{U}_n^{(1)} \simeq \mathbf{U}_A^-(\widehat{\mathfrak{gl}}_n)$ where $\widehat{\mathfrak{gl}}_n$ is defined as the (universal) central extension

$$0 \rightarrow \mathbb{C}c \rightarrow \widehat{\mathfrak{gl}}_n \rightarrow \mathfrak{gl}_n[z, z^{-1}] \rightarrow 0.$$

The algebra \mathbf{K} is then the quantum analogue of the algebra generated by xz^k for $k \geq 1$, where $x \in Z(\mathfrak{gl}_n)$.

ii) The algebra \mathbf{K} admits another description. Let $\mathbf{e}'_i \in \text{End}(\mathbf{U}_n^{(1)})$ be the adjoint (under $\langle \cdot, \cdot \rangle_G$) of the left multiplication by F_i (the Kashiwara operator familiar in crystal graph theory, see e.g. [HK]). Then $\mathbf{K} = \bigcap_i \text{Ker } \mathbf{e}'_i$. In particular, \mathbf{K} corresponds to the “highest weight” subspace (of the crystal graph of $\mathbf{U}_n^{(1)}$) with respect to the action of the quantum group $\mathbf{U}_A(\widehat{\mathfrak{sl}}_n)$. The above results are in accordance with (and were partly motivated by) [LTV] where the crystal graph of $\mathbf{U}_n^{(1)}$ was completely described, and found to be a disjoint union of copies of the crystal graph of $\mathbf{U}_A^-(\widehat{\mathfrak{sl}}_n)$ indexed by partitions.

iii) In the limit $n \rightarrow \infty$ the diagrams $A_n^{(1)}$ and A_n tend to the common limit A_∞ (i.e. the quiver with vertex set $I + \mathbb{Z}$ and edge set $\Omega = \{(i, i+1)\}$). In that situation, $\mathbf{U}_\infty^{(1)} = \mathbf{U}_\infty = \mathbf{C}_\infty = \mathbf{U}_A^-(\mathfrak{sl}_\infty)$.

iv) Ringel also studied a “degenerate” version of the Hall algebra where one considers representations of the given quiver over \mathbb{C} and uses an Euler characteristic analogue of the convolution product (2.1). This yields a construction of the *Lie algebra* of \mathfrak{sl}_n or $\widehat{\mathfrak{gl}}_n$ rather than of the quantum group (see [Ri3]).

5. FLAG VARIETIES AND HECKE ALGEBRAS

In this section we (briefly) introduce and fix some notation for the more classical geometric objects in type A representation theory such as (partial, affine) flag varieties, (affine) Hecke algebras and some related algebras. Good references for these are [G3] or [CG].

5.1. Let \mathfrak{S}_r denote the symmetric group on r letters. We denote by s_1, \dots, s_{r-1} the simple reflections, so that \mathfrak{S}_r may be presented as the group generated by $s_i, i = 1, \dots, r-1$ with relations

$$(5.1) \quad s_i^2 = 1, \quad s_i s_j = s_j s_i \text{ for } i \neq j \pm 1,$$

$$(5.2) \quad s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}.$$

The extended affine symmetric group is $\widehat{\mathfrak{S}}_r = \mathfrak{S}_r \ltimes \mathbb{Z}^r$. It is isomorphic to the group generated by s_i for $i \in \mathbb{Z}/r\mathbb{Z}$ and τ subject to the relations (5.1), (5.2), $\tau^r = 1$ and $\tau s_i = s_{i+1} \tau$.

As usual we will denote by $l : \mathfrak{S}_r \rightarrow \mathbb{N}$ and $l : \widehat{\mathfrak{S}}_r \rightarrow \mathbb{N}$ the length functions. By definition, we have $l(s_i) = 1$ and $l(\tau) = 0$.

. Now fix $\mathbf{d} = (d_1, \dots, d_n) \in \mathbb{N}^n$ such that $\sum_i d_i = r$. Associated to \mathbf{d} are parabolic subgroups $\mathfrak{S}_{\mathbf{d}} \subset \mathfrak{S}_r$ and $\widehat{\mathfrak{S}}_{\mathbf{d}} \subset \widehat{\mathfrak{S}}_r$ generated by s_i for $i \notin \{d_1, d_1 + d_2, \dots, d_1 + \dots + d_n\}$. Observe that $\widehat{\mathfrak{S}}_{\mathbf{d}}$ is always finite.

5.2. Let k be either \mathbb{F}_q or $\overline{\mathbb{F}_q}$. Let L be an r -dimensional k -vector space. We denote by \mathcal{B} the variety of all complete flags

$$0 = L_0 \subset L_1 \subset \dots \subset L_r = L, \quad \dim L_i = i$$

(a smooth projective variety). The group $GL(L)$ acts transitively on \mathcal{B} . The set of $GL(L)$ -orbits on $\mathcal{B} \times \mathcal{B}$ is in bijection with the symmetric group \mathfrak{S}_r . For $w \in \mathfrak{S}_r$ we denote by \mathcal{O}_w the corresponding orbit.

. Alternatively, we may fix one flag $(L_i)_i \in \mathcal{B}$ and denote by $B \subset GL(L)$ its stabilizer (a Borel subgroup). This provides us with an identification $\mathcal{B} \simeq GL(L)/B$. The set of B -orbits on \mathcal{B} is also in natural bijection with \mathfrak{S}_r and we denote by \mathcal{B}_w the orbit corresponding to w (a Schubert cell). It is known that $\dim \mathcal{B}_w = l(w)$.

. Now let $n \in \mathbb{N}$ and fix $\mathbf{d} = (d_1, \dots, d_n) \in \mathbb{N}^n$ such that $\sum_i d_i = r$. Let $\mathcal{F}_{\mathbf{d}}$ be the variety of n -step partial flags

$$0 = L_0 \subset L_1 \subset \dots \subset L_n = L, \quad \dim L_i/L_{i-1} = d_i.$$

This is again a smooth projective variety. As above, let $(L_i)_i$ be a fixed flag in $\mathcal{F}_{\mathbf{d}}$ and let $P \subset GL(L)$ be its stabilizer (a parabolic subgroup). This identifies $\mathcal{F}_{\mathbf{d}}$ with $GL(L)/P$. The P -orbits on $\mathcal{F}_{\mathbf{d}}$ are now parametrized by double cosets $\mathfrak{S}_{\mathbf{d}} \backslash \mathfrak{S}_r / \mathfrak{S}_{\mathbf{d}}$. We again denote by $\mathcal{F}_{\mathbf{d},w}$ the orbit corresponding to $w \in \mathfrak{S}_{\mathbf{d}} \backslash \mathfrak{S}_r / \mathfrak{S}_{\mathbf{d}}$ (a generalized Schubert cell).

. The Zariski closures $\overline{\mathcal{B}_w}$ or $\overline{\mathcal{F}_{\mathbf{d},w}}$ of Schubert cells are called *Schubert varieties*.

5.3. We consider an affine analogue of the above notions. Set $\mathbb{L} = k^r((z))$. By definition, a *lattice* in \mathbb{L} is a free $k[[z]]$ -submodule $L \subset \mathbb{L}$ of rank r . Let \mathcal{G} denote the variety of all lattices. This infinite-dimensional space is a countable union of (finite-dimensional) smooth projective varieties (indeed, let us fix one lattice L_0 ; then for every $l \in \mathbb{N}$ let $X_l \subset \mathcal{G}$ be the smooth projective, finite-dimensional variety of lattices L with $z^l L_0 \subset L \subset z^{-l} L_0$; we have $X_l \subset X_{l+1}$ and $\mathcal{G} = \cup_l X_l$).

. Now let \mathcal{X} be the variety of all infinite flags of lattices $\dots \subset L_i \subset L_{i+1} \subset \dots$ such that $\dim_k L_{i+1}/L_i = 1$ and which satisfy the periodicity condition $L_{i+r} = z^{-1} L_i$. This is again a union of finite-dimensional smooth projective varieties. The map $\mathcal{X} \rightarrow \mathcal{G}, (L_i) \mapsto L_0$ is a smooth locally trivial B -fibration. The group $GL_{k((z))}(\mathbb{L})$ acts transitively on \mathcal{X} and the set of $GL_{k((z))}(\mathbb{L})$ -orbits on $\mathcal{X} \times \mathcal{X}$ is in natural

bijection with $\widehat{\mathfrak{S}}_r$: for $j \in \mathbb{Z}/r\mathbb{Z}$, s_j corresponds to the orbit consisting of pairs $((L_i), (L'_i))$ such that $L_i = L'_i$ for all $i \not\equiv j \pmod{r}$; the orbit $\{((L_i), (L'_i)) \mid L_i = L'_{i+1} \forall i\}$ corresponds to τ . For any $w \in \widehat{\mathfrak{S}}_r$ we will write \mathcal{O}_w for the associated orbit in $\mathcal{X} \times \mathcal{X}$.

As in Section 5.2, we may also fix a flag $(L_i) \in \mathcal{X}$ and denote by $I \subset GL_{k((z))}(\mathbb{L})$ its stabilizer (an Iwahori subgroup). This identifies \mathcal{X} with $GL_{k((z))}(\mathbb{L})/I$. The set of I -orbits on \mathcal{X} is also in natural bijection with $\widehat{\mathfrak{S}}_r$ and we will denote by \mathcal{X}_w the orbits corresponding to $w \in \widehat{\mathfrak{S}}_r$ (an affine Schubert cell). As in the finite type case, we have $\dim \mathcal{X}_w = l(w)$.

. Finally, let $\mathbf{d} = (d_1, \dots, d_n) \in \mathbb{N}^n$ be such that $\sum_i d_i = r$. We set

$$\mathcal{Y}_{\mathbf{d}} = \{(L_i) \mid L_i \subset L_{i+1}, \quad \dim_k L_i/L_{i-1} = d_{i \bmod n}, \quad L_{i+n} = z^{-1}L_i\}.$$

As before, there is a (noncanonical) identification $\mathcal{Y}_{\mathbf{d}} \simeq GL_{k((z))}(\mathbb{L})/P$ for some (parahoric) subgroup P ; the P -orbits \mathcal{Y}_w on $\mathcal{Y}_{\mathbf{d}}$ are indexed by elements $w \in \widehat{\mathfrak{S}}_{\mathbf{d}} \backslash \widehat{\mathfrak{S}}_r / \widehat{\mathfrak{S}}_{\mathbf{d}}$ (see [Lu3] for a formula for $\dim \mathcal{Y}_w$).

5.4. Define the Hecke algebra \mathbf{H}_r (resp. the affine Hecke algebra $\widehat{\mathbf{H}}_r$) as the unital $\mathbb{C}[v, v^{-1}]$ -algebra generated by elements T_w , $w \in \mathfrak{S}_r$ (resp. $w \in \widehat{\mathfrak{S}}_r$) satisfying the following relations

$$\begin{aligned} T_w T_{w'} &= T_{ww'} & \text{if } l(ww') &= l(w) + l(w'), \\ (T_{s_i} + 1)(T_{s_i} - v^{-2}) &= 0, & \text{for all } i. \end{aligned}$$

We set $\tilde{T}_w = v^{l(w)} T_w$ for every $w \in \mathfrak{S}_r$ (resp. $w \in \widehat{\mathfrak{S}}_r$).

In the case of an affine Hecke algebra, there is another very useful presentation, due to Bernstein.

Theorem 5.1 (see [Lu5]). *The algebra $\widehat{\mathbf{H}}_r$ is isomorphic to the algebra generated by $T_i^{\pm 1}$, $i = 1, \dots, r-1$ and $X_j^{\pm 1}$, $j = 1, \dots, r$ with relations*

$$\begin{aligned} T_i T_i^{-1} &= 1 = T_i^{-1} T_i, & (T_i + 1)(T_i - v^{-2}) &= 0, \\ T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}, & |i - j| > 1 &\Rightarrow T_i T_j = T_j T_i, \\ X_i X_i^{-1} &= 1 = X_i^{-1} X_i, & X_i X_j &= X_j X_i, \\ T_i X_i T_i &= v^{-2} X_{i+1}, & j \neq i, i+1 &\Rightarrow X_j T_i = T_i X_j. \end{aligned}$$

The isomorphism between the two presentations is such that $T_{s_i} \mapsto T_i$ and $\tilde{T}_{\lambda}^{-1} \mapsto X_1^{\lambda_1} \dots X_r^{\lambda_r}$ if $\lambda = (\lambda_1, \dots, \lambda_r) \in \mathbb{Z}^r$ is *dominant*, i.e if $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r$. The center of $\widehat{\mathbf{H}}_r$ is $Z(\widehat{\mathbf{H}}_r) = \mathbb{C}[v, v^{-1}][X_1^{\pm 1}, \dots, X_r^{\pm 1}]^{\mathfrak{S}_r}$.

. The relevance of Hecke algebras for us and the link with the previous paragraphs comes from the following geometric construction. Let us assume that $k = \mathbb{F}_q$. Let us set $G = GL_{k((z))}(\mathbb{L})$ and denote by $\mathbb{C}_G[\mathcal{X} \times \mathcal{X}]$ the space of G -invariant functions whose supports lie on a finite union of orbits. We equip $\mathbb{C}_G[\mathcal{X} \times \mathcal{X}]$ with the algebra structure given by the convolution product

$$f \star g = p_{13!}(p_{12}^*(f) \cdot p_{23}^*(g))$$

where $p_{ij} : \mathcal{X} \times \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ is the projection on the i th and j th components. Observe that this product is well-defined since for any two orbits $\mathcal{O}_w, \mathcal{O}_{w'}$, the projection $p_{13} : (\mathcal{O}_w \times \mathcal{X}) \cap (\mathcal{X} \times \mathcal{O}_{w'}) \rightarrow \mathcal{X} \times \mathcal{X}$ is proper.

In the next theorem, we specialize the Hecke algebra by setting $v^{-2} = q$.

Theorem 5.2 ([IM]). *The assignment $T_w \mapsto \mathbf{1}_{\mathcal{O}_w}$ defines an isomorphism of algebras $\widehat{\mathbf{H}}_r \simeq \mathbb{C}_G[\mathcal{X} \times \mathcal{X}]$.*

. It follows from the above Theorem that the structure constants for the convolution algebra $\mathbb{C}_G[\mathcal{X} \times \mathcal{X}]$ are polynomials in q . Hence we may (and we will) consider the convolution algebra as a $\mathbb{C}[v, v^{-1}]$ -algebra.

Naturally, the same result holds in the case of the usual flag variety \mathcal{B} , and we obtain in this way a geometric construction of the finite Hecke algebra \mathbf{H}_r .

5.5. Let us now consider parabolic analogues of the constructions in the previous paragraph. We again assume that $k = \mathbb{F}_q$. Let us set

$$\mathcal{Y} = \sqcup_{\mathbf{d}} \mathcal{Y}_{\mathbf{d}},$$

where in the sum \mathbf{d} runs over all sequences $(d_1, \dots, d_n) \in \mathbb{N}^n$ satisfying $\sum d_i = r$.

The group $G = GL_{k((z))}(\mathbb{L})$ acts on $\mathcal{Y} \times \mathcal{Y}$. It is easy to check that the G -orbits in $\mathcal{Y}_{\mathbf{d}} \times \mathcal{Y}_{\mathbf{d}'}$ are indexed by $\widehat{\mathfrak{S}}_{\mathbf{d}} \backslash \widehat{\mathfrak{S}}_r / \widehat{\mathfrak{S}}_{\mathbf{d}'}$. Without risk of confusion, we denote by $\mathcal{O}_w \subset \mathcal{Y}_{\mathbf{d}} \times \mathcal{Y}_{\mathbf{d}'}$ the orbit corresponding to w .

. For $w \in \widehat{\mathfrak{S}}_{\mathbf{d}} \backslash \widehat{\mathfrak{S}}_r / \widehat{\mathfrak{S}}_{\mathbf{d}'}$, we set $T_w = \sum_{\delta \in w} T_{\delta}$ and we let $\widehat{\mathbf{H}}_{\mathbf{d}\mathbf{d}'} \subset \widehat{\mathbf{H}}_r$ be the $\mathbb{C}[v, v^{-1}]$ -linear span of the elements T_w for $w \in \widehat{\mathfrak{S}}_{\mathbf{d}} \backslash \widehat{\mathfrak{S}}_r / \widehat{\mathfrak{S}}_{\mathbf{d}'}$. Set $e_{\mathbf{d}} = \sum_{\delta \in \widehat{\mathfrak{S}}_{\mathbf{d}}} T_{\delta}$. Then $\widehat{\mathbf{H}}_{\mathbf{d}\mathbf{d}'} = e_{\mathbf{d}} \widehat{\mathbf{H}}_r e_{\mathbf{d}'}$. Put

$$\widehat{\mathbf{S}}_{n,r} = \bigoplus_{\mathbf{d}, \mathbf{d}'} \widehat{\mathbf{H}}_{\mathbf{d}\mathbf{d}'}.$$

We equip this space with the multiplication

$$e_{\mathbf{d}} h e_{\mathbf{d}'} \bullet e_{\mathbf{c}} h' e_{\mathbf{c}'} = \delta_{\mathbf{d}'\mathbf{c}} e_{\mathbf{d}} h e_{\mathbf{d}'} h' e_{\mathbf{c}'} \in \widehat{\mathbf{H}}_{\mathbf{d}\mathbf{c}'} \quad \text{for all } h, h' \in \widehat{\mathbf{H}}_r.$$

. Of course, replacing everywhere \mathcal{Y} by \mathcal{F} and $\widehat{\mathfrak{S}}_r$ by \mathfrak{S}_r , we obtain the definition of the algebra $\mathbf{S}_{n,r}$.

In the next theorem, we specialize the Hecke algebra by setting $v^{-2} = q$.

Theorem 5.3 ([VV1]). *The linear map $\widehat{\mathbf{S}}_{n,r} \rightarrow \mathbb{C}_G[\mathcal{Y} \times \mathcal{Y}]$ such that $T_w \mapsto \mathbf{1}_{\mathcal{O}_w}$ is an algebra isomorphism.*

Again a similar result holds in the finite type case. Finally, as in Section 5.4, we deduce from the above Theorem that $\mathbb{C}_G[\mathcal{Y} \times \mathcal{Y}]$ is naturally a $\mathbb{C}[v, v^{-1}]$ -algebra.

5.6. As for Hall algebras, there is a construction of the Hecke algebras involving (complexes of) equivariant constructible sheaves, rather than G -invariant functions. We will not need such a construction and refer to e.g. [G3] or [V2] for details. A consequence of this construction is the existence of canonical bases for these algebras, which we now define.

Let $w \in \widehat{\mathfrak{S}}_r$. We put

$$\mathbf{c}_w = \sum_{i,x} v^{-i+l(w)-l(x)} \dim \mathcal{H}_{\mathcal{X}_x}^i (IC_{\mathcal{X}_w}) \tilde{T}_x$$

where $IC_{\mathcal{X}_w}$ denotes the intersection cohomology complex associated to $\mathcal{C}_{\mathcal{X}_w}$ and where \mathcal{H}^i stands for local cohomology. The elements $\{\mathbf{c}_w \mid w \in \widehat{\mathfrak{S}}_r\}$ form the canonical (or *Kazhdan-Lusztig*) $\mathbb{C}[v, v^{-1}]$ -basis of $\widehat{\mathbf{H}}_r$.

. Similarly, fix \mathbf{d}, \mathbf{d}' and let $w \in \widehat{\mathfrak{S}}_{\mathbf{d}} \setminus \widehat{\mathfrak{S}}_r / \widehat{\mathfrak{S}}_{\mathbf{d}'}$. Let $L_{\mathbf{d}}$ be any point in $\mathcal{Y}_{\mathbf{d}}$ and denote by $\mathcal{Y}_{w, \mathbf{d}}$ the fiber above $L_{\mathbf{d}}$ of the projection of $\mathcal{O}_w \rightarrow \mathcal{Y}_{\mathbf{d}}$ on the first component. This is an algebraic variety of dimension $y(w)$, say (an explicit formula for $y(w)$ can be found in [Lu3]). For every $w \in \widehat{\mathfrak{S}}_{\mathbf{d}} \setminus \widehat{\mathfrak{S}}_r / \widehat{\mathfrak{S}}_{\mathbf{d}'}$ set

$$\mathbf{c}_w = \sum_{i, x} v^{-i+y(w)-y(x)} \dim \mathcal{H}_{\mathcal{Y}_{x, \mathbf{d}}}^i(IC_{\mathcal{Y}_{w, \mathbf{d}}}) \tilde{T}_x.$$

The elements $\bigsqcup_{\mathbf{d}, \mathbf{d}'} \{\mathbf{c}_w \mid w \in \widehat{\mathfrak{S}}_{\mathbf{d}} \setminus \widehat{\mathfrak{S}}_r / \widehat{\mathfrak{S}}_{\mathbf{d}'}\}$ form the canonical basis of $\widehat{\mathbf{S}}_{n, r}$.

. In addition, as for Hall algebras, there is a purely algebraic characterization of the elements of the canonical bases of the algebras \mathbf{H}_r and $\mathbf{S}_{n, r}$.

Let us first consider the semilinear involution $u \mapsto \bar{u}$ of $\widehat{\mathbf{H}}_r$ defined by $\bar{T}_w = T_{w^{-1}}^{-1}$.

Theorem 5.4 ([KL1], [KL2]). *For each $w \in \widehat{\mathfrak{S}}_r$ the element $\mathbf{c}_w \in \widehat{\mathbf{H}}_r$ is uniquely determined by*

$$\text{i) } \overline{\mathbf{c}_w} = \mathbf{c}_w, \quad \text{ii) } \mathbf{c}_w = \tilde{T}_w + \sum_{x < w} P_{x, w}(v) \tilde{T}_x, \quad P_{x, w}(v) \in v\mathbb{C}[v].$$

The polynomial $P_{x, w}(v)$ is the affine Kazhdan-Lusztig polynomial of type \tilde{A}_{r-1} associated to w and x

Similarly, define a semilinear involution $\tau : \widehat{\mathbf{H}}_{\mathbf{d}\mathbf{d}'} \rightarrow \widehat{\mathbf{H}}_{\mathbf{d}\mathbf{d}'}$ by $\tau(u) = v^{-2l(w_{\mathbf{d}'})} \bar{u}$ where $w_{\mathbf{d}'} \in \widehat{\mathfrak{S}}_{\mathbf{d}'}$ is the longest element. The previous theorem generalizes as follows.

Theorem 5.5. *For each $w \in \widehat{\mathfrak{S}}_{\mathbf{d}} \setminus \widehat{\mathfrak{S}}_r / \widehat{\mathfrak{S}}_{\mathbf{d}'}$ the element \mathbf{c}_w is uniquely determined by the properties :*

$$\text{i) } \tau(\mathbf{c}_w) = \mathbf{c}_w, \quad \text{ii) } \mathbf{c}_w = \tilde{T}_w + \sum_{x < w} Q_{x, w}(v) \tilde{T}_x, \quad Q_{x, w}(v) \in v\mathbb{C}[v].$$

. Again, we have given the results in the affine case but they are all valid in the finite type case as well.

6. CANONICAL BASES AND KAZHDAN-LUSZTIG POLYNOMIALS

We now give the fundamental constructions (due to Lusztig and Zelevinsky) relating representations of quivers of type A to flag varieties. In particular, this will identify the coefficients of the transition matrix between the canonical basis $\{\mathbf{b}_{\mathbf{m}}\}$ and the PBW basis $\{\mathbf{f}_{\mathbf{m}}\}$ with Kazhdan-Lusztig polynomials. The strategy is as follows : we construct a compactification of an orbit closure $\overline{\mathcal{O}_{\mathbf{m}}}$ which is a (generalized) Schubert variety, in such a way that the stratification by Schubert cells restricts on $\overline{\mathcal{O}_{\mathbf{m}}}$ to the stratification by $G_{\mathbf{d}}$ -orbits and use Theorem 5.4. In this section, $k = \mathbb{C}$ or $k = \overline{\mathbb{F}}_q$.

6.1. Let us first treat the finite type case. Let $\mathbf{d} = (d_1, \dots, d_n) \in \mathbb{N}^n$ be a dimension vector, and let us set $\sum d_i = r$ and

$$V_i = k^{d_i}, \quad V^i = \bigoplus_{j \leq i} V_j, \quad L = \bigoplus_i V_i.$$

Then (V^i) is a partial flag belonging to $\mathcal{F}_{\mathbf{d}}$. Let us denote by $P \subset GL(L)$ its stabilizer. We construct a map $\iota : E(\mathbf{d}) \rightarrow \mathcal{F}_{\mathbf{d}}$ as follows. To a representation $(\phi_i) \in E(\mathbf{d})$ we associate the flag defined by

$$U_i = \{(x_1, \dots, x_n) \in V_1 \oplus \dots \oplus V_n \mid x_{j+1} = \phi_j(x_j) \text{ for } j \geq i\}.$$

Proposition 6.1 ([Z1]). *The map ι is an embedding of $E(\mathbf{d})$ as a dense open subset of the (generalized) Schubert variety*

$$\overline{\mathcal{F}_{w_0}} := \{(U_i) \mid U_i \supset V^{i-1} \text{ for all } i\}.$$

Moreover, each $G_{\mathbf{d}}$ -orbit $\mathcal{O}_{\mathbf{m}} \subset E(\mathbf{d})$ embeds as a dense open subset of a P -orbit $\mathcal{F}_{w(\mathbf{m})}$ for some $w(\mathbf{m}) \in \mathfrak{S}_{\mathbf{d}} \backslash \mathfrak{S}_r / \mathfrak{S}_{\mathbf{d}}$.

Proof. It is clear that $\iota(E(\mathbf{d})) \subset \overline{\mathcal{F}_{w_0}}$. Conversely, it is easy to see that $(U_i) \in \overline{\mathcal{F}_{w_0}}$ belongs to $\iota(E(\mathbf{d}))$ if and only if $U_i \cap \bigoplus_{j>i} V_j = 0$ for all i (an open condition). Finally note that $G_{\mathbf{d}}$ embeds as a Levi subgroup of P and that ι is $G_{\mathbf{d}}$ -equivariant. \diamond

Remark. Let $\mathcal{U} \subset \overline{\mathcal{F}_{w_0}}$ be any P -orbit and pick $(U_i) \in \mathcal{U}$. Then

$$(U_i/V^{i-1}, \phi_i = \text{Id} : U_i/V^{i-1} \rightarrow U_{i+1}/V^i)$$

defines a representation of Δ_n . Let \mathbf{m} be the multipartition describing this representation. It is easy to check that $\iota(\mathcal{O}_{\mathbf{m}}) \subset \mathcal{U}$. In particular, the assignment $\mathbf{m} \mapsto w(\mathbf{m})$ is a bijection between the set of $G_{\mathbf{d}}$ -orbits in $E(\mathbf{d})$ and the set of P -orbits (or Schubert cells) in $\overline{\mathcal{F}_{w_0}}$.

. Now let us fix $\mathcal{O}_{\mathbf{m}} \subset E(\mathbf{d})$. By the previous Proposition, the simple perverse sheaf $IC_{\mathcal{O}_{\mathbf{m}}}$ is the restriction to $\overline{\mathcal{O}_{\mathbf{m}}}$ of the simple perverse sheaf $IC_{\overline{\mathcal{F}_{w_0}}}$. In particular, for any $\mathcal{O}_{\mathbf{n}} \subset \overline{\mathcal{O}_{\mathbf{m}}}$ we have

$$\dim \mathcal{H}_{\mathcal{O}_{\mathbf{n}}}^i(IC_{\mathcal{O}_{\mathbf{m}}}) = \dim \mathcal{H}_{\mathcal{F}_{w(\mathbf{n})}}^i(IC_{\mathcal{F}_{w(\mathbf{m})}}).$$

Next, consider the natural map $\pi : \mathcal{B} \simeq G/B \rightarrow G/P \simeq \mathcal{F}_{\mathbf{d}}$ given by

$$(0 = U_0 \subset U_1 \cdots \subset U_r = L) \mapsto (U_{d_1} \subset U_{d_1+d_2} \subset \cdots \subset U_r = L).$$

This is a locally trivial fibration with smooth projective fiber P/B . Thus, for any $w_1, w_2 \in \mathfrak{S}_{\mathbf{d}} \backslash \mathfrak{S}_r / \mathfrak{S}_{\mathbf{d}}$ there holds

$$\dim \mathcal{H}_{\mathcal{F}_{w_1}}^i(IC_{\mathcal{F}_{w_2}}) = \dim \mathcal{H}_{\pi^{-1}(\mathcal{F}_{w_1})}^i(IC_{\pi^{-1}(\mathcal{F}_{w_2})}).$$

Finally, $\pi^{-1}(\mathcal{F}_w) = \sqcup_{y \in w} \mathcal{B}_y$ (a finite union of B -orbits). Since \mathcal{F}_w is irreducible, there exists a unique open B -orbit \mathcal{B}_{y_w} in $\pi^{-1}(\mathcal{F}_w)$ (corresponding to the longest element $y(w) \in w$). Thus we obtain

$$\begin{aligned} \dim \mathcal{H}_{\mathcal{O}_{\mathbf{n}}}^i(IC_{\mathcal{O}_{\mathbf{m}}}) &= \dim \mathcal{H}_{\pi^{-1}(\mathcal{F}_{w(\mathbf{n})})}^i(IC_{\pi^{-1}(\mathcal{F}_{w(\mathbf{m})})}) \\ &= \dim \mathcal{H}_{\mathcal{B}_{y(w(\mathbf{n}))}}^i(IC_{\mathcal{B}_{y(w(\mathbf{m}))}}). \end{aligned}$$

Corollary 6.1. *The coefficients of the transition matrix between the canonical basis $\{\mathbf{b}_{\mathbf{m}}\}$ and the PBW basis $\{\mathbf{f}_{\mathbf{m}}\}$ of \mathbf{U}_n are Kazhdan-Lusztig polynomials (of type $A_{\sum d_i - 1}$), where \mathbf{d} is determined by \mathbf{m} via $\mathcal{O}(\mathbf{m}) \subset E_{\mathbf{d}}$.*

6.2. Let us now briefly describe the affine analogue of the above construction, which is due to Lusztig. Let $\mathbf{d} = (d_1, \dots, d_n) \in \mathbb{N}^n$ and set $r = \sum d_i$. For clarity, we denote by $\bar{i} \in \mathbb{Z}/n\mathbb{Z}$ the class of an integer $i \in \mathbb{Z}$. Set $V_i = k^{d_i}$, $L = \bigoplus_i V_i$ and $\mathbb{L} = L((z))$. Let $\cdots \subset \mathcal{L}_0 \subset \mathcal{L}_1 \subset \cdots$ be the n -periodic flag of lattices defined by

$$\begin{aligned} \mathcal{L}_0 &= L[[z]], \\ \mathcal{L}_i &= \mathcal{L}_0 \oplus z^{-1}V_1 \oplus \cdots \oplus z^{-1}V_i \quad \text{for } 1 \leq i \leq n, \\ \mathcal{L}_{j+n} &= z^{-1}\mathcal{L}_j \quad \text{for all } j. \end{aligned}$$

Let $P \subset GL_k((z))(\mathbb{L})$ be its stabilizer. We construct a map $E(\mathbf{d}) \rightarrow \mathcal{Y}_{\mathbf{d}}$ as follows : let $0 \leq \bar{i} < n$ be the residue of i modulo n and set $l(j) = (j - \bar{j})/n$; let $(\phi_i) \in E(\mathbf{d})$ and set

$$\tilde{\phi}_i = \begin{cases} \phi_i & \text{if } \bar{i} \neq 0 \\ z^{-1}\phi_i & \text{if } \bar{i} = 0, \end{cases}$$

and $\phi^h(x_i) = \tilde{\phi}_{i+h} \cdots \tilde{\phi}_{i+1} \tilde{\phi}_i(x_i)$ if $x_i \in V_i$. Note that for any fixed x_i , $\phi^h(x_i) = 0$ for h big enough. Now we put $\iota((\phi_i)) = (\mathcal{N}_j)$ where $\mathcal{N}_j = \mathcal{L}_{j-1} \oplus z^{-l(j-1)-1}\mathcal{U}_{j-1}$ and

$$\mathcal{U}_j = \left\{ \sum_{h \geq 0} \phi^h(x_{\bar{j}}) \mid x_{\bar{j}} \in V_{\bar{j}} \right\}.$$

. Just as in the finite type case, one may prove :

Proposition 6.2 ([Lu3]). *The map ι is an embedding of $E(\mathbf{d})$ as a dense open subset of the (generalized) affine Schubert variety*

$$\overline{\mathcal{Y}_{w_0}} := \{(\mathcal{N}_i) \mid \mathcal{N}_i \supset \mathcal{L}_{i-1} \text{ for all } i\}.$$

Moreover, each $G_{\mathbf{d}}$ -orbit $\mathcal{O}_{\mathbf{m}} \subset E(\mathbf{d})$ embeds as a dense open subset of a P -orbit $\mathcal{Y}_{w(\mathbf{m})}$ for some $w(\mathbf{m}) \in \widehat{\mathfrak{S}}_{\mathbf{d}} \backslash \widehat{\mathfrak{S}}_r / \widehat{\mathfrak{S}}_{\mathbf{d}}$.

Corollary 6.2. *The coefficients of the transition matrix between the canonical basis $\{\mathbf{b}_{\mathbf{m}}\}$ and the PBW basis $\{\mathbf{f}_{\mathbf{m}}\}$ of $\mathbf{U}_n^{(1)}$ are affine Kazhdan-Lusztig polynomials (of type $A_{\sum d_i - 1}^{(1)}$).*

Remark. Nakajima constructed in [N3] a similar compactification of $E(\mathbf{d})$ for an arbitrary quiver. However, it is only in type A that the coefficients of the canonical basis are given by Kazhdan-Lusztig polynomials (see, however [BZ2] for some descriptions of singularities of orbit closures in type D in terms of *products* of Grassmanians).

7. SCHUR-WEYL DUALITY IN THE GEOMETRIC SETTING

7.1. Let us first briefly recall the usual Schur-Weyl duality. Let V be an n -dimensional \mathbb{C} -vector space. There are two commuting group actions on $V^{\otimes r}$: $GL(V)$ acts diagonally while \mathfrak{S}_r acts by permutation of the factors. Moreover we have

$$\begin{aligned} \mathbb{C}[\mathfrak{S}_r] &= \text{End}_{GL(V)}(V^{\otimes r}) & \text{if } n \geq r, \\ U(\mathfrak{gl}(V)) &\twoheadrightarrow \text{End}_{\mathfrak{S}_r}(V^{\otimes r}) \end{aligned}$$

The algebra $\text{End}_{\mathfrak{S}_r}(V^{\otimes r})$ is called the Schur algebra. There are affine and quantum versions of this duality. The affine version is obtained by replacing \mathfrak{gl}_n by $\widehat{\mathfrak{gl}}_n = \mathfrak{gl}_n[z, z^{-1}] \oplus \mathbb{C}c$, the symmetric group \mathfrak{S}_r by $\widehat{\mathfrak{S}}_r$ and V by $V[z, z^{-1}]$.

. In the quantum version, we replace \mathfrak{gl}_n by the quantum group $\mathbf{U}_v(\mathfrak{gl}_n)$. Recall that this is the $\mathbb{C}(v)$ -Hopf algebra generated by elements $E_i, F_i, i = 1, \dots, n-1$ and $H_i^{\pm 1}, i = 1, \dots, n$ satisfying relations

$$\begin{aligned} H_i H_i^{-1} &= H_i^{-1} H_i = 1, & H_i H_j &= H_j H_i, \\ H_i E_j &= v^{a_{ij}} E_j H_i, & H_i F_j &= v^{-a_{ij}} F_j H_i, \\ [E_i, F_j] &= \delta_{ij} \frac{H_i H_{i+1}^{-1} - H_i^{-1} H_{i+1}}{v - v^{-1}}, \\ [E_i, E_j] &= [F_i, F_j] = 0 & \text{if } |i - j| > 1, \\ E_i^{(2)} E_j - E_i E_j E_i + E_j E_i^{(2)} &= F_i^{(2)} F_j - F_i F_j F_i + F_j F_i^{(2)} = 0 & \text{if } |i - j| = 1. \end{aligned}$$

where (a_{ij}) is the Cartan matrix of type A_{n-1} . The coproduct is defined as in Section 4.1.

The standard representation \mathbb{C}^n of \mathfrak{gl}_n admits a quantization V_q (the “vector” representation). Using the coproduct Δ we define an action of $\mathbf{U}_v(\mathfrak{gl}_n)$ on $V_q^{\otimes r}$. It is well-known that $\mathbf{U}_v(\mathfrak{gl}_n)$ is a *quasitriangular Hopf algebra*, i.e there exists an element $\mathcal{R} \in \mathbf{U}_v(\mathfrak{gl}_n) \otimes \mathbf{U}_v(\mathfrak{gl}_n)$ satisfying the relation $\mathcal{R}\Delta = \Delta^{op}\mathcal{R}$. From this we construct an action of the Artin braid group on r strands B_r on $V_q^{\otimes r}$ by the formula

$$\sigma_i(v_1 \otimes \cdots \otimes v_r) = v_1 \otimes \cdots \otimes P\mathcal{R}(v_i \otimes v_{i+1}) \otimes \cdots \otimes v_r,$$

where σ_i is the i th standard generator of B_r and $P : u \otimes v \mapsto v \otimes u$ is the permutation (see e.g [ES] for more details on quasitriangular Hopf algebras). One shows that the action of B_r factorizes through the Hecke algebra \mathbf{H}_r , and that, as in the classical case, we have

$$\begin{aligned} \mathbf{H}_r &= \text{End}_{\mathbf{U}_v(\mathfrak{gl}_n)}(V_q^{\otimes r}) \quad \text{if } n \geq r, \\ \mathbf{U}_v(\mathfrak{gl}_n) &\twoheadrightarrow \text{End}_{\mathbf{H}_r}(V_q^{\otimes r}) \end{aligned}$$

The algebra $\text{End}_{\mathbf{H}_r}(V_q^{\otimes r})$ is called the q -Schur algebra.

. Finally, the quantum affine version is now obtained by replacing $\mathbf{U}_v(\mathfrak{gl}_n)$ by $\mathbf{U}_v(\widehat{\mathfrak{gl}}_n)$, the space V_q by its affine version $V_q[z, z^{-1}]$, and \mathbf{H}_r by $\widehat{\mathbf{H}}_r$. However, there is a subtle point here: $\mathbf{U}_v(\widehat{\mathfrak{gl}}_n)$ is the quantization of $\mathfrak{gl}_n[z, z^{-1}] \oplus \mathbb{C}c$ rather than the quantization of the Kac-Moody algebra $\widehat{\mathfrak{sl}}_n$ or the quantization of the slightly bigger algebra $\widehat{\mathfrak{gl}}'_n := \mathfrak{gl}_n \oplus \bigoplus_{l \in \mathbb{Z}^*} z^l \mathfrak{sl}_n \oplus \mathbb{C}c$ (unfortunately, this latter algebra is often also denoted $\widehat{\mathfrak{gl}}_n$ in the litterature). In particular, the Hopf algebra $\mathbf{U}_v(\widehat{\mathfrak{gl}}_n)$ is best defined through the *Drinfeld new realization* (see [D]) rather than through quantum Serre-type relations. We refer to [GV] for the precise definition.

Remark. In the affine versions of the Schur-Weyl duality, the modules $V[z, z^{-1}]$ and $V_q[z, z^{-1}]$ are of level zero, i.e the central element c (or its quantum analog) act trivially. Thus, for the purposes of Schur-Weyl duality it is enough to consider the loop algebra $L\mathfrak{gl}_n = \mathfrak{gl}_n[z, z^{-1}]$ or its quantization.

7.2. Following Beilinson-Lusztig-MacPherson in the finite-type case and Ginzburg-Vasserot and Lusztig in the affine case, we will now give a geometric construction of the (quantum) Schur-Weyl dualities. We assume here that $k = \mathbb{F}_q$.

. Recall the notations of Section 5 : L is a fixed r -dimensional k -vector space, \mathcal{B} the flag variety and $\mathcal{F} = \sqcup_{\mathbf{d}} \mathcal{F}_{\mathbf{d}}$ the n -step partial flag variety. For any $i = 1, \dots, n$ we consider the subvarieties of $\mathcal{F} \times \mathcal{F}$

$$\mathcal{O}_i^+ = \{((L_j), (L'_j)) \mid L_j = L'_j \text{ if } j \neq i \text{ and } L'_i \subset L_i, \dim L_i/L'_i = 1\},$$

$$\mathcal{O}_i^- = \{((L_j), (L'_j)) \mid L_j = L'_j \text{ if } j \neq i \text{ and } L_i \subset L'_i, \dim L'_i/L_i = 1\},$$

and for any \mathbf{d} we set $\mathcal{O}_{\mathbf{d}}^0 = \Delta \mathcal{F}_{\mathbf{d}} \subset \mathcal{F}_{\mathbf{d}} \times \mathcal{F}_{\mathbf{d}}$.

Theorem 7.1 ([BLM]). *The assignment $E_i \mapsto \mathbf{1}_{\mathcal{O}_i^+}$, $F_i \mapsto \mathbf{1}_{\mathcal{O}_i^-}$, $H_i \mapsto \sum_{\mathbf{d}} q^{d_i} \mathbf{1}_{\mathcal{O}_{\mathbf{d}}^0}$ extends to a surjective homomorphism of algebras*

$$\mathbf{U}_v(\mathfrak{gl}_n) \twoheadrightarrow \mathbb{C}_G[\mathcal{F} \times \mathcal{F}] \otimes_{\mathbb{C}[v, v^{-1}]} \mathbb{C}(v).$$

. Next we consider the affine case. Thus $\dim L = r$, $\mathbb{L} = L((z))$, \mathcal{X} is the affine flag variety and \mathcal{Y} is the affine n -step partial flag variety. Again we set

$$\mathcal{O}_i^+ = \{((L_j), (L'_j)) \mid L_j = L'_j \text{ if } j \neq i \text{ and } L'_i \subset L_i, \dim_k L_i/L'_i = 1\},$$

$$\mathcal{O}_i^- = \{((L_j), (L'_j)) \mid L_j = L'_j \text{ if } j \neq i \text{ and } L_i \subset L'_i, \dim_k L'_i/L_i = 1\},$$

and $\mathcal{O}_{\mathbf{d}}^0 = \Delta \mathcal{Y}_{\mathbf{d}} \subset \mathcal{Y}_{\mathbf{d}} \times \mathcal{Y}_{\mathbf{d}}$.

Theorem 7.2 ([GV], [Lu4]). *There exists a surjective algebra homomorphism*

$$\mathbf{U}_v(\widehat{\mathfrak{gl}}_n) \rightarrow \mathbb{C}_G[\mathcal{Y} \times \mathcal{Y}] \otimes_{\mathbb{C}[v, v^{-1}]} \mathbb{C}(v).$$

On the subalgebra $U_v(\widehat{\mathfrak{gl}}'_n)$ it is given by $E_i \mapsto \mathbf{1}_{\mathcal{O}_i^+}$, $F_i \mapsto \mathbf{1}_{\mathcal{O}_i^-}$, $H_i \mapsto \sum_{\mathbf{d}} q^{d_i} \mathbf{1}_{\mathcal{O}_{\mathbf{d}}^0}$.

. We are now ready to give the geometric version of Schur-Weyl duality. The convolution product defines a left action of $\mathbb{C}_G[\mathcal{F} \times \mathcal{F}]$ on $\mathbb{C}_G[\mathcal{F} \times \mathcal{B}]$ and a right action of $\mathbb{C}_G[\mathcal{B} \times \mathcal{B}]$ on $\mathbb{C}_G[\mathcal{F} \times \mathcal{B}]$. Moreover, it is clear that these two actions commute. Recall the algebra $\mathbf{S}_{n,r} = \mathbb{C}_G[\mathcal{F} \times \mathcal{F}]$ defined in Section 5.

Theorem 7.3 ([GV]). *There exists an isomorphism of algebras*

$$\mathbf{S}_{n,r} \simeq \text{End}_{\mathbf{H}_r}(V_q^{\otimes r})$$

and an isomorphism of $\mathbf{S}_{n,r} - \mathbf{H}_r$ -bimodules $\mathbb{C}_G[\mathcal{F} \times \mathcal{B}] \simeq V_q^{\otimes r}$ such that the following diagram commutes

$$(7.1) \quad \begin{array}{ccccc} \mathbf{S}_{n,r} = \mathbb{C}_G[\mathcal{F} \times \mathcal{F}] & \rightsquigarrow & \mathbb{C}_G[\mathcal{F} \times \mathcal{B}] & \leftarrow & \mathbb{C}_G[\mathcal{B} \times \mathcal{B}] \\ \sim \downarrow & & \sim \downarrow & & \sim \downarrow \\ \text{End}_{\mathbf{H}_r}(V_q^{\otimes r}) & \rightsquigarrow & V_q^{\otimes r} & \leftarrow & \mathbf{H}_r \end{array}$$

(here $A \rightsquigarrow B$ denotes an action of an algebra A on B).

. In a similar way, we have

Theorem 7.4 ([VV1]). *There exists an isomorphism of algebras*

$$\widehat{\mathbf{S}}_{n,r} \simeq \text{End}_{\widehat{\mathbf{H}}_r}(V_q[z, z^{-1}])^{\otimes r}$$

and an isomorphism of $\widehat{\mathbf{S}}_{n,r} - \widehat{\mathbf{H}}_r$ -bimodules $\mathbb{C}_G[\mathcal{Y} \times \mathcal{X}] \simeq (V_q[z, z^{-1}])^{\otimes r}$ such that the following diagram commutes

$$(7.2) \quad \begin{array}{ccccc} \widehat{\mathbf{S}}_{n,r} = \mathbb{C}_G[\mathcal{Y} \times \mathcal{Y}] & \rightsquigarrow & \mathbb{C}_G[\mathcal{Y} \times \mathcal{X}] & \leftarrow & \mathbb{C}_G[\mathcal{X} \times \mathcal{X}] \\ \sim \downarrow & & \sim \downarrow & & \sim \downarrow \\ \text{End}_{\widehat{\mathbf{H}}_r}((V_q[z, z^{-1}])^{\otimes r}) & \rightsquigarrow & (V_q[z, z^{-1}])^{\otimes r} & \leftarrow & \widehat{\mathbf{H}}_r \end{array}$$

7.3. Using a combination of the constructions in Sections 6 and 7, we can give a conceptual interpretation of Theorem 4.3. Put

$$\mathcal{O}^- = \{((L_i), (L'_i)) \mid L'_i \subset L_i \text{ for all } i\} \subset \mathcal{Y} \times \mathcal{Y}.$$

Observe that to a point $((L_i), (L'_i)) \in \mathcal{O}^-$ we may associate the representation of $\Delta_n^{(1)}$ on $\bigoplus_{i \in \mathbb{Z}/n\mathbb{Z}} L_i/L'_i$ using the natural maps $\phi_i : L_i/L'_i \rightarrow L_{i+1}/L'_{i+1}$ (as in Section 6). Note that such a representation is always nilpotent by the periodicity condition for affine flags. For a multipartition \mathbf{m} we denote by $\mathcal{Z}_{\mathbf{m}} \subset \mathcal{O}^-$ the set of

all points whose associated representation of $\Delta_n^{(1)}$ is of type \mathbf{m} . Let us also denote by $\widehat{\mathbf{S}}_{n,r}^-$ the convolution algebra $\mathbb{C}_G[\mathcal{O}^-]$.

Proposition 7.1 ([VV1]). *The linear map $\varphi : \mathbf{U}_n^{(1)} \rightarrow \widehat{\mathbf{S}}_{n,r}^-$ such that $\varphi(\mathbf{1}_{\mathcal{O}_{\mathbf{m}}}) = \mathbf{1}_{Z_{\mathbf{m}}}$ is a surjective algebra homomorphism.*

Proof. This follows directly from the definitions of the Hall product and of the convolution product. \diamond

. To sum up, we have the following diagram of geometric constructions of algebras :

$$\mathbf{U}_n^{(1)} \longrightarrow \mathbb{C}_G[\mathcal{O}^-] \hookrightarrow \mathbb{C}_G[\mathcal{Y} \times \mathcal{Y}] \rightsquigarrow \mathbb{C}_G[\mathcal{Y} \times \mathcal{X}] \rightsquigarrow \mathbb{C}_G[\mathcal{X} \times \mathcal{X}]$$

On the other hand, the bicommutant Theorem 7.4 gives us a canonical embedding $Z(\widehat{\mathbf{H}}_r) \subset \widehat{\mathbf{S}}_{n,r}^-$. Recall that by Bernstein's theorem,

$$Z(\widehat{\mathbf{H}}_r) = \mathbb{C}[v, v^{-1}][X_1^{\pm 1}, \dots, X_r^{\pm 1}]^{\mathfrak{S}_r}.$$

Let us set $Z^-(\widehat{\mathbf{H}}_r) = \mathbb{C}[v, v^{-1}][X_1^{-1}, \dots, X_r^{-1}]^{\mathfrak{S}_r}$.

Proposition 7.2 ([S1]). *The morphism φ restricts to a surjective algebra homomorphism $i_r : \mathbf{K} \rightarrow Z^-(\widehat{\mathbf{H}}_r)$. Moreover, the maps i_r are compatible with the natural projective system $Z^-(\widehat{\mathbf{H}}_r) \rightarrow Z^-(\widehat{\mathbf{H}}_{r-1})$ and yield in the projective limit an isomorphism*

$$i : \mathbf{K} \xrightarrow{\sim} Z^-(\widehat{\mathbf{H}}_{\infty}) \simeq \Gamma \otimes \mathbb{C}[v, v^{-1}].$$

. In particular, this map gives rise, for each n , to two new natural families of symmetric polynomials, namely $i(\mathbf{b}_{(\lambda, \dots, \lambda)}^*)$ and $i(\pi(\mathbf{b}_{\lambda, \dots, \lambda}))$ where $\pi : \mathbf{U}_n^{(1)} \rightarrow \mathbf{K}$ is the orthogonal projection. When $n = 1$ these families are the dual Schur functions and the Schur functions respectively.

. In the same spirit, we make the following conjecture :

Conjecture. *The restriction of Green's scalar product $\langle \cdot, \cdot \rangle_G$ to $\mathbf{K} \simeq \Gamma \otimes \mathbb{C}[v, v^{-1}]$ is given by the formula*

$$(p_{\lambda}, p_{\mu}) = \delta_{\lambda\mu} z_{\lambda} q^{(n-1)|\lambda|} (1 - q^{-1})^{n|\lambda|} \prod_{i=1}^{l(\lambda)} \frac{1 - q^{n\lambda_i}}{(1 - q^{\lambda_i})^2}.$$

This can be viewed as a generalization of the Hall-Littlewood scalar product.

Remarks. i) All the spaces in diagrams (7.1) and (7.2) are equipped with canonical bases. One can show that all the maps are (in some suitable sense) compatible with these bases (see [VV1], [SV]).

ii) Using the map $i : \mathbf{K} \xrightarrow{\sim} Z^-(\widehat{\mathbf{H}}_{\infty})$ and its compatibility with canonical bases it is possible to give some explicit (but very complicated) expressions for the central elements of $\widehat{\mathbf{H}}_r$ in terms of the Kazhdan-Lusztig basis (see [S2]).

8. REPRESENTATION THEORY OF HECKE ALGEBRAS AND QUANTUM AFFINE ALGEBRAS AND GRADED NILPOTENT ORBITS

. This survey would be very incomplete without mentioning a totally different aspect of quivers of type A and their canonical bases, namely their interpretations as Grothendieck rings of finite-dimensional modules for an affine Hecke algebra and for an affine q -Schur algebra.

8.1. We will first deal with Hecke algebras. We make a few preliminary definitions and remarks. Since $Z(\widehat{\mathbf{H}}_r) = \mathbb{C}[v, v^{-1}][X_1^{\pm 1}, \dots, X_r^{\pm 1}]^{\mathfrak{S}_r}$, any character $\theta : Z(\widehat{\mathbf{H}}_r) \rightarrow \mathbb{C}^*$ is given by an $r+1$ -tuple $(s_1, \dots, s_r; \epsilon)$ such that $\theta(x_i) = s_i$ and $\theta(v) = \epsilon$. It is convenient to view such an $r+1$ -tuple as a pair $(s, \epsilon) \in T \times \mathbb{C}^*$ where $T \subset GL_r(\mathbb{C})$ is the subgroup of diagonal matrices. Note that θ determines s only up to conjugation (i.e permutation). By Schur's lemma, $Z(\widehat{\mathbf{H}}_r)$ acts by a character in any simple finite-dimensional module. Let us denote by $Rep_\theta(\widehat{\mathbf{H}}_r)$ the category of finite-dimensional $\widehat{\mathbf{H}}_r$ -modules of central character θ .

. The link between representation theory of $\widehat{\mathbf{H}}_r$ and quivers of type A stems from the following fundamental theorem.

Theorem 8.1 ([KL3],[G1]). *Fix $\theta = (s, \epsilon)$ and assume that ϵ is not a root of unity. Set $G_s = Stab_{GL_r(\mathbb{C})} s$. There is a bijection between the set of simple modules in $Rep_\theta \widehat{\mathbf{H}}_r$ and the set of G_s -orbits in*

$$\mathcal{N}_\theta = \{x \in \mathfrak{gl}_r, x \text{ nilpotent}, sxs^{-1} = \epsilon^{-2}x\}.$$

. The proof of the above theorem (the so-called Deligne-Langlands conjecture for $\widehat{\mathbf{H}}_r$) is based on a geometric realization of $\widehat{\mathbf{H}}_r$ completely different from the one given in Section 5, as the Grothendieck group of (equivariant) coherent sheaves on a certain variety $Z \subset T^*\mathcal{B} \times T^*\mathcal{B}$ (see [CG],[KL3]). As a part of this realization, two types of modules are constructed for each triple (s, ϵ, x) : the simple module $L_{\theta, x}$ and the so-called *standard module* $M_{\theta, x}$, whose character is easy to compute. The determination of the characters of the simple modules is now a consequence of the following result.

Theorem 8.2. *Let $\theta = (s, \epsilon)$ and assume that ϵ is not a root of unity. For each $x \in \mathcal{N}_\theta$ we set $\mathcal{O}_x = G_s \cdot x$. Then for any two $x, y \in \mathcal{N}_\theta$ the multiplicity of $L_{\theta, x}$ in $M_{\theta, y}$ is*

$$[M_{\theta, x} : L_{\theta, y}] = \sum_k \dim \mathcal{H}_{\mathcal{O}_y}^k(IC_{\mathcal{O}_x}).$$

. Let us reinterpret the above Theorems in terms of quivers A_m . Let us fix ϵ (the value of the deformation parameter v). For simplicity, let us denote by $Rep_\epsilon \widehat{\mathbf{H}}_r$ the category of all finite-dimensional modules of central charge $\theta = (s, \epsilon)$ with $Spec(s) \subset \epsilon^{2\mathbb{Z}}$. For any such s , write $\mathbb{C}^r = \bigoplus_{i \in \mathbb{Z}} V_{\epsilon^{2i}}$ for the corresponding eigenspace decomposition of \mathbb{C}^r . Now observe that $x(V_{\epsilon^{2i}}) \subset V_{\epsilon^{2i-2}}$, i.e x is nothing but a representation on the space \mathbb{C}^r of the quiver A_∞^∞ . Let $\mathbf{m}(s, x)$ be the type of this representation. Altogether, Theorems 8.1 and 8.2 are nicely summarized as follows.

Corollary 8.1 ([A]). *Assume that ϵ is not a root of unity. There exists a linear isomorphism*

$$\Theta_\infty : \bigoplus_r Rep_\epsilon^*(\widehat{\mathbf{H}}_r) \xrightarrow{\sim} \mathbf{U}_{\infty, |v|=1}$$

such that $\Theta_\infty([M_{\theta, x}]^) = \mathbf{f}_{\mathbf{m}(s, x)}$ and $\Theta_\infty([L_{\theta, x}]^*) = \mathbf{b}_{\mathbf{m}(s, x)}$.*

Remark. The above Theorem together with the results of Section 6 shows that, in particular, the multiplicities $[M_{\theta, x} : L_{\theta, y}]$ are equal to evaluation at 1 of some Kazhdan-Lusztig polynomials of type A .

. The case when ϵ^2 is a primitive n th root of unity is more subtle. In that situation, Theorem 8.2 remains true while in Theorem 8.1 only a certain subset of G_s -orbits in \mathcal{N}_θ is relevant. Observe that now if $\text{Spec}(s) \subset \epsilon^{2\mathbb{Z}}$ then to a pair (s, x) is associated a representation of the cyclic quiver $A_n^{(1)}$ rather than A_∞ . The classification of simple modules is best described by the next theorem.

Corollary 8.2 ([A],[Gro]). *Assume that ϵ^2 is a primitive n th root of unity. There exists a linear isomorphism*

$$\Theta_n : \bigoplus_r \text{Rep}_\epsilon^*(\widehat{\mathbf{H}}_r) \xrightarrow{\sim} \mathbf{C}_{n,|v=1} \subset \mathbf{U}_{n,|v=1}^{(1)}$$

such that $\Theta_\infty([L_{\theta,x}]^*) = \mathbf{b}_{\mathbf{m}(s,x)}$.

. In particular, simple modules in $\text{Rep}_\epsilon(\widehat{\mathbf{H}}_r)$ are indexed by aperiodic multipartitions of total dimension r .

Remark. i) Ariki proved much more than the results stated above : he endowed the space $\bigoplus_r \text{Rep}_\epsilon^*(\widehat{\mathbf{H}}_r)$ with an algebra structure (using the restriction functor $\text{Rep}_\epsilon(\widehat{\mathbf{H}}_{r+r'}) \rightarrow \text{Rep}_\epsilon(\widehat{\mathbf{H}}_r) \otimes \text{Rep}_\epsilon(\widehat{\mathbf{H}}_{r'})$) and showed that the map Θ is actually an algebra homomorphism.

ii) A consequence of the above Theorem and of the results in Section 6 is that the multiplicities $[M_{\theta,x} : L_{\theta,y}]$ are now given by evaluation at 1 of *affine* Kazhdan-Lusztig polynomials of type A.

iii) Recently, Vasserot succeeded in extending some of the above results to the case of a *double affine Hecke algebra*, or *Cherednik algebra* [V3].

8.2. Let us now turn to the case of the quantum affine algebra $\mathbf{U}_v(\widehat{\mathfrak{sl}}_k)$. It is known that any finite-dimensional representation of $\mathbf{U}_v(\widehat{\mathfrak{sl}}_k)$ factors through $\widehat{\mathbf{S}}_{k,r}$ for some $r \gg 0$. Hence, by Schur-Weyl duality we expect a very similar classification result as in Section 8.1. A geometric construction of $\widehat{\mathbf{S}}_{k,r}$ in the same spirit as [G1], [KL3] was given in [GV].

Theorem 8.3 ([GV]). *Let $\epsilon \in \mathbb{C}^*$. There is a bijection between the simple finite-dimensional modules of $\mathbf{U}_\epsilon(\widehat{\mathfrak{gl}}_k)$ and the set of pairs*

$$\{(s, x) \mid s \in T, x \in \mathfrak{gl}_r, x^k = 0, sxs^{-1} = \epsilon^{-2}x\} / GL_r(\mathbb{C}).$$

To each such pair is associated a simple module $L_{s,x}$ and a standard module $M_{s,x}$, and we have

$$[M_{s,x} : L_{s',y}] = \delta_{s,s'} \sum_l \dim \mathcal{H}_{\mathcal{O}_y}^l(IC_{\mathcal{O}_x})$$

where \mathcal{O}_z is the G_s -orbit containing z .

. In particular, the classification of simple modules can again be interpreted in terms of quivers of type A. However, observe that, as opposed to Section 8.1., there are no additional restrictions when ϵ is a root of unity. Let $\text{Rep}_\epsilon(\mathbf{U}_v(\widehat{\mathfrak{sl}}_k))$ denote the category of all finite-dimensional modules with a composition series involving only simple modules $L_{s,x}$ with $\text{Spec}(s) \subset \epsilon^{2\mathbb{Z}}$. For notational convenience, we set $A_\infty^{(1)} = A_\infty$ in what follows.

Corollary 8.3 ([VV1]). *Let $n \in \mathbb{N} \cup \{\infty\}$ be the order of ϵ^2 . There exists a linear isomorphism*

$$\Xi_n : \bigoplus_k \text{Rep}_\epsilon^*(\mathbf{U}_v(\widehat{\mathfrak{sl}}_k)) \xrightarrow{\sim} \mathbf{U}_{n,|v=1}^{(1)}$$

such that $\Xi_n([M_{\theta,x}]^*) = \mathbf{f}_{\mathbf{m}(s,x)}$ and $\Xi_n([L_{\theta,x}]^*) = \mathbf{b}_{\mathbf{m}(s,x)}$.

Remark. i) Again, the standard modules $M_{s,x}$ can be explicitly described and their characters computed (see [V1]). Also, as for Hecke algebras, the multiplicities $[M_{\theta,x} : L_{\theta,y}]$ are evaluation at 1 of some Kazhdan-Lusztig polynomials (affine if ϵ is a root of unity).

. One may ask for a similar interpretation for Grothendieck groups of categories of representations of $\mathbf{U}_\epsilon(\mathfrak{sl}_k)$ or \mathbf{H}_r . Although no geometric constructions similar to those in [KL3] or [GV] exist for the *finite type* quantum group or Hecke algebra, there exist evaluation maps $ev : \mathbf{U}_v(\widehat{\mathfrak{sl}}_k) \rightarrow \mathbf{U}_v(\mathfrak{sl}_k)$, $ev : \widehat{\mathbf{H}}_r \rightarrow \mathbf{H}_r$. These in turn induce maps $ev^* : Rep_\epsilon^*(\mathbf{U}_v(\widehat{\mathfrak{sl}}_k)) \rightarrow Rep_\epsilon^*(\mathbf{U}_v(\mathfrak{sl}_k))$ and $ev^* Rep_\epsilon^*(\widehat{\mathbf{H}}_r) \rightarrow Rep_\epsilon^*(\mathbf{H}_r)$. In fact these maps nicely fit in the previous picture, as shown by the following theorems.

Theorem 8.4 ([A],[VV1]). *Let $n \in \mathbb{N} \cup \{\infty\}$ be the order of ϵ^2 . There is a commutative diagram*

$$\begin{array}{ccc} \bigoplus_r Rep_\epsilon^*(\widehat{\mathbf{H}}_r) & \xrightarrow[\Theta_n]{\sim} & \mathbf{C}_{n,v=1} \subset \mathbf{U}_{n|v=1}^{(1)} \\ ev^* \downarrow & & \downarrow \gamma \\ \bigoplus_r Rep_\epsilon^*(\mathbf{H}_r) & \xrightarrow[\theta_n]{\sim} & V_{\Lambda_0} \end{array}$$

where V_{Λ_0} is the highest-weight integrable representation of $\mathbf{C}_n \simeq \mathbf{U}_v^-(\widehat{\mathfrak{sl}}_n)$, and γ is the action on the highest weight vector.

Theorem 8.5 ([VV1]). *Let $n \in \mathbb{N} \cup \{\infty\}$ be the order of ϵ^2 . There is a commutative diagram*

$$\begin{array}{ccc} \bigoplus_k Rep_\epsilon^*(\mathbf{U}_v(\widehat{\mathfrak{sl}}_k)) & \xrightarrow[\Xi_n]{\sim} & \mathbf{U}_{n|v=1}^{(1)} \\ ev^* \downarrow & & \downarrow \gamma \\ \bigoplus_k Rep_\epsilon^*(\mathbf{U}_v(\mathfrak{sl}_k)) & \xrightarrow[\xi_n]{\sim} & \Lambda_{\infty|v=1} \end{array}$$

where Λ_∞ is the Fock space representation of \mathbf{U}_n constructed in [Hay], [VV1], and γ is the action on the vacuum vector of Λ_∞ .

. In addition, the spaces V_{Λ_0} and Λ_∞ are again equipped with two bases: a “canonical basis” and a “PBW basis”. These correspond, under the isomorphisms θ_n and ξ_n to the (dual) bases of simple modules and standard modules respectively. Finally, the map γ is compatible with these canonical bases.

. This beautiful interpretation of character formulae in terms of canonical bases of Hall algebras and Fock spaces or highest weight representations was first conjectured by Lascoux-Leclerc-Thibon in [LLT]. It was proved by Ariki for Hecke algebras (in the more general framework of *cyclotomic Hecke algebras*, which include in particular Hecke algebras of type B) and by Varagnolo and Vasserot for quantum affine algebras. One nice consequence of this construction for quantum affine algebras is that it gives a new proof of Lusztig’s conjecture for the characters of simple $\mathbf{U}_\epsilon(\mathfrak{sl}_k)$ -modules when ϵ is a root of unity.

9. SOME FURTHER GEOMETRIC CONSTRUCTIONS

. To conclude this survey, we briefly mention some other interesting occurrences of quivers of type A in geometry.

9.1. Nakajima quiver varieties. Nakajima recently introduced a new class of algebraic varieties and convolution algebras which play an increasingly important role in geometric representation theory (see [Nak1],[Nak3]). These varieties are defined for an arbitrary quiver, but they have a special interpretation for affine (or finite) Dynkin diagrams as moduli spaces of sheaves on complex surfaces.

. Let us consider the action of the cyclic group $\Gamma = \mathbb{Z}/n\mathbb{Z}$ on \mathbb{C}^2 via the embedding

$$\begin{aligned} \mathbb{Z}/n\mathbb{Z} &\rightarrow SL(2, \mathbb{C}) \\ 1 &\mapsto \begin{pmatrix} \zeta_n & 0 \\ 0 & \zeta_n^{-1} \end{pmatrix} \end{aligned}$$

where ζ_n is a primitive n th root of unity. This action extends to a natural action on $\mathbb{P}^2(\mathbb{C})$, leaving the projective line at infinity l_∞ invariant. The McKay correspondence gives a bijection between the set of irreducible Γ -modules and the set of simple roots (vertices) of the Dynkin diagram $A_{n-1}^{(1)}$. Hence, we may identify the Grothendieck (semi)group $K_0(\text{Rep } \Gamma)$ with the set of dimension vectors of $\Delta_n^{(1)}$. Now let \mathbf{v}, \mathbf{w} be two such dimension vectors, and let V, W be corresponding Γ -modules. The (smooth) Nakajima quiver variety $\mathcal{M}(\mathbf{v}, \mathbf{w})$ associated to $\Delta_n^{(1)}$ is the moduli space of pairs (\mathcal{E}, ϕ) where \mathcal{E} is a Γ -equivariant torsion-free sheaf on $\mathbb{P}^2(\mathbb{C})$ such that $H^1(\mathbb{P}^2, \mathcal{E}(-l_\infty)) \simeq V$ and $\phi : \mathcal{E}_\infty \xrightarrow{\sim} \mathcal{O}_{l_\infty} \otimes W$ is a Γ -invariant isomorphism. Let $\mathcal{M}_{vec}(\mathbf{v}, \mathbf{w}) \subset \mathcal{M}(\mathbf{v}, \mathbf{w})$ be the subvariety consisting of pairs (\mathcal{E}, ϕ) where \mathcal{E} is a vector bundle. The singular Nakajima variety is defined as $\mathcal{M}_0(\mathbf{v}, \mathbf{w}) = \mathcal{M}_{vec}(\mathbf{v}, \mathbf{w}) \times (S\mathbb{C}^2)^\Gamma$. There is a projective map $\pi : \mathcal{M}(\mathbf{v}, \mathbf{w}) \rightarrow \mathcal{M}_0(\mathbf{v}, \mathbf{w})$ given by $(\mathcal{E}, \phi) \mapsto (\mathcal{E}^{**}, \phi, \text{supp}(\mathcal{E}^{**}/\mathcal{E}))$.

. In the McKay correspondence, the extending vertex of the affine Dynkin diagram is associated to the trivial representation ρ_0 . The quiver varieties $\mathcal{M}(\mathbf{v}, \mathbf{w})$ and $\mathcal{M}_0(\mathbf{v}, \mathbf{w})$ for Δ_{n-1} are defined just as in the affine case with the added condition that V and W do not contain ρ_0 .

. Remarkably, the above geometrically defined varieties also admit a very concrete realization in terms of quivers (for arbitrary type). We give it for type $A_{n-1}^{(1)}$, the finite type being (as above) obtained from it as a particular case. Let us consider the *double quiver* of $\Delta_n^{(1)}$, where each arrow is replaced by a pair of arrows going in opposite directions. Let H be the set of oriented edges of this graph. For any $h \in H$ we denote by $o(h)$ and $i(h)$ the outgoing and incoming edges of h respectively. Let $(\mathbf{v}, \mathbf{w}) \in \mathbb{N}^n \times \mathbb{N}^n$ with $\mathbf{v} = (v_a)_{a \in \mathbb{Z}/n\mathbb{Z}}$ and $\mathbf{w} = (w_a)_{a \in \mathbb{Z}/n\mathbb{Z}}$. Fix some $\mathbb{Z}/n\mathbb{Z}$ -graded \mathbb{C} -vector spaces $V = \bigoplus V_a$ and $W = \bigoplus W_a$ such that $\dim V = \mathbf{v}$ and $\dim W = \mathbf{w}$. Set

$$E(V, V) = \bigoplus_{h \in H} \text{Hom}(V_{o(h)}, V_{i(h)}),$$

$$L(V, W) = \bigoplus_k \text{Hom}(V_k, W_k), \quad L(W, V) = \bigoplus_k \text{Hom}(W_k, V_k)$$

and

$$\mathbf{M}(\mathbf{v}, \mathbf{w}) = E(V, V) \oplus L(W, V) \oplus L(V, W).$$

An element of $\mathbf{M}(\mathbf{v}, \mathbf{w})$ will usually be denoted by its components (B, i, j) .

. Let $\epsilon : H \rightarrow \{1, -1\}$ be any function satisfying $\epsilon(h) + \epsilon(\bar{h}) = 0$, where \bar{h} is the edge h with opposite orientation. Consider the subvariety

$$\mathbf{M}^0(\mathbf{v}, \mathbf{w}) = \{(B, i, j) \in \mathbf{M}(\mathbf{v}, \mathbf{w}) \mid \sum_{o(h)=k} \epsilon(h) B_{\bar{h}} B_h + i_k j_k = 0 \text{ for all } k\}.$$

The group $G_V = \prod_k GL(V_k)$ acts on $\mathbf{M}^0(\mathbf{v}, \mathbf{w})$ by conjugation. The quiver varieties are obtained as quotients of the variety by G_V . According to the Geometric Invariant Theory philosophy, there are essentially two ways of doing this. Define an element (B, i, j) to be *semistable* if the following condition holds : the only B -invariant, graded subspace $S \subset \bigoplus_k V_k$ contained in $\text{Ker } j$ is $\{0\}$. Let $\mathcal{M}^{0,ss}(\mathbf{v}, \mathbf{w}) \subset \mathbf{M}^0(\mathbf{v}, \mathbf{w})$ denote the open subvariety of semistable points.

Theorem 9.1 ([N1], [VV2]). *We have $\mathcal{M}_0(\mathbf{v}, \mathbf{w}) = \mathbf{M}^0(\mathbf{v}, \mathbf{w})//G_V$ (categorical quotient) and $\mathcal{M}(\mathbf{v}, \mathbf{w}) = \mathbf{M}^{0,ss}(\mathbf{v}, \mathbf{w})/G_V$ (geometric quotient).*

. Quite remarkably, in type A the quiver varieties are again related to partial flag varieties. Let us consider the case of the quiver Δ_n . Let \mathbf{v}, \mathbf{w} be n -tuples of positive integers, and define another n -tuple \mathbf{a} as follows :

$$a_1 = w_1 + \cdots + w_{n-1} - v_1, \quad a_n = v_{n-1},$$

$$a_i = w_i + \cdots + w_{n-1} - v_i + v_{i-1}, \quad \text{for } i = 2, \dots, n-1.$$

Behold the partial flag variety

$$\mathcal{F}_{\mathbf{a}} = \{U_1 \subset U_2 \subset \cdots \subset U_{n-1} \subset \mathbb{C}^r \mid \dim U_i/U_{i-1} = a_i\},$$

where $r = \sum_i a_i$. The cotangent bundle $T^*(\mathcal{F}_{\mathbf{a}})$ can be identified with the variety of pairs

$$\{((U_i), x) \mid x(U_i) \subset U_{i+1}\} \subset \mathcal{F}_{\mathbf{a}} \times \mathcal{N},$$

where $\mathcal{N} \subset \mathfrak{gl}_r$ denotes the set of nilpotent elements. It is known that the image of the natural projection $p : T^*\mathcal{F}_{\mathbf{a}} \rightarrow \mathcal{N}$ is the Zariski closure of one orbit, say $\mathcal{O}_{\mathbf{a}}$.

. Now let $y \in \mathcal{N}$ be a nilpotent element with w_i Jordan blocks of length i and let $z, h \in \mathfrak{gl}_r$ be such that (y, h, z) is an \mathfrak{sl}_2 -triple. The *Slodowy slice* at y is defined as

$$\mathcal{S}_y = \{u \in \mathcal{N} \mid [u - y, z] = 0\}.$$

The variety \mathcal{S}_y is transverse to all nilpotent orbits containing y in their closure, and plays an important role in the Brieskorn-Slodowy approach to Kleinian singularities. The following was conjectured by Nakajima and proved by Maffei [M].

Theorem 9.2. *We have*

$$\mathcal{M}_0(\mathbf{v}, \mathbf{w}) \simeq \mathcal{S}_y \cap \overline{\mathcal{O}_{\mathbf{a}}}$$

and

$$\mathcal{M}(\mathbf{v}, \mathbf{w}) \simeq p^{-1}(\mathcal{S}_y \cap \overline{\mathcal{O}_{\mathbf{a}}}) \subset T^*\mathcal{F}_{\mathbf{a}}.$$

. We note that a similar link between quiver varieties and transversal slices to nilpotent orbits appears in [MV], and is related there to some infinite Grassmannians.

9.2.Parabolic structures. Let k be an algebraically closed field and let X be a curve defined over k . If $x \in X$ is a smooth point then the category of torsion sheaves (or skyscraper sheaves) on X supported at x is equivalent to the category of finite-dimensional representations of the local ring \mathcal{O}_x at x . Since this is a discrete valuation ring, this category is equivalent to $\text{Mod}_k(\Delta_1^{(1)})$. In particular, the Hall algebra of the category of coherent sheaves on X (over \mathbb{F}_q) contains a copy of $\mathbf{U}_1^{(1)} \simeq \Gamma \otimes \mathbb{C}[v, v^{-1}]$ at each (smooth) point.

. Now let us assume in addition that X is endowed with an action of the cyclic group $G = \mathbb{Z}/n\mathbb{Z}$, and that x is an isolated fixed point for this action. Hence G acts on \mathcal{O}_x as $i \cdot f = \zeta_n^i f$, where $\zeta_n = e^{2i\pi/n}$. In that situation, the category of G -equivariant torsion sheaves supported at x is equivalent to $\text{Mod}_k(\Delta_n^{(1)})$.

. Finally, let us consider the case of a finite group G acting on X , and let $\pi : X \rightarrow X/G \simeq Y$ denote the quotient map. Assume that this map is ramified at points y_1, \dots, y_r with ramification indices p_1, \dots, p_r . Then, locally over each y_i we are in the situation just described with $G = \mathbb{Z}/d_i\mathbb{Z}$. In particular, the category of G -equivariant coherent sheaves on X contains a copy of $\text{Mod}_k(\Delta_{d_i}^{(1)})$ for each i . This has applications in [S3] to the study of Hall algebras of such categories (see also [GL]).

. Closely related to the above is the notion of *parabolic structure*. Let X be a curve and let $x \in X$ be a smooth point. A p -parabolic structure at x on a coherent sheaf \mathcal{F} on X is a flag of coherent sheaves

$$\mathcal{F} = \mathcal{F}_0 \subset \dots \subset \mathcal{F}_p = \mathcal{F}(-x).$$

Note that if \mathcal{F} is locally free at x then we have an induced filtration

$$(\mathcal{F}/\mathcal{F}_0)|_x \subset \dots \subset (\mathcal{F}_p/\mathcal{F}_0)|_x \simeq \mathcal{F}|_x,$$

i.e a parabolic flag in the fiber of \mathcal{F} at x . This notion can be seen as some globalization of the concept of an affine flag as in Section 5. In [FK] the authors consider a similar situation in the case of a surface : using a variant of the Hall algebra, they construct an action of $\widehat{\mathfrak{gl}}_p$ on the cohomology of certain moduli spaces of coherent sheaves on surfaces with p -parabolic structure along a rational curve.

REFERENCES

- [ADK] Abeasis S., Del Fra A., Kraft H., *The geometry of representations of A_m* , Math. Ann. **256**, no. 3, 401-418, (1981).
- [A] Ariki S., *On the decomposition numbers of the Hecke algebra of $G(m, 1, n)$* , J. Math. Kyoto Univ. **36**, no. 4, 789-808, (1996).
- [BLM] Beilinson A., Lusztig G., MacPherson R., *A geometric setting for the quantum deformation of GL_n* , Duke Math. J. **61**, no. 2, 655-677, (1990).
- [BZ1] Bobiński G., Zwara G., *Normality of orbit closures for Dynkin quivers of type A_n* , Manuscripta Math. **105**, no. 1, 103-109, (2001).
- [BZ2] Bobiński G., Zwara G., *Schubert varieties and representations of Dynkin quivers*, preprint.
- [CG] Chriss N., Ginzburg V., *Representation theory and complex geometry*, Birkhuser Boston, Inc., Boston, MA, (1997).
- [D] Drinfeld V., *A new realization of Yangians and of quantum affine algebras. (Russian)* Dokl. Akad. Nauk SSSR **296**, no. 1, 13-17 (1987); translation in Soviet Math. Dokl. **36**, no. 2, 212-216, (1988).
- [ES] Etingof P., Schiffmann O., *Lectures on quantum groups*, Lectures in Mathematical Physics. International Press, Boston, MA, (1998).
- [FK] Finkelberg M., Kuznetsov A., *Parabolic sheaves on surfaces and affine Lie algebra $\widehat{\mathfrak{gl}}_n$* , J. Reine Angew. Math. **529**, 155-203, (2000).
- [GL] Geigle W., Lenzing H., *A class of weighted projective curves arising in the representation theory of finite-dimensional algebras*, Singularities, representations of algebras and vector bundles, Springer LNM **1273**, 265-297, (1987).
- [G1] Ginzburg V., *Deligne-Langlands conjecture and representations of affine Hecke algebras*, preprint Moscow (1985).
- [G2] Ginzburg V., *Perverse sheaves on a Loop group and Langlands' duality*, preprint alg-geom/9511007 (1995).
- [G3] Ginzburg V., *Geometric methods in the representation theory of Hecke algebras and quantum groups (notes by Vladimir Baranovsky)*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., **514**, Representation theories and algebraic geometry (Montreal, PQ, 1997), 127-183, (1998).

- [Gro] Grojnowski I., *Representations of affine Hecke algebras (and affine quantum GL_n) at roots of unity.*, Internat. Math. Res. Notices, no. **5**, (1994).
- [GV] Ginzburg V., Vasserot E., *Langlands reciprocity for affine quantum groups of type A_n .*, Internat. Math. Res. Notices no. **3**, 67-85, (1993).
- [Gre] Green J. A., *Hall algebras, hereditary algebras and quantum groups.*, Invent. Math. **120**, no. 2, 361-377, (1995).
- [Hal] Hall P., *The algebra of partitions*, Proc. 4th Canadian Math. Congress, 147-159, (1959).
- [Hay] Hayashi T., *q -analogues of Clifford and Weyl algebras-spionr and oscillator representations of quantum enveloping algebras*, Comm. Math. Phys. **127**, 129-144, (1990).
- [HK] Hong J., Kang S.-J., *Quantum groups and crystal bases*, Graduate Studies in Mathematics **42**. AMS (2002).
- [IM] Iwahori N., Matsumoto H., *On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups.*, Inst. Hautes Etudes Sci. Publ. Math. No. **25**, 5-48, (1965).
- [KL1] Kazhdan D., Lusztig G., *Representations of Coxeter groups and Hecke algebras.*, Invent. Math. **53**, no. 2, 165-184, (1979).
- [KL2] Kazhdan D., Lusztig G., *Schubert varieties and Poincar duality.* Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), pp. 185-203, Proc. Sympos. Pure Math., XXXVI, Amer. Math. Soc., Providence, R.I., (1980).
- [KL3] Kazhdan D., Lusztig G., *Proof of the Deligne-Langlands conjecture for Hecke algebras.*, Invent. Math. **87**, no. 1, 153-215, (1987).
- [LLT] Lascoux A., Leclerc B., Thibon J.-Y., *Hecke algebras at roots of unity and crystal bases of quantum affine algebras.*, Comm. Math. Phys. **181**, no. 1, 205-263, (1996).
- [LM] Lakshmibai V., Magyar P., *Degeneracy schemes, quiver schemes, and Schubert varieties.*, Internat. Math. Res. Notices, no. **12**, 627-640, (1998).
- [LTV] Leclerc B., Thibon J.-Y., Vasserot E., *Zelevinsky's involution at roots of unity.* J. Reine Angew. Math. **513**, 33-51, (1999).
- [Lu0] Lusztig G., *Introduction to quantum groups*, Progress in Mathematics, **110**. Birkhauser, (1993).
- [Lu1] Lusztig G., *Green polynomials and singularities of unipotent classes.*, Adv. in Math. **42**, no. **2**, 169-178, (1981).
- [Lu2] Lusztig G., *Singularities, character formulas, and a q -analog of weight multiplicities.*, Analysis and topology on singular spaces, II, III (Luminy, 1981), 208-229, Astérisque, **101-102**, Soc. Math. France, Paris, (1983).
- [Lu3] Lusztig G., *Quivers, perverse sheaves, and quantized enveloping algebras.*, J. Amer. Math. Soc. **4**, no. 2, 365-421, (1991).
- [Lu4] Lusztig G., *Aperiodicity in quantum affine \mathfrak{gl}_n .*, Sir Michael Atiyah: a great mathematician of the twentieth century. Asian J. Math. **3**, no. 1, 147-177, (1999).
- [Lu5] Lusztig G., *Transfer maps for quantum affine \mathfrak{gl}_n .*, Representations and quantizations (Shanghai, 1998), 341-356, China High. Educ. Press, Beijing, (2000).
- [Mac] Macdonald I., *Symmetric functions and Hall polynomials*, Oxford press (1979).
- [M] Maffei A., *Quiver varieties of type A*, preprint AG/9812142, (1998).
- [MV] Mirkovic I., Vybornov M., *On quiver varieties and affine Grassmannians of type A*, preprint math/AG0206084 (2002).
- [N1] Nakajima H., *Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras.*, Duke Math. J. **76**, no. 2, 365-416, (1994).
- [N2] Nakajima, H., *Quiver varieties and Kac-Moody algebras.*, Duke Math. J. **91**, no. 3, 515-560, (1998).
- [N3] Nakajima H., *Varieties associated with quivers.*, Representation theory of algebras and related topics (Mexico City, 1994), 139-157, CMS Conf. Proc., 19, (1996).
- [N4] Nakajima H., *Quiver varieties and finite-dimensional representations of quantum affine algebras.*, J. Amer. Math. Soc. **14**, no. 1, 145-238, (2001).
- [Re] Reineke M., *Quivers, desingularizations and canonical bases*, to appear in Studies in memory of Issai Schur, Birkhäuser.
- [Ri1] Ringel C., *Hall algebras and quantum groups*, Invent. Math. **101** no. 3, 583-591, (1990).
- [Ri2] Ringel C., *Hall algebras*, Topics in algebra, Part 1 (Warsaw 1988), PWN, Warsaw, 433-447, (1990).
- [Ri3] Ringel C., *From representations of quivers via Hall and Loewy algebras to quantum groups.* Proceedings of the International Conference on Algebra, Part 2 (Novosibirsk, 1989), 381-401, Contemp. Math., **131**, Part 2, Amer. Math. Soc., Providence, RI, (1992).
- [S1] Schiffmann O., *The Hall algebra of a cyclic quiver and canonical bases of Fock spaces.*, Internat. Math. Res. Notices, no. **8**, 413-440, (2000).
- [S2] Schiffmann O., *On the center of affine Hecke algebras of type A.*, J. Algebra **253**, no. 2, 392-416, (2002).

- [S3] Schiffmann O., *Noncommutative smooth projective curves and quantum loop algebras*, preprint QA/0205267 (2002).
- [SV] Schiffmann O., Vasserot E., *Geometric construction of the global base of the quantum modified algebra of $\widehat{\mathfrak{gl}}_n$* , Transform. Groups **5**, no. 4, 351-360, (2000).
- [V1] Vasserot E., *Affine quantum groups and equivariant K-theory*, Transform. Groups **3**, no. 3, 269-299, (1998).
- [V2] Vasserot E., *On the action of the dual group on the cohomology of perverse sheaves on the affine Grassmannian*, Compositio Math. **131**, no. 1, 51-60, (2002).
- [V3] Vasserot E., *Induced and simple modules of double affine Hecke algebras*, preprint RT/0207127 (2002).
- [VV1] Varagnolo M., Vasserot E., *On the decomposition matrices of the quantized Schur algebra*, Duke Math. J. **100**, no. 2, 267-297, (1999).
- [VV2] Varagnolo M., Vasserot E., *On the K-theory of the cyclic quiver variety*, Internat. Math. Res. Notices, no. **18**, 1005-1028, (1999).
- [Z1] Zelevinsky A., *Two remarks on graded nilpotent classes*, Uspekhi Mat. Nauk **40**, no. **1** (241), 199-200, (1985).
- [Z2] Zelevinsky A., *Representations of finite classical groups. A Hopf algebra approach*, Lecture Notes in Mathematics, **869**, (1981).

Olivier Schiffmann, schiffma@dma.ens.fr,
DMA, École Normale Supérieure, 45 rue d'Ulm, 75230 Paris Cedex 05-FRANCE