

# A simple question about a complicated object

The complicated object is the cohomological induction functor (for which the biblical reference is Knapp & Vogan [KV]). Here is the simple question. For definiteness sake set  $G := PU(n, 1)$ ,  $H := PU(k, 1)$  with  $k \leq n$ , and let  $\mathcal{H}_\rho^n$  (resp.  $\mathcal{H}_\rho^k$ ) be the category of Harish-Chandra modules with the generalized infinitesimal character of the trivial module attached to  $G$  (resp.  $H$ ). I'll surprise nobody by claiming that  $H$  is a subgroup of  $G$ . What's far less obvious, but proved by Khoroshkin [K], is the existence of a full embedding  $F$  of  $\mathcal{H}_\rho^k$  into  $\mathcal{H}_\rho^n$  (Fuser [F, Thm I.4.2] showed that  $F$  is even Ext-full, that is compatible with Ext calculus), prompting the question : is there a geometric interpretation of the embedding  $F$  ? The first candidate for  $F$  is the (ordinary) induction functor ; but this fails miserably — so let's break the pseudo-suspense of this introduction by saying that I claim that  $F$  is (isomorphic to) a certain cohomological induction functor, and conjecture that this phenomenon is general.

## 1. Statements

Let  $G$  be a center free connected semisimple Lie group,  $K \subset G$  a maximal compact subgroup,  $\mathfrak{g} \supset \mathfrak{k}$  the respective complexified Lie algebras. Let's start by recalling the notion of Harish-Chandra module. Say that a  $\mathfrak{g}$ -module  $V$  is  **$\mathfrak{k}$ -finite** if it is a sum of finite dimensional sub- $\mathfrak{k}$ -modules, and that  $V$  is an  **$(\mathfrak{g}, \mathfrak{k})$ -module** if it is  $\mathfrak{k}$ -finite and  $\mathfrak{k}$ -semisimple. The category  $\mathcal{H} = \mathcal{H}(\mathfrak{g}, K)$  of **Harish-Chandra modules** is the full subcategory of  $\mathfrak{g}$ -mod whose objects are those  **$(\mathfrak{g}, \mathfrak{k})$ -modules** of finite length  $V$  such that for any finite dimensional  $\mathfrak{k}$ -invariant subspace  $F \subset V$  the action of  $\mathfrak{k}$  on  $F$  exponentiates to  $K$ . The category  $\mathcal{H}$  is a  $\mathbb{C}$ -category in the sense of Bass [B] page 57. Let  $I$  be the annihilator of the trivial module in the center of  $U(\mathfrak{g})$ , let

$$\mathcal{H}_\rho = \mathcal{H}_\rho(\mathfrak{g}, K)$$

be the full sub- $\mathbb{C}$ -category of  $\mathcal{H}$  whose objects are annihilated by some power of  $I$ , let  $\mathcal{I}$  be the set of isomorphism classes of simple objects of  $\mathcal{H}_\rho$  [it is a finite set] ; for

each  $i \in \mathcal{I}$  choose a representative  $V_i \in i$  and let  $\ell(i)$  be the **projective dimension** of  $V_i$  [*i.e.* the supremum in  $\mathbb{Z} \cup \{+\infty\}$  of the set  $\{n \in \mathbb{Z} \mid \text{Ext}^n(V_i, -) \neq 0\}$ ].

- (1) **Definition.** The  $\mathcal{H}_\rho$ -**ordering** is the smallest partial ordering  $\leq$  on  $\mathcal{I}$  satisfying

$$\left. \begin{array}{l} i, j \in \mathcal{I} \\ \ell(j) = \ell(i) + 1 < \infty \\ \text{Ext}^1(V_j, V_i) \neq 0 \end{array} \right\} \implies i \leq j.$$

- (2) **Definition.** The **sub- $\mathbb{C}$ -category generated by the subset  $\mathcal{J}$  of  $\mathcal{H}_\rho$**  is the full sub- $\mathbb{C}$ -category  $\langle \mathcal{J} \rangle_{\mathcal{H}_\rho}$  of  $\mathcal{H}_\rho$  characterized by the condition that an object  $V$  of  $\mathcal{H}_\rho$  belongs to  $\langle \mathcal{J} \rangle_{\mathcal{H}_\rho}$  iff each simple subquotient of  $V$  is isomorphic to  $V_j$  for some  $j \in \mathcal{J}$ .

- (3) **Definition.** Say that a full sub- $\mathbb{C}$ -category  $\mathcal{C}$  of  $\mathcal{H}_\rho$  is **Ext-full** in  $\mathcal{H}_\rho$  if for all  $V, W \in \mathcal{C}$  the natural morphism

$$\text{Ext}_{\mathcal{C}}^\bullet(V, W) \rightarrow \text{Ext}_{\mathcal{H}_\rho}^\bullet(V, W)$$

is an isomorphism.

For  $i \in \mathcal{I}$  put  $\mathcal{J}_i := \{j \in \mathcal{I} \mid j \leq i\}$ , let  $\theta$  be the Cartan involution of  $(\mathfrak{g}, K)$ , denote by  $d$  the dimension of  $G/K$ , and consider the following

(4) **Property of  $G$ .** For each  $i \in \mathcal{I}$  such that  $V_i$  is unitary the cohomology

$H^{d-\ell(i)}(\mathfrak{g}, K; V_i)$  is nonzero and there is a  $\theta$ -stable parabolic subalgebra of  $\mathfrak{g}$  with Levi subgroup  $L = L_i$  (see Vogan [V2,4.1,4.2] for definitions) satisfying

(a) the corresponding cohomological induction functor  $F$  (see [KV]) sets up an equivalence

$$\mathcal{H}_\rho(\mathfrak{l}, L \cap K) \xrightarrow{\sim} \langle \mathcal{J}_i \rangle_{\mathcal{H}_\rho(\mathfrak{g}, K)} ;$$

(b)  $F\mathbb{C} \simeq V_i$  ;

(c)  $\langle \mathcal{J}_i \rangle_{\mathcal{H}_\rho(\mathfrak{g}, K)}$  is Ext-full in  $\mathcal{H}_\rho(\mathfrak{g}, K)$  ;

(d) if  $a$  is nonzero vector of  $H^{d-\ell(i)}(\mathfrak{g}, K; V_i)$  and  $V$  a simple object of  $\mathcal{H}_\rho(\mathfrak{l}, L \cap K)$ , then the map

$$H^\bullet(\mathfrak{l}, L \cap K; V) \rightarrow H^{d-\ell(i)+\bullet}(\mathfrak{g}, K; FV)$$

$$x \mapsto F(x) \cup a$$

[where  $\cup$  denotes the cup-product] is an isomorphism [of  $H^\bullet(\mathfrak{l}, L \cap K; \mathbb{C})$ -modules] ;

(e) we have  $2\ell(i) = d + \dim L/(L \cap K)$ .

Note once and for all that (e) follows from (a) by the well known argument which consists in setting  $V := \mathbb{C}$  and using Poincaré duality.

(6) **Conjecture.** All center free connected semisimple Lie groups have Property (4).

A partial proof (with explicitly indicated gaps) of the fact that  $PU(n, 1)$ ,  $P\text{Spin}(n, 1)$  and  $SL(3, \mathbb{R})$  have Property (4) is contained in the expanded version of this text, downloadable from

<http://www.iecn.u-nancy.fr/~gaillard/Recherche/Ci/ci.html>

\* \* \*

- [F] Fuser A., Autour de la conjecture d'Alexandru, Thèse de l'Université Nancy 1 (1997).
- [K] Khoroshkin S.M., Category of Harish-Chandra modules of the group  $SU(n,1)$ . *Funct. Anal. Appl.* **14** (1980) 153-155.
- [KV] Knapp A. & Vogan D., **Cohomological induction and unitary representations**, Princeton University Press (1995).
- [V] Vogan D., Cohomology and group representations, *Proc. Symp. Pure Math.* **61** (1997) 219-243.