

Admissible and perfect elements in modular lattices

Rafael Stekolshchik

email: rs2@biu.013.net.il.

ABSTRACT. For the modular lattice $D^4 = \{1 + 1 + 1 + 1\}$ associated with the extended Dynkin diagram \tilde{D}_4 (and also for D^r , where $r > 4$), Gelfand and Ponomarev introduced the notion of *admissible* and *perfect* lattice elements and classified them. In this work, we classify the admissible and perfect elements in the modular lattice $D^{2,2,2} = \{2 + 2 + 2\}$ associated with the extended Dynkin diagram \tilde{E}_6 . Gelfand and Ponomarev constructed admissible elements for D^r recurrently in the length of multi-indices, which they called *admissible sequences*. Here we suggest a direct method for creating admissible elements. Admissible sequences and admissible elements for $D^{2,2,2}$ (resp. D^4) form 14 classes (resp. 11 classes) and possess some periodicity.

If under all indecomposable representations of a modular lattice the image of an element is either zero or the whole representation space, the element is said to be *perfect*. Our classification of perfect elements for $D^{2,2,2}$ is based on the description of admissible elements. The constructed set H^+ of perfect elements is the union of 64-element distributive lattices $H^+(n)$, and H^+ is the distributive lattice itself. The lattice of perfect elements B^+ obtained by Gelfand and Ponomarev for D^4 can be imbedded into the lattice of perfect elements H^+ , associated with $D^{2,2,2}$.

Dedicated to my wife Rosa

Contents

List of Figures	9
List of Tables	11
Chapter 1. Introduction	13
1.1. Modular and linear lattices, perfect elements	13
1.2. Outline of admissible elements	15
1.2.1. The idea of admissible elements of Gelfand and Ponomarev	15
1.2.2. Reduction of the admissible elements to atomic elements	16
1.2.2.1. Finite classification	16
1.2.2.2. φ_i –homomorphism	16
1.2.2.3. Reduction to atomic elements	16
1.2.2.4. Periodicity	17
1.3. Bibliographic notes: word problems, tame problems	18
1.4. Outline of this work	19
1.5. Atomic polynomials	20
1.6. The elementary maps φ_i of Gelfand-Ponomarev	21
1.7. Admissible sequences and admissible elements	21
1.7.1. Admissible sequences for $D^{2,2,2}$	21
1.7.2. Admissible sequences for D^4	22
1.8. Inclusion of admissible elements in the modular lattice	24
1.9. Examples of the admissible elements	25
1.9.1. The lattice $D^{2,2,2}$	25
1.9.2. The lattice D^4	25
1.10. Cumulative polynomials	26
1.11. Examples of the cumulative polynomials	26
1.12. The sublattice $H^+(n)$ of the perfect elements	27
1.13. The union of sublattices $H^+(n)$	28
1.13.1. Perfect cubes in the free modular lattice D^r	28
1.13.2. Perfect sublattices in the lattice $D^{2,2,2}$	28
Chapter 2. Elementary maps φ_i and atomic, admissible, cumulative polynomials	31
2.1. Construction of the Coxeter functor Φ^+ by means of the lattice operations	31
2.2. The associated representations ν^0, ν^1	31
2.3. The elementary maps φ_i	32
2.4. Basic relations for the maps ψ_i and φ_i	33
2.5. Additivity and multiplicativity of the joint maps ψ_i	35
2.6. Basic properties of atomic elements a_n^{ij} and A_n^{ij}	37
2.7. Action of the maps ψ_i and φ_i on atomic elements	39
2.8. φ_i –homomorphic theorem	41
2.9. The perfectness of $y_i + y_j$	42
2.10. The elementary map φ_i : a way to construct new perfect elements	44

2.11.	List of the admissible sequences	45
2.12.	The theorem on the admissible element classes	46
2.13.	The inclusion theorem and the cumulative polynomials	46
Chapter 3.	Perfect polynomials in $D^{2,2,2}$	49
3.1.	The chains $a_i(n) \subseteq b_i(n) \subseteq c_i(n)$. Perfectness and distributivity of $H^+(n)$.	49
3.2.	The multiplicative form and the cardinality of $H^+(n)$	50
3.3.	The additive form for elements of $H^+(n)$. Considerations mod θ	51
3.4.	Sums and intersections	51
3.5.	Other generators: $p_i(n) \subseteq q_i(n) \subseteq s_i(n)$	53
3.6.	Coupling together $H^+(n)$ and $H^+(n+1)$. Considerations mod θ	54
3.7.	Boolean algebras U_n and V_{n+1}	55
3.8.	The cardinality of $U_n \cup V_{n+1}$	55
3.9.	The Boolean algebra $U_n \cup V_{n+1}$	57
3.10.	The theorem on perfect elements. Considerations mod θ	59
Chapter 4.	Atomic and admissible polynomials in D^4	63
4.1.	Admissible sequences in D^4	63
4.2.	Atomic polynomials and elementary maps in the modular lattice D^4	70
4.3.	Basic relations for the elementary and joint maps in the case of D^4	73
4.4.	Additivity and multiplicativity of the joint maps in the case of D^4	74
4.5.	The action of maps ψ_i and φ_i on the atomic elements in D^4	75
4.6.	The fundamental property of the elementary maps	77
4.7.	The φ_i -homomorphic elements in D^4	77
4.8.	The theorem on the classes of admissible elements in D^4	78
4.8.1.	Basic properties of admissible elements in D^4	78
4.8.2.	Coincidence with the Gelfand-Ponomarev polynomials	80
Appendix A.	Modular and linear lattices	87
A.1.	Distributive lattices and Boolean algebras	87
A.2.	Modular lattices	88
A.3.	Hasse diagrams	89
A.4.	A characterization of distributive and modular lattices	90
A.5.	When the union of two 3D cubes is a 4D cube?	92
A.6.	Representations of the modular lattices	94
A.7.	A relation between representations of graphs and representation of lattices	96
A.8.	The Coxeter functor Φ^+ for $D^{2,2,2}$	98
A.9.	Linear lattices	101
A.10.	Grassmann-Cayley algebras and modular lattices	104
A.10.1.	A Peano space and the bracket	104
A.10.2.	Motivation of the usage the symbol \vee	106
A.10.3.	The meet	106
A.10.4.	Note on Howe-Huang projective invariants of quadruples	107
Appendix B.	A proof of the theorem on admissible classes	109
B.1.	A proof of the theorem on admissible elements in lattice $D^{2,2,2}$	109
B.1.1.	Case $z_\gamma \xrightarrow{\varphi_3} z_{3\gamma}$	109
B.1.2.	Case $z_{3\gamma} \xrightarrow{\varphi_1} z_{(13)\gamma}$	110
B.1.3.	Case $z_\beta \xrightarrow{\varphi_3} z_{3\beta}$	112
B.1.4.	Case $z_{3\beta} \xrightarrow{\varphi_1} z_{(13)\beta}$	113

B.1.5.	Case $z_{(21)^{2k}} \xrightarrow{\varphi_1} z_{1(21)^{2k}}$	114
B.1.6.	Case $z_{1(21)^{2k}} \xrightarrow{\varphi_2} z_{(21)^{2k+1}}$	115
B.1.7.	Case $z_{(21)^{2k+1}} \xrightarrow{\varphi_1} z_{1(21)^{2k+1}}$	116
B.1.8.	Case $z_{1(21)^{2k+1}} \xrightarrow{\varphi_2} z_{(21)^{2k+2}}$	118
B.2.	A proof of the theorem on admissible elements in lattice D^4	119
B.2.1.	Line $F21$, $e_\alpha \xrightarrow{\varphi_3} e_{3\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_3} f_{3\alpha 0}$	119
B.2.2.	Line $F21$, $e_\alpha \xrightarrow{\varphi_4} e_{4\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_4} f_{4\alpha 0}$	120
B.2.3.	Line $F21$, $e_\alpha \xrightarrow{\varphi_1} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_1} f_{1\alpha 0}$	120
B.2.4.	Line $F31$, $e_\alpha \xrightarrow{\varphi_1} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_1} f_{1\alpha 0}$	121
B.2.5.	Line $G11$, $e_\alpha \xrightarrow{\varphi_2} e_{2\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_2} f_{2\alpha 0}$	121
B.2.6.	Line $G21$, $e_\alpha \xrightarrow{\varphi_3} e_{2\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_3} f_{3\alpha 0}$	122
B.2.7.	Line $G21$, $e_\alpha \xrightarrow{\varphi_4} e_{4\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_4} f_{4\alpha 0}$	122
B.2.8.	Line $G21$, $e_\alpha \xrightarrow{\varphi_1} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_1} f_{1\alpha 0}$	123
B.2.9.	Line $H11$, $e_\alpha \xrightarrow{\varphi_2} e_{2\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_2} f_{2\alpha 0}$	124
B.2.10.	Line $H21$, $e_\alpha \xrightarrow{\varphi_1} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_1} f_{1\alpha 0}$	124
B.2.11.	Line $H21$, $e_\alpha \xrightarrow{\varphi_3} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_3} f_{3\alpha 0}$	125
B.2.12.	Line $H21$, $e_\alpha \xrightarrow{\varphi_4} e_{4\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_4} f_{4\alpha 0}$	125
Appendix C.	The inclusion theorem and properties of $H^+(n)$	127
C.1.	Proof of the inclusion theorem	127
C.1.1.	More properties of the atomic elements	127
C.1.2.	The a -form and A -form of admissible elements	128
C.1.3.	Admissible sequences $\alpha 1$ and α . The inclusion theorem.	130
C.1.4.	An S -form of $g_{\alpha 0}$	132
C.2.	Properties of $H^+(n)$	136
C.2.1.	The perfectness of $a_i(1)$	136
C.2.2.	The perfectness of $b_i(1)$	136
C.2.3.	The perfectness of $c_i(1)$	137
C.2.4.	The perfectness of $a_i(n), b_i(n), c_i(n)$	137
C.2.5.	The distributivity of $H^+(n)$	137
C.2.6.	The cardinality of $H^+(n)$	138
C.2.7.	A relation between the upper and the lower generators of $H^+(n)$	139
Conjectures		145
Bibliography		147

List of Figures

1.1	Modular, Arguesian and Linear Lattices	14
1.2	Periodicity of admissible elements	17
1.3	How maps φ_i act on the admissible sequences	23
1.4	Action of maps φ_i on the admissible sequences for D^4	24
1.5	The 64-element distributive lattice $H^+(n)$ of perfect elements	28
1.6	Comparison between perfect sublattices in D^4 and $D^{2,2,2}$	29
1.7	The 16-element Boolean algebra $U_n \cup V_{n+1}$	30
3.1	The Boolean algebras U_n and V_{n+1}	56
4.1	Slices of admissible sequences for D^4 , $l = 3$ and $l = 4$	69
4.2	Slices of admissible sequences for D^4 , $l = 4$ and $l = 5$	69
A.1	Hasse diagrams of D_{18} and M_{28}	89
A.2	Two Hasse diagrams: the diamond M_3 and the pentagon N_5	90
A.3	The 4-dimensional cube $D \cup C$	93
A.4	Maps f_1 and g_1 . $K = \ker f_1$, $G = \text{Im} g_1$	97
A.5	The reflection functor $F_{x_0}^+$	100
A.6	The reflection functor $F_{y_3}^+ F_{y_2}^+ F_{y_1}^+ F_{x_0}^+$	100
A.7	The Coxeter functor Φ^+	101

List of Tables

2.1	The basic properties of atomic elements a_n^{ij} and A_n^{ij} in $D^{2,2,2}$	37
2.2	For $D^{2,2,2}$: the admissible sequences starting at $i_1 = 1$	45
2.3	For $D^{2,2,2}$: the admissible elements starting at $i_1 = 1$	47
3.1	The multiplicative and additive forms of $H^+(n)$	54
3.2	The 8-element Boolean algebra U_n	55
3.3	The 8-element Boolean algebra V_{n+1}	56
4.1	Admissible sequences for the modular lattice D^4	67
4.2	Comparison of notions in $D^{2,2,2}$ and D^4	72
4.3	Admissible polynomials in the modular lattice D^4	85
A.1	Preprojective representations	99
C.1	The admissible elements, a -form and A -form	129
C.2	$\alpha 1$, α , $e_{\alpha 1}$ and $g_{\alpha 0}$	131
C.3	The S -forms of $g_{\alpha 0}$	132
C.4	The relation $e_{\alpha 1} = g_{\alpha 0}Z$	135
C.5	The characteristic function on the $H^+(0)$	142
C.6	The characteristic function on the $H^+(1)$. The generators $a_i(1), b_i(1), c_i(1)$	143
C.7	The characteristic function on the $H^+(1)$. The elements 10–64	144

CHAPTER 1

Introduction

... These developments, and several others that I have not mentioned, are a belated validation of Garrett Birkhoff's vision, which we learned in three editions of his Lattice Theory, and they betoken Professor Gelfand's oft-repeated prediction that lattice theory will play a leading role in the mathematics of the twenty-first century.

G-C Rota, [R97, p.1445].

1.1. Modular and linear lattices, perfect elements

The objects considered in this work have diverse applications from quantum logic of quantum mechanics (see Remark 1.1.2) to representation problems of posets and quivers of *tame* type (see §1.3). The potential of these applications is, however, still dormant to a considerable extent, even the basic notions are not widely known. In Appendix A we give definitions and examples of some notions including modular and linear lattices, representations of modular lattices and perfect elements. In this section we briefly remind a number of notions.

A *lattice* is a set L with two commutative and associative operations: a *sum* and an *intersection*. If $a, b \in L$, then we denote the intersection by ab and the sum by $a + b$. Both operations are *idempotent*

$$aa = a, \quad a + a = a$$

and satisfy the *absorption law*

$$a(a + b) = a, \quad a + ab = a.$$

On the lattice L , an *inclusion* \subseteq is defined:

$$a \subseteq b \iff ab = a \text{ or } a + b = b,$$

see §A.1.

A lattice is said to be *modular* if, for every $b, a, c \in L$,

$$a \subseteq b \implies b(a + c) = a + bc, \tag{1.1}$$

see §A.2.

Good examples of modular lattices, bringing us some intuition, are the lattices of subspaces of a given vector space, normal subgroups of a given group, ideals of a given ring.

REMARK 1.1.1. Actually, the above examples of modular lattices are more than modular. In his famous work [Hai85] M. Haiman baptized these lattices the *linear lattices*. The linear lattices are also known as lattices of *commuting equivalence relations*, see §A.9. The term *linear* was suggested by G.-C.Rota, see [Hai85, p.1], [FMR96].

The linear lattices were studied by B. Jónsson in [Jo53], [Jo54], where these lattices were named lattices with *representations of type 1*. In [Jo53] B. Jónsson also introduced

the *Arguesian law* also studied by M. P. Schützenberger [Sch45]. The Arguesian law, see [MY2000, p.3] and [Hai85], is a lattice-theoretic form of the Desargues theorem of projective geometry, see [Gr98, Ch.4]. Lattices satisfying the Arguesian law are called *Arguesian lattices*. For more details about linear and Arguesian lattices, see §A.9. On Fig. 1.1 the hierarchical relations between three classes of lattices: modular, Arguesian, and linear are shown.




FIGURE 1.1. Modular, Arguesian and Linear Lattices

We denote by D^r the free modular lattice with r generators

$$D^r = \{e_1, \dots, e_r\}, \quad (1.2)$$

see [GP74], [Fr80]. The word “free” means that no relation exists between generators $e_i, i = 1, \dots, r$.

Denote by $D^{2,2,2}$ the modular lattice with 6 generators $\{x_1, y_1, x_2, y_2, x_3, y_3\}$ satisfying the following relations:

$$D^{2,2,2} = \{x_1 \subseteq y_1, \quad x_2 \subseteq y_2, \quad x_3 \subseteq y_3\}. \quad (1.3)$$

Symbol D in (1.2) and (1.3) is used in honour of R. Dedekind who derived the main properties of the modular lattices, [De1897]. The *modular law* (1.1) is also known as *Dedekind’s law* and the modular lattices are sometimes called the *Dedekind lattices*, see §A.2.

Let L be a modular lattice, X a finite dimensional vector space, $\mathcal{L}(X)$ the modular lattice of linear subspaces in L . A morphism $\rho : L \rightarrow \mathcal{L}(X)$ is called a *linear representation* of L in the space X , see §A.6. By §A.9 (Proposition A.9.5) the modular lattice $\mathcal{L}(X)$ is a linear lattice. Thus, we consider representations of the modular lattice L in the linear lattices $\mathcal{L}(X)$.

Following [GP74] we introduce now the notions of *perfect elements* and *linear equivalence relation*. The element $a \in L$ is said to be *perfect* if $\rho(a) = 0$ or $\rho(a) = X$ for each

indecomposable representation ρ of the modular lattice L , where X is the representation space of ρ , see §A.6.

Two elements $a, b \in L$ are called *linearly equivalent* if $\rho(a) = \rho(b)$ for all indecomposable representations $\rho : L \rightarrow \mathcal{L}(X)$ and we write

$$a \equiv b \pmod{\theta}. \quad (1.4)$$

The relation θ is called the *linear equivalence relation*, see §A.6.

REMARK 1.1.2. Manin [Man79, p.98] pointed out at a connection between *perfect elements* and *quantum tautologies* in the quantum logic introduced by Birkhoff and von Neumann in [BvN36]. In quantum logic, a *perfect element* is called a *modular question* [Man79]. In [KS67], Kochen and Specker considered related topics in the foundations of the quantum mechanics and the quantum logic.

G.-C.Rota writes in [R97, p.1444] that Haiman's proof theory [Hai85] for linear lattices brings to fruition the program that was set forth in [BvN36]: "The authors¹ did not know that the modular lattices of quantum mechanics are linear lattices. In light of Haiman's proof theory, we may now confidently assert that Birkhoff and von Neumann's logic of quantum mechanics is indeed the long-awaited new "logic"..."

Finberg, Mainetti and Rota in [FMR96] gave a simplified presentation of the work of Haiman [Hai85].

For more details about *linear lattices*, see §A.9.

1.2. Outline of admissible elements

First, in this section, we outline the idea of admissible elements due to Gelfand and Ponomarev [GP74]. In [GP74], [GP76], these elements were used to obtain the distributive sublattice of perfect polynomials. We think today, that apart from being helpful in construction of perfect elements, the admissible elements are interesting in themselves.

Further, we outline properties of admissible elements obtained in this work: *finite classification*, φ -homomorphism, *reduction to atomic elements* and *periodicity*, see 1.2.2.

1.2.1. The idea of admissible elements of Gelfand and Ponomarev. Here, we describe, omitting some details, the idea of constructing the elementary maps and admissible elements of Gelfand-Ponomarev. As we will see below, the admissible elements *grow*, in a sense, from generators of the lattice or from the lattice's unity.

For every representation ρ of the modular lattice L , a new representation $\tilde{\rho}$ is uniquely constructed by means of the Coxeter functor Φ^+ , introduced by Bernstein, Gelfand and Ponomarev in [BGP73] to study representations of graphs, see §A.8. Let X (resp. \tilde{X}) be a space of the representation ρ (resp. $\tilde{\rho}$), and $\mathcal{L}(X)$ (resp. $\mathcal{L}(\tilde{X})$) be the set of subspaces of X (resp. \tilde{X}):

$$\rho : L \rightarrow \mathcal{L}(X), \quad \tilde{\rho} : L \rightarrow \mathcal{L}(\tilde{X}). \quad (1.5)$$

Construction of the elementary map φ . Let there exist a map φ mapping every subspace $A \in \mathcal{L}(\tilde{X})$ to some subspace $B \in \mathcal{L}(X)$:

$$\begin{aligned} \varphi : \mathcal{L}(\tilde{X}) &\rightarrow \mathcal{L}(X), \\ \varphi A &= B, \quad \text{or} \quad A \xrightarrow{\varphi} B. \end{aligned} \quad (1.6)$$

It turns out that for many elements $a \in L$ there exists an element $b \in L$ (at least in $L = D^4$ and $D^{2,2,2}$) such that

$$\varphi \tilde{\rho}(a) = \rho(b) \quad (1.7)$$

¹G. Birkhoff G. and J. von Neuman [BvN36].

for every pair of representations $(\rho, \tilde{\rho})$, where $\tilde{\rho} = \Phi^+ \rho$. The map φ in (1.7) is constructed in such a way, that φ do not depend on ρ . Thus, we can write

$$a \xrightarrow{\varphi} b. \quad (1.8)$$

In particular, eq. (1.7) and (1.8) are true for admissible elements, whose definition will be given later, see §1.7. Eq. (1.7) is the main property characterizing the admissible elements.

For D^4 , Gelfand and Ponomarev constructed 4 maps of form (1.6): $\varphi_1, \varphi_2, \varphi_3, \varphi_4$ and called them *elementary maps*. In this work, for $D^{2,2,2}$, we construct 3 maps φ_i , one per chain $x_i \subseteq y_i$, see (§1.6 and Ch.2). Given a sequence of relations

$$a_{i_1} \xrightarrow{\varphi_{i_1}} a_{i_2} \xrightarrow{\varphi_{i_2}} a_{i_3} \xrightarrow{\varphi_{i_3}} \dots \xrightarrow{\varphi_{i_{n-2}}} a_{i_{n-1}} \xrightarrow{\varphi_{i_{n-1}}} a_n, \quad (1.9)$$

We say that elements $a_2, a_3, \dots, a_{i_{n-2}}, a_{i_{n-1}}, a_n$ *grow* from the element a_{i_1} . By abuse of language, we say that elements growing from generators e_i or unity I are *admissible elements*, and the corresponding sequence of indices $\{i_n i_{n-1} i_{n-2} \dots i_2 i_1\}$ is said to be the *admissible sequence*. For details in the cases $D^{2,2,2}$ (resp. D^4) see §1.7.1, (resp. §1.7.2).

1.2.2. Reduction of the admissible elements to atomic elements. Here, we briefly give steps of creation of the admissible elements in this work.

1.2.2.1. *Finite classification.* The admissible elements (and admissible sequences) can be reduced to a finite number of classes, see §1.7. Admissible sequences for $D^{2,2,2}$ (resp. D^4) are depicted on Fig. 1.3 (resp. Fig. 1.4). The key property of admissible sequences allowing to do this classification is the following relation between maps φ_i .

For $D^{2,2,2}$ (see Proposition 2.10.2):

$$iji = iki, \\ \varphi_i \varphi_j \varphi_i = \varphi_i \varphi_k \varphi_i, \text{ where } \{i, j, k\} = \{1, 2, 3\}.$$

For D^4 (see Proposition 4.6.1):

$$ikj = ilj, \\ \varphi_i \varphi_k \varphi_j = \varphi_i \varphi_l \varphi_j, \text{ where } \{i, j, k, l\} = \{1, 2, 3, 4\}.$$

1.2.2.2. φ_i -homomorphism. In a sense, the elementary maps φ_i are homomorphic with respect to admissible elements. More exactly, introduce a notion of φ_i -homomorphic elements. Let

$$a \xrightarrow{\varphi_i} \tilde{a} \quad \text{and} \quad p \xrightarrow{\varphi_i} \tilde{p}. \quad (1.10)$$

The element $a \in L$ is said to be φ_i -homomorphic, if

$$ap \xrightarrow{\varphi_i} \tilde{a}\tilde{p} \quad (1.11)$$

for all $p \in L$, see §1.6.

1.2.2.3. *Reduction to atomic elements.* We permanently apply the following mechanism of creation of admissible elements from more simple elements. Let φ_i be any elementary map, and

$$a \xrightarrow{\varphi_i} \tilde{a}, \quad b \xrightarrow{\varphi_i} \tilde{b}, \quad c \xrightarrow{\varphi_i} \tilde{c}, \quad p \xrightarrow{\varphi_i} \tilde{p}. \quad (1.12)$$

Suppose $abcp$ is any admissible element and a, b, c are φ_i -homomorphic. By means of relation (1.11) we construct new admissible element

$$\tilde{a}\tilde{b}\tilde{c}\tilde{p},$$

The elements a, b, c are called *atomic*, see §1.5.

1.2.2.4. *Periodicity.* Admissible sequences obtained by reducing in heading (1) possess some *periodicity*. The corresponding admissible elements are also periodic. On Fig. 1.2 we see an example of admissible elements for $D^{2,2,2}$:

$$e_{(213)^p(21)}, \quad e_{3(213)^p(21)}, \quad e_{13(213)^p(21)}.$$

Three vertical lines on Fig. 1.2 correspond to three series of inclusions:

$$\begin{aligned} \cdots &\subseteq e_{(213)^{p+1}(21)} \subseteq e_{(213)^p(21)} \subseteq \cdots \subseteq e_{(213)(21)} \subseteq e_{(21)}, \\ \cdots &\subseteq e_{3(213)^{p+1}(21)} \subseteq e_{3(213)^p(21)} \subseteq \cdots \subseteq e_{3(213)(21)} \subseteq e_{3(21)}, \\ \cdots &\subseteq e_{13(213)^{p+1}(21)} \subseteq e_{13(213)^p(21)} \subseteq \cdots \subseteq e_{13(213)(21)} \subseteq e_{13(21)} \subseteq e_1, \end{aligned} \quad (1.13)$$

Relation (1.13) is easily obtained from Table C.2. From the same table we can see periodicity of indices of atomic elements entering in the decomposition of admissible elements. The triangular helix on Fig. 1.2 corresponds to relations

$$\begin{aligned} e_{(213)^p(21)} &\xrightarrow{\varphi_3} e_{3(213)^p(21)}, \\ e_{3(213)^p(21)} &\xrightarrow{\varphi_1} e_{13(213)^p(21)}, \\ e_{13(213)^p(21)} &\xrightarrow{\varphi_2} e_{(213)^{p+1}(21)}. \end{aligned} \quad (1.14)$$

Relation (1.14) is a particular case of the main theorem on admissible elements (Theorem 2.12.1).




FIGURE 1.2. Periodicity of admissible elements

1.3. Bibliographic notes: word problems, tame problems

Admissible elements for D^r in [GP74], [GP76] are built recurrently in the length of multi-indices named *admissible sequences*. In this work we suggest a direct method for creating admissible elements. For $D^{2,2,2}$ (resp. D^4), the admissible sequences and admissible elements form 14 classes (resp. 11 classes) and possess some periodicity properties, see §1.7.1, Tables 2.2, 2.3 (resp. §1.7.2, Tables 4.1, 4.3).

For the definition of admissible polynomials due to Gelfand and Ponomarev and examples obtained from this definition, see §4.8.2. Dlab and Ringel in [DR80] and Cylke in [Cy182] proved that perfect elements¹ constructed by Gelfand and Ponomarev coincide with all possible perfect elements modulo linear equivalence, see §A.6.

The study of D^r , where $r \geq 4$ might be a rather hard task. We just point to two works on the word problem for the free modular lattices. Herrmann proved in [H83] that the word problem in D^4 is unsolvable, and Freese in [Fr80] proved that the word problem in D^r is unsolvable for $r \geq 5$. Furthermore, Herrmann proved that *the modular lattice freely generated by a partially ordered set P containing one of lattices*


$$1 + 1 + 1 + 1, \quad 2 + 2 + 2, \quad 1 + 2 + 3,$$

has an unsolvable word problem, [H83, p.526]. Thus, the modular lattice $D^{2,2,2}$, which is of interest to us, has also an unsolvable word problem. We mention also the following fact proved by Wille [W73]: *a modular lattice is finite if and only if it does not contain*

$$1 + 1 + 1 + 1 \quad \text{and} \quad 1 + 2 + 2.$$


Gelfand conjectured that the free linear lattice in four generators has the solvable word problem. If this conjecture is true, it would distinguish the class of linear lattices from that of modular lattices, see [KY2003, p.4].

The problem of classifying all indecomposable representations of D^4 (*indecomposable quadruples*) of subspaces of arbitrary finite dimensional spaces is equivalent to that of determining all indecomposable representations of the graph



which corresponds to the extended Dynkin diagram \tilde{D}_4 . This is a representation problem of *tame* type solved by Nazarova [Naz67] and by Gelfand and Ponomarev [GP70] by different means.

The problem of classifying all indecomposable representations of $D^{2,2,2}$ is equivalent to that of determining all indecomposable representations of the graph



¹In the context of this article *perfect element* is the same as *perfect polynomial*, similarly *admissible element* is the same that *admissible polynomial*

which corresponds to the extended Dynkin diagram \tilde{E}_6 . This is also a representation problem of *tame* type solved by Nazarova [Naz73] and by Dlab and Ringel [DR74], [DR76].

Ponomarev considered in detail perfect elements for lattices related to the Dynkin diagram A_n [Pon90]. For every Dynkin diagram, he described *perfect elements* by means of so-called *hereditary subsets* in the corresponding Auslander-Reiten diagram, [Pon90, Theorem 1.6].

Cylke [Cyl93] has constructed a sublattice $B^+(S) \cup B^-(S)$ in the modular lattice $L(S)$ freely generated by a poset S of finite type, or tame type of finite growth. The sublattice $B^+(S) \cup B^-(S)$ is a natural generalization of the sublattices $B^+ \cup B^-$ of perfect elements constructed by Gelfand-Ponomarev [GP74]. The sublattice $B^+(S) \cup B^-(S)$ is built by means of *poset differentiation*. Nazarova and Roiter used this technique to construct indecomposable representations of posets and quivers [NR72], [Naz73], [Ro85], [GR92].

1.4. Outline of this work

In this work we construct *admissible* and *perfect* elements for the modular lattice $D^{2,2,2}$, associated with the extended Dynkin diagram \tilde{E}_6 , see Ch.2 – Ch.3, and *admissible* elements for the modular lattice D^4 , associated with the extended Dynkin diagram \tilde{D}_4 , see Ch.4. For D^4 , the admissible elements constructed in this work and admissible elements constructed by Gelfand and Ponomarev [GP74] coincide at least modulo linear equivalence, see §4.8.2.

The diagrams \tilde{E}_6 and \tilde{D}_4 are *tame quivers*, i.e., their representations can be classified [Naz73], [DR76].

During the construction of perfect elements in the lattice $D^{2,2,2}$ various lattice polynomials in $D^{2,2,2}$ naturally appear: *atomic*, φ -*homomorphic*, *admissible*, *cumulative* and *perfect*. *Atomic polynomials* form a basis for constructing *admissible elements*. The *cumulative polynomials* are constructed by means of admissible elements, and *perfect* elements are constructed from the *cumulative* elements:

$$\mathbf{Atomic} \implies \mathbf{Admissible} \implies \mathbf{Cumulative} \implies \mathbf{Perfect}$$

In the sequel to this paper we intend to show that admissible sequences play an important role in the study of the preprojective algebras, [GP79], [DR79], [Rin96].

The results of the work were partially published in [St89].

In Ch.2 we introduce *atomic*, *admissible* and *cumulative* lattice polynomials and consider basic properties of these polynomials. Table 2.3 gives a full list of admissible polynomials for $D^{2,2,2}$, see also §1.7.

In Ch.3 we obtain sublattices of perfect elements $H^+(n)$ in $D^{2,2,2}$ and show that the lattice

$$H^+ = \bigcup_{n=0}^{\infty} H^+(n)$$

is a distributive lattice of perfect elements. The union H^+ has an interesting architecture and contains distributive lattice B^+ of perfect elements of D^4 , see Fig. 1.6, §1.12.

In Ch.4, by repeating the technique developed for $D^{2,2,2}$ in Ch.2, we construct admissible elements in D^4 .

For the convenience of the reader, we recall in Appendix A some properties of the modular lattices, their representations, and give a brief introduction to linear lattices.

Proofs of a number of theorems and verification of several properties are moved to Appendices B and C.

In this work we consider the diagram \tilde{E}_6 only with central orientation in which all arrows are directed to the vertex x_0 . The modular lattice $D^{2,2,2}$ is generated by partially ordered set $2 + 2 + 2$. The generators $x_i, y_i, (i = 1, 2, 3)$ satisfy the inclusions $x_i \subseteq y_i$. For convenience, we include unity I in the lattice $D^{2,2,2}$:

$$\begin{array}{ccccccc}
 x_1 & \rightarrow & y_1 & \rightarrow & x_0 & \leftarrow & y_2 \leftarrow x_2 \\
 & & \tilde{E}_6 & & \uparrow & & \\
 & & & & y_3 & & \\
 & & & & \uparrow & & \\
 & & & & x_3 & & \\
 & & & & & & D^{2,2,2} \\
 & & & & & & \begin{array}{c} I \\ \cup \mid \\ y_3 \\ \cup \mid \\ x_3 \end{array}
 \end{array} \quad (1.15)$$

Following Bernstein-Gelfand-Ponomarev [BGP73] and Gelfand-Ponomarev [GP74] we use the Coxeter functor Φ^+ , and the elementary linear maps φ_i .

1.5. Atomic polynomials

Atomic lattice polynomials have a simple lattice definition. They are called *atomic* because *admissible*, *cumulative* and *perfect* lattice polynomials (as well as *invariant* in [St92]) are defined by means of these polynomials.

For $D^{2,2,2}$, the definition of $a_n^{ij}, A_n^{ij}, n \in \{0, 1, 2, \dots\}$, is cyclic through the indices i, j, k , where the triple $\{i, j, k\}$ is a permutation of $\{1, 2, 3\}$. We set

$$\begin{aligned}
 a_n^{ij} &= \begin{cases} a_n^{ij} = I & \text{for } n = 0, \\ a_n^{ij} = x_i + y_j a_{n-1}^{jk} & \text{for } n \geq 1, \end{cases} \\
 A_n^{ij} &= \begin{cases} A_n^{ij} = I & \text{for } n = 0, \\ A_n^{ij} = y_i + x_j A_{n-1}^{ki} & \text{for } n \geq 1, \end{cases}
 \end{aligned} \quad (1.16)$$

The elements a_n^{ij}, A_n^{ij} are said to be *atomic elements* in $D^{2,2,2}$:

Examples of atomic elements in $D^{2,2,2}$.

$$\begin{array}{ll}
 a_1^{12} = x_1 + y_2, & A_1^{12} = y_1 + x_2, \\
 a_2^{12} = x_1 + y_2(x_2 + y_3), & A_2^{12} = y_1 + x_2(y_3 + x_1), \\
 a_3^{12} = x_1 + y_2(x_2 + y_3(x_3 + y_1)), & A_3^{12} = y_1 + x_2(y_3 + x_1(y_2 + x_3)), \\
 a_4^{12} = x_1 + y_2(x_2 + y_3(y_3 + y_1(x_1 + y_2))), & A_4^{12} = y_1 + x_2(y_3 + x_1(y_2 + x_3(y_1 + x_2))), \\
 \dots & \dots
 \end{array}$$

For D^4 , we define *atomic* lattice polynomials a_n^{ij} , where $i, j \in \{1, 2, 3, 4\}$, $n \in \mathbb{Z}_+$, as follows

$$a_n^{ij} = \begin{cases} a_n^{ij} = I & \text{for } n = 0, \\ a_n^{ij} = e_i + e_j a_{n-1}^{kl} = e_i + e_j a_{n-1}^{lk} & \text{for } n \geq 1, \end{cases} \quad (1.17)$$

where $\{i, j, k, l\}$ is the permutation of the quadruple $\{1, 2, 3, 4\}$, and $\{e_1, e_2, e_3, e_4\}$ are generators in D^4 , see §4.2.

Examples of the atomic elements in D^4 .

$$\begin{aligned}
 a_1^{12} &= e_1 + e_2, \\
 a_2^{12} &= e_1 + e_2(e_3 + e_4), \\
 a_3^{12} &= e_1 + e_2(e_3 + e_4(e_1 + e_2)), \\
 a_4^{12} &= e_1 + e_2(e_2 + e_3(e_1 + e_2(e_3 + e_4))), \\
 &\dots
 \end{aligned}$$

1.6. The elementary maps φ_i of Gelfand-Ponomarev

Throughout the work the maps φ_i play a central role. For exact definition of the maps φ_i , see §2.3. Gelfand and Ponomarev introduced the maps φ_i in [GP74, p.27] for the modular lattices D^r and called them *elementary maps*.

REMARK 1.6.1 (An explanation of indices). Let X_0 be the representation space of the representation ρ , X_0^n be the representation space of the representation $\rho^n = (\Phi^+)^n \rho$, where $n \geq 1$ and Φ^+ is the Coxeter transformation. The lower index 0 means that X_0^n is the image of the generator x_0 of the lattice $D^{2,2,2}$ under the representation $(\Phi^+)^n \rho$, see (1.15):

$$\begin{aligned} (\Phi^+)^n \rho(x_0) &= X_0^n \text{ for } n \geq 1, \text{ where } \rho(x_0) = X_0. \\ (\Phi^+)^n \rho(y_i) &= Y_i^n \text{ for } n \geq 1, \text{ where } \rho(y_i) = Y_i, (i = 1, 2, 3). \\ (\Phi^+)^n \rho(x_i) &= X_i^n \text{ for } n \geq 1, \text{ where } \rho(x_i) = X_i, (i = 1, 2, 3). \end{aligned} \quad (1.18)$$

The indices $i = 1, 2, 3$ of φ_i mean that φ_i operates to the subspace $Y_i \subseteq X_0$, where Y_i is the image of the generator y_i under the representation $\Phi^+ \rho$, see (1.19):

$$\begin{array}{ccccccc} X_1 & \subseteq & Y_1 & \subseteq & X_0 & \supseteq & Y_2 \supseteq X_2 \\ & & \cup & & \cup & & \\ & & \rho & & Y_3 & & \\ & & \cup & & \cup & & \\ & & & & X_3 & & \end{array} \quad \begin{array}{ccccccc} X_1^1 & \subseteq & Y_1^1 & \subseteq & X_0^1 & \supseteq & Y_2^1 \supseteq X_2^1 \\ & & \cup & & \cup & & \\ & & \Phi^+ \rho & & Y_3^1 & & \\ & & \cup & & \cup & & \\ & & & & X_3^1 & & \end{array} \quad (1.19)$$

We introduce now φ_i -homomorphic polynomials playing an important role in our further considerations.

An element $a \subseteq D^{2,2,2}$ is said to be φ_i -homomorphic, if

$$\varphi_i \Phi^+ \rho(ap) = \varphi_i \Phi^+ \rho(a) \varphi_i \Phi^+ \rho(p) \text{ for all } p \subseteq D^{2,2,2}. \quad (1.20)$$

An element $a \subseteq D^{2,2,2}$ is said to be (φ_i, y_k) -homomorphic, if

$$\varphi_i \Phi^+ \rho(ap) = \varphi_i \Phi^+ \rho(y_k a) \varphi_i \Phi^+ \rho(p) \text{ for all } p \subseteq y_k. \quad (1.21)$$

Here $\Phi^+ \rho(a)$, $\Phi^+ \rho(p)$ are the images of $a, p \in D^{2,2,2}$ under representation $\Phi^+ \rho$ (see Remark 1.6.1). For the detailed definition of φ_i and related notions, see §2.2–§2.3.

The notion of φ_i -homomorphic elements in D^4 is similarly introduced in §4.7.

All *atomic* polynomials are φ_i -homomorphic. More exactly, for $D^{2,2,2}$, we have (see Theorem 2.8.1, §2.8):

- 1) The polynomials a_n^{ij} are φ_i - and φ_j -homomorphic.
- 2) The polynomials A_n^{ij} are φ_i -homomorphic.
- 3) The polynomials A_n^{ij} are (φ_j, y_k) -homomorphic.

For D^4 , we have (Theorem 4.7.1, §4.7):

- 1) The polynomials a_n^{ij} are φ_i -homomorphic.
- 2) The polynomials a_n^{ij} (φ_j, e_k) -homomorphic.

1.7. Admissible sequences and admissible elements

1.7.1. Admissible sequences for $D^{2,2,2}$. The *admissible* lattice polynomials are introduced in §2.12. They appear when we use different maps φ_i , where $i = 1, 2, 3$. The admissible polynomials can be indexed by means of a finite number of types of index sequences, called

admissible sequences. Consider a finite sequence of indices $s = i_n \dots i_1$, where $i_p \in \{1, 2, 3\}$. The sequence s is said to be *admissible* if

- (a) Adjacent indices are distinct ($i_p \neq i_{p+1}$).
- (b) In each subsequence iji , we can replace index j by k . In other words:
 $\dots iji \dots = \dots iki \dots$, where all indices i, j, k are distinct.

The admissible sequence with $i_1 = 1$ for $D^{2,2,2}$ may be transformed to one of the seven types (Proposition 2.11.1, see §2.11),

$$\begin{aligned}
 & 1) (213)^m(21)^n, \quad 2) 3(213)^m(21)^n, \quad 3) 13(213)^m(21)^n, \\
 & 4) (312)^m(31)^n, \quad 5) 2(312)^m(31)^n, \quad 6) 12(312)^m(31)^n, \\
 & 7) 1(21)^n = 1(31)^n.
 \end{aligned} \tag{1.22}$$

similarly for $i_1 = 2, 3$, for example, for $i_1 = 2$, the admissible sequences are:

$$\begin{aligned}
 & 1) (123)^m(12)^n, \quad 2) 3(123)^m(12)^n, \quad 3) 23(123)^m(12)^n, \\
 & 4) (321)^m(32)^n, \quad 5) 1(321)^m(32)^n, \quad 6) 21(321)^m(32)^n, \\
 & 7) 2(12)^n = 2(32)^n.
 \end{aligned} \tag{1.23}$$

A description of all *admissible sequences* for $D^{2,2,2}$ is given in Table 2.2, and a description of all *admissible elements* is given in Table 2.3. For every

$$z_\alpha = e_\alpha, f_\alpha, g_{\alpha 0}$$

from Table 2.3, where α is an admissible sequence, the following statement takes place:


Let α be an admissible sequence $i_n \dots i_1$ and $i \neq i_n$. Then $i\alpha$ is also an admissible sequence and

$$\varphi_i \Phi^+ \rho(z_\alpha) = \rho(z_{i\alpha}) \text{ for each representation } \rho$$

(Theorem 2.12.1, §2.12; see Fig. 1.3).

1.7.2. Admissible sequences for D^4 . The admissible sequence for D^4 is defined as follows. Consider a finite sequence of indices $s = i_n \dots i_1$, where $i_p \in \{1, 2, 3\}$. The sequence s is said to be *admissible* if

- (a) Adjacent indices are distinct ($i_p \neq i_{p+1}$).
- (b) In each subsequence ijl , we can replace index j by k . In other words:
 $\dots ijl \dots = \dots ikl \dots$, where all indices i, j, k, l are distinct.

FIGURE 1.3. How maps φ_i act on the admissible sequences

Any admissible sequence with $i = 1$ for D^4 may be transformed to one the next 11 types (Proposition 4.1.2, Table 4.1):

- 1) $(21)^t(41)^r(31)^s = (21)^t(31)^s(41)^r$,
- 2) $(31)^t(41)^r(21)^s = (31)^t(21)^s(41)^r$,
- 3) $(41)^t(31)^r(21)^s = (41)^t(21)^s(31)^r$,
- 4) $1(41)^t(31)^r(21)^s = 1(31)^t(41)^s(21)^r = 1(21)^t(31)^s(41)^r$,
- 5) $2(41)^r(31)^s(21)^t = 2(31)^{s+1}(41)^{r-1}(21)^t$,
- 6) $3(41)^r(21)^s(31)^t = 3(21)^{s+1}(41)^{r-1}(31)^t$,
- 7) $4(21)^r(31)^s(41)^t = 3(31)^{s+1}(21)^{r-1}(41)^t$,
- 8) $(14)^r(31)^s(21)^t = (14)^r(21)^{t+1}(31)^{s-1} = (13)^s(41)^r(21)^t =$
 $(13)^s(21)^{t+1}(41)^{r-1} = (12)^{t+1}(41)^r(31)^{s-1} = (12)^{t+1}(31)^{s-1}(41)^r$,
- 9) $2(14)^r(31)^s(21)^t$,
- 10) $3(14)^r(31)^s(21)^t$,
- 11) $4(14)^r(31)^s(21)^t$.

Let α be an admissible sequence $i_n \dots i_1$ and $i \neq i_n$, where $i_s \in \{1, 2, 3, 4\}$. Let $z_\alpha = e_\alpha, f_{\alpha 0}$ be admissible elements for D^4 from Table 4.3. Then $i\alpha$ is also an admissible sequence and

$$\varphi_i \Phi^+ \rho(z_\alpha) = \rho(z_{i\alpha}) \text{ for each representation } \rho, \quad (1.25)$$

see Theorem 4.8.1, Fig. 1.4. The admissible elements for D^4 form a finite family and are directly constructed, see Table 4.3.

Admissible polynomials given by Table 4.3 coincide $\bmod \theta^1$ with polynomials constructed by Gelfand and Ponomarev. For small lengths of the admissible sequences, we prove that these polynomials coincide without restriction $\bmod \theta$, see Propositions 4.8.3, 4.8.4, 4.8.5 from §4.8.2.




FIGURE 1.4. Action of maps φ_i on the admissible sequences for D^4

REMARK 1.7.1. In §A.10 we mention a connection between some Howe-Huang projective invariants of quadruples [HH96], the Grassmann-Cayley algebra [BBR85] and *admissible elements* in D^4 .

1.8. Inclusion of admissible elements in the modular lattice

One can easily check, with definitions from Table 2.3, that

$$f_\alpha \subseteq e_\alpha, \quad g_{\alpha 0} \subseteq e_\alpha. \quad (1.26)$$

It is more difficult to prove the following theorem (see Theorem 2.13.1, §2.13, Appendix C):

Theorem. *For every admissible sequence α and αi , where $i = 1, 2, 3$ from Table 2.2, we have*

$$e_{\alpha i} \subseteq g_{\alpha 0}, \quad i = 1, 2, 3.$$

A proof of the theorem is given in Appendix C. This proof uses case by case consideration of *admissible sequences* α from Table 2.3.

¹For definition of $\bmod \theta$, see (A.34) from §A.6

1.9. Examples of the admissible elements

1.9.1. The lattice $D^{2,2,2}$. Let n be the length of an admissible sequence α , let k and p be indices defined by Table 2.3.

For $n = 1$: $\alpha = 1$, (Table 2.3, Line 7, $k = 0$). We have

$$e_1 = y_1, \quad f_1 = x_1 \subseteq e_1, \quad g_{10} = y_1(y_2 + y_3).$$

For $n = 2$: $\alpha = 21$, (Table 2.3, Line 9, $k = 0, p = 0$). Here,

$$e_{21} = y_2 a_1^{13} = y_2(x_1 + y_3), \quad e_{12} = y_1(x_2 + y_3) \subseteq g_{10}, \quad (\text{permutation } (21) \rightarrow (12)).$$

$$f_{21} = y_2 y_3 \subseteq e_{21}, \quad g_{210} = e_{21}(x_2 + A_1^{13}) = y_2(x_1 + y_3)(x_2 + x_3 + y_1).$$

For $n = 3$: $\alpha = 321$, (Table 2.3, Line 10, $k = 0, p = 0$). In this case:

$$e_{321} = y_3 A_1^{12} A_1^{23} = y_3(y_1 + x_2)(y_2 + x_3),$$

$$e_{213} = y_2(y_3 + x_1)(x_1 + x_2) \subseteq g_{210}, (\text{permutation } (321) \rightarrow (213)),$$

$$f_{321} = x_3 A_1^{12} = x_3(y_1 + x_2) \subseteq e_{321},$$

$$g_{3210} = e_{321}(y_1 y_3 + a_1^{12}) = y_3(y_1 + x_2)(y_2 + x_3)(x_1 + y_2 + y_1 y_3).$$

For $n = 4$: $\alpha = 1321$, (Table 2.3, Line 11, $k = 0, p = 0$). Here, we have

$$e_{1321} = y_1 a_1^{13} a_2^{32} = y_1(x_1 + y_3)(x_3 + y_2(x_2 + y_1)),$$

$$e_{3213} = y_3(x_3 + y_2)(x_2 + y_1(x_1 + y_3)), (\text{permutation } (1321) \rightarrow (3213)).$$

By the modular law (A.3) of $D^{2,2,2}$ and since $x_1 \subseteq y_1$, we have

$$y_1(x_1 + y_3) = x_1 + y_1 y_3 \quad \text{and}$$

$$e_{3213} = y_3(x_3 + y_2)(x_2 + x_1 + y_1 y_3) \subseteq g_{3210}.$$

Further,

$$f_{1321} = y_1 y_2 a_1^{13} = y_1 y_2(x_1 + y_3) \subseteq e_{1321}$$

and

$$g_{13210} = e_{1321}(x_1 + a_1^{32} A_1^{32}) = e_{1321}(x_1 + (x_3 + y_2)(y_3 + x_2)) = \\ y_1(x_1 + y_3)(x_3 + y_2(x_2 + y_1))(x_1 + (x_3 + y_2)(y_3 + x_2)).$$

1.9.2. The lattice D^4 . Admissible sequences and admissible elements are taken from Table 4.3.

For $n = 1$: $\alpha = 1$, (Table 4.3, Line G11, $r = 0, s = 0, t = 0$). We have

$$e_1 = e_1 \quad (\text{admissible element } e_1 \text{ coincides with generator } e_1),$$

$$f_{10} = e_1(e_2 + e_3 + e_4) \subseteq e_1.$$

For $n = 2$: $\alpha = 21$, (Table 4.3, Line G21, $r = 0, s = 0, t = 1$). We have

$$e_{21} = e_2(e_3 + e_4),$$

$$f_{210} = e_2(e_3 + e_4)(e_4 + e_3(e_1 + e_2) + e_1) =$$

$$e_2(e_3 + e_4)(e_4 + e_2(e_1 + e_3) + e_1) =$$

$$e_2(e_3 + e_4)(e_2(e_1 + e_4) + e_2(e_1 + e_3)) \subseteq e_{21}.$$

For $n = 3$: Consider two admissible sequences: $\alpha = 121$ and $\alpha = 321 = 341$.

1) $\alpha = 121$, (Table 4.3, Line $G11$, $r = 0, s = 0, t = 1$). We have

$$\begin{aligned} e_{121} &= e_1 a_2^{34} = e_1(e_3 + e_4(e_1 + e_2)) = \\ &= e_1(e_3(e_1 + e_2) + e_4(e_1 + e_2)), \\ f_{1210} &= e_{121}(e_1 a_1^{32} + a_1^{24} a_1^{34}) = \\ &= e_{121}(e_1(e_2 + e_3) + (e_2 + e_4)(e_3 + e_4)) = \\ &= e_1(e_3 + e_4(e_1 + e_2))(e_1(e_2 + e_3) + (e_2 + e_4)(e_3 + e_4)) \subseteq e_{121}. \end{aligned}$$

2) $\alpha = 321 = 341$, (Table 4.3, Line $G31$, $t = 0, s = 0, r = 1$). We have

$$\begin{aligned} e_{321} &= e_{341} = e_3 a_1^{21} a_1^{14} = e_3(e_2 + e_1)(e_4 + e_1), \\ f_{3210} &= f_{3410} = e_{321}(a_2^{14} + e_3 a_1^{24}) = \\ &= e_{321}(e_1 + e_4(e_3 + e_2) + e_3(e_2 + e_4)) = \\ &= e_{321}(e_1 + (e_3 + e_2)(e_4 + e_2)(e_3 + e_4)) = \\ &= e_3(e_2 + e_1)(e_4 + e_1)(e_1 + (e_3 + e_2)(e_4 + e_2)(e_3 + e_4)) \subseteq e_{321}. \end{aligned}$$

For $n = 4$: $\alpha = 2341 = 2321 = 2141$, (Table 4.3, Line $F21$, $s = 0, t = 1, r = 1$). We have

$$\begin{aligned} e_{2141} &= e_2 a_2^{41} a_1^{34} = e_2(e_4 + e_1(e_3 + e_2))(e_3 + e_4), \\ f_{21410} &= e_{2141}(a_2^{34} + e_1 a_2^{24}) = \\ &= e_{2141}(e_2(e_4 + e_1(e_3 + e_2)) + e_1(e_2 + e_4(e_1 + e_3))) \subseteq e_{21}. \end{aligned}$$

1.10. Cumulative polynomials

The cumulative polynomials $x_t(n), y_t(n)$, where $t = 1, 2, 3$, and $x_0(n)$ (all of length n) are sums of all admissible elements of the same length n , where n is the length of the multi-index. In other words, cumulative polynomials are as follows

$$\begin{aligned} x_t(n) &= \sum f_{i_n \dots i_2 t}, \quad t = 1, 2, 3, \\ y_t(n) &= \sum e_{i_n \dots i_2 t}, \quad t = 1, 2, 3, \\ x_0(n) &= \sum g_{i_n \dots i_2 0}. \end{aligned}$$

From (1.26) and Theorem 2.13.1 we deduce that:

$$x_t(n) \subseteq y_t(n) \subseteq x_0(n), \quad t = 1, 2, 3.$$

The *cumulative polynomials* satisfy the same inclusions as the corresponding generators in $D^{2,2,2}$.

1.11. Examples of the cumulative polynomials

For $n = 1$:

$$\begin{aligned} x_1(1) &= f_1 = x_1, \\ y_1(1) &= e_1 = y_1, \\ x_0(1) &= g_0 = I. \end{aligned}$$

i.e., the *cumulative polynomials* for $n = 1$ coincide with the generators of $D^{2,2,2}$.

For $n = 2$: By the modular law (A.3) and since $y_3(x_1 + y_2) \subseteq x_1 + y_3$, we have

$$\begin{aligned} x_1(2) &= f_{21} + f_{31} = y_2 y_3, & (f_{21} = f_{31} = y_2 y_3), \\ y_1(2) &= e_{21} + e_{31} = y_2(x_1 + y_3) + y_3(x_1 + y_2) = \\ & (x_1 + y_3)(y_2 + y_3(x_1 + y_2)) = (x_1 + y_3)(y_2 + y_3)(x_1 + y_2), \\ x_0(2) &= g_{10} + g_{20} + g_{30} = y_1(y_2 + y_3) + y_2(y_1 + y_3) + y_3(y_2 + y_3) = \\ & (y_2 + y_3)(y_1 + y_3)(y_1 + y_2). \end{aligned}$$

For $n = 3$: (see Fig. 1.3)

$$\begin{aligned} x_1(3) &= f_{321} + f_{121} + f_{231}, \\ y_1(3) &= e_{321} + e_{121} + e_{231}. \end{aligned}$$

From Table 2.3 (Line 8, $k = 0$) $f_{121} = y_1(x_2 + x_3)$ and by §1.9.1 it follows that

$$x_1(3) = x_3(y_1 + x_2) + y_1(x_2 + x_3) + x_2(y_1 + x_3) = (x_2 + x_3)(y_1 + x_2)(y_1 + x_3).$$

Again, from Table 2.3 (Line 8, $k = 0$) $e_{121} = y_1(x_2 + y_3)(y_2 + x_3)$ and by §1.9.1, we have

$$y_1(3) = y_3(y_1 + x_2)(y_2 + x_3) + y_1(x_2 + y_3)(y_2 + x_3) + y_2(y_1 + x_3)(y_3 + x_2).$$

Finally,

$$\begin{aligned} x_0(3) &= g_{210} + g_{310} + g_{320} + g_{120} + g_{230} + g_{130} = \\ & \sum y_i(x_j + y_k)(y_j + x_i + x_k), \end{aligned}$$

where the sum runs over all permutations $\{i, j, k\}$ of $\{1, 2, 3\}$.

1.12. The sublattice $H^+(n)$ of the perfect elements

The perfect elements (see §A.6) are constructed in the following way:

For $n = 0$:

$$\begin{aligned} a_i(0) &= y_j + y_k, \\ b_i(0) &= x_i + y_j + y_k, \\ c_1(0) &= c_2(0) = c_3(0) = \sum y_i. \end{aligned} \tag{1.27}$$

For $n \geq 1$:

$$\begin{aligned} a_i(n) &= x_j(n) + x_k(n) + y_j(n+1) + y_k(n+1), \\ b_i(n) &= a_i(n) + x_i(n+1) = x_j(n) + x_k(n) + x_i(n+1) + y_j(n+1) + y_k(n+1), \\ c_i(n) &= a_i(n) + y_i(n+1) = x_j(n) + x_k(n) + y_i(n+1) + y_j(n+1) + y_k(n+1). \end{aligned}$$

Proposition (see Proposition 2.13.3, §2.12). *If $z_t(n)$ is one of the cumulative polynomials $x_t(n)$, $y_t(n)$, where $t = 1, 2, 3$, or x_0 , then*

$$\sum_{i=1,2,3} \varphi_i \Phi^+ \rho(z_t(n)) = \rho(z_t(n+1)).$$

We will also show (Proposition 2.10.1) that the property of element z to be *perfect* follows from the same property of element u if

$$\sum_{i=1,2,3} \varphi_i \Phi^+ \rho(z) = \rho(u).$$




FIGURE 1.5. The 64-element distributive lattice $H^+(n)$ of perfect elements. Two generator systems: $\{a_i \subseteq b_i \subseteq c_i\}$, §1.12 and $\{p_i \subseteq q_i \subseteq s_i\}$, §3.5.

The elements $z_t(n)$ are not *perfect*, but *perfect* elements are expressed as sums of *cumulative* polynomials.

Proposition (see Proposition 3.1.1, §3.1). *The elements $a_i(n)$, $b_i(n)$, $c_i(n)$ are perfect for all n .*

The elements $a_i(n) \subseteq b_i(n) \subseteq c_i(n)$, where $i = 1, 2, 3$, generate the 64-element distributive sublattice $H^+(n)$ of *perfect elements* for $n \geq 1$ (see Fig. 1.5) and the 27-element distributive sublattice $H^+(n)$ of *perfect elements* for $n = 0$. The sublattice $H^+(0)$ contains only 27 elements because $c_1(0) = c_2(0) = c_3(0)$.

1.13. The union of sublattices $H^+(n)$

1.13.1. Perfect cubes in the free modular lattice D^r . In [GP74], Gelfand and Ponomarev constructed the sublattice B of perfect elements for the free modular lattice D^r with r generators:

$$B = B^+ \bigcup B^-, \quad \text{where } B^+ = \bigcup_{n=1}^{\infty} B^+(n), \quad B^- = \bigcup_{n=1}^{\infty} B^-(n).$$

They proved that every sublattice $B^+(n)$ (resp. $B^-(n)$) is 2^r -element Boolean algebra, so-called *Boolean cube* (which can be also named *perfect Boolean cube*) and these *cubes* are ordered in the following way. Every element of the cube $B^+(n)$ is included in every element of the cube $B^+(n+1)$, i.e.,

$$\left. \begin{array}{l} v^+(n) \in B^+(n) \\ v^+(n+1) \in B^+(n+1) \end{array} \right\} \implies v^+(n+1) \subseteq v^+(n).$$

By analogy, the dual relation holds:

$$\left. \begin{array}{l} v^-(n) \in B^-(n) \\ v^-(n+1) \in B^-(n+1) \end{array} \right\} \implies v^-(n) \subseteq v^-(n+1).$$

1.13.2. Perfect sublattices in the lattice $D^{2,2,2}$. Let us return to the lattice $D^{2,2,2}$. Consider the modular lattice H^+ generated by the union $\bigcup_{n=0}^{\infty} H^+(n)$ and the modular lattice

H^- generated by the union $\bigcup_{n=0}^{\infty} H^-(n)$, where $H^-(n)$ is the dual modular lattice for $H^+(n)$.

The union $\bigcup_{n=0}^{\infty} H^+(n)$ is not organized as simple as $\bigcup_{n=0}^{\infty} B^+(n)$ in the case of D^r . Denote by

U_n the lower 8-element cube of $H^+(n)$ and by V_{n+1} the upper 8-element cube of $H^+(n+1)$, see Fig. 1.7.

Proposition (see Proposition 3.9.1, §3.9). *The union $U_n \cup V_{n+1}$ is the 16-element Boolean algebra.*

It is necessary to draw 8 additional edges in the places where $H^+(n)$ joins $H^+(n+1)$.

Theorem (see Theorem 3.10.2, §3.10). *The union $\bigcup_{n=0}^{\infty} H^+(n)$ is a distributive lattice mod θ . The diagram H^+ is obtained by uniting the diagrams $H^+(n)$ and joining the cubes U_n and V_{n+1} for all $n \geq 0$, i.e., it is necessary to draw 8 additional edges for all $n \geq 0$, see Fig. 1.7.*

Denote by H^- the lattice dual to the lattice H^+ .

REMARK 1.13.1. For a comparison between sublattices of the perfect elements

$$H^+ \subseteq D^{2,2,2} \text{ and } B^+ \subseteq D^4,$$

compare the Hasse diagrams of H^+ and B^+ on Fig. 1.6. From Fig. 1.6 we see that the perfect sublattice B^+ can be injectively mapped (imbedded) into the perfect sublattice H^+ mod θ . Indeed,

$$p_1(n) + p_2(n) + p_3(n) = b_1(n)b_2(n)b_3(n) \subseteq c_1(n)c_2(n)c_3(n),$$

see §3.5, Proposition 3.5.1, Table 3.5.1.

CONJECTURE 1.13.2. The lattice $H^+ \cup H^-$ contains all perfect elements of $D^{2,2,2}$ mod θ .




FIGURE 1.6. Comparison between perfect sublattices in D^4 and $D^{2,2,2}$




FIGURE 1.7. The 16-element Boolean algebra $U_n \cup V_{n+1}$

CHAPTER 2

Elementary maps φ_i and atomic, admissible, cumulative polynomials

2.1. Construction of the Coxeter functor Φ^+ by means of the lattice operations

Following Gelfand and Ponomarev [GP74, GP76] we construct the representation $\Phi^+\rho$ by means of the lattice operations. Our construction is different from the classic definition of the Coxeter functor of Bernstein-Gelfand-Ponomarev [BGP73].

The representation $\Phi^+\rho$ is constructed in the space

$$X_0^1 = \{(\eta_1, \eta_2, \eta_3) \mid \eta_i \in Y_i, \quad \sum \eta_i = 0\},$$

where i runs over the set $\{1, 2, 3\}$. Set

$$R = \bigoplus_{i=1,2,3} Y_i,$$

i.e.,

$$R = \{(\eta_1, \eta_2, \eta_3) \mid \eta_i \in Y_i, \quad i = 1, 2, 3\}.$$

Then we see that $X_0^1 \subseteq R$. Hereafter η_i denotes a vector from the space Y_i and ξ_i denotes a vector from the space X_i .

Introduce the spaces $G_i, H_i, G'_i, H'_i \subseteq R$, where $i = 1, 2, 3$:

$$\begin{aligned} G_1 &= \{(\eta_1, 0, 0) \mid \eta_1 \in Y_1\}, & H_1 &= \{(\xi_1, 0, 0) \mid \xi_1 \in X_1\}, \\ G_2 &= \{(0, \eta_2, 0) \mid \eta_2 \in Y_2\}, & H_2 &= \{(0, \xi_2, 0) \mid \xi_2 \in X_2\}, \\ G_3 &= \{(0, 0, \eta_3) \mid \eta_3 \in Y_3\}, & H_3 &= \{(0, 0, \xi_3) \mid \xi_3 \in X_3\}, \\ G'_1 &= \{(\xi_1, \eta_2, \eta_3) \mid \xi_1 \in X_1, \eta_i \in Y_i\}, & H'_1 &= \{(0, \eta_2, \eta_3) \mid \eta_i \in Y_i\}, \\ G'_2 &= \{(\eta_1, \xi_2, \eta_3) \mid \xi_2 \in X_2, \eta_i \in Y_i\}, & H'_2 &= \{(\eta_1, 0, \eta_3) \mid \eta_i \in Y_i\}, \\ G'_3 &= \{(\eta_1, \eta_2, \xi_3) \mid \xi_3 \in X_3, \eta_i \in Y_i\}, & H'_3 &= \{(\eta_1, \eta_3, 0) \mid \eta_i \in Y_i\}. \end{aligned} \tag{2.1}$$

Now, let

$$Y_i^1 = G'_i X_0^1, \tag{2.2}$$

$$X_i^1 = H'_i X_0^1, \tag{2.3}$$

$$\rho^1 = \Phi^+\rho = [X_i^1 \subseteq Y_i^1 \subseteq X_0^1 \mid i = 1, 2, 3]. \tag{2.4}$$

This construction of the Coxeter functor Φ^+ is well-defined as follows from §A.8.

2.2. The associated representations ν^0, ν^1

With representations ρ and ρ^1 we associate the following representations ν^0, ν^1 in R :

$$\nu^0(y_i) = X_0^1 + G_i, \quad \nu^0(x_i) = X_0^1 + H_i, \quad i = 1, 2, 3, \tag{2.5}$$

$$\nu^1(y_i) = X_0^1 G'_i, \quad \nu^1(x_i) = X_0^1 H'_i, \quad i = 1, 2, 3. \tag{2.6}$$

Let $\mu : X_0^1 \longrightarrow R$ be an inclusion. Then

$$\nu^1(a) = \mu\rho^1(a) \quad \text{for every } a \in D^{2,2,2}. \quad (2.7)$$

Let $\nabla : R \longrightarrow X_0$ be a projection, $\nabla : (\eta_1, \eta_2, \eta_3) \longrightarrow \Sigma\eta_i$. Then $\ker \nabla = X_0^1$.

LEMMA 2.2.1. *For every $a \in D^{2,2,2}$ and for every subspace $B \subseteq R$, we have*

$$\nabla(\nu^0(a)B) = \nabla(\nu^0(a))\nabla(B). \quad (2.8)$$

Proof. The inclusion

$$\nabla(\nu^0(a)B) \subseteq \nabla(\nu^0(a))\nabla(B)$$

is clear, because $\nu^0(a)B \subseteq \nu^0(a)$ and $\nu^0(a)B \subseteq B$. Conversely, let $z \in \nabla(\nu^0(a))\nabla(B)$, then there exist vectors

$$u \in \nu^0(a) \text{ and } v \in B \text{ such that } \nabla(u) = \nabla(v). \quad (2.9)$$

According to (2.5)

$$\nu^0(a) = X_0^1 + A$$

for some subspace $A \subseteq R$. From (2.9), we have

$$v - u = w \in \ker \nabla = X_0^1, \text{ i.e.,}$$

$$v = u + w \in X_0^1 + A = \nu^0(a), \text{ and hence } v \in \nu^0(a)B.$$

Thus, $z = \nabla(v) \in \nabla(\nu^0(a)B)$ and (2.8) is proved. \square

From (2.8) it follows that the map $\nabla : R \longrightarrow R/X_0^1$ gives a representation by the formula

$$a \longmapsto \nabla(\nu^0(a)).$$

We have

$$\nabla(\nu^0(y_i)) = \nabla G_i = Y_i = \rho(y_i), \quad i = 1, 2, 3, \quad (2.10)$$

$$\nabla(\nu^0(x_i)) = \nabla H_i = X_i = \rho(x_i), \quad i = 1, 2, 3, \quad (2.11)$$

$$\nabla(\nu^0(a)) = \rho(a) \text{ for every } a \in D^{2,2,2}. \quad (2.12)$$

2.3. The elementary maps φ_i

Following [GP74], we introduce the linear maps

$$\varphi_i : X_0^1 \longrightarrow X_0, (\eta_1, \eta_2, \eta_3) \longmapsto \eta_i.$$

Let π_i be a projection R on G_i . Then

$$\varphi_i = \nabla\pi_i\mu. \quad (2.13)$$

It follows from the definition that

$$\varphi_1 + \varphi_2 + \varphi_3 = 0. \quad (2.14)$$

From (2.14) we immediately deduce that

$$\varphi_1 B + \varphi_2 B = \varphi_1 B + \varphi_3 B = \varphi_1 B + \varphi_2 B \quad (2.15)$$

for every subspace $B \subseteq X_0^1$.

2.4. Basic relations for the maps ψ_i and φ_i

We introduce the maps $\psi_i: D^{2,2,2} \longrightarrow \mathcal{L}(R)$ useful in the study of maps φ_i . Set

$$\psi_i(a) = X_0^1 + G_i(H'_i + \nu^1(a)). \quad (2.16)$$

The map ψ_i is said to be a *joint* map.

PROPOSITION 2.4.1. *The joint maps ψ_i satisfy the following basic relations:*

- (1) $\psi_i(x_i) = X_0^1$,
- (2) $\psi_i(x_j) = \psi_k(x_j) = \nu^0(y_i y_k)$,
- (3) $\psi_i(y_i) = \nu^0(x_i(y_j + y_k))$,
- (4) $\psi_i(y_j) = \nu^0(y_i(x_j + y_k))$,
- (5) $\psi_i(I) = \nu^0(y_i(y_j + y_k))$,
- (6) $\psi_i(y_j y_k) = \nu^0(y_i(x_j + x_k))$.

Proof. 1) From (2.6) and (2.16), we have $\psi_i(x_i) = X_0^1 + G_i H'_i$. Since $G_i H'_i = 0$, we have $\psi_i(x_i) = X_0^1$.

2) From (2.6) and (2.16), we have $\psi_i(x_j) = X_0^1 + G_i(H'_i + X_0^1 H'_j)$. Since $H'_j = G_i + G_k$ (where i, j, k are distinct), it follows that

$$\psi_i(x_j) = X_0^1 + G_i(G_j + G_k + X_0^1(G_i + G_k)).$$

By the permutation property (A.4)

$$\psi_i(x_j) = X_0^1 + G_i(X_0^1 + (G_j + G_k)(G_i + G_k)) = X_0^1 + G_i(X_0^1 + G_k + G_j(G_i + G_k)).$$

Since $G_j(G_i + G_k) = 0$, it follows that

$$\psi_i(x_j) = X_0^1 + G_i(X_0^1 + G_k) = (X_0^1 + G_i)(X_0^1 + G_k) = \nu^0(y_i y_k).$$

3) As above, $\psi_i(y_i) = X_0^1 + G_i(H'_i + X_0^1 G'_i)$. Since $H'_i \subseteq G'_i$, and by the modular law (A.3), we have $\psi_i(y_i) = X_0^1 + G_i G'_i(H'_i + X_0^1)$. Since $G_i G'_i = H_i$ and $H'_j = G_i + G_k$ (where i, j, k are distinct), we have

$$\psi_i(y_i) = X_0^1 + H'_i(G_j + G_k + X_0^1) = (X_0^1 + H'_i)(G_j + G_k + X_0^1) = \nu^0(x_i(y_j + y_k)).$$

4) Again, $\psi_i(y_j) = X_0^1 + G_i(H'_i + X_0^1 G'_j)$. From $G_i \subseteq G'_j$ for $i \neq j$ and by the permutation property (A.4), we have

$$\psi_i(y_j) = X_0^1 + G_i(H'_i G'_j + X_0^1).$$

Since $H'_i G'_j = H_j + G_k$, we have

$$\psi_i(y_j) = X_0^1 + G_i(X_0^1 + H_j + G_k) = (X_0^1 + G_i)(X_0^1 + H_j + G_k) = \nu^0(y_i(x_j + y_k)).$$

5) Similarly,

$$\begin{aligned} \psi_i(I) &= X_0^1 + G_i(H'_i + X_0^1) = X_0^1 + G_i(G_j + G_k + X_0^1) = \\ &= (X_0^1 + G_i)(G_j + G_k + X_0^1) = \nu^0(y_i(y_j + y_k)). \end{aligned}$$

6) As above,

$$\psi_i(y_j y_k) = X_0^1 + G_i(H'_i + X_0^1 G'_j G'_k).$$

From $G'_j G'_k = H_j + H_k + G_i$ and by the permutation (A.4), we have

$$\begin{aligned}\psi_i(y_j y_k) &= X_0^1 + G_i((G_j + G_k)(H_j + H_k + G_i) + X_0^1) = \\ &= X_0^1 + G_i(H_j + H_k + G_i(G_j + G_k) + X_0^1).\end{aligned}$$

The intersection vanishes: $G_i(G_j + G_k) = 0$, and hence

$$\psi_i(y_j y_k) = X_0^1 + G_i(H_j + H_k + X_0^1) = \nu^0(y_i(x_j + x_k)). \quad \square$$

REMARK 2.4.2 (The lattice description of the projection, [GP74]). If $\pi: R \longrightarrow R$ is a projection, and A is a subspace of R , then

$$\pi A = \text{Im} \pi(\ker \pi + A). \quad (2.17)$$

Indeed,

$$v \in \text{Im} \pi(\ker \pi + A) \implies v = \pi v \in \pi(\ker \pi + A), \quad \text{i.e., } v \in \pi A.$$

Conversely,

$$v \in \pi A \implies v \in \text{Im} \pi, \text{ besides } v = \pi v \implies \pi v \in \pi A \implies v \in \ker \pi + A. \quad \square$$

PROPOSITION 2.4.3 (A relation between the φ_i and ψ_i). *The following properties establish a relation between the φ_i and ψ_i :*

- (1) Let $a, b \subseteq D^{2,2,2}$. If $\psi_i(a) = \nu^0(b)$, then $\varphi_i \Phi^+ \rho(a) = \rho(b)$.
- (2) Let $c \subseteq D^{2,2,2}$. Then $\nabla \psi_i(c) = \varphi_i \Phi^+ \rho(c)$.
- (3) Let $a, b, c \subseteq D^{2,2,2}$. If $\psi_i(a) = \nu^0(b)$ and $\psi_i(ac) = \psi_i(a)\psi_i(c)$, then

$$\varphi_i \Phi^+ \rho(ac) = \varphi_i \Phi^+ \rho(a) \varphi_i \Phi^+ \rho(c). \quad (2.18)$$

Proof. (1) From (2.13) we deduce that $\varphi_i \Phi^+ \rho(a) = \nabla \pi_i \mu \Phi^+ \rho(a)$. By (2.7), we have $\varphi_i \Phi^+ \rho(a) = \nabla \pi_i \nu^1(a)$. By (2.17) for $\text{Im} \pi_i = G_i$ and $\ker \pi_i = H'_i$, we have

$$\pi_i \nu^1(a) = G_i(H'_i + \nu^1(a)),$$

and

$$\nabla \pi_i \nu^1(a) = \nabla(G_i(H'_i + \nu^1(a))) = \nabla(X_0^1 + G_i(H'_i + \nu^1(a))) = \nabla(\psi_i(a)) = \nabla(\nu^0(b)).$$

From (2.12) $\nabla(\psi_i(a)) = \rho(b)$.

- (2) By definition (2.16) of ψ_i , we have

$$\nabla \psi_i(c) = \nabla(X_0^1 + G_i(H'_i + \nu^1(c))).$$

Since $X_0^1 \subseteq \ker \nabla$, we have $\nabla \psi_i(c) = \nabla(G_i(H'_i + \nu^1(c)))$. By lattice description of the projection (2.17), we have $\nabla \psi_i(c) = \nabla(\pi_i \nu^1(c))$. By (2.7) $\nabla \psi_i(c) = \nabla(\pi_i \mu \Phi^+ \rho(c))$. Finally, by definition (2.13) of φ_i , we have $\nabla \psi_i(c) = \varphi_i \Phi^+ \rho(c)$.

- (3) By the hypothesis $\psi_i(ac) = \nu^0(a)\psi_i(c)$. By Lemma 2.2.1 and eq.(2.8), we have

$$\nabla \psi_i(ac) = \nabla \psi_i(a) \nabla \psi_i(c).$$

The proof follows from (2). \square

From Propositions 2.4.1, 2.4.3 we have

COROLLARY 2.4.4. *For the elementary map φ_i , the following basic relations hold:*

- (1) $\varphi_i \Phi^+ \rho(x_i) = 0$,
- (2) $\varphi_i \Phi^+ \rho(x_j) = \varphi_k \Phi^+ \rho(x_j) = \rho(y_i y_k)$,
- (3) $\varphi_i \Phi^+ \rho(y_i) = \rho(x_i(y_j + y_k))$,
- (4) $\varphi_i \Phi^+ \rho(y_j) = \rho(y_i(x_j + y_k))$,
- (5) $\varphi_i \Phi^+ \rho(I) = \rho(y_i(y_j + y_k))$,
- (6) $\varphi_i \Phi^+ \rho(y_j y_k) = \rho(y_i(x_j + x_k))$.

Proof. It is necessary to explain only (1). Since the i th coordinate in the space H'_i is 0, it follows that

$$\varphi_i \Phi^+ \rho(x_i) = \nabla \pi_i \mu \Phi^+ \rho(x_i) = \nabla \pi_i \nu^1(x_i) = \nabla \pi_i (X_0^1 H'_i) = 0. \quad \square$$

2.5. Additivity and multiplicativity of the joint maps ψ_i

PROPOSITION 2.5.1. *The map ψ_i is additive and quasimultiplicative¹ with respect to the lattice operations $+$ and \cap . The following relations hold for every $a, b \in D^{2,2,2}$:*

- (1) $\psi_i(a) + \psi_i(b) = \psi_i(a + b)$,
- (2) $\psi_i(a)\psi_i(b) = \psi_i((a + x_i)(b + x_j x_k))$,
- (3) $\psi_i(a)\psi_i(b) = \psi_i(a(b + x_i + x_j x_k))$.

Proof. 1) By the modular law (A.3)

$$\begin{aligned} \psi_i(a) + \psi_i(b) &= X_0^1 + G_i(H'_i + \nu^1(a)) + G_i(H'_i + \nu^1(b)) = \\ &= X_0^1 + G_i(H'_i + \nu^1(a) + G_i(H'_i + \nu^1(b))) = X_0^1 + G_i((H'_i + \nu^1(b))(H'_i + G_i) + \nu^1(a)). \end{aligned}$$

Since $H'_i + G_i = R$, it follows that

$$\psi_i(a) + \psi_i(b) = X_0^1 + G_i(H'_i + \nu^1(b) + \nu^1(a)) = X_0^1 + G_i(H'_i + \nu^1(b + a)).$$

2) By definition (2.6) $\nu^1(b) \subseteq X_0^1$, and by the permutation property (A.5) we have

$$X_0^1 + G_i(H'_i + \nu^1(a)) = X_0^1 + H'_i(G_i + \nu^1(a)).$$

By the modular law (A.3) and by (A.5) we have

$$\begin{aligned} \psi_i(a)\psi_i(b) &= \\ &= (X_0^1 + G_i(H'_i + \nu^1(a))) (X_0^1 + G_i(H'_i + \nu^1(b))) = \\ &= X_0^1 + G_i(H'_i + \nu^1(a)) (X_0^1 + H'_i(G_i + \nu^1(b))). \end{aligned}$$

Further,

$$\psi_i(a)\psi_i(b) = X_0^1 + G_i(X_0^1(H'_i + \nu^1(a)) + H'_i(G_i + \nu^1(b))).$$

Since

$$X_0^1(H'_i + \nu^1(a)) = X_0^1 H'_i + \nu^1(a) \quad \text{and} \quad X_0^1 H'_i = \nu^1(x_i),$$

we have

$$\psi_i(a)\psi_i(b) = X_0^1 + G_i(\nu^1(x_i) + \nu^1(a) + H'_i(G_i + \nu^1(b))). \quad (2.19)$$

By the permutation property (A.4) and by (2.19) we have

$$\psi_i(a)\psi_i(b) = X_0^1 + G_i(H'_i + (\nu^1(x_i) + \nu^1(a))(G_i + \nu^1(b))). \quad (2.20)$$

¹The map ψ_i is multiplicative if $\psi_i(a)\psi_i(b) = \psi_i(ab)$. We will say that ψ_i is *quasimultiplicative* if element ab in this relation is a little deformed, or more exactly, if ab is deformed to $(a + x_i)(b + x_j x_k)$ or to $a(b + x_i + x_j x_k)$ as in (2) and (3) of Proposition 2.5.1.

Since

$$G_i = H'_j H'_k \quad \text{and} \quad \nu^1(x_i) + \nu^1(a) = \nu^1(x_i + a) = X_0^1(\nu^1(x_i + a)),$$

it follows that

$$\begin{aligned} \psi_i(a)\psi_i(b) &= X_0^1 + G_i (H'_i + \nu^1(x_i + a)(X_0^1 H'_j H'_k + \nu^1(b))) = \\ &= X_0^1 + G_i (H'_i + \nu^1(x_i + a)(\nu^1(x_j x_k) + \nu^1(b))) = \\ &= X_0^1 + G_i (H'_i + \nu^1(x_i + a)\nu^1(x_j x_k + b)) = \psi_i((a + x_i)(b + x_j x_k)). \end{aligned}$$

3) From (2.19) and since $\nu^1(x_i) = X_0^1 H'_i \subseteq H'_i$, we have

$$\psi_i(a)\psi_i(b) = X_0^1 + G_i (\nu^1(a) + H'_i(G_i + \nu^1(b) + \nu^1(x_i))). \quad (2.21)$$

Again, by (A.4) we have

$$\begin{aligned} \psi_i(a)\psi_i(b) &= X_0^1 + G_i (H'_i + \nu^1(a)(G_i + \nu^1(b) + \nu^1(x_i))) = \\ &= X_0^1 + G_i (H'_i + \nu^1(a)(X_0^1 H'_j H'_k + \nu^1(b) + \nu^1(x_i))). \end{aligned}$$

Thus, $\psi_i(a)\psi_i(b) = \psi_i(a(b + x_i + x_j x_k))$. \square

COROLLARY 2.5.2 (Atomic multiplicativity). *1) Let one of the following inclusions hold:*

- (i) $x_i + x_j x_k \subseteq a$,
- (ii) $x_i + x_j x_k \subseteq b$,
- (iii) $x_j \subseteq a$, $x_j x_k \subseteq b$,
- (iv) $x_i \subseteq b$, $x_j x_k \subseteq a$.

Then the joint map ψ_i operates as a homomorphism on the elements a and b with respect to the lattice operations $+$ and \cap , i.e.,

$$\psi_i(a) + \psi_i(b) = \psi_i(a + b), \quad \psi_i(a)\psi_i(b) = \psi_i(a \cap b).$$

2) The joint map ψ_i applied to the following atomic elements is the intersection preserving map, i.e., multiplicative with respect to the operation \cap :

$$\psi_i(ba_n^{ij}) = \psi_i(b)\psi_i(a_n^{ij}) \text{ for every } b \subseteq D^{2,2,2}, \quad (2.22)$$

$$\psi_j(ba_n^{ij}) = \psi_j(b)\psi_j(a_n^{ij}) \text{ for every } b \subseteq D^{2,2,2}, \quad (2.23)$$

$$\psi_i(bA_n^{ij}) = \psi_i(b)\psi_i(A_n^{ij}) \text{ for every } b \subseteq D^{2,2,2}. \quad (2.24)$$

3) The element x_i can be inserted in any additive expression under the action of the joint map ψ_i :

$$\psi_i(b + x_i) = \psi_i(b) \text{ for every } b \subseteq D^{2,2,2}. \quad (2.25)$$

Proof. 1) Follows from Proposition 2.5.1.

2) Let us consider a_n^{ij} . For $n \geq 2$, we have

$$a_n^{ij} = x_i + y_j a_{n-1}^{jk} = x_i + y_j(x_j + y_k a_{n-2}^{ki}) = x_i + x_j + y_j y_k a_{n-2}^{ki},$$

thus $a_n^{ij} \supseteq x_i + x_j$. Now, we consider atomic element A_n^{ij} . For $n \geq 2$, we have

$$A_n^{ij} = y_i + x_j A_{n-1}^{ki} = y_i + x_j(y_k + x_i A_{n-2}^{ji}) \supseteq x_i + x_j x_k.$$

For $n = 1$, the relation is proved directly.

3) Follows from Corollary 2.4.4, (1) since we have

$$\psi_i(b + x_i) = \psi_i(b) + \psi_i(x_i) \text{ and } \psi_i(x_i) = X_0^1 \subseteq \psi_i(b)$$

for every $b \in D^{2,2,2}$. \square

2.6. Basic properties of atomic elements a_n^{ij} and A_n^{ij}

The *atomic elements* are defined in §1.5. These polynomials are building bricks in the further constructions.

1.1	$a_n^{ij} = x_i + y_j a_{n-1}^{jk}$ (definition)	1.2	$A_n^{ij} = y_i + x_j A_{n-1}^{ki}$ (definition)
2.1	$a_0^{ij} \supseteq a_1^{ij} \supseteq \cdots a_n^{ij} \supseteq \cdots \supseteq x_i + x_j$	2.2	$A_0^{ij} \supseteq A_1^{ij} \supseteq \cdots A_n^{ij} \supseteq \cdots$
3.1	$x_k A_n^{ij} = x_k a_n^{ji}$	3.2	$A_n^{ij} = y_i + x_j a_{n-1}^{ik}$
4.1	$y_i y_j A_n^{ki} = y_i y_j a_n^{ik}$	4.2	$y_i a_{n+1}^{ij} = y_i (x_i + y_j A_n^{kj})$
5.1	$a_n^{ij} a_m^{jk} = A_n^{ji} a_m^{jk}$ for $n \leq m+1$	5.2	$y_i (x_j + x_k) a_n^{ij} = y_i (x_j + x_k) A_n^{ji}$
6.1	$A_n^{ij} a_m^{ki} = y_i a_{m-1}^{ij} + x_k A_n^{ij}$	6.2	$A_n^{ij} a_m^{jk} = y_i a_m^{jk} + x_j A_{n-1}^{ki}$
7.1	$y_i (x_j + x_k) + y_i y_j a_{n-2}^{ik} = y_i a_n^{kj}$	7.2	$y_i y_j + y_i (x_j + x_k) A_{n-2}^{ki} = y_i A_n^{jk}$
8.1	$x_i (y_j + y_k) + y_i y_j A_{n-1}^{kj} =$ $y_i (y_j + y_k) a_n^{ij}$	8.2	$y_i y_j + x_j (y_i + y_k) A_{n-1}^{ki} =$ $y_j (y_i + y_k) A_n^{ij}$

TABLE 2.1. The basic properties of atomic elements a_n^{ij} and A_n^{ij} in $D^{2,2,2}$

The basic properties of atomic elements are given in Table 2.1. These properties, which will be used in further considerations. For convenience, we also give definition of a_n^{ij}, A_n^{ij} in relations (1.1)–(1.2) of Table 2.1. Indices i, j, k in Table 2.1 are distinct and $\{i, j, k\} = \{1, 2, 3\}$. Properties 3.1, 3.2, 4.1, 4.2, 5.1, 5.2 from Table 2.1 mean that atomic elements a_n^{ij} and A_n^{ji} coincide for many lattice polynomials and can substitute one another in many cases.

Proof of Properties 2.1 – 8.2 (Table 2.1).

2.1) First, $a_{n-1}^{ij} = x_i + y_j a_{n-2}^{jk}$, $a_n^{ij} = x_i + y_j a_{n-1}^{jk}$. By induction, if $a_{n-2}^{jk} \supseteq a_{n-1}^{jk}$, then $a_{n-1}^{ij} \supseteq a_n^{ij}$. For $n = 1$, we have $a_0^{ij} = I \supseteq a_1^{ij}$. Further,

$$a_n^{ij} = x_i + y_j (x_j + y_k a_{n-2}^{ki}) = x_i + x_j + y_j y_k a_{n-2}^{ki} \supseteq x_i + x_j. \quad \square$$

2.2) Similarly,

$$A_{n-1}^{ij} = y_i + x_j A_{n-2}^{ki} \supseteq y_i + x_j A_{n-1}^{ki} = A_n^{ij}. \quad \square$$

3.1) By the induction hypothesis $x_j A_{n-1}^{ki} = x_j a_{n-1}^{ik}$, then

$$x_k A_n^{ij} = x_k (y_i + x_j A_{n-1}^{ki}) = x_k (y_i + x_j a_{n-1}^{ik}).$$

By property 1.1 from Table 2.1 we have $a_{n-1}^{ik} \supseteq x_k$, thus by the permutation property (A.4) we deduce

$$x_k A_n^{ij} = x_k (x_j + y_i a_{n-1}^{ik}) = x_k a_n^{ji}. \quad \square$$

3.2) By property 3.1 from Table 2.1 we have $A_n^{ij} = y_i + x_j A_{n-1}^{ki} = y_i + x_j a_{n-1}^{ik}$. \square

4.1) We have

$$y_i y_j a_n^{ik} = y_i y_j (x_i + y_k a_{n-1}^{kj}) = y_i y_j (x_i + y_k (x_k + y_j a_{n-2}^{ji})),$$

i.e.,

$$y_i y_j a_n^{ik} = y_i y_j (x_i + x_k + y_k y_j a_{n-2}^{ji}). \quad (2.26)$$

On the other hand,

$$y_i y_j A_n^{ki} = y_{ij} (y_k + x_i A_{n-1}^{jk}) = y_i y_j (y_k + x_i (y_j + x_k A_{n-2}^{ij})).$$

Since $A_{n-2}^{ij} \supseteq x_i$, we have by the permutation property (A.4):

$$y_i y_j A_n^{ki} = y_i y_j (y_k + x_i (x_k + y_j A_{n-2}^{ij})).$$

Further, since $y_j A_{n-2}^{ij} \supseteq y_i y_j$, again, by (A.4), we have

$$y_i y_j A_n^{ki} = y_i y_j (x_i + y_k (x_k + y_j A_{n-2}^{ij})) = y_i y_j (x_i + x_k + y_k y_j A_{n-2}^{ij}).$$

By the induction hypothesis

$$y_i y_j A_n^{ki} = y_i y_j (x_i + x_k + y_k y_j a_{n-2}^{ji})$$

and by (2.26) we have $y_i y_j A_n^{ki} = y_i y_j a_n^{ik}$. \square

4.2) By property 4.1 we have

$$y_i a_{n+1}^{ij} = y_i (x_i + y_j a_n^{jk}) = x_i + y_i y_j a_n^{jk} = x_i + y_i y_j A_n^{kj} = y_i (x_i + y_j A_n^{kj}). \quad \square$$

5.1) By property 2.1 and (A.4) we have

$$a_n^{ij} a_m^{jk} = (x_i + y_j a_{n-1}^{jk}) a_m^{jk} = (y_j + x_i a_{n-1}^{jk}) a_m^{jk}.$$

By property 3.1 $a_n^{ij} a_m^{jk} = (y_j + x_i A_{n-1}^{kj}) a_m^{jk} = A_n^{ji} a_m^{jk}$. \square

5.2) We have

$$y_i (x_j + x_k) a_n^{ij} = y_i (x_j + x_k) (x_i + y_j a_{n-1}^{jk}).$$

Since $a_{n-1}^{jk} \supseteq x_j + x_k$, it follows that

$$\begin{aligned} y_i (x_j + x_k) a_n^{ij} &= y_i (x_j + x_k) (y_j + x_i a_{n-1}^{jk}) = \\ y_i (x_j + x_k) (y_j + x_i A_{n-1}^{kj}) &= y_i (x_j + x_k) A_n^{ji}. \quad \square \end{aligned}$$

6.1) Since $y_i a_{m-1}^{ij} \subseteq A_n^{ij}$, it follows that

$$A_n^{ij} a_m^{ki} = A_n^{ij} (x_k + y_i a_{m-1}^{ij}) = x_k A_n^{ij} + y_i a_{m-1}^{ij}. \quad \square$$

6.2) Since $x_j A_{n-1}^{ki} \subseteq a_m^{jk}$, it follows that

$$A_n^{ij} a_m^{jk} = a_m^{jk} (y_i + x_j A_{n-1}^{ki}) = y_i a_m^{jk} + x_j A_{n-1}^{ki}. \quad \square$$

7.1) Since

$$y_i (x_j + x_k) + y_i y_j a_{n-2}^{ik} = y_i (x_j + x_k + y_i y_j a_{n-2}^{ik}) = y_i (x_k + y_j (x_j + y_i a_{n-2}^{ik})),$$

by definition (1.1, Table 2.1) we have $y_i (x_k + y_j (x_j + y_i a_{n-2}^{ik})) = y_i a_n^{kj}$. \square

7.2) As above, using the modular law (A.3) and the permutation property (A.4) we have

$$\begin{aligned} y_i y_j + y_i (x_j + x_k) A_{n-2}^{ki} &= y_i ((x_j + x_k) A_{n-2}^{ki} + y_i y_j) = \\ y_i (x_k + x_j A_{n-2}^{ki} + y_i y_j) &= y_i (x_k + y_j (y_i + x_j A_{n-2}^{ki})) = y_i (x_k + y_j A_{n-1}^{ji}) = \\ y_i (y_j + x_k A_{n-1}^{ij}) &= y_i A_n^{jk}. \quad \square \end{aligned}$$

8.1) Since

$$x_i(y_j + y_k) + y_i y_j A_{n-1}^{kj} = (y_j + y_k)(x_i + y_i y_j A_{n-1}^{kj}),$$

by property 4.1 we have

$$\begin{aligned} x_i(y_j + y_k) + y_i y_j A_{n-1}^{kj} &= (y_j + y_k)(x_i + y_i y_j A_{n-1}^{kj}) = \\ y_i(y_j + y_k)(x_i + y_j A_{n-1}^{jk}) &= y_i(y_j + y_k) a_n^{ij}. \quad \square \end{aligned}$$

8.2) As above,

$$y_i y_j + x_j(y_i + y_k) A_{n-1}^{ki} = y_j(y_i + y_k)(y_i + x_j A_{n-1}^{ki}) = y_i(y_j + y_k) A_n^{ij}. \quad \square$$

2.7. Action of the maps ψ_i and φ_i on atomic elements

PROPOSITION 2.7.1 (Action of joint maps on atomic elements). *The joint maps ψ_i applied to the atomic elements a_n^{ij} and A_n^{ij} ($n \geq 1$) satisfy the following relations:*

- (1) $\psi_i(a_n^{ij}) = \nu^0(y_i A_n^{kj})$,
- (2) $\psi_j(a_n^{ij}) = \begin{cases} \nu^0(y_j A_n^{kj}) & \text{for } n > 1, \\ \nu^0(y_j(y_i + y_k) A_1^{kj}) & \text{for } n = 1, \end{cases}$
- (3) $\psi_k(y_i a_n^{ij}) = \nu^0(y_k A_{n+1}^{ji})$,
- (4) $\psi_i(A_n^{ij}) = \nu^0(y_i(y_j + y_k) a_n^{ik})$,
- (5) $\psi_j(y_k A_n^{ij}) = \nu^0(y_j(x_k + y_i) a_n^{ik})$,
- (6) $\psi_k(y_j A_n^{ij}) = \nu^0(y_k a_{n+1}^{ji})$.

REMARK 2.7.2. The *joint* map ψ_k increases the lower index n of the atomic element and swaps indices i and j in headings (3) and (6). The new atomic elements appear in this way.

$$ij \implies ji, \quad n \implies n + 1.$$

In headings (1) and (2), the lower index n does not grow and the upper pair ij becomes kj . The first index of the pair is changed and the atomic elements “ a ” are transformed to atomic elements “ A ”:

$$ij \implies kj, \quad a \implies A.$$

In headings (4) and (5), the lower index n does not grow and the upper pair ij becomes ik . The second index of the pair is changed and the atomic elements “ A ” are transformed to atomic elements “ a ”:

$$ij \implies ik, \quad A \implies a.$$

Map ψ_k transforms $a \implies A$ in headings (1), (2), (3) and $A \implies a$ in headings (4), (5), (6).

REMARK 2.7.3. It is not so interesting to consider $\psi_j(A_n^{ij})$ because

$$\psi_j(A_n^{ij}) = \psi_j(y_i + x_j A_n^{ki}) = \psi_j(y_i).$$

By basic relations (Proposition 2.4.1) $\psi_j(y_i) = \nu^0(y_j(x_i + y_k))$.

For further moving of indices, see §2.11-§2.12 devoted to the admissible elements.

Proof Proposition 2.7.1 uses a simultaneous induction on (1)–(6). For convenience, without loss of generality, we let $i = 1, j = 2$.

1) For $n = 1$, by Proposition 2.4.1 we have

$$\psi_1(a_1^{12}) = \psi_1(x_1 + y_2) = \psi_1(y_2) = \nu^0(y_1(x_2 + y_3)) = \nu^0(y_1 A_1^{32}).$$

For $n \geq 2$, by Corollary 2.4.4 we have

$$\psi_1(a_n^{12}) = \psi_1(x_1 + y_2 a_{n-1}^{23}) = \psi_1(y_2 a_{n-1}^{23}).$$

Since

$$y_2 a_{n-1}^{23} = y_2(x_2 + y_3 a_{n-2}^{31}) = x_2 + y_2 y_3 a_{n-2}^{31}$$

and $a_{n-2}^{31} \supseteq x_1$, we see by multiplicativity (Corollary 2.5.2) that

$$\psi_1(a_n^{12}) = \psi_1(x_2) + \psi_1(y_2 y_3) \psi_1(a_{n-2}^{31}).$$

By basic relations (Proposition 2.4.1)

$$\psi_1(a_n^{12}) = \nu^0(y_1 y_3) + \nu^0(y_1(x_2 + x_3)) \psi_1(a_{n-2}^{31}).$$

By the induction hypothesis from (2) we have $\psi_1(a_{n-2}^{31}) = \nu^0(y_1 A_{n-2}^{21})$ and

$$\psi_1(a_n^{12}) = \nu^0(y_1 y_3 + y_1(x_2 + x_3) A_{n-2}^{21}).$$

By property 7.2 from Table 2.1 we have

$$\psi_1(a_n^{12}) = \nu^0(y_1 A_n^{32}). \quad \square$$

2) For $n = 1$, we have

$$\begin{aligned} \psi_2(a_1^{12}) &= \psi_2(x_1) + \psi_2(y_2) = \nu^0(y_2 y_3 + x_2(y_1 + y_3)) = \\ &= \nu^0(y_2(y_1 + y_3)(y_3 + x_2)) = \nu^0(y_2(y_1 + y_3) A_1^{32}). \end{aligned}$$

For $n \geq 2$, we have $\psi_2(a_n^{12}) = \psi_2(x_1) + \psi_2(y_2 a_{n-1}^{23})$. By the multiplicativity property (Corollary 2.5.2) we have

$$\psi_2(a_n^{12}) = \nu^0(y_2 y_3) + \psi_2(y_2) \psi_2(a_{n-1}^{23}).$$

By the induction hypothesis, from (1) we have $\psi_2(a_{n-1}^{23}) = \nu^0(y_2 A_{n-1}^{13})$ and

$$\psi_2(a_n^{12}) = \nu^0(y_2 y_3 + x_2(y_1 + y_3 A_{n-1}^{13})).$$

Since $A_{n-1}^{13} \subseteq y_1 + y_3$, we see that

$$\psi_2(a_n^{12}) = \nu^0(y_2 y_3 + x_2 A_{n-1}^{13}) = \nu^0(y_2(y_3 + x_2 A_{n-1}^{13})) = \nu^0(y_2 A_n^{32}). \quad \square$$

3) We have

$$\psi_3(y_1 a_n^{12}) = \psi_3(y_1(x_1 + y_2 a_{n-1}^{23})) = \psi_3(x_1 + y_1 y_2 a_{n-1}^{23}).$$

Further, by multiplicativity (Corollary 2.5.2) we have

$$\psi_3(y_1 a_n^{12}) = \psi_3(x_1) + \psi_3(y_1 y_2) \psi_3(a_{n-1}^{23}).$$

From basic relations (Proposition 2.4.1) it follows that

$$\psi_3(y_1 a_n^{12}) = \nu^0(y_3 y_2) + \nu^0(y_3(x_1 + x_2)) \psi_3(a_{n-1}^{23}).$$

By induction, from (2) $\psi_3(y_1 a_n^{12}) = \nu^0(y_3 y_2 + y_3(x_1 + x_2) A_{n-1}^{13})$. Again, from Table 2.1, property 7.2 we have $\psi_3(y_1 a_n^{12}) = \nu^0(y_3 A_{n+1}^{21})$. In this case ψ_k increases the lower index of the atomic element a_n^{ij} . \square

4) As above,

$$\psi_1(A_n^{12}) = \psi_1(y_1 + x_2 A_{n-1}^{31}) = \psi_1(y_1) + \psi_1(x_2 A_{n-1}^{31}).$$

From Table 2.1, property 3.1 we have

$$\psi_1(A_n^{12}) = \psi_1(y_1) + \psi_1(x_2 a_{n-1}^{13}) = \psi_1(y_1) + \psi_1(x_2) \psi_1(a_{n-1}^{13}).$$

By basic relations (Proposition 2.4.1) and by induction hypothesis, from (1) we have

$$\psi_1(A_n^{12}) = \nu^0(x_1(y_2 + y_3) + y_1 y_3 A_{n-1}^{23}).$$

From Table 2.1, property 8.1 we have

$$\psi_1(A_n^{12}) = \nu^0(y_1(y_2 + y_3)a_n^{13}). \quad \square$$

5) Again, $\psi_2(y_3A_n^{12}) = \psi_2(y_3(y_1 + x_2A_{n-1}^{31}))$. Since $y_3 \subseteq A_{n-1}^{31}$, we have by the permutation property (A.4):

$$\psi_2(y_3A_n^{12}) = \psi_2(y_3(x_2 + y_1A_{n-1}^{31})).$$

By (2.22)-(2.25) and Corollary 2.5.2 we have

$$\psi_2(y_3A_n^{12}) = \psi_2(y_3)(\psi_2(x_2) + \psi_2(y_1A_{n-1}^{31})) = \psi_2(y_3)\psi_2(y_1A_{n-1}^{31}).$$

By the induction hypothesis, from (6) it follows that

$$\psi_2(y_3A_n^{12}) = \nu^0(y_2(x_3 + y_1)a_n^{13}))$$

because $\psi_2(y_1A_{n-1}^{31}) = \nu^0(y_2a_n^{13})$. \square

6) We have

$$\psi_2(y_1A_n^{31}) = \psi_2(y_1(y_3 + x_1A_{n-1}^{23})) = \psi_2(y_1y_3 + x_1a_{n-1}^{32}) = \psi_2(y_1y_3) + \psi_2(x_1)\psi_2(a_{n-1}^{32}).$$

By the induction hypothesis, from (2) it follows that

$$\psi_2(y_1A_n^{31}) = \nu^0(y_2(x_1 + x_3) + y_2y_3A_{n-1}^{12}).$$

From Table 2.1, property 4.1 we have

$$\psi_2(y_1A_n^{31}) = \nu^0(y_2(x_1 + x_3) + y_2y_3a_{n-1}^{21}).$$

Finally, from Table 2.1, property 7.1 $\psi_2(y_1A_n^{31}) = \nu^0(y_2a_{n+1}^{13})$. In this case ψ_k increases the lower index of the atomic element A_n^{ij} . \square

From Propositions (2.4.3) and (2.7.1) we have

COROLLARY 2.7.4 (Action of the elementary maps). *The elementary maps φ_i applied to the atomic elements a_n^{ij} and A_n^{ij} ($n \geq 1$) satisfy the following relations:*

- (1) $\varphi_i\Phi^+\rho(a_n^{ij}) = \rho(y_iA_n^{kj}),$
- (2) $\varphi_j\Phi^+\rho(a_n^{ij}) = \begin{cases} \rho(y_jA_n^{kj}) & \text{for } n > 1, \\ \rho(y_j(y_i + y_k)A_1^{kj}) & \text{for } n = 1, \end{cases}$
- (3) $\varphi_k\Phi^+\rho(y_ia_n^{ij}) = \rho(y_kA_{n+1}^{ji}),$
- (4) $\varphi_i\Phi^+\rho(A_n^{ij}) = \rho(y_i(y_j + y_k)a_n^{ik}),$
- (5) $\varphi_j\Phi^+\rho(y_kA_n^{ij}) = \rho(y_j(x_k + y_i)a_n^{ik}),$
- (6) $\varphi_k\Phi^+\rho(y_jA_n^{ij}) = \rho(y_ka_{n+1}^{ji}).$

2.8. φ_i -homomorphic theorem

THEOREM 2.8.1. 1) *The polynomials a_n^{ij} are φ_i - and φ_j -homomorphic.*

2) *The polynomials A_n^{ij} are φ_i -homomorphic.*

3) *The polynomials A_n^{ij} are (φ_j, y_k) -homomorphic.*

Proof. 1), 2) Follow from multiplicativity (2.22) – (2.24), Proposition 2.7.1 and Proposition 2.4.3. \square

3) For convenience, without loss of generality, we let $i = 1, j = 2, k = 3$. We need to prove that

$$\varphi_2 \Phi^+ \rho(A_n^{12} p) = \varphi_2 \Phi^+ \rho(y_3 A_n^{12}) \varphi_2 \Phi^+ \rho(p) \text{ for every } p \subseteq y_3. \quad (2.27)$$

Since $\psi_2(A_n^{12} p) = \psi_2((y_1 + x_2 A_{n-1}^{31}) p)$ and $p \subseteq y_3 \subseteq A_{n-1}^{31}$, we have by the permutation property (A.4):

$$\psi_2(A_n^{12} p) = \psi_2((x_2 + y_1 A_{n-1}^{31}) p) = \psi_2(x_2 + y_1 A_{n-1}^{31}) \psi_2(p).$$

The element x_2 can be dropped according to (2.25), and hence,

$$\psi_2(A_n^{12} p) = \psi_2(y_1 A_{n-1}^{31}) \psi_2(p).$$

By property (6) from Proposition 2.7.1 we have

$$\psi_2(A_n^{12} p) = \nu^0(y_2 a_n^{13}) \psi_2(p). \quad (2.28)$$

Since $y_2 a_n^{13} = y_2(y_1 + y_3) a_n^{13}$, we have $\nu^0(y_2 a_n^{13}) = \nu^0(y_2(y_1 + y_3)) \nu^0(a_n^{13})$ and by (5) from Proposition 2.4.1 we have $\nu^0(y_2 a_n^{13}) = \psi_2(I) \nu^0(a_n^{13})$. Further, $\psi_2(I) \supseteq \psi_2(p)$ and (2.28) is equivalent to

$$\psi_2(A_n^{12} p) = \nu^0(a_n^{13}) \psi_2(p). \quad (2.29)$$

We have $\psi_2(p) \subseteq \psi_2(y_3)$ together with $p \subseteq y_3$, and therefore $\psi_2(p) = \psi_2(p) \psi_2(y_3)$. From (2.29) and Proposition 2.4.1 we have

$$\begin{aligned} \psi_2(A_n^{12} p) &= \nu^0(a_n^{13}) \psi_2(p) \psi_2(y_3) = \\ &= \nu^0(a_n^{13}) \psi_2(p) \nu^0(y_2(x_1 + y_3)) = \nu^0(y_2(x_1 + y_3) a_n^{13}) \psi_2(p). \end{aligned}$$

By (5) from Proposition 2.7.1 $\psi_2(A_n^{12} p) = \psi_2(y_3 A_n^{12}) \psi_2(p)$. Applying projection ∇ from §2.2 we get

$$\nabla \psi_2(A_n^{12} p) = \nabla \psi_2(y_3 A_n^{12}) \nabla \psi_2(p) \quad (2.30)$$

because $\psi_2(y_3 A_n^{12}) = \nu^0(y_2(x_1 + y_3) a_n^{13})$, see Lemma 2.2.1, relation (2.8).

By Proposition 2.4.3 $\nabla \psi_i(c) = \varphi_i \Phi^+ \rho(c)$ for every $c \subseteq D^{2,2,2}$, thus we get (2.27), and therefore heading (3) of the φ -homomorphic Theorem 2.8.1 is proven. \square

The final proposition of this section shows when the lower indices of atomic elements are increased by action some elementary mappings φ_i :

COROLLARY 2.8.2. *The index growth takes place in the following cases:*

- (1) $\varphi_k \Phi^+ \rho(y_i a_n^{ij}) = \rho(y_k A_{n+1}^{ji});$
- (2) $\varphi_k \Phi^+ \rho(y_j a_n^{ij}) = \rho(y_k a_{n+1}^{ji}).$

2.9. The perfectness of $y_i + y_j$

To prove the *perfectness* of the element $y_1 + y_2$, two simple lemmas are necessary.

LEMMA 2.9.1. *Let $A, B \subseteq X, A, B$ be subspaces in the finite-dimensional vector space X . There exists a subspace $C \subseteq B$, such that*

$$C + AB = B \text{ and } CA = 0. \quad (2.31)$$

Indeed, take any direct complement C of AB , i.e., $C \oplus AB = B$. If $v \in CA$ is non-zero, then $v \in CA \subseteq BA$ and the sum $C + AB$ is not direct. \square

LEMMA 2.9.2. *If $U, V, W \subseteq X$ and $(U + V)W = 0$, then*

$$U(V + W) = UV \text{ and } V(U + W) = UV. \quad (2.32)$$

Indeed,

$$U(V + W) \subseteq (U + V)(V + W) = (U + V)W + V = V,$$

i.e., $U(V + W) \subseteq UV$. The inverse inclusion is obvious. Similarly,

$$V(U + W) \subseteq (U + V)(U + W) = U + W(U + V) = U,$$

i.e., $V(U + W) \subseteq UV$ and the inverse inclusion is obvious. \square

PROPOSITION 2.9.3. 1) *The element $y_i + y_j$ where $i \neq j$, is perfect in $D^{2,2,2}$.*

2) *If $\rho(y_i + y_j) = 0$ for some indecomposable representation ρ , where $i \neq j$, then ρ is one from the seven projective representations (Table A.1).*

Proof. 1) Consider $y_1 + y_2$. Let $B = X_3$, $A = Y_1 + Y_2$ in (2.31), then there exists a subspace $C \subseteq X_3$ such that

$$C + X_3(Y_1 + Y_2) = X_3, \quad C(Y_1 + Y_2) = 0. \quad (2.33)$$

Therefore $Y_3 \supseteq C + Y_3(Y_1 + Y_2)$. Again use (2.31) with

$$B = Y_3, A = C + Y_3(Y_1 + Y_2).$$

Then, there exists a subspace $D \subseteq Y_3$ such that

$$D + C + Y_3(Y_1 + Y_2) = Y_3, \quad D[C + Y_3(Y_1 + Y_2)] = 0. \quad (2.34)$$

Using (2.32) with $U = Y_3(Y_1 + Y_2)$, $V = C$ and $W = D$, we obtain from (2.32) and (2.34):

$$U(V + W) = Y_3(Y_1 + Y_2)(C + D) = CY_3(Y_1 + Y_2).$$

By (2.33) we have $Y_3(Y_1 + Y_2)(C + D) = 0$. Since $D + C \subseteq Y_3$, we see that $(Y_1 + Y_2)(C + D) = 0$. Therefore, $D + C$ and $Y_1 + Y_2$ form a direct sum. We complement this sum to X_0 :

$$(D + C) \oplus (Y_1 + Y_2) \oplus F = X_0. \quad (2.35)$$

We will show that the decomposition (2.35) forms a decomposition of the representation ρ in X_0 .

First,

$$Z = Z(Y_1 + Y_2) + Z(D + C + F) \text{ for } Z = Y_1, Y_2, X_1, X_2 \quad (2.36)$$

because $Z \subseteq Y_1 + Y_2$ and the sum (2.35) is direct. By (2.33) we have

$$X_3 \subseteq X_3(Y_1 + Y_2) + X_3(D + C + F) \subseteq X_3,$$

i.e.,

$$X_3 = X_3(Y_1 + Y_2) + X_3(D + C + F). \quad (2.37)$$

Similarly, by (2.34) we have

$$Y_3 \subseteq Y_3(Y_1 + Y_2) + Y_3(D + C + F) \subseteq Y_3,$$

i.e.,

$$Y_3 = Y_3(Y_1 + Y_2) + Y_3(D + C + F). \quad (2.38)$$

So, by (2.36)–(2.38) $Y_1 + Y_2$ and $D + C + F$ form a decomposition of the representation ρ . Since $\rho(y_1 + y_2) = Y_1 + Y_2$, we have either $\rho(y_1 + y_2) = X_0$ or $\rho(y_1 + y_2) = 0$, i.e., $y_1 + y_2$ is a perfect element. \square

2) If $\rho(y_1 + y_2) = 0$, then $Y_1 = X_1 = Y_2 = X_2 = 0$ and ρ is one of $\rho_{x_0}, \rho_{y_3}, \rho_{x_3}$, see Table A.1. \square

COROLLARY 2.9.4. *The elements $y_1 + y_2 + x_3$ and $y_1 + y_2 + y_3$ are perfect.*

Proof. Let $Y_1 + Y_2 = X_0$. Then $Y_1 + Y_2 + X_3 = X_0$. If $Y_1 + Y_2 = 0$, then, in the indecomposable representation $\rho : D^{2,2,2} \rightarrow \mathcal{L}(X_0)$, we have $X_0 = Y_3 = X_3$. The same for the element $y_1 + y_2 + y_3$.

If $\rho(y_1 + y_2) = 0$, then $Y_1 = X_1 = Y_2 = X_2 = 0$ and ρ is one of $\rho_{x_0}, \rho_{y_3}, \rho_{x_3}$, see Table A.1. \square

2.10. The elementary map φ_i : a way to construct new perfect elements

In this section we list three fundamental properties of the elementary map φ_i . Proposition 2.10.1 gives a way to construct new *perfect* elements from already existing ones. Proposition 2.10.2 motivates the construction of *admissible* sequences and *admissible* elements.

PROPOSITION 2.10.1. *Let z be a perfect element in $D^{2,2,2}$ and, for every indecomposable representation ρ , let*

$$\varphi_i \Phi^+ \rho(z) + \varphi_j \Phi^+ \rho(z) = \rho(u), \text{ where } i \neq j. \quad (2.39)$$

Then u is also the perfect element.

Proof. If $\Phi^+ \rho(z) = 0$, then $\rho(u) = 0$. If $\Phi^+ \rho(z) = X_0^1$, then by Corollary 2.4.4 we have

$$\begin{aligned} \rho(u) &= \varphi_i X_0^1 + \varphi_j X_0^1 = \varphi_i \Phi^+ \rho(I) + \varphi_j \Phi^+ \rho(I) = \\ &Y_i(Y_j + Y_k) + Y_j(Y_i + Y_k) = (Y_i + Y_j)(Y_i + Y_k)(Y_j + Y_k) \end{aligned}$$

and by Proposition 2.9.3 the element u is also perfect. \square

PROPOSITION 2.10.2. *For $\{i, j, k\} = \{1, 2, 3\}$, the following relations hold*

$$\varphi_i \varphi_j \varphi_i + \varphi_i \varphi_k \varphi_i = 0, \quad (2.40)$$

$$\varphi_i^3 = 0. \quad (2.41)$$

Proof. For every vector $v \in X_0^1$, by definition of φ_i we have $(\varphi_i + \varphi_j + \varphi_k)(v) = 0$, see eq. (2.14). In other words, $\varphi_i + \varphi_j + \varphi_k = 0$. Therefore,

$$\varphi_i \varphi_j \varphi_i + \varphi_i \varphi_k \varphi_i = \varphi_i(\varphi_j + \varphi_k) \varphi_i = -\varphi_i^3.$$

So, it suffices to prove that $\varphi_i^3 = 0$. For every $z \subseteq D^{2,2,2}$, by headings (1),(3) and (5) of Corollary 2.4.4 we have

$$\begin{aligned} \varphi_i^3((\Phi^+)^3 \rho)(z) &\subseteq \varphi_i^2[\varphi_i((\Phi^+)^3 \rho)(I)] = \\ \varphi_i^2[(\Phi^+)^2 \rho)(y_i(y_j + y_k))] &\subseteq \varphi_i^2[(\Phi^+)^2 \rho)(y_i)] = \\ \varphi_i[(\Phi^+ \rho)(x_i(y_j + y_k))] &\subseteq \varphi_i[(\Phi^+ \rho)(x_i)] = \varphi_i \Phi^+ \rho(x_i) = 0. \quad \square \end{aligned}$$

COROLLARY 2.10.3. *The relation*

$$\varphi_i \varphi_j \varphi_i(B) = \varphi_i \varphi_k \varphi_i(B) \quad (2.42)$$

takes place for every subspace $B \subseteq X_0^3$, where X_0^3 is the representation space of $(\Phi^+)^3 \rho$.

2.11. List of the admissible sequences

Recall that the *admissible sequences* are introduced in §1.7. Let us construct new admissible sequences acting by the *elementary* map φ_i .

Let the action of φ_i on an admissible sequence α be defined so that

- (1) The index i is added in front of the sequence $\alpha = i_1 i_2 \dots i_n$, i.e.,

$$\varphi_i(i_1 i_2 \dots i_n) = i i_1 i_2 \dots i_n. \quad (2.43)$$

- (2) New sequence $\varphi_i(\alpha)$ should also be an admissible sequence, in other words, $i \neq i_1$.

The rule $i \neq i_1$ in the definition of admissible sequence is motivated by the property $\varphi_i = -(\varphi_j + \varphi_k)$, so that φ_i can be always replaced by $\varphi_j + \varphi_k$.

PROPOSITION 2.11.1. *The admissible sequence starting at $i_1 = 1$ may be transformed to one of the seven types given in Table 2.2, column 1 (similarly for $i_1 = 2$ and $i_1 = 3$, see (1.23)). Here, $n \in \{1, 2, 3, \dots\}$, $m \in \{0, 1, 2, 3, \dots\}$. In case 7, $n \geq 0$. In cases 1 and 4, $m > 0$.*

For $m = 0$ cases 1 and 4 are not given in the table: in case 1 (resp. case 4), action φ_1 transforms $(21)^n$ to $1(21)^n$ (resp. $(31)^n$ to $1(31)^n$).

N	Admissible Sequence	Action φ_1	Action φ_2	Action φ_3
1	$(213)^m(21)^n$	$13(213)^{m-1}(21)^{n+1}$	—	$3(213)^m(21)^n$
2	$3(213)^m(21)^n$	$13(213)^m(21)^n$	$(213)^m(21)^{n+1}$	—
3	$13(213)^m(21)^n$	—	$(213)^{m+1}(21)^n$	$3(213)^m(21)^{n+1}$
4	$(312)^m(31)^n$	$12(312)^{m-1}(31)^{n+1}$	$2(312)^m(31)^n$	—
5	$2(312)^m(31)^n$	$12(312)^m(31)^n$	—	$(312)^m(31)^{n+1}$
6	$12(312)^m(31)^n$	—	$2(312)^m(31)^{n+1}$	$(312)^{m+1}(31)^n$
7	$1(21)^n = 1(31)^n$	—	$(21)^{n+1}$	$(31)^{n+1}$

TABLE 2.2. For $D^{2,2,2}$: the admissible sequences starting at $i_1 = 1$

Proof. It suffices to prove that every action of φ_i on any admissible sequence from the list (Table 2.2, column 1) gives us again an admissible sequence from the same list.

Line 1, action φ_1 . Applying φ_1 to the sequence $(213)^m(21)^n$ we get

$$1(213)^m(21)^n = 1(213)(213)^{m-1}(21)^n.$$

Since $1213(2\dots) = 1321(2\dots)$, we see that $1(213)^m(21)^n = 132[1(213)^{m-1}(21)^n]$. By the induction hypothesis

$$1(213)^{m-1}(21)^n = 13(213)^{m-2}(21)^{n+1},$$

so

$$1(213)^m(21)^n = 132[13(213)^{m-2}(21)^{n+1}] = 13(213)^{m-1}(21)^{n+1},$$

i.e., we get a sequence from Line 3. \square

Line 1, action φ_3 . Applying φ_3 to the sequence $(213)^m(21)^n$ we just get a sequence from Line 2. \square

Line 2, action φ_1 . Action φ_1 on the $3(213)^m(21)^n$ leads to Line 3. \square

Line 2, action φ_2 . Applying φ_2 to the sequence $3(213)^m(21)^n$, we get

$$23(213)^m(21)^n = 23(213)(213)^{m-1}(21)^n.$$

Since $23213\dots = 21323\dots$, we have $23(213)^m(21)^n = 213[23(213)^{m-1}(21)^n]$. Again, by the induction hypothesis

$$23(213)^{m-1}(21)^n = (213)^{m-1}(21)^{n+1},$$

so

$$23(213)^m(21)^n = 213[(213)^{m-1}(21)^{n+1}] = (213)^m(21)^{n+1},$$

i.e., a sequence from Line 1. \square

Line 3, action φ_2 . Applying φ_2 we immediately get Line 1. \square

Line 3, action φ_3 . Apply φ_3 to the sequence $13(213)^m(21)^n$. Again,

$$313(213)^m(21)^n = 313(213)(213)^{m-1}(21)^n = 321[313(213)^{m-1}(21)^n].$$

By the induction hypothesis

$$313(213)^m(21)^n = 321[3(213)^{m-1}(21)^{n+1}] = 3(213)^m(21)^{n+1}. \quad \square$$

Lines 4, 5, 6. Similarly as for Lines 1, 2, 3. \square

Line 7. Applying φ_2 and φ_3 we get Line 1 and Line 4 with $m = 0$, respectively: $(21)^{n+1}$ and $(31)^{n+1}$. \square

2.12. The theorem on the admissible element classes

The lattice polynomials indexed by admissible sequences are also said to be *admissible*. We define admissible elements $f_\alpha, e_\alpha, g_{\alpha 0}$ in Table 2.3. By definition $g_0 = I$.

THEOREM 2.12.1 (On classes of admissible elements). *Let $\alpha = i_n i_{n-1} \dots 1$ be an admissible sequence and $i \neq i_n$. Then $i\alpha$ is admissible and, for $z_\alpha = f_\alpha, e_\alpha, g_{\alpha 0}$ from Table 2.3, the following relation holds:*

$$\varphi_i \Phi^+ \rho(z_\alpha) = \rho(z_{i\alpha}). \quad (2.44)$$

For the proof of Theorem 2.12.1, see Appendix B.

2.13. The inclusion theorem and the cumulative polynomials

THEOREM 2.13.1. *For every admissible sequences $\alpha, \alpha i$, ($i = 1, 2, 3$) from Table 2.2, the following inclusion holds¹*

$$e_{\alpha i} \subseteq g_{\alpha 0}, \quad i = 1, 2, 3.$$

For the proof of Theorem 2.13.1, see §C.1.3.

It is easy to check by the definitions from Table 2.3 that

$$f_\alpha \subseteq e_\alpha, \quad g_{\alpha 0} \subseteq e_\alpha. \quad (2.45)$$

So, from Theorem 2.13.1 (*Inclusion Theorem*) and inclusions 2.45 we get

$$f_{\alpha i} \subseteq e_{\alpha i} \subseteq g_{\alpha 0}. \quad (2.46)$$

¹Please do not confuse polynomials $e_{\alpha i}$ considered here with polynomials $e_{i\alpha}$ considered, for example, in Theorem 2.12.1.

N	α	f_α	e_α	$g_{\alpha 0}$
1	γ	$y_1 y_2 a_q^{13} A_{k-1}^{32}$	$y_2 A_{k-1}^{32} a_q^{21} A_k^{13} a_q^{13}$	$e_\alpha(x_1 + a_q^{32} A_k^{32})$
2	3γ	$y_3(x_1 + x_2) A_q^{23} a_{k-1}^{31}$	$y_3 a_{k-1}^{31} A_{q+1}^{12} a_k^{12} A_q^{23}$	$e_\alpha(y_2 y_3 + A_q^{12} a_k^{31})$
3	13γ	$y_3 y_1 a_{q+1}^{32} A_{k-1}^{21}$	$y_1 A_{k-1}^{21} a_{q+1}^{13} A_k^{32} a_{q+1}^{32}$	$e_\alpha(x_3 + a_{q+1}^{21} A_k^{21})$
4	$(213)\gamma$	$y_2(x_3 + x_1) A_{q+1}^{12} a_{k-1}^{23}$	$y_2 a_{k-1}^{23} A_{q+2}^{31} a_k^{31} A_{q+1}^{12}$	$e_\alpha(y_1 y_2 + A_{q+1}^{31} a_k^{23})$
5	$3(213)\gamma$	$y_2 y_3 a_{q+2}^{21} A_{k-1}^{13}$	$y_3 A_{k-1}^{13} a_{q+2}^{32} A_k^{21} a_{q+2}^{21}$	$e_\alpha(x_2 + a_{q+2}^{13} A_k^{13})$
6	$13(213)\gamma$	$y_1(x_2 + x_3) A_{q+2}^{31} a_{k-1}^{12}$	$y_1 a_{k-1}^{12} A_{q+3}^{23} a_k^{23} A_{q+2}^{31}$	$e_\alpha(y_1 y_3 + A_{q+2}^{23} a_k^{12})$
7	$1(21)^{2k}$	$x_1 A_k^{23} a_k^{23}$	$y_1 A_k^{31} a_k^{12} A_k^{23} a_k^{23}$	$e_\alpha(y_2 a_k^{21} + y_3 a_k^{31})$
8	$1(21)^{2k+1}$	$y_1(x_2 + x_3) A_k^{31} a_k^{12}$	$y_1 A_k^{31} a_k^{12} A_{k+1}^{23} a_{k+1}^{23}$	$e_\alpha(y_1 a_{k+1}^{12} + y_1 a_{k+1}^{13})$
9	β	$y_2 y_3 a_q^{21} A_k^{13}$	$y_2 A_k^{32} a_q^{21} A_k^{13} a_{q+1}^{13}$	$e_\alpha(x_2 + a_q^{13} A_{k+1}^{13})$
10	3β	$x_3 A_{q+1}^{12} a_k^{12}$	$y_3 a_k^{31} A_{q+1}^{12} a_k^{12} A_{q+1}^{23}$	$e_\alpha(y_1 y_3 + A_q^{23} a_{k+1}^{12})$
11	13β	$y_1 y_2 a_{q+1}^{13} A_k^{32}$	$y_1 A_k^{21} a_{q+1}^{13} A_k^{32} a_{q+2}^{32}$	$e_\alpha(x_1 + a_{q+1}^{32} A_{k+1}^{32})$
12	$(213)\beta$	$x_2 A_{q+2}^{31} a_k^{31}$	$y_2 a_k^{23} A_{q+2}^{31} a_k^{31} A_{q+2}^{12}$	$e_\alpha(y_2 y_3 + A_{q+1}^{12} a_{k+1}^{31})$
13	$3(213)\beta$	$y_3 y_1 a_{q+2}^{32} A_k^{21}$	$y_3 A_k^{13} a_{q+2}^{32} A_k^{21} a_{q+3}^{21}$	$e_\alpha(x_3 + a_{q+2}^{21} A_{k+1}^{21})$
14	$13(213)\beta$	$x_1 A_{q+3}^{23} a_k^{23}$	$y_1 a_k^{12} A_{q+3}^{23} a_k^{23} A_{q+3}^{31}$	$e_\alpha(y_1 y_2 + A_{q+2}^{31} a_{k+1}^{23})$

TABLE 2.3. For $D^{2,2,2}$: the admissible elements starting at $i_1 = 1$

In Lines 1–6: $\gamma = (213)^{2p}(21)^{2k}$, $k > 0$, $p \geq 0$. In Lines 7–8: $k \geq 0$.

In Lines 9–14: $\beta = (213)^{2p}(21)^{2k+1}$, $k \geq 0$, $p \geq 0$. In all lines, $q = k + 3p$.

Recall the definition of *cumulative polynomials* from §1.10. *Cumulative polynomials* of length n are constructed as sums of all admissible elements of the same length n , where n is the length of the multi-index:

$$\begin{aligned}
x_t(n) &= \sum f_{i_n \dots i_2 t}, & t = 1, 2, 3, \\
y_t(n) &= \sum e_{i_n \dots i_2 t}, & t = 1, 2, 3, \\
x_0(n) &= \sum g_{i_n \dots i_2 0}.
\end{aligned}$$

From (2.46) and Theorem 2.12.1 the next inclusions take place:

COROLLARY 2.13.2.

$$x_t(n) \subseteq y_t(n) \subseteq x_0(n), \quad t = 1, 2, 3. \quad (2.47)$$

Thus, *cumulative polynomials* satisfy the same inclusions as the corresponding generators in $D^{2,2,2}$.

PROPOSITION 2.13.3 (On the cumulative polynomials). *If $z_t(n)$ is one of the cumulative polynomials $x_t(n), y_t(n) (t = 1, 2, 3)$ or $x_0(n)$, then*

$$\sum_{i=1,2,3} \varphi_i \Phi^+ \rho(z_t(n)) = \rho(z_t(n+1)). \quad (2.48)$$

Proof. By Theorem 2.12.1

$$\varphi_i \Phi^+ \rho(z_\alpha) = \rho(z_{i\alpha}) \text{ for every } z_\alpha = f_\alpha, e_\alpha, g_{\alpha 0}.$$

The latter relation is true only under the condition that $i\alpha$ is *admissible*, i.e., $i \neq i_n$ for $\alpha = i_n i_{n-1} \dots i_1$. By (2.15) in the sum of (2.48) we can always exclude one of the elementary maps $\varphi_1, \varphi_2, \varphi_3$ such that $i \neq i_n$. \square

CHAPTER 3

Perfect polynomials in $D^{2,2,2}$

3.1. The chains $a_i(n) \subseteq b_i(n) \subseteq c_i(n)$. Perfectness and distributivity of $H^+(n)$.

Recall the definition of the elements $a_i(n), b_i(n), c_i(n)$. The perfect elements (see §A.6) are constructed in the following way, see Definition 1.27 from §1.12:

For $n = 0$:

$$\begin{aligned} a_i(0) &= y_j + y_k, \\ b_i(0) &= x_i + y_j + y_k, \\ c_1(0) &= c_2(0) = c_3(0) = \sum y_i. \end{aligned} \tag{3.1}$$

For $n \geq 1$:

$$\begin{aligned} a_i(n) &= x_j(n) + x_k(n) + y_j(n+1) + y_k(n+1), \\ b_i(n) &= a_i(n) + x_i(n+1) = x_j(n) + x_k(n) + x_i(n+1) + y_j(n+1) + y_k(n+1), \\ c_i(n) &= a_i(n) + y_i(n+1) = x_j(n) + x_k(n) + y_i(n+1) + y_j(n+1) + y_k(n+1). \end{aligned}$$

By definition we have

$$a_i(n) \subseteq b_i(n) \subseteq c_i(n). \tag{3.2}$$

The perfectness of the elements $a_i(0), b_i(0), c_i(0)$ is proved in Proposition 2.9.3 and Corollary 2.9.4. The perfectness of the elements $a_i(1), b_i(1), c_i(1)$ is proved in §C.2.1-§C.2.3.

PROPOSITION 3.1.1. *1) For every element $v_i(n) = a_i(n), b_i(n), c_i(n)$, the following relation holds:*

$$\rho(v_i(n)) = \sum_{k=1,2,3} \varphi_k \Phi^+ \rho(v_i(n-1)). \tag{3.3}$$

2) The elements $a_i(n), b_i(n), c_i(n)$ are perfect for every $n \geq 1$ and $i = 1, 2, 3$.

The proof of Proposition 3.1.1 is given in §C.2.4.

Let us consider sublattice $H^+(0)$ generated by 3 chains

$$s_i = \{a_i(n) \subseteq b_i(n)\}, \quad i = 1, 2, 3 \tag{3.4}$$

and sublattice $H^+(n)$ ($n \geq 1$) generated by 3 chains

$$s_i = \{a_i(n) \subseteq b_i(n) \subseteq c_i(n)\}, \quad i = 1, 2, 3. \tag{3.5}$$

According to definition of perfect elements (see §A.6), $H^+(n)$ forms a sublattice of perfect elements in $D^{2,2,2}$.

PROPOSITION 3.1.2. *$H^+(n)$ is a distributive sublattice for every $n \geq 0$.*

A proof of Proposition 3.1.2 is given in §C.2.5.

3.2. The multiplicative form and the cardinality of $H^+(n)$

PROPOSITION 3.2.1. *Every element of the lattice $H^+(n)$, where $n \geq 0$, is of the form $v_1 v_2 v_3$, where $v_i \in \{a_i(n), b_i(n), c_i(n), I_n\}$.*

Proof. Let $v_1 v_2 v_3$ and $v'_1 v'_2 v'_3$ be two elements of $H^+(n)$. Then

$$v_1 v_2 v_3 + v'_1 v'_2 v'_3 = \bigcap_{i=1}^3 \sup(v_i, v'_i). \quad (3.6)$$

Indeed, if $v_i \supseteq v'_i$ for every $i = 1, 2, 3$, then $v_1 v_2 v_3 + v'_1 v'_2 v'_3 = v_1 v_2 v_3$. If $v_1 \supseteq v'_1$ and $v_2 \subseteq v'_2$, $v_3 \subseteq v'_3$, then by (C.13) we have

$$v_1 v_2 v_3 + v'_1 v'_2 v'_3 = v_1 v'_2 v'_3 (v_2 v_3 + v'_1) = v_1 v'_2 v'_3. \quad \square$$

COROLLARY 3.2.2. 1) *The lattice $H^+(0)$ contains ≤ 27 elements.*

2) *The lattice $H^+(n)$ for $n > 0$ contains ≤ 64 elements.*

PROPOSITION 3.2.3. *The lattice $H^+(0)$ contains 27 distinct elements.*

PROPOSITION 3.2.4. *The lattice $H^+(1)$ contains 64 distinct elements.*

For proof of Propositions 3.2.3 and 3.2.4, see §C.2.6.

Proposition 3.2.4 is true for arbitrary n :

PROPOSITION 3.2.5. *The lattice $H^+(n)$ for $n \in \{1, 2, 3, \dots\}$ contains 64 distinct elements.*

Proof. Let $H^+(n)$ contain 64 distinct elements. We will prove that $H^+(n+1)$ also contains 64 distinct elements. Let $v_1(n) v_2(n) v_3(n)$ and $v'_1(n) v'_2(n) v'_3(n)$ be two distinct elements in the $H^+(n)$ and ρ is the preprojective representation separating them. For $n = 1$, the representation ρ is given by Tables C.6, C.7. For example,

$$\begin{aligned} \rho(v_1(n) v_2(n) v_3(n)) &= 0, \\ \rho(v'_1(n) v'_2(n) v'_3(n)) &= X_0^1. \end{aligned} \quad (3.7)$$

Therefore,

$$\text{there exists an } i \text{ such that } \rho(v_i(n)) = 0, \quad (3.8)$$

and

$$\rho(v'_i(n)) = X_0^1 \text{ for every } i. \quad (3.9)$$

For every preprojective representation ρ , there exists preprojective representation $\tilde{\rho}$ such that $\Phi^+ \tilde{\rho} = \rho$. (Naturally, $\tilde{\rho} = \Phi^- \rho$, see §A.7, [BGP73]). Then, by (3.3) for i from (3.8) we have

$$\tilde{\rho}(v_i(n+1)) = \sum_{p=1,2,3} \varphi_p \Phi^+ \tilde{\rho}(v_i(n)) = \sum_{p=1,2,3} \varphi_p \rho(v_i(n)) = 0, \quad (3.10)$$

i.e.,

$$\tilde{\rho}(v_1(n+1) v_2(n+1) v_3(n+1)) = 0. \quad (3.11)$$

On the other hand,

$$\tilde{\rho}(v'_i(n+1)) = \sum_{p=1,2,3} \varphi_p \Phi^+ \tilde{\rho}(v'_i(n)) = \sum_{p=1,2,3} \varphi_p \rho(v'_i(n)) = \sum_{p=1,2,3} \varphi_p X_0^1. \quad (3.12)$$

By Corollary 2.4.4 we have

$$\begin{aligned} \sum_{p=1,2,3} \varphi_p X_0^1 &= \sum_{p=1,2,3} \varphi_p \Phi^+ \tilde{\rho}(I) = \\ \tilde{\rho}(y_1(y_2 + y_3) + y_2(y_1 + y_3) + y_3(y_1 + y_2)) &= \tilde{\rho}\left(\bigcap_{i \neq j} (y_i + y_j)\right). \end{aligned} \quad (3.13)$$

Since $y_i + y_j$ is perfect and $\tilde{\rho} = \Phi^- \rho$ is non-projective, by Proposition 2.9.3 we have

$$\tilde{\rho}(y_i + y_j) = X_0 \text{ for every } \{i, j\}, i \neq j. \quad (3.14)$$

Thus, by (3.12), (3.13) and 3.14

$$\tilde{\rho}(v'_i(n+1)) = X_0 \text{ for every } i, \quad (3.15)$$

and therefore

$$\tilde{\rho}(v'_1(n+1)v'_2(n+1)v'_3(n+1)) = X_0. \quad (3.16)$$

Thus, by (3.11) and (3.16) the representation $\tilde{\rho}$ separates $v_1(n+1)v_2(n+1)v_3(n+1)$ and $v'_1(n+1)v'_2(n+1)v'_3(n+1)$. \square

COROLLARY 3.2.6. $H^+(n)$ is a distributive 64-element lattice for $n \geq 1$. It is the direct product (Fig.1.5) of three chains $\{a_i(n) \subseteq b_i(n) \subseteq c_i(n)\}$.

3.3. The additive form for elements of $H^+(n)$. Considerations mod θ

3.4. Sums and intersections

PROPOSITION 3.4.1. 1) For every perfect elements v_i, v_j, v_k , the following relation holds

$$\sum_{p=1,2,3} \varphi_p \Phi^+ \rho\left(\bigcap_{t=i,j,k} v_t\right) = \bigcap_{t=i,j,k} \left(\sum_{p=1,2,3} \varphi_p \Phi^+ \rho(v_t)\right). \quad (3.17)$$

2) For every perfect element v and every u , the following relation holds

$$\sum_{p=1,2,3} \varphi_p \Phi^+ \rho(vu) = \sum_{p=1,2,3} \varphi_p \Phi^+ \rho(v) \sum_{p=1,2,3} \varphi_p \Phi^+ \rho(u). \quad (3.18)$$

Proof. 1) If $\Phi^+ \rho(v_t) = 0$ for some t , then

$$\Phi^+ \rho\left(\bigcap_{t=i,j,k} v_t\right) = 0 \text{ and } \sum_{p=1,2,3} \varphi_p \Phi^+ \rho(v_t) = 0$$

and (3.17) is true. If $\Phi^+ \rho(v_t) = X_0^1$ for every t , then by Corollary 2.4.4 and (3.13) we have

$$\begin{aligned} \Phi^+ \rho\left(\bigcap_{t=i,j,k} v_t\right) &= X_0^1 = \Phi^+ \rho(I), \\ \sum_{p=1,2,3} \varphi_p \Phi^+ \rho(v_t) &= \rho\left(\bigcap_{i \neq j} (y_i + y_j)\right). \end{aligned} \quad (3.19)$$

Hereafter

$$\bigcap_{i \neq j} (y_i + y_j) \text{ means } (y_1 + y_2)(y_1 + y_3)(y_2 + y_3).$$

From (3.19) we have

$$\begin{aligned} \sum_{p=1,2,3} \varphi_p \Phi^+ \rho\left(\bigcap_{t=i,j,k} v_t\right) &= \sum_{p=1,2,3} \varphi_p \Phi^+ \rho(I) = \rho\left(\bigcap_{i \neq j} (y_i + y_j)\right), \\ \bigcap_{t=i,j,k} \left(\sum_{p=1,2,3} \varphi_p \Phi^+ \rho(v_t)\right) &= \rho\left(\bigcap_{i \neq j} (y_i + y_j)\right), \end{aligned} \quad (3.20)$$

and again (3.17) is true. Note that by Proposition 2.9.3 in (3.20) $\rho(\bigcap(y_i + y_j)) = X_0$ except for the case where ρ is a *projective* representation (Table A.7), for which $\rho(\bigcap(y_i + y_j)) = 0$. \square

2) As in 1), if $\Phi^+\rho(v) = 0$, then

$$\Phi^+\rho(vu) = 0 \text{ and } \sum_{p=1,2,3} \varphi_p \Phi^+\rho(v) = 0$$

and (3.18) is true.

If ρ is *projective*, then $\Phi^+\rho = 0$ and (3.18) is true as well.

Consider the case $\Phi^+\rho \neq 0$, i.e., ρ is not a projective representation. If $\Phi^+\rho(v) = X_0^1$, then $\Phi^+\rho(vu) = \Phi^+\rho(u)X_0^1 = \Phi^+\rho(u)$ and

$$\sum_{p=1,2,3} \varphi_p \Phi^+\rho(vu) = \sum_{p=1,2,3} \varphi_p \Phi^+\rho(u). \quad (3.21)$$

On the other hand,

$$\sum_{p=1,2,3} \varphi_p \Phi^+\rho(v) = \sum_{p=1,2,3} \varphi_p \Phi^+\rho(I) = \rho\left(\bigcap_{i \neq j} (y_i + y_j)\right) \quad (3.22)$$

and by Proposition 2.9.3 we have $\sum \varphi_p \Phi^+\rho(v) = X_0$ and

$$\sum_{p=1,2,3} \varphi_p \Phi^+\rho(v) \sum_{p=1,2,3} \varphi_p \Phi^+\rho(u) = \left(\sum_{p=1,2,3} \varphi_p \Phi^+\rho(u) \right) X_0 = \sum_{p=1,2,3} \varphi_p \Phi^+\rho(u). \quad (3.23)$$

So, (3.18) is true. \square

All considerations given below up to the end of §3.3 are represented *modulo linear equivalence*, see §A.6.

PROPOSITION 3.4.2. *The next relation takes place modulo linear equivalence*

$$x_0(n+2) = \bigcap_{i=1,2,3} a_i(n) \quad \text{mod } \theta. \quad (3.24)$$

Proof. We will prove (3.24) for $n = 0, 1$ without restriction mod θ .

Case $n = 0$. By Table 2.3, Line 7, $k = 0$, we have

$$x_0(2) = g_{10} + g_{20} + g_{30} = \sum_{r,s,t} y_r(y_s + y_t) = \bigcap_{r \neq s} (y_r + y_s). \quad (3.25)$$

The sum is taken over all permutations $\{r, s, t\}$ of $\{1, 2, 3\}$. On the other hand, by Table C.5

$$\bigcap_{i=1,2,3} a_i(0) = \bigcap_{r \neq s} (y_r + y_s). \quad (3.26)$$

Case $n = 1$. We have

$$x_0(3) = g_{210} + g_{120} + g_{310} + g_{130} + g_{230} + g_{320}. \quad (3.27)$$

By Table 2.3, Line 9, $k = 0, p = 0, q = 0$, we have, for example,

$$g_{210} = y_2 a_1^{13}(x_2 + A_1^{13}) = y_2(x_1 + y_3)(x_2 + x_3 + y_1).$$

From here,

$$\begin{aligned} g_{210} + g_{310} &= y_2(x_1 + y_3)(x_2 + x_3 + y_1) + y_3(x_1 + y_2)(x_2 + x_3 + y_1) = \\ &= (x_1 + y_3)(x_1 + y_2)[y_2(x_2 + x_3 + y_1) + y_3(x_2 + x_3 + y_1)] = \\ &= (x_1 + y_3)(x_1 + y_2)[x_2 + y_2(x_3 + y_1) + x_3 + y_3(x_2 + y_1)] = \\ &= (x_1 + y_3)(x_1 + y_2)[(x_3 + y_2)(x_3 + y_1) + (x_2 + y_3)(x_2 + y_1)]. \end{aligned}$$

Thus,

$$\begin{aligned} g_{210} + g_{310} &= (x_1 + y_3)(x_1 + y_2)[(x_3 + y_2)(x_3 + y_1) + (x_2 + y_3)(x_2 + y_1)], \\ g_{120} + g_{320} &= (x_2 + y_3)(x_2 + y_1)[(x_3 + y_1)(x_3 + y_2) + (x_1 + y_3)(x_1 + y_2)], \\ g_{130} + g_{230} &= (x_3 + y_2)(x_3 + y_1)[(x_2 + y_1)(x_2 + y_3) + (x_1 + y_2)(x_1 + y_3)]. \end{aligned} \quad (3.28)$$

On the other hand, by (C.16) we have

$$\begin{aligned} t_1(1) &= x_1(1) + y_1(2) = x_1 + e_{21} + e_{31} = \\ &= x_1 + y_2(x_1 + y_3) + y_3(x_1 + y_2) = (x_1 + y_2)(x_1 + y_3), \end{aligned}$$

i.e.,

$$\begin{aligned} g_{210} + g_{310} &= t_1(1)(t_2(1) + t_3(1)), \\ g_{120} + g_{320} &= t_2(1)(t_1(1) + t_3(1)), \\ g_{130} + g_{230} &= t_3(1)(t_1(1) + t_2(1)), \end{aligned} \quad (3.29)$$

and therefore

$$x_0(3) = \bigcap_{i \neq j} (t_i(1) + t_j(1)) = \bigcap_{k=1,2,3} a_k(1). \quad (3.30)$$

Induction step. As in (3.10),

$$\tilde{\rho}(x_0(n+1)) = \sum_{p=1,2,3} \varphi_p \Phi^+ \tilde{\rho}(x_0(n)) = \sum_{p=1,2,3} \varphi_p \Phi^+ \tilde{\rho} \left(\bigcap_{i=1,2,3} a_i(n-2) \right). \quad (3.31)$$

By (3.17), Proposition 3.4.1

$$\tilde{\rho}(x_0(n+1)) = \bigcap_{i=1,2,3} \left(\sum_{p=1,2,3} \varphi_p \Phi^+ \tilde{\rho} a_i(n-2) \right). \quad (3.32)$$

From here, by (3.3), Proposition 3.1.1

$$\tilde{\rho}(x_0(n+1)) = \bigcap_{i=1,2,3} \tilde{\rho}(a_i(n-1)), \quad (3.33)$$

which is to be proved. \square

COROLLARY 3.4.3. *For $n \geq 2$, the elements $x_0(n)$ are perfect.*

Proof. It follows from the fact that the intersection of perfect elements is also perfect. \square

CONJECTURE 3.4.4. *Relation (3.24) takes place without restriction mod θ .*

For $n = 0$, this conjecture was proven above in (3.25), (3.26) and, for $n = 1$, it was proven in (3.30).

3.5. Other generators: $p_i(n) \subseteq q_i(n) \subseteq s_i(n)$

Let us introduce the following polynomials:

$$\begin{aligned} p_i(n) &= x_0(n+2) + x_i(n+1), \\ q_i(n) &= x_0(n+2) + y_i(n+1), \\ s_i(n) &= x_0(n+2) + y_i(n+1) + x_i(n). \end{aligned} \quad (3.34)$$

Compare two generator systems on Fig. 1.5. The generators $a_i(n) \subseteq b_i(n) \subseteq c_i(n)$ are given by the upper edges of $H^+(n)$. The generators $p_i(n) \subseteq q_i(n) \subseteq s_i(n)$ are given by the lower edges of $H^+(n)$.

PROPOSITION 3.5.1. *Table 3.1 gives relations between the upper and lower generators. These relations are true mod θ .*

N	The multiplicative form	The additive form	The number of Polynomials
1	$a_i(n)$	$s_j(n) + s_k(n)$	3
2	$b_i(n)$	$s_j(n) + s_k(n) + p_i(n)$	3
3	$c_i(n)$	$s_j(n) + s_k(n) + q_i(n)$	3
4	$a_i(n)a_j(n)$	$s_k(n)$	3
5	$b_i(n)b_j(n)$	$s_k(n) + p_i(n) + p_j(n)$	3
6	$c_i(n)c_j(n)$	$s_k(n) + q_i(n) + q_j(n)$	3
7	$a_i(n)b_j(n)$	$s_k(n) + p_j(n)$	6
8	$a_i(n)c_j(n)$	$s_k(n) + p_j(n)$	6
9	$b_i(n)c_j(n)$	$s_k(n) + p_i(n) + q_j(n)$	6
10	$a_i(n)a_j(n)b_k(n)$	$p_k(n)$	3
11	$a_i(n)a_j(n)c_k(n)$	$q_k(n)$	3
12	$b_i(n)b_j(n)a_k(n)$	$p_i(n) + p_j(n)$	3
13	$b_i(n)b_j(n)c_k(n)$	$p_i(n) + p_j(n) + q_k(n)$	3
14	$c_i(n)c_j(n)a_k(n)$	$q_i(n) + q_j(n)$	3
15	$c_i(n)c_j(n)b_k(n)$	$q_i(n) + q_j(n) + p_k(n)$	3
16	$a_i(n)b_j(n)c_k(n)$	$p_i(n) + q_j(n)$	6
17	$a_i(n)a_j(n)a_k(n)$	$x_0(n+2)$	1
18	$b_i(n)b_j(n)b_k(n)$	$p_i(n) + p_j(n) + p_k(n)$	1
19	$c_i(n)c_j(n)c_k(n)$	$q_i(n) + q_j(n) + q_k(n)$	1
20	$I_n = a_i(n) + a_j(n)$ for all $\{i, j\}, i \neq j$	$s_i(n) + s_j(n) + s_k(n)$	1

TABLE 3.1. The multiplicative and additive forms of $H^+(n)$

For a proof, see §C.2.7.

3.6. Coupling together $H^+(n)$ and $H^+(n+1)$. Considerations mod θ

We consider ordering of $H^+(n)$ and $H^+(n+1)$ relative to each other. The way $H^+(n)$ is connected with $H^+(n+1)$ is not as simply as free modular lattice D^r in [GP76], [GP77], see §1.13.1, §1.13.2. For the way $H^+(n)$ is connected with $H^+(n+1)$, see the Hasse diagram on Fig. 1.6.

3.7. Boolean algebras U_n and V_{n+1}

Consider the lower cube $U_n \subseteq H^+(n)$ and the upper cube $V_{n+1} \subseteq H^+(n+1)$:

$$\begin{aligned} U_n &= \{b_1b_2b_3, a_1b_2b_3, b_1a_2b_3, b_1b_2a_3, a_1a_2b_3, a_1b_2a_3, b_1a_2a_3, a_1a_2a_3\}, \\ V_{n+1} &= \{\tilde{c}_1, \tilde{c}_2, \tilde{c}_3, \tilde{c}_1\tilde{c}_2, \tilde{c}_2\tilde{c}_3, \tilde{c}_1\tilde{c}_3, \tilde{c}_1\tilde{c}_2\tilde{c}_3, \tilde{c}_1 + \tilde{c}_2 + \tilde{c}_3\}, \end{aligned} \quad (3.35)$$

where $a_i = a_i(n)$, $b_i = b_i(n)$, $\tilde{c}_i = c_i(n+1)$.

N	The multiplicative form of z	The additive form of z	The complement element z'
1	$I_{U_n} = b_1(n)b_2(n)b_3(n)$	$p_1(n) + p_2(n) + p_3(n)$	$a_1(n)a_2(n)a_3(n)$
2	$a_1(n)b_2(n)b_3(n)$	$p_2(n) + p_3(n)$	$b_1(n)a_2(n)a_3(n)$
3	$b_1(n)a_2(n)b_3(n)$	$p_1(n) + p_3(n)$	$a_1(n)b_2(n)a_3(n)$
4	$b_1(n)b_2(n)a_3(n)$	$p_1(n) + p_2(n)$	$a_1(n)a_2(n)b_3(n)$
5	$a_1(n)a_2(n)b_3(n)$	$p_3(n)$	$b_1(n)b_2(n)a_3(n)$
6	$a_1(n)b_2(n)a_3(n)$	$p_2(n)$	$b_1(n)a_2(n)a_3(n)$
7	$b_1(n)a_2(n)b_3(n)$	$p_1(n)$	$a_1(n)b_2(n)a_3(n)$
8	$O_{U_n} = a_1(n)a_2(n)a_3(n)$	$x_0(n+2)$	$b_1(n)b_2(n)b_3(n)$

TABLE 3.2. The 8-element Boolean algebra U_n

In Table 3.2, the elements I_{U_n} and O_{U_n} are, respectively, the maximal and minimal elements of U_n . The distributivity of U_n follows from the distributivity of $H^+(n)$, Proposition 3.1.2. It is easy to see that U_n is an 8-element Boolean lattice, where

$$\begin{aligned} O_{U_n} &= a_1(n)a_2(n)a_3(n), & I_{U_n} &= b_1(n)b_2(n)b_3(n), \\ (a_ia_jb_k)' &= b_ib_ja_k, \text{ i.e., } a_ia_jb_k + b_ib_ja_k = I_{U_n} \text{ and } a_ia_jb_k \cap b_ib_ja_k = O_{U_n}. \end{aligned} \quad (3.36)$$

In Table 3.3, the elements $I_{V_{n+1}}$ and $O_{V_{n+1}}$ are, respectively, the maximal and minimal elements of V_{n+1} . The distributivity of V_{n+1} follows from the distributivity of $H^+(n+1)$. It is easy to see that V_{n+1} is an 8-element Boolean lattice, where

$$\begin{aligned} O_{V_{n+1}} &= c_1(n+1)c_2(n+1)c_3(n+1), & I_{V_{n+1}} &= c_i(n+1) + c_j(n+1), \\ \tilde{c}'_i &= \tilde{c}_j\tilde{c}_k, \text{ i.e., } \tilde{c}_i + \tilde{c}'_i = \tilde{c}_i + \tilde{c}_j\tilde{c}_k = I_{V_{n+1}}, \text{ and } \tilde{c}_i \cap \tilde{c}'_i = \tilde{c}_i \cap \tilde{c}_j\tilde{c}_k = O_{V_{n+1}}. \end{aligned} \quad (3.37)$$


3.8. The cardinality of $U_n \cup V_{n+1}$

PROPOSITION 3.8.1. *The union $U_n \cup V_{n+1}$ contains 16 distinct elements.*

Consider case $n = 1$. We will prove that $(\Phi^-)^2\rho_{x_0}$ (see Table A.1) separates U_1 and V_2 . It is sufficient to prove that

$$\begin{aligned} (\Phi^-)^2\rho_{x_0}(z) &= X_0 \text{ for every } z \subseteq U_1, \\ (\Phi^-)^2\rho_{x_0}(z) &= 0 \text{ for every } z \subseteq V_2. \end{aligned} \quad (3.38)$$

N	The multiplicative form of z	The additive form of z	The complement element z'
1	$I_{V_{n+1}} = c_i(n+1) + c_j(n+1), i \neq j$	$s_1(n+1) + s_2(n+1) + s_3(n+1)$	$c_1(n+1)c_2(n+1)c_3(n+1)$
2	$c_1(n+1)$	$q_1(n+1) + s_2(n+1) + s_3(n+1)$	$c_2(n+1)c_3(n+1)$
3	$c_2(n+1)$	$s_1(n+1) + q_2(n+1) + s_3(n+1)$	$c_1(n+1)c_3(n+1)$
4	$c_3(n+1)$	$s_1(n+1) + s_2(n+1) + q_3(n+1)$	$c_2(n+1)c_3(n+1)$
5	$c_1(n+1)c_2(n+1)$	$q_1(n+1) + q_2(n+1) + s_3(n+1)$	$c_3(n+1)$
6	$c_1(n+1)c_3(n+1)$	$q_1(n+1) + s_2(n+1) + q_3(n+1)$	$c_2(n+1)$
7	$c_2(n+1)c_3(n+1)$	$s_1(n+1) + q_2(n+1) + q_3(n+1)$	$c_1(n+1)$
8	$O_{V_{n+1}} = c_1(n+1)c_2(n+1)c_3(n+1)$	$q_1(n+1) + q_2(n+1) + q_3(n+1)$	$c_1(n+1) + c_2(n+1) + c_3(n+1)$

TABLE 3.3. The 8-element Boolean algebra V_{n+1} FIGURE 3.1. The Boolean algebras U_n and V_{n+1}

For the generators $a_i(1)$ and $b_i(1)$ ¹, we have

$$(\Phi^-)^2 \rho_{x_0}(a_i(1)) = X_j + X_k + \dots \neq 0.$$

The generators $a_i(1), b_i(1)$ are perfect, i.e.,

$$(\Phi^-)^2 \rho_{x_0}(a_i(1)) = X_0, \quad (\Phi^-)^2 \rho_{x_0}(b_i(1)) = X_0,$$

and therefore (3.38) is true.

On the other hand, since

$$c_i(2) = x_j(2) + x_k(2) + \sum_{q=1,2,3} y_q(3) \quad (3.39)$$

¹The generators $a_i(n)$ and $b_i(n)$ are defined in (1.27); for $n = 1$, these generators are explicitly written in (C.19), (C.20)

and $\Phi^+(\Phi^-)^2\rho_{x_0} = \Phi^-\rho_{x_0}$, we have

$$(\Phi^-)^2\rho_{x_0}(c_i(2)) = \sum_{p=1,2,3} \varphi_p \Phi^-\rho_{x_0}(x_j + x_k + \sum_{q=1,2,3} y_q(2)). \quad (3.40)$$

According to Table A.1 we have $\Phi^-\rho_{x_0}(x_j) = 0$ for every j , i.e.,

$$(\Phi^-)^2\rho_{x_0}(c_i(2)) = \sum_{p=1,2,3} \varphi_p \Phi^-\rho_{x_0}(\sum_{q=1,2,3} y_q(2)). \quad (3.41)$$

By Table A.1 we have $\rho_{x_0}(y_j) = 0$. Further,

$$\Phi^-\rho_{x_0}(y_j(2)) = \sum \varphi_q \Phi^+ \Phi^-\rho_{x_0}(y_j) = \sum \varphi_q \rho_{x_0}(y_j) = 0.$$

Thus, we have $(\Phi^-)^2\rho_{x_0}(c_i(2)) = 0$ and (3.38) is true. The induction step is proved as in Proposition 3.2.5 above. \square

3.9. The Boolean algebra $U_n \cup V_{n+1}$

PROPOSITION 3.9.1. $U_n \cup V_{n+1}$ is a 16-element Boolean algebra.

Proof. According to Proposition A.5.1 it is sufficient to prove relations (A.13), (A.14), (A.15) for

$$c_i = c_i(n+1), \quad d_i = a_i(n)b_j(n)b_k(n) = p_j(n) + p_k(n)$$

(see Fig. 3.1 and Table 3.2), i.e.,

$$c_i(n+1) \subseteq p_j(n) + p_k(n), \quad (3.42)$$

$$p_j(n) + p_k(n) \subseteq c_i(n+1) + p_i(n) + p_k(n), \quad (3.43)$$

$$(p_j(n) + p_k(n))c_j(n+1) \subseteq c_i(n+1), \quad (3.44)$$

where $\{i, j, k\}$ is a permutation of $\{1, 2, 3\}$.

According to the definition of perfect elements from §A.6

$$c_i(n+1) = x_j(n+1) + x_k(n+1) + \sum_{t=1,2,3} y_t(n+2).$$

By definition (3.34)

$$p_j(n) = x_0(n+2) + x_j(n+1),$$

$$p_j(n) + p_k(n) = x_0(n+2) + x_j(n+1) + x_k(n+1).$$

1) Then the inclusion (3.42) follows from

$$\sum_{t=1,2,3} y_t(n+2) \subseteq x_0(n+2), \quad (3.45)$$

which, in turn, comes from the Corollary 2.13.2 concerning *cumulative* polynomials.

2) Inclusion (3.43) follows from the relation

$$x_j(n+1) \subseteq c_i(n+1).$$

3) The last inclusion (3.44) is a little more difficult. It is equivalent to

$$\begin{aligned} (p_j(n) + p_k(n))(x_i(n+1) + x_k(n+1) + \sum_{t=1,2,3} y_t(n+2)) \subseteq \\ x_j(n+1) + x_k(n+1) + \sum_{t=1,2,3} y_t(n+2). \end{aligned} \quad (3.46)$$

From (3.45) and since $x_k(n+1) \subseteq p_j(n)$, we see that (3.46) is equivalent to

$$\begin{aligned} x_k(n+1) + \sum_{t=1,2,3} y_t(n+2) + (p_j(n) + p_k(n))x_i(n+1) \subseteq \\ x_j(n+1) + x_k(n+1) + \sum_{t=1,2,3} y_t(n+2). \end{aligned} \quad (3.47)$$

To prove (3.47), it suffices to check that

$$(p_j(n) + p_k(n))x_i(n+1) \subseteq y_i(n+2), \quad (3.48)$$

i.e.,

$$a_i(n)b_j(n)b_k(n)x_i(n+1) \subseteq y_i(n+2). \quad (3.49)$$

Since $x_i(n+1) \subseteq b_j(n)b_k(n)$ (by relation A.13), we see that (3.49) is equivalent to

$$a_i(n)x_i(n+1) \subseteq y_i(n+2). \quad (3.50)$$

We will prove (3.49) modulo linear equivalence

$$a_i(n)x_i(n+1) \subseteq y_i(n+2) \pmod{\theta}. \quad (3.51)$$

CONJECTURE 3.9.2. *Relation (3.51) takes place without restriction mod θ .*

So, let us prove (3.51). Let $n=1, i=1$. We have to prove that

$$a_1(1)x_1(2) \subseteq y_1(3). \quad (3.52)$$

According to (1.27) or (C.19), (C.20) we have

$$\begin{aligned} a_1(1) = & x_2 + x_3 + y_3(x_2 + y_1) + y_1(x_2 + y_3) + y_1(x_3 + y_2) + y_2(x_3 + y_1) = \\ & (x_2 + y_3)(x_2 + y_1) + (x_3 + y_2)(x_3 + y_1). \end{aligned}$$

Further, according to §1.11, we have

$$x_1(2) = f_{21} + f_{31} = y_2y_3$$

and

$$\begin{aligned} y_1(3) = & y_3(y_1 + x_2)(y_2 + x_3) + y_1(x_2 + y_3)(x_3 + y_2) + y_2(y_1 + x_3)(y_3 + x_2) = \\ & (x_2 + y_3)(x_3 + y_2)[y_3(y_1 + x_2)(y_2 + x_3) + y_1 + y_2(y_1 + x_3)(y_3 + x_2)] = \\ & (x_2 + y_3)(x_3 + y_2)[(y_1 + x_2)(x_3 + y_2y_3) + y_1 + (y_1 + x_3)(x_2 + y_2y_3)] = \\ & (x_2 + y_3)(x_3 + y_2)[(y_1 + x_2)(x_3 + y_1 + y_2y_3) + (y_1 + x_3)(x_2 + y_1 + y_2y_3)] = \\ & (x_2 + y_3)(x_3 + y_2)(x_3 + y_1 + y_2y_3)(x_2 + y_1 + y_2y_3)(y_1 + x_2 + x_3). \end{aligned}$$

Now, (3.52) is true since

$$x_1(2) = y_2y_3 \subseteq (x_2 + y_3)(x_3 + y_2)(x_3 + y_1 + y_2y_3)(x_2 + y_1 + y_2y_3)$$

and

$$a_1(1) = (x_2 + y_3)(x_2 + y_1) + (x_3 + y_2)(x_3 + y_1) \subseteq y_1 + x_2 + x_3.$$

Therefore, (3.52) is also true mod θ , i.e., (3.51) is true for $n=1$.

Induction step. Let (3.51) be true for n , i.e.,

$$\Phi^+\rho(a_i(n)x_i(n+1)) \subseteq \Phi^+\rho(y_i(n+2)) \quad (3.53)$$

for every indecomposable representation ρ . Applying $\sum \varphi_i$ we have

$$\sum_{p=1,2,3} \varphi_p \Phi^+ \rho(a_i(n)x_i(n+1)) \subseteq \sum_{p=1,2,3} \varphi_p \Phi^+ \rho(y_i(n+2)). \quad (3.54)$$

Since $a_i(n)$ is perfect, we have according to (3.18) and Proposition 3.4.1

$$\sum_{p=1,2,3} \varphi_i \Phi^+ \rho(a_i(n)) \sum_{p=1,2,3} \varphi_i \Phi^+ \rho(x_i(n+1)) \subseteq \sum_{p=1,2,3} \varphi_i \Phi^+ \rho(y_i(n+2)). \quad (3.55)$$

By Propositions 2.13.3 and 3.1.1

$$\rho(a_i(n+1)x_i(n+2)) \subseteq \rho(y_i(n+3)). \quad (3.56)$$

So, the induction step and Proposition 3.9.1 are proven. \square

COROLLARY 3.9.3 (Connection edges). *The following 8 inclusions hold:*

$$c_i(n+1) \subseteq a_i(n)b_j(n)b_k(n), \quad i = 1, 2, 3, \quad (3.57)$$

$$c_i(n+1)c_j(n+1) \subseteq a_i(n)a_j(n)b_k(n), \quad k = 1, 2, 3, \quad (3.58)$$

$$\bigcap_{i=1,2,3} c_i(n+1) \subseteq \bigcap_{i=1,2,3} a_i(n), \quad (3.59)$$

$$\sum_{i=1,2,3} c_i(n+1) \subseteq \bigcap_{i=1,2,3} b_i(n). \quad (3.60)$$

The 8 inclusions (3.57) – (3.60) correspond to the 8 edges shown on Fig. 1.7 or Fig. A.3.

Proof. Inclusions (3.57) follow from (3.42), inclusions (3.58), (3.59) and (3.60) follows from (3.57), see Table 3.1. \square

3.10. The theorem on perfect elements. Considerations mod θ

We will prove the theorem describing connection construction of sublattices $H^+(n)$.

PROPOSITION 3.10.1. 1) For every $u \in U_n$ and every $v \in H^+(n+1)$, we have $u+v \in U_n$.
2) For every $u \in H^+(n)$ and every $v \in V_{n+1}$, we have $uv \in V_{n+1}$.

Proof. 1) According to Table 3.2, the elements of U_n are

$$p_i(n), \quad p_i(n) + p_j(n), \quad \sum p_i(n), \quad x_0(n+2). \quad (3.61)$$

It is sufficient to verify heading 1) for the generators $p_i(n+1), q_i(n+1), s_i(n+1)$ of $H^+(n+1)$. According to definition (3.34) it is sufficient to verify that adding the elements

$$x_0(n+3), \quad x_i(n+2), \quad y_i(n+2)$$

does not lead out of the elements of type (3.61). According to (2.45) and Corollary (2.13.2) we have

$$x_0(n+3) \subseteq y_i(n+2) \subseteq x_0(n+2) \quad (3.62)$$

and adding the element $x_0(n+3)$ does not lead out of U_n . From (3.62) it follows that

$$p_i(n) = x_0(n+2) + x_i(n+1) \supseteq y_i(n+2) \supseteq x_i(n+2). \quad (3.63)$$

Therefore adding $y_i(n+2)$ and $x_i(n+2)$ does not lead out of U_n . Finally,

$$p_i(n) \supseteq x_i(n+1), \quad p_j(n) + x_i(n+1) = p_i(n) + p_j(n), \quad (3.64)$$

i.e., addition $x_i(n+1)$ does not go lead of the U_n either. \square

2) Every $u \in H^+(n)$ is of the form $u_1 u_2 u_3$, where (Fig. 1.5)

$$u_i \in \{a_i(n) \subseteq b_i(n) \subseteq c_i(n) \subseteq I\}.$$

If $u_i \neq a_i(n)$ for every i , then $u \supseteq \bigcap b_i(n) \supseteq \sum c_i(n+1) \supseteq v$ and $uv = v$. So, it is sufficient to consider the elements $u \in H^+(n)$ of the type $a_i(n)u'$, $a_i(n)a_j(n)u'$ and $a_i(n)a_j(n)a_k(n)$, where $u' \supseteq v$, i.e., it is sufficient to prove that

$$va_i(n) \in V_{n+1} \text{ for every } v \subseteq V_{n+1}. \quad (3.65)$$

By Corollary 3.9.3 we have $c_i(n+1) \subseteq a_i(n)$ and

$$c_i(n+1)a_i(n) = c_i(n+1), \quad (3.66)$$

and so (3.65) is proven for $v = c_i(n+1)$.

Now we consider the case $i \neq j$, for example, $a_2(n)c_1(n+1)$. Further, we have

$$\begin{aligned} c_1(n+1) &= x_3(n+1) + x_2(n+1) + \sum_{i=1,2,3} y_i(n+2), \\ x_3(n+1) &\subseteq a_2(n), \\ \sum_{i=1,2,3} y_i(n+2) &\subseteq x_0(n+2) = \bigcap_{i=1,2,3} a_i(n) \subseteq a_2(n). \end{aligned} \quad (3.67)$$

The last relation follows from the (3.24) and it is true mod θ . Therefore

$$c_1(n+1)a_2(n) = x_3(n+1) + \sum_{i=1,2,3} y_i(n+2) + x_2(n+1)a_2(n). \quad (3.68)$$

According to (3.51) we have $x_2(n+1)a_2(n) \subseteq c_2(n+1)$ and

$$\begin{aligned} c_1(n+1)a_2(n) &\subseteq x_3(n+1) + \sum_{i=1,2,3} y_i(n+2) + x_2(n+1)c_2(n+1) = \\ c_2(n+1)[x_3(n+1) + \sum_{i=1,2,3} y_i(n+2) + x_2(n+1)] &= \\ c_2(n+1)c_1(n+1) &\subseteq a_2(n)c_1(n+1). \end{aligned}$$

Thus,

$$c_1(n+1)a_2(n) = c_1(n+1)c_2(n+1) \quad (3.69)$$

and (3.65) is proven for $v = c_j(n+1)$, $i \neq j$. Finally, by (3.66), (3.69) we have

$$\begin{aligned} a_j(n) \sum_{i=1,2,3} c_i(n+1) &= a_j(n)[c_i(n+1) + c_j(n+1)] = \\ c_j(n+1) + c_i(n+1)a_j(n) &= c_j(n+1) + c_i(n+1)c_j(n+1), \end{aligned}$$

i.e.,

$$a_j(n) \sum_{i=1,2,3} c_i(n+1) = c_j(n+1). \quad \square \quad (3.70)$$

THEOREM 3.10.2. *The union $\bigcup_{n=0}^{\infty} H^+(n)$ is a distributive lattice mod θ . The diagram H^+ is obtained by uniting the diagrams of $H^+(n)$ for $n \geq 0$ and joining the cubes U_n and V_{n+1} for all $n \geq 0$, i.e., it is necessary to draw 8 additional edges for all $n \geq 0$. (Fig. 1.7).*

Proof. We have to show that the *sum* and the *intersection* do not lead out of H^+ . The distributivity follows from the absence of diamonds M_3 , see Proposition A.4.2, [Bir48]. If

$$u \in H^+(n), \quad v \in H^+(m)$$

and these lattices are not adjacent, i.e., $n \leq m - 2$, then $v \subseteq u$. Indeed, by Proposition 3.9.1 (Fig. 1.7) we have

$$v \subseteq \sum_{i=1,2,3} c_i(m) \subseteq \bigcap_{i=1,2,3} b_i(m-1) \subseteq \bigcap_{i=1,2,3} c_i(m-1) \subseteq \bigcap_{i=1,2,3} a_i(m-2) \subseteq u.$$

So, we consider only the case $u \in H^+(n), v \in H^+(n+1)$. Every element $u \in H^+(n)$ is of the form $u_1 + u_2$, where $u_1 \in H^+(n)$ and $u_2 \in U_n$. By Proposition 3.10.1 we have

$$u + v = u_1 + u_2 + v = u_1 + (u_2 + v) \in H^+(n). \quad (3.71)$$

Similarly, every element v from $H^+(n+1)$ is of the form $v_1 v_2$, where $v_1 \in H^+(n+1)$ and $v_2 \in V_{n+1}$. By Proposition 3.10.1

$$vu = v_1 v_2 u = v_1 (v_2 u) \in H^+(n+1). \quad (3.72)$$

So, Theorem 3.10.2 follows from (3.71) and (3.72). This concludes the proof. \square

CHAPTER 4

Atomic and admissible polynomials in D^4

4.1. Admissible sequences in D^4

For definition of admissible sequences in the case of the modular lattice D^4 , see §1.7.2. Essentially, the fundamental property of this definition is

$$ijk = ilk \quad \text{for all } \{i, j, k, l\} = \{1, 2, 3, 4\} \quad (4.1)$$

Relation (4.1) is our main tool in all further calculations of admissible sequences of D^4 .

Without loss of generality only sequences starting at 1 can be considered. The following proposition will be used for the classification of admissible sequences in D^4 .

PROPOSITION 4.1.1. *The following relations hold*

- 1) $(31)^r(32)^s(31)^t = (32)^s(31)^{r+t}$,
- 2) $(31)^r(21)^s(31)^t = (31)^{r+t}(21)^s$,
- 3) $(42)^r(41)^s = (41)^r(31)^s$, $s \geq 1$,
- 4) $2(41)^r(31)^s = 2(31)^{s+1}(41)^{r-1}$, $r \geq 1$,
- 5) $(43)^r(42)^s(41)^t = (41)^r(21)^s(31)^t$,
- 6) $1(41)^r(21)^s = 1(21)^s(41)^r$, $1(i1)^r(j1)^s = 1(j1)^s(i1)^r$, $i, j \in \{2, 3, 4\}$, $i \neq j$,
- 7) $(41)^r(21)^t(31)^s = (41)^r(31)^s(21)^t$,
- 8) $(13)^s(21)^r = (12)^r(31)^s$,
- 9) $12(41)^r(31)^s(21)^t = (14)^r(31)^{s+1}(21)^t = (14)^r(21)^{t+1}(31)^s$,
- 10) $12(14)^r(31)^s(21)^t = (14)^r(31)^s(21)^{t+1}$,
- 11) $13(14)^r(31)^s(21)^t = (14)^r(31)^{s+2}(21)^{t-1}$,
- 12) $32(14)^r(31)^s(21)^t = (31)^s(21)^{t+1}(41)^r = 34(14)^r(31)^s(21)^t$,
- 13) $42(14)^r(31)^s(21)^t = (41)^s(21)^{t+1}(31)^r = 43(14)^r(31)^s(21)^t$,
- 14) $23(14)^r(31)^s(21)^t = (21)^{t+1}(31)^s(41)^r = 24(14)^r(31)^s(21)^t$.

Proof. 1) For $r = 1$, we have

$$\begin{aligned} (31)(32)^s(31)^t &= 31(32)(32)\dots(32)(32)(31)^t = 31(41)(32)\dots(32)(32)(31)^t = \\ &= 31(41)(41)\dots(41)(41)(31)^t = 32(41)(41)\dots(41)(41)(31)^t = 32(32)(41)\dots(41)(41)(31)^t = \\ &= 32(32)(32)\dots(32)(31)(31)^t = (32)^s(31)^{t+1}. \end{aligned}$$

Applying induction on r we get the relation

$$(31)^r(32)^s(31)^t = (32)^s(31)^{r+t}. \quad \square$$

2) For $t = 1$, we have

$$\begin{aligned} (31)^r(21)^s(31) &= (31)^r(21)(21)\dots(21)(21)31 = (31)^r(21)(21)\dots(21)(34)31 = \\ &= (31)^r(21)(21)\dots(34)(34)31 = (31)^r(34)(34)\dots(34)(34)21 = (31)^r(34)(34)\dots(34)(21)21 = \\ &= (31)^r(34)(34)\dots(21)(21)21 = (31)^r(31)(21)\dots(21)(21)21 = (31)^{r+1}(21)^s. \end{aligned}$$

Applying induction on t we get the relation

$$(31)^r(21)^s(31)^t = (31)^{r+t}(21)^s. \quad \square$$

3) For $s = 1$, we have

$$\begin{aligned} (42)^r 41 &= 42(42)(42)\dots(42)(42)41 = 42(42)(42)\dots(42)(42)31 = \\ &= 42(42)(42)\dots(42)(31)31 = 42(31)(31)\dots(31)(31)31 = \\ &= 41(31)(31)\dots(31)(31)31 = 41(31)^r. \end{aligned}$$

Thus, by heading 2) we have

$$(42)^r (41)^s = 41(31)^r (41)^{s-1} = (41)^s (31)^r. \quad \square$$

4) For $s = 0$, we have

$$2(41)^r = 2(41)(41)^{r-1} = 2(31)(41)^{r-1}$$

and by heading 2):

$$2(41)^r (31)^s = 2(31)(41)^{r-1} (31)^s = 2(31)^{s+1} (41)^{r-1}. \quad \square$$

5) Applying heading 3) we get

$$(43)^r (42)^s (41)^t = (43)^r (41)^s (31)^t.$$

Again, applying heading 3) to $(43)^r (41)^s$ we get

$$(43)^r (41)^s = (41)^r (21)^s$$

and

$$(43)^r (42)^s (41)^t = (41)^r (21)^s (31)^t. \quad \square$$

6) For $s = 1$, we have

$$\begin{aligned} 1(41)^r (21) &= 1(41)(41)\dots(41)(41)(21) = 1(41)(41)\dots(41)(23)(21) = \\ &= 1(41)(41)\dots(23)(23)(21) = 1(23)(23)\dots(23)(23)(21) = \\ &= 1(23)(23)\dots(23)(23)(41) = 1(23)(23)\dots(23)(41)(41) = \\ &= 1(23)(41)\dots(41)(41)(41) = 1(21)(41)\dots(41)(41)(41) = \\ &= 1(21)(41)^r. \end{aligned}$$

Thus, by heading 2) we have

$$1(41)^r (21)^s = 1(21)(41)^r (21)^{s-1} = 1(21)^s (41)^r. \quad \square$$

7) Follows from 6). \square

8) First,

$$1321 = 1421 = 1431 = 1231 = 1241 = 1341 \tag{4.2}$$

For $r = 1$, by (4.2) we have

$$\begin{aligned} 13(21)^s &= 13(21)(21)\dots(21)(21) = 12(31)(21)(21)\dots(21)(21)(21) = \\ &= 12(34)(21)(21)\dots(21)(21)(21) = 12(34)(34)(21)\dots(21)(21)(21) = \\ &= 12(34)(34)(34)\dots(34)(34)(31) = 12(12)(12)(12)\dots(12)(34)(31) = \\ &= 12(12)(12)(12)\dots(12)(12)(31) = (12)^s (31). \end{aligned}$$

Suppose

$$(13)^r (21)^s = (12)^s (31)^r,$$

then we have

$$\begin{aligned}
 (13)^{r+1}(21)^s &= 13(12)^s(31)^r = 13(12)(12)\dots(12)(12)(31)^r = \\
 &13(43)(43)\dots(43)(12)(31)^r = 13(43)(43)\dots(43)(41)(31)^r = \\
 &12(43)(43)\dots(43)(41)(31)^r = 12(12)(43)\dots(43)(41)(31)^r = \\
 &12(12)(12)\dots(12)(41)(31)^r = 12(12)(12)\dots(12)(31)(31)^r = (12)^s(31)^{r+1}.
 \end{aligned}$$

Thus, by induction the following relation holds:

$$(13)^r(21)^s = (12)^s(31)^r \quad \square$$

9) First of all,

$$\begin{aligned}
 12(41)^r(31)^s(21)^t &= 12(41)(41)\dots(41)(41)(31)^s(21)^t = \\
 12(32)(41)\dots(41)(41)(31)^s(21)^t &= 12(32)(32)\dots(32)(31)(31)^s(21)^t = \\
 14(14)(32)\dots(32)(31)(31)^s(21)^t &= 14(14)(14)\dots(14)(31)(31)^s(21)^t = \\
 (14)^r(31)^{s+1}(21)^t.
 \end{aligned}$$

By heading 4) we have

$$4(31)^{s+1}(21)^t = 4(21)^{t+1}(31)^s.$$

Thus,

$$(14)^r(31)^{s+1}(21)^t = (14)^r(21)^{t+1}(31)^s. \quad \square$$

10) First, we have

$$12(14)^r(31) = (14)^r(21)^2 \tag{4.3}$$

since

$$\begin{aligned}
 12(14)^r(31) &= 12(14)(14)\dots(14)(14)(31) = 12(32)(14)\dots(14)(14)(31) = \\
 12(32)(32)\dots(32)(34)(31) &= 14(14)(32)\dots(32)(34)(31) = \\
 14(14)(14)\dots(14)(34)(21) &= 14(14)(14)\dots(14)(21)(21) = (14)^r(21)^2.
 \end{aligned}$$

By (4.3) we have

$$12(14)^r(31)^s(21)^t = 12(14)^r(31)(31)^{s-1}(21)^t = (14)^r(21)^2(31)^{s-1}(21)^t.$$

By headings 2) and 4):

$$(21)^2(31)^{s-1}(21)^t = (21)^{t+2}(31)^{s-1}$$

and

$$12(14)^r(31)^s(21)^t = (14)^r(21)^{t+2}(31)^{s-1} = (14)^r(31)^s(21)^{t+1}. \quad \square$$

11) Applying permutation $2 \leftrightarrow 3$ to heading 10), we get

$$13(14)^r(21)^t(31)^s = (14)^r(21)^t(31)^{s+1}.$$

By heading 4)

$$13(14)^r(21)^t(31)^s = (14)^r(31)^{s+2}(21)^{t-1}. \quad \square$$

12) By heading 8)

$$(14)^r(31)^s = (13)^s(41)^r$$

and

$$\begin{aligned} 32(14)^r(31)^s(21)^t &= 32(13)^s(41)^r(21)^t = \\ 32(13)(13)\dots(13)(13)(41)^r(21)^t &= 34(24)(13)\dots(13)(13)(41)^r(21)^t = \\ 34(24)(24)\dots(24)(21)(41)^r(21)^t &= 31(24)(24)\dots(24)(21)(41)^r(21)^t = \\ 31(31)(31)\dots(31)(21)(41)^r(21)^t &= (31)^s(21)(41)^r(21)^t. \end{aligned}$$

By heading 2) we have

$$(31)^s(21)(41)^r(21)^t = (31)^s(21)^{t+1}(41)^r$$

and

$$32(14)^r(31)^s(21)^t = (31)^s(21)^{t+1}(41)^r \quad \square$$

13) By heading 8):

$$(14)^r(31)^s = (13)^s(41)^r$$

and

$$42(14)^r(31)^s(21)^t = 42(13)^s(41)^r(21)^t.$$

Applying permutation $4 \leftrightarrow 3$ to heading 12) we get

$$42(13)^r(41)^s(21)^t = (41)^s(21)^{t+1}(31)^r. \quad \square$$

14) By heading 4) we have

$$23(14)^r(31)^s(21)^t = 23(14)^r(21)^{t+1}(31)^{s-1}.$$

Applying permutation $2 \leftrightarrow 3$ to heading 12) we get

$$23(14)^r(21)^{t+1}(31)^{s-1} = (21)^{t+1}(31)^s(41)^r. \quad \square$$

PROPOSITION 4.1.2. *Full list of admissible sequences starting at 1 is given by Table 4.1.*

Note to Table 4.1. Type Fij (resp. Gij , Hij) denotes the admissible sequence starting at j and ending at i . Sequences of type Fij and $H11$ contain an even number of symbols, sequences of type Gij and Hij , ($i > 1$) contain an odd number of symbols. For differences in types Fij , Gij , Hij , see the table.

Proof. It suffices to prove that maps φ_i , where $i = 1, 2, 3, 4$, do not lead out of Table 4.1. The exponents r, s, t may be any non-negative integer number. The proof is based on the relations from Proposition 4.1.1. We refer only to number of relation and drop reference to Proposition 4.1.1 itself.

Lines F21–F41. Consider, for example, φ_3 . By heading 4) we get

$$\varphi_3((21)^t(41)^r(31)^s) = 3(41)^{r+1}(21)^{t-1}(31)^s, \quad (4.4)$$

i.e., we get $G31$. \square

Line G11. Consider the action φ_2 . By heading 6) we get

$$\varphi_2(1(41)^r(31)^s(21)^t) = 2(1(41)^r(21)^t(31)^s) = 21(41)^r(21)^t(31)^s.$$

By heading 2) we have

$$\varphi_2(1(41)^r(31)^s(21)^t) = 21(41)^r(21)^t(31)^s = (21)^{t+1}(41)^r(31)^s, \quad (4.5)$$

i.e., we get $F21$. \square

Line G21. By heading 9) applying φ_1 we get $H11$. For the action φ_3 , we have

$$32(41)^r(31)^s(21)^t = 31(41)^r(31)^s(21)^t = (31)^{s+1}(41)^r(21)^t, \quad (4.6)$$

	Admissible Sequence	Action φ_1	Action φ_2	Action φ_3	Action φ_4
$F21$	$(21)^t(41)^r(31)^s = (21)^t(31)^s(41)^r$	$G11$	-	$G31$	$G41$
$F31$	$(31)^s(41)^r(21)^t = (31)^s(21)^t(41)^r$	$G11$	$G21$	-	$G41$
$F41$	$(41)^r(31)^s(21)^t = (41)^r(21)^t(31)^s$	$G11$	$G21$	$G31$	-
$G11$	$1(41)^r(31)^s(21)^t =$ $1(31)^s(41)^r(21)^t =$ $1(21)^t(31)^s(41)^r$	-	$F21$	$F31$	$F41$
$G21$	$2(41)^r(31)^s(21)^t = 2(31)^{s+1}(41)^{r-1}(21)^t$	$H11$	-	$F31$	$F41$
$G31$	$3(41)^r(21)^t(31)^s = 3(21)^{t+1}(41)^{r-1}(31)^s$	$H11$	$F21$	-	$F41$
$G41$	$4(21)^t(31)^s(41)^r = 4(31)^{s+1}(21)^{t-1}(41)^r$	$H11$	$F21$	$F31$	-
$H11$	$(14)^r(31)^s(21)^t = (14)^r(21)^{t+1}(31)^{s-1} =$ $(13)^s(41)^r(21)^t = (13)^s(21)^{t+1}(41)^{r-1} =$ $(12)^{t+1}(41)^r(31)^{s-1} = (12)^{t+1}(31)^{s-1}(41)^r$	-	$H21$	$H31$	$H41$
$H21$	$2(14)^r(31)^s(21)^t$	$H11$	-	$F31$	$F41$
$H31$	$3(14)^r(31)^s(21)^t$	$H11$	$F21$	-	$F41$
$H41$	$4(14)^r(31)^s(21)^t$	$H11$	$F21$	$F31$	-

TABLE 4.1. Admissible sequences for the modular lattice D^4

i.e., we get $F31$. For φ_4 from heading 4), we have

$$\begin{aligned}
 42(41)^r(31)^s(21)^t &= 42(31)^{s+1}(41)^{r-1}(21)^t = \\
 41(31)^{s+1}(41)^{r-1}(21)^t &= (41)^r(31)^{s+1}(21)^t,
 \end{aligned} \tag{4.7}$$

i.e., we get $F41$. For φ_1 from heading 9), we have

$$\begin{aligned}
 12(41)^r(31)^s(21)^t &= (14)^r(31)^{s+1}(21)^t = \\
 (14)^r(21)^{t+1}(31)^s,
 \end{aligned} \tag{4.8}$$

i.e., we get $H11$. \square

Line G31. For the action φ_1 by relation $13(41)^t = 12(41)^t$ we get $H11$. For the action φ_2 and by heading 6), we have

$$23(41)^r(31)^s(21)^t = 23(41)^r(21)^t(31)^s = 21(41)^r(21)^t(31)^s = (21)^{t+1}(41)^r(31)^s$$

i.e. we get $F21$.

For the action φ_4 from headings 6), 4), we have

$$43(41)^r(31)^s(21)^t = 43(41)^r(21)^t(31)^s = 43(21)^{t+1}(41)^{r-1}(31)^s = 41(21)^{t+1}(41)^{r-1}(31)^s$$

So, by heading 2) we have

$$43(41)^r(31)^s(21)^t = (41)^r(21)^{t+1}(31)^s,$$

i.e., we get $F41$. \square

Line G41. By heading 9) we get $H11$. For action φ_2 we have from 4)

$$24(21)^t(31)^s(41)^r = 24(31)^{s+1}(21)^{t-1}(41)^r = 21(31)^{s+1}(21)^{t-1}(41)^r = (21)^t(31)^{s+1}(41)^r$$

i.e., we get $F21$. For the action φ_3 we have from 2)

$$34(21)^t(31)^s(41)^r = 31(21)^t(31)^s(41)^r = (31)^{s+1}(21)^t(41)^r$$

i.e., we get $F31$ \square

Line H11. Actions are rather trivial. \square

Line H21. For action φ_1 the relation follows from heading 10), for φ_3 – follows from heading 12), and for φ_4 – follows from heading 13). \square

Line H31. Follows from headings 9), 14) and 13). \square

Line H41. Action φ_1 is trivial since $14(14)^r = (14)^{r+1}$. Action φ_2 follows from heading 14) and action φ_3 follows from 12). \square

Thus, all cases are considered and the proposition is proved. \square



The pyramid on the Fig. 1.4 has internal points. We consider the slice $S(n)$ containing all sequences of the same length n . The slices $S(3)$ and $S(4)$ are shown on the Fig. 4.1. The slice $S(4)$ contains only one internal point

$$14(21) = 13(21) = 13(41) = 12(41) = 14(31) = 12(31) \quad (4.9)$$

The slices $S(4)$ and $S(5)$ are shown on the Fig. 4.2. The slice $S(5)$ contains 3 internal points

$$\begin{aligned} 2(31)(21) &= 2(41)(21), \\ 3(21)(31) &= 3(41)(31), \\ 4(21)(41) &= 4(31)(41). \end{aligned} \quad (4.10)$$

REMARK 4.1.3. The slice $S(n)$ contains $\frac{1}{2}n(n+1)$ different admissible sequences. Actions of φ_i , where $i = 1, 2, 3, 4$ move every line in the triangle $S(n)$, which is parallel to some edge of the triangle, to the edge of $S(n+1)$. The edge containing k points is moved to $k+1$ points in the $S(n+1)$.

FIGURE 4.1. Slices of admissible sequences for D^4 , $l = 3$ and $l = 4$ FIGURE 4.2. Slices of admissible sequences for D^4 , $l = 4$ and $l = 5$

4.2. Atomic polynomials and elementary maps in the modular lattice D^4

The free modular lattice D^4 is generated by 4 generators:

$$D^4 = \{e_1, e_2, e_3, e_4\}.$$

Recall, that *atomic* lattice polynomials a_n^{ij} , where $i, j \in \{1, 2, 3, 4\}$, $n \in \{0, 1, 2, 3, \dots\}$, for the case of D^4 , are defined as follows

$$a_n^{ij} = \begin{cases} a_n^{ij} = I & \text{for } n = 0, \\ a_n^{ij} = e_i + e_j a_{n-1}^{kl} = e_i + e_j a_{n-1}^{lk} & \text{for } n \geq 1, \end{cases}$$

where $\{i, j, k, l\}$ is the permutation of $\{1, 2, 3, 4\}$, see §1.5.

PROPOSITION 4.2.1. 1) *The following property of the atomic elements take place*

$$e_j a_n^{kl} = e_j a_n^{lk} \text{ for } n \geq 1, \text{ and distinct indices } j, k, l. \quad (4.11)$$

2) *The definition of the atomic elements a_n^{ij} in (1.17) is well-defined.*

3) *We have*

$$a_n^{ij} \subseteq a_{n-1}^{ij} \subseteq \dots \subseteq a_2^{ij} \subseteq a_1^{ij} \subseteq a_0^{ij} = I \text{ for all } i \neq j. \quad (4.12)$$

4) *To equalize the lower indices of the admissible polynomials $f_{\alpha 0}$ (see Table 4.3 and Theorem 4.8.1) we will use the following relation:*

$$e_j + e_i a_{t+1}^{jk} a_{s-1}^{lk} = e_j + e_k a_t^{ij} a_s^{il} \quad \text{for all } \{i, j, k, l\} = \{1, 2, 3, 4\}. \quad (4.13)$$

Proof. 1) Suppose (4.11) is true for the index $n - 1$:

$$e_j a_{n-1}^{kl} = e_j a_{n-1}^{lk}.$$

Since $e_j \subseteq a_{n-1}^{ji}$, by the permutation property (A.4), we have

$$e_j a_n^{kl} = e_j (e_k + e_l a_{n-1}^{ij}) = e_j (e_k + e_l a_{n-1}^{ji}) = e_j (e_l + e_k a_{n-1}^{ji}) = e_j a_n^{lk}. \quad \square \quad (4.14)$$

2) follows from 1). \square

3) By induction hypothesis we have $a_{n-1}^{kl} \subseteq a_{n-2}^{kl}$, and therefore

$$a_n^{ij} = e_i + e_j a_{n-1}^{kl} \subseteq e_i + e_j a_{n-2}^{kl} = a_{n-1}^{ij}. \quad \square$$

4) Without loss of generality, we will show that

$$e_2 + e_1 a_{t+1}^{24} a_{s-1}^{34} = e_2 + e_4 a_t^{12} a_s^{13}. \quad (4.15)$$

By permutation property (A.4) we have

$$\begin{aligned} e_2 + e_1 a_{t+1}^{24} a_{s-1}^{34} &= e_2 + e_1 (e_4 + e_2 a_t^{13}) a_{s-1}^{43} = \\ e_2 + e_1 (e_4 + e_2 a_t^{13} a_{s-1}^{43}) &= e_2 + e_4 (e_1 + e_2 a_t^{13} a_{s-1}^{43}) = \\ e_2 + e_4 a_t^{13} (e_1 + e_2 a_{s-1}^{43}) &= e_2 + e_4 a_t^{13} a_s^{12}. \quad \square \end{aligned}$$

Now we briefly recall definitions due to Gelfand and Ponomarev [GP74], [GP76], [GP77] of spaces G_i , G_i' , representations ν^0 , ν^1 , joint maps ψ_i , and elementary maps φ_i , where $i = 1, 2, 3, 4$. To compare these definitions with a case of the modular lattice $D^{2,2,2}$, see Ch.2.

We denote by

$$\{Y_1, Y_2, Y_3, Y_4 \mid Y_i \subseteq X_0, i = 1, 2, 3, 4\}$$

the representation ρ of D^4 in the finite dimensional vector space X_0 , and by

$$\{Y_1^1, Y_2^1, Y_3^1, Y_4^1 \mid Y_i^1 \subseteq X_0^1, i = 1, 2, 3, 4\}$$

the representation $\Phi^+\rho$ of D^4 . Here Y_i (resp. Y_i^1) is the image of the generator e_i under the representation ρ (resp. $\Phi^+\rho$).

$$\begin{array}{ccc} & Y_3 & Y_3^1 \\ & \cap \mid & \cap \mid \\ Y_1 \subseteq X_0 \supseteq Y_2 & & Y_1^1 \subseteq X_0^1 \supseteq Y_2^1 \\ & \cup \mid & \cup \mid \\ \rho & Y_4 & \Phi^+\rho & Y_4^1 \end{array} \quad (4.16)$$

The space X_0^1 — the space of the representation $\Phi^+\rho$ — is

$$X_0^1 = \{(\eta_1, \eta_2, \eta_3, \eta_4) \mid \eta_i \in Y_i, \quad \sum \eta_i = 0\},$$

where $i \in \{1, 2, 3, 4\}$. As in the case of $D^{2,2,2}$ §2.1, we set

$$R = \bigoplus_{i=1,2,3,4} Y_i,$$

i.e.,

$$R = \{(\eta_1, \eta_2, \eta_3, \eta_4) \mid \eta_i \in Y_i, \quad i = 1, 2, 3\}.$$

Then, $X_0^1 \subseteq R$.

For the case of D^4 , the spaces G_i and G'_i are introduced as follows:

$$\begin{aligned} G_1 &= \{(\eta_1, 0, 0, 0) \mid \eta_1 \in Y_1\}, & G'_1 &= \{(0, \eta_2, \eta_3, \eta_4) \mid \eta_i \in Y_i\}, \\ G_2 &= \{(0, \eta_2, 0, 0) \mid \eta_2 \in Y_2\}, & G'_2 &= \{(\eta_1, 0, \eta_3, \eta_4) \mid \eta_i \in Y_i\}, \\ G_3 &= \{(0, 0, \eta_3, 0) \mid \eta_3 \in Y_3\}, & G'_3 &= \{(\eta_1, \eta_2, 0, \eta_4) \mid \eta_i \in Y_i\}, \\ G_4 &= \{(0, 0, 0, \eta_4) \mid \eta_4 \in Y_4\}, & G'_4 &= \{(\eta_1, \eta_2, \eta_3, 0) \mid \eta_i \in Y_i\}. \end{aligned} \quad (4.17)$$

For details, see [GP74, p.43]¹.

The associated representations ν_0, ν_1 in R are defined by Gelfand and Ponomarev [GP74, eq.(7.2)]²:

$$\begin{aligned} \nu^0(e_i) &= X_0^1 + G_i, & i &= 1, 2, 3, \\ \nu^1(e_i) &= X_0^1 G'_i, & i &= 1, 2, 3. \end{aligned} \quad (4.18)$$

Following [GP74], we introduce the elementary maps φ_i :

$$\varphi_i : X_0^1 \longrightarrow X_0, (\eta_1, \eta_2, \eta_3, \eta_4) \longmapsto \eta_i.$$

From the definition we have

$$\varphi_1 + \varphi_2 + \varphi_3 + \varphi_4 = 0. \quad (4.19)$$

¹Compare with definition (2.1) of the spaces G_i, G'_i, H_i, H'_i in the case of $D^{2,2,2}$; see also Table 4.2.

²Compare with definitions for the case of $D^{2,2,2}$ in §2.2.

Notions	$D^{2,2,2}$	D^4
Generators	$\{x_1 \subseteq y_1, x_2 \subseteq y_2, x_3 \subseteq y_3\}$	$\{e_1, e_2, e_3, e_4\}$
Atomic elements	$a_n^{ij} = x_i + y_j a_{n-1}^{jk},$ $A_n^{ij} = y_i + x_j A_{n-1}^{ki},$ $\{i, j, k\} = \{1, 2, 3\}$	$a_n^{ij} = e_i + e_j a^{kl},$ $\{i, j, k, l\} = \{1, 2, 3, 4\}$
Representation ρ	$\rho(x_i) = X_i, \rho(y_i) = Y_i$	$\rho(e_i) = Y_i$
Space X_0^1	$X_0^1 = \{(\eta_1, \eta_2, \eta_3) \mid \eta_i \in Y_i\},$ where $\eta_1 + \eta_2 + \eta_3 = 0$	$X_0^1 = \{(\eta_1, \eta_2, \eta_3, \eta_4) \mid \eta_i \in Y_i\},$ where $\eta_1 + \eta_2 + \eta_3 + \eta_4 = 0$
Representation $\Phi^+ \rho$	$\rho(x_i) = X_i^1, \rho(y_i) = Y_i^1$	$\rho(y_i) = Y_i^1$
Spaces G_i, H_i	$G_1 = \{(\eta_1, 0, 0) \mid \eta_1 \in Y_1\},$ $H_1 = \{(\xi_1, 0, 0) \mid \xi_1 \in X_1\},$ $G_2 = \{(0, \eta_2, 0) \mid \eta_2 \in Y_2\},$ $H_2 = \{(0, \xi_2, 0) \mid \xi_2 \in X_2\},$ $G_3 = \{(0, 0, \eta_3) \mid \eta_3 \in Y_3\},$ $H_3 = \{(0, 0, \xi_3) \mid \xi_3 \in X_3\},$	$G_1 = \{(\eta_1, 0, 0, 0) \mid \eta_1 \in Y_1\}$ $G_2 = \{(0, \eta_2, 0, 0) \mid \eta_2 \in Y_2\}$ $G_3 = \{(0, 0, \eta_3, 0) \mid \eta_3 \in Y_3\}$ $G_4 = \{(0, 0, 0, \eta_4) \mid \eta_4 \in Y_4\}$
Joint maps ψ_i	$\psi_i(a) = X_0^1 + G_i(H_i' + \nu^1(a))$	$\psi_i(a) = X_0^1 + G_i(G_i' + \nu^1(a))$
Quasi-multiplicativity	$\psi_i(a)\psi_i(b) =$ $\psi_i((a + e_i)(b + x_j x_k))$	$\psi_i(a)\psi_i(b) =$ $\psi_i((a + e_i)(b + e_j e_k e_l))$
Elementary maps φ_i	$\varphi_i : X_0^1 \longrightarrow X_0,$ $(\eta_1, \eta_2, \eta_3) \longmapsto \eta_i$	$\varphi_i : X_0^1 \longrightarrow X_0,$ $(\eta_1, \eta_2, \eta_3, \eta_4) \longmapsto \eta_i$
Fundamental properties of elementary maps φ_i	$\varphi_i \varphi_j \varphi_i + \varphi_i \varphi_k \varphi_i = 0,$ $\varphi_i^3 = 0$	$\varphi_i \varphi_k \varphi_j + \varphi_i \varphi_l \varphi_j = 0$ $\varphi_i^2 = 0$
Fundamental properties of indices (admissible sequences)	$\boxed{iji = iki}$	$\boxed{ikj = ilj}$

TABLE 4.2. Comparison of notions in $D^{2,2,2}$ and D^4

4.3. Basic relations for the elementary and joint maps in the case of D^4

We define *joint* maps $\psi_i: D_4 \longrightarrow \mathcal{L}(R)$ as in the case of $D^{2,2,2}$:

$$\psi_i(a) = X_0^1 + G_i(G'_i + \nu^1(a)). \quad (4.20)$$

PROPOSITION 4.3.1. *In the case D^4 the joint maps ψ_i satisfy the following basic relations¹:*

- (1) $\psi_i(e_i) = X_0^1$,
- (2) $\psi_i(e_j) = \nu^0(e_i(e_k + e_l))$,
- (3) $\psi_i(I) = \nu^0(e_i(e_j + e_k + e_l))$,
- (4) $\psi_i(e_k e_l) = \psi_j(e_k e_l) = \nu^0(e_i e_j)$.

Proof. 1) From (4.20) and (4.18) we have $\psi_i(e_i) = X_0^1 + G_i G'_i = X_0^1$. \square

2) We have $\psi_i(y_j) = X_0^1 + G_i(G'_i + X_0^1 G'_j)$. From $G_i \subseteq G'_j$ for $i \neq j$ and by the permutation property (A.4), we get

$$\psi_i(y_j) = X_0^1 + G_i(G'_i G'_j + X_0^1).$$

Since $G'_i G'_j = G_k + G_l$, where i, j, k, l are distinct indices, we have

$$\psi_i(y_j) = X_0^1 + G_i(X_0^1 + G_k + G_l) = (X_0^1 + G_i)(X_0^1 + G_k + G_l) = \nu^0(e_i(e_k + e_l)). \quad \square$$

3) Again,

$$\begin{aligned} \psi_i(I) &= X_0^1 + G_i(G'_i + X_0^1) = \\ &= X_0^1 + G_i(G_j + G_k + G_l + X_0^1) = \nu^0(e_i(e_j + e_k + e_l)). \end{aligned} \quad (4.21) \quad \square$$

4) Since $G_i \subseteq G'_k$ for all $i \neq k$, we have

$$\begin{aligned} \psi_i(e_k e_l) &= X_0^1 + G_i(G'_i + X_0^1 G'_k G'_l) = \\ &= X_0^1 + G_i(G'_i G'_k G'_l + X_0^1). \end{aligned} \quad (4.22)$$

Since $G_j(G_i + G_k + G_l) = 0$, then

$$G'_i G'_j = G_k + G_l. \quad (4.23)$$

Indeed,

$$\begin{aligned} G'_i G'_j &= (G_j + G_k + G_l)(G_i + G_k + G_l) = \\ &= G_k + G_l + G_j(G_i + G_k + G_l) = G_k + G_l. \end{aligned} \quad (4.24)$$

From (4.22) and (4.23), we see that

$$\begin{aligned} \psi_i(e_k e_l) &= X_0^1 + G_i(G'_i G'_k G'_l + X_0^1) = \\ &= X_0^1 + G_i(X_0^1 + G_j) = \nu_0(e_i e_j). \end{aligned} \quad \square$$

The main relation between φ_i and ψ_i (Proposition 2.4.3) holds also for the case of D^4 . Namely, let $a, b, c \subseteq D^{2,2,2}$, then

- (i) If $\psi_i(a) = \nu^0(b)$, then $\varphi_i \Phi^+ \rho(a) = \rho(b)$.
 - (ii) If $\psi_i(a) = \nu^0(b)$ and $\psi_i(ac) = \psi_i(a)\psi_i(c)$, then
- $$\varphi_i \Phi^+ \rho(ac) = \varphi_i \Phi^+ \rho(a) \varphi_i \Phi^+ \rho(c). \quad (4.25)$$

From Proposition 4.3.1 and eq.(4.25) we have

¹Compare with Proposition 2.4.1 for $D^{2,2,2}$

COROLLARY 4.3.2. *For the elementary map φ_i the following basic relations hold¹:*

- (1) $\varphi_i \Phi^+ \rho(e_i) = 0$,
- (2) $\varphi_i \Phi^+ \rho(e_j) = \rho(e_i(e_k + e_l))$,
- (3) $\varphi_i \Phi^+ \rho(I) = \rho(e_i(e_j + e_k + e_l))$,
- (4) $\varphi_i \Phi^+ \rho(e_k y_l) = \rho(e_i e_j)$.

4.4. Additivity and multiplicativity of the joint maps in the case of D^4

PROPOSITION 4.4.1. *The map ψ_i is additive and quasimultiplicative with respect to the lattice operations $+$ and \cap , namely²:*

- (1) $\psi_i(a) + \psi_i(b) = \psi_i(a + b)$,
- (2) $\psi_i(a)\psi_i(b) = \psi_i((a + e_i)(b + x_j x_k x_l))$,
- (3) $\psi_i(a)\psi_i(b) = \psi_i(a(b + e_i + x_j x_k x_l))$.

Proof. 1) By the modular law (A.3)

$$\begin{aligned} \psi_i(a) + \psi_i(b) &= X_0^1 + G_i(G'_i + \nu^1(a)) + G_i(G'_i + \nu^1(b)) = \\ &= X_0^1 + G_i(G'_i + \nu^1(a) + G_i(G'_i + \nu^1(b))) = X_0^1 + G_i((G'_i + \nu^1(b))(G'_i + G_i) + \nu^1(a)). \end{aligned}$$

Since $G'_i + G_i = R$, it follows that

$$\psi_i(a) + \psi_i(b) = X_0^1 + G_i(G'_i + \nu^1(b) + \nu^1(a)) = X_0^1 + G_i(G'_i + \nu^1(b + a)). \quad \square$$

2) By definition (2.6) $\nu^1(b) \subseteq X_0^1$, and by the permutation property (A.5) we have

$$X_0^1 + G_i(G'_i + \nu^1(a)) = X_0^1 + G'_i(G_i + \nu^1(a)).$$

By the modular law (A.3) and by (A.5)

$$\begin{aligned} \psi_i(a)\psi_i(b) &= \\ &= X_0^1 + G_i(G'_i + \nu^1(a))(X_0^1 + G'_i(G_i + \nu^1(b))) = \\ &= X_0^1 + G_i(X_0^1(G'_i + \nu^1(a)) + G'_i(G_i + \nu^1(b))). \end{aligned}$$

Since

$$X_0^1(G'_i + \nu^1(a)) = X_0^1 G'_i + \nu^1(a) \quad \text{and} \quad X_0^1 G'_i = \nu^1(e_i),$$

we see that

$$\psi_i(a)\psi_i(b) = X_0^1 + G_i(\nu^1(e_i) + \nu^1(a) + G'_i(G_i + \nu^1(b))). \quad (4.26)$$

By the permutation property (A.4) and by (2.19) we have

$$\psi_i(a)\psi_i(b) = X_0^1 + G_i(G'_i + (\nu^1(e_i) + \nu^1(a))(G_i + \nu^1(b))). \quad (4.27)$$

Since

$$G_i = G'_j G'_k G'_l \quad \text{and} \quad \nu^1(e_i) + \nu^1(a) = \nu^1(e_i + a) = X_0^1(\nu^1(e_i + a)),$$

it follows that

$$\begin{aligned} \psi_i(a)\psi_i(b) &= X_0^1 + G_i(G'_i + \nu^1(e_i + a)(X_0^1 G'_j G'_k G'_l + \nu^1(b))) = \\ &= X_0^1 + G_i(G'_i + \nu^1(e_i + a)(\nu^1(e_j e_k e_l) + \nu^1(b))) = \\ &= X_0^1 + G_i(G'_i + \nu^1(e_i + a)\nu^1(e_j e_k e_l + b)) = \psi_i((a + e_i)(b + e_j e_k e_l)). \quad \square \end{aligned}$$

¹Compare with Corollary 2.4.4 — a similar proposition for $D^{2,2,2}$.

²Compare with Proposition 2.5.1, case $D^{2,2,2}$

3) From (4.26) and since $\nu^1(e_i) = X_0^1 G'_i \subseteq G'_i$, we have

$$\psi_i(a)\psi_i(b) = X_0^1 + G_i (\nu^1(a) + G'_i(G_i + \nu^1(b) + \nu^1(e_i))) . \quad (4.28)$$

Again, by (A.4) we have

$$\begin{aligned} \psi_i(a)\psi_i(b) &= X_0^1 + G_i (G'_i + \nu^1(a)(G_i + \nu^1(b) + \nu^1(e_i))) = \\ &= X_0^1 + G_i (G'_i + \nu^1(a)(X_0^1 G'_j G'_k G'_l + \nu^1(b) + \nu^1(e_i))) . \end{aligned}$$

Thus, $\psi_i(a)\psi_i(b) = \psi_i(a(b + x_i + x_j x_k x_l))$. \square

We need the following corollary (atomic multiplicativity) from Proposition 4.4.1

COROLLARY 4.4.2. 1) Suppose one of the following inclusions holds¹:

- (i) $e_i + e_j e_k e_l \subseteq a$,
- (ii) $e_i + e_j e_k e_l \subseteq b$,
- (iii) $e_i \subseteq a$, $e_j e_k e_l \subseteq b$,
- (iv) $e_i \subseteq b$, $e_j e_k e_l \subseteq a$.

Then the joint map ψ_i operates as a homomorphism on the elements a and b with respect to the lattice operations $+$ and \cap , i.e.,

$$\psi_i(a) + \psi_i(b) = \psi_i(a + b), \quad \psi_i(a)\psi_i(b) = \psi_i(a \cap b).$$

2) The joint map ψ_i applied to the following atomic elements is the intersection preserving map, i.e., multiplicative with respect to the operation \cap :

$$\psi_i(b a_n^{ij}) = \psi_i(b) \psi_i(a_n^{ij}) \text{ for every } b \subseteq D^4. \quad (4.29)$$

4.5. The action of maps ψ_i and φ_i on the atomic elements in D^4

PROPOSITION 4.5.1. The joint maps ψ_i applied to the atomic elements a_n^{ij} satisfy the following relations²

- (1) $\psi_i(a_n^{ij}) = \nu^0(e_i a_n^{kl})$,
- (2) $\psi_j(a_n^{ij}) = \nu^0(e_j(e_k + e_l))$,
- (3) $\psi_j(e_i a_n^{kl}) = \nu^0(e_j a_{n+1}^{kl})$.

Proof. 1) Since $\psi_i(e_i) = X_0^1$ (Proposition 4.3.1, heading (1)), we have

$$\psi_i(a_n^{ij}) = \psi_i(e_i + e_j a_{n-1}^{kl}) = \psi_i(e_i) + \psi_i(e_j a_{n-1}^{kl}) = \psi_i(e_j a_{n-1}^{kl}).$$

We suppose that heading (3) of Proposition 4.5.1 for $n - 1$ is true (induction hypothesis), and we get

$$\psi_i(a_n^{ij}) = \psi_i(e_j a_{n-1}^{kl}) = \nu^0(e_j a_n^{kl}). \quad \square$$

2) Here,

$$\psi_j(a_n^{ij}) = \psi_j(e_i + e_j a_{n-1}^{kl}) = \psi_j(e_i) + \psi_j(e_j a_{n-1}^{kl}) = \psi_j(e_i).$$

Further, by Proposition 4.3.1, heading (2), we have

$$\psi_j(a_n^{ij}) = \psi_j(e_i) = \nu^0(e_j(e_k + e_l)). \quad \square$$

¹Compare with Corollary 2.5.2, case $D^{2,2,2}$.

²Compare with Proposition 2.7.1, the modular lattice $D^{2,2,2}$.

3) For convenience, without loss of generality, we will show that

$$\psi_1(e_2 a_n^{34}) = \nu^0(e_1 a_{n+1}^{34}). \quad (4.30)$$

By permutation property (A.4) and since $\nu^1(e_4 a_{n-1}^{12}) \subseteq X_0^1$, we have

$$\begin{aligned} \nu^1(e_2 a_n^{34}) &= \nu^1(e_2(e_3 + e_4 a_{n-1}^{12})) = \\ &X_0^1 G_2'(X_0^1 G_3' + \nu^1(e_4 a_{n-1}^{12})) = X_0^1 G_2'(G_3' + \nu^1(e_4 a_{n-1}^{12})). \end{aligned} \quad (4.31)$$

By (4.31) we have

$$\begin{aligned} \psi_1(e_2 a_n^{34}) &= X_0^1 + G_1(G_1' + \nu^1(e_2 a_n^{34})) = \\ &X_0^1 + G_1(G_1' + X_0^1 G_2'(G_3' + \nu^1(e_4 a_{n-1}^{12}))). \end{aligned}$$

Since $G_1 \subseteq G_2'$ and $G_1 \subseteq G_3'$, by permutation property (A.4), we have

$$\begin{aligned} \psi_1(e_2 a_n^{34}) &= X_0^1 + G_1(G_1' G_2' + X_0^1(G_3' + \nu^1(e_4 a_{n-1}^{12}))) = \\ &X_0^1 + G_1(G_3' + G_4' + X_0^1(G_3' + \nu^1(e_4 a_{n-1}^{12}))) = \\ &X_0^1 + G_1((G_3' + G_4')(G_3' + \nu^1(e_4 a_{n-1}^{12}))) + X_0^1. \end{aligned}$$

Since $G_4 \subseteq G_3'$, we have

$$\begin{aligned} \psi_1(e_2 a_n^{34}) &= X_0^1 + G_1(G_4' + G_3'(G_3' + \nu^1(e_4 a_{n-1}^{12}))) + X_0^1 = \\ &X_0^1 + G_1((X_0^1 + G_4') + (X_0^1 + G_3'(G_3' + \nu^1(e_4 a_{n-1}^{12}))))). \end{aligned}$$

Pay attention to the fact

$$\psi_3(e_4 a_{n-1}^{12}) = X_0^1 + G_3'(G_3' + \nu^1(e_4 a_{n-1}^{12})),$$

and therefore

$$\begin{aligned} \psi_1(e_2 a_n^{34}) &= X_0^1 + G_1((X_0^1 + G_4') + \psi_3(e_4 a_{n-1}^{12})) = \\ &(X_0^1 + G_1)((X_0^1 + G_4') + \psi_3(e_4 a_{n-1}^{12})) = \\ &\nu^0(e_1)(\nu^0(e_4) + \psi_3(e_4 a_{n-1}^{12})). \end{aligned} \quad (4.32)$$

By induction hypothesis for $n - 1$ in heading (3) we have

$$\psi_3(e_4 a_{n-1}^{12}) = \nu^0(e_3 a_n^{12}),$$

and by (4.32) we have

$$\psi_1(e_2 a_n^{34}) = \nu^0(e_1)(\nu^0(e_4) + \nu^0(e_3 a_n^{12})) = \nu^0(e_1(e_4 + e_3 a_n^{12})) = \nu^0(e_1 a_{n+1}^{43}). \quad \square$$

From Propositions (2.4.3) and (4.5.1) we have

COROLLARY 4.5.2. *The elementary maps φ_i applied to the atomic elements a_n^{ij} , where $n \geq 1$, satisfy the following relations¹*

- (1) $\varphi_i \Phi^+ \rho(a_n^{ij}) = \rho(e_i a_n^{kl}),$
- (2) $\varphi_j \Phi^+ \rho(a_n^{ij}) = \rho(e_j(e_k + e_l)),$
- (3) $\varphi_j \Phi^+ \rho(e_i a_n^{kl}) = \rho(e_j a_{n+1}^{kl}).$

¹Compare with Corollary 2.7.4, case $D^{2,2,2}$

4.6. The fundamental property of the elementary maps

PROPOSITION 4.6.1. For $\{i, j, k, l\} = \{1, 2, 3, 4\}$ the following relations hold¹

$$\varphi_i \varphi_k \varphi_j + \varphi_i \varphi_l \varphi_j = 0, \quad (4.33)$$

$$\varphi_i^2 = 0. \quad (4.34)$$

Proof. For every vector $v \in X_0^1$, by definition of φ_i we have $(\varphi_i + \varphi_j + \varphi_k + \varphi_l)(v) = 0$, see eq. (4.19). In other words, $\varphi_i + \varphi_j + \varphi_k + \varphi_l = 0$. Therefore,

$$\varphi_i \varphi_k \varphi_j + \varphi_i \varphi_l \varphi_j = \varphi_i(\varphi_i + \varphi_j) \varphi_j = \varphi_i^2 \varphi_j + \varphi_i \varphi_j^2.$$

So, it suffices to prove that $\varphi_i^2 = 0$. For every $z \subseteq D^4$, by Corollary 4.3.2, headings (3) and (1), we have

$$\begin{aligned} \varphi_i^2((\Phi^+)^2 \rho)(z) &\subseteq \varphi_i(\varphi_i((\Phi^+)^2 \rho(I))) = \\ \varphi_i(\Phi^+ \rho(e_i(e_j + e_k + e_l))) &\subseteq \varphi_i(\Phi^+ \rho(e_i)) = 0. \quad \square \end{aligned}$$

COROLLARY 4.6.2. The relation

$$\varphi_i \varphi_k \varphi_j(B) = \varphi_i \varphi_l \varphi_j(B) \quad (4.35)$$

takes place² for every subspace $B \subseteq X_0^2$, where X_0^2 is the representation space of $(\Phi_i^+)^2 \rho$.

Essentially, relations (4.33) and (4.35) are fundamental and motivate the construction of the admissible sequences satisfying the following relation:

$$ikj = ilj, \quad (4.36)$$

where indices i, j, k, l are all distinct, see Table 4.2 and §4.1.

4.7. The φ_i -homomorphic elements in D^4

By analogy with the modular lattice $D^{2,2,2}$ (see §1.6), we introduce now φ_i -homomorphic polynomials in D^4 .

An element $a \subseteq D^4$ is said to be φ_i -homomorphic, if

$$\varphi_i \Phi^+ \rho(ap) = \varphi_i \Phi^+ \rho(a) \varphi_i \Phi^+ \rho(p) \text{ for all } p \subseteq D^4. \quad (4.37)$$

An element $a \subseteq D^4$ is said to be (φ_i, e_k) -homomorphic, if

$$\varphi_i \Phi^+ \rho(ap) = \varphi_i \Phi^+ \rho(e_k a) \varphi_i \Phi^+ \rho(p) \text{ for all } p \subseteq e_k. \quad (4.38)$$

THEOREM 4.7.1. 1) The polynomials a_n^{ij} are φ_i -homomorphic³.

2) The polynomials a_n^{ij} are (φ_j, e_k) -homomorphic for distinct indices $\{i, j, k\}$.

Proof. 1) Follows from multiplicativity (4.29), Proposition 4.5.1 and the analog of Proposition 2.4.3 for D^4 .

2) For convenience, without loss of generality, we will show that

$$\varphi_2 \Phi^+ \rho(a_n^{12} p) = \varphi_2 \Phi^+ \rho(e_3 a) \varphi_2 \Phi^+ \rho(p) \text{ for all } p \subseteq e_3. \quad (4.39)$$

Let $p \subseteq e_3$. First,

$$pa_s^{12} = pe_3 a_s^{12} = pe_3 a_s^{21} = pa_s^{21}.$$

¹Compare with Proposition 2.10.2, case $D^{2,2,2}$

²Compare with Corollary 2.10.3, case $D^{2,2,2}$

³Compare with Theorem 2.8.1, case $D^{2,2,2}$.

By Corollary 4.4.2 and Proposition 4.5.1, heading (3) we have

$$\begin{aligned}\psi_2(pa_n^{12}) &= \psi_2(pe_3a_n^{21}) = \psi_2(p(e_2 + e_1a_{n-1}^{34})) = \\ \psi_2(p(e_1a_{n-1}^{34})) &= \psi_2(p)\psi_2(e_1a_{n-1}^{34}) = \\ \psi_2(p)\nu^0(e_2a_n^{34}).\end{aligned}\tag{4.40}$$

Since $e_2a_n^{34} = e_2(e_1 + e_3 + e_4)a_n^{34}$ and

$$\begin{aligned}\nu^0(e_2a_n^{34}) &= \nu^0(e_2(e_1 + e_3 + e_4)a_n^{34}) = \\ \nu^0(e_2(e_1 + e_3 + e_4))\nu^0(a_n^{34}) &= \psi_2(I)\nu^0(a_n^{34}),\end{aligned}$$

by (4.40) we have

$$\psi_2(pa_n^{12}) = \psi_2(p)\psi_2(I)\nu^0(a_n^{34}) = \psi_2(p)\nu^0(a_n^{34}).\tag{4.41}$$

We have $\psi_2(p) \subseteq \psi_2(e_3)$ together with $p \subseteq e_3$, and therefore $\psi_2(e_3p) = \psi_2(e_3)\psi_2(p)$. From (4.41) we get

$$\begin{aligned}\psi_2(pa_n^{12}) &= \psi_2(p)\psi_2(e_3)\nu^0(a_n^{34}) = \\ \psi_2(p)\nu^0(e_2(e_1 + e_4))\nu^0(a_n^{34}) &= \\ \psi_2(p)\nu^0(e_2(e_1 + e_4)(a_n^{34})) &= \psi_2(p)\psi_2(e_3a_n^{21}),\end{aligned}$$

i.e.,

$$\psi_2(pa_n^{12}) = \psi_2(p)\psi_2(e_3a_n^{21}).\tag{4.42}$$

Applying projection ∇ to (4.42) as in Theorem 2.8.1 we get (4.39) and theorem is proved. \square

4.8. The theorem on the classes of admissible elements in D^4

THEOREM 4.8.1. *Let $\alpha = i_n i_{n-1} \dots 1$ be an admissible sequence for D^4 and $i \neq i_n$. Then $i\alpha$ is admissible and, for $z_\alpha = e_\alpha$ or $f_{\alpha 0}$ from Table 4.3, the following relation holds¹:*

$$\varphi_i \Phi^+ \rho(z_\alpha) = \rho(z_{i\alpha}).\tag{4.43}$$

For the proof of the theorem on admissible elements in D^4 , see §B.2. The proof repeatedly uses the basic properties of the admissible sequences in D^4 considered in §4.8.1, Lemma 4.8.2.

4.8.1. Basic properties of admissible elements in D^4 . We prove here a number of basic properties² of the atomic elements in D^4 used in the proof of the theorem on admissible elements (Theorem 4.8.1). In particular, in some cases the lower indices of polynomials a_s^{ij} entering in the admissible elements $f_{\alpha 0}$ can be transformed as in the following

LEMMA 4.8.2. *1) Every polynomial $f_{\alpha 0}$ from Table 4.3 can be represented as an intersection of e_α and P . For every $i \neq i_n$ (see §1.7.2), we select P to be some φ_i -homomorphic polynomial.*

2) The lower indices of polynomials a_s^{ij} entering in the admissible elements $f_{\alpha 0}$ can be equalized as follows:

$$\begin{aligned}f_{\alpha 0} &= f_{(21)^t(41)^r(31)^s 0} = \\ e_\alpha(e_2a_{2t}^{34} + a_{2r+1}^{41}a_{2s-1}^{31}) &= e_\alpha(a_{2t}^{43} + e_1a_{2r}^{24}a_{2s}^{23}).\end{aligned}\tag{4.44}$$

¹Compare with Theorem 2.12.1, case $D^{2,2,2}$

²Compare with Table 2.1 and §2.6, case $D^{2,2,2}$.

The generic relation¹ is the following:

$$\begin{aligned} e_i(e_i a_t^{jl} + a_{r+1}^{kj} a_{s-1}^{kl}) = \\ e_i(a_t^{jl} + e_i a_{r+1}^{kj} a_{s-1}^{kl}) = e_i(a_t^{jl} + e_k a_r^{ij} a_s^{il}). \end{aligned} \quad (4.45)$$

3) The substitution

$$\begin{cases} r \mapsto r - 2, \\ s \mapsto s + 2 \end{cases} \quad (4.46)$$

does not change the polynomial $e_i(e_i a_t^{jl} + a_{r+1}^{kj} a_{s-1}^{kl})$, namely:

$$e_i(e_i a_t^{jl} + a_{r+1}^{kj} a_{s-1}^{kl}) = e_i(e_i a_t^{jl} + a_{r-1}^{kj} a_{s+1}^{kl}). \quad (4.47)$$

4) The substitution (4.46) does not change the polynomial $e_i a_s^{kj} a_r^{kl}$:

$$e_i a_{r+1}^{kj} a_{s-1}^{kl} = e_i a_{r-1}^{kj} a_{s+1}^{kl}. \quad (4.48)$$

Proof. 1) We consider here only Line $F21$ from Table 4.3, all other cases are similarly considered. Every polynomial $f_{\alpha 0}$ from Table 4.3 is the intersection e_α and P , where P is the sum contained in the parantheses, and polynomial $f_{\alpha 0}$ can be represented in the different equivalent forms, such that only the polynomial P is changed. For Line $F21$, we have

$$\begin{aligned} f_{\alpha 0} &= f_{(21)^t(41)^r(31)^s 0} = \\ e_\alpha(e_2 a_{2t}^{34} + a_{2r+1}^{41} a_{2s-1}^{31}) &= e_\alpha(e_2 a_{2t}^{43} + a_{2r+1}^{41} a_{2s-1}^{31}) = & (i) \\ e_\alpha(a_{2t}^{34} + e_2 a_{2r+1}^{41} a_{2s-1}^{31}) &= e_\alpha(a_{2t}^{43} + e_2 a_{2r+1}^{41} a_{2s-1}^{31}) = & (ii) \\ e_\alpha(a_{2t}^{34} + e_2 a_{2r+1}^{14} a_{2s-1}^{13}) &= e_\alpha(a_{2t}^{43} + e_2 a_{2r+1}^{14} a_{2s-1}^{13}) = & (iii) \\ e_\alpha(e_2 a_{2t}^{34} + a_{2r+1}^{14} a_{2s-1}^{13}) &= e_\alpha(e_2 a_{2t}^{43} + a_{2r+1}^{14} a_{2s-1}^{13}) = & (iv) \\ e_\alpha(e_2 a_{2t}^{34} a_{2s-1}^{13} + a_{2r+1}^{14}) &= e_\alpha(e_2 a_{2t}^{43} a_{2s-1}^{13} + a_{2r+1}^{14}) = \end{aligned} \quad (4.49)$$

Relation (i) in (4.49) follows from the definition of Table 4.3 and (4.11). Relations (ii) in (4.49) follow from the inclusion $e_\alpha \subseteq e_2$ and permutation property (A.4). Relations (iii) and (iv) follow from (4.11). Further, by (4.12)

$$e_\alpha \subseteq a_{2s}^{31} \subseteq a_{2s-1}^{31}$$

and by permutation property (A.4) we have the relation (v).

The polynomial P in (iv) is φ_1 -homomorphic since

$$e_1 \subseteq a_{2r+1}^{14} a_{2s-1}^{13} \text{ and } e_2 e_3 e_4 \subseteq e_2 a_{2t}^{34}.$$

The polynomial P in (iii) is φ_3 -homomorphic since

$$e_3 \subseteq a_{2t}^{34} \text{ and } e_1 e_2 e_4 \subseteq e_1 e_2 \subseteq e_2 a_{2r+1}^{14} a_{2s-1}^{13}.$$

The polynomial P in (iii) is φ_4 -homomorphic since

$$e_4 \subseteq a_{2t}^{43} \text{ and } e_1 e_2 e_3 \subseteq e_1 e_2 \subseteq e_2 a_{2r+1}^{14} a_{2s-1}^{13}.$$

Essentially, heading 1) allows to select an appropriate form of the polynomial P for every given φ_i . \square

¹Throughout this lemma we suppose that $\{i, j, k, l\} = \{1, 2, 3, 4\}$.

2) Let us prove (4.44). By (4.13) we have

$$\begin{aligned} e_\alpha(e_2 a_{2t}^{34} + a_{2r+1}^{41} a_{2s-1}^{31}) &= e_\alpha(a_{2t}^{43} + e_2 a_{2r+1}^{41} a_{2s-1}^{31}) = \\ e_\alpha(a_{2t}^{43} + (e_4 + e_2 a_{2r+1}^{41} a_{2s-1}^{31})) &= e_\alpha(a_{2t}^{34} + e_4 + e_1 a_{2r}^{24} a_{2s}^{23}) = \\ e_\alpha(a_{2t}^{34} + e_1 a_{2r}^{24} a_{2s}^{23}). \end{aligned}$$

Now consider (4.45). Since $e_j \subseteq a_t^{jl}$, by (4.13) we have

$$\begin{aligned} e_i(a_t^{jl} + e_i a_{r+1}^{kj} a_{s-1}^{kl}) &= e_i(a_t^{jl} + (e_j + e_i a_{r+1}^{kj} a_{s-1}^{kl})) = \\ e_i(a_t^{jl} + e_k a_{r+1}^{ij} a_{s-1}^{il}). \quad \square \end{aligned}$$

3) By (4.45) we have

$$e_i(e_i a_t^{jl} + a_{r+1}^{kj} a_{s-1}^{kl}) = e_i(a_t^{jl} + e_k a_r^{ij} a_s^{il}),$$

On the other hand

$$\begin{aligned} e_i(e_i a_t^{jl} + a_{r-1}^{kj} a_{s+1}^{kl}) &= \\ e_i(e_i a_t^{lj} + a_{s+1}^{kl} a_{r-1}^{kj}) &= e_i(a_t^{lj} + e_k a_s^{il} a_r^{ij}), \end{aligned}$$

i.e.,

$$e_i(e_i a_t^{jl} + a_{r+1}^{kj} a_{s-1}^{kl}) = e_i(e_i a_t^{jl} + a_{r-1}^{kj} a_{s+1}^{kl}).$$

and (4.47) is true.

4) By (4.13) we have

$$\begin{aligned} e_i a_{r+1}^{kj} a_{s-1}^{kl} &= \\ e_i(e_k + e_j a_r^{il}) a_{s-1}^{kl} &= e_i(e_k + e_j a_r^{il} a_{s-1}^{kl}) = \\ e_i(e_k + e_l a_{r-1}^{ij} a_s^{kj}) &= e_i(e_k + e_l a_{r-1}^{ij}) a_s^{kj} = e_i a_r^{kl} a_s^{kj}, \end{aligned}$$

i.e.,

$$e_i a_{r+1}^{kj} a_{s-1}^{kl} = e_i a_r^{kl} a_s^{kj}. \quad (4.50)$$

By repeating (4.50), we see that

$$e_i a_{r+1}^{kj} a_{s-1}^{kl} = e_i a_{r-1}^{kj} a_{s+1}^{kl}.$$

The lemma is proved. \square

4.8.2. Coincidence with the Gelfand-Ponomarev polynomials. Let $e_\alpha, f_{\alpha 0}$ be admissible elements constructed in this work and $\tilde{e}_\alpha, \tilde{f}_{\alpha 0}$ be the admissible elements constructed by Gelfand and Ponomarev [GP74]. Since Theorem 4.8.1 takes place for both $e_\alpha, f_{\alpha 0}$ and for $\tilde{e}_\alpha, \tilde{f}_{\alpha 0}$ [GP74, Th.7.2, Th.7.3] we have

PROPOSITION 4.8.3. *The elements e_α , (resp. $f_{\alpha 0}$) and \tilde{e}_α (resp. $\tilde{f}_{\alpha 0}$) coincide mod θ .*

We will prove that this coincidence (without restriction mod θ) takes place for the admissible sequences of the small length. Recall definitions of \tilde{e}_α and $\tilde{f}_{\alpha 0}$ from [GP74].

The definition of \tilde{e}_α , [GP74, p.6].

$$\tilde{e}_{i_n i_{n-1} \dots i_2 i_1} = \tilde{e}_{i_n} \sum_{\beta \in \Gamma_e(\alpha)} \tilde{e}_\beta, \quad (4.51)$$

where

$$\begin{aligned} \Gamma_e(\alpha) = \{\beta = (k_{n-1}, \dots, k_2, k_1) \mid k_{n-1} \notin \{i_n, i_{n-1}\}, \dots, k_1 \notin \{i_2, i_1\}, \text{ and} \\ k_1 \neq k_2, \dots, k_{n-2} \neq k_{n-1}\}. \end{aligned} \quad (4.52)$$

The definition of $\tilde{f}_{\alpha 0}$, [GP74, p.53].

$$\tilde{f}_{i_n i_{n-1} \dots i_2 i_1 0} = \tilde{e}_{i_n} \sum_{\beta \in \Gamma_f(\alpha)} \tilde{e}_\beta, \quad (4.53)$$

where

$$\Gamma_f(\alpha) = \{\beta = (k_n, \dots, k_2, k_1) \mid k_n \notin \{i_n, i_{n-1}\}, \dots, k_2 \notin \{i_2, i_1\}, k_1 \notin \{i_1\} \text{ and } k_1 \neq k_2, \dots, k_{n-2} \neq k_{n-1}\}. \quad (4.54)$$

PROPOSITION 4.8.4 (The elements \tilde{e}_α). *Consider elements \tilde{e}_α for $\alpha = 21, 121, 321, 2341$ (see §1.9.2). The relation*

$$e_\alpha = \tilde{e}_\alpha$$

takes place without restriction mod θ .

Proof. For $n = 2$: $\alpha = 21$. We have

$$\tilde{e}_{21} = e_2 \sum_{j \neq 1, 2} e_j = e_2(e_3 + e_4). \quad (4.55)$$

According to §1.9.2 we see that $e_{21} = \tilde{e}_{21}$. \square

For $n = 3$: 1) $\alpha = 121$,

$$\begin{aligned} \Gamma_e(\alpha) &= \{(k_2 k_1) \mid k_2 \in \{3, 4\}, k_1 \in \{3, 4\}, k_1 \neq k_2\}, \\ \tilde{e}_{121} &= e_1 \sum_{\beta \in \Gamma_e(\alpha)} e_\beta = e_1(e_{34} + e_{43}) = e_1(e_3(e_1 + e_2) + e_4(e_1 + e_2)) = e_1 a_2^{34}. \end{aligned} \quad (4.56)$$

By §1.9.2 we have $e_{121} = \tilde{e}_{121}$. \square

2) $\alpha = 321 = 341$. We have

$$\Gamma_e(\alpha) = \{(k_2 k_1) \mid k_2 \in \{3, 2\}, k_1 \in \{2, 1\}, k_1 \neq k_2\} = \{14, 13, 43\}.$$

$$\begin{aligned} \tilde{e}_{321} &= e_1 \sum_{\beta \in \Gamma_e(\alpha)} e_\beta = e_3(e_{14} + e_{13} + e_{43}) = \\ &= e_3(e_1(e_2 + e_4) + e_1(e_2 + e_3) + e_4(e_1 + e_2)) = \\ &= e_3((e_1 + e_2)(e_2 + e_4)(e_1 + e_4) + e_1(e_2 + e_3)) = \\ &= e_3((e_1 + e_2)(e_1 + e_4)(e_2 + e_4 + e_1(e_2 + e_3))) = \\ &= e_3((e_1 + e_2)(e_1 + e_4)(e_4 + (e_1 + e_2)(e_2 + e_3))) = \\ &= e_3((e_1 + e_2)(e_1 + e_4)(e_4(e_2 + e_3) + (e_1 + e_2))). \end{aligned}$$

Since $e_1 + e_2 \subseteq e_4(e_2 + e_3) + e_1 + e_2$, we have

$$\tilde{e}_{321} = e_3(e_{14} + e_{13} + e_{43}) = e_3(e_1 + e_2)(e_1 + e_4). \quad (4.57)$$

Since \tilde{e}_{321} is symmetric with respect to transposition $2 \leftrightarrow 4$ we have

$$\tilde{e}_{321} = \tilde{e}_{341} = e_3(e_{14} + e_{13} + e_{43}) = e_3(e_{12} + e_{13} + e_{23}) = e_3(e_1 + e_2)(e_1 + e_4). \quad (4.58)$$

By §1.9.2 we have $e_{321} = \tilde{e}_{321}$. \square

For $n = 4$: $\alpha = 2341 = 2321 = 2141$. We have

$$\begin{aligned} \Gamma_e(\alpha) &= \{(k_3 k_2 k_1) \mid k_3 \in \{1, 4\}, k_2 \in \{1, 2\}, k_1 \in \{2, 3\}, \quad k_1 \neq k_2, k_2 \neq k_3\} = \\ &= \{(123), (412), (413), (423)\}, \end{aligned} \quad (4.59)$$

and

$$\begin{aligned}
\tilde{e}_{2341} &= e_2 \sum_{\beta \in \Gamma_e(\alpha)} e_\beta = e_2(e_{123} + e_{412} + e_{413}) = \\
&e_2(e_1(e_2 + e_3)(e_4 + e_3) + e_4(e_1 + e_2)(e_3 + e_2) + e_4(e_1 + e_3)(e_2 + e_3)) = \\
&e_2(e_4 + e_3)(e_1(e_2 + e_3) + e_4(e_1 + e_2)(e_3 + e_2) + e_4(e_1 + e_3)(e_3 + e_2)) = \\
&e_2(e_4 + e_3)(e_1(e_2 + e_3) + e_4 + e_4(e_1 + e_3)(e_1 + e_2)(e_3 + e_2)) = \\
&e_2(e_4 + e_3)(e_1(e_2 + e_3) + e_4).
\end{aligned} \tag{4.60}$$

By §1.9.2 we have $e_{2341} = \tilde{e}_{2341}$. The proposition is proved. \square

PROPOSITION 4.8.5 (The elements $\tilde{f}_{\alpha 0}$). *Consider elements $\tilde{f}_{\alpha 0}$ for $\alpha = 21, 121, 321$ (see §1.9.2). The following relation*

$$f_{\alpha 0} = \tilde{f}_{\alpha 0}$$

takes place without restriction mod θ .

Proof. For $n = 2$: $\alpha = 21$. We have

$$\begin{aligned}
\Gamma_f(\alpha) &= \{(k_2 k_1) \mid k_2 \in \{3, 4\}, k_1 \in \{2, 3, 4\}, k_1 \neq k_2\}, \\
\tilde{f}_{210} &= e_2 \sum_{\beta \in \Gamma_f(\alpha)} e_\beta = e_2(e_{32} + e_{34} + e_{42} + e_{43}) = \\
&e_2(e_3(e_1 + e_4) + e_3(e_2 + e_1) + e_4(e_1 + e_3) + e_4(e_1 + e_2)) = \\
&e_2((e_3 + e_4)(e_1 + e_4)(e_1 + e_3) + e_3(e_1 + e_2) + e_4(e_1 + e_2)) = \\
&e_2(e_3 + e_4)((e_1 + e_4)(e_1 + e_3) + e_3(e_1 + e_2) + e_4(e_1 + e_2)) = \\
&e_2(e_3 + e_4)(e_1 + e_4(e_1 + e_3) + e_3(e_1 + e_2) + e_4(e_1 + e_2)) = \\
&e_2(e_3 + e_4)(e_4(e_1 + e_3) + (e_1 + e_3)(e_1 + e_2) + e_4(e_1 + e_2)) = \\
&e_2(e_3 + e_4)(e_4(e_1 + e_3)(e_1 + e_2) + (e_1 + e_3)(e_1 + e_2) + e_4) = \\
&e_2(e_3 + e_4)(e_4 + (e_1 + e_3)(e_1 + e_2)) = \\
&e_2(e_3 + e_4)(e_4 + e_1 + e_3(e_1 + e_2)).
\end{aligned} \tag{4.61}$$

By §1.9.2 we have $f_{210} = \tilde{f}_{210}$. \square

For $n = 3$: 1) $\alpha = 121$. For this case, we have

$$\Gamma_f(\alpha) = \{(k_3 k_2 k_1) \mid k_3 \in \{3, 4\}, k_2 \in \{3, 4\}, k_1 \in \{2, 3, 4\}, k_1 \neq k_2, k_2 \neq k_3\}, \tag{4.62}$$

and

$$\begin{aligned}
\tilde{f}_{1210} &= e_1 \sum_{\beta \in \Gamma_f(\alpha)} e_\beta = e_1(e_{342} + e_{343} + e_{432} + e_{434}) = \\
&e_1[e_3(e_1 + e_2)(e_4 + e_2) + e_3(e_1(e_4 + e_3) + e_2(e_4 + e_3)) + \\
&\quad e_4(e_1 + e_2)(e_3 + e_2) + e_4(e_1(e_4 + e_3) + e_2(e_4 + e_3))] = \\
&e_1[e_3(e_1 + e_2)(e_4 + e_2 + e_3(e_1 + e_2(e_4 + e_3))) + \\
&\quad e_4(e_1 + e_2)(e_3 + e_2 + e_4(e_1 + e_2(e_4 + e_3)))] = \\
&e_1[e_3(e_1 + e_2)(e_4 + e_2 + e_1(e_3 + e_2(e_4 + e_3))) + \\
&\quad e_4(e_1 + e_2)(e_3 + e_2 + e_1(e_4 + e_2(e_4 + e_3)))] = \\
&e_1[(e_4 + e_2 + e_1(e_3 + e_2)(e_4 + e_3))] \\
&\quad [(e_3 + e_2 + e_1(e_4 + e_2)(e_4 + e_3))] \\
&\quad [e_3(e_1 + e_2) + e_4(e_1 + e_2)] = \\
&e_1[(e_4 + e_2)(e_4 + e_3) + e_1(e_3 + e_2)] \\
&\quad [(e_3 + e_2 + e_1(e_4 + e_2)(e_4 + e_3))] \\
&\quad [e_3(e_1 + e_2) + e_4(e_1 + e_2)] = \\
&e_1[(e_4 + e_2)(e_4 + e_3) + e_1(e_3 + e_2)] \\
&\quad [e_1(e_3 + e_2) + (e_4 + e_2)(e_4 + e_3)] \\
&\quad [e_3(e_1 + e_2) + e_4(e_1 + e_2)].
\end{aligned} \tag{4.63}$$

Since the first two intersection polynomials in the last expression of \tilde{f}_{1210} coincide with

$$(e_4 + e_2)(e_4 + e_3) + e_1(e_3 + e_2),$$

we have

$$\begin{aligned}
\tilde{f}_{1210} &= \\
&e_1((e_4 + e_2)(e_4 + e_3) + e_1(e_3 + e_2))(e_3 + e_4(e_1 + e_2)).
\end{aligned} \tag{4.64}$$

By §1.9.2 we have $f_{1210} = \tilde{f}_{1210}$. \square

2) $\alpha = 321$. Here we have

$$\begin{aligned}
\Gamma_f(\alpha) &= \{(k_3 k_2 k_1) \mid k_3 \in \{1, 4\}, k_2 \in \{3, 4\}, k_1 \in \{2, 3, 4\}, k_1 \neq k_2, k_2 \neq k_3\} = \\
&\{(132), (142), (134), (143), (432), (434)\}
\end{aligned} \tag{4.65}$$

and

$$\begin{aligned}
\tilde{f}_{3210} &= e_1 \sum_{\beta \in \Gamma_f(\alpha)} e_\beta = e_3(e_{132} + e_{134} + e_{143} + e_{432} + e_{434}) = \\
&e_3[e_1(e_4 + e_2)(e_3 + e_2) + e_1(e_4 + e_2)(e_3 + e_4) + \\
&\quad e_1(e_3 + e_2)(e_3 + e_4) + e_4(e_3 + e_2)(e_1 + e_2) + \\
&\quad e_4(e_1 + e_2)(e_3 + e_4)].
\end{aligned} \tag{4.66}$$

Since

$$\begin{aligned}
&e_1(e_4 + e_2)(e_3 + e_4) + e_4(e_1 + e_2)(e_3 + e_4) = \\
&(e_3 + e_4)(e_4 + e_2)(e_1 + e_4(e_1 + e_2(e_3 + e_4))) = \\
&(e_3 + e_4)(e_4 + e_2)(e_1 + e_4)(e_1 + e_2(e_3 + e_4)),
\end{aligned} \tag{4.67}$$

by (4.66) we have

$$\begin{aligned}
\tilde{f}_{3210} = & e_3[e_1(e_4 + e_2)(e_3 + e_2) + \\
& (e_3 + e_4)(e_4 + e_2)(e_1 + e_4)(e_1 + e_2(e_3 + e_4)) + \\
& e_1(e_3 + e_2)(e_3 + e_4) + e_4(e_3 + e_2)(e_1 + e_2)] = \\
& e_3(e_1 + e_2)(e_1 + e_4)[e_1(e_4 + e_2)(e_3 + e_2) + \\
& (e_3 + e_4)(e_4 + e_2)(e_1 + e_2(e_3 + e_4)) + \\
& e_1(e_3 + e_2)(e_3 + e_4) + e_4(e_3 + e_2)] = \\
& e_3(e_1 + e_2)(e_1 + e_4)[e_1(e_4 + e_2)(e_3 + e_2)(e_3 + e_4) + \\
& (e_3 + e_4)(e_4 + e_2)(e_1 + e_2(e_3 + e_4)) + \\
& e_1(e_3 + e_2) + e_4(e_3 + e_2)].
\end{aligned} \tag{4.68}$$

Since

$$e_1(e_4 + e_2)(e_3 + e_2)(e_3 + e_4) \subseteq e_1(e_3 + e_2),$$

by (4.68) we have

$$\begin{aligned}
\tilde{f}_{3210} = & e_3(e_1 + e_2)(e_1 + e_4)[(e_3 + e_4)(e_4 + e_2)(e_1 + e_2(e_3 + e_4)) + \\
& e_1(e_3 + e_2) + e_4(e_3 + e_2)] = \\
& e_3(e_1 + e_2)(e_1 + e_4)[e_1(e_3 + e_4)(e_4 + e_2) + e_2(e_3 + e_4)) + \\
& e_1(e_3 + e_2) + e_4(e_3 + e_2)] = \\
& e_3(e_1 + e_2)(e_1 + e_4)[e_1(e_3 + e_4)(e_4 + e_2)(e_3 + e_2) + e_2(e_3 + e_4)) + \\
& e_1 + e_4(e_3 + e_2)].
\end{aligned} \tag{4.69}$$

Since

$$e_1(e_3 + e_4)(e_4 + e_2)(e_3 + e_2) \subseteq e_1,$$

by (4.69) we have

$$\begin{aligned}
\tilde{f}_{3210} = & e_3(e_1 + e_2)(e_1 + e_4)(e_2(e_3 + e_4)) + e_1 + e_4(e_3 + e_2) = \\
& e_3(e_1 + e_2)(e_1 + e_4)(e_1 + (e_2 + e_4)(e_3 + e_2)(e_3 + e_4)).
\end{aligned} \tag{4.70}$$

By §1.9.2 we have $f_{3210} = \tilde{f}_{3210}$. The proposition is proved. \square

CONJECTURE 4.8.6. For every admissible sequence α , the elements e_α (resp. $f_{\alpha 0}$) and \tilde{e}_α (resp. $\tilde{f}_{\alpha 0}$) coincide without restriction mod θ (see Proposition 4.8.3).

In Propositions 4.8.4 and 4.8.5, this conjecture was proven for small values of lengths of the admissible sequence α .

	Admissible sequence α	Admissible polynomial e_α	Admissible polynomial $f_{\alpha 0}$
$F21$	$(21)^t(41)^r(31)^s =$ $(21)^t(31)^s(41)^r$	$e_2 a_{2s}^{31} a_{2r}^{41} a_{2t-1}^{34}$	$e_\alpha(e_2 a_{2t}^{34} + a_{2r+1}^{41} a_{2s-1}^{31}) =$ $e_\alpha(a_{2t}^{43} + e_1 a_{2r}^{24} a_{2s}^{23})$
$F31$	$(31)^s(41)^r(21)^t =$ $(31)^s(21)^t(41)^r$	$e_3 a_{2t}^{21} a_{2r}^{41} a_{2s-1}^{24}$	$e_\alpha(e_3 a_{2s}^{42} + a_{2r+1}^{41} a_{2t-1}^{21}) =$ $e_\alpha(a_{2s}^{42} + e_1 a_{2r}^{34} a_{2t}^{32})$
$F41$	$(41)^r(31)^s(21)^t =$ $(41)^r(21)^t(31)^s$	$e_4 a_{2t}^{21} a_{2s}^{31} a_{2r-1}^{32}$	$e_\alpha(e_4 a_{2r}^{32} + a_{2s+1}^{31} a_{2t-1}^{21}) =$ $e_\alpha(a_{2r}^{32} + e_1 a_{2s}^{43} a_{2t}^{42})$
$G11$	$1(41)^r(31)^s(21)^t =$ $1(31)^s(41)^r(21)^t =$ $1(21)^t(31)^s(41)^r$	$e_1 a_{2s}^{24} a_{2t}^{34} a_{2r}^{32}$	$e_\alpha(e_1 a_{2r+1}^{32} + a_{2s+1}^{24} a_{2t-1}^{34}) =$ $e_\alpha(a_{2r+1}^{32} + e_4 a_{2s}^{21} a_{2t}^{31})$
$G21$	$2(41)^r(31)^s(21)^t =$ $2(31)^{s+1}(41)^{r-1}(21)^t$	$e_2 a_{2t}^{34} a_{2s+1}^{31} a_{2r-1}^{14} =$ $e_2 a_{2s}^{34} a_{2t-1}^{31} a_{2r+1}^{14}$	$e_\alpha(e_2 a_{2r}^{14} + a_{2s+2}^{31} a_{2t-1}^{34}) =$ $e_\alpha(a_{2r}^{14} + e_3 a_{2s+1}^{21} a_{2t}^{24})$
$G31$	$3(41)^r(21)^t(31)^s =$ $3(21)^{t+1}(41)^{r-1}(31)^s$	$e_3 a_{2s}^{24} a_{2t+1}^{21} a_{2r-1}^{14} =$ $e_3 a_{2s}^{24} a_{2t-1}^{21} a_{2r+1}^{14}$	$e_\alpha(e_3 a_{2r}^{14} + a_{2t+2}^{21} a_{2s-1}^{24}) =$ $e_\alpha(a_{2r}^{14} + e_2 a_{2t+1}^{31} a_{2s}^{34})$
$G41$	$4(21)^t(31)^s(41)^r =$ $4(31)^{s+1}(21)^{t-1}(41)^r$	$e_4 a_{2r}^{32} a_{2s+1}^{31} a_{2t-1}^{12} =$ $e_4 a_{2r}^{32} a_{2s-1}^{31} a_{2t+1}^{12}$	$e_\alpha(e_4 a_{2t}^{12} + a_{2s}^{31} a_{2r+1}^{32}) =$ $e_\alpha(a_{2t}^{12} + e_3 a_{2s+1}^{41} a_{2r}^{42})$
$H11$	$(14)^r(31)^s(21)^t =$ $(14)^r(21)^{t+1}(31)^{s-1} =$ $(13)^s(41)^r(21)^t =$ $(13)^s(21)^{t+1}(41)^{r-1}$ $(12)^{t+1}(41)^r(31)^{s-1} =$ $(12)^{t+1}(31)^{s-1}(41)^r$	$e_1 a_{2r+1}^{23} a_{2s-1}^{24} a_{2t-1}^{34} =$ $e_1 a_{2r-1}^{23} a_{2s-1}^{24} a_{2t+1}^{34} =$ $e_1 a_{2r-1}^{23} a_{2s+1}^{24} a_{2t-1}^{34}$	$e_\alpha(e_1 a_{2r}^{23} + a_{2s}^{24} a_{2t}^{34}) =$ $e_\alpha(e_1 a_{2s}^{24} + a_{2t}^{34} a_{2r}^{23}) =$ $e_\alpha(e_1 a_{2t}^{34} + a_{2r}^{23} a_{2s}^{24})$
$H21$	$2(14)^r(31)^s(21)^t$	$e_2 a_{2r-1}^{14} a_{2s-1}^{31} a_{2t+2}^{34} =$ $e_2 a_{2r-1}^{14} a_{2s+1}^{31} a_{2t}^{34}$	$e_\alpha(e_2 a_{2r}^{14} + a_{2s}^{31} a_{2t+1}^{34}) =$ $e_\alpha(e_2 a_{2t+1}^{34} + a_{2s}^{31} a_{2r}^{14})$
$H31$	$3(14)^r(31)^s(21)^t$	$e_3 a_{2r-1}^{14} a_{2t-1}^{21} a_{2s+2}^{24} =$ $e_3 a_{2r-1}^{14} a_{2t+1}^{21} a_{2s}^{24}$	$e_\alpha(e_3 a_{2r}^{14} + a_{2s+1}^{24} a_{2t}^{21}) =$ $e_\alpha(e_3 a_{2s+1}^{24} + a_{2r}^{14} a_{2t}^{21})$
$H41$	$4(14)^r(31)^s(21)^t$	$e_4 a_{2s-1}^{13} a_{2t-1}^{21} a_{2r+2}^{23} =$ $e_4 a_{2s-1}^{13} a_{2t+1}^{21} a_{2r}^{23}$	$e_\alpha(e_4 a_{2r+1}^{23} + a_{2s}^{13} a_{2t}^{21}) =$ $e_\alpha(e_4 a_{2t}^{21} + a_{2r+1}^{23} a_{2s}^{13})$

TABLE 4.3. Admissible polynomials in the modular lattice D^4

Notes to Table:

- 1) For more details about admissible sequences, see Proposition 4.1.2 and Table 4.1.
- 2) For relations given in two last columns (definitions of admissible polynomials e_α and $f_{\alpha 0}$), see Lemma 4.8.2.
- 3) In each line, each low index should be non-negative. For example, for Line $F21$, we have: $s \geq 0, r \geq 0, t \geq 1$; for Line $G21$, we have: $s \geq 0, r \geq 0, t \geq 0$.

APPENDIX A

Modular and linear lattices

The notion of modular lattice misled logicians and mathematicians for decades. . . most examples of modular lattices occurring in algebra and combinatorics enjoy the stronger property of being linear lattices. Unlike modular lattices, it is not known whether linear lattices can not be defined by identities alone . . . This lack of an abstract definition is perhaps the reason why in the past the theory of linear lattices was subsumed into the theory of modular lattices.

Mainetti M., Yan C. H., [MY99, p.12].

A.1. Distributive lattices and Boolean algebras

A *lattice* is a set L with two commutative and associative operations: a *sum* and an *intersection*. If $a, b \in L$, then we denote the intersection by ab and the sum by $a + b$. More frequently the sum of a and b is denoted by $a \cup b$ or $a \vee b$ and the intersection is denoted by $a \cap b$ or $a \wedge b$. Both operations are *idempotent* and satisfy the *absorbtion law*:

$$\begin{array}{ll} aa = a, & \text{idempotency of intersection,} \\ a + a = a, & \text{idempotency of sum,} \\ a(a + b) = a, & \text{absorbtion law,} \\ a + ab = a, & \text{absorbtion law.} \end{array}$$

The sum and intersection of n elements a_1, \dots, a_n is denoted, respectively, by

$$\sum_{i=1}^n a_i \text{ and } \bigcap_{i=1}^n a_i$$

On the lattice L , an order relation \subseteq (*an inclusion*) is defined:

$$a \subseteq b \iff ab = a \text{ or } a + b = b.$$

A lattice L is said to be *distributive* if the following two *distributive laws* hold

$$\begin{array}{ll} a(b + c) = ab + ac & \text{for every } a, b, c \subseteq L, \\ a + bc = (a + b)(a + c) & \text{for every } a, b, c \subseteq L. \end{array} \tag{A.1}$$

A distributive lattice L is said to be a *Boolean algebra* if L contains a *minimal* element O and a *maximal* element I and, for each element $a \in L$, there exists a unique *complement*

a' such that

$$\begin{aligned} aa' &= O, \\ a + a' &= I, \\ (a')' &= a, \\ (ab)' &= a' + b', \\ (a + b)' &= a'b'. \end{aligned} \tag{A.2}$$

An *atom* a of a Boolean algebra is a minimal nonzero element, i.e., if $x \subseteq a$ then, either $x = O$ or $x = a$. An element y is *atomless* if there is no atom x such that $x \subseteq y$. A Boolean algebra is *atomic* if there are no atomless elements.

FACT A.1.1. *Some infinite Boolean algebras do not contain any atoms. All finite Boolean algebras are atomic, [BS81, p.135], [Hal63].*

A *coatom* a of a Boolean algebra is a maximal nonzero element, i.e., if $x \supseteq a$, then either $x = I$ or $x = a$.

FACT A.1.2 (Stone, [Stn36]). *Every finite Boolean algebra is isomorphic to the Boolean algebra of all subsets of a finite set; in particular, the number of elements of every finite Boolean algebra is a power of two, [Gr98, Ch.2].*

A.2. Modular lattices

A lattice is said to be *modular* if, for every $b, a, c \in L$,

$$a \subseteq b \implies b(a + c) = a + bc. \tag{A.3}$$

The relation (A.2) is said to be *the modular law*. The *modular law* is sometimes referred to as *Dedekind's law*. Every distributive lattice is modular but not conversely.

A set of subspaces of the given vector space R (and, of course, submodules of the given module M) form a modular lattice, namely if $A, B, C \subseteq R$ and $A \subseteq B$, then

$$B(A + C) = A + BC.$$

Indeed, the inclusion $A + BC \subseteq B(A + C)$ holds, because $A + BC \subseteq A + C$ and $A + BC \subseteq B$. Conversely, let $x \in B(A + C)$, i.e., $x \in B$ and $x = x_a + x_c$, where vectors x_a and x_c are components of the vector x , such that $x_a \in A$ and $x_c \in C$. Then $x_c = x - x_a \in B$, i.e., $x_c \in BC$ and $x_a + x_c \in A + BC$.

The main properties of the modular lattices were derived by R. Dedekind [De1897], [Bir48]. We need the two features of the modular lattice, which are called *permutation properties of the modular lattice* or just *permutation properties*.

PROPOSITION A.2.1 (Permutation properties). *In any modular lattice L , for every four elements $A, B, C, D \subseteq L$, the following two permutation properties hold*

$$A(B + CD) = A(CB + D) = A(CB + CD) \text{ for every } A \subseteq C. \tag{A.4}$$

$$A + D(B + C) = A + B(D + C) = A + (D + C)(B + C) \text{ for every } C \subseteq A. \tag{A.5}$$

Proof. By the modular law (A.2) we have

$$\begin{aligned} A(B + CD) &= AC(B + CD) = A(CB + CD) \text{ and} \\ A(CB + D) &= AC(CB + D) = A(CB + CD), \end{aligned}$$

which proves (A.4). Similarly,

$$A + D(B + C) = A + C + D(B + C) = A + (D + C)(B + C) \text{ and}$$

$$A + B(D + C) = A + C + B(D + C) = A + (B + C)(D + C),$$

which proves (A.5). \square

A.3. Hasse diagrams

Hasse diagrams (also called *upward drawings*) are constructed to clear up the lattice by eliminating obvious edges. The vertices are arranged vertically, so the directional arrows are implied (always upwards) and omitted. Any edge implied by transitivity is not shown, i.e., if there is some element x such that $a \subseteq x \subseteq b$, then the edge $[a, b]$ is not shown on the Hasse diagram. On Fig. A.1, two Hasse diagrams are depicted: free distributive lattice D_{18} generated by three generators and free modular lattice M_{28} generated by three generators. Lattices D_{18} and M_{28} are finite, D_{18} contains 18 elements and M_{28} contains 28 elements.





FIGURE A.1. Hasse diagrams of two lattices: the free distributive 3-generated lattice D_{18} and the free modular 3-generated lattice M_{28}

For other examples of Hasse diagrams, see

- The 64-element distributive lattice $H^+(n)$, Fig. 1.5,
- Perfect sublattices in D^4 and $D^{2,2,2}$, Fig. 1.6,
- The 16-element Boolean algebra $U_n \cup V_{n+1}$, Fig. 1.7,
- The Boolean algebras U_n and V_{n+1} , Fig. 3.1,
- The diamond M_3 and the pentagon N_5 , Fig. A.2.
- 4-dimensional cube $D \cup C$, Fig. A.3.

FIGURE A.2. Two Hasse diagrams: the diamond M_3 and the pentagon N_5

A.4. A characterization of distributive and modular lattices

Every distributive lattice L is also modular. Indeed, if $a \subseteq b$, then

$$b(a + c) = ba + bc = a + bc,$$

i.e., (A.3) holds. The converse is false. Consider the diamond lattice M_3 , see Fig. A.2. It easily follows from the modular law (A.3) that the lattice M_3 is modular. However, M_3 is not distributive, because

$$c(a + b) = c, \text{ but } ca + cb = ab, \text{ i.e., } c(a + b) \neq ca + cb, \quad (\text{A.6})$$


see Fig. A.2. Further, the pentagon N_5 is not a modular lattice, because

$$x \subseteq y, \quad y(x + z) = y, \quad x + yz = x, \text{ i.e., } y(x + z) \neq x + yz. \quad (\text{A.7})$$

The following characterization of modular lattices is well-known [Bir48].

PROPOSITION A.4.1. *A lattice L is modular if and only if it does not contain pentagon N_5 as a sublattice.*

Proof. According to (A.7) L does not contain N_5 , since N_5 is not modular.



We will show that, conversely, if L is not modular, then it must contain a pentagon. Let us suppose that there are three elements x, y, z such that

$$\begin{aligned} x &\subseteq y, \\ x + yz &\neq y(x + z), \\ x + yz &\subseteq y(x + z). \end{aligned} \quad (\text{A.8})$$

The last inclusion in (A.8) is true in every lattice.

Consider the sublattice generated by the following elements:

$$yz \subseteq x + yz \subseteq y(x + z) \subseteq x + z \text{ and } z.$$

We will show that this sublattice is isomorphic to N_5 . For sums, we have

$$x + z \subseteq (x + yz) + z \subseteq y(x + z) + z \subseteq x + z,$$

and therefore

$$y(x + z) + z = x + z, \quad (x + yz) + z = x + z.$$

For intersections, we have

$$zy \subseteq z(x + yz) \subseteq zy$$

since $x \subseteq y$. Therefore $z(x + yz) = zy$. Besides, $z(y(x + z)) = zy$. Finally, let us show that

$$yz \neq x + yz.$$


Suppose that $yz = x + yz$; then $x \subseteq y$ and

$$yz = x + yz \subseteq y(x + z) \subseteq y(yz + z) = yz,$$

i.e., $x + yz = y(x + z)$ which contradicts the hypothesis (A.8). \square

PROPOSITION A.4.2 ([Bir48]). *A lattice L is distributive if and only if it does not contain sublattices M_3 and N_5 .*

Proof. According to (A.7), (A.6) any distributive lattice L does not contain M_3 and N_5 , since M_3 and N_5 are not distributive. Suppose, conversely, that L does not contain N_5 . Then by Proposition A.4.1 the lattice L is modular. Let L be not distributive.



We will show that, if L is not distributive, then it must contain a diamond. Suppose that there are three elements a, b, c that sublattice $\{a, b, c\}$ is not distributive. We consider the sublattice generated by the following elements p, r, s :

$$\begin{aligned} p &= (ab + c)(a + b), \\ r &= (ac + b)(a + c), \\ s &= (bc + a)(b + c). \end{aligned} \tag{A.9}$$

Since the modular law holds in L , we have

$$\begin{aligned} p + r &= (ab + c)(a + b) + (ac + b)(a + c) = (a + b)(a + c)(ab + c + ac + b) = \\ &= (a + b)(a + c)(c + b). \end{aligned}$$

Similarly,

$$p + s = r + s = p + r = (a + b)(a + c)(b + c). \tag{A.10}$$

In just the same way,

$$\begin{aligned} pr &= (ab + c)(a + b)(ac + b)(a + c) = (ab + c)(ac + b) = \\ &= ac + b(ab + c) = ac + ab + bc, \end{aligned}$$

and

$$ps = pr = rs = ac + ab + bc. \tag{A.11}$$

We will show now that the elements p, r, s are distinct. Suppose, for example, that $p = r$. Then

$$p + r = p = r = pr$$

and

$$(a + b)(a + c)(b + c) = ac + ab + bc \tag{A.12}$$

Consider now the intersection of both sides of the (A.12) with a :

$$a(b + c) = a(ac + ab + bc) = ac + ab + abc = ac + ab.$$

Similarly,

$$b(a + c) = ba + bc, \quad c(a + b) = ca + cb,$$

which contradicts the hypothesis that $\{a, b, c\}$ is not distributive. \square

The following proposition shows the equivalence of the two distributivity laws.

PROPOSITION A.4.3. *Two following distributivity laws (see (A.1)) are equivalent for an arbitrary lattice L :*

$$\begin{aligned} (i) \quad & a(b + c) = ab + ac \quad \text{for every } a, b, c \in L, \\ (ii) \quad & a + bc = (a + b)(a + c) \quad \text{for every } a, b, c \in L \end{aligned}$$

Proof. Let us prove the implication (i) \implies (ii):

$$(a + b)(a + c) = (a + b)a + (a + b)c = a + (a + b)c = a + ac + bc = a + bc.$$

Conversely, (ii) \implies (i):

$$ab + ac = (ab + c)(ab + a) = (ab + c)a = (a + c)(b + c)a = a(b + c). \quad \square$$

The following criterion of distributivity of the modular lattice is proved by B. Jonnson:

PROPOSITION A.4.4. ([Jo55, Th.5]) *Let A be a modular lattice, p a positive integer, and X_1, X_2, \dots, X_p nonempty linearly ordered subsets of A . For the sublattice of A generated by the set*

$$X_1 \cup X_2 \cup \dots \cup X_p$$

to be distributive it is necessary and sufficient that, for any

$$x_1 \in X_1, x_2 \in X_2, \dots, x_p \in X_p,$$

the sublattice of A generated by the set $\{x_1, x_2, \dots, x_p\}$ be distributive. \square

We use this result for $p = 3$ in the proof of the distributivity of the sublattice of perfect elements $H^+(n)$ in Proposition 3.1.2.

A.5. When the union of two 3D cubes is a 4D cube?

PROPOSITION A.5.1. *Let D and C be two 8-element Boolean algebras with coatoms $d_1, d_2, d_3 \in D$, coatoms $c_1, c_2, c_3 \in C$, maximal elements $I_d \in D$, $I_c \in C$ and minimal elements $O_d \in D$, $O_c \in C$. If the following inclusions hold:*

$$c_i \subseteq d_i, \quad i = 1, 2, 3, \tag{A.13}$$

$$d_i \subseteq c_i + d_j, \quad i, j \in \{1, 2, 3\}, \tag{A.14}$$

$$d_i c_j \subseteq c_i, \quad i \neq j, i, j \in \{1, 2, 3\}, \tag{A.15}$$

then the union $D \cup C$ is a 16-element Boolean algebra with coatoms d_1, d_2, d_3 and I_c .

Proof. Let us prove that the sum does not lead out of D .

$$c \in C, \quad d \in D \implies c + d \in D. \tag{A.16}$$

Case $c = c_i, 1) - 4)$

1) By (A.14) $d_i + d_j \subseteq c_i + d_j = c_i + d_i d_j + d_j = d_i(c_i + d_j) + d_j \subseteq d_i + d_j$, i.e.,

$$c_i + d_j = d_i + d_j = I_d. \tag{A.17}$$

2) From (A.17) we have

$$c_i + d_i d_j = d_i(c_i + d_j) = d_i. \tag{A.18}$$

3) Since $c_j c_k \subseteq d_j d_k$, we get

$$\begin{aligned} c_i + d_j d_k &= c_i c_j + c_i + c_j c_k + d_j d_k = \\ c_j(c_i + c_k) + c_i + d_j d_k &= c_j + c_i + d_j d_k = c_i + (c_j + d_j d_k). \end{aligned}$$

According to (A.18) we have

$$c_i + d_j d_k = c_i + d_j = I_d. \quad (\text{A.19})$$

4) From (A.19) for $O_d = d_i d_j d_k$ we have

$$c_i + d_i d_j d_k = d_i(c_i + d_j d_k) = d_i. \quad (\text{A.20})$$




FIGURE A.3. The 4-dimensional cube $D \cup C$

Case $c = c_i c_j$, 5) – 8)

5) Since $c_j \subseteq d_j$, we get

$$c_i c_j + d_j = d_j. \quad (\text{A.21})$$

6) From (A.21) we obtain

$$c_i c_j + d_k = c_i c_j + c_i c_k + d_k = c_i + d_k = I_d. \quad (\text{A.22})$$

7) Similarly,

$$c_i c_j + d_i d_k = d_i(c_i c_j + d_k) = d_i. \quad (\text{A.23})$$

8) For $O_d = d_i d_j d_k$, we have

$$c_i c_j + d_i d_j d_k = d_i d_j(c_i c_j + d_k) = d_i d_j \quad (\text{A.24})$$

Case $c = c_i c_j c_k = O_c$.

In this case we have $c \subseteq d$ for every $d \in D$ and

$$c_i c_j c_k + d = d \text{ for every } d \in D. \quad (\text{A.25})$$

Case $c = I_c$.

In this case c is the sum of two coatoms $c_i + c_j$, here we have also

$$I_c + d \in D \text{ for every } d \in D. \quad (\text{A.26})$$

We will prove that the intersection does not lead out of C .

$$c \in C, \quad d \in D \implies dc \in C. \quad (\text{A.27})$$

Cases $c = c_i, c = I_c, c = c_i c_j, 9) - 11)$.

9) From (A.15) we have $d_i c_j \subseteq c_i c_j$. From (A.13) we have $c_i c_j \subseteq d_i c_j$, so

$$d_i c_j = c_i c_j. \quad (\text{A.28})$$

10) From (A.28) we deduce that

$$d_i I_c = d_i (c_i + c_j) = c_i + d_i c_j = c_i + c_i c_j = c_i. \quad (\text{A.29})$$

11) Similarly,

$$d_i c_j c_k = (d_i c_j)(d_i c_k) = c_i c_j c_k = O_c. \quad (\text{A.30})$$

12) Finally,

$$d_i d_j c_k = (d_i c_k)(d_j c_k) = c_i c_j c_k = O_c. \quad (\text{A.31})$$

For all other $d = d_i d_j$ or for $d = d_i d_j d_k$ we also have $dc \in C$ for every $c \in C$.

Complementarity in $D \cup C$.

From (A.19) and (A.31) we obtain

$$c'_i = d_j d_k. \quad (\text{A.32})$$

From (A.22) and (A.30) we have

$$d'_i = c_j c_k. \quad \square \quad (\text{A.33})$$

A.6. Representations of the modular lattices

In this section we mostly follow definitions of Gelfand and Ponomarev [GP74].

Let L be a modular lattice, X a finite dimensional vector space, $\mathcal{L}(X)$ the modular lattice of linear subspaces in X . A morphism $\rho : L \rightarrow \mathcal{L}(X)$ is called a *linear representation* of L in the space X . Thus, the representation $\rho : L \rightarrow \mathcal{L}(X)$ maps every element $a \in L$ to the subspace $\rho(a) \subseteq X$ such that, for every $a, b \in L$, we have:

$$\rho(ab) = \rho(a)\rho(b), \quad \rho(a+b) = \rho(a) + \rho(b).$$

Let $\rho_1 : L \rightarrow \mathcal{L}(U)$ and $\rho_2 : L \rightarrow \mathcal{L}(V)$ be representations of L in the spaces U and V . We put

$$\rho(a) = \rho_1(a) \oplus \rho_2(a)$$

for every $a \in L$, where the subspace $\rho(a) \subseteq U \oplus V$ is the set of all pairs

$$\{(u, v) \mid u \in \rho_1(a) \text{ and } v \in \rho_2(a)\}.$$

This correspondence gives the representation ρ in the space $U \oplus V$. The representation ρ is called the *direct sum of representations* $\rho_1 \oplus \rho_2$.

The representation ρ is said to be *decomposable* if it is isomorphic to a direct sum $\rho_1 \oplus \rho_2$ of non-zero representations ρ_1 and ρ_2 . It is easy to prove that the representation ρ in the space X is decomposable if and only if there are subspaces V and U such that $X = U \oplus V$ and

$$\rho(a) = U\rho(a) + V\rho(a) \quad \text{for every}^1 \quad a \in L.$$

The representation ρ is said to be *indecomposable* if there exists an element $a \in L$ such that

$$\rho(a) \neq U\rho(a) + V\rho(a)$$

for every decomposition X into the direct sum $X = U \oplus V$.

Following [GP74] we introduce now the notions of *perfect elements* and *linear equivalence relation*. The element $a \in L$ is said to be *perfect* if either $\rho(a) = 0$ or $\rho(a) = X$ for each

¹Recall that $U\rho(a)$ denotes the intersection of subspaces: $U\rho(a) = U \cap \rho(a)$.

indecomposable representation ρ of the modular lattice L , where X is the representation space of ρ .

Obviously, the sum and intersection of perfect elements is also a perfect element. Thus, perfect elements form a sublattice in the lattice L .

Two elements $a, b \in L$ are called *linearly equivalent* if $\rho(a) = \rho(b)$ for all indecomposable representations $\rho : L \rightarrow \mathcal{L}(X)$ and we write

$$a \equiv b \pmod{\theta}. \quad (\text{A.34})$$

The relation θ is called the *linear equivalence relation*.

We say that a property of L is true *modulo linear equivalence* if the property is true for each indecomposable representation ρ of L .

Let ρ be the representation of the modular lattice L in the vector space X . The subspace $U \subseteq X$ is said to be *admissible with respect to ρ* , if one of following two relations hold

$$\begin{aligned} \text{(i)} \quad & U(\rho(a) + \rho(b)) = U\rho(a) + U\rho(b), \\ \text{(ii)} \quad & U + \rho(a)\rho(b) = (U + \rho(a))(U + \rho(b)) \end{aligned} \quad (\text{A.35})$$

for every $a, b \in L$. The equivalence of (i) and (ii) follows from Proposition A.4.3.

PROPOSITION A.6.1. ([GP74, Prop.2.1]) *Let ρ be a representation of the lattice L in a vector space X , let U be the subspace of X , and $\nabla : X \rightarrow X/U$ the canonical map. Then the following relations are equivalent:*

- (1) *The subspace U is admissible with respect to ρ .*
- (2) *The map $x \mapsto U\rho(x)$ defines a representation in the subspace U .*
- (3) *The mapping $x \mapsto \nabla\rho(x)$ defines a representation in the quotient space X/U .*

Proof. Let us prove the implication (1) \implies (2). If U is admissible with respect to ρ , then

$$U\rho(a + b) = U(\rho(a) + \rho(b)) = U\rho(a) + U\rho(b).$$

Besides,

$$U\rho(ab) = U\rho(a)\rho(b) = (U\rho(a))(U\rho(b)).$$

Thus, we have a representation in the subspace U . The representation in the subspace U is called a *subrepresentation* of ρ and is denoted by ρ_U . \square

Conversely, (2) \implies (1):

$$\begin{aligned} \rho_U(a) + \rho_U(b) &= \rho_U(a + b), \\ U\rho(a) + U\rho(b) &= U\rho(a + b) = U(\rho(a) + \rho(b)). \end{aligned}$$

So, we have (A.35,(i)). \square

Now, consider (1) \implies (3). Set

$$\nu(a) = \rho(a) + U,$$

Then (A.35,(ii)) is equivalent to the relation

$$\nu(a)\nu(b) = \nu(ab). \quad (\text{A.36})$$

Since, $\ker \nabla = U$, we have $\nabla\rho(a) = \nabla\nu(a)$. Further,

$$\nabla\nu(a + b) = \nabla\rho(a + b) = \nabla(\rho(a) + \rho(b)) = \nabla\rho(a) + \nabla\rho(b) = \nabla\nu(a) + \nabla\nu(b),$$

and (3) from Proposition A.6.1 is equivalent to the following relation:

$$(\nabla\nu(a))(\nabla\nu(b)) = \nabla\nu(ab). \quad (\text{A.37})$$

Thus, we just need to prove that (A.37) follows from (A.36). The inclusion

$$(\nabla\nu(a))(\nabla\nu(b)) \supseteq \nabla\nu(ab)$$

is obvious, since $\nu(ab) \subseteq \nu(a), \nu(b)$. So, it is sufficient to prove that

$$(\nabla\nu(a))(\nabla\nu(b)) \subseteq \nabla\nu(ab). \quad (\text{A.38})$$

Let $z \in \nabla\nu(a)$ and $z \in \nabla\nu(b)$. Then there exist vectors $v_a \in \nu(a)$ and $v_b \in \nu(b)$ such that

$$\nabla(v_a) = \nabla(v_b) = z.$$

Then, $w = v_a - v_b \in \ker \nabla = U$, so

$$v_b = v_a - w \in \nu(a) + U = \nu(a),$$

i.e., $v_b \in \nu(a)$, so

$$v_b \in \nu(a)\nu(b) = \nu(ab)$$

and $z \in \nabla\nu(ab)$, and hence (A.38) and (A.37) are proved. \square

Finally, consider the implication (3) \implies (1). We need to prove that (A.36) follows from (A.37). Let $v \in \nu(a)\nu(b)$, i.e.,

$$v \in \nu(a) = \rho(a) + U, \text{ and } v \in \nu(b) = \rho(b) + U.$$

By (A.37) there exists $w \in \nu(ab)$ such that $\nabla(v) = \nabla(w)$. Therefore, $v - w \in U$ and $v \in \nu(ab) + U = \nu(ab)$. Thus, $\nu(a)\nu(b) \subseteq \nu(ab)$. The inverse inclusion is obvious and (A.36) holds. \square

A.7. A relation between representations of graphs and representation of lattices

In a sense, representations of lattices are equivalent to the representations of graphs, [BGP73], [Gab72]. We demonstrate this fact on the example of the graph \tilde{E}_6 .

PROPOSITION A.7.1. *Let ρ be the indecomposable representation of the diagram \tilde{E}_6 , (A.39).*

$$\begin{array}{ccccccc} X_1 & \xrightarrow{g_1} & Y_1 & \xrightarrow{f_1} & X_0 & \xleftarrow{f_2} & Y_2 \xleftarrow{g_2} X_2 \\ & & & & \uparrow & f_3 & \\ & & \rho & & Y_3 & & \\ & & & & \uparrow & g_3 & \\ & & & & X_3 & & \end{array} \quad (\text{A.39})$$

If $X_0 \neq 0$, then maps f_i and g_i are monomorphisms.


Proof. 1) Let g_1 be not a monomorphism and $G = \ker g_1$. Consider the subspace M in X complementary to G in the direct sum $X_1 = G \oplus M$. Then ρ is the direct sum of the following two subrepresentations:

$$\begin{array}{ccccccccc} M & Y_1 & X_0 & Y_2 & X_2 & & G & 0 & 0 & 0 & 0 \\ & & Y_3 & & & \oplus & & 0 & & & \\ & & X_3 & & & & & 0 & & & \end{array} \quad (\text{A.40})$$

Since ρ is an indecomposable representation, we see that ρ coincides with either the first or the second summand in (A.40). Since $X_0 \neq 0$, we deduce that ρ coincides with the first summand in (A.40) and $G = 0$ in the second summand. Thus, g_1 is a monomorphism.

2) Let us prove that f_1 is a monomorphism. Let $K = \ker f_1$ and $G = \text{Im} g_1$. Consider a complementary subspace M to KG in K and a complementary subspace P to KG in G :

$$K = KG \oplus M, \quad G = KG \oplus P. \quad (\text{A.41})$$

FIGURE A.4. Maps f_1 and g_1 . $K = \ker f_1$, $G = \text{Im } g_1$

Then, we have

$$MG = MKG = 0, \quad PK = PKG = 0. \quad (\text{A.42})$$

Since $PK = P(KG + M) = 0$, we see that the subspaces P, M and KG form the direct sum

$$M \oplus P \oplus KG = K + G \quad (\text{A.43})$$

Further, there is a complementary subspace D to $K + G$ in Y_1 :

$$\begin{aligned} Y_1 &= (K + G) \oplus D, \text{ i.e.,} \\ Y_1 &= M \oplus P \oplus KG \oplus D \text{ or} \\ Y_1 &= M \oplus G \oplus D. \end{aligned} \quad (\text{A.44})$$

3) Since we already proved that g_1 is monomorphism and $g_1^{-1}(G \oplus D) = X_1$, we see that ρ in the direct sum of following two subrepresentations:

$$\begin{array}{ccccc} 0 & M & 0 & 0 & 0 \\ & 0 & & & \\ & 0 & & & \end{array} \oplus \begin{array}{ccccc} X_1 & G \oplus D & X_0 & Y_2 & X_2 \\ & & Y_3 & & \\ & & X_3 & & \end{array} \quad (\text{A.45})$$

Since the representation ρ is indecomposable with $X_0 \neq 0$, it follows that $M = 0$. From (A.41) and (A.44) we see that $K = KG$ and $K \subseteq G$, i.e., $\ker f_1 \subseteq \text{Im } g_1$, and therefore

$$\begin{aligned} G &= K \oplus P, \\ Y_1 &= K \oplus P \oplus D. \end{aligned} \quad (\text{A.46})$$

Since

$$X_1 = g^{-1}(K \oplus P) = g^{-1}(K) \oplus g^{-1}(P),$$

we see that the representation ρ is the direct sum of the following two subrepresentations:

$$\begin{array}{ccc} g^{-1}(K) & K & \begin{matrix} 0 & 0 & 0 \\ 0 & & \\ 0 & & \end{matrix} \\ & & \oplus \\ & & \begin{matrix} g^{-1}(P) & P \oplus D & \begin{matrix} X_0 & 0 & 0 \\ Y_3 & & \\ X_3 & & \end{matrix} \end{matrix} \end{array} \quad (\text{A.47})$$

As above, the first summand is zero, $K = \ker f_1 = 0$, $G = P = \text{Im} g_1$ and f_1 is monomorphism. \square

An analog of this proposition for an arbitrary star graph with orientation in which all arrows directed to the branch vertex space X_0 is also true, as one can show.

COROLLARY A.7.2. *The category of representations of the graph \tilde{E}_6 with orientation (A.39) coincides with the category of representations of the lattice $D^{2,2,2}$ in non-zero spaces:*

$$\rho : L \longrightarrow \mathcal{L}(X_0), \quad X_0 \neq 0.$$

In particular, the Coxeter functors Φ^+ , Φ^- which appeared in the representation theory of graphs ([BGP73], [GP74]) can be applied to the representations of lattices. The reflection functors from [BGP73] can not, however, be used because these functors change the orientation of the graph.

The representation ρ for which $\Phi^+\rho = 0$ (resp. $\Phi^-\rho = 0$) is called the *projective* (resp. *injective*). For every indecomposable representation ρ , the new indecomposable representation $\Phi^+\rho$ (resp. $\Phi^-\rho$) can be constructed except for the case where ρ is *projective* (resp. *injective*).

For example, we can construct a new indecomposable representation $\Phi^+\rho$ of \tilde{E}_6 except for seven indecomposable representations ρ_{x_0} and $\rho_{y_i}, \rho_{x_i}, i = 1, 2, 3$; for them $\Phi^+\rho = 0$, see Table A.7.

By [GP79, Prop. 8,9] *the projective indecomposable representations of any graph are naturally enumerated by the vertices of the graph and can be recovered from the orientation of the graph.*

If ρ is indecomposable and not *projective*, i.e., $\Phi^+\rho \neq 0$, then $\rho = \Phi^-\Phi^+\rho$.

If ρ is indecomposable and not *injective*, i.e., $\Phi^-\rho \neq 0$, then $\rho = \Phi^+\Phi^-\rho$.

If ρ is indecomposable and $(\Phi^+)^k\rho \neq 0$, then $\rho = (\Phi^-)^k(\Phi^+)^k\rho$.

If ρ is indecomposable and $(\Phi^-)^k\rho \neq 0$, then $\rho = (\Phi^+)^k(\Phi^-)^k\rho$.

The representation ρ is called *preprojective* if, for some *projective* representation $\tilde{\rho}$,

$$(\Phi^+)^k\rho = (\Phi^+)^k(\Phi^-)^k\tilde{\rho} = \tilde{\rho}, \quad (\Phi^+)^{k+1}\rho = \Phi^+\tilde{\rho} = 0.$$

The representation ρ is called *preinjective* if, for some *injective* representation $\tilde{\rho}$,

$$(\Phi^-)^k\rho = (\Phi^-)^k(\Phi^+)^k\tilde{\rho} = \tilde{\rho}, \quad (\Phi^-)^{k+1}\rho = \Phi^-\tilde{\rho} = 0.$$

The representation ρ is called *regular* if $(\Phi^+)^k\rho \neq 0$ and $(\Phi^-)^k\rho \neq 0$ for every $k \in \mathbb{Z}$.

A.8. The Coxeter functor Φ^+ for $D^{2,2,2}$

Let ρ be an indecomposable representation of \tilde{E}_6 in a space X_0 and $\rho^1 = \Phi^+\rho$ an indecomposable representation of \tilde{E}_6 in the space X_0^1 . According to [BGP73], the Coxeter functor Φ^+ is constructed as the sequence of reflection functors F_z^+ , where z runs over the vertices of \tilde{E}_6 :

$$\Phi^+\rho = F_{x_1}^+ F_{x_2}^+ F_{x_3}^+ F_{y_1}^+ F_{y_2}^+ F_{y_3}^+ F_{x_0}^+ \rho.$$

	ρ	$\Phi^-\rho$	$(\Phi^-)^2\rho$
ρ_{x_0}	0 0 1 0 0 0 0	0 1 2 1 0 1 0	1 2 4 2 1 2 1
ρ_{y_1}	0 1 1 0 0 0 0	1 1 2 1 0 1 0	0 1 3 2 1 2 1
ρ_{y_2}	0 0 1 1 0 0 0	0 1 2 1 1 1 0	1 1 3 2 0 2 1
ρ_{y_3}	0 0 1 0 0 1 0	0 1 2 1 0 1 1	1 1 3 2 1 2 0
ρ_{x_1}	1 1 1 0 0 0 0	0 0 1 1 0 1 0	0 1 2 1 1 1 1
ρ_{x_2}	0 0 1 1 1 0 0	0 1 1 0 0 1 0	1 1 2 1 0 1 1
ρ_{x_3}	0 0 1 0 0 1 1	0 1 1 1 0 0 0	1 1 2 1 1 1 0

TABLE A.1. Preprojective representations. Seven representations in the first column are projective

The reflection functor $F_{x_0}^+$ changes only the space X_0 to X_0^1 and maps $Y_i \longrightarrow X_0$ to maps $\delta_i : X_0^1 \longrightarrow Y_i$ for each $i = 1, 2, 3$. Let

$$\nabla : \{(\eta_1, \eta_2, \eta_3) \mid \eta_i \in Y_i\} \longrightarrow \sum J_i(\eta_i),$$


see Fig. A.5. Then by [BGP73], we have $X_0^1 = \ker \nabla$ from the exact sequence

$$0 \longrightarrow \ker \nabla \longrightarrow \oplus Y_i \xrightarrow{\nabla} X_0 \longrightarrow 0 \quad (\text{A.48})$$

and

$$\delta_i : \{(\eta_1, \eta_2, \eta_3) \mid \eta_i \in Y_i, \sum J_i(\eta_i) = 0\} \longrightarrow \eta_i. \quad (\text{A.49})$$

Now, we have $X_0^1 \xrightarrow{\delta_i} Y_i \xleftarrow{I_i} X_i$. Applying the reflection functor $F_{y_i}^+$, we only change the space Y_i to the space Y_i^1 , the corresponding maps δ_i to μ_i and I_i to P_i for each $i = 1, 2, 3$.

FIGURE A.5. The reflection functor $F_{x_0}^+$

So

$$\begin{aligned} \nabla_i : \{((\eta_1, \eta_2, \eta_3), \xi_i) \mid \eta_i \in Y_i, \Sigma J_i(\eta_i) = 0, \xi_i \in X_i\} &\longrightarrow \\ &\longrightarrow \delta_i(\eta_1, \eta_2, \eta_3) + I_i(\xi) = \eta_i + I_i(\xi_i). \end{aligned}$$

Then $Y_i^1 = \ker \nabla_i$ from the following exact sequences, where $i = 1, 2, 3$.


$$0 \longrightarrow \ker \nabla_i \longrightarrow X_0^1 \oplus X_i \xrightarrow{\nabla_i} Y_i \longrightarrow 0. \quad (\text{A.50})$$

We have,

$$\begin{aligned} Y_i^1 &= \ker \nabla_i = \\ &= \{((\eta_1, \eta_2, \eta_3), \xi_i) \mid \eta_i \in Y_i, \quad \Sigma J_k(\eta_k) = 0, \quad \xi_i \in X_i, \quad \eta_i + I_i(\xi_i) = 0\} \end{aligned} \quad (\text{A.51})$$

and $X_0^1 \xleftarrow{\mu_i} Y_i^1 \xrightarrow{P_i} X_i$. Here $\mu_i : Y_i^1 \longrightarrow X_0^1$,

$$\begin{aligned} \mu_1((\eta_1, \eta_2, \eta_3), \xi_1) &= ((\eta_1, \eta_2, \eta_3) \mid \eta_k \in Y_k, \quad \eta_1 + I_1(\xi_1) = 0, \quad \Sigma J_k(\eta_k) = 0), \\ \mu_2((\eta_1, \eta_2, \eta_3), \xi_2) &= ((\eta_1, \eta_2, \eta_3) \mid \eta_k \in Y_k, \quad \eta_2 + I_1(\xi_2) = 0, \quad \Sigma J_k(\eta_k) = 0), \\ \mu_3((\eta_1, \eta_2, \eta_3), \xi_3) &= ((\eta_1, \eta_2, \eta_3) \mid \eta_k \in Y_k, \quad \eta_3 + I_1(\xi_3) = 0, \quad \Sigma J_k(\eta_k) = 0). \end{aligned} \quad (\text{A.52})$$

FIGURE A.6. The reflection functor $F_{y_3}^+ F_{y_2}^+ F_{y_1}^+ F_{x_0}^+$

REMARK A.8.1. In any representation of the lattice $D^{2,2,2}$, the maps I_k and J_k are monomorphisms and we have $\eta_i = -\xi_i$ in (A.51), i.e., $\eta_i \in X_i$. In other words,

$$\begin{aligned} \mu_1((\eta_1, \eta_2, \eta_3), \xi_1) &= ((-\xi_1, \eta_2, \eta_3) \mid \eta_k \in Y_k, \quad \xi_1 \in X_1, \quad \Sigma \eta_k = 0), \\ \mu_2((\eta_1, \eta_2, \eta_3), \xi_2) &= ((\eta_1, -\xi_2, \eta_3) \mid \eta_k \in Y_k, \quad \xi_2 \in X_2, \quad \Sigma \eta_k = 0), \\ \mu_3((\eta_1, \eta_2, \eta_3), \xi_3) &= ((\eta_1, \eta_2, -\xi_3) \mid \eta_k \in Y_k, \quad \xi_3 \in X_3, \quad \Sigma \eta_k = 0). \end{aligned} \quad (\text{A.53})$$

and $\text{Im}\mu_i = G'_i X_0^1$. This proves that (2.2) from §2.1 is well-defined.

Further,

$$0 \longrightarrow \ker P_i \xrightarrow{Q_i} Y_i^1 \xrightarrow{P_i} X_i \longrightarrow 0 \quad (\text{A.54})$$

and $X_i^1 = \ker P_i$, where

$$\begin{aligned} P_1((\eta_1, \eta_2, \eta_3), \xi_1) &= (\xi_1 \mid \eta_k \in Y_k, \quad \eta_1 + I_1(\xi_1) = 0, \quad \Sigma J_k(\eta_k) = 0)), \\ P_2((\eta_1, \eta_2, \eta_3), \xi_2) &= (\xi_2 \mid \eta_k \in Y_k, \quad \eta_2 + I_1(\xi_2) = 0, \quad \Sigma J_k(\eta_k) = 0)), \\ P_3((\eta_1, \eta_2, \eta_3), \xi_3) &= (\xi_3 \mid \eta_k \in Y_k, \quad \eta_3 + I_1(\xi_3) = 0, \quad \Sigma J_k(\eta_k) = 0)). \end{aligned} \quad (\text{A.55})$$

REMARK A.8.2. For any representation of the lattice $D^{2,2,2}$, the maps J_k are monomorphisms, therefore

$$\begin{aligned} \ker P_1 &= \{((0, \eta_2, \eta_3), 0) \mid \eta_k \in Y_k, \quad \Sigma \eta_k = 0\}, \\ \ker P_2 &= \{((\eta_1, 0, \eta_3), 0) \mid \eta_k \in Y_k, \quad \Sigma \eta_k = 0\}, \\ \ker P_3 &= \{((\eta_1, \eta_2, 0), 0) \mid \eta_k \in Y_k, \quad \Sigma \eta_k = 0\}. \end{aligned} \tag{A.56}$$

Since

$$\begin{array}{ccccccc}
X_1^1 & \xrightarrow{Q_1} & Y_1^1 & \xrightarrow{\mu_1} & X_0^1 & \xleftarrow{\mu_2} & Y_2^1 \xleftarrow{Q_2} X_2^1 \\
& & & & \downarrow \mu_3 & & \uparrow \mu_3 \\
& & & & Y_3^1 & & Y_3^1 \\
& & & & \uparrow Q_3 & & \downarrow P_3 \\
& & & & X_3^1 & & X_3
\end{array}$$

$\Phi^+ \rho = F_{x_3}^+ F_{x_2}^+ F_{x_1}^+ F_{y_3}^+ F_{y_2}^+ F_{y_1}^+ F_{x_0}^+ \rho$

FIGURE A.7. The Coxeter functor Φ^+

$$\begin{aligned}\mu_1 Q_1(\ker P_1) &= \{(0, \eta_2, \eta_3) \mid \eta_k \in Y_k, \quad \Sigma \eta_k = 0\}, \\ \mu_2 Q_2(\ker P_2) &= \{(\eta_1, 0, \eta_3) \mid \eta_k \in Y_k, \quad \Sigma \eta_k = 0\}, \\ \mu_3 Q_3(\ker P_3) &= \{(\eta_1, \eta_2, 0) \mid \eta_k \in Y_k, \quad \Sigma \eta_k = 0\}\end{aligned}\tag{A.57}$$

and $\text{Im}(\mu_i Q_i) = H'_i X_0^1$, it follows that (2.3) from §2.1 is well-defined.

A.9. Linear lattices

In this section, we mostly follow the definitions of the *linear lattices* from works of M. Haiman [Hai85] and M. Mainetti, C. H. Yan [MY99].

A set π of non-empty pairwise disjoint subsets such that their union coincides with A is said to be a *partition* of the set A . The elements of π are called *blocks* of the partition π . If elements $a, b \in A$ lie in the same block π , we write

$$a \equiv b \pmod{\pi}, \text{ or } a \equiv b(\pi), \text{ or } a\pi b.$$

PROPOSITION A.9.1. *There exists a bijection between partitions π of the set A and equivalence relations R_π on the set A .*

Proof. Given a partition π , we define the equivalence relation R_π on the set A as follows:

$$\begin{aligned} \langle a, b \rangle \in R_\pi &\iff a\pi b, \quad \text{or} \\ aR_\pi b &\iff a\pi b. \end{aligned} \quad (\text{A.58})$$

Conversely, if R is an equivalence relation, then the partition π is defined to be:

$$\pi = \{[a]R \mid a \in A\}. \quad \square \quad (\text{A.59})$$

PROPOSITION A.9.2. *The set of partitions $\text{Part}(A)$ of set A constitutes a lattice called the partition lattice.*

Proof. Let π_1, π_2 be partitions of A :

$$\begin{aligned} \pi_1 &= \bigcup_{s \in S} \pi_1^s, \text{ where } \pi_1^{s_1} \cap \pi_1^{s_2} = \emptyset \text{ for all } s_1, s_2 \in S, \\ \pi_2 &= \bigcup_{t \in T} \pi_2^t, \text{ where } \pi_2^{t_1} \cap \pi_2^{t_2} = \emptyset \text{ for all } t_1, t_2 \in T. \end{aligned}$$

For two partitions π_1 and π_2 , their intersection, inclusion and union are defined as follows. Blocks of intersection are pairwise intersections of blocks of π_1 and π_2 .

$$\pi_1 \cap \pi_2 = \bigcup_{s \in S, t \in T} (\pi_1^s \cap \pi_2^t), \text{ for all } s \in S, t \in T. \quad (\text{A.60})$$

Every block of $\pi_1 \subseteq \pi_2$ is included in some block of π_2 .

$$\pi_1 \subseteq \pi_2 = \{x \equiv y(\pi_1) \implies x \equiv y(\pi_2)\}.$$

The union $\pi_1 \cup \pi_2$ is the intersection of all partitions π containing both π_1 and π_2 .

$$\pi_1 \cup \pi_2 = \bigcap_{\pi_1, \pi_2 \subseteq \pi} \pi. \quad \square$$

For more details concerning the *partition lattice*, see [Gr98, Ch.4].

COROLLARY A.9.3. *The set of equivalence relations $\{R_\pi\}$ constitutes the lattice of equivalence relations $\text{Rel}(A)$ isomorphic to the partition lattice*

$$\text{Rel}(A) \simeq \text{Part}(A). \quad (\text{A.61})$$

Proof. It follows from Propositions A.9.1 and A.9.2 \square

Consider now *commuting equivalence relations*. Composition $R \circ T$ of relations R and T is defined as follows:

$$x(R \circ T)y \text{ if and only if there exists } z \in A \text{ such that } xRz \text{ and } zTy. \quad (\text{A.62})$$

Two equivalence relations R and T are said to *commute*¹ if

$$R \circ T = T \circ R.$$

DEFINITION A.9.4 ([MY99], [MY2000]). A *linear lattice* is a sublattice of the lattice of equivalence relations (\simeq partition lattice) on a set, with the property that any two equivalence relations in the sublattice commute, in the sense of composition of relations.

¹It is easy to give an example of non-commuting equivalence relations: if R is “father of” and T is “mother of”, then R and T do not commute, since RT is “maternal grandfather of” and TR is “paternal grandmother of”, [Bz2001].

The classic example of the linear lattice is the lattice of subspaces of a vector space V . Any subspace $S \subseteq V$ determines an equivalence relation R_S on our vector space, given by letting

$$x(R_S)y \text{ if and only if } x - y \in S. \quad (\text{A.63})$$

PROPOSITION A.9.5. 1) All equivalence relations produced, as in (A.63), by subspaces of the vector space V , commute.

2) Commuting equivalence relations given by (A.63) constitute the linear lattice.

Proof. 1) We have

$$x(R_{S_1} \circ R_{S_2})z \text{ if and only if } x - y \in S_1 \text{ and } y - z \in S_2 \text{ for some } y. \quad (\text{A.64})$$

By (A.64), we have

$$x - z \in S_1 + S_2. \quad (\text{A.65})$$

Conversely, suppose $x - z = s_1 + s_2$, where $s_1 \in S_1$ and $s_2 \in S_2$. Let $y = z - s_1$. Then

$$\begin{aligned} x - y &= x - z + s_1 = s_2 \in S_2, \quad \text{and} \\ y - z &= z - s_1 - z = -s_1 \in S_1. \end{aligned} \quad (\text{A.66})$$

Thus, by (A.65), (A.65) and (A.66), we see that

$$x(R_{S_1} \circ R_{S_2})z \text{ if and only if } x - z \in S_1 + S_2.$$

Similarly,

$$x(R_{S_2} \circ R_{S_1})z \text{ if and only if } x - z \in S_2 + S_1,$$

and therefore

$$x(R_{S_1} \circ R_{S_2})z \text{ if and only if } x(R_{S_2} \circ R_{S_1})z,$$

i.e.,

$$R_{S_1} \circ R_{S_2} = R_{S_2} \circ R_{S_1} = R_{S_1 + S_2}. \quad \square \quad (\text{A.67})$$

2) By definition (A.9.4) it suffices to prove that the commuting equivalence relations defined by (A.63) constitute a lattice. The equivalence relation corresponding to the sum of subspaces $S_1 + S_2$ is defined by (A.67).

By (A.9.4) blocks of the partition π_S , corresponding to the equivalence relation R_S are cosets V/S . In other words, the partition π_S consists of the blocks π_S^α as follows

$$\pi_S = \{\pi_S^\alpha \mid x - y \in \pi_S^\alpha \iff x - y \in S\},$$

By (A.60) the partition corresponding to the intersection of two relations S_1 and S_2 consists of cosets $V/(S_1 \cap S_2)$, and

$$\pi_{S_1} \cap \pi_{S_2} = \pi_{S_1 \cap S_2},$$

i.e.,

$$R_{S_1} \cap R_{S_2} = R_{S_1 \cap S_2}. \quad \square$$

REMARK A.9.6. There are lots of linear lattices. Proposition A.9.5 can be generalized to groups. Given any group G , each normal subgroup H determines an equivalence relation on G , which we call R_H , such that

$$x(R_H)y \quad \text{if and only if} \quad xy^{-1} \in H.$$

So the lattice of normal subgroups of G becomes a linear lattice. Similarly, lattices of ideals of a ring are also linear lattices, see Fig. 1.1.

DEFINITION A.9.7 ([Jo53], [Sch45]). Let $a_0, a_1, a_2, b_0, b_1, b_2$ be elements of the modular lattice L , and let

$$\begin{aligned} z_{ij} &= (a_i + a_j)(b_i + b_j) \text{ for } 0 \leq i < j < 3, \\ z &= z_{01}(z_{02} + z_{12}). \end{aligned} \tag{A.68}$$

The identity

$$(a_0 + b_0)(a_1 + b_1)(a_2 + b_2) \subseteq a_0(z + a_1) + b_0(z + b_1) \tag{A.69}$$

is called the Arguesian law and lattices satisfying the Arguesian law are called *Arguesian lattices*. For another version of the Arguesian law, see [MY2000, p.3] and [Hai85]. Identity (A.69) is a lattice-theoretic form of the Desargues theorem of projective geometry, see [Sch45], [MY2000, p.2], [Gr98, Ch.4].

REMARK A.9.8. Every Arguesian lattice is modular, see [Jo54, Th.1.9]. There exists modular lattices which are not Arguesian, see [Gr98, Ch.4.5]. Linear lattices are Arguesian, but the converse is not true: in [Hai91] M. Haiman constructed a family of Arguesian lattices which cannot be represented as linear lattices, see [KY2003], see Fig. 1.1.

A.10. Grassmann-Cayley algebras and modular lattices

A.10.1. A Peano space and the bracket. A Grassmann-Cayley algebra (or *double algebra*) is essentially the exterior algebra of a vector space, equipped with **two** operations, *join*¹ \vee and *meet* \wedge , which are the algebraic analogs of the *union*² \cup and *intersection* \cap of subspaces of a vector space:

$$\begin{array}{lll} \text{the join } \vee & \longleftrightarrow & \text{the union } \cup \text{ (the sum } + \text{)} \\ \text{the meet } \wedge & \longleftrightarrow & \text{the intersection } \cap \end{array} \tag{A.70}$$

These operations were introduced by Grassmann [Gra11], under the name of *progressive* and *regressive* products. In works of G.-C.Rota and his coworkers, the Grassmann work has been recognized and extended, see [DRS76], [BBR85].

Barnabei, Brini and Rota [BBR85, p. 121] write “To the best of our knowledge of published work, the first mathematicians to understand, albeit imperfectly, the program *Ausdehnungslehre*³ were Clifford and Schröder, first, and later A. N. Whitehead, Élie Cartan, and best of all Giuseppe Peano. ... It was Schröder, in an appendix to his *Algebra and Logik*, who first stressed the analogy between the algebra of *progressive* and *regressive* products, and the algebra of sets with *union* and *intersection*.”

By analogy to a *Hilbert space* endowed with a symmetric bilinear form (x, y) , called the inner product, and to a *symplectic space* endowed with an antisymmetric bilinear form $[x, y]$, a *Peano space* is a vector space V of dimension n , endowed with an antisymmetric non-degenerate multilinear form $[x_1, \dots, x_n]$ defined for $x_i \in V$, called the *bracket* [BBR85].

¹The classical Grassmann algebra is the exterior algebra of a vector space, equipped with only one operation, the wedge product \wedge . Barnabei, Brini and Rota [BBR85, p. 127] write “The product in the exterior algebra is called the *join* and is denoted by the symbol \vee . We note that the usage is at variance with ordinary usage, where exterior multiplication is denoted by the wedge \wedge . This departure from common usage will be amply justified in the sequel.” For motivation of the usage the symbol \vee , see §A.10.2.

²This lattice-theoretic operation union \cup does not coincide with the set-theoretic operation union \sqcup . For vector spaces, the operation union \cup is the linear span of the union \sqcup as follows:

$$A \cup B = \overline{A \sqcup B}.$$

The equivalent notation frequently used instead the operation union \cup is the operation *sum* $+$.

³Grassmann’s program *Ausdehnungslehre* refers to 1844, see [Gra11]. The English translation of *Ausdehnungslehre* is *geometric calculus*. Grassmann’s idea was to develop a calculus for the *join* and *meet*. In [BBR85] authors consider Grassmann’s geometric calculus and its relevance in invariant theory.

The group of all linear transformations preserving the bracket is $SL(V)$, the special linear group. The algebra of $SL(V)$ -invariants is generated by brackets (the first fundamental theorem of invariant theory, see [Weyl46, Th. II.6.A]) or [Dol2003, Th. 2.1]).

Let K be the ground field. A *bracket* (of step n) over the space V is a non-degenerate alternating n -linear form, i.e., a function

$$x_1, \dots, x_n \mapsto [x_1, \dots, x_n] \in K,$$

where vectors x_1, \dots, x_n lie in the vector space V , with the following properties:

- (1) $[x_1, \dots, x_n] = 0$ if at least two among the x 's coincide;
- (2) for every $x, y \in V$ and $\alpha, \beta \in K$

$$[x_1, \dots, x_{i-1}, \alpha x + \beta y, x_{i+1}, \dots, x_n] =$$

$$\alpha [x_1, \dots, x_{i-1}, x, x_{i+1}, \dots, x_n] + \beta [x_1, \dots, x_{i-1}, y, x_{i+1}, \dots, x_n];$$
- (3) there exists a basis $\{b_1, \dots, b_n\}$ of V such that $[b_1, \dots, b_n] \neq 0$.

Let $F(V)$ be the free associative algebra with unity over K generated by the elements of a Peano space V . The *exterior algebra* $E(V)$ of the Peano space V is obtained as the quotient of $F(V)$ modulo the ideal generated by v^2 , for all $v \in V$, see, for example, [MY99, Def. 2.1] or [BBR85, Prop. 3.6]. For further properties of the exterior algebra see, e.g., [Gre67]. For exterior Grassmann algebra in the context of supersymmetries, see, e.g., [Ber87] or [L05].

Let

$$\Phi : F(V) \rightarrow E(V),$$

denote the canonical projection. If $x_1 x_2 \dots x_k$ is a word in $F(V)$, we denote its image by

$$\Phi(x_1 x_2 \dots x_k) = x_1 \vee x_2 \vee \dots \vee x_k. \quad (\text{A.72})$$

The element $x_1 \vee x_2 \vee \dots \vee x_k$ is called an *extensor of step k* , or *decomposable antisymmetric tensor* or *decomposable k -vector*. The join of two extensors is an extensor. A linear combination of extensors of step k is called an *antisymmetric tensor of step k* , [BBR85].

PROPOSITION A.10.1. (Properties of the join, [BBR85, Prop. 3.1-3.4]) 1) For every permutation σ of $\{1, 2, \dots, k\}$ we have

$$x_{\sigma(1)} \vee x_{\sigma(2)} \vee \dots \vee x_{\sigma(k)} = \text{sign}(\sigma) x_1 \vee x_2 \vee \dots \vee x_k. \quad (\text{A.73})$$

2) Let $E_k(V)$ be the subspace of $E(V)$ generated by all extensors of step k . Let $A \in E_j(V)$, $B \in E_k(V)$. Then

$$B \vee A = (-1)^{jk} A \vee B. \quad (\text{A.74})$$

3) $\dim E_k(V) = \binom{n}{k}$ for $0 \leq k \leq n$, and hence $\dim E(V) = 2^n$.

4) Let B be a dimension $k > 0$ subspace of V , let $\{x_1, x_2, \dots, x_k\}$ and $\{y_1, y_2, \dots, y_k\}$ are two bases of B . Then

$$x_1 \vee x_2 \vee \dots \vee x_k = C y_1 \vee y_2 \vee \dots \vee y_k \quad (\text{A.75})$$

for some non-zero scalar C .

By Proposition A.10.1, heading 4) there exists one-to-one correspondence (up to non-zero scalar multiple) between non-trivial subspaces of V and non-zero extensor uniquely representing this subspace. Such a representing extensor is said to be *associated* to the subspace.

A.10.2. Motivation of the usage the symbol \vee . Here we formulate the main property of the operation join \vee connecting it with the lattice-theoretic operation sum $+$ (or, equivalently, with the union \cup), see [BBR85, Prop. 3.5].

PROPOSITION A.10.2. *Let A, B be two subspaces of V associated to extensors F and G , respectively. Then*

- (i) $F \vee G = 0$ if and only if $A \cap B \neq 0$;
- (ii) if $A \cap B = 0$, then the extensor $F \vee G$ is associated to the subspace generated by $A + B$.

Reformulate Proposition A.10.2 as follows. Let

$$\begin{aligned} F &= f_1 \vee f_2 \vee \cdots \vee f_j, \\ G &= g_1 \vee g_2 \vee \cdots \vee g_k, \end{aligned} \tag{A.76}$$

and \overline{F} (resp. \overline{G}) denotes the linear span of extensor F (resp. G).

If vectors

$$f_1, f_2, \dots, f_j, g_1, g_2, \dots, g_k$$

are distinct and linearly independent, i.e. $\overline{F} \cap \overline{G} = 0$, then

$$\overline{F \vee G} = \overline{F} + \overline{G}. \tag{A.77}$$

Thus, the join operation \vee on extensors represents the geometric operation of the lattice sum of subspaces of V at least for independent vectors. This is one of motivations to use the symbol \vee .

Another motivation is a discovery of a class of identities that hold in the exterior algebras, which turn into the classical theorems of projective geometry on the incidence of subspaces of projective spaces, and therefore to identities in modular lattices, see (A.70).

For more details concerning these identities, see [Haw94], [Haw96], [MY2000], [MY99].

Theorems such as Desargues, Pappus, Bricard, and Fontené of classical projective geometry are revealed to be expressible as simple and elegant identities holding among joins and meets of extensors in the framework of the Grassmann-Cayley algebra, [BBR85], [MY99], [MY2000]. We suggest to call the transformation of modular lattice polynomials to Grassmann-Cayley algebra polynomials and *vice versa* by the substitution (A.70) the *Rota rendering*. The Rota rendering establishes a connection between the theory of Grassmann-Cayley algebras and the modular (and even linear) lattices.

A.10.3. The meet. A similar operation on extensors should be defined for the lattice operation intersection of two subspaces. This is the *meet* operation. Let extensors F and G are given by (A.76) with $j + k \geq n$, where n is dimension of a *Peano space* V . Then

$$F \wedge G = \sum_{\sigma} \text{sign}(\sigma) [a_{\sigma(1)}, \dots, a_{\sigma(d-k)}, b_1, \dots, b_k] a_{\sigma(d-k+1)} \cdots a_{\sigma(j)}. \tag{A.78}$$

The meet is associative and anti-commutative in the following sense ([BBR85, Prop. 4.4]):

$$F \wedge G = (-1)^{(d-k)(d-j)} G \wedge F. \tag{A.79}$$

Both the join and meet are defined on extensors only, but the definitions are extended to arbitrary elements of $E(V)$ by distributivity.

PROPOSITION A.10.3. ([BBR85, Prop. 4.3], [MY99, Prop. 2.5]) 1) We have

$$F \wedge G = 0 \text{ if and only if } F + G \text{ does not span } V.$$

2) If $F + G$ spans V , then¹

$$\overline{F \wedge G} = \overline{F} \cap \overline{G}. \quad (\text{A.80})$$

Thus the meet operation corresponds to the intersection \cap . A basis $\{e_1, \dots, e_n\}$ in the space V is said to be *unimodular* if $[e_1, \dots, e_n] = 1$. The extensor

$$E = e_1 \vee e_2 \vee \dots \vee e_n$$

in $E(V)$ is called the *integral*. Indeed, it corresponds to the Berezin integral on supermanifolds, cf. [Ber87], [L05]. The integral is well defined and does not depend on the choice of a unimodular basis.

A.10.4. Note on Howe-Huang projective invariants of quadruples. Projective invariants of quadruples were obtained by Howe R., Huang R. in [HH96]. They gave an explicit set of generators for the ring of invariants of a set of four subspaces in a projective space.

REMARK A.10.4. For other results in this direction, see [Gro98], [Rin80]. In [Gro98], Grosshans gave a complete description of the relationship between invariants and quadruples. Invariants of the tame quivers were obtained by Ringel in [Rin80].

Among projective invariants of quadruples obtained by Howe-Huang, the invariants of type III ([HH96, Th. 14]) can be transformed to the polynomials $f_{GC}(E_1, E_2, E_3, E_4)$ in four subspaces E_1, E_2, E_3, E_4 under the join \vee and meet \wedge in the framework of the Grassmann-Cayley algebra. The transformation of Howe-Huang projective invariants to polynomials $f_{GC}(E_1, E_2, E_3, E_4)$ is performed by means of the *Cayley factorization* in the sense of [Wh91].

Further, applying the Rota rendering (A.70) to $f_{GC}(E_1, E_2, E_3, E_4)$, we get polynomials $f_{ML}(E_1, E_2, E_3, E_4)$ in the modular lattice D^4 . It may be shown that *polynomials* $f_{ML}(E_1, E_2, E_3, E_4)$ are *admissible elements* in D^4 .

¹Compare with (A.77).

APPENDIX B

A proof of the theorem on admissible classes

We will briefly write

$$\begin{aligned}
 a &\xrightarrow{\varphi_i} b \quad \text{instead of} \quad \varphi_i \Phi^+ \rho(a) = \rho(b), \\
 a &\xrightarrow{\varphi_i} b \xrightarrow{\varphi_j} c \quad \text{instead of} \quad \varphi_i \Phi^+ \rho(a) = \rho(b), \quad \varphi_j \Phi^+ \rho(b) = \rho(c), \\
 &\dots \\
 a &\xrightarrow{\varphi_i} b \dots c \xrightarrow{\varphi_i} d \quad \text{instead of} \quad \varphi_i \Phi^+ \rho(a) = \rho(b), \dots, \varphi_j \Phi^+ \rho(c) = \rho(d),
 \end{aligned} \tag{B.1}$$

see (1.8), (1.9).

B.1. A proof of the theorem on admissible elements in lattice $D^{2,2,2}$

We will now prove the main property of admissible elements.

Theorem 2.12.1 *Let $\alpha = i_n i_{n-1} \dots 1$ be an admissible sequence and $i \neq i_n$. Then $i\alpha$ is admissible and, for $z_\alpha = f_\alpha, e_\alpha, g_{\alpha 0}$ from the Table 2.3, the following relation holds:*

$$\varphi_i \Phi^+ \rho(z_\alpha) = \rho(z_{i\alpha}). \tag{B.2}$$

Proof of Theorem 2.12.1 For the cases of Lines 1–6, it suffices to prove only

$$z_\gamma \xrightarrow{\varphi_3} z_{3\gamma} \quad \text{and} \quad z_{3\gamma} \xrightarrow{\varphi_1} z_{(13)\gamma}, \tag{B.3}$$

because the relations

$$z_{(13)\gamma} \xrightarrow{\varphi_2} z_{(213)\gamma} \xrightarrow{\varphi_3} z_{3(213)\gamma} \xrightarrow{\varphi_1} z_{13(213)\gamma} \xrightarrow{\varphi_2} z_{(213)(213)\gamma}$$

follow from (B.3) as the permutation $1 \rightarrow 3 \rightarrow 2 \rightarrow 1$ is applied to Lines 1–3 of Table 2.3 from §2.12.

B.1.1. Case $z_\gamma \xrightarrow{\varphi_3} z_{3\gamma}$.

(F) Line 1. We will show that

$$\begin{aligned}
 f_\gamma &\xrightarrow{\varphi_3} f_{3\gamma}, \\
 f_\gamma &= y_1 y_2 a_q^{13} A_{k-1}^{32} \xrightarrow{\varphi_3} y_3 (x_1 + x_2) A_q^{23} a_{k-1}^{31} = f_{3\gamma}.
 \end{aligned} \tag{B.4}$$

Applying Corollary 2.4.4 to $y_1 y_2$ and Corollary 2.7.4 to a_q^{13} and A_{k-1}^{32} we have

$$\begin{aligned}
 y_1 y_2 &\xrightarrow{\varphi_3} y_3 (y_1 + y_2), \\
 a_q^{13} &\xrightarrow{\varphi_3} y_3 A_q^{23}, \\
 A_{k-1}^{32} &\xrightarrow{\varphi_3} y_3 (y_1 + y_2) a_{k-1}^{31}.
 \end{aligned} \tag{B.5}$$

Since a_q^{13} and A_{k-1}^{32} are the φ_3 –homomorphic polynomials (Theorem 2.8.1), we see that “intersection of relations” (B.5) gives us relation (B.4). If $q = 1$, then

$$a_q^{13} \xrightarrow{\varphi_3} y_3 (y_1 + y_2) A_q^{23} \subseteq y_3 (y_1 + y_2)$$

and we have the same “intersection of relations”. \square

(E) Line 1. We want to show that

$$\begin{aligned} e_\gamma &\xrightarrow{\varphi_3} e_{3\gamma}, \\ e_\gamma &= y_2 A_{k-1}^{32} a_q^{21} A_k^{13} a_q^{13} \xrightarrow{\varphi_3} y_3 a_{k-1}^{31} A_{q+1}^{12} a_k^{12} A_q^{23} = e_{3\gamma}. \end{aligned} \quad (\text{B.6})$$

By Corollary 2.7.4 we have

$$\begin{aligned} y_2 a_q^{21} &\xrightarrow{\varphi_3} y_3 A_{q+1}^{12}, \\ A_{k-1}^{32} &\xrightarrow{\varphi_3} y_3 (y_1 + y_2) a_{k-1}^{31}, \\ A_k^{13} &\xrightarrow{\varphi_3} y_3 (y_1 + x_2) a_k^{12}, \\ a_q^{13} &\xrightarrow{\varphi_3} y_3 A_q^{23}. \end{aligned} \quad (\text{B.7})$$

Since a_q^{13}, A_{k-1}^{32} are φ_3 -homomorphic and A_k^{13} is (φ_3, y_2) -homomorphic (Theorem 2.8.1), and $y_3 A_{q+1}^{12} \subseteq y_3 (y_1 + x_2)$, we see that (B.6) comes out from (B.7). As in the previous case, we have the same result for $q = 1$. \square

(G) Line 1. It will be shown that

$$\begin{aligned} g_{\gamma 0} &\xrightarrow{\varphi_3} g_{3\gamma 0}, \\ g_{\gamma 0} &= e_\gamma (x_1 + a_q^{32} A_k^{32}) \xrightarrow{\varphi_3} e_{3\gamma} (y_2 y_3 + A_q^{12} a_k^{31}) = g_{3\gamma 0}. \end{aligned} \quad (\text{B.8})$$

By (B.6) we have $e_\gamma \xrightarrow{\varphi_3} e_{3\gamma}$. By basic relations (2.4.1) and atomic multiplicativity (2.5.2) we have

$$\psi_3(x_1 + a_q^{32} A_k^{32}) = \psi_3(x_1) + \psi_3(a_q^{32}) \varphi_3(A_k^{32}).$$

According to Proposition 2.7.1 (the joint map actions) we obtain

$$\varphi_3(x_1 + a_q^{32} A_k^{32}) = \nu^0(y_2 y_3) + \nu^0(y_3 A_q^{12} (y_3 (y_1 + y_2) a_k^{31})) = \nu^0(y_2 y_3 + y_3 A_q^{12} a_k^{31}).$$

Therefore by Proposition 2.4.3 (the connection) we see that $x_1 + a_q^{32} A_k^{32}$ is φ_3 -homomorphic and

$$x_1 + a_q^{32} A_k^{32} \xrightarrow{\varphi_3} y_2 y_3 + y_3 A_q^{12} a_k^{31} = y_3 (y_2 y_3 + A_q^{12} a_k^{31}).$$

By (B.6) $e_{3\gamma} \subseteq y_3$, hence we get (B.8). \square

B.1.2. Case $z_{3\gamma} \xrightarrow{\varphi_1} z_{(13)\gamma}$.

(F) Line 2. We will show that

$$\begin{aligned} f_{3\gamma} &\xrightarrow{\varphi_1} f_{(13)\gamma}, \\ f_{3\gamma} &= y_3 (x_1 + x_2) A_q^{23} a_{k-1}^{31} \xrightarrow{\varphi_1} y_3 y_1 a_{q+1}^{32} A_{k-1}^{21} = f_{(13)\gamma} \end{aligned} \quad (\text{B.9})$$

By Corollary 2.7.4

$$\begin{aligned} x_1 + x_2 &\xrightarrow{\varphi_1} y_1 y_3, \\ y_3 A_q^{23} &\xrightarrow{\varphi_1} y_1 a_{q+1}^{32}, \\ a_{k-1}^{31} &\xrightarrow{\varphi_1} y_1 A_{k-1}^{21}. \end{aligned} \quad (\text{B.10})$$

As above, a_{k-1}^{31} and $x_1 + x_2$ are φ_1 -homomorphic (Theorem 2.8.1) then “intersection of relations” (B.10) gives us relation (B.9). For $k - 1 = 1$ we have $a_{k-1}^{31} \xrightarrow{\varphi_1} y_1 (y_2 + y_3) A_{k-1}^{21}$. Since $a_{q+1}^{32} \subseteq y_2 + y_3$, we have the same result. \square

(E) Line 2. It will be shown that

$$\begin{aligned} e_{3\gamma} &\xrightarrow{\varphi_1} e_{(13)\gamma}, \\ e_{3\gamma} &= y_3 a_{k-1}^{31} A_{q+1}^{12} a_k^{12} A_q^{23} \xrightarrow{\varphi_1} y_1 A_{k-1}^{21} a_{q+1}^{13} A_k^{32} a_{q+1}^{32} = e_{(13)\gamma}. \end{aligned} \quad (\text{B.11})$$

By Corollary 2.7.4 we have

$$\begin{aligned} y_3 A_q^{23} &\xrightarrow{\varphi_1} y_1 a_{q+1}^{32}, \\ a_{k-1}^{31} &\xrightarrow{\varphi_1} y_1 A_{k-1}^{21}, \\ a_k^{12} &\xrightarrow{\varphi_1} y_1 A_k^{32} \subseteq y_1(y_2 + y_3), \\ A_{q+1}^{12} &\xrightarrow{\varphi_1} y_1(y_2 + y_3) a_{q+1}^{13}. \end{aligned} \quad (\text{B.12})$$

The atomic elements a_{q-1}^{13} , a_k^{12} and A_{q+1}^{12} are φ_1 -homomorphic (Theorem 2.8.1). Then (B.11) follows from (B.12). As in the previous case for $k - 1 = 1$ we have the same result. \square

(G) Line 2. It will be shown that

$$\begin{aligned} g_{3\gamma 0} &\xrightarrow{\varphi_1} g_{(13)\gamma 0}, \\ g_{3\gamma} &= e_{3\gamma}(y_2 y_3 + A_q^{12} a_k^{31}) \xrightarrow{\varphi_1} e_{(13)\gamma}(x_3 + a_{q+1}^{21} A_k^{21}) = g_{(13)\gamma}. \end{aligned} \quad (\text{B.13})$$

Relation (B.13) follows from

$$\begin{aligned} y_2 y_3 &\xrightarrow{\varphi_1} y_1(x_2 + x_3), \\ A_q^{12} &\xrightarrow{\varphi_1} y_1(y_2 + y_3) a_q^{13}, \\ a_k^{31} &\xrightarrow{\varphi_1} y_1 A_k^{21} \end{aligned} \quad (\text{B.14})$$

since

$$\begin{aligned} y_1(x_2 + x_3) + y_1(y_2 + y_3) a_q^{13} A_k^{21} &= y_1(y_2 + y_3)(x_2 + x_3 + y_1 a_q^{13} A_k^{21}) = \\ y_1(y_2 + y_3)(A_k^{21}(x_2 + y_1 a_q^{13}) + x_3) &= y_1(y_2 + y_3)(A_k^{21} a_{q+1}^{21} + x_3) \end{aligned}$$

and $e_{(13)\gamma} \subseteq y_1(y_2 + y_3)$. Again, as in two previous cases if $k = 1$ for a_k^{31} , then

$$a_k^{31} \xrightarrow{\varphi_1} y_1(y_2 + y_3) A_k^{21}$$

with the same result. \square

We will omit the case $a_1^{ij} \xrightarrow{\varphi_j} y_j(y_i + y_k) A_1^{kj}$ (Corollary 2.7.4, (2), $k = 1$) for all further cases of the theorem. As above, for the cases of Lines 9 – 14 it suffices to prove only

$$z_\beta \xrightarrow{\varphi_3} z_{3\beta} \quad \text{and} \quad z_{3\beta} \xrightarrow{\varphi_i} z_{(13)\beta}, \quad (\text{B.15})$$

because the following relation

$$z_{(13)\beta} \xrightarrow{\varphi_2} z_{(213)\beta} \xrightarrow{\varphi_3} z_{3(213)\beta} \xrightarrow{\varphi_1} z_{13(213)\beta} \xrightarrow{\varphi_2} z_{(213)(213)\beta}$$

follows from (B.15) as the permutation $1 \rightarrow 3 \rightarrow 2 \rightarrow 1$ is applied to Lines 9 – 11 of Table 2.3. We will omit references to the φ_i -homomorphic Theorem (Theorem 2.8.1) and “intersection of relations” in §§ B.1.1-B.1.2.

B.1.3. Case $z_\beta \xrightarrow{\varphi_3} z_{3\beta}$.

(F) Line 9. It should be shown that

$$\begin{aligned} f_\beta &\xrightarrow{\varphi_3} f_{3\beta}, \\ f_\beta &= y_2 y_3 a_q^{21} A_k^{13} \xrightarrow{\varphi_3} x_3 A_{q+1}^{12} a_k^{12} = f_{3\beta}. \end{aligned} \tag{B.16}$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} y_3 &\xrightarrow{\varphi_3} x_3(y_1 + y_2), \\ y_2 a_q^{21} &\xrightarrow{\varphi_3} y_3 A_{q+1}^{12} \subseteq x_2 + y_1, \\ y_2 A_k^{13} &\xrightarrow{\varphi_3} y_3(x_2 + y_1) a_k^{12}. \end{aligned} \tag{B.17}$$

Then B.16 follows from B.17 \square

(E) Line 9. We want to prove that

$$\begin{aligned} e_\beta &\xrightarrow{\varphi_3} e_{3\beta}, \\ e_\beta &= y_2 A_k^{32} a_q^{21} A_k^{13} a_{q+1}^{13} \xrightarrow{\varphi_3} y_3 a_k^{31} A_{q+1}^{12} a_k^{12} A_{q+1}^{23} = e_{3\beta}. \end{aligned} \tag{B.18}$$

It follows from the following relations (Corollary 2.7.4):

$$\begin{aligned} A_k^{32} &\xrightarrow{\varphi_3} y_3(y_1 + y_2) a_k^{31}, \\ y_2 a_q^{21} &\xrightarrow{\varphi_3} y_3 A_{q+1}^{12} \subseteq x_2 + y_1, \\ y_2 A_k^{13} &\xrightarrow{\varphi_3} y_3(x_2 + y_1) a_k^{12}, \\ a_{q+1}^{13} &\xrightarrow{\varphi_3} y_3 A_{q+1}^{23}. \quad \square \end{aligned} \tag{B.19}$$

(G) Line 9. We need to show

$$\begin{aligned} g_\beta 0 &\xrightarrow{\varphi_3} f_{3\beta 0}, \\ g_{\beta 0} &= e_\beta(x_2 + a_q^{13} A_{k+1}^{13}) \xrightarrow{\varphi_3} e_{3\beta}(y_1 y_3 + A_q^{23} a_{k+1}^{12}) = g_{3\beta 0}. \end{aligned} \tag{B.20}$$

We have

$$e_\beta(x_2 + a_q^{13} A_{k+1}^{13}) = e_\beta(x_2 + y_2 a_q^{13} A_{k+1}^{13})$$

and

$$e_{3\beta}(y_1 y_3 + A_q^{23} a_{k+1}^{12}) = e_{3\beta}(y_1 y_3 + y_3(x_2 + y_1) A_q^{23} a_{k+1}^{12}),$$

and the following relations (Corollary 2.7.4) are true:

$$\begin{aligned} x_2 &\xrightarrow{\varphi_3} y_1 y_3, \\ a_q^{13} &\xrightarrow{\varphi_3} y_3 A_q^{23}, \\ y_2 A_{k+1}^{13} &\xrightarrow{\varphi_3} y_3(x_2 + y_1) a_{k+1}^{12} \subseteq A_1^{12} \subseteq e_{3\beta}. \end{aligned} \tag{B.21}$$

Then relation (B.20) follows from the (B.21). \square

B.1.4. Case $z_{3\beta} \xrightarrow{\varphi_1} z_{(13)\beta}$.

(F) Line 10. It will be shown that

$$\begin{aligned} f_{3\beta} &\xrightarrow{\varphi_1} f_{(13)\beta}, \\ f_{3\beta} &= x_3 A_{q+1}^{12} a_k^{12} \xrightarrow{\varphi_1} y_1 y_2 a_{q+1}^{13} A_k^{32} = f_{(13)\beta}. \end{aligned} \quad (\text{B.22})$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} x_3 &\xrightarrow{\varphi_1} y_1 y_2, \\ A_{q+1}^{12} &\xrightarrow{\varphi_1} y_1 (y_2 + y_3) A_{q+1}^{13}, \\ a_k^{12} &\xrightarrow{\varphi_1} A_k^{32} \subseteq y_2 + y_3. \end{aligned} \quad (\text{B.23})$$

Then (B.22) follows from (B.23). \square

(E) Line 10. It will be shown that

$$\begin{aligned} e_{3\beta} &\xrightarrow{\varphi_1} e_{(13)\beta}, \\ e_{3\beta} &= y_3 a_k^{31} A_{q+1}^{12} a_k^{12} A_{q+1}^{23} \xrightarrow{\varphi_1} y_1 A_k^{21} a_{q+1}^{13} A_k^{32} a_{q+2}^{32} = e_{(13)\beta}. \end{aligned} \quad (\text{B.24})$$

The following relations are true (Corollary 2.7.4)

$$\begin{aligned} a_k^{31} &\xrightarrow{\varphi_1} y_1 A_k^{21}, \\ A_{q+1}^{12} &\xrightarrow{\varphi_1} y_1 (y_2 + y_3) A_{q+1}^{13}, \\ a_k^{12} &\xrightarrow{\varphi_1} A_k^{32} \subseteq y_2 + y_3, \\ y_3 A_{q+1}^{23} &\xrightarrow{\varphi_1} y_1 a_{q+2}^{32}. \end{aligned} \quad (\text{B.25})$$

Then (B.24) follows from (B.25). \square

(G) Line 10. It will be shown that

$$\begin{aligned} g_{3\beta 0} &\xrightarrow{\varphi_1} g_{(13)\beta 0}, \\ g_{3\beta 0} &= e_{3\beta} (y_1 y_3 + A_q^{23} a_{k+1}^{12}) \xrightarrow{\varphi_1} e_{(13)\beta} (x_1 + a_{q+1}^{32} A_{k+1}^{32}) = g_{(13)\beta 0}. \end{aligned} \quad (\text{B.26})$$

We have

$$e_{3\beta} (y_1 y_3 + A_q^{23} a_{k+1}^{12}) = e_{3\beta} (y_1 y_3 + y_3 A_q^{23} a_{k+1}^{12})$$

and

$$e_{(13)\beta} (x_1 + a_{q+1}^{32} A_{k+1}^{32}) = e_{(13)\beta} (x_1 (x_3 + y_2) + y_1 a_{q+1}^{32} A_{k+1}^{32}),$$

so the following relations (Corollary 2.7.4) are true:

$$\begin{aligned} y_1 y_3 &\xrightarrow{\varphi_1} x_1 (y_2 + y_3) y_1 (x_3 + y_2) = x_1 (x_3 + y_2), \\ y_3 A_q^{23} &\xrightarrow{\varphi_1} y_1 a_{q+1}^{32} \subseteq x_3 + y_2, \\ a_{k+1}^{12} &\xrightarrow{\varphi_1} y_1 A_{k+1}^{32}. \end{aligned} \quad (\text{B.27})$$

Then (B.26) follows from (B.27). \square

Now, we will prove

$$\begin{aligned} \text{Line 1}(p=0) &\xrightarrow{\varphi_1} \text{Line 7} \xrightarrow{\varphi_2} \text{Line 9}(p=0) \xrightarrow{\varphi_1} \text{Line 8} \xrightarrow{\varphi_2} \text{Line 1}(p=0), \\ z_{(21)^{2k}} &\xrightarrow{\varphi_1} z_{1(21)^{2k}} \xrightarrow{\varphi_2} z_{(21)^{2k+1}} \xrightarrow{\varphi_1} z_{1(21)^{2k+1}} \xrightarrow{\varphi_2} z_{(21)^{2k+2}} \text{ for } z_\alpha = f_\alpha, e_\alpha, g_{\alpha 0}. \end{aligned} \quad (\text{B.28})$$

B.1.5. Case $z_{(21)^{2k}} \xrightarrow{\varphi_1} z_{1(21)^{2k}}$.

(F) Line 1 ($p = 0$) $\xrightarrow{\varphi_1}$ Line 7. We will prove that

$$\begin{aligned} f_{(21)^{2k}} &\xrightarrow{\varphi_1} f_{1(21)^{2k}}, \\ f_{(21)^{2k}} &= y_1 y_2 a_k^{13} A_{k-1}^{32} \xrightarrow{\varphi_1} x_1 A_k^{23} a_k^{23} = f_{1(21)^{2k}}. \end{aligned} \quad (\text{B.29})$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} y_1 &\xrightarrow{\varphi_1} x_1(y_2 + y_3), \\ a_k^{13} &\xrightarrow{\varphi_1} y_1 A_k^{23} \subseteq y_2 + y_3, \\ y_2 A_k^{32} &\xrightarrow{\varphi_1} y_1 A_k^{23}. \end{aligned} \quad (\text{B.30})$$

Then (B.29) follows from (B.30). \square

(E) Line 1 ($p = 0$) $\xrightarrow{\varphi_1}$ Line 7. We will prove that

$$\begin{aligned} e_{(21)^{2k}} &\xrightarrow{\varphi_1} e_{1(21)^{2k}}, \\ e_{(21)^{2k}} &= y_2 A_{k-1}^{32} a_k^{21} A_k^{13} a_k^{13} \xrightarrow{\varphi_1} y_1 A_k^{31} a_k^{12} A_k^{23} a_k^{23} = e_{1(21)^{2k}}. \end{aligned} \quad (\text{B.31})$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} y_2 A_{k-1}^{32} &\xrightarrow{\varphi_1} y_1 a_k^{23}, \\ a_k^{21} &\xrightarrow{\varphi_1} y_1 A_k^{31}, \\ A_k^{13} &\xrightarrow{\varphi_1} y_1(y_2 + y_3) a_k^{12}, \\ a_k^{13} &\xrightarrow{\varphi_1} y_1 A_k^{23} \subseteq y_2 + y_3. \end{aligned} \quad (\text{B.32})$$

Then (B.31) follows from (B.32). \square

(G) Line 1 ($p = 0$) $\xrightarrow{\varphi_1}$ Line 7. We need to show

$$\begin{aligned} g_{(21)^{2k}0} &\xrightarrow{\varphi_1} g_{1(21)^{2k}0}, \\ g_{(21)^{2k}0} &= e_{(21)^{2k}}(x_1 + a_k^{32} A_k^{32}) \xrightarrow{\varphi_1} e_{1(21)^{2k}}(y_2 a_k^{21} + y_3 a_k^{31}) = g_{1(21)^{2k}0}. \end{aligned} \quad (\text{B.33})$$

Since $A_k^{32} = y_3 + x_2 A_{k-1}^{13}$ and $x_2 \subseteq a_k^{32}$, we see that $a_k^{32} A_k^{32} = x_2 A_{k-1}^{13} + y_3 a_k^{32}$ and

$$x_1 + a_k^{32} A_k^{32} = (x_1 + y_3 a_k^{32}) + x_2 A_{k-1}^{13} = a_{k+1}^{13} + x_2 A_{k-1}^{13}.$$

The transformation is well-defined because $k > 0$ (see Table 2.3). Further, since $e_{(21)^{2k}} \subseteq A_{k-1}^{13}$ (by Table 2.3), we have

$$e_{(21)^{2k}}(x_1 + a_k^{32} A_k^{32}) = e_{(21)^{2k}}(a_{k+1}^{13} + x_2 A_{k-1}^{13}) = e_{(21)^{2k}}(x_2 + a_{k+1}^{13} A_{k-1}^{13}). \quad (\text{B.34})$$

Apply now φ_1 . The polynomial $x_2 + a_{k+1}^{13} A_{k-1}^{13}$ is φ_1 -homomorphic because $x_1 \subseteq a_{k+1}^{13} A_{k-1}^{13}$. We get

$$\begin{aligned} e_{(21)^{2k}} &\xrightarrow{\varphi_1} e_{1(21)^{2k}}, \\ x_2 &\xrightarrow{\varphi_1} y_1 y_3, \\ a_{k+1}^{13} &\xrightarrow{\varphi_1} y_1 A_{k+1}^{23} \subseteq y_2 + y_3, \\ A_{k-1}^{13} &\xrightarrow{\varphi_1} y_1(y_2 + y_3) a_{k-1}^{12}. \end{aligned} \quad (\text{B.35})$$

Since $e_{1(21)^{2k+1}} \subseteq y_1$, we get

$$e_{(21)^{2k}}(x_1 + a_k^{32} A_k^{32}) \xrightarrow{\varphi_1} e_{1(21)^{2k+1}}(y_1 y_3 + y_1 A_{k+1}^{23} a_{k-1}^{12}) = e_{1(21)^{2k+1}}(y_1 y_3 + A_{k+1}^{23} a_{k-1}^{12}). \quad (\text{B.36})$$

Since $e_{1(21)^{2k+1}} \subseteq a_{k-1}^{12}$, we have

$$\begin{aligned} e_{1(21)^{2k+1}}(y_1 y_3 + A_{k+1}^{23} a_{k-1}^{12}) &= e_{1(21)^{2k+1}}(A_{k+1}^{23} + y_1 y_3 a_{k-1}^{12}) = \\ e_{1(21)^{2k+1}}(y_2 + x_3 A_k^{12} + y_1 y_3 a_{k-1}^{12}) &= e_{1(21)^{2k+1}}(y_2 + y_3 A_k^{12} (x_3 + y_1 a_{k-1}^{12})) = \\ e_{1(21)^{2k+1}}(y_2 + y_3 A_k^{12} a_k^{31}). \end{aligned}$$

Again, by Table 2.3 $e_{1(21)^{2k+1}} \subseteq y_1 \subseteq A_k^{12}$, therefore the last polynomial is equal to

$$e_{1(21)^{2k+1}}(y_2 A_k^{12} + y_3 a_k^{31}),$$

i.e.,

$$e_{1(21)^{2k+1}}(y_1 y_3 + A_{k+1}^{23} a_{k-1}^{12}) = e_{1(21)^{2k+1}}(y_2 A_k^{12} + y_3 a_k^{31}). \quad (\text{B.37})$$

(B.33) follows from (B.37) and from the next lemma.

LEMMA B.1.1. *The following relation takes place*

$$e_{1(21)^{2k+1}}(y_2 A_k^{12} + y_3 a_k^{31}) = e_{1(21)^{2k+1}}(y_2 a_k^{21} + y_3 a_k^{31}). \quad (\text{B.38})$$

Proof. Indeed,

$$\begin{aligned} e_{1(21)^{2k+1}}(y_2 A_k^{12} + y_3 a_k^{31}) &= e_{1(21)^{2k+1}}(y_2 (y_1 + x_2 A_{k-1}^{31}) + y_3 a_k^{31}) = \\ e_{1(21)^{2k+1}}(y_2 y_1 + x_2 A_{k-1}^{31} + y_3 a_k^{31}). \end{aligned}$$

Since $e_{1(21)^{2k+1}} \subseteq A_{k-1}^{31}$, we have

$$\begin{aligned} e_{1(21)^{2k+1}}(y_2 A_k^{12} + y_3 a_k^{31}) &= e_{1(21)^{2k+1}}(x_2 + A_{k-1}^{31} (y_1 y_2 + y_3 a_k^{31})) = \\ e_{1(21)^{2k+1}}(x_2 + y_1 y_2 A_{k-1}^{31} + y_3 a_k^{31}). \end{aligned}$$

Further, by basic relations (Table 2.1, 4.1) we have

$$\begin{aligned} e_{1(21)^{2k+1}}(y_2 A_k^{12} + y_3 a_k^{31}) &= e_{1(21)^{2k+1}}(x_2 + y_1 y_2 a_{k-1}^{13} + y_3 a_k^{31}) = \\ e_{1(21)^{2k+1}}(y_2 (x_2 + y_2 a_{k-1}^{13}) + y_3 a_k^{31}) &= e_{1(21)^{2k+1}}(y_2 a_k^{21} + y_3 a_k^{31}). \end{aligned}$$

Thus, the lemma is proven, (B.38) and so (B.33) are true. \square

Action φ_2 transforms Line 7 to Line 9 with $p = 0$. For $z_\alpha = f_\alpha, e_\alpha, g_{\alpha 0}$, we are going to prove

B.1.6. Case $z_{1(21)^{2k}} \xrightarrow{\varphi_2} z_{(21)^{2k+1}}$.

(F) Line 7 $\xrightarrow{\varphi_2}$ Line 9 ($p = 0$). We want to prove that

$$\begin{aligned} f_{1(21)^{2k}} &\xrightarrow{\varphi_2} f_{(21)^{2k+1}}, \\ f_{1(21)^{2k}} &= x_1 A_k^{23} a_k^{23} \xrightarrow{\varphi_2} y_2 y_3 a_k^{21} A_k^{13} = f_{(21)^{2k+1}}. \end{aligned} \quad (\text{B.39})$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} x_1 &\xrightarrow{\varphi_2} y_2 y_3, \\ A_k^{23} &\xrightarrow{\varphi_2} y_2 (y_1 + y_3) a_k^{21}, \\ a_k^{23} &\xrightarrow{\varphi_2} y_2 A_k^{13} \subseteq y_1 + y_3. \end{aligned} \quad (\text{B.40})$$

Then (B.39) follows from (B.40). \square

(E) Line 7 $\xrightarrow{\varphi_2}$ Line 9 ($p = 0$). We want to prove that

$$\begin{aligned} e_{1(21)^{2k}} &\xrightarrow{\varphi_2} e_{(21)^{2k+1}}, \\ e_{1(21)^{2k}} &= y_1 A_k^{31} a_k^{12} A_k^{23} a_k^{23} \xrightarrow{\varphi_2} y_2 A_k^{32} a_k^{21} A_k^{13} a_{k+1}^{13} = e_{(21)^{2k+1}}. \end{aligned} \quad (\text{B.41})$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} y_1 A_k^{31} &\xrightarrow{\varphi_2} y_1 a_{k+1}^{13}, \\ a_k^{12} &\xrightarrow{\varphi_2} y_2 A_k^{32}, \\ A_k^{23} &\xrightarrow{\varphi_2} y_2 (y_1 + y_3) a_k^{21}, \\ a_k^{23} &\xrightarrow{\varphi_2} y_2 A_k^{13} \subseteq y_1 + y_3. \end{aligned} \quad (\text{B.42})$$

Thus, (B.41) follows from (B.42). \square

(G) Line 7 $\xrightarrow{\varphi_2}$ Line 9 ($p = 0$). It will be shown that

$$\begin{aligned} g_{1(21)^{2k}0} &\xrightarrow{\varphi_2} g_{(21)^{2k+1}0}, \\ g_{1(21)^{2k}0} &= e_{1(21)^{2k+1}} (y_2 a_k^{21} + y_3 a_k^{31}) \xrightarrow{\varphi_2} e_{(21)^{2k+1}} (x_2 + a_k^{13} A_k^{13}) = g_{(21)^{2k+1}0}. \end{aligned} \quad (\text{B.43})$$

The following relations are true:

$$\begin{aligned} y_2 a^{21} &\xrightarrow{\varphi_2} x_2 (y_1 + y_3) a^{13} y_2 k A_k^{31} = x_2 A_k^{31}, \\ y_3 a^{31} &\xrightarrow{\varphi_2} y_2 A_k^{13}. \end{aligned} \quad (\text{B.44})$$

Therefore,

$$y_2 a_k^{21} + y_3 a_k^{31} \xrightarrow{\varphi_2} y_2 (x_2 A_k^{31} + A_k^{13}) = y_2 (x_2 a_k^{13} + A_k^{13}) = x_2 a_k^{13} + y_2 A_k^{13}. \quad (\text{B.45})$$

Since

$$e_{1(21)^{2k+1}} \xrightarrow{\varphi_2} e_{(21)^{2k+1}} \subseteq a_{k+1}^{13} \subseteq a_k^{13}, \quad (\text{B.46})$$

we see by (B.45) that

$$e_{1(21)^{2k+1}} (y_2 a_k^{21} + y_3 a_k^{31}) \xrightarrow{\varphi_2} e_{(21)^{2k+1}} (x_2 a_k^{13} + y_2 A_k^{13}) = e_{(21)^{2k+1}} (x_2 + y_2 a_k^{13} A_k^{13}). \quad \square \quad (\text{B.47})$$

Action φ_1 transforms Line 9 with $p = 0$ to Line 8 with $p = 0$. For $z_\alpha = f_\alpha, e_\alpha, g_{\alpha 0}$, we are going to prove

B.1.7. Case $z_{(21)^{2k+1}} \xrightarrow{\varphi_1} z_{1(21)^{2k+1}}$.

(F) Line 9 ($p = 0$) $\xrightarrow{\varphi_1}$ Line 8. It will be shown that

$$\begin{aligned} f_{(21)^{2k+1}} &\xrightarrow{\varphi_1} f_{1(21)^{2k+1}}, \\ f_{(21)^{2k+1}} &= y_2 y_3 a_k^{21} A_k^{13} \xrightarrow{\varphi_1} y_1 (x_2 + x_3) A_k^{31} a_k^{12} = f_{1(21)^{2k+1}}. \end{aligned} \quad (\text{B.48})$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} y_2 y_3 &\xrightarrow{\varphi_1} y_1 (x_2 + x_3) \subseteq y_2 + y_3, \\ a_k^{21} &\xrightarrow{\varphi_1} y_1 A_k^{31}, \\ A_k^{13} &\xrightarrow{\varphi_1} y_1 (y_2 + y_3) a_k^{12}. \end{aligned} \quad (\text{B.49})$$

Then (B.48) follows from (B.49). \square

(E) Line 9 ($p = 0$) $\xrightarrow{\varphi_1}$ Line 8. Here, it should be proven that

$$\begin{aligned} e_{(21)^{2k+1}} &\xrightarrow{\varphi_1} e_{1(21)^{2k+1}}, \\ e_{(21)^{2k+1}} &= y_2 A_k^{32} a_k^{21} A_k^{13} a_{k+1}^{13} \xrightarrow{\varphi_1} y_1 A_k^{31} a_k^{12} A_{k+1}^{23} a_{k+1}^{23} = e_{1(21)^{2k+1}}. \end{aligned} \quad (\text{B.50})$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} y_2 A_k^{32} &\xrightarrow{\varphi_1} y_1 a_{k+1}^{23} \subseteq y_2 + y_3, \\ a_k^{21} &\xrightarrow{\varphi_1} y_1 A_k^{31}, \\ A_k^{13} &\xrightarrow{\varphi_1} y_1 (y_2 + y_3) a_k^{12}, \\ a_{k+1}^{13} &\xrightarrow{\varphi_1} y_1 A_{k+1}^{23}. \end{aligned} \quad (\text{B.51})$$

Therefore, (B.50) follows from (B.51). \square

(G) Line 9 ($p = 0$) $\xrightarrow{\varphi_1}$ Line 8. We will show that

$$\begin{aligned} g_{(21)^{2k+1}0} &\xrightarrow{\varphi_1} g_{1(21)^{2k+1}0}, \\ g_{(21)^{2k+1}0} &= e_{(21)^{2k+1}}(x_2 + a_k^{13} A_{k+1}^{13}) \xrightarrow{\varphi_1} e_{1(21)^{2k+1}}(y_1 a_{k+1}^{12} + y_1 a_{k+1}^{13}) = g_{1(21)^{2k+1}0}. \end{aligned} \quad (\text{B.52})$$

First, from the relations

$$\begin{aligned} e_{(21)^{2k+1}} &\xrightarrow{\varphi_1} e_{1(21)^{2k+1}} \subseteq y_1, \\ x_2 &\xrightarrow{\varphi_1} y_1 y_3, \\ a_k^{13} &\xrightarrow{\varphi_1} y_1 A_k^{23} \subseteq y_2 + y_3, \\ A_{k+1}^{13} &\xrightarrow{\varphi_1} y_1 (y_2 + y_3) a_{k+1}^{12} \end{aligned} \quad (\text{B.53})$$

we have

$$\begin{aligned} e_{(21)^{2k+1}}(x_2 + a_k^{13} A_{k+1}^{13}) &\xrightarrow{\varphi_1} e_{1(21)^{2k+1}}(y_1 y_3 + y_1 A_k^{23} a_{k+1}^{12}) = \\ &= e_{1(21)^{2k+1}}(y_1 y_3 + A_{k+1}^{23} a_{k+1}^{12}). \end{aligned} \quad (\text{B.54})$$

Transform the last polynomial from (B.54):

$$\begin{aligned} y_1 y_3 + A_k^{23} a_{k+1}^{12} &= y_1 y_3 + A_k^{23} (x_1 + y_2 a_k^{23}) = \\ y_1 y_3 + x_1 A_k^{23} + y_2 a_k^{23} &= y_1 (y_3 + x_1 A_k^{23}) + y_2 a_k^{23} = y_1 A_{k+1}^{31} + y_2 a_k^{23}. \end{aligned}$$

Thus,

$$e_{(21)^{2k+1}}(x_2 + a_k^{13} A_{k+1}^{13}) \xrightarrow{\varphi_1} e_{1(21)^{2k+1}}(y_1 A_{k+1}^{31} + y_2 a_k^{23}). \quad (\text{B.55})$$

Then (B.52) follows from the next lemma.

LEMMA B.1.2. *We have*

$$e_{1(21)^{2k+1}}(y_1 A_{k+1}^{31} + y_2 a_k^{23}) = e_{1(21)^{2k+1}}(y_1 a_{k+1}^{12} + y_1 a_{k+1}^{13}). \quad (\text{B.56})$$

Proof. Indeed,

$$\begin{aligned} e_{1(21)^{2k+1}}(y_1 A_{k+1}^{31} + y_2 a_k^{23}) &= e_{1(21)^{2k+1}}(y_1 (y_3 + x_1 A_k^{23}) + y_2 a_k^{23}) = \\ e_{1(21)^{2k+1}}(y_1 y_3 + x_1 A_k^{23} + y_2 a_k^{23}) &= e_{1(21)^{2k+1}}(x_1 + y_1 y_3 A_k^{23} + y_2 a_k^{23}). \end{aligned}$$

Since $e_{1(21)^{2k+1}} \subseteq A_{k+1}^{23} \subseteq A_k^{23}$, it follows that

$$\begin{aligned} e_{1(21)^{2k+1}}(y_1 A_{k+1}^{31} + y_2 a_k^{23}) &= e_{1(21)^{2k+1}}(x_1 + A_k^{23} (y_1 y_3 + y_2 a_k^{23})) = \\ e_{1(21)^{2k+1}}(x_1 + y_1 y_3 A_k^{23} + y_2 a_k^{23}) &= e_{1(21)^{2k+1}}(x_1 + y_1 y_3 A_{k+1}^{23} + y_2 a_{k+1}^{23}). \end{aligned}$$

By basic relations of atomic elements (Table 2.1, 4.1) we have

$$\begin{aligned} e_{1(21)^{2k+1}}(x_1 + A_k^{23}(y_1 y_3 + y_2 a_k^{23})) &= e_{1(21)^{2k+1}}(x_1 + y_1 y_3 a_k^{32} + y_2 a_k^{23}) = \\ e_{1(21)^{2k+1}}(y_1(x_1 + y_3 a_k^{32}) + y_2 a_k^{23}) &= e_{1(21)^{2k+1}}(y_1 a_{k+1}^{13} + y_2 a_k^{23}). \end{aligned}$$

Now,

$$\begin{aligned} e_{1(21)^{2k+1}}(y_1 a_{k+1}^{13} + y_2 a_k^{23}) &= e_{1(21)^{2k+1}}(y_1 a_{k+1}^{13} + x_1 + y_2 a_k^{23}) = \\ e_{1(21)^{2k+1}}(y_1 a_{k+1}^{13} + a_{k+1}^{12}) &= e_{1(21)^{2k+1}}(y_1 a_{k+1}^{13} + y_1 a_{k+1}^{12}). \end{aligned}$$

The lemma is proved. \square

Action φ_2 transforms Line 8 to the Line 1 with $p = 0$. For $z_\alpha = f_\alpha, e_\alpha, g_{\alpha 0}$ we are going to prove

B.1.8. Case $z_{1(21)^{2k+1}} \xrightarrow{\varphi_2} z_{(21)^{2k+2}}.$

(F) Line 8 $\xrightarrow{\varphi_2}$ Line 1 ($p = 0$). We want to prove that

$$\begin{aligned} f_{1(21)^{2k+1}} &\xrightarrow{\varphi_2} f_{(21)^{2k+2}}, \\ f_{1(21)^{2k+1}} &= y_1(x_2 + x_3)A_k^{31}a_k^{12} \xrightarrow{\varphi_1} y_1 y_2 a_{k+1}^{13} A_k^{32} = f_{(21)^{2k+2}}. \end{aligned} \tag{B.57}$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} x_2 + x_3 &\xrightarrow{\varphi_2} y_1 y_2, \\ y_1 A_k^{31} &\xrightarrow{\varphi_2} y_2 a_{k+1}^{13}, \\ a_k^{12} &\xrightarrow{\varphi_2} y_1 A_k^{32}. \end{aligned} \tag{B.58}$$

Then (B.57) follows from (B.58). \square

(E) Line 8 $\xrightarrow{\varphi_2}$ Line 1 ($p = 0$). We need to show

$$\begin{aligned} e_{1(21)^{2k+1}} &\xrightarrow{\varphi_2} e_{(21)^{2k+2}}, \\ e_{1(21)^{2k+1}} &= y_1 A_k^{31} a_k^{12} A_{k+1}^{23} a_{k+1}^{23} \xrightarrow{\varphi_2} y_2 A_k^{32} a_{k+1}^{21} A_{k+1}^{13} a_{k+1}^{13} = e_{(21)^{2k+2}}. \end{aligned} \tag{B.59}$$

The following relations are true (Corollary 2.7.4):

$$\begin{aligned} y_1 A_k^{31} &\xrightarrow{\varphi_2} y_2 a_{k+1}^{13} \subseteq y_1 + y_3, \\ a_k^{12} &\xrightarrow{\varphi_2} y_2 A_k^{32}, \\ A_{k+1}^{23} &\xrightarrow{\varphi_2} y_2(y_1 + y_3)a_{k+1}^{21}, \\ a_{k+1}^{23} &\xrightarrow{\varphi_2} y_1 A_{k+1}^{13}. \end{aligned} \tag{B.60}$$

Thus, (B.59) follows from (B.60). \square

(G) Line 8 $\xrightarrow{\varphi_2}$ Line 1 ($p = 0$). It will be shown that

$$\begin{aligned} g_{1(21)^{2k+1}0} &\xrightarrow{\varphi_2} g_{(21)^{2k+2}0}, \\ g_{1(21)^{2k+1}0} &= e_{1(21)^{2k+1}}(y_1 a_{k+1}^{12} + y_1 a_{k+1}^{13}) \xrightarrow{\varphi_2} e_{(21)^{2k+2}}(x_1 + a_{k+1}^{32} A_{k+2}^{32}) = g_{(21)^{2k+2}0}. \end{aligned} \tag{B.61}$$

First, since $e_{1(21)^{2k+1}} \subseteq y_1$, we have

$$e_{1(21)^{2k+1}}(y_1 a_{k+1}^{12} + y_1 a_{k+1}^{13}) = e_{1(21)^{2k+1}}(x_1 + y_2 a_k^{23} + y_1 y_3 a_k^{32}).$$

Since $x_2 \subseteq y_2 a_k^{23}$ and the polynomial $x_1 + y_2 a_k^{23} + y_1 y_3 a_k^{32}$ meets the requirements of atomic multiplicativity (Corollary 2.5.2), we have

$$e_{1(21)^{2k+1}}(x_1 + y_2 a_k^{23} + y_1 y_3 a_k^{32}) \xrightarrow{\varphi_2} e_{(21)^{2k+2}}(y_2 y_3 + x_2(y_1 + y_3)A_k^{13} + y_2(x_1 + x_3)A_k^{12}).$$

We transform the last polynomial. Since $e_{(21)^{2k+2}} \subseteq y_2$, it follows that

$$\begin{aligned} e_{(21)^{2k+2}}(y_2 y_3 + x_2(y_1 + y_3)A_k^{13} + y_2(x_1 + x_3)A_k^{12}) &= \\ e_{(21)^{2k+2}}(y_2 y_3 + x_2 A_k^{13} + (x_1 + x_3)A_k^{12}) &= \\ e_{(21)^{2k+2}}(y_2 y_3 + x_2 A_k^{13} + x_1 + x_3 A_k^{12}) &= e_{(21)^{2k+2}}(y_2 y_3 + x_2 A_k^{13} + x_1 + x_3 a_k^{12}). \end{aligned}$$

Since $e_{(21)^{2k+2}} \subseteq a_k^{12}$, we see by permutation property (A.4) that

$$\begin{aligned} e_{(21)^{2k+2}}(y_2 y_3 + x_2 A_k^{13} + x_1 + x_3 a_k^{12}) &= e_{(21)^{2k+2}}(x_1 + x_3 + a_k^{12}(y_2 y_3 + x_2 A_k^{13})) = \\ e_{(21)^{2k+2}}(x_1 + x_3 + x_2 A_k^{13} + y_2 y_3 a_k^{12}) &= e_{(21)^{2k+2}}(x_1 + y_3(x_3 + y_2 a_k^{12}) + x_2 A_k^{13}) = \\ e_{(21)^{2k+2}}(x_1 + y_3 a_{k+1}^{32} + x_2 A_k^{13}) &= e_{(21)^{2k+2}}(x_1 + a_{k+1}^{32}(y_3 + x_2 A_k^{13}) = e_{(21)^{2k+2}}(x_1 + a_{k+1}^{32} A_{k+1}^{32}), \end{aligned}$$

which was to be proved. \square

The proof of Theorem 2.12.1 is finished. \square

B.2. A proof of the theorem on admissible elements in lattice D^4

Every element e_α from Table 4.3 is the intersection of three polynomials. In all cases, two of these polynomials are φ_i -homomorphic under action ψ_i . The third one is not φ_i -homomorphic, but satisfies the following relation:

$$e_j e_n^{kl} \xrightarrow{\varphi_i} e_i e_{n+1}^{kl},$$

see Corollary 4.5.2, heading(3).

In this proof, we repeatedly use the basic properties of admissible elements from Lemma 4.8.2, §4.8.1.

B.2.1. Line $F21$, $e_\alpha \xrightarrow{\varphi_3} e_{3\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_3} f_{3\alpha 0}$. Here,

$$\alpha = (21)^t (41)^r (31)^s, e_\alpha = e_{(21)^t (41)^r (31)^s},$$

and

$$e_\alpha = e_2 a_{2s}^{31} a_{2r}^{41} a_{2t-1}^{34} \xrightarrow{\varphi_3} e_3 a_{2s}^{24} a_{2t-1}^{21} a_{2r+1}^{41} = e_{3\alpha}, \quad (\text{B.62})$$

where 3α is obtained by eq.(4.4). By Corollary 4.5.2 we have

$$\begin{aligned} a_{2s}^{31} &\xrightarrow{\varphi_3} e_3 a_{2s}^{24}, \\ a_{2t-1}^{34} &\xrightarrow{\varphi_3} e_3 a_{2t-1}^{21}, \\ e_2 a_{2r}^{41} &\xrightarrow{\varphi_3} e_3 a_{2r+1}^{41}. \end{aligned}$$

Since a_{2s}^{31} and a_{2t-1}^{34} are φ_3 -homomorphic (Corollary 4.4.2), we get (B.62). The polynomial $e_{3\alpha}$ corresponds to Line $G31$ under the substitution

$$\begin{aligned} r &\mapsto r + 1, \\ t &\mapsto t - 1, \end{aligned} \quad (\text{B.63})$$

see relation (4.4). For the case $f_{\alpha 0}$, we have

$$e_\alpha (e_2 a_{2t}^{34} + a_{2r+1}^{14} a_{2s-1}^{13}) = e_\alpha (a_{2t}^{34} + e_2 a_{2r+1}^{14} a_{2s-1}^{13}),$$

and we select

$$P = a_{2t}^{34} + e_2 a_{2r+1}^{14} a_{2s-1}^{13},$$

see Lemma 4.8.2. By Corollary 4.5.2 we have

$$\begin{aligned} a_{2t}^{34} &\xrightarrow{\varphi_3} e_3 a_{2t}^{21}, \\ a_{2s-1}^{13} &\xrightarrow{\varphi_3} e_3 a_{2s-1}^{42}, \\ e_2 a_{2r+1}^{14} &\xrightarrow{\varphi_3} e_3 a_{2r+2}^{14}, \end{aligned}$$

and we get

$$P = a_{2t}^{34} + e_2 a_{2r+1}^{14} a_{2s-1}^{13} \xrightarrow{\varphi_3} e_3 a_{2t}^{21} + e_3 a_{2r+2}^{14} a_{2s-1}^{42} = \tilde{P}. \quad (\text{B.64})$$

Since $a_{2s}^{42} \subseteq a_{2s-1}^{42}$, we have

$$\begin{aligned} e_{3\alpha} \tilde{P} &= e_3 a_{2s}^{24} a_{2t-1}^{21} a_{2r+1}^{41} (e_3 a_{2t}^{21} + e_3 a_{2r+2}^{14} a_{2s-1}^{42}) = \\ &= e_3 a_{2s}^{24} a_{2t-1}^{21} a_{2r+1}^{41} (a_{2r+2}^{14} + e_3 a_{2t}^{21} a_{2s-1}^{42}), \end{aligned}$$

which corresponds to Line $G31$ with substitution (B.63), see relation (4.4). \square

B.2.2. Line $F21$, $e_\alpha \xrightarrow{\varphi_4} e_{4\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_4} f_{4\alpha 0}$. We have $\alpha = (21)^t(41)^r(31)^s$, $e_\alpha = e_{(21)^t(41)^r(31)^s}$,

$$e_\alpha = e_2 a_{2s}^{31} a_{2r}^{41} a_{2t-1}^{34} \xrightarrow{\varphi_4} e_4 a_{2s+1}^{31} a_{2r}^{32} a_{2t-1}^{12} = e_{4\alpha}, \quad (\text{B.65})$$

because by Corollary 4.5.2 we have

$$\begin{aligned} a_{2r}^{41} &\xrightarrow{\varphi_4} e_4 a_{2r}^{32}, \\ a_{2t-1}^{43} &\xrightarrow{\varphi_4} e_4 a_{2t-1}^{12}, \\ e_4 a_{2s}^{31} &\xrightarrow{\varphi_4} e_4 a_{2s+1}^{31}. \end{aligned}$$

The polynomial $e_{4\alpha}$ corresponds to Line $G41$. For the case $f_{\alpha 0}$, we select

$$P = a_{2t}^{43} + e_2 a_{2r+1}^{14} a_{2s-1}^{13}.$$

We have

$$P = a_{2t}^{43} + e_2 a_{2r+1}^{14} a_{2s-1}^{13} \xrightarrow{\varphi_4} e_4 a_{2t}^{21} + e_4 a_{2r+1}^{23} a_{2s}^{13} = \tilde{P}, \quad (\text{B.66})$$

because by Corollary 4.5.2 we have

$$\begin{aligned} a_{2r+1}^{41} &\xrightarrow{\varphi_4} e_4 a_{2r+1}^{32}, \\ a_{2t}^{43} &\xrightarrow{\varphi_4} e_4 a_{2t}^{12}, \\ e_4 a_{2s-1}^{31} &\xrightarrow{\varphi_4} e_4 a_{2s}^{31}. \end{aligned}$$

The polynomial \tilde{P} corresponds to Line $G41$. \square

B.2.3. Line $F21$, $e_\alpha \xrightarrow{\varphi_1} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_1} f_{1\alpha 0}$. The admissible sequence α and polynomial e_α are given as in §B.2.2. Here,

$$e_\alpha = e_2 a_{2s}^{31} a_{2r}^{41} a_{2t-1}^{34} \xrightarrow{\varphi_1} e_1 a_{2s}^{42} a_{2r}^{23} a_{2t}^{34} = e_{1\alpha}, \quad (\text{B.67})$$

because by Corollary 4.5.2 we have

$$\begin{aligned} a_{2s}^{31} &\xrightarrow{\varphi_1} e_1 a_{2s}^{42}, \\ a_{2r}^{41} &\xrightarrow{\varphi_1} e_1 a_{2r}^{23}, \\ e_2 a_{2t-1}^{34} &\xrightarrow{\varphi_1} e_1 a_{2t}^{34}. \end{aligned}$$

The polynomial $e_{1\alpha}$ corresponds to Line $G11$. For the case $f_{\alpha 0}$, we select

$$P = e_2 a_{2t}^{43} + a_{2r+1}^{14} a_{2s-1}^{13},$$

and we have

$$P = e_2 a_{2t}^{43} + a_{2r+1}^{14} a_{2s-1}^{13} \xrightarrow{\varphi_1} e_1 a_{2t+1}^{43} + e_1 a_{2r+1}^{23} a_{2s-1}^{24} = \tilde{P}, \quad (\text{B.68})$$

because by Corollary 4.5.2 we have

$$\begin{aligned} a_{2r+1}^{41} &\xrightarrow{\varphi_1} e_1 a_{2r+1}^{32}, \\ a_{2s-1}^{31} &\xrightarrow{\varphi_1} e_1 a_{2s-1}^{24}, \\ e_2 a_{2t}^{43} &\xrightarrow{\varphi_1} e_1 a_{2t+1}^{43}, \end{aligned}$$

Since $a_{2s}^{24} \subseteq a_{2s-1}^{24}$ (see Line G11), by (4.45) we have

$$\begin{aligned} e_{1\alpha} \tilde{P} &= e_1 a_{2s}^{42} a_{2r}^{23} a_{2t}^{34} (e_1 a_{2t+1}^{43} + a_{2r+1}^{23} a_{2s-1}^{24}) = \\ &= e_1 a_{2s}^{42} a_{2r}^{23} a_{2t}^{34} (e_1 a_{2t+1}^{43} a_{2s-1}^{24} + a_{2r+1}^{23}) = e_{1\alpha} (a_{2t-1}^{43} a_{2s+1}^{24} + e_1 a_{2r+1}^{23}), \end{aligned}$$

which corresponds to Line G11. \square

B.2.4. Line F31, $e_\alpha \xrightarrow{\varphi_1} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_1} f_{1\alpha 0}$. First, we have

$$e_\alpha = e_3 a_{2t}^{21} a_{2r}^{41} a_{2s-1}^{24} \xrightarrow{\varphi_1} e_1 a_{2t}^{43} a_{2r}^{23} a_{2s}^{24} = e_{1\alpha}. \quad (\text{B.69})$$

For the case $f_{\alpha 0}$, select

$$P = e_3 a_{2s}^{42} + a_{2r+1}^{14} a_{2t-1}^{12},$$

see Lemma 4.8.2. We have

$$P = e_3 a_{2s}^{42} + a_{2r+1}^{14} a_{2t-1}^{12} \xrightarrow{\varphi_1} e_1 a_{2s+1}^{42} + e_1 a_{2r+1}^{23} a_{2t-1}^{34}. \quad (\text{B.70})$$

Since $a_{2t}^{34} \subseteq a_{2t-1}^{34}$, we get

$$\begin{aligned} e_{1\alpha} (e_1 a_{2s+1}^{42} + e_1 a_{2r+1}^{23} a_{2t-1}^{34}) &= e_1 a_{2t}^{43} a_{2r}^{23} a_{2s}^{24} (e_1 a_{2s+1}^{42} + e_1 a_{2r+1}^{23} a_{2t-1}^{34}) = \\ &= e_1 a_{2t}^{43} a_{2r}^{23} a_{2s}^{24} (e_1 a_{2s+1}^{42} a_{2t-1}^{34} + e_1 a_{2r+1}^{23}) = e_{1\alpha} (e_1 a_{2r+1}^{23} + a_{2s+1}^{42} a_{2t-1}^{34}) = \tilde{P}, \end{aligned}$$

we get Line G11. \square

Cases F31 (actions ψ_2, ψ_4), and F41 (actions ψ_1, ψ_2, ψ_3) are similarly proved. \square

B.2.5. Line G11, $e_\alpha \xrightarrow{\varphi_2} e_{2\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_2} f_{2\alpha 0}$. We have

$$\alpha = 1(21)^t(31)^r(41)^s, \quad e_\alpha = e_{1(21)^t(31)^r(41)^s}.$$

Here $2\alpha = (21)^{t+1}(31)^r(41)^s$, and by Corollary 4.5.2 we get

$$\begin{aligned} a_{2s}^{24} &\xrightarrow{\varphi_2} e_2 a_{2s}^{13}, \\ a_{2r}^{32} &\xrightarrow{\varphi_2} e_2 a_{2r}^{14}, \\ e_1 a_{2t}^{34} &\xrightarrow{\varphi_2} e_2 a_{2t+1}^{34}, \end{aligned}$$

hence,

$$e_\alpha = e_1 a_{2s}^{24} a_{2t}^{34} a_{2r}^{32} \xrightarrow{\varphi_2} e_2 a_{2s}^{13} a_{2t+1}^{34} a_{2r}^{14} = e_{2\alpha}. \quad (\text{B.71})$$

The polynomial $e_{2\alpha}$ corresponds to Line G21 under the substitution $t \mapsto t+1$. For the case $f_{\alpha 0}$,

$$\begin{aligned} e_\alpha (e_1 a_{2r+1}^{32} + a_{2s+1}^{24} a_{2t-1}^{34}) &= \\ e_\alpha (e_1 a_{2r+1}^{23} + a_{2s+1}^{24} a_{2t-1}^{34}) &= e_\alpha (a_{2r+1}^{23} + e_1 a_{2s+1}^{24} a_{2t-1}^{34}), \end{aligned}$$

and we select

$$P = a_{2r+1}^{23} + e_1 a_{2s+1}^{24} a_{2t-1}^{34}.$$

We have

$$P = e_1 a_{2r+1}^{23} + e_1 a_{2s+1}^{24} a_{2t-1}^{34} \xrightarrow{\varphi_2} e_2 a_{2r+1}^{14} + e_2 a_{2s+1}^{13} a_{2t}^{34} = \tilde{P}. \quad (\text{B.72})$$

Since

$$e_{2\alpha} \subseteq e_2 a_{2s}^{13} \subseteq e_2 a_{2s-1}^{13},$$

by the equalizing property (4.47) we have

$$\begin{aligned} e_{2\alpha} \tilde{P} &= e_{2\alpha} (e_2 a_{2r+1}^{14} + e_2 a_{2s+1}^{13} a_{2t}^{34}) = \\ &= e_{2\alpha} (e_2 a_{2r+1}^{14} + e_2 a_{2s-1}^{13} a_{2t+2}^{34}) = e_{2\alpha} (a_{2t+2}^{34} + e_2 a_{2s-1}^{13} a_{2r+1}^{14}), \end{aligned}$$

i.e., we get Line $F21$ under substitution the $t \mapsto t + 1$. \square

Cases $G11$ (actions ψ_3, ψ_4) are similarly proved. \square

B.2.6. Line $G21$, $e_\alpha \xrightarrow{\varphi_3} e_{2\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_3} f_{3\alpha 0}$. We have

$$\alpha = 2(41)^r (31)^s (21)^t, \quad e_\alpha = e_{2(41)^r (31)^s (21)^t}.$$

By eq.(4.6) $3\alpha = (31)^{s+1} (41)^r (21)^t$, and by Corollary 4.5.2 we have

$$\begin{aligned} a_{2t}^{34} &\xrightarrow{\varphi_3} e_3 a_{2t}^{12}, \\ a_{2s+1}^{31} &\xrightarrow{\varphi_3} e_3 a_{2s+1}^{24}, \\ e_2 a_{2r-1}^{14} &\xrightarrow{\varphi_3} e_3 a_{2r}^{14}, \end{aligned}$$

i.e.,

$$e_\alpha = e_2 a_{2t}^{34} a_{2s+1}^{31} a_{2r-1}^{14} \xrightarrow{\varphi_3} e_3 a_{2t}^{12} a_{2s+1}^{24} a_{2r}^{14} = e_{3\alpha}. \quad (\text{B.73})$$

The polynomial $e_{3\alpha}$ corresponds to Line $F31$ for substitution $s \mapsto s + 1$. For the case $f_{\alpha 0}$, we select

$$P = e_2 a_{2r}^{14} + a_{2s+2}^{31} a_{2t-1}^{34},$$

and we have

$$P = e_2 a_{2r}^{14} + a_{2s+2}^{31} a_{2t-1}^{34} \xrightarrow{\varphi_3} e_3 a_{2r+1}^{14} + a_{2s+2}^{24} a_{2t-1}^{12} = \tilde{P}. \quad (\text{B.74})$$

Since $e_{3\alpha} \subseteq a_{2t}^{12} \subseteq a_{2t-1}^{12}$, we get

$$\begin{aligned} e_{3\alpha} \tilde{P} &= e_{3\alpha} (e_3 a_{2r+1}^{14} + a_{2s+2}^{24} a_{2t-1}^{12}) = \\ &= e_{3\alpha} (a_{2r+1}^{14} a_{2t-1}^{12} + e_3 a_{2s+2}^{24}) \end{aligned} \quad (\text{B.75})$$

which corresponds to Line $F31$. By (B.74) and (B.75) $f_{3\alpha}$ and $e_{3\alpha}$ correspond to Line $F31$ under substitution the $s \mapsto s + 1$. \square

B.2.7. Line $G21$, $e_\alpha \xrightarrow{\varphi_4} e_{4\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_4} f_{4\alpha 0}$. The admissible sequence α and the polynomial e_α are given as above in §B.2.6.

By eq.(4.7) $4\alpha = (41)^r (31)^{s+1} (21)^t$, and by Corollary 4.5.2 we have

$$\begin{aligned} a_{2t}^{34} &\xrightarrow{\varphi_4} e_4 a_{2t}^{12}, \\ a_{2r-1}^{14} &\xrightarrow{\varphi_4} e_4 a_{2r-1}^{23}, \\ e_2 a_{2s+1}^{31} &\xrightarrow{\varphi_4} e_4 a_{2s+2}^{31}, \end{aligned}$$

and therefore

$$e_\alpha = e_2 a_{2t}^{34} a_{2s+1}^{31} a_{2r-1}^{14} \xrightarrow{\varphi_4} e_4 a_{2t}^{12} a_{2r-1}^{23} a_{2s+2}^{31} = e_{4\alpha}. \quad (\text{B.76})$$

The polynomial $e_{4\alpha}$ corresponds to Line $F41$ under the substitution $s \mapsto s+1$. For the case $f_{\alpha 0}$, we have

$$\begin{aligned} e_{\alpha}(a_{2r}^{14} + e_2 a_{2s+2}^{31} a_{2t-1}^{34}) &= e_{\alpha}(e_2 a_{2r}^{14} + e_2 a_{2s+2}^{31} a_{2t-1}^{34}) = \\ e_{\alpha}(e_2 a_{2r}^{41} + e_2 a_{2s+2}^{31} a_{2t-1}^{34}) &= e_{\alpha}(a_{2r}^{41} + e_2 a_{2s+2}^{31} a_{2t-1}^{34}), \end{aligned}$$

and we select

$$P = a_{2r}^{41} + e_2 a_{2s+2}^{31} a_{2t-1}^{34}.$$

Here, we get

$$P = a_{2r}^{41} + e_2 a_{2s+2}^{31} a_{2t-1}^{34} \xrightarrow{\varphi_4} e_4 a_{2r}^{23} + e_4 a_{2s+3}^{31} a_{2t-1}^{12} = \tilde{P}. \quad (\text{B.77})$$

Since

$$e_{4\alpha} \tilde{P} = e_{4\alpha}(e_4 a_{2r}^{32} + a_{2s+3}^{31} a_{2t-1}^{12}),$$

we see that $e_{4\alpha} \tilde{P}$ corresponds to Line $F41$ under the substitution $s \mapsto s+1$. \square

B.2.8. Line $G21$, $e_{\alpha} \xrightarrow{\varphi_1} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_1} f_{1\alpha 0}$. The admissible sequence α and the polynomial e_{α} are given as above in §B.2.6.

By eq.(4.8) $1\alpha = (14)^r(31)^{s+1}(21)^t$, and by Corollary 4.5.2 we have

$$\begin{aligned} a_{2s+1}^{31} &\xrightarrow{\varphi_1} e_1 a_{2s+1}^{24}, \\ a_{2r-1}^{14} &\xrightarrow{\varphi_1} e_1 a_{2r-1}^{23}, \\ e_2 a_{2t}^{34} &\xrightarrow{\varphi_1} e_1 a_{2t+1}^{34}, \end{aligned}$$

and therefore

$$e_{\alpha} = e_2 a_{2s+1}^{31} a_{2r-1}^{14} a_{2t}^{34} \xrightarrow{\varphi_1} e_1 a_{2s+1}^{24} a_{2r-1}^{23} a_{2t+1}^{34} = e_{1\alpha}. \quad (\text{B.78})$$

The polynomial $e_{1\alpha}$ corresponds to Line $H11$ under the substitution $t \mapsto t+1$. For the case $f_{\alpha 0}$, we have

$$\begin{aligned} e_{\alpha} P &= e_2 a_{2s+1}^{31} a_{2r-1}^{14} a_{2t}^{34} (e_2 a_{2r}^{14} + a_{2s+2}^{31} a_{2t-1}^{34}) = \\ e_2 a_{2s+1}^{31} a_{2r-1}^{14} a_{2t}^{34} &(a_{2r}^{14} + e_2 a_{2s+2}^{31} a_{2t-1}^{34}), \end{aligned}$$

and we select

$$P = a_{2r}^{14} + e_2 a_{2s+2}^{31} a_{2t-1}^{34}.$$

Thus,

$$P = a_{2r}^{14} + e_2 a_{2s+2}^{31} a_{2t-1}^{34} \xrightarrow{\varphi_1} e_1 a_{2r}^{23} + e_1 a_{2s+2}^{24} a_{2t}^{34} = \tilde{P}. \quad (\text{B.79})$$

By (4.48)

$$e_1 a_{2s+2}^{24} a_{2t}^{34} = e_1 a_{2s+2}^{24} a_{2t}^{34},$$

and by (B.79), we get

$$\tilde{P} = e_1 a_{2r}^{23} + e_1 a_{2s+2}^{24} a_{2t+2}^{34}. \quad (\text{B.80})$$

By (B.80) and (B.78) $f_{1\alpha}$ and $e_{1\alpha}$ correspond to Line $H11$ under the substitution $t \mapsto t+1$. \square

Cases $G31$ (actions ψ_1, ψ_2, ψ_4), and $G41$ (actions ψ_1, ψ_2, ψ_3) are similarly proved. \square

B.2.9. Line $H11$, $e_\alpha \xrightarrow{\varphi_2} e_{2\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_2} f_{2\alpha 0}$. Here,

$$\alpha = (14)^r(31)^s(21)^t, e_\alpha = e_{(23)^{2r-1}(24)^{2s-1}(34)^{2t+1}}.$$

Since $2\alpha = 2(14)^r(31)^s(21)^t$, and by Corollary 4.5.2 we have

$$\begin{aligned} a_{2r-1}^{23} &\xrightarrow{\varphi_2} e_2 a_{2r-1}^{14}, \\ a_{2s-1}^{24} &\xrightarrow{\varphi_2} e_2 a_{2s-1}^{31}, \\ e_1 a_{2t+1}^{34} &\xrightarrow{\varphi_2} e_2 a_{2t+2}^{34}, \end{aligned}$$

i.e.,

$$e_\alpha = e_1 a_{2r-1}^{23} a_{2s-1}^{24} a_{2t+1}^{34} \xrightarrow{\varphi_2} e_2 a_{2r-1}^{14} a_{2s-1}^{31} a_{2t+2}^{34} = e_{2\alpha}. \quad (\text{B.81})$$

The polynomial $e_{2\alpha}$ corresponds to Line $H21$. For the case $f_{\alpha 0}$, select

$$P = e_1 a_{2t}^{34} + a_{2s}^{24} a_{2r}^{23},$$

and we have

$$P = e_1 a_{2t}^{34} + a_{2s}^{24} a_{2r}^{23} \xrightarrow{\varphi_2} e_2 a_{2t+1}^{34} + e_2 a_{2s}^{13} a_{2r}^{14} = \tilde{P}. \quad (\text{B.82})$$

By (B.82) and (B.81) $f_{2\alpha}$ and $e_{2\alpha}$ correspond to Line $H21$. \square

Cases $H11$ (actions ψ_3, ψ_4) are similarly proved. \square

B.2.10. Line $H21$, $e_\alpha \xrightarrow{\varphi_1} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_1} f_{1\alpha 0}$. We have

$$\alpha = 2(14)^r(31)^s(21)^t, e_\alpha = e_{2(14)^r(31)^s(21)^t}.$$

Since $1\alpha = 12(14)^r(31)^s(21)^t = (14)^r(31)^s(21)^{t+1}$ (see heading (10) of Proposition 4.1.1), and by Corollary 4.5.2 we have

$$\begin{aligned} a_{2r-1}^{14} &\xrightarrow{\varphi_1} e_1 a_{2r-1}^{23}, \\ a_{2s-1}^{13} &\xrightarrow{\varphi_1} e_1 a_{2s-1}^{24}, \\ e_2 a_{2t+2}^{34} &\xrightarrow{\varphi_1} e_1 a_{2t+3}^{34}, \end{aligned}$$

i.e.,

$$e_\alpha = e_2 a_{2r-1}^{14} a_{2s-1}^{13} a_{2t+2}^{34} \xrightarrow{\varphi_1} e_1 a_{2r-1}^{23} a_{2s-1}^{24} a_{2t+3}^{34} = e_{1\alpha}. \quad (\text{B.83})$$

The polynomial $e_{2\alpha}$ corresponds to Line $H11$ under the substitution $t \mapsto t+1$. For the case $f_{\alpha 0}$, we select

$$P = a_{2r}^{14} + e_2 a_{2s}^{31} a_{2t+1}^{34},$$

and we have

$$P = a_{2r}^{14} + e_2 a_{2s}^{31} a_{2t+1}^{34} \xrightarrow{\varphi_1} e_1 a_{2r}^{23} + e_1 a_{2s}^{24} a_{2t+2}^{34} = \tilde{P}. \quad (\text{B.84})$$

By (B.84) and (B.83) $f_{1\alpha}$ and $e_{1\alpha}$ correspond to Line $H11$ under the substitution $t \mapsto t+1$. \square

B.2.11. Line $H21$, $e_\alpha \xrightarrow{\varphi_3} e_{1\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_3} f_{3\alpha 0}$. The admissible sequence α and the polynomial e_α are given as above in §B.2.10.

Since $3\alpha = 32(14)^r(31)^s(21)^t = (31)^s(21)^{t+1}(41)^r$ (see heading (12) of Proposition 4.1.1), and by Corollary 4.5.2 we have

$$\begin{aligned} a_{2s-1}^{13} &\xrightarrow{\varphi_3} e_3 a_{2s-1}^{24}, \\ a_{2t+2}^{34} &\xrightarrow{\varphi_3} e_3 a_{2t+2}^{12}, \\ e_2 a_{2r-1}^{14} &\xrightarrow{\varphi_3} e_3 a_{2r}^{14}, \end{aligned}$$

i.e.,

$$e_\alpha = e_2 a_{2r-1}^{14} a_{2s-1}^{13} a_{2t+2}^{34} \xrightarrow{\varphi_3} e_3 a_{2r}^{14} a_{2s-1}^{24} a_{2t+2}^{12} = e_{3\alpha}. \quad (\text{B.85})$$

The polynomial $e_{3\alpha}$ corresponds to Line $F31$ under the substitution $t \mapsto t+1$. For the case $f_{\alpha 0}$, we select

$$P = e_2 a_{2r}^{14} + a_{2s}^{31} a_{2t+1}^{34},$$

and we have

$$P = e_2 a_{2r}^{14} + a_{2s}^{31} a_{2t+1}^{34} \xrightarrow{\varphi_3} e_3 a_{2r+1}^{14} + e_3 a_{2s}^{24} a_{2t+1}^{12} = \tilde{P}. \quad (\text{B.86})$$

By (B.85)

$$e_{3\alpha} \tilde{P} = e_{3\alpha} (e_3 a_{2r+1}^{14} + a_{2s}^{24} a_{2t+1}^{12}) = e_{3\alpha} (e_3 a_{2s}^{24} + a_{2r+1}^{14} a_{2t+1}^{12}),$$

and therefore, by (B.86) and (B.85), we see that $f_{1\alpha}$ and $e_{1\alpha}$ correspond to Line $F31$ under the substitution $t \mapsto t+1$. \square

B.2.12. Line $H21$, $e_\alpha \xrightarrow{\varphi_4} e_{4\alpha}$, $f_{\alpha 0} \xrightarrow{\varphi_4} f_{4\alpha 0}$. The admissible sequence α and the polynomial e_α are given as above in §B.2.10.

Since $4\alpha = 42(14)^r(31)^s(21)^t = (41)^s(21)^{t+1}(31)^r$ (see heading (13) of Proposition 4.1.1), and by Corollary 4.5.2 we have

$$\begin{aligned} a_{2t+2}^{34} &\xrightarrow{\varphi_4} e_4 a_{2t+2}^{12}, \\ a_{2r-1}^{14} &\xrightarrow{\varphi_4} e_4 a_{2r-1}^{23}, \\ e_2 a_{2s-1}^{13} &\xrightarrow{\varphi_4} e_4 a_{2s}^{13}, \end{aligned}$$

i.e.,

$$e_\alpha = e_2 a_{2r-1}^{14} a_{2s-1}^{13} a_{2t+2}^{34} \xrightarrow{\varphi_4} e_4 a_{2r-1}^{23} a_{2s}^{13} a_{2t+2}^{12} = e_{4\alpha}. \quad (\text{B.87})$$

The polynomial $e_{4\alpha}$ corresponds to Line $F41$ under the substitution $t \mapsto t+1$. For the case $f_{\alpha 0}$, we select

$$P = a_{2r}^{41} + e_2 a_{2s}^{31} a_{2t+1}^{34},$$

and we have

$$P = a_{2r}^{41} + e_2 a_{2s}^{31} a_{2t+1}^{34} \xrightarrow{\varphi_4} e_4 a_{2r}^{23} + e_4 a_{2s+1}^{31} a_{2t+1}^{12}. \quad (\text{B.88})$$

By (B.88) and (B.87) we see that $f_{4\alpha}$ and $e_{4\alpha}$ correspond to Line $F41$ under the substitution $t \mapsto t+1$. \square

Cases $H31$, (actions ψ_1, ψ_3, ψ_4) and $H41$, (actions ψ_1, ψ_2, ψ_4) are similarly proved. \square

Theorem 4.8.1 on the admissible elements in D^4 is proved for all polynomials e_α , see Table 4.3. \square

APPENDIX C

The inclusion theorem and properties of $H^+(n)$

C.1. Proof of the inclusion theorem

C.1.1. More properties of the atomic elements. The atomic elements a_n^{ij} and A_n^{ji} can frequently substitute each other. This has already been shown in Lines 3.1 – 5.2 of Table 2.1.

PROPOSITION C.1.1. *The following relations hold for $n - 2 \leq m$:*

- (1) $y_i a_m^{ik} a_n^{kj} = y_i a_m^{ik} A_n^{jk}$,
- (2) $y_i A_m^{ji} a_n^{jk} = y_i A_m^{ji} A_n^{kj}$.

Proof. 1) Since

$$y_i a_m^{ik} a_n^{kj} = y_i a_m^{ik} (x_k + y_j (x_j + y_i a_{n-2}^{ik}))$$

and $y_i a_{n-2}^{ik} \supseteq y_i a_m^{ik}$ for $n - 2 \leq m$, it follows that

$$y_i a_m^{ik} a_n^{kj} = y_i a_m^{ik} (y_j + x_k (x_j + y_i a_{n-2}^{ik})).$$

Since $a_{n-2}^{ik} \supseteq x_k$, we have

$$y_i a_m^{ik} a_n^{kj} = y_i a_m^{ik} (y_j + x_k (y_i + x_j a_{n-2}^{ik})).$$

Further, by Table 2.1, Line 3.1, we have $x_j a_{n-2}^{ik} = x_j A_{n-2}^{ki}$ and

$$y_i a_m^{ik} a_n^{kj} = y_i a_m^{ik} (y_j + x_k (y_i + x_j A_{n-2}^{ki})) = y_i a_m^{ik} A_n^{jk}. \quad \square$$

2) Since

$$y_i A_m^{ji} a_n^{jk} = y_i A_m^{ji} (x_j + y_k (x_k + y_i a_{n-2}^{ij})) = y_i A_m^{ji} (x_j + x_k + y_k y_i a_{n-2}^{ij})$$

and by Table 2.1, Line 4.1, we have $y_k y_i a_{n-2}^{ij} = y_k y_i A_{n-2}^{ji}$, and

$$y_i A_m^{ji} a_n^{jk} = y_i A_m^{ji} (x_j + x_k + y_k y_i A_{n-2}^{ji}).$$

Since $A_{n-2}^{ji} \supseteq A_m^{ji}$, we see that

$$\begin{aligned} y_i A_m^{ji} a_n^{jk} &= y_i A_m^{ji} (y_k y_i + A_{n-2}^{ji} (x_j + x_k)) = \\ y_i A_m^{ji} (y_k y_i + x_j + x_k A_{n-2}^{ji}) &= y_i A_m^{ji} (x_j + y_k (y_i + x_k A_{n-2}^{ji})). \end{aligned}$$

Further,

$$y_i + x_k A_{n-2}^{ji} \supseteq y_i A_m^{ji},$$

and we have

$$y_i A_m^{ji} a_n^{jk} = y_i A_m^{ji} (y_k + x_j (y_i + x_k A_{n-2}^{ji})) = y_i A_m^{ji} A_n^{kj}. \quad \square$$

C.1.2. The a -form and A -form of admissible elements.

PROPOSITION C.1.2. For all admissible elements f_α and e_α of Table 2.3,

- (1) Every symbol “ A ” can be substituted by the symbol “ a ”.
- (2) Every symbol “ a ” can be substituted by the symbol “ A ”.

The substitutions $A \longleftrightarrow a$ are collected in Table C.1. The original form of admissible elements is given by Table 2.3. Two other forms of admissible elements are given by Proposition C.1.2. The substitution (1) (resp. (2)) from Proposition C.1.2 is said to be an a -form (resp. A -form). (Table C.1).

Proof of Proposition C.1.2. For elements f_α , both statements (1) and (2) of Proposition C.1.2 follow from Table 2.1, headings (3.1), (4.1) and (5.2). For convenience, we give these properties here:

$$\begin{aligned} x_k A_n^{ij} &= x_k a_n^{ji} && \text{(Table 2.1, Line 3.1),} \\ y_i y_j A_n^{ki} &= y_i y_j a_n^{ik} && \text{(Table 2.1, Line 4.1),} \\ y_i (x_j + x_k) A_n^{ij} &= y_i (x_j + x_k) a_n^{ji} && \text{(Table 2.1, Line 5.2).} \end{aligned}$$

Consider elements e_α . We will use Proposition C.1.2 and property 5.1 from Table 2.1. We have

$$a_n^{ij} a_m^{jk} = A_n^{ji} a_m^{jk} \text{ for } n \leq m+1 \quad \text{(Table 2.1, Line 5.1).}$$

Line 1, Line 9 of Table 2.3. Take Line 1:

$$e_\alpha = y_2 A_{k-1}^{32} a_q^{21} A_k^{13} a_q^{13}.$$

According to Line 5.1 of Table 2.1, we have

$$a_q^{21} a_q^{13} = A_q^{12} a_q^{13} \text{ and } e_\alpha = y_2 A_{k-1}^{32} A_q^{12} A_k^{13} a_q^{13}.$$

Further, by Proposition C.1.1, (2) we have

$$y_2 A_k^{12} a_q^{13} = y_2 A_k^{12} A_q^{31} \text{ and } e_\alpha = y_2 A_{k-1}^{32} A_q^{12} A_k^{13} A_q^{31}.$$

This is the A -form of e_α .

On the other hand, by Proposition C.1.1, (2) we have

$$y_2 A_{k-1}^{32} A_k^{13} = y_2 A_{k-1}^{32} a_k^{31} \text{ and } e_\alpha = y_2 A_{k-1}^{32} a_q^{21} a_k^{31} a_q^{13}.$$

Again, by Line 5.1 of Table 2.1, we have

$$A_{k-1}^{32} a_k^{31} = a_{k-1}^{23} a_k^{31} \text{ and } e_\alpha = y_2 a_{k-1}^{23} a_q^{21} a_k^{31} a_q^{13}.$$

It is the a -form of e_α . The same for Line 9 of Table 2.3. \square

Line 2, Line 10 of Table 2.3. For Line 2, we have

$$e_\alpha = y_3 a_{k-1}^{31} A_{q+1}^{12} a_k^{12} A_q^{23}.$$

By Table 2.1, Line 5.1 we have

$$a_{k-1}^{31} a_k^{12} = A_{k-1}^{13} a_k^{12} \text{ and } e_\alpha = y_3 A_{k-1}^{13} A_{q+1}^{12} a_k^{12} A_q^{23}.$$

By proposition C.1.1, (2) we obtain

$$y_3 A_{k-1}^{13} a_k^{12} = y_3 A_{k-1}^{13} A_k^{21} \text{ and } e_\alpha = y_3 A_{k-1}^{13} A_{q+1}^{12} A_k^{21} A_q^{23} \quad (A\text{-form}).$$

On the other hand, by proposition C.1.1, (2) we have

$$y_3 A_q^{23} A_{q+1}^{12} = y_3 A_q^{23} a_{q+1}^{21} \text{ and } e_\alpha = y_3 a_{k-1}^{31} a_{q+1}^{21} a_k^{12} A_q^{23}.$$

N	α	f_α , $a\text{-form} = A\text{-form}$	e_α , $a\text{-form} = A\text{-form}$	$g_{\alpha 0}$
1	$(213)^{2p}(21)^{2k}$	$y_1 y_2 a_q^{13} a_{k-1}^{23} =$ $y_1 y_2 A_q^{31} A_{k-1}^{32}$	$y_2 a_{k-1}^{23} a_q^{21} a_k^{31} a_q^{13} =$ $y_2 A_{k-1}^{32} A_q^{12} A_k^{13} A_q^{31}$	$e_\alpha(x_1 + a_q^{32} A_k^{32})$
2	$3(213)^{2p}(21)^{2k}$	$y_3(x_1 + x_2) a_q^{32} a_{k-1}^{31} =$ $y_3(x_1 + x_2) A_q^{23} A_{k-1}^{13}$	$y_3 a_{k-1}^{31} a_{q+1}^{21} a_k^{12} a_q^{32} =$ $y_3 A_{k-1}^{13} A_{q+1}^{12} A_k^{21} A_q^{23}$	$e_\alpha(y_2 y_3 + A_q^{12} a_k^{31})$
3	$13(213)^{2p}(21)^{2k}$	$y_3 y_1 a_{q+1}^{32} a_{k-1}^{12} =$ $y_3 y_1 A_{q+1}^{23} A_{k-1}^{21}$	$y_1 a_{k-1}^{12} a_{q+1}^{13} a_k^{23} a_{q+1}^{32} =$ $y_1 A_{k-1}^{21} A_{q+1}^{31} A_k^{32} A_{q+1}^{23}$	$e_\alpha(x_3 + a_{q+1}^{21} A_k^{21})$
4	$(213)^{2p+1}(21)^{2k}$	$y_2(x_3 + x_1) a_{q+1}^{21} a_{k-1}^{23} =$ $y_2(x_3 + x_1) A_{q+1}^{12} A_{k-1}^{32}$	$y_2 a_{k-1}^{23} a_{q+2}^{13} a_k^{31} a_{q+1}^{21} =$ $y_2 A_{k-1}^{32} A_{q+2}^{31} A_k^{13} A_{q+1}^{12}$	$e_\alpha(y_1 y_2 + A_{q+1}^{31} a_k^{23})$
5	$3(213)^{2p+1}(21)^{2k}$	$y_2 y_3 a_{q+2}^{21} a_{k-1}^{31} =$ $y_2 y_3 A_{q+2}^{12} A_{k-1}^{13}$	$y_3 a_{k-1}^{31} a_{q+2}^{32} a_k^{12} a_{q+2}^{21} =$ $y_3 A_{k-1}^{13} A_{q+2}^{23} A_k^{21} A_{q+2}^{12}$	$e_\alpha(x_2 + a_{q+2}^{13} A_k^{13})$
6	$13(213)^{2p+1}(21)^{2k}$	$y_1(x_2 + x_3) a_{q+2}^{13} a_{k-1}^{12} =$ $y_1(x_2 + x_3) A_{q+2}^{31} A_{k-1}^{21}$	$y_1 a_{k-1}^{12} a_{q+3}^{32} a_k^{23} a_{q+2}^{13} =$ $y_1 A_{k-1}^{21} A_{q+3}^{23} A_k^{32} A_{q+2}^{31}$	$e_\alpha(y_1 y_3 + A_{q+2}^{23} a_k^{12})$
7	$1(21)^{2k}$	$x_1 a_k^{32} a_k^{23} =$ $x_1 A_k^{23} A_k^{32}$	$y_1 a_k^{13} a_k^{12} a_k^{32} a_k^{23} =$ $y_1 A_k^{31} A_k^{21} A_k^{23} A_k^{32}$	$e_\alpha(y_2 a_k^{21} + y_3 a_k^{31})$
8	$1(21)^{2k+1}$	$y_1(x_2 + x_3) a_k^{13} a_k^{12} =$ $y_1(x_2 + x_3) A_k^{31} A_k^{21}$	$y_1 a_k^{13} a_k^{12} a_{k+1}^{32} a_{k+1}^{23} =$ $y_1 A_k^{31} A_k^{21} A_{k+1}^{23} A_{k+1}^{32}$	$e_\alpha(y_1 a_{k+1}^{12} + y_1 a_{k+1}^{13})$
9	$(213)^{2p}(21)^{2k+1}$	$y_2 y_3 a_q^{21} a_k^{31} =$ $y_2 y_3 A_q^{12} A_k^{13}$	$y_2 a_k^{23} a_q^{21} a_k^{31} a_{q+1}^{13} =$ $y_2 A_k^{32} A_q^{12} A_k^{13} A_{q+1}^{31}$	$e_\alpha(x_2 + a_q^{13} A_{k+1}^{13})$
10	$3(213)^{2p}(21)^{2k+1}$	$x_3 a_{q+1}^{21} a_k^{12} =$ $x_3 A_{q+1}^{12} A_k^{21}$	$y_3 a_k^{31} a_{q+1}^{21} a_k^{12} a_{q+1}^{32} =$ $y_3 A_k^{13} A_{q+1}^{12} A_k^{21} A_{q+1}^{23}$	$e_\alpha(y_1 y_3 + A_q^{23} a_{k+1}^{12})$
11	$13(213)^{2p}(21)^{2k+1}$	$y_1 y_2 a_{q+1}^{13} a_k^{23} =$ $y_1 y_2 A_{q+1}^{31} A_k^{32}$	$y_1 a_k^{12} a_{q+1}^{13} a_k^{23} a_{q+2}^{32} =$ $y_1 A_k^{21} A_{q+1}^{31} A_k^{32} A_{q+2}^{23}$	$e_\alpha(x_1 + a_{q+1}^{32} A_{k+1}^{32})$
12	$(213)^{2p+1}(21)^{2k+1}$	$x_2 a_{q+2}^{13} a_k^{31} =$ $x_2 A_{q+2}^{31} A_k^{13}$	$y_2 a_k^{23} a_{q+2}^{13} a_k^{31} a_{q+2}^{21} =$ $y_2 A_k^{32} A_{q+2}^{31} A_k^{13} A_{q+2}^{12}$	$e_\alpha(y_2 y_3 + A_{q+1}^{12} a_{k+1}^{31})$
13	$3(213)^{2p+1}(21)^{2k+1}$	$y_3 y_1 a_{q+2}^{32} a_k^{12} =$ $y_3 y_1 A_{q+2}^{23} A_k^{21}$	$y_3 a_k^{31} a_{q+2}^{32} a_k^{12} a_{q+3}^{21} =$ $y_3 A_k^{13} A_{q+2}^{23} A_k^{21} A_{q+3}^{12}$	$e_\alpha(x_3 + a_{q+2}^{21} A_{k+1}^{21})$
14	$13(213)^{2p+1}(21)^{2k+1}$	$x_1 a_{q+3}^{32} a_k^{23} =$ $x_1 A_{q+3}^{23} A_k^{32}$	$y_1 a_k^{12} a_{q+3}^{32} a_k^{23} a_{q+3}^{13} =$ $y_1 A_k^{21} A_{q+3}^{23} A_k^{32} A_{q+3}^{31}$	$e_\alpha(y_1 y_2 + A_{q+2}^{31} a_{k+1}^{23})$

TABLE C.1. The admissible elements, a -form and A -form

In Lines 1 – 6: $k > 0$ and $p \geq 0$. In Lines 7 – 8: $k > 0$.

In Lines 9 – 14: $k \geq 0, p \geq 0$. In all Lines: $q = k + 3p$.

By Table 2.1, Line 5.1 we get

$$A_q^{23} a_{q+1}^{21} = a_q^{32} a_{q+1}^{21} \text{ and } e_\alpha = y_3 a_{k-1}^{31} a_{q+1}^{21} a_k^{12} a_q^{32}, \quad (a\text{-form}).$$

The same for Line 10 of Table 2.3. \square

Lines 3–6, Lines 11–14 of Table 2.3. The proof for Line 3 follows from Line 1 under substitution $1 \rightarrow 3 \rightarrow 2 \rightarrow 1$. Under the same substitution the case of Line 11 follows from the case of Line 10. Similarly, by the same substitution we have

$$\begin{aligned} \text{Line 1} &\implies \text{Line 3} \implies \text{Line 5}, \\ \text{Line 9} &\implies \text{Line 11} \implies \text{Line 13}, \\ \text{Line 2} &\implies \text{Line 4} \implies \text{Line 6}, \\ \text{Line 10} &\implies \text{Line 12} \implies \text{Line 14}. \quad \square \end{aligned}$$

Lines 7–8, Table 2.3. Let us take Line 7, where $e_\alpha = y_1 A_k^{31} a_k^{12} A_k^{23} a_k^{23}$. By Proposition C.1.1, (2) we have

$$y_1 A_k^{23} a_k^{23} = y_1 A_k^{23} A_k^{32} \text{ and } e_\alpha = y_1 A_k^{31} a_k^{12} A_k^{23} A_k^{32}.$$

Again,

$$y_1 A_k^{32} a_k^{12} = y_1 A_k^{32} A_k^{21}, \text{ so } e_\alpha = y_1 A_k^{31} A_k^{21} A_k^{23} A_k^{32} \quad (A\text{-form}).$$

On the other hand, by Proposition C.1.1, (2) we have

$$y_1 A_k^{31} A_k^{23} = y_1 A_k^{31} a_k^{32} \text{ and } e_\alpha = y_1 A_k^{31} a_k^{12} a_k^{32} a_k^{23}.$$

By Table 2.1, Line 5.1 we have

$$A_k^{31} a_k^{32} = a_k^{13} a_k^{32}, \text{ so } e_\alpha = y_1 a_k^{13} a_k^{12} a_k^{32} a_k^{23} \quad (a\text{-form}).$$

The same for Line 8 of Table 2.3. \square

REMARK C.1.3 (to Table C.2). Transformation T^{123} substitutes $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ and T^{12} substitutes $1 \rightarrow 2 \rightarrow 1$. These transformations are needed for getting corresponding admissible elements $g_{\alpha 0}$ from the Table 2.3 or their a -forms and A -forms from Table C.1. For example, consider Line 2. In this case $\alpha = (321)^{2p}(32)^{2k}$. This sequence can be obtained by applying T^{123} to the sequence $\tilde{\alpha} = (213)^{2p}(21)^{2k}$ from Table 2.3, i.e.,

$$\alpha = (321)^{2p}(32)^{2k} = T^{123}[\tilde{\alpha}] = T^{123}[(213)^{2p}(21)^{2k}].$$

C.1.3. Admissible sequences $\alpha 1$ and α . The inclusion theorem. Our intention is to prove the following theorem.

Theorem 2.13.1 *For every admissible sequence α or αi , where $i = 1, 2, 3$ from Table 2.2, the following inclusion holds:*

$$e_{\alpha i} \subseteq g_{\alpha 0}, \quad i = 1, 2, 3.$$

Without loss of generality it suffices to prove Theorem 2.13.1 for $i = 1$. Thus, we need description of admissible sequences $\alpha 1$ and α in the form of sequences of Table 2.3.

PROPOSITION C.1.4. *For the corresponding indices $\alpha 1$ and α from Table C.2 (columns 2 and 3), we have*

$$\varphi_\alpha(1) = \alpha 1.$$

Proof. Line 1. Let $p > 0$. We have

$$\begin{aligned} \varphi_{21(321)^{2p-1}(32)^{2k}}(1) &= 21(321)^{2p-1}(32)^{2k}1 = \\ &= 21(321)(321)^{2p-2}(32)^{2k}1 = 213((21)(321)^{2p-2}(32)^{2k}1). \end{aligned}$$

By induction, from Line 4, Table C.2 we get

$$\varphi_{21(321)^{2p-1}(32)^{2k}}(1) = 213[(213)^{2p-1}(21)^{2k}] = (213)^{2p}(21)^{2k} = \alpha 1.$$

If $p = 0$, then $2(12)^{2k}1 = (21)^{2k+1}$. The same for Line 9. \square

N	$\alpha 1$ index of $e_{\alpha 1}$	α (index of $g_{\alpha 0}$) = $T^{123}[\tilde{\alpha}]$ or $T^{12}[\tilde{\alpha}]$	$e_{\alpha 1}$ A -form	$g_{\alpha 0}$ = $T^{123}[g_{\alpha 0}]$ or $T^{12}[g_{\alpha 0}]$
1	$(213)^{2p}(21)^{2k}$	$21(321)^{2p-1}(32)^{2k} =$ $T^{123}[13(213)^{2p-1}(21)^{2k}]$	$y_2 A_{k-1}^{32} A_q^{12} A_k^{13} A_q^{31}$	$y_2 A_{k-1}^{32} A_q^{31} A_k^{13} A_{q-1}^{12}$ $\cap (y_2 y_1 + A_{q-1}^{31} a_k^{23})$
1'	$(21)^{2k}$	$2(12)^{2k-1} =$ $T^{12}[1(21)^{2k-1}]$	$y_2 A_{k-1}^{32} A_k^{12} A_k^{13} A_k^{31}$	$y_2 A_{k-1}^{32} A_{k-1}^{12} A_k^{13} A_k^{31}$ $\cap (y_2 a_k^{21} + y_2 a_k^{23})$
2	$3(213)^{2p}(21)^{2k}$	$(321)^{2p}(32)^{2k} =$ $T^{123}[(213)^{2p}(21)^{2k}]$	$y_3 A_{k-1}^{13} A_{q+1}^{12} A_k^{21} A_q^{23}$	$y_3 A_{k-1}^{13} A_q^{23} A_k^{21} A_q^{12}$ $\cap (x_2 + a_q^{13} A_k^{13})$
3	$13(213)^{2p}(21)^{2k}$	$1(321)^{2p}(32)^{2k} =$ $T^{123}[3(213)^{2p}(21)^{2k}]$	$y_1 A_{k-1}^{21} A_{q+1}^{31} A_k^{32} A_{q+1}^{23}$	$y_1 A_{k-1}^{21} A_{q+1}^{23} A_k^{32} A_q^{31}$ $\cap (y_3 y_1 + A_q^{23} a_k^{12})$
4	$(213)^{2p+1}(21)^{2k}$	$21(321)^{2p}(32)^{2k} =$ $T^{123}[13(213)^{2p}(21)^{2k}]$	$y_2 A_{k-1}^{32} A_{q+2}^{31} A_k^{13} A_{q+1}^{12}$	$y_2 A_{k-1}^{32} A_{q+1}^{12} A_k^{13} A_{q+1}^{31}$ $\cap (x_1 + a_{q+1}^{32} A_k^{32})$
5	$3(213)^{2p+1}(21)^{2k}$	$(321)^{2p+1}(32)^{2k} =$ $T^{123}[(213)^{2p+1}(21)^{2k}]$	$y_3 A_{k-1}^{13} A_{q+2}^{23} A_k^{21} A_{q+2}^{12}$	$y_3 A_{k-1}^{13} A_{q+2}^{12} A_k^{21} A_{q+1}^{23}$ $\cap (y_2 y_3 + A_{q+1}^{12} a_k^{31})$
6	$13(213)^{2p+1}(21)^{2k}$	$1(321)^{2p+1}(32)^{2k} =$ $T^{123}[3(213)^{2p+1}(21)^{2k}]$	$y_1 A_{k-1}^{21} A_{q+3}^{23} A_k^{32} A_{q+2}^{31}$	$y_1 A_{k-1}^{21} A_{q+2}^{31} A_k^{32} A_{q+2}^{23}$ $\cap (x_3 + a_{q+2}^{21} A_k^{21})$
7	$1(21)^{2k}$	$(12)^{2k} =$ $T^{12}[(21)^{2k}]$	$y_1 A_k^{31} A_k^{21} A_k^{23} A_k^{32}$	$y_1 A_{k-1}^{31} A_k^{21} A_k^{23} A_k^{32}$ $\cap (x_2 + a_k^{31} A_k^{31})$
8	$1(21)^{2k+1}$	$(12)^{2k+1} =$ $T^{12}[(21)^{2k+1}]$	$y_1 A_k^{31} A_k^{21} A_{k+1}^{23} A_{k+1}^{32}$	$y_1 A_k^{31} A_k^{21} A_k^{23} A_{k+1}^{32}$ $\cap (x_1 + a_k^{23} A_{k+1}^{23})$
9	$(213)^{2p}(21)^{2k+1}$	$21(321)^{2p-1}(32)^{2k+1} =$ $T^{123}[13(213)^{2p-1}(21)^{2k+1}]$	$y_2 A_k^{32} A_q^{12} A_k^{13} A_{q+1}^{31}$	$y_2 A_k^{32} A_q^{31} A_k^{13} A_q^{12}$ $\cap (y_2 y_3 + A_{q-1}^{12} a_{k+1}^{12})$
9'	$(21)^{2k+1}$	$2(12)^{2k} =$ $T^{12}[1(21)^{2k}]$	$y_2 A_k^{32} A_k^{12} A_k^{13} A_{k+1}^{31}$	$y_2 A_k^{32} A_{k+1}^{12} A_k^{13} A_k^{31}$ $\cap (y_2 a_k^{21} + y_2 a_k^{23})$
10	$3(213)^{2p}(21)^{2k+1}$	$(321)^{2p}(32)^{2k+1} =$ $T^{123}[(213)^{2p}(21)^{2k+1}]$	$y_3 A_k^{13} A_{q+1}^{12} A_k^{21} A_{q+1}^{23}$	$y_3 A_k^{13} A_q^{23} A_k^{21} A_{q+1}^{12}$ $\cap (x_3 + a_q^{21} A_{k+1}^{21})$
11	$13(213)^{2p}(21)^{2k+1}$	$1(321)^{2p}(32)^{2k+1} =$ $T^{123}[3(213)^{2p}(21)^{2k+1}]$	$y_1 A_k^{21} A_{q+1}^{31} A_k^{32} A_{q+2}^{23}$	$y_1 A_k^{21} A_{q+1}^{23} A_k^{32} A_{q+1}^{31}$ $\cap (y_2 y_1 + A_q^{31} a_{k+1}^{23})$
12	$(213)^{2p+1}(21)^{2k+1}$	$21(321)^{2p}(32)^{2k+1} =$ $T^{123}[(213)^{2p+1}(21)^{2k+1}]$	$y_2 A_k^{32} A_{q+2}^{31} A_k^{13} A_{q+2}^{12}$	$y_2 A_k^{32} A_{q+2}^{12} A_k^{13} A_{q+2}^{31}$ $\cap (x_2 + a_{q+1}^{13} A_{k+1}^{13})$
13	$3(213)^{2p+1}(21)^{2k+1}$	$(321)^{2p+1}(32)^{2k+1} =$ $T^{123}[(213)^{2p+1}(21)^{2k+1}]$	$y_3 A_k^{13} A_{q+2}^{23} A_k^{21} A_{q+3}^{12}$	$y_3 A_k^{13} A_{q+2}^{12} A_k^{21} A_{q+2}^{23}$ $\cap (y_3 y_1 + A_{q+1}^{23} a_{k+1}^{12})$
14	$13(213)^{2p+1}(21)^{2k+1}$	$1(321)^{2p+1}(32)^{2k+1} =$ $T^{123}[3(213)^{2p+1}(21)^{2k+1}]$	$y_1 A_k^{21} A_{q+3}^{23} A_k^{32} A_{q+3}^{31}$	$y_1 A_k^{21} A_{q+2}^{31} A_k^{32} A_{q+3}^{23}$ $\cap (x_1 + a_{q+2}^{32} A_{k+1}^{32})$

TABLE C.2. $\alpha 1$, α , $e_{\alpha 1}$ and $g_{\alpha 0}$ For Line 1 and Line 9: $p > 0$. For Lines 1–6: $k > 0, p \geq 0$.For Lines 7–8: $k > 0$. For Lines 9–14: $k \geq 0, p \geq 0$. For all lines: $q = k + 3p$.Line 4. Here,

$$21(321)^{2p}(32)^{2k}1 = 21(321)(321)^{2p-1}(32)^{2k}1 = 213((21)(321)^{2p-1}(32)^{2k}1).$$

By induction, from heading 1) we get

$$21(321)^{2p}(32)^{2k}1 = (213)(213)^{2p}(21)^{2k} = (213)^{2p+1}(21)^{2k} = \alpha 1.$$

The same for Line 12. \square

Other lines of Table C.2 are considered by analogy with these two cases. \square

C.1.4. An S -form of $g_{\alpha 0}$.

DEFINITION C.1.5. The form of the elements $g_{\alpha 0}$ from the right column of Table C.3 is said to be an S -form (symmetric).

PROPOSITION C.1.6. Both forms of the elements $g_{\alpha 0}$ from Table C.3 coincide.

N	$g_{\alpha 0}$ from Table C.2	$g_{\alpha 0}$ S -form
1	$y_2 A_{k-1}^{32} A_q^{31} A_k^{13} A_{q-1}^{12} (y_2 y_1 + A_{q-1}^{31} a_k^{23})$	$y_2 A_{k-1}^{32} A_q^{31} A_k^{13} A_{q-1}^{12} (a_q^{21} + a_{k+1}^{12})$
1'	$y_2 A_{k-1}^{32} A_{k-1}^{12} A_k^{13} A_k^{31} (y_2 a_k^{21} + y_2 a_k^{23})$	$y_2 A_{k-1}^{32} A_{k-1}^{12} A_k^{13} A_k^{31} (a_k^{21} + a_{k+1}^{12})$
2	$y_3 A_{k-1}^{13} A_q^{23} A_k^{21} A_q^{12} (x_2 + a_q^{13} A_k^{13})$	$y_3 A_{k-1}^{13} A_q^{23} A_k^{21} A_q^{12} (A_{q+1}^{12} + A_k^{13})$
3	$y_1 A_{k-1}^{21} A_{q+1}^{23} A_k^{32} A_q^{31} (y_3 y_1 + A_q^{23} a_k^{12})$	$y_1 A_{k-1}^{21} A_{q+1}^{23} A_k^{32} A_q^{31} (a_{q+1}^{13} + a_{k+1}^{31})$
4	$y_2 A_{k-1}^{32} A_{q+1}^{12} A_k^{13} A_{q+1}^{31} (x_1 + a_{q+1}^{32} A_k^{32})$	$y_2 A_{k-1}^{32} A_{q+1}^{12} A_k^{13} A_{q+1}^{31} (A_{q+2}^{31} + A_k^{32})$
5	$y_3 A_{k-1}^{13} A_{q+2}^{12} A_k^{21} A_{q+1}^{23} (y_2 y_3 + A_{q+1}^{12} a_k^{31})$	$y_3 A_{k-1}^{13} A_{q+2}^{12} A_k^{21} A_{q+1}^{23} (a_{q+2}^{32} + a_{k+1}^{23})$
6	$y_1 A_{k-1}^{21} A_{q+2}^{31} A_k^{32} A_{q+2}^{23} (x_3 + a_{q+2}^{21} A_k^{21})$	$y_1 A_{k-1}^{21} A_{q+2}^{31} A_k^{32} A_{q+2}^{23} (A_{q+3}^{23} + A_k^{21})$
7	$y_1 A_{k-1}^{31} A_k^{21} A_k^{23} A_k^{32} (x_2 + a_k^{31} A_k^{31})$	$y_1 A_{k-1}^{31} A_k^{21} A_k^{23} A_k^{32} (A_{k+1}^{32} + A_k^{31})$
8	$y_1 A_k^{31} A_k^{21} A_k^{23} A_{k+1}^{32} (x_1 + a_k^{23} A_{k+1}^{23})$	$y_1 A_k^{31} A_k^{21} A_k^{23} A_{k+1}^{32} (A_{k+1}^{21} + A_{k+1}^{23})$
9	$y_2 A_k^{32} A_q^{31} A_k^{13} A_q^{12} (y_2 y_3 + A_{q-1}^{12} a_{k+1}^{31})$	$y_2 A_k^{32} A_q^{31} A_k^{13} A_q^{12} (a_{q+1}^{13} + a_{k+1}^{31})$
9'	$y_2 A_k^{32} A_k^{12} A_k^{13} A_k^{31} (y_1 a_k^{12} + y_3 a_k^{32})$	$y_2 A_k^{32} A_k^{12} A_k^{13} A_k^{31} (a_{k+1}^{31} + a_{k+1}^{13})$
10	$y_3 A_k^{13} A_q^{23} A_k^{21} A_{q+1}^{12} (x_3 + a_q^{21} A_{k+1}^{21})$	$y_3 A_k^{13} A_q^{23} A_k^{21} A_{q+1}^{12} (A_{q+1}^{23} + A_{k+1}^{21})$
11	$y_1 A_k^{21} A_{q+1}^{23} A_k^{32} A_{q+1}^{31} (y_2 y_1 + A_q^{31} a_{k+1}^{23})$	$y_1 A_k^{21} A_{q+1}^{23} A_k^{32} A_{q+1}^{31} (a_{q+2}^{32} + a_{k+1}^{23})$
12	$y_2 A_k^{32} A_{q+1}^{12} A_k^{13} A_{q+2}^{31} (x_2 + a_{q+1}^{13} A_{k+1}^{13})$	$y_2 A_k^{32} A_{q+1}^{12} A_k^{13} A_{q+2}^{31} (A_{q+2}^{12} + A_{k+1}^{13})$
13	$y_3 A_k^{13} A_{q+2}^{12} A_k^{21} A_{q+2}^{23} (y_3 y_1 + A_{q+1}^{23} a_{k+1}^{12})$	$y_3 A_k^{13} A_{q+2}^{12} A_k^{21} A_{q+2}^{23} (a_{q+3}^{21} + a_{k+1}^{12})$
14	$y_1 A_k^{21} A_{q+2}^{31} A_k^{32} A_{q+3}^{23} (x_1 + a_{q+2}^{32} A_{k+1}^{32})$	$y_1 A_k^{21} A_{q+2}^{31} A_k^{32} A_{q+3}^{23} (A_{q+3}^{31} + A_{k+1}^{32})$

TABLE C.3. The S -forms of $g_{\alpha 0}$

Proof. 1) Since $e_\alpha \subseteq A_q^{31} \subseteq A_{q-1}^{31}$, we see that

$$e_\alpha (y_2 y_1 + A_{q-1}^{31} a_k^{23}) = e_\alpha (y_2 y_1 A_{q-1}^{31} + a_k^{23}).$$

Since $y_2 y_1 A_{q-1}^{31} = y_2 y_1 a_{q-1}^{13}$ and $e_\alpha \subseteq y_2$, it follows that

$$e_\alpha(y_2 y_1 + A_{q-1}^{31} a_k^{23}) = e_\alpha(y_2 y_1 a_{q-1}^{13} + a_k^{23}) = e_\alpha(y_1 a_{q-1}^{13} + y_2 a_k^{23}).$$

Finally, since $x_2 \subseteq y_2 a_k^{23}$ and $x_1 \subseteq y_1 a_{q-1}^{13}$, it follows that

$$e_\alpha(y_2 y_1 + A_{q-1}^{31} a_k^{23}) = e_\alpha((x_2 + y_1 a_{q-1}^{13}) + (x_1 + y_2 a_k^{23})) = e_\alpha(a_q^{21} + a_{k+1}^{12}). \quad \square$$

1') Since $e_\alpha \subseteq y_2$, we have

$$e_\alpha(y_2 a_k^{21} + y_2 a_k^{23}) = e_\alpha(a_k^{21} + y_2 a_k^{23}).$$

Since $x_1 \subseteq a_k^{21}$, we have

$$e_\alpha(y_2 a_k^{21} + y_2 a_k^{23}) = e_\alpha(a_k^{21} + (x_1 + y_2 a_k^{23})) = e_\alpha(a_k^{21} + a_{k+1}^{12}). \quad \square$$

2) Since

$$a_q^{13} = x_1 + y_3 a_{q-1}^{32} \supseteq y_3 a_{q-1}^{32} \supseteq y_3 a_q^{32} \supseteq e_\alpha$$

(Table C.1, Line 2, a -form), it follows that

$$e_\alpha(x_2 + a_q^{13} A_k^{13}) = e_\alpha(x_2 a_q^{13} + A_k^{13}) = e_\alpha(x_2 A_q^{31} + A_k^{13}).$$

Further, since $y_1 \subseteq A_k^{13}$, it follows that

$$e_\alpha(x_2 + a_q^{13} A_k^{13}) = e_\alpha(y_1 + x_2 A_q^{31} + A_k^{13}) = e_\alpha(A_{q+1}^{12} + A_k^{13}). \quad \square$$

3) Since $e_\alpha \subseteq A_{q+1}^{23} \subseteq A_q^{23}$, we see that

$$e_\alpha(y_3 y_1 + A_q^{23} a_k^{12}) = e_\alpha(y_3 y_1 A_q^{23} + a_k^{12}) = e_\alpha(y_3 y_1 a_q^{32} + a_k^{12}).$$

Since $e_\alpha \subseteq y_1$, it follows that

$$\begin{aligned} e_\alpha(y_3 y_1 a_q^{32} + a_k^{12}) &= e_\alpha(y_3 a_q^{32} + y_1 a_k^{12}) = \\ e_\alpha(x_1 + y_3 a_q^{32} + x_3 + y_1 a_k^{12}) &= e_\alpha(a_{q+1}^{13} + a_{k+1}^{31}). \quad \square \end{aligned}$$

4) Since $a_{q+1}^{32} = x_3 + y_2 a_q^{21} \supseteq y_2 a_{q+1}^{21} \supseteq e_\alpha$ (Table C.1, Line 4, a -form), we see that

$$\begin{aligned} e_\alpha(x_1 + a_{q+1}^{32} A_k^{32}) &= e_\alpha(x_1 a_{q+1}^{32} + A_k^{32}) = e_\alpha(x_1 A_{q+1}^{23} + A_k^{32}) = \\ e_\alpha(y_3 + x_1 A_{q+1}^{23} + A_k^{32}) &= e_\alpha(A_{q+2}^{31} + A_k^{32}). \quad \square \end{aligned}$$

5) Since $e_\alpha \subseteq y_3 A_{q+2}^{12} \subseteq y_3 A_{q+1}^{12}$ and $y_3 \supseteq e_\alpha$, we see that

$$\begin{aligned} e_\alpha(y_2 y_3 + A_{q+1}^{12} a_k^{31}) &= e_\alpha(y_2 y_3 A_{q+1}^{12} + a_k^{31}) = e_\alpha(y_2 a_{q+1}^{21} + y_3 a_k^{31}) = \\ e_\alpha((x_3 + y_2 a_{q+1}^{21}) + (x_2 + y_3 a_k^{31})) &= e_\alpha(a_{q+2}^{32} + a_{k+1}^{23}). \quad \square \end{aligned}$$

6) Since $a_{q+2}^{21} = x_2 + y_1 a_{q+1}^{13} \supseteq y_1 a_{q+1}^{13} \supseteq e_\alpha$ (Table C.1, Line 6, a -form), it follows that

$$\begin{aligned} e_\alpha(x_3 + a_{q+2}^{21} A_k^{21}) &= e_\alpha(x_3 a_{q+2}^{21} + A_k^{21}) = e_\alpha(x_3 A_{q+2}^{12} + A_k^{21}) = \\ e_\alpha(y_2 + x_3 A_{q+2}^{12} + A_k^{21}) &= e_\alpha(A_{q+3}^{23} + A_k^{21}). \quad \square \end{aligned}$$

7) By (Table C.1, Line 7, a -form) $a_k^{31} = x_3 + y_1 a_{k-1}^{12} \supseteq y_1 a_{k-1}^{12} \supseteq y_1 a_k^{12} \supseteq e_\alpha$, and therefore

$$\begin{aligned} e_\alpha(x_2 + a_k^{31} A_k^{31}) &= e_\alpha(x_2 a_k^{31} + A_k^{31}) = e_\alpha(x_2 A_k^{13} + A_k^{31}) = \\ e_\alpha(y_3 + x_2 A_k^{13} + A_k^{31}) &= e_\alpha(A_{k+1}^{32} + A_k^{31}). \quad \square \end{aligned}$$

8) By (Table C.1, Line 8, a -form) $a_k^{23} \supseteq a_{k+1}^{23} \supseteq e_\alpha$ and we have

$$\begin{aligned} e_\alpha(x_1 + a_k^{23} A_{k+1}^{23}) &= e_\alpha(x_1 a_k^{23} + A_{k+1}^{23}) = \\ e_\alpha(y_2 + x_1 A_{k+1}^{23} + A_{k+1}^{23}) &= e_\alpha(A_{k+1}^{21} + A_{k+1}^{23}). \quad \square \end{aligned}$$

9) Since $A_{q-1}^{12} \supseteq A_q^{12} \supseteq e_\alpha$, we see that

$$\begin{aligned} e_\alpha(y_2y_3 + A_{q-1}^{12}a_{k+1}^{31}) &= e_\alpha(y_2y_3A_{q-1}^{12} + a_{k+1}^{31}) = e_\alpha(y_2y_3a_{q-1}^{21} + a_{k+1}^{31}) = \\ e_\alpha(x_3 + y_2y_3a_{q-1}^{21} + a_{k+1}^{31}) &= e_\alpha(y_3(x_3 + y_2a_{q-1}^{21}) + a_{k+1}^{31}) = e_\alpha(y_3a_q^{32} + a_{k+1}^{31}) = \\ e_\alpha(x_1 + y_3a_q^{32} + a_{k+1}^{31}) &= e_\alpha(a_{q+1}^{13} + a_{k+1}^{31}). \quad \square \end{aligned}$$

9') Here,

$$e_\alpha(y_1a_k^{12} + y_3a_k^{32}) = e_\alpha((x_3 + y_1a_k^{12}) + (x_1 + y_3a_k^{32})) = e_\alpha(a_{k+1}^{31} + a_{k+1}^{13}). \quad \square$$

10) By (Table C.1, Line 10, a -form) $a_q^{21} \supseteq a_{q+1}^{21} \supseteq e_\alpha$, thus,

$$\begin{aligned} e_\alpha(x_3 + a_q^{21}A_{k+1}^{21}) &= e_\alpha(x_3a_q^{21} + A_{k+1}^{21}) = e_\alpha(x_3A_q^{12} + A_{k+1}^{21}) = \\ e_\alpha(y_2 + x_3A_q^{12} + A_{k+1}^{21}) &= e_\alpha(A_{q+1}^{23} + A_{k+1}^{21}). \quad \square \end{aligned}$$

11) Since $A_q^{31} \supseteq A_{q+1}^{31} \supseteq e_\alpha$, we get

$$\begin{aligned} e_\alpha(y_2y_1 + A_q^{31}a_{k+1}^{23}) &= e_\alpha(y_2y_1A_q^{31} + a_{k+1}^{23}) = e_\alpha(y_2y_1a_q^{13} + a_{k+1}^{23}) = \\ e_\alpha(x_2 + y_2y_1a_q^{13} + a_{k+1}^{23}) &= e_\alpha(y_2(x_2 + y_1a_q^{13}) + a_{k+1}^{23}) = e_\alpha(y_2a_{q+1}^{21} + a_{k+1}^{23}) = \\ e_\alpha(x_3 + y_2a_{q+1}^{21} + a_{k+1}^{23}) &= e_\alpha(a_{q+2}^{32} + a_{k+1}^{23}). \quad \square \end{aligned}$$

12) By (Table C.1, Line 12, a -form) $a_{q+1}^{13} \supseteq a_{q+2}^{13} \supseteq e_\alpha$ and we get

$$\begin{aligned} e_\alpha(x_2 + a_{q+1}^{13}A_{k+1}^{13}) &= e_\alpha(x_2a_{q+1}^{13} + A_{k+1}^{13}) = \\ e_\alpha(y_1 + x_2a_{q+1}^{13} + A_{k+1}^{13}) &= e_\alpha(A_{q+2}^{12} + A_{k+1}^{32}). \quad \square \end{aligned}$$

13) Here,

$$\begin{aligned} e_\alpha(y_3y_1 + A_{q+1}^{23}a_{k+1}^{12}) &= e_\alpha(y_3y_1A_{q+1}^{23} + a_{k+1}^{12}) = e_\alpha(y_3y_1a_{q+1}^{32} + a_{k+1}^{12}) = \\ e_\alpha(x_1 + y_3y_1a_{q+1}^{32} + a_{k+1}^{12}) &= e_\alpha(y_1(x_1 + y_3a_{q+1}^{32}) + a_{k+1}^{12}) = e_\alpha(y_1a_{q+2}^{13} + a_{k+1}^{12}) = \\ e_\alpha((x_2 + y_1a_{q+2}^{13}) + a_{k+1}^{12}) &= e_\alpha(a_{q+3}^{21} + a_{k+1}^{12}). \quad \square \end{aligned}$$

14) By (Table C.1, Line 14, a -form) $a_{q+2}^{32} \supseteq a_{q+3}^{32} \supseteq e_\alpha$ and we have

$$\begin{aligned} e_\alpha(x_1 + a_{q+2}^{32}A_{k+1}^{32}) &= e_\alpha(x_1a_{q+2}^{32} + A_{k+1}^{32}) = \\ e_\alpha(y_3 + x_1a_{q+2}^{32} + A_{k+1}^{32}) &= e_\alpha(y_3 + x_1A_{q+2}^{23} + A_{k+1}^{32}) = e_\alpha(A_{q+3}^{31} + A_{k+1}^{32}). \quad \square \end{aligned}$$

After all previous preparations the proof of Theorem 2.13.1 is rather trivial. Instead of inclusion

$$e_{\alpha i} \subseteq g_{\alpha 0}, \text{ where } i = 1, 2, 3,$$

we will prove a sharper statement

PROPOSITION C.1.7. *For the lattice elements $e_{\alpha i}$ and $g_{\alpha 0}$, we have*

$$e_{\alpha i} = g_{\alpha 0}Z, \quad i = 1, 2, 3,$$

where the element Z for $i = 1$ is given in Table C.4.

Proof of Proposition C.1.7 follows from the S -form of elements $g_{\alpha 0}$ from Table C.3 and the a -form and A -form of the elements $e_{\alpha i}$ from Table C.1. The polynomial $g_{\alpha 0}$ from Table C.3 is the intersection of M and P :

$$g_{\alpha 0} = M \cap P, \tag{C.1}$$

N	$\alpha 1$ index of $e_{\alpha 1}$	α index of $g_{\alpha 0}$	$e_{\alpha 1} = g_{\alpha 0} Z$
1	$(213)^{2p}(21)^{2k}$	$21(321)^{2p-1}(32)^{2k}$	$e_{\alpha 1} = g_{\alpha 0} A_q^{12}$
2	$3(213)^{2p}(21)^{2k}$	$(321)^{2p}(32)^{2k}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+1}^{12}$
3	$13(213)^{2p}(21)^{2k}$	$1(321)^{2p}(32)^{2k}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+1}^{31}$
4	$(213)^{2p+1}(21)^{2k}$	$21(321)^{2p}(32)^{2k}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+2}^{31}$
5	$3(213)^{2p+1}(21)^{2k}$	$(321)^{2p+1}(32)^{2k}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+2}^{23}$
6	$13(213)^{2p+1}(21)^{2k}$	$1(321)^{2p+1}(32)^{2k}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+3}^{23}$
7	$1(21)^{2k}$	$(12)^{2k}$	$e_{\alpha 1} = g_{\alpha 0} A_k^{31}$
8	$1(21)^{2k+1}$	$(12)^{2k+1}$	$e_{\alpha 1} = g_{\alpha 0} A_{k+1}^{23}$
9	$(213)^{2p}(21)^{2k+1}$	$21(321)^{2p-1}(32)^{2k+1}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+1}^{31}$
10	$3(213)^{2p}(21)^{2k+1}$	$(321)^{2p}(32)^{2k+1}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+1}^{23}$
11	$13(213)^{2p}(21)^{2k+1}$	$1(321)^{2p}(32)^{2k+1}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+2}^{23}$
12	$(213)^{2p+1}(21)^{2k+1}$	$21(321)^{2p}(32)^{2k+1}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+2}^{12}$
13	$3(213)^{2p+1}(21)^{2k+1}$	$(321)^{2p+1}(32)^{2k+1}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+3}^{12}$
14	$13(213)^{2p+1}(21)^{2k+1}$	$1(321)^{2p+1}(32)^{2k+1}$	$e_{\alpha 1} = g_{\alpha 0} A_{q+3}^{31}$

TABLE C.4. The relation $e_{\alpha 1} = g_{\alpha 0} Z$

where M is the expression located on the left of the brackets in the S -form of Table C.1 and P is the sum contained in the parantheses of this S -form. Then, for every line from Table C.1, we have

$$MZ = e_{\alpha 1} \text{ and } e_{\alpha 1} \subseteq P. \quad (\text{C.2})$$

From (C.1) and (C.2) it follows that

$$g_{\alpha 0} Z = MPZ = e_{\alpha 1} P = e_{\alpha 1}.$$

For example, consider Lines 1 and 2 of Table C.4.

Line 1. We have

$$\begin{aligned} g_{\alpha 0} &= y_2 A_{k-1}^{32} A_q^{31} A_k^{13} A_{q-1}^{12} (a_q^{21} + a_{k+1}^{12}), \\ M &= y_2 A_{k-1}^{32} A_q^{31} A_k^{13} A_{q-1}^{12}, \quad P = a_q^{21} + a_{k+1}^{12}, \quad Z = A_q^{12}. \end{aligned}$$

Since $A_q^{12} \subseteq A_{q-1}^{12}$, it follows that

$$MZ = y_2 A_{k-1}^{32} A_q^{31} A_k^{13} A_q^{12} = e_{\alpha 1}$$

and by a -form from Table C.1 we obtain $e_{\alpha 1} \subseteq a_q^{21} \subseteq P$. \square

Line 2. We have

$$\begin{aligned} g_{\alpha 0} &= y_3 A_{k-1}^{13} A_q^{23} A_k^{21} A_q^{12} (A_{q+1}^{12} + A_k^{13}), \\ M &= y_3 A_{k-1}^{13} A_q^{23} A_k^{21} A_q^{12}, \quad P = A_{q+1}^{12} + A_k^{13}, \quad Z = A_{q+1}^{12}. \end{aligned}$$

Since $A_{q+1}^{12} \subseteq A_q^{12}$, we see that

$$MZ = y_3 A_{k-1}^{13} A_q^{23} A_k^{21} A_{q+1}^{12} = e_{\alpha 1}.$$

In addition, $e_{\alpha 1} \subseteq A_{q+1}^{12} \subseteq P$. \square

Proposition C.1.7 together with Theorem 2.13.1 (Inclusion Theorem) are proven.

C.2. Properties of $H^+(n)$

C.2.1. The perfectness of $a_i(1)$. Without loss of generality we consider $i = 1$; we have

$$\begin{aligned} \rho(a_1(1)) &= \rho(x_2(1) + x_3(1)) + \rho(y_2(2) + y_3(2)) = \\ &= \rho(x_2 + x_3) + \sum \varphi_i \Phi^+ \rho(y_2(1) + y_3(1)) = \\ &= X_2 + X_3 + \sum \varphi_i \Phi^+ \rho(y_2 + y_3). \end{aligned}$$

(a) Case $\Phi^+ \rho(y_2 + y_3) \neq 0$. In this case, $\rho(a_1(1)) \neq 0$. Since $y_2 + y_3$ is perfect (Proposition 2.9.3), we see that $\Phi^+ \rho(y_2 + y_3) = X_0^1$ and by Corollary 2.4.4 we obtain

$$\begin{aligned} \rho(a_1(1)) &= X_2 + X_3 + \sum \varphi_i X_0^1 = X_2 + X_3 + \sum \varphi_i \Phi^+ \rho(I) = \\ &= X_2 + X_3 + (Y_1 + Y_2)(Y_1 + Y_3)(Y_2 + Y_3). \end{aligned} \quad (C.3)$$

If $X_2 + X_3 = 0$, then

$$\rho(a_1(1)) = (Y_1 + Y_2)(Y_1 + Y_3)(Y_2 + Y_3)$$

and $a_1(1)$ is perfect because $y_i + y_j$ is perfect. If $X_2 + X_3 \neq 0$ then $Y_2 + Y_3 = X_0$, and therefore

$$\begin{aligned} \rho(a_1(1)) &= X_2 + X_3 + (Y_1 + Y_2)(Y_1 + Y_3) = \\ &= (Y_1 + Y_2 + X_3)(Y_1 + Y_3 + X_2) = \\ &= \rho((y_1 + y_2 + x_3)(y_1 + y_3 + x_2)). \end{aligned} \quad (C.4)$$

Thus, $a_1(1)$ is perfect by Corollary 2.9.4.

(b) Case $\Phi^+ \rho(y_2 + y_3) = 0$. Here, we have $Y_2^1 + Y_3^1 = 0$. Then $\Phi^+ \rho$ is of the form

$$\begin{array}{ccccc} X_1^1 & Y_1^1 & X_0^1 & 0 & 0 \\ & & 0 & & \\ & & 0 & & \end{array}$$

For any indecomposable representation $\Phi^+ \rho$, it can be true only for

$$\Phi^+ \rho = \begin{cases} \rho_{x_0} \text{ or} \\ \rho_{y_1} \text{ or} \\ \rho_{x_1}, \end{cases} \quad \text{i.e.,} \quad \rho = \begin{cases} \Phi^{-1} \rho_{x_0} \text{ or} \\ \Phi^{-1} \rho_{y_1} \text{ or} \\ \Phi^{-1} \rho_{x_1}, \end{cases} \quad (C.5)$$

see Table A.1. For these 3 representations, $\rho(x_2 + x_3) = 0$, and therefore $\rho(a_1(1)) = 0$. \square

C.2.2. The perfectness of $b_i(1)$. To prove the *perfectness* of $b_1(1) \supseteq a_1(1)$, it suffices to consider only the representation ρ , for which $\rho(a_1(1)) = 0$, i.e., the case §C.2.1 (b). In this case, ρ satisfies eq. (C.5). Since $b_i(1) = a_i(1) + x_1(2)$, we get

$$\rho(b_1(1)) = \rho(a_1(1) + y_3 y_2) = Y_3 Y_2.$$

Case $\rho = \Phi^{-1} \rho_{x_0}$ (similarly, for $\Phi^{-1} \rho_{y_1}$). In this case, ρ is the direct sum

$$\begin{array}{ccccc} 0 & Y_1 & Y_1 & 0 & 0 \\ & & 0 & & \\ & & 0 & & \end{array} \oplus \begin{array}{ccccc} 0 & 0 & Y_2 & Y_2 & 0 \\ & & Y_2 & & \\ & & 0 & & \end{array}$$

where $\dim Y_1 = \dim Y_2 = \dim Y_3 = 1$, which contradicts the indecomposability of ρ .

Case $\rho = \Phi^{-1}\rho_{x_1}$. Then $\rho(b_1(1)) = Y_3Y_2 = \mathbb{R}^1$, i.e., $\rho(b_1(1)) = X_0$ and $b_1(1)$ is perfect. \square

C.2.3. The perfectness of $c_i(1)$. Since $c_1(1) \supseteq b_1(1)$, we see as above, that it suffices to consider only the case $\rho(b_1(1)) = \rho(a_1(1)) = 0$, i.e., again, ρ satisfies eq. (C.5). Since $c_i(1) = a_i(1) + y_1(2)$, it follows that

$$\rho(c_1(1)) = \rho(a_i(1) + y_2(x_1 + y_3) + y_3(x_1 + y_2)) = Y_2(X_1 + Y_3) + Y_3(X_1 + Y_2).$$

For $\rho = \Phi^{-1}\rho_{x_0}$ and $\rho = \Phi^{-1}\rho_{x_1}$, we have $X_1 = 0$. Then

$$Y_2(X_1 + Y_3) + Y_3(X_1 + Y_2) = Y_2Y_3$$

and we are back to the argumentation of §C.2.2. Let us consider $\rho = \Phi^{-1}\rho_{y_1}$.

Here, $X_1 + Y_2 = X_0$, otherwise $\dim(X_1 + Y_2) = 1$ and $X_1 = Y_2$ and the representation $\Phi^{-1}\rho_{y_1}$ is decomposable. Since $X_1 + Y_2 = X_0$, we get $Y_3(X_1 + Y_2) = Y_3$; by analogy we have $Y_2(X_1 + Y_3) = Y_2$, and hence $\rho(c_1(1)) = Y_3 + Y_2$. Since $y_2 + y_3$ is perfect, it follows that $c_1(1)$ is also a perfect element.

The perfectness of elements $a_i(1), b_i(1), c_i(1)$ for $i = 2, 3$ is similarly proved. \square

C.2.4. The perfectness of $a_i(n), b_i(n), c_i(n)$. Now, we will prove the perfectness of the elements $a_i(1), b_i(1), c_i(1)$.

Proposition 3.1.1 1) For every element $v_i(n) = a_i(n), b_i(n), c_i(n)$, we have

$$\rho(v_i(n)) = \sum_{k=1,2,3} \varphi_k \Phi^+ \rho(v_i(n-1)). \quad (\text{C.6})$$

2) The elements $a_i(n), b_i(n), c_i(n)$ are perfect for every $n \geq 1$ and $i = 1, 2, 3$.

Proof. 1) follows from the definition of the cumulative polynomials from §1.10, Proposition 2.13.3 and the definition of the perfect elements $v_i(n)$ from §A.6. \square

2) For $n = 1$, it was only just proved in §§ C.2.1, C.2.2, C.2.3. We will prove, for example, the perfectness of $a_1(n)$. From eq. (C.6) we have

$$\rho(a_1(n)) = \sum_{i=1,2,3} \varphi_i \Phi^+ \rho(a_1(n-1)). \quad (\text{C.7})$$

By (C.7), Proposition 2.10.1 and relation (2.15) we see that the perfectness of $a_1(n)$ follows from the perfectness of $a_1(n-1)$. \square

C.2.5. The distributivity of $H^+(n)$. Our proof of the following proposition is founded on the well-known result of B. Jonsson [Jo55] (see Proposition A.4.4):

A modular lattice generated by the chains s_1, s_2, s_3 is distributive if and only if every sublattice $[v_1, v_2, v_3]$, where v_i is some element from s_i , is distributive.

Proposition 3.1.2 $H^+(n)$ is distributive sublattices for every $n \geq 0$.

Proof. By Jonsson's criterion it suffices to prove that

$$v_i v_k + v_j v_k = (v_i + v_j) v_k \quad (\text{C.8})$$

for distinct i, j, k . In our case, every two generators from different chains compose the same sum

$$v_i + v_j = I_n, \text{ where } i \neq j. \quad (\text{C.9})$$

For $n \geq 1$, we have

$$I_n = \sum_{i=1,2,3} x_i(n) + \sum_{i=1,2,3} y_i(n+1). \quad (\text{C.10})$$

For $n = 0$, we have

$$I_0 = \sum_{i=1,2,3} y_i. \quad (\text{C.11})$$

By (C.8) and (C.9) it suffices to prove

$$v_i v_k + v_j v_k = v_k \quad (\text{C.12})$$

for distinct i, j, k . We will omit index n in the polynomials $a_i(n), b_i(n), c_i(n)$.

Suppose that

$$a_i a_j + a_k = I_n. \quad (\text{C.13})$$

Then, $b_i a_j + a_k = I_n$ and $c_i a_j + a_k = I_n$. Further, by the modular law (A.3) we have

$$\begin{aligned} a_i &= a_i(a_i a_j + a_k) = a_i a_j + a_i a_k, \\ b_i &= b_i(a_i a_j + a_k) = b_i a_j + b_i a_k, \\ c_i &= c_i(a_i a_j + a_k) = c_i a_j + c_i a_k. \end{aligned} \quad (\text{C.14})$$

Now, if $a_i \subseteq v_i$, where $v_i = b_i$ or c_i , then

$$a_i = a_i a_j + a_i a_k \subseteq a_i v_j + a_i v_k \subseteq a_i(v_j + a_i v_k) \subseteq a_i,$$

i.e., $a_i = a_i v_j + a_i v_k$. The same for b_i, c_i :

$$\begin{aligned} a_i &= a_i v_j + a_i v_k, \\ b_i &= b_i v_j + b_i v_k, \\ c_i &= a_i v_j + c_i v_k. \end{aligned} \quad (\text{C.15})$$

So, (C.15) and (C.12) follow from (C.13). Therefore, it suffices to prove (C.13). For $n = 0$ we have $a_i = y_j + y_k$ and

$$I_n \supseteq a_i a_j + a_k = (y_j + y_k)(y_i + y_k) + y_i + y_j \supseteq y_k + y_i + y_j = I_n.$$

Set

$$t_i(n) = x_i(n) + y_i(n+1). \quad (\text{C.16})$$

Obviously, $a_i(n) = t_j(n) + t_k(n)$. By (C.10), for $n \geq 1$, we have (parameter n is dropped)

$$I_n \supseteq a_i a_j + a_k = (t_j + t_k)(t_i + t_k) + t_i + t_j \supseteq t_k + t_i + t_j = I_n.$$

The distributivity of $H^+(n)$ is proven. \square

C.2.6. The cardinality of $H^+(n)$. Now we will prove that the distributive lattice $H^+(0)$ is a 27-element sublattice in $D^{2,2,2}$ and the distributive lattice $H^+(n)(n \geq 1)$ is the 64-element sublattice in $D^{2,2,2}$.

We introduce *characteristic function* χ_ρ^v to be

$$\chi_\rho^v = \begin{cases} 1 & \text{for } \rho(v) = X_0, \\ 0 & \text{for } \rho(v) = 0. \end{cases} \quad (\text{C.17})$$

Proposition 3.2.3 *The lattice $H^+(0)$ contains 27 distinct elements.*

Proof. It suffices to enumerate all 27 elements in the lattice $H^+(0)$ and find representation which separates each pair of elements. Table C.5 contains values of the characteristic function (C.17) on 6 representations ρ_{y_i}, ρ_{x_i} , where $i = 1, 2, 3$ (see Table A.1), and all 27 elements of $H^+(0)$. All signatures in Table C.5 are distinct. \square

Proposition 3.2.4 *The lattice $H^+(1)$ contains 64 distinct elements.*

As above, for the sublattice $H^+(0)$, it suffices to enumerate all 64 elements in the lattice $H^+(1)$ and find representation which separates each pair of elements. Tables C.6, C.7 contain

values of the characteristic function (C.17) on 9 representations $\rho_{x_i}, \Phi^{-1}\rho_{y_i}, \Phi^{-1}\rho_{x_i}$, where $i = 1, 2, 3$, (see Table A.1) and all 64 elements of $H^+(1)$.

By (1.27) the generators $a_i, b_i, c_i, i = 1, 2, 3$ from the sublattice $H^+(1)$ are:

$$\begin{aligned} a_1(1) &= x_2 + x_3 + y_3(x_2 + y_1) + y_1(x_2 + y_3) + y_1(x_3 + y_2) + y_2(x_3 + y_1), \\ a_2(1) &= x_1 + x_3 + y_3(x_1 + y_2) + y_2(x_1 + y_3) + y_1(x_3 + y_2) + y_2(x_3 + y_1), \\ a_3(1) &= x_2 + x_1 + y_3(x_2 + y_1) + y_1(x_2 + y_3) + y_3(x_1 + y_2) + y_2(x_1 + y_3), \end{aligned} \quad (\text{C.19})$$

$$\begin{aligned} b_1(1) &= a_1(1) + y_2y_3, & c_1(1) &= a_1(1) + y_2(x_1 + y_3) + y_3(x_1 + y_2), \\ b_2(1) &= a_2(1) + y_1y_3, & c_2(1) &= a_2(1) + y_1(x_2 + y_3) + y_3(x_2 + y_1), \\ b_3(1) &= a_3(1) + y_1y_2, & c_3(1) &= a_3(1) + y_1(x_3 + y_2) + y_2(x_3 + y_1), \end{aligned} \quad (\text{C.20})$$

see Table C.6. By (C.18) Lines 10 – 64 of Table C.7 are obtained by multiplying of appropriate Lines 1 – 9 of Table C.6.

No two signatures in Tables C.6 coincide, which was to be proved. \square

C.2.7. A relation between the upper and the lower generators of $H^+(n)$.

Proposition 3.5.1 *Table 3.1 from §3.34 gives relations between the upper and the lower generators. These relations are true mod θ .*

Proof. For definition of $a_i(n), b_i(n), c_i(n)$ (resp. $p_i(n), q_i(n), s_i(n)$), see (1.27) (resp. (3.34)).

1) By (3.24) we have $x_0(n+2) \subseteq a_i(n)$ and

$$\begin{aligned} s_j(n) + s_k(n) &= \\ x_j(n) + y_j(n+1) + x_k(n) + y_k(n+1) + x_0(n+2) &= \\ a_i(n) + x_0(n+2) &= a_i(n). \end{aligned}$$

2) Here, we have

$$\begin{aligned} s_j(n) + s_k(n) + p_i(n) &= \\ a_i(n) + p_i(n) &= \\ x_j(n) + y_j(n+1) + x_k(n) + y_k(n+1) + x_0(n+2) + x_i(n+1) &= \\ a_i(n) + x_i(n+1) &= b_i(n). \end{aligned}$$

3) Further,

$$\begin{aligned} s_j(n) + s_k(n) + q_i(n) &= a_i(n) + q_i(n) = \\ x_j(n) + y_j(n+1) + x_k(n) + y_k(n+1) + x_0(n+2) + y_i(n+1) &= \\ a_i(n) + y_i(n+1) &= c_i(n). \end{aligned}$$

4) Consider the intersection $a_k(n)a_j(n)$:

$$\begin{aligned} a_k(n)a_j(n) &= \\ [x_i(n) + y_i(n+1) + x_j(n) + y_j(n+1)][x_i(n) + y_i(n+1) + x_k(n) + y_k(n+1)] &= \\ x_i(n) + y_i(n+1) + (x_j(n) + y_j(n+1))a_j(n). \end{aligned}$$

Since $x_j(n) + y_j(n+1) \subseteq a_i(n)a_k(n)$, we see that

$$\begin{aligned} a_k(n)a_j(n) &\subseteq x_i(n) + y_i(n+1) + a_i(n)a_j(n)a_k(n) = \\ x_i(n) + y_i(n+1) + x_0(n+2) &= s_i(n). \end{aligned}$$

On the other hand,

$$s_i(n) = x_i(n) + y_i(n+1) + x_0(n+2) \subseteq a_j(n)a_k(n).$$

Therefore,

$$s_i(n) = a_j(n)a_k(n).$$

5) Since $x_i(n+1) \subseteq a_j(n)$ and $x_j(n+1) \subseteq a_i(n)$, by heading 4) we have

$$\begin{aligned} b_i(n)b_j(n) &= (a_i(n) + x_i(n+1))(a_j(n) + x_j(n+1)) = \\ &= x_i(n+1) + x_j(n+1) + a_i(n)a_j(n) = \\ &= x_i(n+1) + x_j(n+1) + a_i(n)a_j(n) + x_0(n+2) = p_i(n) + p_j(n) + s_k(n). \end{aligned}$$

6) Since $y_i(n+1) \subseteq a_j(n)$ and $y_j(n+1) \subseteq a_i(n)$, by heading 4) we get

$$\begin{aligned} c_i(n)c_j(n) &= (a_i(n) + y_i(n+1))(a_j(n) + y_j(n+1)) = \\ &= y_i(n+1) + y_j(n+1) + a_i(n)a_j(n) = \\ &= y_i(n+1) + y_j(n+1) + a_i(n)a_j(n) + x_0(n+2) = q_i(n) + q_j(n) + s_k(n). \end{aligned}$$

7) Since $x_i(n+1) \subseteq b_j(n)$, by heading 4) we see that

$$\begin{aligned} a_i(n)b_j(n) &= a_i(n)(a_j(n) + x_j(n+1)) = x_j(n+1) + a_i(n)a_j(n) = \\ &= x_j(n+1) + a_i(n)a_j(n) + x_0(n+2) = p_j(n) + s_k(n). \end{aligned}$$

8) Since $y_i(n+1) \subseteq c_j(n)$, by heading 4) we have

$$\begin{aligned} a_i(n)c_j(n) &= a_i(n)(a_j(n) + y_j(n+1)) = y_j(n+1) + a_i(n)a_j(n) = \\ &= y_j(n+1) + a_i(n)a_j(n) + x_0(n+2) = q_j(n) + s_k(n). \end{aligned}$$

9) Since $x_i(n+1) \subseteq c_j(n)$ and $y_j(n+1) \subseteq b_i(n)$, by heading 4) we have

$$\begin{aligned} b_i(n)c_j(n) &= (a_i(n) + x_i(n+1))(a_j(n) + y_j(n+1)) = \\ &= x_i(n+1) + y_j(n+1) + a_i(n)a_j(n) = \\ &= x_i(n+1) + y_j(n+1) + a_i(n)a_j(n) + x_0(n+2) = p_j(n) + q_j(n) + s_k(n). \end{aligned}$$

10) Since $x_k(n+1) \subseteq a_i(n)a_j(n)$, we see that

$$\begin{aligned} a_i(n)a_j(n)b_k(n) &= a_i(n)a_j(n)(a_k(n) + x_k(n+1)) = \\ &= x_k(n+1) + a_i(n)a_j(n)a_k(n) = x_k(n+1) + x_0(n+2) = p_k(n). \end{aligned}$$

11) Since $y_k(n+1) \subseteq a_i(n)a_j(n)$, we have

$$\begin{aligned} a_i(n)a_j(n)c_k(n) &= a_i(n)a_j(n)(a_k(n) + y_k(n+1)) = \\ &= y_k(n+1) + a_i(n)a_j(n)a_k(n) = y_k(n+1) + x_0(n+2) = q_k(n). \end{aligned}$$

12) Here we have

$$\begin{aligned} b_i(n)b_j(n)a_k(n) &= (a_i(n) + x_i(n+1))(a_j(n) + x_j(n+1))a_k(n) = \\ &= x_i(n+1) + x_j(n+1) + a_i(n)a_j(n)a_k(n) = \\ &= x_i(n+1) + x_j(n+1) + x_0(n+2) = p_i(n) + p_j(n). \end{aligned}$$

13) Further,

$$\begin{aligned} b_i(n)b_j(n)c_k(n) &= (a_i(n) + x_i(n+1))(a_j(n) + x_j(n+1))(a_k(n) + y_k(n+1)) = \\ &= x_i(n+1) + x_j(n+1) + y_k(n+1) + a_i(n)a_j(n)a_k(n) = \\ &= x_i(n+1) + x_j(n+1) + y_k(n+1) + x_0(n+2) = p_i(n) + p_j(n) + q_k(n). \end{aligned}$$

14) As above,

$$\begin{aligned} c_i(n)c_j(n)a_k(n) &= (a_i(n) + y_i(n+1))(a_j(n) + y_j(n+1))a_k(n) = \\ y_i(n+1) + y_j(n+1) + a_i(n)a_j(n)a_k(n) &= y_i(n+1) + y_j(n+1) + x_0(n+2) = \\ q_i(n) + q_j(n). \end{aligned}$$

15) Similarly, we have

$$\begin{aligned} c_i(n)c_j(n)b_k(n) &= (a_i(n) + y_i(n+1))(a_j(n) + y_j(n+1))(a_k(n) + x_k(n+1)) = \\ y_i(n+1) + y_j(n+1) + x_k(n+1) + a_i(n)a_j(n)a_k(n) &= \\ y_i(n+1) + y_j(n+1) + x_k(n+1) + x_0(n+2) &= q_i(n) + q_j(n) + p_k(n). \end{aligned}$$

16) Similarly,

$$\begin{aligned} a_i(n)b_j(n)c_k(n) &= a_i(n)(a_j(n) + x_j(n+1))(a_k(n) + y_k(n+1)) = \\ x_j(n+1) + y_k(n+1) + a_i(n)a_j(n)a_k(n) &= \\ x_j(n+1) + y_k(n+1) + x_0(n+2) &= p_j(n) + q_k(n). \end{aligned}$$

17) It is just (3.24):

$$a_i(n)a_j(n)a_k(n) = x_0(n+2).$$

18) Here we have

$$\begin{aligned} b_i(n)b_j(n)b_k(n) &= (a_i(n) + x_i(n+1))(a_j(n) + x_j(n+1))(a_k(n) + x_k(n+1)) = \\ x_i(n+1) + x_j(n+1) + x_k(n+1) + a_i(n)a_j(n)a_k(n) &= \\ x_i(n+1) + x_j(n+1) + x_k(n+1) + x_0(n+2) &= p_i(n) + p_j(n) + p_k(n). \end{aligned}$$

19) Similarly,

$$\begin{aligned} c_i(n)c_j(n)c_k(n) &= (a_i(n) + y_i(n+1))(a_j(n) + y_j(n+1))(a_k(n) + y_k(n+1)) = \\ y_i(n+1) + y_j(n+1) + y_k(n+1) + a_i(n)a_j(n)a_k(n) &= \\ y_i(n+1) + y_j(n+1) + y_k(n+1) + x_0(n+2) &= q_i(n) + q_j(n) + q_k(n). \end{aligned}$$

20) As follows from Line 1) of this table,

$$a_i(n) + a_j(n) = (s_j(n) + s_k(n)) + (s_i(n) + s_k(n)) = s_i(n) + s_j(n) + s_k(n).$$

Proposition 3.5.1 is proven. \square

N	Polynomial	ρ_{y_1}	ρ_{y_2}	ρ_{y_3}	ρ_{x_1}	ρ_{x_2}	ρ_{x_3}	(Zeros, Units)
1	$a_1(0) = y_2 + y_3$	0	1	1	0	1	1	(2,4)
2	$a_2(0) = y_1 + y_3$	1	0	1	1	0	1	(2,4)
3	$a_3(0) = y_2 + y_1$	1	1	0	1	1	0	(2,4)
4	$b_1(0) = x_1 + y_2 + y_3$	0	1	1	1	1	1	(1,5)
5	$b_2(0) = x_2 + y_1 + y_3$	1	0	1	1	1	1	(1,5)
6	$b_3(0) = x_3 + y_2 + y_1$	1	1	0	1	1	1	(1,5)
7	$a_1(0)a_2(0)$	0	0	1	0	0	1	(4,2)
8	$a_1(0)a_3(0)$	0	1	0	0	1	0	(4,2)
9	$a_2(0)a_3(0)$	1	0	0	1	0	0	(4,2)
10	$b_1(0)b_2(0)$	0	0	1	1	1	1	(2,4)
11	$b_1(0)b_3(0)$	0	1	0	1	1	1	(2,4)
12	$b_2(0)b_3(0)$	1	0	0	1	1	1	(2,4)
13	$a_1(0)b_2(0)$	0	0	1	0	1	1	(3,3)
14	$a_1(0)b_3(0)$	0	1	0	0	1	1	(3,3)
15	$a_2(0)b_1(0)$	0	0	1	1	0	1	(3,3)
16	$a_2(0)b_3(0)$	1	0	0	1	0	1	(3,3)
17	$a_3(0)b_1(0)$	0	1	0	1	1	0	(3,3)
18	$a_3(0)b_2(0)$	1	0	0	1	1	0	(3,3)
19	$a_1(0)a_2(0)b_3(0)$	0	0	0	0	0	1	(5,1)
20	$a_1(0)b_2(0)a_3(0)$	0	0	0	0	1	0	(5,1)
21	$b_1(0)a_2(0)a_3(0)$	0	0	0	1	0	0	(5,1)
22	$a_1(0)b_2(0)b_3(0)$	0	0	0	0	1	1	(4,2)
23	$b_1(0)a_2(0)b_3(0)$	0	0	0	1	0	1	(4,2)
24	$b_1(0)b_2(0)a_3(0)$	0	0	0	1	1	0	(4,2)
25	$a_1(0)a_2(0)a_3(0)$	0	0	0	0	0	0	(6,0)
26	$b_1(0)b_2(0)b_3(0)$	0	0	0	1	1	1	(3,3)
27	$I_0 = y_1 + y_2 + y_3$	1	1	1	1	1	1	(6,0)

TABLE C.5. The characteristic function on the $H^+(0)$

Notes: (a) Lines 7 – 27 are obtained by multiplying the appropriate Lines 1 – 6, since

$$\chi_\rho^{v_1 v_2} = \chi_\rho^{v_1} \chi_\rho^{v_2}. \quad (\text{C.18})$$

(b) We call a sequence of “zeros” and “units” in the set $\rho_{y_1}, \rho_{y_2}, \dots, \rho_{x_3}$ the *signature* of the corresponding polynomial.

N	Poly- nomial	ρ_{x_1}	ρ_{x_2}	ρ_{x_3}	$\Phi^{-1}\rho_{y_1}$	$\Phi^{-1}\rho_{y_2}$	$\Phi^{-1}\rho_{y_3}$	$\Phi^{-1}\rho_{x_1}$	$\Phi^{-1}\rho_{x_2}$	$\Phi^{-1}\rho_{x_3}$	(Zeros, Units)
1	$a_1(1)$	0	1	1	0	1	1	0	1	1	(3,6)
2	$a_2(1)$	1	0	1	1	0	1	1	0	1	(3,6)
3	$a_3(1)$	1	1	0	1	1	0	1	1	0	(3,6)
4	$b_1(1)$	0	1	1	0	1	1	1	1	1	(2,7)
5	$b_2(1)$	1	0	1	1	0	1	1	1	1	(2,7)
6	$b_3(1)$	1	1	0	1	1	0	1	1	1	(2,7)
7	$c_1(1)$	0	1	1	1	1	1	1	1	1	(1,8)
8	$c_2(1)$	1	0	1	1	1	1	1	1	1	(1,8)
9	$c_3(1)$	1	1	0	1	1	1	1	1	1	(1,8)

TABLE C.6. The characteristic function on the $H^+(1)$. The generators $a_i(1), b_i(1), c_i(1)$

N	Poly- nomial	ρ_{x_1}	ρ_{x_2}	ρ_{x_3}	$\Phi^{-1}\rho_{y_1}$	$\Phi^{-1}\rho_{y_2}$	$\Phi^{-1}\rho_{y_3}$	$\Phi^{-1}\rho_{x_1}$	$\Phi^{-1}\rho_{x_2}$	$\Phi^{-1}\rho_{x_3}$	(Zeros, Units)
10	$a_1(1)a_2(1)$	0	0	1	0	0	1	0	0	1	(6,3)
11	$a_1(1)a_3(1)$	0	1	0	0	1	0	0	1	0	(6,3)
12	$a_2(1)a_3(1)$	1	0	0	1	0	0	1	0	0	(6,3)
13	$b_1(1)b_2(1)$	0	0	1	0	0	1	1	1	1	(4,5)
14	$b_1(1)b_3(1)$	0	1	0	0	1	0	1	1	1	(4,5)
15	$b_2(1)b_3(1)$	1	0	0	1	0	0	1	1	1	(4,5)
16	$c_1(1)c_2(1)$	0	0	1	1	1	1	1	1	1	(2,7)
17	$c_1(1)c_3(1)$	0	1	0	1	1	1	1	1	1	(2,7)
18	$c_2(1)c_3(1)$	1	0	0	1	1	1	1	1	1	(2,7)
19	$a_1(1)b_2(1)$	0	0	1	0	0	1	0	1	1	(5,4)
20	$a_1(1)b_3(1)$	0	1	0	0	1	0	0	1	1	(5,4)
21	$a_2(1)b_1(1)$	0	0	1	0	0	1	1	0	1	(5,4)
22	$a_2(1)b_3(1)$	1	0	0	1	0	0	1	0	1	(5,4)
23	$a_3(1)b_1(1)$	0	1	0	0	1	0	1	1	0	(5,4)
24	$a_3(1)b_2(1)$	1	0	0	1	0	0	1	1	0	(5,4)
25	$a_1(1)c_2(1)$	0	0	1	0	1	1	0	1	1	(4,5)
26	$a_1(1)c_3(1)$	0	1	0	0	1	1	0	1	1	(4,5)
27	$a_2(1)c_1(1)$	0	0	1	1	0	1	1	0	1	(4,5)
28	$a_2(1)c_3(1)$	1	0	0	1	0	1	1	0	1	(4,5)
29	$a_3(1)c_1(1)$	0	1	0	1	1	0	1	1	0	(4,5)
30	$a_3(1)c_2(1)$	1	0	0	1	1	0	1	1	0	(4,5)
31	$b_1(1)c_2(1)$	0	0	1	0	1	1	1	1	1	(3,6)
32	$b_1(1)c_3(1)$	0	1	0	0	1	1	1	1	1	(3,6)
33	$b_2(1)c_1(1)$	0	0	1	1	0	1	1	1	1	(3,6)
34	$b_2(1)c_3(1)$	1	0	0	1	0	1	1	1	1	(3,6)
35	$b_3(1)c_1(1)$	0	1	0	1	1	0	1	1	1	(3,6)
36	$b_3(1)c_2(1)$	1	0	0	1	1	0	1	1	1	(3,6)
37	$a_1(1)a_2(1)b_3(1)$	0	0	0	0	0	0	0	0	1	(8,1)
38	$a_1(1)a_3(1)b_2(1)$	0	0	0	0	0	0	0	1	0	(8,1)
39	$a_2(1)a_3(1)b_1(1)$	0	0	0	0	0	0	1	0	0	(8,1)
40	$a_1(1)a_2(1)c_3(1)$	0	0	0	0	0	1	0	0	1	(7,2)
41	$a_1(1)a_3(1)c_2(1)$	0	0	0	0	1	0	0	1	0	(7,2)
42	$a_2(1)a_3(1)c_1(1)$	0	0	0	1	0	0	1	0	0	(7,2)
43	$b_1(1)b_2(1)a_3(1)$	0	0	0	0	0	0	1	1	0	(7,2)
44	$b_1(1)b_3(1)a_2(1)$	0	0	0	0	0	0	1	0	1	(7,2)
45	$b_2(1)b_3(1)a_1(1)$	0	0	0	0	0	0	0	1	1	(7,2)
46	$b_1(1)b_2(1)c_3(1)$	0	0	0	0	0	1	1	1	1	(5,4)
47	$b_1(1)b_3(1)c_2(1)$	0	0	0	0	1	0	1	1	1	(5,4)
48	$b_2(1)b_3(1)c_1(1)$	0	0	0	1	0	0	1	1	1	(5,4)
49	$c_1(1)c_2(1)a_3(1)$	0	0	0	1	1	0	1	1	0	(5,4)
50	$c_1(1)c_3(1)a_2(1)$	0	0	0	1	0	1	1	0	1	(5,4)
51	$c_2(1)c_3(1)a_1(1)$	0	0	0	0	1	1	0	1	1	(5,4)
52	$c_1(1)c_2(1)b_3(1)$	0	0	0	1	1	0	1	1	1	(4,5)
53	$c_1(1)c_3(1)b_2(1)$	0	0	0	1	0	1	1	1	1	(4,5)
54	$c_2(1)c_3(1)b_1(1)$	0	0	0	0	1	1	1	1	1	(4,5)
55	$a_1(1)b_2(1)c_3(1)$	0	0	1	0	0	1	0	1	1	(7,2)
56	$a_1(1)b_3(1)c_2(1)$	0	0	0	0	1	0	0	1	1	(7,2)
57	$a_2(1)b_1(1)c_3(1)$	0	0	0	0	0	1	1	0	1	(7,2)
58	$a_2(1)b_3(1)c_1(1)$	0	0	0	1	0	0	1	0	1	(7,2)
59	$a_3(1)b_1(1)c_2(1)$	0	0	0	0	1	0	1	1	0	(3,6)
60	$a_3(1)b_2(1)c_1(1)$	0	0	0	1	0	0	1	1	0	(3,6)
61	$a_1(1)a_2(1)a_3(1)$	0	0	0	0	0	0	0	0	0	(9,0)
62	$b_1(1)b_2(1)b_3(1)$	0	0	0	0	0	0	1	1	1	(6,3)
63	$c_1(1)c_2(1)c_3(1)$	0	0	0	1	1	1	1	1	1	(3,6)
64	I_1	1	1	1	1	1	1	1	1	1	(1,8)

TABLE C.7. The characteristic function on the $H^+(1)$. The elements 10–64

Conjectures

1. **Conjecture 1.13.2** *The lattice $H^+ \cup H^-$ contains all perfect elements of $D^{2,2,2} \bmod \theta$.*

For D^4 , the similar conjecture (due to Gelfand-Ponomarev) was proved by Dlab and Ringel in [DR80] and Cylke in [Cyl82], see §1.3.

2. **Conjecture 3.4.4** *The relation*

$$x_0(n+2) = \bigcap_{i=1,2,3} a_i(n) \quad \bmod \theta,$$

see (3.24, Proposition 3.4.2) takes place without restriction $\bmod \theta$.

Here, $x_0(n+2)$ is the cumulative polynomial defined in §1.10, and the $a_i(n)$ are perfect elements defined in §1.27. The elements $x_0(n+2)$ are also perfect, see Corollary 3.4.3.

3. **Conjecture 3.9.2** *The relation*

$$a_i(n)x_i(n+1) \subseteq y_i(n+2) \quad \bmod \theta,$$

(see (3.51)), takes place without restriction $\bmod \theta$.

Here, $x_i(n+1), y_i(n+2)$ are the cumulative polynomial defined in §1.10, and the $a_i(n)$ are perfect elements defined in §1.27.

4. **Conjecture 4.8.6** *For every admissible sequence α , the elements e_α (resp. $f_{\alpha 0}$) and \tilde{e}_α (resp. $\tilde{f}_{\alpha 0}$) coincide without restriction $\bmod \theta$ (see Proposition 4.8.3).*

Here, the elements e_α (resp. $f_{\alpha 0}$) given by Table 4.3, and the elements \tilde{e}_α (resp. $\tilde{f}_{\alpha 0}$) determined by Gelfand and Ponomarev, see §4.8.2. For small admissible sequences, this coincidence is proved in Propositions 4.8.4, 4.8.5.

Bibliography

- [Ber87] Berezin F. A., *Introduction to superanalysis*. Edited and with a foreword by A. A. Kirillov. With an appendix by V. I. Ogievetsky. Translated from the Russian by J. Niederle and R. Kotecký. Translation edited by Dimitri Leites. D. Reidel Publishing Co., Dordrecht, 1987. xii+424 pp.
- [BGP73] Bernstein I. N., Gelfand I. M., Ponomarev V. A., *Coxeter functors and Gabriel's theorem*. Uspehi Mat. Nauk 28 (1973), no. 2(170), 19–33. English translation: Russian Math. Surveys, 28(1973), 17–32.
- [BBR85] Barnabei M., Brini A., and Rota G.-C., *On the exterior calculus of invariant theory*. J. Algebra 96 (1985), 120–160.
- [BvN36] Birkhoff G., von Neuman J., *The logic of quantum mechanics*. Annals of Mathematics, 37 (1936), no. 4, 823–843.
- [Bir48] Birkhoff G. *Lattice theory*. New York, 1948.
- [Bz2001] Baez J., *This Week's Finds in Mathematical Physics*.
<http://math.ucr.edu/home/baez/week173.html> , or
<http://www.lns.cornell.edu/spr/2001-11/msg0037095.html> , 2001.
- [BS81] Burris S., Sankappanavar H. P. *A Course in Universal Algebra*. Graduate Texts in Mathematics, 78. Springer-Verlag, New York-Berlin, 1981. xvi+276
<http://www.math.uwaterloo.ca/~snburris/htdocs/ualg.html>
- [Cyl82] Cylke A. A., *Perfect elements in free modular lattices*. Funktsional. Anal. i Prilozhen. 16 (1982) no. 1, 87–88. English translation: Funct. Anal. Appl. 16, No.1, 73–74 (1982).
- [Cyl93] Cylke A. A., *Perfect and linearly equivalent elements of modular lattices*. CMS Conference Proceedings, (1993), vol. 14, 125–148.
- [De1897] Dedekind R., *Über die von drei Moduln erzeugte Dualgruppe*. Math. Ann., 53(1897), 371–403.
- [DRS76] Doubilet P., Rota G.-C., Stein J., *On the foundation of combinatorial theory. IX. Combinatorial methods in invariant theory*. Stud. Appl. Math. 53 (1976), 185–216.
- [DR74] Dlab V., Ringel C. M. *Representations of graphs and algebras*. Carleton Mathematical Lecture Notes, No. 8. Department of Mathematics, Carleton University, Ottawa, Ont., (1974). iii+86 pp.
- [DR76] Dlab V., Ringel C. M., *Indecomposable representations of graphs and algebras*. Mem. Amer. Math. Soc. 173 (1976).
- [DR79] Dlab V., Ringel C. M., *The preprojective algebra of a modulated graph*. Proceedings 2nd Int. Conf., Ottawa, (1979), Lect. Notes in Math. 832 (1980), 216–231.
- [DR80] Dlab V., Ringel C. M., *Perfect elements in the free modular lattices*. Math. Ann. 247 (1980), 95–100.
- [Dol2003] Dolgachev I., *Lectures on Invariant Theory*. London Math. Soc. LN Series 296 (2003) University of Michigan.
- [FMR96] Finberg D., Mainetti M., Rota G.-C., *The logic of commuting equivalence relations*. Logic and algebra (Pontignano, 1994), 69–96, Lecture Notes in Pure and Appl. Math., 180, Dekker, New York, (1996).
- [Fr80] Freese R., *Free modular lattices*. Trans. Amer. Math. Soc. 261 (1980), no. 1, 81–91.
- [Gab72] Gabriel P., *Unzerlegbare Darstellungen I*, *Manuscripta Math.* 6 (1972), 71–103.
- [GP70] Gelfand I. M., Ponomarev V. A., *Problems of linear algebra and classification of quadruples of subspaces in a finite-dimensional vector space*. Hilbert space operators and operator algebras (Proc. Internat. Conf., Tihany, 1970), pp. 163–237. Colloq. Math. Soc. Janos Bolyai, 5, North-Holland, Amsterdam, 1972.
- [GP74] Gelfand I. M., Ponomarev V. A., *Free modular lattices and their representations*. Collection of articles dedicated to the memory of Ivan Georgievic Petrovskii (1901-1973), IV. Uspehi Mat. Nauk 29 (1974), no. 6(180), 3–58. English translation: Russian Math. Surveys 29 (1974), no. 6, 1–56.

- [GP76] Gelfand I. M., Ponomarev V. A., *Lattices, representations, and their related algebras*. I. Uspehi Mat. Nauk 31 (1976), no. 5(191), 71–88. English translation: Russian Math. Surveys 31 (1976), no. 5, 67–85.
- [GP77] Gelfand I. M., Ponomarev V. A., *Lattices, representations, and their related algebras*. II. Uspehi Mat. Nauk 32 (1977), no. 1(193), 85–106. English translation: Russian Math. Surveys 32 (1977), no. 1, 91–114.
- [GP79] Gelfand I. M., Ponomarev V. A., *Model algebras and representations of graphs*. Funkts. Anal. i Prilozhen. 13 (1979) no.3, 1–12, English translation: Funct. Anal. Appl. 13 (1980), no.3, 157–166.
- [Gra11] Grassmann H. *Gesammelte Mathematische und Physikalische Werke. Ersten Bandes. Erster Theil: Die Ausdehnungslehre von 1844 und die geometrische Analyse* (German). [Collected mathematical and physical works. First volume. First Theil: The geometric calculus of 1844 and the geometrical analysis]. Teubner, Leipzig, 1911.
- [Gr98] Grätzer G. *General Lattice Theory*, 2nd ed. Boston, MA: Birkhäuser, (1998).
- [GR92] Gabriel P., Roiter A. V., *Representations of Finite-Dimensional Algebras*. Enc. of Math. Sci., 73, Algebra VIII, (Springer, 1992).
- [Gro98] Grosshans F. D. *Invariant theory, orbits and non-decomposable quadruples of subspaces having non-zero defect*. Linear Algebra Appl. 282 (1998), no. 1-3, 63–81.
- [Gre67] Greub W. H. *Multilinear algebra*. Die Grundlehren der Mathematischen Wissenschaften, Band 136 Springer-Verlag, New York, Inc., New York 1967 x+225 pp.
- [Hai85] Haiman M., *Proof theory for linear lattices*. Adv. Math. 58 (1985), 209–242.
- [Hai91] Haiman M., *Arguesian lattices which are not type-1*. Algebra Universalis. 28 (1991), 128–137.
- [Hal63] Halmos P. R., *Lectures on Boolean Algebras*, Van Nostrand Co., Princeton, NJ, (1963).
- [Haw94] Hawrylycz M. J., *Geometric Identities in Invariant Theory*. Ph.D. thesis, Massachusetts Institute of Technology, 1994.
- [Haw96] Hawrylycz M. J., *Arguesian identities in invariant theory*. Adv. Math. 122, No. 1 (1996), 1–48.
- [H83] Herrmann C., *On the word problem for the modular lattice with four free generators*. Math. Ann. 265(1983), 513–527.
- [HH96] Howe R., Huang R. *Projective invariants of four subspaces*. Adv. Math. 118 (1996), no. 2, 295–336.
- [Jo53] Jónsson B., *On the representation of lattices*. Math. Scand. 1 (1953), 193–206.
- [Jo54] Jónsson B., *Modular lattices and Desargues’ theorem*. Math. Scand. 2 (1954), 295–314.
- [Jo55] Jónsson B., *Distributive sublattices of a modular lattices*. Proc. Amer. Math. Soc., 6(1955), 682–688.
- [KS67] Kochen S., and Specker E. P. (1967). *The problem of hidden variables in quantum mechanics*. Journal of Mathematics and Mechanics, 17, 59–87.
- [KY2003] Kung J., Yan C. H., *Six Problems of Gian-Carlo Rota in Lattice Theory and Universal Algebra*. Algebra Universalis, vol. 49(2003), 113–127.
- [L05] Bernstein J., Leites D., Shander V., *Seminar on Supersymmetries. vol. 1-1. Algebra and Calculus*, ed. by Leites D., 2005, to appear.
- [Man79] Manin Yu. I., *Provable and unprovable*. (Dokazuemoe i nedokazuemoe.) (Russian) Sovetskoe radio, (1979) (in Russian), 168 p.
- [MY99] Mainetti M., Yan C. H., *Arguesian Identities in Linear Lattices, Advances in Mathematics*. vol. 144, (1999), 50–93.
- [MY2000] Mainetti M., Yan C. H., *Geometric identities in lattice theory*. In memory of Gian-Carlo Rota. J. Combin. Theory Ser. A 91 (2000), no. 1-2, 411–450.
- [Naz67] Nazarova L. A. *Representations of quadruples*. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 31 (1967) 1361–1378.
- [Naz73] Nazarova L. A., *Representations of quivers of infinite type*. Izv. AN SSSR, (1973), 37, N 4. p. 752–791.
- [NR72] Nazarova L. A., Roiter A. V., *Representations of partially ordered sets*. Zap. Nauchn. Semin, Leningr. Otd. Mat. Inst. Steklova, 1972, 28, p. 5–31 (English transl. J. Sov. Math., 1975., 3., p. 585–606).
- [Pon90] Ponomarev V. A., *Equipped graphs and modular lattices attached to them*. Topics in Algebra, Banach Center Publications, vol 26., part 1, Warsaw (1990), 385–405.
- [Rin80] Ringel C. M., *The rational invariants of the tame quivers*. Invent. Math. 58 (1980), no. 3, 217–239.
- [Rin82] Ringel C. M., *Four papers on linear algebras. Introduction to: Representation theory*. Selected papers by I.M. Gelfand et al. London Math. Soc. LN Series 69 (1982), 141–156.
- [Rin96] Ringel C. M., *The preprojective algebra of a quiver. Algebras and modules*. II (Geiranger, 1996), 467–480, CMS Conf. Proc., 24, Amer. Math. Soc., Providence, RI, 1998.

- [Ro85] Roiter A. V., *Representations of posets and tame matrix problems*. Proc. Durham Simp.(1985) - Cambridge, (1986), 91–109.
- [R97] Rota G-C., *The many lives of lattice theory*. Notices Amer. Math. Soc. 44 (11) (1997), 1440–1445.
- [Sch45] Schützenberger M. P. *Sur certains axiomes de la théorie des structures*. C.R. Acad. Sci. Paris 221 (1945), 218–220.
- [Sik64] Sikorski R., *Boolean Algebras*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 25, Springer-Verlag, Berlin, 1964.
- [St84] Stekolshchik R. B., *Invariant elements in modular lattice*. Functional analysis and its applications, 18 (1984) no.1, 82–83. English translation: Funct.Anal. Appl. 18 (1984), no. 1, 73–75.
- [St89] Stekolshchik R. B., *Perfect elements in modular lattice, associated with extended Dynkin diagram \tilde{E}_6* . Functional analysis and its applications, 23 (1989) no.3, 90–92. English translation: Funct.Anal. Appl. 3(1989), p.251–254.
- [St92] Stekolshchik R. B., *Invariant elements in modular lattice, associated with extended Dynkin diagram \tilde{E}_6* . Functional analysis and its applications, 26 (1992) no.1, 42–53. English translation: Funct.Anal. Appl.(1992), p.33–42.
- [Stn36] Stone M. H., *The theory of representation for Boolean algebras*. Trans. Amer. Math. Soc. 40 (1936), no. 1, 37–111.
- [Weyl46] Weyl H., *The classical groups, their invariants and representations*. Princeton Univ. Press, Princeton, N.J., 1946.
- [W73] Wille R. *Über modulare Verbände, die von einer endlichen halbgeordneten Menge frei erzeugt werden*. (German) Math. Z. 131 (1973), 241–249.
- [Wh91] White L., *Multilinear Cayley factorization*. Invariant-theoretic algorithms in geometry (Minneapolis, MN, 1987). J. Symbolic Comput. 11 (1991), no. 5-6, 421–438.