THE INVERSE OF A REAL GENERALIZED CARTAN MATRIX

HECHUN ZHANG

Dedicated to Professor T. H. Koornwinder on the occasion of his 60th birthday

ABSTRACT. The purpose of this note is to give explicit criteria to determine whether a real generalized Cartan matrix is of finite type, affine type or of hyperbolic type by considering the principal minors and the inverse of the matrix. In particular, it will be shown that a real generalized Cartan matrix is of finite type if and only if it is invertible and the inverse is a positive matrix. A real generalized Cartan matrix is of hyperbolic type if and only if it is invertible and the inverse is non-positive.

AMS Classification: 15A09, 15A48, 17B67.

Keywords: Real generalized Cartan matrix, finite type, affine type, indefinite type.

1. Introduction

For any complex $n \times n$ matrix A, one can associate with it a complex Lie algebra $\mathfrak{g}(A)$, see [1] or [5]. The properties of A determine the structure of the Lie algebra $\mathfrak{g}(A)$. In [1], some results are given for generalized Cartan matrices which are integral matrices. In particular, it was proved in [1] that any generalized Cartan matrix of finite type or affine type is symmetrizable. However, for any real generalized Cartan matrix, this is not true any more, see the example in the next section. Hence, we need to develop a new approach to deal with real generalized Cartan matrices. In this paper, we generalize some results in [1] to arbitrary real generalized Cartan matrices. Moreover, we give some give explicit criteria to judge whether a real generalized Cartan matrix is of finite type or of hyperbolic type. In particular, it will be proved that a real generalized Cartan matrix is of finite type if and only if it is invertible and the inverse is a positive matrix. A generalized Cartan matrix is of hyperbolic type if and only if it is invertible and the inverse is non-positive. Furthermore, we obtain a similar result for real generalized Cartan matrices of affine type. Some of the results are even new for generalized Cartan matrices.

This work was done during the author's visit at the University of Amsterdam. The author would like to thank Professor T. H. Koornwinder for his warm hospitality and to Netherlands Organization for Scientific Research (NWO) for financial support. The author would like also to thank Professor P. Moree for encouragement and valuable discussion.

¹The author is partially supported by NSF of China and NWO of the Netherlands.

1

2. Vinberg's classification theorem

The main tool of the paper is Vinberg's classification theorem. A matrix A is called decomposable if, after reordering the indices (i.e. a permutation of its rows and the same permutation of its columns), A decomposes into a non-trivial direct sum. A matrix that is not decomposable will be called indecomposable.

Let $u = (u_1, u_2, \dots, u_n)^T$ be a real column vector. We write u > 0 if $u_i > 0$ for all i, and $u \ge 0$ if $u_i \ge 0$ for all i.

Definition 2.1. Let $A = (a_{ij})_{n \times n} \in M_n(\mathbb{R})$. The matrix A is called a real generalized Cartan matrix if it satisfies the following properties:

- (1) A is indecomposable,
- (2) $a_{ij} \leq 0 \text{ for } i \neq j$,
- (3) $a_{ij} = 0$ if and only if $a_{ji} = 0$.

The following lemma was proved in [1].

Lemma 2.2. Let A be a real generalized cartan matrix and let $u \ge 0$ be a real column vector. Then $Au \ge 0$ implies that either u > 0 or u = 0.

The following theorem is known as Vinberg's classification theorem see [1], [4].

Theorem 2.3. Let A be a real generalized Cartan matrix. Then one and only one of the following possibilities holds for both A and A^T .

- (1) $det A \neq 0$; there exists u > 0 such that Au > 0; $Av \geq 0$ implies v > 0 or v = 0. In this case, A is called of finite type.
- (2) det A = 0; there exists u > 0 such that Au = 0; $Av \ge 0$ implies Av = 0. In this case, A is called of affine type.
- (3) There exists u > 0 such that Au < 0; $Av \ge 0$, $v \ge 0$ imply v = 0. In this case, A is called of indefinite type.

It is easy to see that Vinberg's classification theorem can be simplified as follows.

Corollary 2.4. Let A be a $n \times n$ real generalized Cartan matrix. Then

- (1) A is of finite type if and only if there exists $u \in \mathbb{R}^n$, u > 0 such that Au > 0.
- (2) A is of affine type if and only if there exists $u \in \mathbb{R}^n$, u > 0 such that Au = 0.
- (3) A is of indefinite type if and only if there exists $u \in \mathbb{R}^n$, u > 0 such that Au < 0.

Example. Let
$$A = \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & -1 \\ -2 & -1 & 4 \end{pmatrix}$$
. Then

$$A \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}.$$

Hence, A is of finite type. Clearly, A is non-symmetrizable.

3. THE MAIN RESULTS

Denote by I the $n \times n$ identity matrix.

Theorem 3.1. For any real generalized Cartan matrix A, there exists a unique real number $d \in \mathbb{R}$ such that A + dI is of affine type. Furthermore, A + bI is of finite type if b > d and A + bI is of indefinite type if b < d.

Proof. Let

$$I_{fin}(A) = \{a \in \mathbb{R} | A + aI \text{ is of finite type } \},$$

$$I_{aff}(A) = \{a \in \mathbb{R} | A + aI \text{ is of affine type } \},$$

$$I_{ind}(A) = \{a \in \mathbb{R} | A + aI \text{ is of indefinite type } \}.$$

By Vinberg's classification theorem, we have

$$\mathbb{R} = I_{fin}(A) \dot{\cup} I_{aff}(A) \dot{\cup} I_{ind}(A).$$

If $a \in I_{fin}(A)$ and $b \ge a$, there exists u > 0 such that (A + aI)u > 0. Hence, (A + bI)u > 0 and so $b \in I_{fin}(A)$. For this given real number a, we can find a sufficient small $\epsilon > 0$ such that $(A + (a - \epsilon)I)u > 0$. Hence, $a - \epsilon \in I_{fin}(A)$ and $I_{fin}(A)$ is an open set. Similarly, if $a \in I_{ind}(A)$ and $b \le a$, then there exists u > 0 such that (A + aI)u < 0, then (A + bI)u < 0 and so $b \in I_{ind}(A)$. For this given real number a, we can find a sufficient small $\epsilon > 0$ such that $(A + (a + \epsilon)I)u < 0$. Hence, $a + \epsilon \in I_{ind}(A)$ and $I_{ind}(A)$ is an open set.

If $a \in I_{aff}$, then -a is a real eigenvalue of the matrix A. Hence, $I_{aff}(A)$ is a finite set. Since $I_{fin}(A)$ and $I_{ind}(A)$ are open sets, $I_{aff}(A)$ is non-empty. Assume that $a_1, a_2 \in I_{aff}(A)$ and $a_1 < a_2$. Then there exists b, $a_1 < b < a_2$, such that A + bI is of finite type or indefinite type. In either cases, it is a contradiction. Hence, $I_{aff}(A)$ only contains one element, say d. Moreover,

$$I_{fin}(A) = \{ a \in \mathbb{R} | a > d \}$$

and

$$I_{ind}(A) = \{ a \in \mathbb{R} | a < d \}.$$

The following lemma was proved in [1].

Lemma 3.2. Let A be a real generalized Cartan matrix. If A is of finite or affine type, then any proper principal submatrix decomposes into a direct sum of matrices of finite type.

Theorem 3.3. The real generalized Cartan matrix A is of finite type if and only if $\det A \neq 0$ and $A^{-1} > 0$.

Proof. Let A be a real generalized Cartan matrix of finite type. Then $det A \neq 0$ by Vinberg's theorem. Let $A^{-1} = (u_1, u_2, \dots, u_n)$ where u_i 's are the column vectors of A^{-1} . Then $Au_i \geq 0$ for all i. Again by Vinberg's theorem, we have $u_i > 0$ for all i, i.e. $A^{-1} > 0$.

Conversely, assume that A is a generalized Cartan matrix such that $det A \neq 0$ and $A^{-1} > 0$. Of course, A is not of affine type. If A is of indefinite type, then there exists u > 0 such that Au = w < 0. Since $A^{-1} > 0$, we have $u = A^{-1}w < 0$ which is a contradiction. Therefore, A is of finite type.

Definition 3.4. A real generalized Cartan matrix A is called a matrix of hyperbolic type if it is of indefinite type and any proper principal submatrix of A is a direct sum of matrices of finite type or affine type. A hyperbolic generalized Cartan matrix A is called strictly hyperbolic if any proper principal submatrix of A is a direct sum of matrices of finite type.

Theorem 3.5. Let A be a real generalized Cartan matrix. Then A is of hyperbolic type if and only if for any column vector $u \in \mathbb{R}^n$, $Au \leq 0$ implies $u \geq 0$.

Proof. Let A be a generalized Cartan matrix of hyperbolic type. Let $0 \neq u \in \mathbb{R}^n$ such that $Au \leq 0$. We claim that $u \geq 0$. If u is not non-negative, then u has both positive and negative entries since $u \leq 0$ contradicts to that A is of indefinite type. By reordering, we may assume that

$$u_1 > 0, \dots, u_{s-1} > 0, u_s \le 0, \dots, u_{n-1} \le 0, u_n < 0.$$

We divide A into block form

$$A = \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix}$$

where A_1 is a $(s-1)\times(s-1)$ submatrix, A_2 is a $(s-1)\times(n-s+1)$ submatrix, A_3 is a $(n-s+1)\times(s-1)$ submatrix and A_4 is a $(n-s+1)\times(n-s+1)$ submatrix. Let $u'=(u_1,\dots,u_{s-1})^T$ and let $w'=(u_s,\dots,u_n)^T$. Then we have

$$A_1u' + A_2w' \le 0.$$

Hence, $A_1u' \leq -A_2w' \leq 0$. Since A_1 is a direct sum of matrices of finite type or affine type, we conclude that A_1 is of affine type and so s = n and $A_2u_n = 0$. Therefore, $A_2 = 0$ and so $A_3 = 0$ which contradicts to that A is indecomposable. Hence, $u \geq 0$.

If A is of indefinite type but is not of hyperbolic type, then A has a $(n-1) \times (n-1)$ principal submatrix A_1 which is of indefinite type. We may assume that

$$A = \begin{pmatrix} A_1 & -b_1 \\ -c_1^T & a_{nn} \end{pmatrix},$$

where $b_1 \ge 0$, $c_1 \ge 0$ are column vectors in \mathbb{R}^{n-1} . There exists u > 0 in \mathbb{R}^{n-1} such that $A_1 u < 0$. For $a \in \mathbb{R}_+$, set

$$w = \begin{pmatrix} au \\ -1 \end{pmatrix}.$$

Then we have Aw < 0 for sufficient large a. Contradiction!

Theorem 3.6. Let A be a real generalized Cartan matrix. Then A is of hyperbolic type if and only if $\det A \neq 0$ and $A^{-1} \leq 0$.

Proof. Assume that A is of hyperbolic type. If det A = 0, then there exist $0 \neq u \in \mathbb{R}^n$ such that Au = A(-u) = 0. By Theorem 3.5, $u \geq 0$ and $-u \geq 0$ which force u = 0. Hence, A is nondegenerate. Assume that

$$A^{-1} = (u_1, u_2, \cdots, u_n),$$

where $u_1, u_2, \dots, u_n \in \mathbb{R}^n$. Then $Au_i \geq 0$ for all i. By theorem 3.5, $u_i \leq 0$.

Conversely, assume that $det A \neq 0$ and $A^{-1} \leq 0$. Let $u \in \mathbb{R}^n$ be such that $Au = w \geq 0$. Then $u = A^{-1}w \leq 0$. Again by Theorem 3.5, A is of hyperbolic type.

Theorem 3.7. The real generalized Cartan matrix A is of finite type if and only if all of the principal minors of A are positive. The real generalized Cartan matrix A is of affine type if and only if $\det A = 0$ and all of the principal minors with $\operatorname{size} \leq n-1$ are positive.

Proof. Let B be an arbitrary real generalized Cartan matrix of finite type. Then for any $a \ge 0$, B + aI is also of finite type. Hence, $det(B + aI) \ne 0$ which means the real eigenvalues (if there are any) of B are positive. Therefore, detB > 0.

Now assume that $A = (a_{ij})_{n \times n}$ is a generalized Cartan matrix and that all of the principal minors of A are positive. We then use induction on n to show that the matrix A is of finite type. The case n = 1 is trivial. If A is not of finite type, then A must be of indefinite type because A can not be of affine type. By the induction hypothesis, any principal indecomposable submatrix is of finite type. Hence, A is of strictly hyperbolic type. Therefore, $A^{-1} \leq 0$. But the (1,1)-entry of A^{-1} is a $(n-1) \times (n-1)$ principal minor of A over det A which means that det A < 0. Contradiction! Hence, A must be of finite type.

If A is of affine type, then any proper principal submatrix of A is a direct sum of matrices of finite type. Hence, the determinant of the proper principal matrix is positive. HECHUN ZHANG

6

Conversely, assume that det A = 0 and all of the principal minors with size $\leq (n-1)$ are all positive. Clearly, A is not of finite type. If A is of indefinite type, then A must be of strictly hyperbolic type, and so $det A \neq 0$ which is a contradiction. Therefore, A is of affine type.

Remark 3.8. If a real generalized Cartan matrix A is of finite type (resp. affine type) and is symmetrizable, then it is easy to see that there exist positive real numbers d_1, d_2, \dots, d_n (see [1] 2.3) and a symmetric matrix B such that

$$A = diag(d_1, d_2, \cdots, d_n)B.$$

From the above theorem, one can see that B is positive definite (resp. semi positive definite).

Corollary 3.9. Let A be a RGCM of hyperbolic type. Then det A < 0.

Proof. We already proved that det A < 0 if A is of strictly hyperbolic type. Assume that A is of hyperbolic type. For a sufficient small $\epsilon > 0$, $A + \epsilon I$ is still of indefinite type. However, any principle indecomposable submatrix is of finite type. Hence, $det(A + \epsilon I) < 0$. Note that $det A \neq 0$. Let ϵ tend to zero. Then det A < 0.

References

- 1. Kac, V. G., Infinite-dimensional Lie algebras. Third edition. Cambridge University Press, Cambridge, 1990.
- 2. Li, W., Classification of generalized Cartan matrices of hyperbolic type. A Chinese summary appears in Chinese Ann. Math. Ser. A 9 (1988), no. 1, 103. Chinese Ann. Math. Ser. B 9 (1988), no. 1, 68–77.
- 3. Lusztig, G. and Tits, J., The inverse of a Cartan matrix. An. Univ. Timişoara Ser. Ştiinţ. Mat. 30 (1992), no. 1, 17–23.
- 4. Vinberg, E. B., Discrete linear groups that are generated by reflections. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 35 (1971), 1072–1112.
- 5. Wan, Z. X., Introduction to Kac-Moody algebras. World Scientific Publishing Co., Inc., Teaneck, NJ, 1991.
- 6. Zhang, H. C., Weights of integrable modules. (Chinese) Acta Math. Sinica 38 (1995), no. 1, 30-37.

Department of Mathematical Sciences, Tsinghua University, Beijing, 100084, P. R. China, and KdV Institute, Plantage Muidergracht 24, 1018 TV, Amsterdam, Netherlands, Department of Mathematical Sciences,

E-mail address: hzhang@math.tsinghua.edu.cn,