

## CASIMIR OPERATORS FOR LIE SUPERALGEBRAS

DIMITRY LEITES<sup>1</sup>, ALEXANDER SERGEEV<sup>2</sup>

ABSTRACT. Casimir operators — the generators of the center of the enveloping algebra — are described for simple or close to them “classical” finite dimensional Lie superalgebras with nondegenerate symmetric even bilinear form in: Sergeev A., The invariant polynomials on simple Lie superalgebras. Represent. Theory 3 (1999), 250–280; math-rt/9810111 and for the “queer” series in: Sergeev A., The centre of enveloping algebra for Lie superalgebra  $Q(n, C)$ . Lett. Math. Phys. 7, no. 3, 1983, 177–179.

Here we consider the remaining cases, and state conjectures proved for small values of parameter.

Under deformation (quantization) the Poisson Lie superalgebra  $\mathfrak{po}(0|2n)$  on purely odd superspace turns into  $\mathfrak{gl}(2^{n-1}|2^{n-1})$  and, conjecturally, the lowest terms of the Taylor series expansion with respect to the deformation parameter (Planck’s constant) of the Casimir operators for  $\mathfrak{gl}(2^{n-1}|2^{n-1})$  are the Casimir operators for  $\mathfrak{po}(0|2n)$ .

Similarly, quantization sends  $\mathfrak{po}(0|2n-1)$  into  $\mathfrak{q}(2^{n-1})$  and the above procedure makes Casimir operators for  $\mathfrak{q}(2^{n-1})$  into same for  $\mathfrak{po}(0|2n-1)$ .

Casimir operators for the Lie superalgebra  $\mathfrak{vect}(0|m)$  of vector fields on purely odd superspace are only constants for  $m > 2$ . Conjecturally, same is true for the Lie superalgebra  $\mathfrak{svect}(0|m)$  of divergence free vector fields, and its deform, for  $m > 3$ .

Invariant polynomials on  $\mathfrak{po}(0|2n-1)$  are also described. They do not correspond to Casimir operators.

This is a version of the paper published in: Ivanov E. et. al. (eds.) *Supersymmetries and Quantum Symmetries* (SQS’99, 27–31 July, 1999), Dubna, JINR, 2000, 409–411. Meanwhile we have understood how to prove our conjecture for  $\mathfrak{vect}(0|m)$  and precisely same idea was used by Noam Shomron [Sh], who additionally described, to an extent, indecomposable  $\mathfrak{vect}(0|m)$ -modules. The only difference with the Dubna version is this and similar remarks.

As always, speaking about invariant polynomials on Lie superalgebras (or their duals), one should bear in mind a more natural from super point of view approach due to Shander [Sha]: description of nonpolynomial invariant functions. Actually, even in non-super cases people did consider non-polynomial elements, say, localizations of the enveloping algebras, but there it does not lead to totally new invariants.

**0. Recapitulations.** Recall that by the *Casimir elements* for the Lie algebra or superalgebra  $\mathfrak{g}$  we mean the generators of the center  $Z(\mathfrak{g})$  of  $U(\mathfrak{g})$ . Their applications in description of physical phenomena are well-known and numerous.

If  $\mathfrak{g}$  possesses an invariant nondegenerate supersymmetric bilinear form  $B$ , then  $\mathfrak{g} \simeq \mathfrak{g}^*$ . In this case, the description of the Casimir elements for  $\mathfrak{g}$  is equivalent to the description of the algebra  $I(\mathfrak{g})$  of invariant polynomials on  $\mathfrak{g}$  because  $Z(\mathfrak{g})$  is isomorphic to  $I(\mathfrak{g})$  (as vector spaces). To describe  $I(\mathfrak{g})$  is, however, easier than  $Z(\mathfrak{g})$ . For the most complete description of  $I(\mathfrak{g})$  for simple finite dimensional Lie superalgebras and/or their “relatives”, i.e., close to them “classical” algebras, such as central extensions, etc., see [S1].

1991 *Mathematics Subject Classification*. 17A70 (Primary) 17B01, 17B70 (Secondary).

*Key words and phrases*. Lie superalgebras, Casimir operators.

The financial support of NFR is gratefully acknowledged.

This description of  $I(\mathfrak{g})$  does not cover  $\mathfrak{q}(n)$ , and  $\mathfrak{po}(0|m)$ , and their relatives: it is a much more difficult problem to be solved by other means and ideas than those of [S1].

Thus, Casimir operators are described at the moment for

- simple Lie superalgebras with Cartan matrix and their “relatives” [S1]; (the first results are due to Berezin [B] and Kac [K1, K2]);
- $\mathfrak{q}(n)$  (but not for relatives), see [S3];
- $\mathfrak{pe}(n)$  (no Casimir operators) and  $\mathfrak{spe}(n)$ , see [Sch] (implicitly) and [S2] (for  $n = 3$ ); now we can refer to an interesting paper of Maria Gorelik [G], where the  $\mathfrak{spe}(n)$ -invariants are completely and explicitly described (though the author herself does not like the result and did not even put on arXive, to say nothing about publishing, we find it a very nice one; it is based on her previous papers [G2]);
- $\mathfrak{vect}(0|m)$  (for  $m > 2$ ), see Shomron’s paper [Sh].

There remain the following cases:

- the divergence free sublagebras: the “standard” one,  $\mathfrak{svect}(0|m)$ , that preserves the volume element  $vol$  and its deformation  $\widetilde{\mathfrak{svect}(0|m)}$  preserving  $(1 + \theta_1 \cdot \theta_m)vol$  for which, conjecturally, there are no Casimir operators (for  $m > 3$ );
- $\mathfrak{sq}(n)$  and  $\mathfrak{psq}(n)$ ;
- $\mathfrak{po}(0|m)$  considered below (but the general proof is still lacking),  $\mathfrak{spo}(0|m)$  as well as  $\mathfrak{h}(0|m)$  and  $\mathfrak{sh}(0|m)$ .

**Example 1.** For  $\mathfrak{g} = \mathfrak{gl}(n)$  the Casimir operators corresponding to the first  $n$  invariant polynomials  $\text{tr}(X^k)$ , where  $X \in \mathfrak{g}$ , i.e., for  $k = 1, \dots, n$ , generate the center of  $U(\mathfrak{g})$ . The form  $B$  is defined by the trace.

**Example 2.** The Lie superalgebra  $\mathfrak{gl}(m|n)$  is a super analog of  $\mathfrak{gl}(n)$ ; the form  $B$  is defined by the supertrace. It is known that the center of its universal enveloping algebra is NOT finitely generated. For the generators we may take  $\text{str}(X^k)$  for all positive integer values of  $k$ . (A comment: suppose we somehow proved that apart from traces there are no invariant polynomials on  $\mathfrak{gl}(n)$ . To prove that only finitely many of them generate the algebra, consider the power series expansion of  $\det(X - \lambda 1_n)$  with respect to  $\lambda$ . The coefficients of  $\lambda$  are polynomials in  $\text{tr}(X^k)$ ; the characteristic polynomial yields a recursive formula for  $\text{tr}(X^{n+1})$ . Contrarywise, the superdeterminant, the Berezinian, is a rational function in  $\text{str}(X^k)$ , to express the Berezinian in terms of supertraces we need all of them.)

On the other hand, any *rational* function on  $\mathfrak{gl}(m|n)$  can be expressed as a rational function of the first  $n + m$  supertraces, cf. [S1] with [B] and [K1, K2]).

**Example 3.** The Lie superalgebra  $\mathfrak{po}(0|2n)$ , the Poisson superalgebra, is one more super analog of  $\mathfrak{gl}(n)$ . Indeed, there exists a well-known deformation (quantization) with parameter  $\hbar$  such that at  $\hbar = 0$  we have  $\mathfrak{po}(0|2n)$  whereas at  $\hbar \neq 0$  the Lie superalgebras from the parametric family are isomorphic to  $\mathfrak{gl}(2^{n-1}|2^{n-1})$ .

Recall, that the superspace of  $\mathfrak{po}(0|2n)$  is isomorphic to the Grassmann superalgebra on  $2n$  generators  $\xi, \eta$  and the bracket is given by the formula

$$\{f, g\} = (-1)^{p(f)} \sum \left( \frac{\partial f}{\partial \xi_i} \frac{\partial g}{\partial \eta_i} + \frac{\partial f}{\partial \eta_i} \frac{\partial g}{\partial \xi_i} \right).$$

The form  $B$  on  $\mathfrak{po}(0|m)$  for any  $m$  is given by the integral  $B(f, g) = \int fg \, vol$ , where  $vol$  is the volume element, and, clearly, the following are invariant polynomials on  $\mathfrak{po}(0|m)$ :

$$r_k = \int f^k vol(\xi, \eta), \text{ where } k = 1, 2, \dots \quad (*)$$

Observe, however, that, unlike  $\mathfrak{gl}(m|n)$  case, these polynomials  $r_k$  do not generate the whole algebra of invariant polynomials for  $n \geq 2$ . A counterexample of least degree is given for  $n = 2$  by the Casimir operator whose radial part is equal to  $x_1^2 x_2^2 (x_1^2 - x_2^2)$ .

Quantization turns the integral on functions that generate  $\mathfrak{po}(0|m)$  into the supertrace on supermatrices from  $\mathfrak{gl}(2^{n-1}|2^{n-1})$  for  $m = 2n$  and into queertrace on the general queer superalgebra  $\mathfrak{q}(2^{n-1})$  for  $m = 2n - 1$ . This and other arguments, see [BL], indicates that  $\mathfrak{q}(n)$  is one more super analog of  $\mathfrak{gl}(n)$ . Recall that  $\text{qtr} \left( \begin{pmatrix} A & B \\ B & A \end{pmatrix} \right) = \text{tr} B$ , see [BL].

*Remark .* For any Lie superalgebra  $\mathfrak{g}$  and its finite dimensional representation  $\rho$ , the polynomials

$$P_{k,\rho} = \text{str} \rho(X)^k$$

are invariant ones. For  $\mathfrak{gl}(m|n)$  any degree  $k$  homogeneous invariant polynomial can be represented as

$$\sum_{\rho} c_{\rho} P_{k,\rho}, \quad (**)$$

where the sum runs over a finite number of finite dimensional representations, see [S1]. For Lie superalgebras  $\mathfrak{po}(0|2n)$  the situation is different: when  $n \geq 2$  and  $k \geq 6$  the polynomial  $(*)$  can not be obtained as any finite sum of the form  $(**)$ , so we have strict inclusions  $I(\mathfrak{g}) \supset I_{(*)}(\mathfrak{g}) \supset I_{(**)}(\mathfrak{g})$ , where  $I_{(*)}$  and  $I_{(**)}$  are algebras of polynomials of the form  $(*)$  and  $(**)$ , respectively.

To describe the complete set of  $\mathfrak{po}(0|2n)$ -invariants, recall that the Grassmann superalgebra  $\mathbb{C}[\xi, \eta]$  can be deformed into the Clifford superalgebra  $\text{Cliff}_{\hbar}(\hat{\xi}, \hat{\eta})$ , where

$$\hat{\xi}_i \hat{\eta}_j + \hat{\eta}_j \hat{\xi}_i = \delta_{ij} \hbar.$$

For any subset  $I = \{i_1 < \dots < i_l\} \subset \{1, 2, \dots, n\}$  set  $\xi_I = \xi_{i_1} \dots \xi_{i_l}$  and  $\eta_I = \eta_{i_1} \dots \eta_{i_l}$ ; notations  $\hat{\xi}_I$  and  $\hat{\eta}_I$  are similar. Define the linear map  $Q : \mathbb{C}[\xi, \eta] \rightarrow \text{Cliff}_{\hbar}(\hat{\xi}, \hat{\eta})$  by setting  $Q(\xi_I \eta_J) = \hat{\xi}_I \hat{\eta}_J$ ; before applying  $Q$  to a monomial one has to reduce it to a normal form, say, to  $\xi \eta$ -form.

**4. Lemma .**  $[Q(f), Q(g)] = \hbar Q(\{f, g\}) + \mathcal{O}(\hbar^2)$ , where in the left hand side stands the supercommutator in the Clifford superalgebra, and  $\{\cdot, \cdot\}$  in the right hand side is the Poisson bracket.

The Lie superalgebra associated with the associative superalgebra  $\text{Cliff}_{\hbar}(\hat{\xi}, \hat{\eta})$  is  $\mathfrak{gl}(2^{n-1}|2^{n-1})$ ; for it, the invariant polynomials are described above and in [S1]. The Lie superalgebra associated with the Clifford superalgebra generated by  $2n - 1$  elements is  $\mathfrak{q}(2^{n-1})$  (see [L]); for it, the Casimir elements are described in [S3].

**5. Lemma .** Let  $F$  be an invariant polynomial on  $\text{Cliff}_{\hbar}$  considered as Lie superalgebra, i.e., on  $\mathfrak{gl}(2^{n-1}|2^{n-1})$  or on its subalgebra  $\mathfrak{q}(2^{n-1})$ . Let us represent it in the form  $F = \sum_{k \geq k_0} F_k \hbar^k$ .

If  $P \in I(\mathfrak{po}(0|2n))$  is such that  $Q(P) = F_{k_0}$ , then  $P$  is an invariant polynomial on  $\mathfrak{po}(0|m)$ .

In other words, the lowest (with respect to  $\hbar$ -degree) components of invariant polynomials in the operators, i.e., on  $\mathfrak{gl}(2^{n-1}|2^{n-1})$  or  $\mathfrak{q}(2^{n-1})$ , are invariant polynomials in their symbols — the elements from  $\mathfrak{po}(0|m)$  for  $m = 2n$  or  $2n - 1$ , respectively. In particular, it is easy to see that

$$\text{str}(Q(f)^k) \quad (\text{or } \text{qtr}(Q(f)^k)) = \frac{1}{k} \left( \int f^k \text{vol} \right) \hbar^n + \mathcal{O}(\hbar^n). \quad (***)$$

**6. Conjecture .** ([S2]) All the invariant polynomials on  $\mathfrak{po}(0|m)$  for any  $m$  (odd as well as even) may be obtained in the way indicated in Lemma 5.

This 15 years old conjecture is now proven for  $m$  even and  $\leq 6$ . To prove it in full generality, we need new ideas as compared with those we know and successfully used in [S1]: here the degree of complexity of computations performed à la [S1] grows steeply with  $m$ .

#### REFERENCES

- [B] Berezin, F. Representations of the supergroup  $U(p, q)$ . Funkcional. Anal. i Prilozhen. 10 (1976), n. 3, 70–71 (in Russian); Berezin, F. Laplace–Cazimir operators on Lie supergroups. The general theory. Preprints ITEPh 77, Moscow, ITEPh, 1977; Berezin F. *Analysis with anticommuting variables*, Kluwer, 1987
- [BL] Bernstein J. N., Leites D. A., The superalgebra  $Q(n)$ , the odd trace and the odd determinant. C. R. Acad. Bulgare Sci. 35 (1982), no. 3, 285–286
- [G] Gorelik M., The center of a simple  $P$ -type Lie superalgebra. (preprint, 23 Sep 2001)
- [G2] Gorelik M., On the Ghost Centre of Lie Superalgebras, math.RT/9910114; id., Annihilation Theorem and Separation Theorem for basic classical Lie superalgebras, math.RA/0008143
- [K1] Kac V. G., Characters of typical representations of classical Lie superalgebras. Commun. Alg. v. 5, 1977, 889–897
- [K2] Kac V. G., Representations of classical Lie superalgebras. In: Bleuler K. et al. (eds.) *Differential geometrical methods in mathematical physics*, II (Proc. Conf., Univ. Bonn, Bonn, 1977), Lecture Notes in Math., 676, Springer, Berlin, 1978, 597–626.
- [L] Leites, D. A. Clifford algebras as superalgebras, and quantization. (Russian. English summary) Teoret. Mat. Fiz. 58 (1984), no. 2, 229–232. (English translation: Theoret. and Math. Phys. 58 (1984), no. 2, 150–152.)
- [Sch] Scheunert M., Invariant supersymmetric multilinear forms and the Casimir elements of  $P$ -type Lie superalgebras. J. Math. Phys. 28, no. 5, 1987, 1180–1191; id., Casimir elements of the general linear and the  $P$ -type Lie superalgebras. Proceedings of the XV International Conference on Differential Geometric Methods in Theoretical Physics (Clausthal, 1986), 275–288, World Sci. Publishing, Teaneck, NJ, 1987
- [S1] Sergeev A., The invariant polynomials on simple Lie superalgebras. Represent. Theory 3 (1999), 250–280; math-rt/9810111
- [S2] Sergeev A., Laplace operators and Lie superalgebra representations. Ph.D. thesis, Moscow University, 1985. (an expanded English translation in: Leites D. (ed.) *Seminar on Supermanifolds*, Reports of Stockholm University, n. 32/1988–15, 44–95
- [S3] Sergeev A., The centre of enveloping algebra for Lie superalgebra  $Q(n, C)$ . Lett. Math. Phys. 7, no. 3, 1983, 177–179
- [Sha] Shander V., Invariant functions on supermatrices, math.RT/9810112 (a detailed version of Shander, V. N. Orbits and invariants of the supergroup  $GQ_n$ . (Russian) Funktsional. Anal. i Prilozhen. 26 (1992), no. 1, 69–71 translation in Funct. Anal. Appl. 26 (1992), no. 1, 55–56)
- [Sh] Shomron N., Blocks of Lie Superalgebras of Type  $W(n)$ , math.RT/0009103

<sup>1</sup>(CORRESPONDENCE): DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STOCKHOLM, ROSLAGSV. 101, KRÄFTRIKET HUS 6, SE-106 91, STOCKHOLM, SWEDEN; MLEITES@MATEMATIK.SU.SE; <sup>2</sup>BALAKOVO INST. OF TECHNIQUE OF TECHNOLOGY AND CONTROL, BRANCH OF SARATOV TECHNICAL UNIVERSITY, BALAKOVO, SARATOV REGION, RUSSIA