

THE HOWE DUALITY AND THE PROJECTIVE REPRESENTATIONS OF SYMMETRIC GROUPS

ALEXANDER SERGEEV

ABSTRACT. The symmetric group \mathfrak{S}_n possesses a nontrivial central extension, whose irreducible representations, different from the irreducible representations of \mathfrak{S}_n itself, coincide with the irreducible representations of the algebra \mathfrak{A}_n generated by indeterminates $\tau_{i,j}$ for $i \neq j$, $1 \leq i, j \leq n$ subject to the relations

$$\begin{aligned} \tau_{i,j} &= -\tau_{j,i}, \quad \tau_{i,j}^2 = 1, \quad \tau_{i,j}\tau_{k,l} = -\tau_{k,l}\tau_{i,j} \text{ if } \{i,j\} \cap \{k,l\} = \emptyset; \\ \tau_{i,j}\tau_{j,k}\tau_{i,j} &= \tau_{j,k}\tau_{i,j}\tau_{j,k} = -\tau_{i,k} \text{ for any } i, j, k, l. \end{aligned}$$

Recently M. Nazarov realized irreducible representations of \mathfrak{A}_n and Young symmetrizers by means of the Howe duality between the Lie superalgebra $\mathfrak{q}(n)$ and the Hecke algebra $H_n = \mathfrak{S}_n \circ C_n$, the semidirect product of \mathfrak{S}_n with the Clifford algebra C_n on n indeterminates.

Here I construct one more analog of Young symmetrizers in H_n as well as the analogs of Specht modules for \mathfrak{A}_n and H_n .

§1. INTRODUCTION

Lately we witness an increase of interest in the study of representations of symmetric groups. In particular, in their projective representations.

Recall that the symmetric group \mathfrak{S}_n has a nontrivial central extension whose irreducible representations do not reduce to those of \mathfrak{S}_n but coincide (may be identified) with the irreducible representations of the algebra \mathfrak{A}_n determined by generators $\tau_{i,j}$ for $i \neq j$, $1 \leq i, j \leq n$ subject to the relations

$$\begin{aligned} \tau_{i,j} &= -\tau_{j,i}, \quad \tau_{i,j}^2 = 1, \quad \tau_{i,j}\tau_{k,l} = -\tau_{k,l}\tau_{i,j} \text{ if } \{i,j\} \cap \{k,l\} = \emptyset; \\ \tau_{i,j}\tau_{j,k}\tau_{i,j} &= \tau_{j,k}\tau_{i,j}\tau_{j,k} = -\tau_{i,k} \text{ for any } i, j, k, l. \end{aligned} \tag{1.1}$$

In [N1] Nazarov realized irreducible representations of \mathfrak{A}_n by means of an orthogonal basis constructed in each of the spaces of the representations and indicating the action of the generators $\tau_{i,i+1}$ on them (an analog of the Young orthogonal form). In [N2], with the help of an “odd” analog of the degenerate affine Hecke algebra Nazarov constructed elements of the algebra $H_n = \mathfrak{S}_n \circ C_n$, the semidirect product of \mathfrak{S}_n with the Clifford algebra C_n on n indeterminates. The elements of H_n serve as analogs of Young symmetrizers.

Here I construct one more analog of Young symmetrizers in H_n as well as analogs of Specht modules (cf. [Ja]) for the algebras \mathfrak{A}_n and H_n . This construction is based on another form of expression of Young symmetrizers for \mathfrak{S}_n .

Namely, let t be a Young tableau (i.e., a Young diagram filled in with numbers 1 to n), R_t and C_t the row and column stabilizers of t ;

$$\rho_t = \sum_{\sigma \in R_t} \sigma, \quad \kappa_t = \sum_{\sigma \in C_t} \varepsilon(\sigma) \sigma.$$

1991 *Mathematics Subject Classification.* 20C30, 20C25, 17A70.

Key words and phrases. Projective representations, symmetric group, Howe duality.

I am thankful to D. Leites for support and help.

Then, up to a constant factor, the Young symmetrizer of t is

$$e_t = \kappa_t \rho_t. \quad (1.2)$$

Let us represent the Young symmetrizer differently. Let I be the sequence obtained by reading the tableau t along columns left to right and downwards. For each $i \in I$ the *Jucys–Murphy elements* ($[J]$, $[M]$) are defined to be

$$x_i = \sum_{\alpha \text{ preceeds } i} s_{\alpha i}, \text{ where the } s_{\alpha i} \in \mathfrak{S}_n \text{ are transpositions.}$$

One can verify that the x_i commute with each other. Set

$$\tilde{\kappa}_t = \prod_{i \in I} (j - x_i), \text{ where } j \text{ is the number of the column occupied by } i.$$

It is subject to a direct verification that e_t can be also expressed as

$$e_t = \tilde{\kappa}_t \rho_t. \quad (1.3)$$

I use this other representation of the Young symmetrizers to construct the corresponding elements in H_n and prove that theyse corresponding elements are idempotents that generate isotypical ideals. With the help of these elements and their analogs for \mathfrak{A}_n I construct a realization of multiples of irreducible modules similar to Specht modules, cf. [Ja].

The proofs are based on the Howe duality between Lie superalgebra $\mathfrak{q}(n)$ and H_n .

§2. BACKGROUND

Let \mathfrak{S}_n be the symmetric group, C_n the Clifford algebra generated by n indeterminates p_1, \dots, p_n subject to the relations

$$p_i^2 = -1, \quad p_i p_j + p_j p_i = 0 \text{ for } i \neq j.$$

The symmetric group acts on C_n permuting the generators, so we can form a semidirect product $H_n = \mathfrak{S}_n \circ C_n$. Set

$$\tau_{i,j} = \frac{1}{\sqrt{2}}(p_i - p_j)s_{i,j}.$$

As is not difficult to verify, the relations (1.1) hold; hence, the algebra generated by the $\tau_{i,j}$ is isomorphic to \mathfrak{A}_n . Besides, \mathfrak{A}_n supercommutes with C_n ; hence, $H_n = \mathfrak{A}_n \otimes C_n$, as *superalgebras* if we define parity in \mathfrak{A}_n by setting $p(\tau_{i,j}) = \bar{1}$.

Let V be a superspace of superdimension (n, n) with the fixed basis $\{e_i\}_{i=1}^n \cup \{e_{\bar{i}}\}_{\bar{i}=1}^{\bar{n}}$ and the odd operator

$$Q : e_i \mapsto (-1)^{\bar{i}} e_{\bar{i}}; \quad e_{\bar{i}} \mapsto e_i.$$

The (super)centrilizer of Q in $\text{Mat}(V)$ is denoted by $Q(V)$, cf. [BL]. We denote the Lie superalgebras associated with the associative superalgebras $\text{Mat}(V)$ and $Q(V)$ by $\mathfrak{gl}(V)$ and $\mathfrak{q}(V)$, respectively.

As is shown in [S1], the algebras $\mathfrak{q}(V)$ and H_k constitute a Howe-dual pair in the superspace $W = V^{\otimes k}$. Therefore, the following decomposition takes place:

$$W = \bigoplus_{\lambda} 2^{-\delta(\lambda)} T^{\lambda} \otimes V^{\lambda}, \quad (2.1)$$

where λ runs over strict partitions of k , T^{λ} is an irreducible (in supersence) H_k -module, V^{λ} an irreducible $\mathfrak{q}(n)$ -module and $\lambda_{n+1} = 0$ and where

$$\delta(\lambda) = \begin{cases} 0 & \text{if the number of nonzero parts of } \lambda \text{ is even} \\ 1 & \text{otherwise.} \end{cases}$$

Select a basis in $\mathfrak{q}(n)$: let the e_i^* be the left dual basis to the $\{e_i\}$, $i = 1, \dots, n; \bar{1} \dots \bar{n}$; set

$$E_{i,j} = e_i \otimes e_j^* + e_{\bar{i}} \otimes e_{\bar{j}}^*; \quad F_{i,j} = e_i \otimes e_j^* + e_{\bar{i}} \otimes e_j^*.$$

Then $\mathfrak{h} = \text{Span}(E_{i,i} \text{ and } F_{i,i} : i = 1, \dots, n)$ is a Cartan subalgebra in $\mathfrak{q}(n)$ and $\mathfrak{b} = \text{Span}(E_{i,j} \text{ and } F_{i,j} : i \leq j)$ is a Borel subalgebra, cf. [Pe].

If λ is a strict partition and $\lambda_{n+1} = 0$, then λ can be interpreted as a linear functional on \mathfrak{h}_0 : set

$$\lambda(E_{ii}) = \lambda_i. \quad (2.2)$$

Let R^λ be the H_k -module equal to the direct sum of $2^{(l(\lambda) - \delta(\lambda))/2}$ copies of T^λ . It is not difficult to see that R^λ coincides with the set of \mathfrak{b} -highest weight vectors of weight λ in W .

2.1. Lemma . *Let μ be a strict partition, $\mu_{n+1} = 0$. Let V^μ be an isotypical modul of type μ and n_μ the multiplicity of the highsest weight vector in V^μ . Then the highest weights of $V^\mu \otimes V$ are of the form $\mu + \varepsilon_i$, where the ε_i are the weights of V , and their multiplicity is equal to $2n_\mu$.*

Proof follows easily from the multiplication table of the projective Schur functions, see [P]. \square

2.2. Lemma . *Let $\mathfrak{g} = \mathfrak{q}(n)$; \mathfrak{b} and \mathfrak{h} be defined as above and V a \mathfrak{g} -module. Let V_λ^+ be the set of \mathfrak{b} -highest vector of weight λ .*

If $u \in U(\mathfrak{g})$ and $uV_\lambda^+ \subset V_\lambda^+$, then there exists $w \in U(\mathfrak{h})$ such that $u|_{V_\lambda^+} = w|_{V_\lambda^+}$.

Proof. Let $u = \sum u_\alpha$ be the weight decomposition of $u \in U(\mathfrak{g})$ with respect to \mathfrak{h}_0 , the even part of the Cartan subalgebra \mathfrak{h} . Therefore, if $v \in V_\lambda$, then $uv = \sum_\alpha u_\alpha v$ and, if $u_\alpha v \neq 0$, then the weight of $u_\alpha v$ is equal to $\lambda + \alpha$. Thus, we may assume that $u = u_0 \in U(\mathfrak{g})^{\mathfrak{h}_0}$, where $U(\mathfrak{g})^{\mathfrak{h}_0}$ is the centralizer of \mathfrak{h} .

Thanks to [S2], we know that $U(\mathfrak{g})^{\mathfrak{h}_0} \cong U(\mathfrak{h}) \oplus L$, where $L = U(\mathfrak{g})^{\mathfrak{h}_0} \cap U(\mathfrak{g})\mathfrak{b}^+$, where \mathfrak{b}^+ is the linear span of the positive roots in \mathfrak{b} , is a twosided ideal in $U(\mathfrak{g})^{\mathfrak{h}_0}$.

Hence, $u = w + u_1$, where $w \in U(\mathfrak{h})$ and $u_1 \in L$; this implies that $uv = wv$ for $v \in V_\lambda$. \square

§3. SPECHT MODULES OVER H_k

Let λ be a strict partition and t the shifted tableau of the form λ , where $\lambda_1 > \lambda_2 > \dots > \lambda_n > 0$ and $\sum \lambda_i = k$. Let us fill in the tableau with the numbers 1 to k and define the functional λ on the Cartan subalgebra \mathfrak{h} by setting

$$\lambda(E_{ii}) = \lambda_i; \quad \lambda(F_{ii}) = 0.$$

In $W = V^{\otimes k}$, where V is the standard $\mathfrak{q}(V)$ -module of dimension (n, n) , consider the submodule M^λ consisting of the vectors of weight λ .

Again, let R_t be the row stabilizer of t and $\rho_t = \sum_{\sigma \in R_t} \sigma$. Let I be the sequence obtained by reading the tableau t downwards and from left to right. For $i \in I$ define: $\pi_i = \sum \tau_{\alpha, i}$, where the sum runs over all the α 's, $\alpha \in I$, that preceed $i \in I$.

It is subject to a direct verification that

$$\pi_i \pi_j + \pi_j \pi_i = 0 \text{ for } i \neq j.$$

These are odd analogs of the Jucys-Murphy elements. Set

$$\kappa_t = \prod_{i \in I} \left(\frac{1}{2} j(j+1) - \pi_i^2 \right),$$

where j is the number of the column occupied by i .

3.1. Theorem . *Let $v_t \in M^\lambda$ be the vector whose stabilizer is R_t . Then $\kappa_t(v_t) \neq 0$, $\kappa_t(v_t) \in R^\lambda$ and $\kappa_t(M^\lambda) = C_k \kappa_t(v_t)$.*

Proof. Induction on $\sum \lambda_i = k$. It suffices to assume that t is filled in consequently columnwise, from left to right and downwards. Let s be the tableau obtained from t by deleting the last cell and μ the corresponding partition while l is the length of the last column and r is its number. Then $\kappa_t = \kappa_s \cdot \kappa_k$, where κ_s corresponds to tableau s and $\kappa_k = \frac{1}{2}r(r+1) - \pi_k^2$.

By induction, $\kappa_s(M^\mu) = C_{k-1} \kappa_s(v_s) \subset R^\mu$. Therefore, $\kappa_s(M^\nu) = 0$ for any $\nu > \mu$ ordered with respect to *dominance*, cf. [M]. Hence,

$$\kappa_s(M^\lambda) = \bigoplus_{i=1}^n \kappa_s(M^{\lambda-\varepsilon_i}) \otimes e_i.$$

If $i > l$, then $\lambda - \varepsilon_i > \lambda - \varepsilon_l = \mu$ and, by induction, $\kappa_s(M^{\lambda-\varepsilon_i}) = 0$. Hence,

$$\kappa_s(M^\lambda) = \bigoplus_{i=1}^l \kappa_s(M^{\lambda-\varepsilon_i}) \otimes e_i.$$

The inequality $\nu \geq \lambda - \varepsilon_i$ true for $i < l$ implies that $\nu \geq \mu$, so the same irreducible H_{k-1} -modules enter the decomposition of $M^{\lambda-\varepsilon_i}$ as those that enter that of M^μ . Therefore, if $m_i \in M^{\lambda-\varepsilon_i}$ and $\kappa_s(m_i) \neq 0$, then there exist $m \in M^\mu$ and a homomorphism $\varphi_i : M^\mu \rightarrow M^{\lambda-\varepsilon_i}$ such that

$$\varphi_i(m) = m_i.$$

Applying κ_s to this identity we get

$$\kappa_s(m_i) = \kappa_s(\varphi_i(m)) = \varphi_i(\kappa_s(m)) \in C_{k-1} \kappa_s(v_s).$$

The Howe duality between $\mathfrak{q}(n)$ and H_{k-1} allows us to assume that $\varphi_i \in U(\mathfrak{q}(n))$. This proves that $\kappa_s(M^\lambda) \subset V^\mu \otimes V$, where V^μ is the $\mathfrak{q}(n)$ -submodule of $V^{\otimes(k-1)}$ generated by $\kappa_s(M^\mu)$. By Lemma 2.1 the highest weights of $V^\mu \otimes V$ are of the form $\mu + \varepsilon_i$, so the possible weights are only

$$\mu + \varepsilon_1, \mu + \varepsilon_l = \lambda \text{ and } \mu + \varepsilon_i < \mu + \varepsilon_l \text{ if } i > l.$$

So the submodule generated by the weights $\mu + \varepsilon_i$ for $i > l$ does not contain weight λ ; hence, neither does it contain $\kappa_s(M^\lambda)$.

Therefore, $\kappa_s(M^\lambda)$ is contained in the submodule generated by the highest weight vectors of weight $\mu + \varepsilon_1$ and $\mu + \varepsilon_l$. By Lemma 2.1 all the highest weight vectors form a free C_k -module with two generators whose weights are $\mu + \varepsilon_1$ and $\mu + \varepsilon_l$; so in each of these submodules the operator π_k^2 acts by multiplying by a constant. Let $v \in V^{\otimes(k-1)}$ and $e_i \in V$. Then it is not difficult to verify that

$$\pi_k(v \otimes e_i) = \frac{1}{\sqrt{2}} \sum_{1 \leq i \leq n} (F_{ij} - p_k E_{ij})(v \otimes e_i),$$

where p_i is the change of parity operator in the i -th factor of $V^{\otimes k}$.

If v is a highest weight vector of weight μ , then $v \otimes e_1$ is a highest weight vector of weight $\mu + \varepsilon_1$ and

$$\pi_k(v \otimes e_1) = \frac{1}{\sqrt{2}} (F_{11} - p_k E_{11})(v \otimes e_1),$$

hence,

$$\begin{aligned} \pi_k^2(v \otimes e_1) &= -\frac{1}{2} (F_{11} - p_k E_{11})^2 (v \otimes e_1) = \\ &= \frac{1}{2} (E_{11}^2 - E_{11})^2 (v \otimes e_1) = \\ &= \frac{1}{2} ((r+1)^2 - (r+1)) (v \otimes e_1) = \frac{1}{2} r(r+1) (v \otimes e_1). \end{aligned}$$

Therefore, it suffices to demonstrate that

$$\kappa_t(v_l) = \kappa_t(v_s \otimes e_l) = \kappa_k(\kappa_s(v_s \otimes e_l)) \neq 0.$$

It is not difficult to verify that if v is the highest weight vector, then

$$\pi_k^2(v \otimes e_l) = -\frac{1}{2} \sum_{k \leq i} [(E_{ik}^{(2)} v) \otimes e_k + p_k(F_{ik}^{(2)} v) \otimes e_k] + 2(\sum E_{kk} v) \otimes e_i,$$

where

$$E_{ik}^{(2)} = \sum_{k \leq j \leq i} (E_{ij} E_{jk} - F_{ij} F_{jk}), \quad F_{ik}^{(2)} = \sum_{k \leq j \leq i} (E_{ij} F_{jk} - F_{ij} E_{jk}).$$

Therefore,

$$\pi_k^2(v \otimes e_l) = \frac{1}{2} (E_{ll}^{(2)} - E_{ll} + 2(E_{11} + \dots + E_{ll})) v \otimes e_k + \dots,$$

where ... replace the sum of terms of the form $u \otimes e_i$ with $i < l$ and

$$\begin{aligned} \pi_k^2(v \otimes e_l) &= \frac{1}{2} (\lambda_l^2 - \lambda_l + 2(\lambda_1 + \dots + \lambda_l)) v \otimes e_l + \dots = \\ &= \frac{1}{2} ((r-l)(r-l-1) + 2(r + \dots + r - (l-1) + 1 + r - l)) v \otimes e_l + \dots = \\ &= \frac{1}{2} (r^2 - r + 2l - 2) v \otimes e_l + \dots \end{aligned}$$

Consequently,

$$\kappa_k(v \otimes e_l) = \frac{1}{2} (r^2 + r - r^2 + r - 2l + 2) (v \otimes e_l) + \dots = (r - l + 1) (v \otimes e_l) + \dots \neq 0.$$

□

3.2. Remark . By [N2] (Th. 7.2) for any strict partition μ there exist elements $\psi_S \in R^\mu$, where S is the standard tableau of type μ , such that $p\psi_S$ constitute a basis of R^μ while p runs over C_n . Moreover, the action of the elements

$$x_k = \sum_{i < k} (s_{ik} + s_{ik} p_i p_k)$$

is given by the formula

$$x_k \psi_S = C_S(k) \psi_\Lambda,$$

where $C_S(k) = \sqrt{(j-i)(j-i+1)}$ and k occupies the (i, j) -th slot.

In the above notations

$$x_k = \sqrt{2} p_k \pi_k,$$

hence, $x_k^2 = 2\pi_k$ and

$$\begin{aligned} \kappa_t &= \prod_i \frac{1}{2} (j(j+1) - \pi_i^2) = \\ &= \prod_i \frac{1}{2} (j(j+1) - x_i^2) = 2^{-k} \prod_i (j(j+1) - x_i^2). \end{aligned}$$

As earlier, we assume that the tableau is filled in left to right and downwards along columns.

Let us show that if $\mu \geq \lambda$, then $\kappa_t \psi_S = 0$ for any standard tableau S of type μ and distinct from t .

Let k be the number occupying the last place of the last column in the tableau t (filled in left to right and downwards). Let t^* be the tableau obtained from t by deleting the box with k ; let S^* be similarly constructed from S . If $\mu^* \geq \lambda^*$ (shapes of S^* and t^* , respectively), then the induction hypothesis applies. If $\mu^* < \lambda^*$, then $\mu_1 = \lambda_1 + 1$ and k occupies the last

position of the first row of S . Hence, if j is the number of the column of t occupied by k , then

$$\begin{aligned}\pi_k^2(v_S) &= \frac{1}{2}(j+1-1)(j+1-1+1)v_S = \\ &= \frac{1}{2}j(j+1)v_S.\end{aligned}$$

Hence, $(\frac{1}{2}j(j+1) - \pi_k^2)\psi_S = 0$ and $\kappa_t\psi_S = 0$.

This argument shows that κ_t is the projection in M^λ onto the subspace generated by ψ_t . At the same time

$$\begin{aligned}\kappa_k\psi_t &= \frac{1}{2}(j(j+1) - (j-i)(j-i+1))\psi_t = \\ &= (ij - \frac{1}{2}i(i-1))\psi_t \neq 0.\end{aligned}$$

3.3. Specht modules. Recall that the *Specht module* for a strict partition λ is the submodule in M^λ generated by the vectors κ_tv_t for all tableaux t , where v_t is the vector whose stabilizer is equal to R_t .

3.3.1. Theorem . *Specht module is equal to R^λ . It is isotypical and its H_k -centralizer is isomorphic to the Clifford algebra with $l(\lambda)$ generators.*

Proof. Let us realize R^λ as the set of highest weight vectors in $V^{\otimes k}$. By the Howe duality between $U(\mathfrak{q}(V))$ and H_k the algebra of H_k -homomorphisms is generated by $U(\mathfrak{q}(V))$. But thanks to Lemma 2.2 we may assume that the algebra of H_k -homomorphisms is generated by $U(\mathfrak{h})$ for the Cartan subalgebra \mathfrak{h} of $\mathfrak{q}(V)$. If $\lambda_i = 0$, then in our notations for the basis of \mathfrak{h} we have

$$F_{ii}^2 = E_{ii} = \lambda_i = 0 \text{ in } R^\lambda$$

and, therefore, $\ker F_{ii}$ is an \mathfrak{h} -submodule in the set $(V^\lambda)^+$ of highest weight vectors. But $(V^\lambda)^+$ is irreducible as \mathfrak{h} -module; hence, $F_{ii}|_{(V^\lambda)^+} = 0$ which implies $F_{ii}|_{R^\lambda} = 0$. Thus, the algebra of H_k -homomorphisms is generated by the F_{ii} for i such that $\lambda_i \neq 0$.

By Theorem 3.1 the Specht module is contained in R^λ and any homomorphism of M^λ sends R^λ into itself. Hence, the Specht module coincides with R^λ . \square

3.3.2. Corollary . *Set $p_t^i = \sum p_\alpha$, where the α belongs to the i -th column of tableau t . Then for any H_k -module endomorphism φ of M^λ we have*

$$\varphi(\kappa_tv_t) = f \cdot \kappa_tv_t,$$

where f belongs to the subalgebra of C_k generated by $p_t^1, \dots, p_t^{l(\lambda)}$.

Proof. Let us realize M^λ as the subset of vectors of weight λ in $V^{\otimes k}$. Then any endomorphism of M^λ may be identified with an element of $U(\mathfrak{q}(V))$; the restriction of this endomorphism on R^λ may be identified with an element of $U(\mathfrak{h})$.

By Theorem 3.3.1 the endomorphism algebra of R^λ is generated by the F_{ii} and to prove the corollary, it suffices to verify it for these elements. We have

$$F_{ii}(\kappa_tv_t) = \kappa_t(F_{ii}v_t) = \kappa_t(p_t^i v_t) = p_t^i \kappa_t(v_t)$$

\square

3.3.3. Corollary . *Let $\varphi : M^\lambda \rightarrow M^\lambda$ be an H_k -module endomorphism given by the formula $\varphi(v_t) = \rho_t \kappa_t(v_t)$. Then $\varphi|_{R^\lambda} = c \in \mathbb{C}$.*

Proof. Let us show that φ commutes with the endomorphisms F_{ii} . Indeed,

$$\begin{aligned}\varphi \cdot F_{ii}(v_t) &= \varphi(p_t^i v_t) = p_t^i \varphi(v_t) = p_t^i \rho_t \kappa_t(v_t); \\ F_{ii} \cdot \varphi(v_t) &= F_{ii} \rho_t \kappa_t(v_t) = \rho_t \kappa_t(F_{ii}v_t) = \\ \rho_t \kappa_t(p_t^i v_t) &= \rho_t p_t^i \kappa_t(v_t) = p_t^i \rho_t \kappa_t(v_t).\end{aligned}$$

The latter identity holds thanks to the fact that p_t^i commutes with ρ_t .

Thus, $\varphi \cdot F_{ii}(v_t) = F_{ii} \cdot \varphi(v_t)$ and, since the elements v_t generate the H_k -module M^λ , we have

$$\varphi \cdot F_{ii} = F_{ii} \cdot \varphi.$$

Since $\varphi(R^\lambda) \subset R^\lambda$ for any endomorphism φ of M^λ , it follows that $\varphi|_{R^\lambda}$ is an element from the centralizer of R^λ . But by Theorem 3.3.1 the centralizer is the Clifford algebra $C_{l(\lambda)}$. But φ is an even central element of $C_{l(\lambda)}$, hence, φ is a constant. \square

3.3.4. Corollary . *Set $e_t = \kappa_t \rho_t$. Then*

$$e_t^2 = c \cdot e_t \text{ for } c \in \mathbb{C}, c \neq 0$$

and the algebra $e_t H_k e_t$ is isomorphic to $C_{l(\lambda)}$ and is generated by the p_t^i for $1 \leq i \leq l(\lambda)$.

Proof. Thanks to Corollary 3.3.3

$$\varphi(\kappa_t(v_t)) = \kappa_t \rho_t \kappa_t(v_t) = c \kappa_t(v_t)$$

or, equivalently,

$$e_t^2 = c \cdot e_t.$$

Since the constant term of $\kappa_t \rho_t$, equal to the coefficient of v_t in $\kappa_t(v_t)$, is nonzero, as follows from the proof of Theorem 3.3.1, then the routine arguments with the help of the bases (cf. [W]) shows that $c \neq 0$. \square

Moreover, the algebra $e_t H_k e_t$ is *anti*-isomorphic to the algebra of H_k -endomorphism of the submodule of M^λ generated by $\kappa_t(v_t)$. Denote the latter module by R_1^λ . Thanks to Corollary 3.3.2 if φ is an endomorphism of R^λ , then $\varphi(R_1^\lambda) \subset R_1^\lambda$ and the restriction map $\varphi|_{\kappa_t(v_t)}$ determines an antiisomorphism of $\text{End}_{H_k}(R^\lambda)$ into the algebra generated by the p_t^i for $1 \leq i \leq l(\lambda)$. But the latter algebra is isomorphic to $C_{l(\lambda)}$; hence, $\varphi|_{\kappa_t(v_t)}$ is an antiisomorphism and $R_1^\lambda = R^\lambda$.

§4. THE SPECHT MODULES OVER \mathfrak{A}_k

In this section we construct analogs of the modules M^λ and the Specht modules R^λ for \mathfrak{A}_k .

First, we need the following statement.

4.1. Lemma . *Let $\pi_1 = \tau_{12}, \dots, \pi_2 = \tau_{13} + \tau_{23}, \dots, \pi_k = \sum_{\alpha < k} \tau_{\alpha k}$ be odd analogs of Jucys–Murphy’s elements. Then*

$$e_k = \prod_{i \geq 2} \frac{2}{i(i-1)} \pi_i^2$$

is an idempotent and $e_k \mathfrak{A}_k e_k$ is isomorphic to the Clifford algebra with $k-1$ generators.

Proof. It is easy to verify by induction that

$$e_k = \frac{1}{k!} \sum_{\alpha \geq 0} (-1)^\alpha 2^{k-\alpha-1} \Sigma_{2\alpha+1},$$

where $\Sigma_{2\alpha+1}$ is the sum of all elements from \mathfrak{A}_k of the form $\tau_{i_1 i_2} \tau_{i_2 i_3} \dots \tau_{i_{2\alpha} i_{2\alpha+1}}$. This implies that e_k is a central element that does not vary under the replacement of the sequence $\pi_i = \sum_{\alpha < i} \tau_{\alpha i}$ with $\pi_{\sigma(i)} = \sum_{\alpha < i} \tau_{\sigma(\alpha) \sigma(i)}$ for any i and $\sigma \in \mathfrak{S}_k^*$. It is not difficult to verify that

$$(\tau_{12} + \tau_{23} + \tau_{31}) \cdot (2 - \tau_{12} \tau_{23} + \tau_{13} \tau_{32}) = 0.$$

Since

$$2 - \tau_{12}\tau_{23} + \tau_{13}\tau_{32} = \pi_2^2 = (\tau_{13} + \tau_{23})^2$$

is a factor in the expression for e_k , it follows that

$$(\tau_{12} + \tau_{23} + \tau_{31})e_k = 0.$$

From symmetry considerations

$$(\tau_{ij} + \tau_{jl} + \tau_{li})e_k = 0 \text{ for any distinct } i, j, l \in \{1, \dots, k\}.$$

Further on,

$$\begin{aligned} \pi_i^2 &= (\sum \tau_{\alpha i})^2 = i - 1 - \sum (\tau_{\alpha\beta}\tau_{\beta i} + \tau_{\beta\alpha}\tau_{\alpha i}) = \\ &= i - 1 - (\sum (1 + \tau_{\alpha\beta}\tau_{\beta i} + \tau_{\beta\alpha}\tau_{\alpha i}) - 1) = \\ &= i - 1 + \frac{1}{2}(i - 1)(i - 2) - \sum \tau_{\alpha\beta}(\tau_{\alpha\beta} + \tau_{\beta i} + \tau_{i\alpha}). \end{aligned}$$

Hence, $\pi_i^2 e_k = \frac{1}{2}(i - 1)(i - 2)e_k$ and, therefore,

$$e_k^2 = \left(\prod_{i \geq 2} \frac{2}{i(i - 1)} \pi_i^2 \right) e_k = e_k.$$

Furthermore, since e_k is a central element, then $e_k \mathfrak{A}_k e_k = \mathfrak{A}_k e_k$. Let I be the ideal in \mathfrak{A}_k generated by the elements $\tau_{ij} + \tau_{jl} + \tau_{li}$; then $e_n I = 0$. Let $\bar{\mathfrak{A}} = \mathfrak{A}_k / I$. In $\bar{\mathfrak{A}}$, then, the following relations hold:

$$\tau_{12} = \pi_2, \tau_{23} = \frac{1}{2}(\pi_3 - \pi_2), \tau_{34} = \frac{1}{3}(\pi_4 - \pi_3), \dots, \tau_{k-1,k} = \frac{1}{k-1}(\pi_k - \pi_{k-1}).$$

Hence, $\bar{\mathfrak{A}}$ is generated by the π_i . Above we showed that $\pi_i^2 = \frac{1}{2}(i - 1)(i - 2)$ in $\bar{\mathfrak{A}}$. So $\bar{\mathfrak{A}}$ is the Clifford algebra generated by the images of the π_i for $2 \leq i \leq k$.

Further on, $(1 - e_k)I = I$; so $\mathfrak{A}_k(1 - e_k) \supset I$, but since $\bar{\mathfrak{A}}$ is a Clifford algebra (and, in particular, is simple), then $\mathfrak{A}_k(1 - e_k) = I$ and, therefore, $\mathfrak{A}_k e_k \cong \mathfrak{A}_k / (1 - e_k)\mathfrak{A}_k \cong \mathfrak{A}_k / I \cong \bar{\mathfrak{A}}$, the Clifford algebra with $k - 1$ generators. \square

Let λ be a strict partition and t a λ -tableau. For $1 \leq i \leq l(\lambda)$ set

$$p_t^i = \sum p_\alpha, \text{ where } \alpha \text{ runs over the entries of the } i\text{-th row of } t$$

and let σ_i be the element from \mathfrak{A}_k constructed as in Lemma 4.1 from the sequence of numbers that stand in the i -th row.

Let V be an (n, n) -dimensional superspace with $n \geq l(\lambda)$ and $W = V^{\otimes k}$, where $k = \sum \lambda_i$. Let M^t be the subspace of W spanned by the vectors $w \in W$ such that

$$E_{ii}w = \lambda_i w; \quad F_{ii}w = p_t^i w \text{ for } 1 \leq i \leq l(\lambda).$$

Set also $\sigma_t = \prod_{1 \leq i \leq l(\lambda)} \sigma_i$; let R^t be the set of highest weight vectors that belong to M^t .

4.2. Theorem . a) As \mathfrak{A}_k -module, M^t is isomorphic to $\mathfrak{A}_k \sigma_t$.

b) The \mathfrak{A}_k -submodule in M^t generated by the $\kappa_{t'} v_{t'}$, where t' has the same rows as t , is an isotypical one and its centralizer is isomorphic to the Clifford algebra $C_{k-l(\lambda)}$.

Proof. Since $\sigma_t v_t = c \cdot v_t$, heading a) is clear.

The facts $v_t \in M^t$ and $\mathfrak{A}_k v_t \subset R^t$ are also clear.

Let us show first of all that the centralizer of R^t is isomorphic to the Clifford algebra $C_{k-l(\lambda)}$.

Obviously, \mathfrak{A}_k and $C_k \otimes U(\mathfrak{g}(V))$ form a Howe-dual pair in W . By Theorem 3.3.1 the centralizer of the H_k -module R^λ is the Clifford algebra generated by the F_{ii} for $1 \leq i \leq l(\lambda)$.

Therefore, the centralizer of the \mathfrak{A}_k -module R^λ is the Clifford algebra $C_{k+l(\lambda)}$ generated by the F_{ii} for $1 \leq i \leq l(\lambda)$ and p_1, \dots, p_k .

The condition $F_{ii}w = p_t^i w$ is equivalent to another one: $ew = w$ for

$$e = \prod_{1 \leq i \leq l(\lambda)} \frac{1}{2} \left(1 - \frac{1}{\lambda_i} p_t^i F_{ii} \right) \text{ and } e^2 = e.$$

Therefore, the centralizer of R^t is isomorphic to $eC_{k+l(\lambda)}e$.

Let $C_{k-l(\lambda)}$ be the subalgebra of C_k generated by the $p_i - p_j$ for i, j that belong to the same row of t . Then it is not difficult to see that $eC_{k+l(\lambda)}e \cong C_{k-l(\lambda)}$ and this proves that the centralizer of R^t is isomorphic to $C_{k-l(\lambda)}$.

Let us prove now that the submodule R_1^t of R^t generated by the elements $\kappa_{t'}v_t$, where t' has, up to permutations, the same rows as t , is isomorphic to R^t . To this end it suffices to prove that any endomorphism of R^t sends R_1^t into itself.

But, indeed, any endomorphism of R^t is a multiplication by $p_i - p_j$ for i, j that belong to the same row of t . Now, it suffices to verify that $(p_i - p_j)\kappa_{t'}v_t \in R_1^t$. But

$$(p_i - p_j)\kappa_{t'}v_t = \sqrt{2}\tau_{ij}s_{ij}\kappa_{t'}v_t = -\sqrt{2}\tau_{ij}\kappa_{s_{ij}(t')}v_{t'}.$$

But the rows of $s_{ij}t'$ consist of the same elements that constitute the rows of t . □

REFERENCES

- [BL] Bernstein J., Leites D., The superalgebra $Q(n)$, the odd trace and the odd determinant. C.R. de l'acad bulg de Sci. v. 35, 1982, n.3, 285–286
- [Ja] James G., *The representation theory of the symmetric groups*, Lect. Notes Math. 682, 1978
- [J] Jucys A., Symmetric polynomials and the center of the symmetric group ring, Report Math. Phys. 5, 1974, 107–112
- [M] Murphy G., A new construction of the Young seminormal representations of the symmetric group, J. Algebra 69, 1981, 287–291
- [N1] Nazarov M., An orthogonal basis in the irreducible projective representations of the symmetric group, Funct. Anal. Appl., 22, 1988, n.2, 66–68
- [N2] Nazarov M., Young's symmetrizers for projective representations of the symmetric group, Adv. Math., 127, 1997, n.2, 190–257
- [Pe] Penkov I., Characters of the typical irreducible representations of finite dimensional $\mathfrak{q}(n)$ -modules (Russian), Funktsional. Anal. i Prilozhen. 20 (1986), n. 1, 37–45
- [P] Pragacz P., Algebro-geometric applications of Schur S - and Q - polynomials, Lect. Notes Math. 1478, 1991, 130–191
- [S1] Sergeev A., The tensor algebra of the standard representation as a module over the Lie superalgebras $\mathfrak{gl}(n, m)$ and $\mathfrak{q}(n)$, Math. Sbornik, 123 (165), 1984, n.3, 422–430
- [S2] Sergeev A., The center of the enveloping algebra for the Lie superalgebra $\mathfrak{q}(n)$, Lett. Math. Phys., 7, 1983, 177–179
- [W] Weyl H., *Classical groups, their invariants and representations*, Princeton Univ. Press, Princeton, 1946

DEPT. OF MATH., UNIV. OF STOCKHOLM, ROSLAGSV. 101, KRÄFTRIKET HUS 6, S-106 91, STOCKHOLM, SWEDEN. ON LEAVE OF ABSENCE FROM BALAKOVO INST. OF TECHNIQUE TECHNOLOGY AND CONTROL, E-MAIL: MLEITES@MATEMATIK.SU.SE (SUBJECT: FOR SERGEEV)