

ad -NILPOTENT IDEALS OF A BOREL SUBALGEBRA III

PAOLA CELLINI
PAOLO PAPI

ABSTRACT. This paper is devoted to a detailed study of certain remarkable posets which form a natural partition of all abelian ideals of a Borel subalgebra. Our main result is a nice uniform formula for the dimension of maximal ideals in these posets. We also obtain results on ad -nilpotent ideals which complete the analysis started in [CP2], [CP3].

§1 INTRODUCTION

In this paper we investigate in detail certain remarkable posets of abelian ideals $\mathcal{I}_{ab}(\tau, \alpha)$ of a Borel subalgebra of a simple Lie algebra. These posets depend upon a long root α and an element τ of the coroot lattice (subject to certain natural restrictions), and form a partition of all abelian ideals of a Borel subalgebra. As a consequence of our analysis we may deduce uniform formulas for the dimensions of maximal and minimal ideals in $\mathcal{I}_{ab}(\tau, \alpha)$. Our work generalizes and refines the work of Panyushev and Suter on similar topics, using as a key tool the combinatorics of affine Weyl groups and root systems. Our formulas also provide a uniform explanation of classical Malcev's results on the maximal dimension of an abelian subalgebra of a simple Lie algebra. In order to explain more precisely our results we need to set the basic framework.

Let \mathfrak{g} be a complex simple Lie algebra of rank n . Let $\mathfrak{b} \subset \mathfrak{g}$ be a fixed Borel subalgebra, with Cartan component \mathfrak{h} ; let Δ^+ be the positive system of the root system Δ of \mathfrak{g} corresponding to the previous choice, and $\Pi = \{\alpha_1, \dots, \alpha_n\}$ be the associated set of simple roots. Let Q, Q^\vee be the root and coroot lattices, and W be the Weyl group of \mathfrak{g} . For each $\alpha \in \Delta^+$ let \mathfrak{g}_α denote the root space of \mathfrak{g} relative to α . Denote by \mathcal{I}_{ab} the set of abelian ideals of \mathfrak{b} .

The class \mathcal{I}_{ab} has been introduced in [K1], but the paper that started the recent research activity on the subject is the long research announcement [K2]. In this paper Kostant established deep connections between the combinatorics of abelian ideals, the machinery of Cartan decompositions and the theory of discrete series (for developments in this direction see also [CMP]). His work is partly based on a simple but surprising result due to D. Peterson (the so-called Peterson's $2^{rk(\mathfrak{g})}$ abelian ideals theorem): the abelian ideals may be encoded and in turn enumerated through elements in a certain subset \mathcal{W}_{ab} of the affine Weyl group \widehat{W} of \mathfrak{g} . In [CP2] we extended this encoding, say $\mathfrak{i} \mapsto w_i$, to the set \mathcal{I} of ideals of \mathfrak{b} contained in $[\mathfrak{b}, \mathfrak{b}]$. These ideals are characterized, among the ideals of \mathfrak{b} , by the following property:

1991 *Mathematics Subject Classification*. Primary: 17B20; Secondary: 20F55.

Key words and phrases. Borel subalgebra, abelian ideal, Lie algebra, affine Weyl group.

for any $x \in \mathfrak{i}$, $ad(x) : \mathfrak{g} \rightarrow \mathfrak{g}$ is a nilpotent endomorphism, hence we call them *ad-nilpotent ideals*. Moreover we gave a characterization of the image of the map $\mathfrak{i} \mapsto w_{\mathfrak{i}}$ inside \widehat{W} . Reformulating this characterization in terms of the affine representation of \widehat{W} allowed us to describe a suitable simplex D in $\mathfrak{h}_{\mathbb{R}}^*$ whose Q^\vee -points parametrize \mathcal{I} (see [CP3]). In such a way we obtained a bijection between \mathcal{I} and certain W -orbits in the coroot lattice of \mathfrak{g} , which affords a uniform enumeration of \mathcal{I} . A short part of section 3 of this paper is devoted to give results on \mathcal{I} which complete the analysis started in our previous papers and which have been announced in [CP3]: we characterize the ideals in \mathcal{I} having a fixed class of nilpotence in terms of elements of $D \cap Q^\vee$ and the generators of $\mathfrak{i} \in \mathcal{I}$ as a \mathfrak{b} -module in terms of $w_{\mathfrak{i}}$.

The remaining and more extended part of the paper deals with abelian ideals. Recall from [CP2] that if $\mathfrak{i} \in \mathcal{I}_{ab}$ and $w_{\mathfrak{i}} = t_\tau v$, where t_τ is the translation by $\tau \in Q^\vee$, and $v \in W$, then τ belongs to set X of Q^\vee -points lying in twice the closure of the fundamental alcove of \widehat{W} . We provide a general description of the elements of X as orthogonal sums of highest roots of nested standard irreducible parabolic subsystems of Δ . This result turns out to be a very useful technical device for our goals.

We begin our analysis decomposing \mathcal{I}_{ab} w.r.t. X : the map $\mathcal{W}_{ab} \rightarrow X$, $t_\tau v \mapsto \tau$ is surjective, thus we have a partition into non-empty sets

$$\mathcal{I}_{ab} = \bigcup_{\tau \in X} \mathcal{I}_{ab}(\tau),$$

$\mathcal{I}_{ab}(\tau)$ being the set of ideals \mathfrak{i} whose $w_{\mathfrak{i}}$ has t_τ as translational part. In [K2] and [CMP] it is shown that $\mathcal{I}_{ab}(\tau)$ is in canonical bijection with $W_\tau \backslash W$, where, if \mathfrak{k}_τ is the equal rank symmetric subalgebra of \mathfrak{g} formed by the fixed points of $Ad(\exp(\sqrt{-1}\pi\tau))$, W_τ denotes the Weyl group of $(\mathfrak{k}_\tau, \mathfrak{h})$. In the following discussion we list our main results.

We prove that, for all $\tau \in X$, $\mathcal{I}_{ab}(\tau)$ has a unique minimal element; moreover we describe explicitly the maximal elements of $\mathcal{I}_{ab}(\tau)$. If $\tau = 0$, then $\mathcal{I}_{ab}(\tau)$ consists just of the zero ideal, so assume $\tau \neq 0$. If \mathfrak{i} is maximal in $\mathcal{I}_{ab}(\tau)$, then $w_{\mathfrak{i}}^{-1}(-\theta + 2\delta)$ is a long simple root (here θ is the highest root of Δ and δ is the fundamental imaginary root of the affine system $\widehat{\Delta}$ attached to Δ). So we are led to consider, for any α in the set Π_ℓ of long simple roots, the subposets

$$\mathcal{I}_{ab}(\tau, \alpha) = \{\mathfrak{i} \in \mathcal{I}_{ab} \mid w_{\mathfrak{i}} = t_\tau v, w_{\mathfrak{i}}^{-1}(-\theta + 2\delta) = \alpha\}.$$

For any $\mathfrak{i} \in \mathcal{I}_{ab}(\tau)$, $w_{\mathfrak{i}}^{-1}(-\theta + 2\delta)$ belongs to Δ and is a long positive root. We denote by Δ_ℓ^+ the set of long positive roots in Δ and extend the definition of $\mathcal{I}_{ab}(\tau, \alpha)$ to all α in Δ_ℓ^+ . Thus we have

$$\mathcal{I}_{ab}(\tau) = \bigcup_{\alpha \in \Delta_\ell^+} \mathcal{I}_{ab}(\tau, \alpha).$$

$\mathcal{I}_{ab}(\tau, \alpha)$ is possibly empty; we give an explicit criterion for $\mathcal{I}_{ab}(\tau, \alpha)$ to be non-empty. We prove that if $\mathcal{I}_{ab}(\tau, \alpha) \neq \emptyset$, then it has unique minimal and maximal elements. More precisely, we prove that the poset $\mathcal{I}_{ab}(\tau, \alpha)$ is order isomorphic to the coset space $W((\Pi \cap \alpha^\perp) \backslash P_{\tau, \alpha}) \backslash W(\Pi \cap \alpha^\perp)$ (see 7.6 for the precise definition of $P_{\tau, \alpha}$

and 2.8, 2.9 for the definition of the poset of cosets.) This result is a generalization of [S, Theorem 11]. We explicitly describe elements $w_\tau, y_{\tau,\alpha}, \check{y}_{\tau,\alpha} \in \widehat{W}$ such that

$$i(w_\tau) = \min \mathcal{I}_{ab}(\tau), \quad i(w_\tau y_{\tau,\alpha}) = \min \mathcal{I}_{ab}(\tau, \alpha), \quad i(w_\tau \check{y}_{\tau,\alpha}) = \max \mathcal{I}_{ab}(\tau, \alpha).$$

The ideals $\max \mathcal{I}_{ab}(\tau, \alpha)$ for all simple roots α such that $\mathcal{I}_{ab}(\tau, \alpha) \neq \emptyset$ are exactly the maximal elements of $\mathcal{I}_{ab}(\tau)$.

For $\alpha \in \Delta_\ell^+$ set $X_\alpha = \{\tau \in X \mid \mathcal{I}_{ab}(\tau, \alpha) \neq \emptyset\}$. Then X_α is chain in Q^\vee , in the standard partial order, and its minimal element is θ^\vee . Assume $X_\alpha = \{\tau_1, \dots, \tau_{k(\alpha)}\}$, $\theta^\vee = \tau_1 < \dots < \tau_{k(\alpha)}$, and denote by $\mathfrak{m}_h(\alpha)$ the maximal ideal in $\mathcal{I}_{ab}(\tau_h, \alpha)$. Then a careful analysis of the elements of \widehat{W} corresponding to the ideals $\mathfrak{m}_h(\alpha)$ allows us to prove that

$$\mathfrak{m}_1(\alpha) \subset \dots \subset \mathfrak{m}_{k(\alpha)}(\alpha).$$

The proof of these inclusion relations gives a clear explanation of the “shell-like” structure of the elements of \mathcal{W}_{ab} noted in [S, §5]. Moreover it gives as a by-product a case-free proof of the following nice result, due to Suter [S, Corollary 13], which holds for the maximal abelian ideals $\mathfrak{m}(\alpha) = \mathfrak{m}_{k(\alpha)}(\alpha)$, $\alpha \in \Pi_\ell$ of \mathfrak{b} :

$$\dim(\mathfrak{m}(\alpha)) = g - 1 + \frac{1}{2} \left(|\widehat{\Delta}_\alpha| - |\Delta_\alpha| \right)$$

where $\widehat{\Delta}_\alpha, \Delta_\alpha$ are the (finite) subsystems generated by the simple roots orthogonal to α in $\widehat{\Delta}, \Delta$ respectively, and g is the dual Coxeter number of Δ .

In particular we obtain a uniform version of Malcev’s formulas for the dimension of maximal commutative subalgebras of \mathfrak{g} : if d denotes the dimension of such a subalgebra, we have

$$d = \dim \mathfrak{m}_{k(\bar{\alpha})}(\bar{\alpha}).$$

where $\bar{\alpha}$ is a long simple root having maximal distance from the affine node in the extended Dynkin diagram of Δ .

Then we show that, when $\alpha \in \Pi_\ell$, $\mathfrak{m}_{k(\alpha)}(\alpha)$ is a maximal abelian ideal of \mathfrak{b} and that each maximal abelian ideal of \mathfrak{b} occurs in this way for a unique $\alpha \in \Pi_\ell$. Hence the map $\alpha \mapsto \mathfrak{m}_{k(\alpha)}(\alpha)$ is a bijection between Π_ℓ and maximal ideals in \mathcal{I}_{ab} ; the same bijection has been shown via a case by case verification in [PR] and in a uniform way in [P, Corollary 3.8].

Finally, we find a general dimension formula for all the ideals $\mathfrak{m}_h(\alpha)$. In the above notation, to each pair (α, h) , $1 \leq h < k(\alpha)$, $\alpha \in \Delta_\ell^+$ we associate a finite irreducible subsystem $\Delta_{h,\alpha}$ of $\widehat{\Delta}$ and we prove that

$$\dim(\mathfrak{m}_h(\alpha)) = g - 1 + \sum_{i=1}^{h-1} (g_h(\alpha) - 1)$$

where $g_h(\alpha)$ is the dual Coxeter number of $\Delta_{h,\alpha}$.

The paper is organized as follows. In section 2 we recall some basic facts about affine root systems and Coxeter groups. In section 3 we recall our encoding of ad-nilpotent ideals of \mathfrak{g} via the affine Weyl group and we prove the two results on \mathcal{I}

stated above. In section 4 we characterize the maximal abelian ideals of \mathfrak{b} in terms of the corresponding elements in \widehat{W} . In section 5 we give a structure theorem for the elements of X . In section 6 we introduce $\mathcal{I}_{ab}(\tau)$, we describe its unique minimal element and begin the study of the maximal elements, proving that they should belong to $\mathcal{I}_{ab}(\tau, \alpha)$ with $\alpha \in \Pi_\ell$. Section 7 is devoted to the proof of the existence of unique minimal and maximal elements in $\mathcal{I}_{ab}(\tau, \alpha)$. In section 8 we give the explicit description of the maximal elements $\mathfrak{m}_h(\alpha)$ which leads to our dimension formulas. The last section deals with the explicit computation of $\dim(\mathfrak{m}_h(\alpha))$ for any admissible pair (h, α) , $\alpha \in \Pi_\ell$.

Notational conventions. If R is a root system we shall denote by $\Delta^+(R)$, $Q(R)$, $Q^+(R)$, $W(R)$ the positive roots, the root lattice, the positive root lattice and the Weyl group of R , respectively. If Y is a subset of R , we shall also use the notation $\Delta(Y)$, and $W(Y)$, to denote the subsystem generated by Y , and its Weyl group, respectively. If R is irreducible, we set $\theta(R)$ to be its highest root. We reserve the symbols Δ^+ , Q , Q^+ , θ , W for the same data relative to the root system Δ of \mathfrak{g} .

If R is simply laced we shall regard all roots as long.

The maximal irreducible subsystems of a root system R will be called the *components* of R .

For $\alpha \in Q$, $\alpha = \sum_{i=1}^n \lambda_i \alpha_i$ we set $\text{Supp}(\alpha) = \{\alpha_i \mid \lambda_i \neq 0\}$.

We shall mainly use two partial orderings throughout the paper, both denoted by \leq : we shall endow Q (or Q^\vee) with the usual order: $\alpha < \beta$ if $\beta - \alpha$ is a sum of positive roots (resp. coroots). Further we shall regard W and \widehat{W} as posets via the weak order (more precisely the weak left Bruhat order with respect to the standard set of Coxeter generators, see [H, 5.9]). Subsets of W or \widehat{W} will be endowed with the restriction of \leq . Only in one instance we shall use the Bruhat order: it will be denoted by $<_B$.

We shall denote by $A \setminus B$ the set difference of A and B , and by $A + B$ the symmetric difference of the sets A , B . For $k \in \mathbb{N}^+$, we set $[k] = \{1, \dots, k\}$.

§2 AFFINE ROOT SYSTEMS AND COMBINATORICS OF ROOTS

Our basic tools are the standard theory of affine Weyl groups and some combinatorial properties of root systems. This section is devoted to recollect these topics.

We set $V \equiv \mathfrak{h}_{\mathbb{R}}^* = \bigoplus_{i=1}^n \mathbb{R}\alpha_i$ and denote by $(\ , \)$ the positive symmetric bilinear form induced on V by the Killing form. The affine root system associated to Δ can be introduced as follows [Kac, Chapter 6]. We extend V and its inner product setting $\widehat{V} = V \oplus \mathbb{R}\delta \oplus \mathbb{R}\lambda$, $(\delta, \delta) = (\delta, V) = (\lambda, \lambda) = (\lambda, V) = 0$, and $(\delta, \lambda) = 1$. We still denote by $(\ , \)$ the resulting (non-degenerate) bilinear form. The affine root system associated to Δ is $\widehat{\Delta} = \Delta + \mathbb{Z}\delta = \{\alpha + k\delta \mid \alpha \in \Delta, k \in \mathbb{Z}\}$; remark that the affine roots are non isotropic vectors. The set of positive affine roots is $\widehat{\Delta}^+ = (\Delta^+ + \mathbb{N}\delta) \cup (\Delta^- + \mathbb{N}^+\delta)$, where $\Delta^- = -\Delta^+$. We denote by θ the highest root of Δ and set $\alpha_0 = -\theta + \delta$, $\widehat{\Pi} = \{\alpha_0, \dots, \alpha_n\}$. We sometimes call the roots of Δ inside $\widehat{\Delta}$ the “finite roots”. For each $\alpha \in \widehat{\Delta}^+$ we denote by s_α the corresponding reflection of \widehat{V} , $s_\alpha(x) = x - \frac{2(\alpha, x)}{(\alpha, \alpha)}x$ for $x \in \widehat{V}$. The affine Weyl group associated to Δ is the group \widehat{W} generated by $\{s_\alpha \mid \alpha \in \widehat{\Delta}^+\}$; the set $\{s_\alpha \mid \alpha \in \widehat{\Pi}\}$ is a set of

Coxeter generators for \widehat{W} ; we denote by ℓ the corresponding length function. Note that $w(\delta) = \delta$ for any $w \in \widehat{W}$. It turns out that \widehat{W} is a semidirect product $T \rtimes W$, where $T = \{t_\tau \mid \tau \in Q^\vee\} \cong Q^\vee$ is the subgroup of *translations*, and the action of W on T is $vt_\tau v^{-1} = t_{v(\tau)}$. The action of t_τ on $V \oplus \mathbb{R}\delta$, in particular on the roots, is given by $t_\tau(x) = x - (x, \tau)\delta$, for each $x \in V \oplus \mathbb{R}\delta$. (See [Kac] for the general definition of t_τ on \widehat{V}).

Now consider the \widehat{W} -invariant affine subspace $E = \{x \in \widehat{V} \mid (x, \delta) = 1\}$. Remark that $E = V \oplus \mathbb{R}\delta + \lambda$; let $\pi : E \rightarrow V$ be the natural projection. For $w \in \widehat{W}$ we set $\overline{w} = \pi \circ w|_E$. We have that the map $w \mapsto \overline{w}$ gives an isomorphism of \widehat{W} onto a group W_{af} of affine transformations of V [Kac, 6.6]. Under this isomorphism each element t_τ of T maps to the true translation by τ . So W_{af} is the usual affine representation of the affine Weyl group [B, VI, §2]. In fact, for $\alpha \in \Delta^+$ and $k \in \mathbb{Z}$ set

$$H_{\alpha,k} = \{x \in V \mid (x, \alpha) = k\}$$

and let $s_{\alpha,k}$ denote the orthogonal reflection with respect to $H_{\alpha,k}$ in V . For $\beta \in \widehat{V}$, let \widehat{H}_β be the hyperplane orthogonal to β in \widehat{V} . Then it is easy to see that, for each $\alpha \in \Delta^+$, $k > 0$ and $h \geq 0$, we have

$$\begin{aligned} \pi(\widehat{H}_{-\alpha+k\delta} \cap E) &= H_{\alpha,k}, & \overline{s_{-\alpha+k\delta}} &= s_{\alpha,k}, \\ \pi(\widehat{H}_{\alpha+h\delta} \cap E) &= H_{\alpha,-h}, & \overline{s_{\alpha+h\delta}} &= s_{\alpha,-h}. \end{aligned}$$

Hence we have a transparent correspondence between the linear and the affine context. Henceforward we shall omit the bar and use the same notation for the elements in \widehat{W} and the corresponding elements W_{af} . We denote by C is the fundamental chamber of \widehat{W} and by C_1 the fundamental alcove of W_{af} ,

$$\begin{aligned} C &= \{x \in \widehat{V} \mid (x, \alpha_i) > 0 \text{ for } i = 0, \dots, n\}, \\ C_1 &= \{x \in V \mid (x, \alpha) > 0 \forall \alpha \in \Pi, (x, \theta) < 1\} \\ &= \{x \in V \mid 0 < (x, \alpha) < 1 \forall \alpha \in \Delta^+\}. \end{aligned}$$

Then we have $C_1 = \pi(C \cap E)$. This implies the following result.

2.1. For any $\alpha \in \Delta^+$, $k \in \mathbb{N}^+$, $h \in \mathbb{N}$

$$\begin{aligned} w^{-1}(-\alpha + k\delta) < 0 &\text{ if and only if } H_{\alpha,k} \text{ separates } C_1 \text{ and } w(C_1), \\ w^{-1}(\alpha + h\delta) < 0 &\text{ if and only if } H_{\alpha,-h} \text{ separates } C_1 \text{ and } w(C_1). \end{aligned}$$

2.2. For $j > 0$ set

$$C_j = \{x \in V \mid (x, \alpha) > 0 \forall \alpha \in \Pi, (x, \theta) < j\};$$

also set

$$C_\infty = \{x \in V \mid (x, \alpha_i) > 0 \text{ for } i = 1, \dots, n\}.$$

Thus C_∞ is the fundamental chamber of W . The walls of C_j are the hyperplanes, $H_{\alpha_i,0}$, $1 \leq i \leq n$ and $H_{\theta,j}$, so that $C_j = jC_1$. For all $\beta \in \Delta^+$ and $x \in C_\infty$ we have $(x, \beta) \leq (x, \theta)$. In particular we obtain that if $H_{\beta,k} \cap C_2 \neq \emptyset$, then $k = 1$.

2.3. Let R be a finite or affine root system. Choose the set R^+ of positive roots, and denote by R_Σ the corresponding set of simple roots. Let G be the Weyl group of R ; for $\alpha \in R^+$ we denote by s_α the reflection with respect to α . For $v \in G$ we set

$$N(v) = \{\beta \in R^+ \mid v^{-1}(\beta) < 0\}.$$

Moreover, we denote by L_v the set of left descents of v :

$$L_v = \{\alpha \in R_\Sigma \mid \ell(s_\alpha v) < \ell(v)\}.$$

Similarly we define the set of *right descents* of v as $R_v = \{\alpha \in R_\Sigma \mid \ell(vs_\alpha) < \ell(v)\}$.

The following facts are well known; proofs and details about (3) and (4) can be found in [D], [CP1]. Item (5) follows from (3).

- (1) $L_v = N(v) \cap R_\Sigma$;
- (2) $\alpha \in L_v$ if and only if there exists a reduced expression of v starting with s_α .
- (3) $N(v)$ is finite, closed under root addition and, moreover, if $\beta_1, \dots, \beta_k \in R^+$, $c_1, \dots, c_k \in \mathbb{R}^+$, and $c_1\beta_1 + \dots + c_k\beta_k \in N(v)$, then $\beta_i \in N(v)$ for at least one $i \in \{1, \dots, k\}$.
- (4) Conversely, if $R \not\cong \widehat{A}_1$, and $N \subseteq R^+$ is a finite set which satisfies the following conditions:
 - (a) N is closed under root addition,
 - (b) for all $\beta_1, \beta_2 \in R^+$ such that $\beta_1 + \beta_2 \in N$, at least one of β_1, β_2 belongs to N ,

then there exists $v \in W(R)$ such that $N = N(v)$. If $R \cong \widehat{A}_1$, then the sets $N(v)$, $v \in W(R)$, are exactly all the sets $\{-\alpha + i\delta \mid 1 \leq i \leq k\}$, for all $k \in \mathbb{N}^+$, and $\{\alpha + i\delta \mid 0 \leq i \leq k\}$, for all $k \in \mathbb{N}$, where α is the finite simple root.

- (5) Let $I \subseteq R_\Sigma$. Then $L_v \subseteq I$ if and only if for all $\alpha \in N(v)$ we have $\text{Supp}(\alpha) \cap I \neq \emptyset$.

We describe explicitly the set $N(v)$. Part (1) of the following statement is clear; part (2) is well-known [B, VI, 1.6].

2.4. Let $w \in G$ and let $w = \sigma_1 \cdots \sigma_k$ be any reduced expression of w with $\sigma_i = s_{\beta_i}$, $\beta_i \in R_\Sigma$. If we set $\gamma_1 = \beta_1, \gamma_2 = \sigma_1(\beta_2), \dots, \gamma_k = \sigma_1 \cdots \sigma_{k-1}(\beta_k)$, then we have:

- (1) $w = s_{\gamma_k} \cdots s_{\gamma_2} s_{\gamma_1}$.
- (2) $N(w) = \{\gamma_1, \dots, \gamma_k\}$.

2.5 We also set $\overline{N}(v) = N(v) \cup -N(v)$. It is easy to see that for any $u, v \in G$ we have

$$\overline{N}(vu) = \overline{N}(v) + v\overline{N}(u),$$

where we denote by $+$ the symmetric difference of sets.

We recall that the weak order \leq of G is defined as follows: for $v, v' \in G$, we have $v \leq v'$ if and only if there exists an element $u \in G$ such that $v' = vu$, $\ell(v') = \ell(v) + \ell(u)$. From 2.4 and 2.5 we easily obtain the following characterization of the weak order.

2.6. $v \leq v'$ if and only if $N(v) \subseteq N(v')$.

Moreover, we have the following results.

2.7. The following conditions (1), (2), (3) and (4) are equivalent:

(1) $N(vu) \supseteq N(v)$; (2) $vN(u) \subseteq \Delta^+$; (3) $\ell(vu) = \ell(v) + \ell(u)$; (4) $v \leq vu$.

Similarly, the following conditions (1'), (2'), (3') and (4') are equivalent:

(1') $N(vu) \subseteq N(v)$; (2') $vN(u) \subseteq \Delta^-$; (3') $\ell(vu) = \ell(v) - \ell(u)$; (4') $v \geq vu$.

Let G' be a standard parabolic subgroup of G , and let R' be its root system. Recall that any $v \in G$ can be decomposed as $v = v'v''$ where $v' \in G'$ and v'' is an element of minimal length in the right coset $G'v$; moreover, v' , v'' are unique. It follows that

2.8. For all $v \in G$ there exists $v' \in G'$ such that $N(v) \cap R' = N(v')$.

Set moreover $R'' = R \setminus R'$. It is well known that $y \in G'x$ is the minimal length representative of the right coset $G'x$ if and only if $L_y \subseteq R''$. Therefore, $\{y \in G \mid L_y \subseteq R''\}$ is the set of minimal length representatives for the set of right cosets $G' \setminus G$. Thus the restriction of the weak order of G on $\{y \in G \mid L_y \subseteq R''\}$ induces a partial ordering on $G' \setminus G$. When saying *the poset* $G' \setminus G$, we shall always refer to this ordering.

2.9. The poset $G' \setminus G$ has a unique minimal and a unique maximal element. The identity 1 of G clearly corresponds to the minimum of $G' \setminus G$. If w_0 is the longest element of G and w'_0 is the longest element of G' , then we have that $N(w'_0 w_0) = \Delta^+(R) \setminus \Delta^+(R')$. By 2.3 (5), if $y \in G$ is such that $L_y \subseteq R''$, then $N(y) \subseteq \Delta^+(R) \setminus \Delta^+(R')$; therefore we obtain that $w'_0 w_0$ is the unique maximal element in $\{y \in G \mid L_y \subseteq R''\}$, with respect to the weak order. Thus $w'_0 w_0$ corresponds to the unique maximal element of $G' \setminus G$.

§3 ad-NILPOTENT IDEALS VIA AFFINE WEYL GROUPS

We turn now to *ad*-nilpotent ideals. Recall from [CP2, §2] that to any *ad*-nilpotent ideal $\mathfrak{i} = \sum_{\alpha \in \Phi} \mathfrak{g}_\alpha$ we can associate a set of positive roots in the affine root system attached to \mathfrak{g} in the following way:

$$L(\mathfrak{i}) = \bigcup_{k \geq 1} (-\Phi_{\mathfrak{i}}^k + k\delta),$$

where $\Phi_{\mathfrak{i}}^1 = \Phi$ and $\Phi_{\mathfrak{i}}^k = (\Phi_{\mathfrak{i}}^{k-1} + \Phi) \cap \Delta$, $k \geq 2$. In [CP2, Prop. 2.4] we proved that there exists a (unique) $w_{\mathfrak{i}} \in \widehat{W}$ such that $N(w_{\mathfrak{i}}) = L(\mathfrak{i})$. Set

$$\mathcal{W} = \{w_{\mathfrak{i}} \mid \mathfrak{i} \in \mathcal{I}\}, \quad \mathcal{W}_{ab} = \{w_{\mathfrak{i}} \mid \mathfrak{i} \in \mathcal{I}_{ab}\},$$

so that $\mathfrak{i} \mapsto w_{\mathfrak{i}}$ is a bijection $\mathcal{I} \rightarrow \mathcal{W}$, restricting to a bijection $\mathcal{I}_{ab} \rightarrow \mathcal{W}_{ab}$. The inverse map will be denoted by $w \mapsto \mathfrak{i}(w)$.

The sets \mathcal{W} , \mathcal{W}_{ab} have been characterized in our previous papers.

Proposition 3.1. (1) [CP3, Prop. 2] *Let $w \in \widehat{W}$, $w = t_\tau v$, $\tau \in Q^\vee$, $v \in W$. Then $w \in \mathcal{W}$ if and only if the following conditions hold:*

- (i) $w(C_1) \subset C_\infty$;
 - (ii) $(v^{-1}(\tau), \alpha_i) \leq 1$ for each $i \in [n]$ and $(\tau, v(\theta)) \geq -2$.
- (2) [CP2, Th. 2.9] $w \in \mathcal{W}_{ab}$ if and only if $w(C_1) \subset C_2$.

An obvious but useful observation is the following remark (see 2.6).

3.2. *For $\mathfrak{i}, \mathfrak{j} \in \mathcal{I}$, the following conditions are equivalent:*

- (1) $\mathfrak{i} \subseteq \mathfrak{j}$;
- (2) $N(w_{\mathfrak{i}}) \subseteq N(w_{\mathfrak{j}})$;
- (3) $w_{\mathfrak{i}} \leq w_{\mathfrak{j}}$.

Finally, we remark that the construction given at the beginning of the section is quite simple in the special case of abelian ideals. Indeed, we have the following result.

Proposition 3.3. (Peterson, see also [CP2, section 2]) *Let $\mathfrak{i} \in \mathcal{I}_{ab}$. Then $N(w_{\mathfrak{i}}) = \{-\alpha + \delta \mid \mathfrak{g}_\alpha \subseteq \mathfrak{i}\}$. In particular, $\dim(\mathfrak{i}) = \ell(w_{\mathfrak{i}})$.*

Conversely, assume that $w \in \widehat{W}$ is such that $N(w) \subseteq \Delta^- + \delta$ and set $\mathfrak{i} = \sum_{-\alpha + \delta \in N(w)} \mathfrak{g}_\alpha$. Then $\mathfrak{i} \in \mathcal{I}_{ab}$.

The next results complete the analysis developed in [CP2], [CP3], showing once again that all the combinatorial information regarding *ad*-nilpotent ideals can be recovered from $w_{\mathfrak{i}}$.

Remark that $\sum_{\alpha \in \Phi} \mathfrak{g}_\alpha \in \mathcal{I}$ if and only if Φ is a dual order ideal of Δ^+ endowed with the usual partial order, i.e., if and only if, for all $\alpha \in \Phi$ and $\beta \in \Delta^+$ such that $\alpha \leq \beta$, we have that $\beta \in \Phi$.

For all $\mathfrak{i} \in \mathcal{I}$ we set

$$\Phi_{\mathfrak{i}} = \{\beta \in \Delta^+ \mid \mathfrak{g}_\beta \subseteq \mathfrak{i}\},$$

and denote by $\mathcal{A}_{\mathfrak{i}}$ the set of minimal elements of $\Phi_{\mathfrak{i}}$. It is clear that $\mathcal{A}_{\mathfrak{i}}$ is an antichain of Δ^+ , i.e. a set of incomparable elements. As explained in [CP3, §4 (1)], $\Phi_{\mathfrak{i}}$ is completely determined by $\mathcal{A}_{\mathfrak{i}}$.

Proposition 3.4. *For $\mathfrak{i} \in \mathcal{I}$ the following statements are equivalent:*

- (1) $w_{\mathfrak{i}}(\alpha_i) = \beta - \delta$, $\beta \in \Delta^+$ for some i , $0 \leq i \leq n$;
- (2) $\beta \in \mathcal{A}_{\mathfrak{i}}$.

Proof. (1) \implies (2). We have $w_{\mathfrak{i}}(\alpha_i) = \beta - \delta < 0$; hence, $w_{\mathfrak{i}}s_i(\alpha_i) = -\beta + \delta > 0$. By 2.7, it follows that $N(w_{\mathfrak{i}}) = N(w_{\mathfrak{i}}s_i) \cup \{-\beta + \delta\}$. If there exists $\gamma \in \Phi_{\mathfrak{i}}$ such that $\beta > \gamma$, then $-\gamma + \delta$ is equal to $-\beta + \delta$ plus a sum of roots in Δ^+ . But $-\gamma + \delta \in N(w_{\mathfrak{i}})$, hence $-\gamma + \delta \in N(w_{\mathfrak{i}}s_i)$, and since $-\beta + \delta \notin N(w_{\mathfrak{i}}s_i)$, by 2.3 (3) we obtain that some root in Δ^+ belongs to $N(w_{\mathfrak{i}}s_i)$. This is not possible, since $N(w_{\mathfrak{i}}s_i) \subseteq N(w_{\mathfrak{i}}) \subseteq \Delta^- + \mathbb{N}^+\delta$.

(2) \implies (1). Let $\beta \in \Phi_{\mathfrak{i}}$ be minimal. We shall prove that $L(\mathfrak{i}) \setminus \{-\beta + \delta\}$ verifies conditions (a) and (b) of 2.3 (4). This will imply that there exists $w \in \widehat{W}$ such that $L(\mathfrak{i}) \setminus \{-\beta + \delta\} = N(w)$; hence, by 2.6 and 2.5, $w_{\mathfrak{i}} = ws_i$ for some $i \in \{0, \dots, n\}$ and

$w_i(\alpha_i) = -w(\alpha_i) = \beta - \delta$. Set $N = L(\mathfrak{i}) \setminus \{-\beta + \delta\}$. Since $N \subseteq \Delta^- + \mathbb{N}^+\delta$, no pair of roots in N sum up to $-\beta + \delta$; since, moreover, $L(\mathfrak{i})$ is closed under root addition we obtain that N is closed, too. It remains to prove that (b) holds. Assume $\eta \in N$ and $\eta = \eta_1 + \eta_2$, with $\eta_1, \eta_2 \in \widehat{\Delta}^+$. If $\eta_1, \eta_2 \neq -\beta + \delta$, we are done since (b) holds for $L(\mathfrak{i})$. We assume that $\eta_1 = -\beta + \delta$ and prove that then $\eta_2 \in N$. Set $\eta = -\bar{\eta} + k\delta$, $k > 0$, $\bar{\eta} \in \Delta^+$. We prove the claim by induction on k . First notice that $\eta_2 \notin \Delta^+ + \mathbb{N}\delta$, since, if $\eta_2 = \alpha + (k-1)\delta$ with $\alpha \in \Delta^+$, then $-\bar{\eta} = -\beta + \alpha$, which implies $\beta > \bar{\eta}$ against the minimality of β in Φ_i . So we may assume $\eta_2 = -\alpha + (k-1)\delta$, $k > 1$, $\alpha \in \Delta^+$. We first assume $k = 2$, so that $\eta = (-\beta + \delta) + (-\alpha + \delta)$. By assumption $\bar{\eta} \in \Phi_i^2$, hence $\bar{\eta} = \beta' + \alpha'$ with $\beta', \alpha' \in \Phi_i$. Moreover we have that $\bar{\eta} = \beta + \alpha$, hence $0 < (\alpha' + \beta', \alpha + \beta) = (\alpha', \alpha) + (\alpha', \beta) + (\beta', \alpha) + (\beta', \beta)$. It follows that at least one of the scalar products in the right hand side of the above identity is positive. If $(\alpha', \alpha) > 0$, then $\alpha' - \alpha \in \Delta$, and since $\beta = \beta' + (\alpha' - \alpha)$, we obtain that $\alpha' - \alpha < 0$, by the minimality of β in Φ_i . Since $-\alpha' + \delta = -\alpha + \delta + (\alpha - \alpha')$, $-\alpha' + \delta \in L(\mathfrak{i})$, $\alpha - \alpha' \in \Delta^+ \setminus L(\mathfrak{i})$, and $L(\mathfrak{i})$ has property (b), we obtain that $-\alpha + \delta \in L(\mathfrak{i})$, hence $-\alpha + \delta \in N$. If $(\beta', \alpha) > 0$, we obtain similarly that $-\alpha + \delta \in N$. So we assume that $(\alpha', \beta) > 0$. Then by minimality of β we have that $\alpha' - \beta \in \Delta^+$, hence $\alpha - \beta' = \alpha' - \beta \in \Delta^+$ and we can argue as in the previous case, so obtaining that $-\alpha + \delta \in N$. The case $(\beta', \beta) > 0$ is similar. Thus we have proved the claim for $k = 2$. Now we assume $k > 2$ and the claim true for $k - 1$. By assumption we have $\bar{\eta} = \beta' + \alpha'$ with $\alpha' \in \Phi_i^{k-1}$ and $\beta' \in \Phi_i$. Arguing as above, we find that at least one of (α', α) , (α', β) , (β', α) , (β', β) , is positive. If $(\alpha', \alpha) > 0$, then by minimality of β we obtain that $\alpha - \alpha' > 0$; moreover $-\alpha' + (k-1)\delta = -\alpha + (k-1)\delta + (\alpha - \alpha')$, and since $-\alpha' + (k-1)\delta \in L(\mathfrak{i})$ and $\alpha - \alpha' \notin L(\mathfrak{i})$ we obtain that $-\alpha + (k-1)\delta \in L(\mathfrak{i})$, hence $-\alpha + (k-1)\delta \in N$. If $(\beta', \beta) > 0$, we obtain as above that $\alpha - \alpha' > 0 \in \Delta^+$, and we are reduced to the previous case. So we assume that $(\alpha', \beta) > 0$. As above, $\alpha' - \beta > 0$, hence there exists $\gamma \in \Delta^+$ such that $\alpha' = \beta + \gamma$ and $\alpha = \beta' + \gamma$. Hence we have that $-\alpha' + (k-1)\delta = (-\beta + \delta) + (-\gamma + (k-2)\delta)$. By the induction assumption we obtain that $-\gamma + (k-2)\delta \in L(\mathfrak{i})$, hence $-\alpha + (k-1)\delta = (-\beta' + \delta) + (-\gamma + (k-2)\delta) \in L(\mathfrak{i})$ and therefore $-\alpha + (k-1)\delta \in N$. The case $(\beta', \alpha) > 0$ is similar. \square

For each $\beta \in \mathcal{A}_i$, let x_β be any non zero element in \mathfrak{g}_β . Then $\{x_\beta \mid \beta \in \mathcal{A}_i\}$ is a minimal set of generators of \mathfrak{i} as a \mathfrak{b} -module. Hence Proposition 3.4, together with [CP2, Prop. 2.12, ii], imply the following result.

Corollary 3.5. *For any $\mathfrak{i} \in \mathcal{I}$, the minimal number of generators of \mathfrak{i} as a \mathfrak{b} -module equals the number of right descents of w_i and the cardinality of the antichain \mathcal{A}_i .*

To proceed further, we need to recall one of the main results from [CP3]. Set

$$D = \{\sigma \in Q^\vee \mid (\sigma, \alpha_i) \leq 1 \text{ for each } i \in \{1, \dots, n\} \text{ and } (\sigma, \theta) \geq -2\}.$$

Set moreover

$$D_{ab} = \{\sigma \in Q^\vee \mid (\sigma, \alpha) \in \{0, 1, -1, -2\} \text{ for each } \alpha \in \Delta^+\}.$$

It is clear that $D_{ab} \subseteq D$. The first part of next statement is a result of [CP3], generalizing the second one, which appears in [K2].

Proposition 3.6. [CP3, Prop. 3], [K2, Theorem 2.6] *The map $F : t_\tau v \mapsto v^{-1}(\tau)$ is a bijection between \mathcal{W} and D . Moreover, F restricts to a bijection between \mathcal{W}_{ab} and D_{ab} .*

Set now, for $1 \leq j \leq ht(\theta)$:

$$D'_j = \{ \sigma \in Q^\vee \mid -(j+1) \leq (\sigma, \beta) < j+1 \forall \beta \in \Delta^+ \}.$$

Set $D_i = D'_i \cap D$. Note that $D_{ab} = D_1 = D'_1$. Next lemma follows by standard arguments (see [IM]); for the sake of completeness we include a proof.

Lemma 3.7. *Assume $\mathfrak{i} \in \mathcal{I}$ and $w_{\mathfrak{i}} = t_\tau v$, with $\tau \in Q^\vee$ and $v \in W$. The following conditions are equivalent:*

- (i) $w_{\mathfrak{i}}(C_1) \subset C_{j+1}$;
- (ii) *for each $\alpha \in \Delta^+$, $0 \leq (\tau, \alpha) \leq j+1$; moreover, if $(\tau, \alpha) = 0$, then $v^{-1}(\alpha) > 0$ and if $(\tau, \alpha) = j+1$, then $v^{-1}(\alpha) < 0$.*

Proof. Assume that (i) holds; then the hyperplanes $H_{\alpha,0}$, $H_{\alpha,j+1}$ do not separate C_1 and $w_{\mathfrak{i}}(C_1)$ for any $\alpha \in \Delta^+$. By direct computation, we obtain

$$w_{\mathfrak{i}}^{-1}(-\alpha + (j+1)\delta) = v^{-1}t_{-\tau}(-\alpha + (j+1)\delta) = -v^{-1}(\alpha) + (j+1 - (\alpha, \tau))\delta.$$

By 2.1, the above root should be positive, hence $(\alpha, \tau) \leq j+1$, and, if equality holds, then necessarily $v^{-1}(\alpha) < 0$. Similarly, $w_{\mathfrak{i}}^{-1}(\alpha) = v^{-1}(\alpha) - (\alpha, \tau)\delta > 0$, hence $(\alpha, \tau) \geq 0$, and $v^{-1}(\alpha) > 0$ if $(\alpha, \tau) = 0$. Hence (ii) holds. The converse is similar. \square

For $\mathfrak{i} \in \mathcal{I}$, let $n(\mathfrak{i})$ denote the nilpotence index of $\mathfrak{i} \in \mathcal{I}$, i.e. the minimal $n \in \mathbb{N}^+$ such that $\mathfrak{i}^{n+1} = 0$, where \mathfrak{i}^n is the n th term of the descending central series of \mathfrak{i} ($\mathfrak{i}^1 = \mathfrak{i}$).

Proposition 3.8. *F induces by restriction bijections between $\{w_{\mathfrak{i}} \mid \mathfrak{i} \in \mathcal{I}, n(\mathfrak{i}) \leq j\}$ and D_j .*

Proof. By [CP2, Corollary 2.5] the ad-nilpotent ideal \mathfrak{i} of \mathfrak{b} has nilpotence index at most j if and only if $-\alpha + (j+1)\delta \notin L_{\mathfrak{i}}$ for any $\alpha \in \Delta^+$; in turn, this condition is equivalent to

$$w_{\mathfrak{i}}(C_1) \subset C_{j+1}$$

(this follows, e.g., by formulas 2.1). By Lemma 3.7, we assume that condition 3.7(ii) holds and consider $(v^{-1}(\tau), \beta)$ for $\beta \in \Delta^+$. We have either $\beta = v^{-1}(\alpha)$ or $\beta = -v^{-1}(\alpha)$ for some $\alpha \in \Delta^+$. In the former case $v^{-1}(\alpha) > 0$ and by our assumption $0 \leq (v^{-1}(\tau), \beta) = (\tau, \alpha) < j+1$; in the latter case $v^{-1}(\alpha) < 0$, whence $-j-1 \leq (v^{-1}(\tau), \beta) = -(\tau, \alpha) < 0$. In either cases we have $-j-1 \leq (v^{-1}(\tau), \beta) < j+1$. In a similar way we see that, conversely, if $-j-1 \leq (v^{-1}(\tau), \beta) < j+1$, then 3.7(ii) holds. \square

§4 A CHARACTERIZATION OF MAXIMAL ABELIAN IDEALS

In this section we study maximal abelian ideals of \mathfrak{b} , obtaining a characterization for them in terms of w_i .

We shall need the following easy remark.

4.1. If $\tau \in Q^\vee$ is dominant, then either $(\tau, \theta) = 0$, or $(\tau, \theta) \geq 2$. This is immediate since 0 is the unique element in $Q^\vee \cap \overline{C}_1$.

Lemma 4.2. *Let $i \in \mathcal{I}_{ab}$ and $\beta \in \Delta$. If $\beta + k\delta \in w_i(\widehat{\Pi})$, for some $k \geq 2$, then $\beta = -\theta$, $k = 2$, and $w_i^{-1}(-\theta + 2\delta) \in \Pi$.*

Proof. Let $\beta + k\delta = w_i(\alpha)$, $\alpha \in \widehat{\Pi}$. Then $\alpha \notin L_{w_i^{-1}}$, hence $\ell(w_i s_\alpha) > \ell(w_i)$ and by 2.7 $N(w_i s_\alpha) = N(w_i) \cup \{w_i(\alpha)\} = N(w_i) \cup \{\beta + k\delta\}$. If $\beta > 0$, then by 2.3 (3) we should obtain that $\beta \in N(w_i s_\alpha)$, which is impossible since $\beta \notin N(w_i)$; hence $\beta < 0$. In particular, $\beta + k\delta \notin \Pi + \mathbb{N}\delta$, hence, by 3.3, no root in $\Pi + \mathbb{N}\delta$ belongs to $N(w_i s_\alpha)$. But it is clear that $\beta + k\delta$ is equal to $-\theta + 2\delta$ plus a linear combination of roots in $\Pi + \mathbb{N}\delta$ with integral non-negative coefficients; hence by 2.3 (3) we obtain that $-\theta + 2\delta \in N(w_i s_\alpha)$. Using 3.3 again, we obtain that $-\theta + 2\delta = \beta + k\delta$. It remains to prove that $\alpha \neq -\theta + \delta$. Let $w_i = t_\tau v$, $\tau \in Q^\vee$, $v \in W$. If by contradiction $\alpha = -\theta + \delta$, then $v(\theta) = \theta$ and $t_\tau v(-\theta + \delta) = -\theta + (1 + (\theta, \tau))\delta$. Since τ is dominant, by 4.1 we obtain that either $t_\tau v(-\theta + \delta) = -\theta + \delta$ or $t_\tau v(-\theta + \delta) = -\theta + k\delta$ with $k > 2$. \square

Lemma 4.3. *Let $i \in \mathcal{I}_{ab}$. Then $w_i(\widehat{\Pi}) \subseteq \pm(\Delta^- + \delta) \cup \Pi \cup \{-\theta + 2\delta\}$.*

Proof. Let $\alpha \in \widehat{\Pi}$. If $w_i(\alpha) < 0$, then by 3.3 $w_i(\alpha) \in \Delta^+ - \delta$. Assume $w_i(\alpha) > 0$, so that, by 2.7, $N(w_i s_\alpha) = N(w_i) \cup \{w_i(\alpha)\}$. If $w_i(\alpha) = \beta + k\delta$ with $\beta \in \Delta^+$ and $k \geq 0$, then by 2.3 (3) there exists some $\gamma \in \Pi$ such that $\gamma \in N(w_i s_\alpha)$, whence, by 3.3, $\gamma = w_i(\alpha)$. Finally, if $w_i(\alpha) = -\beta + k\delta$ with $\beta \in \Delta^+$ and $k > 0$, then either $k = 1$, or, by 4.2, $\beta = \theta$ and $k = 2$. \square

Theorem 4.4. *$i \in \mathcal{I}_{ab}$ is maximal if and only if $w_i(\widehat{\Pi}) \cap (\Delta^- + \delta) = \emptyset$. In this case $-\theta + 2\delta \in w_i(\Pi_\ell)$.*

Proof. The abelian ideal i is maximal in \mathcal{W}_{ab} if and only if, for all $w \in \widehat{W}$ such that $w_i \leq w$ in the weak order, $w \notin \mathcal{W}_{ab}$. By 3.3, if $u \in \mathcal{W}_{ab}$ and $u' \leq u$ in the weak order of \widehat{W} , then $u' \in \mathcal{W}_{ab}$ as well. This implies that i is maximal in \mathcal{I}_{ab} if and only if, for all $\alpha \in \widehat{\Pi}$ such that $w_i(\alpha) > 0$, we have that $w_i s_\alpha \notin \mathcal{W}_{ab}$. Since for $w_i(\alpha) > 0$ we have $N(w_i s_\alpha) = N(w_i) \cup \{w_i(\alpha)\}$, this happens if and only if $w_i(\alpha) \notin \Delta^- + \delta$. This proves the first statement.

Now if $w_i(\widehat{\Pi}) \cap (\Delta^- + \delta) = \emptyset$, we have in particular that w_i is not the identity of \widehat{W} . Notice, moreover, that by 4.1 no translation may correspond to some non zero abelian ideal. Hence if $w_i = t_\tau v$, then v is not the identity; therefore $v(\alpha) < 0$ for at least one $\alpha \in \Pi$. But by 4.3, $t_\tau v(\widehat{\Pi}) \subseteq \Delta^+ - \delta \cup \Pi \cup \{-\theta + 2\delta\}$, hence for such an α , $v(\alpha) = -\theta$ and $w_i(\alpha) = -\theta + 2\delta$. Moreover, since θ is long, α is long too. \square

4.5. Remark. If $w \in \mathcal{W}_{ab}$, and $w = t_\tau v$ with $\tau \in Q^\vee$, $v \in W$, then $w^{-1}(-\theta + 2\delta) \in \Pi_\ell$ if and only if $v^{-1}(-\theta) \in \Pi_\ell$. In fact, if $v(\alpha) = -\theta$, then by 4.1 we obtain that either $w(\alpha) = -\theta$, or $w(\alpha) = -\theta + 2\delta$, but by 4.3 the first case cannot occur.

In 3.6 we recalled Kostant's bijection between \mathcal{I}_{ab} and a certain subset D_{ab} of Q^\vee . Theorem 4.4 translates into the following characterization of maximal abelian ideals inside D_{ab} .

Proposition 4.6. *Let $\mathfrak{i} \in \mathcal{I}_{ab}$ and let $w_i = t_\tau v$, with $\tau \in Q^\vee$ and $v \in W$. Then \mathfrak{i} is maximal if and only if*

- (1) $(v^{-1}(\tau), \alpha_i) \neq -1$ for $1 \leq i \leq n$;
- (2) $(v^{-1}(\tau), \theta) \neq 0$.

Proof. For all $\alpha \in \widehat{\Delta}$ we have $w_i(\alpha) = v(\alpha) - (v(\alpha), \tau)\delta$; moreover, $v(-\theta + \delta) = -v(\theta) + \delta$ and $(\delta, \tau) = 0$. Therefore by a direct computation we obtain that $w_i(\widehat{\Pi}) \cap \Delta^- + \delta = \emptyset$ if and only if (1) and (2) hold. \square

§5 THE STRUCTURE OF X

As explained in the introduction, the set $X = Q^\vee \cap \overline{C}_2$ plays a prominent role in the description of the abelian ideals. In this section we provide a technical characterization of X . Indeed we shall give an orthogonal decomposition of each element of X as a sum of highest roots of nested irreducible parabolic subsystems of Δ .

We need preliminarily the analysis of which roots are dominant; this fact is known to the experts, nevertheless we prefer to provide a proof (which is different from the one given in [St1, Proposition 2.1]).

It is clear that the highest root θ is dominant. Moreover, θ^\vee is a short root in the dual system Δ^\vee and is clearly dominant relatively to Δ^\vee . Symmetrically, if $\theta(\Delta^\vee)$ is the highest root of Δ^\vee and we set $\theta_s = \frac{2\theta(\Delta^\vee)}{(\theta(\Delta^\vee), \theta(\Delta^\vee))}$, we obtain that θ_s is a short dominant root in Δ . Therefore θ, θ_s are dominant; the converse is also true.

Lemma 5.1. *The unique dominant roots in Δ are θ and θ_s .*

Proof. Let $\alpha \in \Delta$ be a long dominant root. Then $(\alpha, \beta) \geq 0$ for all $\beta \in \Delta^+$; indeed, since α is long, we have $(\beta, \alpha^\vee) \in \{0, 1\}$ for all $\beta \in \Delta^+$. It follows that $\beta - \alpha \notin \Delta^+$ for all $\beta \in \Delta^+$, otherwise $(\beta - \alpha, \alpha^\vee) = (\beta, \alpha^\vee) - (\alpha, \alpha^\vee) < 0$, against the assumption that α is dominant. Therefore $\alpha = \theta$. Now assume that α is a short dominant root. Then α^\vee is a long dominant root in Δ^\vee , therefore $\alpha^\vee = \theta(\Delta^\vee)$ and $\alpha = \theta_s$. \square

Lemma 5.2. *If $\tau \in Q$ is dominant and non zero then $\tau \geq \theta_s$. Equivalently, if $\tau \in Q^\vee$ is dominant and non zero then $\tau \geq \theta^\vee$.*

Proof. It is well known that the fundamental weights are nonnegative linear combinations of Π , hence any dominant weight σ is a nonnegative linear combination

of Π ; it is also clear that $\text{Supp}(\sigma) = \Pi$. In particular, if $\sigma \in -Q^+$, then there exists some $\alpha \in \Pi$ such that $(\sigma, \alpha) < 0$. Indeed, if $0 \neq \sigma = \sum_{i=1}^n \lambda_i \alpha_i$ with $\lambda_i \leq 0$, then there exists an i such that $\lambda_i < 0$ and $(\sigma, \alpha_i) < 0$. This is immediate, since $(\alpha_i, \alpha_j) \leq 0$ for $1 \leq i < j \leq n$, hence $(\sigma, \alpha_j) \geq 0$ for all j such that $\lambda_j = 0$.

Now let $\tau \in Q$ be dominant and assume, by contradiction, that $\tau - \theta_s \notin Q^+$. Then $\tau - \theta_s$ is an integral linear combination of Π with some negative coefficient, say $\tau = \theta_s - \sum_{j=1}^h \lambda_j \alpha_{i_j} + \sum_{j=h+1}^n \lambda_j \alpha_{i_j}$, with $\{i_1, \dots, i_n\} = [n]$, $1 \leq h \leq n$, $\lambda_j \in \mathbb{N}$ and $\lambda_j > 0$ for some $1 \leq j \leq h$. Set $\sigma = \sum_{j=1}^h \lambda_j \alpha_{i_j}$, and $v = \sum_{j=h+1}^n \lambda_j \alpha_{i_j}$. Let $\sigma' \in Q^+$ be a maximal element, with respect to the standard partial order, such that $\sigma' \leq \sigma$, and $\theta_s - \sigma' \in \Delta$. Set $\theta'_s = \theta_s - \sigma'$ and $\sigma'' = \sigma - \sigma'$, so that $\tau = \theta'_s - \sigma'' + v$. By the above discussion, if $\sigma'' \neq 0$ then there exist some $k \in [h]$ such that $\sigma'' > \alpha_{i_k}$ and $(-\sigma'', \alpha_{i_k}) < 0$. But it is clear that $(v, \alpha_{i_k}) \leq 0$; moreover, $(\theta'_s, \alpha_{i_k}) \leq 0$, otherwise $\theta'_s - \alpha_{i_k} = \theta_s - (\sigma' + \alpha_{i_k}) \in \Delta$, and $\sigma' + \alpha_{i_k} \leq \sigma$, against the maximality of σ' . It follows that $(\tau, \alpha_{i_k}) < 0$, a contradiction. Hence $\tau = \theta'_s + v$. By 5.1, θ'_s is not dominant, hence $(\theta'_s, \alpha_i) < 0$ for some $i \in [n]$. Since $\theta'_s = \theta_s - \sigma'$, θ_s is dominant, and $(-\sigma', \alpha_{i_j}) \geq 0$ for $j \in [n] \setminus [h]$, we obtain that $(\theta'_s, \alpha_{i_k}) < 0$ for some $k \in [h]$. But then $(v, \alpha_{i_k}) \leq 0$, hence $(\tau, \alpha_{i_k}) < 0$, against the assumption. \square

Notation. We set

$$\Delta_\theta = \{\alpha \in \Delta \mid (\alpha, \theta) = 0\}, \quad \Pi'_\theta = \{\alpha \in \Pi \mid (\alpha, \theta) \neq 0\}.$$

Lemma 5.3.

- (1) Δ_θ is the standard parabolic subsystem of Δ generated by $\Pi \setminus \Pi'_\theta$.
- (2) The longest element w_0 of W preserves the irreducible components of Δ_θ and acts on them as their longest element.
- (3) Let Γ be any irreducible component of Δ_θ , $\theta(\Gamma)$ its highest root and $\alpha \in \Pi'_\theta$. Then $(\alpha, \theta(\Gamma)) < 0$.

Proof.

- (1) This is well known, since θ is dominant.
- (2) Let w_0^θ be the longest element of $W(\Delta_\theta)$. Then w_0^θ is the commutative product of the longest elements relative to each irreducible component of Δ_θ , hence it preserves Δ_θ componentwise and acts on each component as its longest element. Consider $s_\theta w_0^\theta$. For all $\alpha \in \Delta^+$ we have $s_\theta(\alpha) = \alpha - (\alpha, \theta^\vee)\theta$, hence $s_\theta(\alpha) < 0$ if and only if $\alpha \not\leq \theta$; it follows that $N(s_\theta) = \Delta^+ \setminus \Delta_\theta^+$. Moreover, s_θ fixes Δ_θ pointwise, thus we have $N(s_\theta w_0^\theta) = N(s_\theta) + s_\theta N(w_0^\theta) = (\Delta^+ \setminus \Delta_\theta^+) + s_\theta \Delta_\theta^+ = (\Delta^+ \setminus \Delta_\theta^+) + \Delta_\theta^+ = \Delta^+$. It follows that $s_\theta w_0^\theta = w_0$, which clearly implies the claim.
- (3) This follows from the fact that if $\alpha \in \Pi'_\theta$, then α is connected to each component of the Dynkin graph of $\Pi \setminus \Pi'_\theta$. This implies that $(\alpha, \beta) \leq 0$ for all $\beta \in \Gamma$ and $(\alpha, \beta) < 0$ for at least one $\beta \in \Gamma \cap \Pi$. Since $\text{Supp}(\theta(\Gamma)) = \Gamma \cap \Pi$, we obtain $(\alpha, \theta(\Gamma)) < 0$. \square

Definition 5.4. Let

$$\Delta = \Delta_1 \supset \Delta_2 \supset \dots \supset \Delta_k$$

be a chain of standard parabolic irreducible subsystems of Δ and set $\theta_i = \theta(\Delta_i)$ for $i = 1, \dots, k$. We shall say that $\Delta = \Delta_1 \supset \Delta_2 \supset \dots \supset \Delta_k$ is a good chain if:

- (1) Δ_i is an irreducible component of $\{\alpha \in \Delta_{i-1} \mid (\alpha, \theta_{i-1}) = 0\}$.
- (2) If $\alpha \in \Pi$ and $(\alpha, \theta_i) > 0$, then $\alpha \perp \Delta_j$ for all $j \geq i + 2$.
- (3) Δ_i contains at least a long root.

Notation. For a good chain $\{\Delta = \Delta_1 \supset \Delta_2 \supset \dots \supset \Delta_k\}$ we set

$$\theta_i = \theta(\Delta_i); \quad \Pi_i = \Delta_i \cap \Pi; \quad \Pi'_i = \{\alpha \in \Pi_i \mid (\alpha, \theta_i) \neq 0\},$$

for $i = 1, \dots, k$.

Lemma 5.5. Let $\{\Delta = \Delta_1 \supset \Delta_2 \supset \dots \supset \Delta_k\}$ be a good chain.

- (1) For $i = 1, \dots, k$, θ_i is a long root, in particular $(\theta_i^\vee, \beta) \in \{-1, 0, 1, 2\}$, for all $\beta \in \Delta^+$ and $(\theta_i^\vee, \beta) = 2$ if and only if $\beta = \theta_i$.
- (2) We have $(\theta_i^\vee, \beta) = 0$ for all $\beta \in \Delta_j$ with $1 \leq i < j \leq k$.
- (3) Let $\alpha \in \Pi$. Then $(\theta_i^\vee, \alpha) = -1$ if and only if $\alpha \in \Pi'_{i-1}$, and $(\theta_i^\vee, \alpha) > 0$ if and only if $\alpha \in \Pi'_i$.

Proof. The first statement follows from property (3) in Definition 5.4. The second statement follows directly from the definitions, too. We prove (3). Assume $\alpha \in \Pi$. By Lemma 5.3 (3), if $\alpha \in \Pi'_{i-1}$, we have $(\theta_i^\vee, \alpha) = -1$. Moreover, it is clear that $(\theta_i^\vee, \alpha) \in \{1, 2\}$ if $\alpha \in \Pi'_i$ and that $(\theta_i^\vee, \alpha) = 0$ if $\alpha \in \Pi_i \setminus \Pi'_i$, so it remains to prove that $(\alpha, \theta_i) = 0$ if $\alpha \notin \Pi'_{i-1} \cup \Pi'_i$. Assume $\alpha \in \Pi_{i-j} \setminus (\Pi'_{i-1} \cup \Pi'_i)$, with $j \geq 1$. If $\alpha \in \Pi'_{i-j}$, then $j > 1$ and we are done by property (2) of Definition 5.4. If $\alpha \notin \Pi'_{i-j}$, then α lies in a connected component of $\Delta_{\theta_{i-j}}$. If such component is not Δ_{i-j+1} , then $\alpha \perp \Pi_{i-j+1}$, hence $(\alpha, \theta_i) = 0$, since $\theta_i \in \Delta_{i-j+1}$. If $\alpha \in \Pi_{i-j+1}$, we conclude by induction. \square

Set $X^+ = X \setminus \{0\}$.

Proposition 5.6. Let $\{\Delta = \Delta_1 \supset \dots \supset \Delta_k\}$ be a good chain of root subsystems of Δ and set $\tau = \sum_{i=1}^k \theta_i^\vee$. Then $\tau \in X^+$. Moreover, if $\alpha \in \Pi$, we have $(\tau, \alpha) \neq 0$ if and only if $\alpha \in \Pi'_k$.

Conversely, for each $\tau \in X^+$ there exists a good chain $\{\Delta = \Delta_1 \supset \dots \supset \Delta_k\}$ such that $\tau = \sum_{i=1}^k \theta_i^\vee$.

Proof. Clearly $\sum_{i=1}^k \theta_i^\vee \in Q^\vee$, so we only have to prove that $\sum_{i=1}^k \theta_i^\vee \in \overline{C}_2$. We have

$\left(\sum_{i=1}^k \theta_i^\vee, \theta \right) = (\theta^\vee, \theta) + \sum_{i=2}^k (\theta_i^\vee, \theta) = 2$ since, for $i \geq 2$, $\theta_i \in \Delta_\theta$. Thus it remains to prove that $\sum_{i=1}^k \theta_i^\vee$ is dominant. Let $\alpha \in \Pi$. By Lemma 5.5, $(\alpha, \theta_i) \neq 0$ if and only if either $\alpha \in \Pi'_i$ or $\alpha \in \Pi'_{i-1}$. Set $i^* = \max \{i \mid \alpha \in \Pi_i\}$. Then $\alpha \notin \Pi'_i$ for all $i < i^*$ and $i > i^*$. If also $\alpha \notin \Pi'_{i^*}$, we obtain $\left(\alpha, \sum_{i=1}^k \theta_i^\vee \right) = 0$. Assume $\alpha \in \Pi'_{i^*}$. If $i^* < k$, then

clearly $\alpha \neq \theta_{i^*}$ and by Lemma 5.5 we obtain $\left(\alpha, \sum_{i=1}^k \theta_i^\vee\right) = (\alpha, \theta_{i^*}^\vee) + (\alpha, \theta_{i^*+1}^\vee) = 1 - 1 = 0$. If $i^* = k$, then $\left(\alpha, \sum_{i=1}^k \theta_i^\vee\right) = (\alpha, \theta_{i^*}^\vee) \in \{1, 2\}$.

Now we prove the converse statement. Assume $\tau \in X^+$. As remarked in 4.1 we have $(\tau, \theta) = 2$ and, by Lemma 5.2, $\tau \geq \theta^\vee$. If $\tau = \theta^\vee$ we are done, so we assume $\tau - \theta^\vee \neq 0$. We have $(\tau - \theta^\vee, \theta) = 0$ and $\tau - \theta^\vee \in (Q^\vee)^+$. Since θ is dominant, this implies that all the simple coroots which support $\tau - \theta^\vee$ must be perpendicular to θ , so that, indeed, $\tau - \theta^\vee \in Q^\vee(\Delta_\theta)$. It is clear that $\tau - \theta^\vee$ is still dominant relatively to Δ_θ ; we claim that, moreover, $\text{Supp}(\tau - \theta^\vee)$ is included in a single irreducible component of Δ_θ . By Lemma 5.3 (3), for any irreducible component Γ of Δ_θ and $\alpha \in \Pi'_\theta$ we have $(\alpha, \theta(\Gamma)^\vee) = -1$. Let $\Gamma_1, \dots, \Gamma_j$ be irreducible components of Δ_θ . Then $\tau - \theta^\vee$ is a sum of dominant elements in $Q^\vee(\Gamma_1), \dots, Q^\vee(\Gamma_j)$. If, by contradiction, $\text{Supp}(\tau - \theta^\vee)$ has non-trivial intersection with two distinct irreducible components, say Γ_1 and Γ_2 , of Δ_θ , then, by Lemma 5.2, $\tau - \theta^\vee \geq \theta(\Gamma_1)^\vee + \theta(\Gamma_2)^\vee$. It follows that, for $\alpha \in \Pi'_\theta$, we have $(\alpha, \tau - \theta^\vee) \leq -2$, and therefore that $(\alpha, \tau) \leq -1$, which is impossible. So there exists an irreducible component Δ_2 of Δ such that $\tau - \theta^\vee$ lies in $Q^\vee(\Delta_2)$. By Lemma 5.2, $\tau - \theta^\vee \geq \theta_2^\vee$. By induction we can find a chain of irreducible subsystems $\Delta = \Delta_1 \supset \Delta_2 \supset \dots \supset \Delta_k$, such that Δ_i is an irreducible component of $(\Delta_i)_{\theta_i}$ and $\tau = \sum_{i=1}^k \theta_i^\vee$.

It remains to prove that $\Delta = \Delta_1 \supset \Delta_2 \supset \dots \supset \Delta_k$ satisfies conditions (2) and (3) of Definition 5.4.

First we prove (3). Assume by contradiction that Δ_i consists entirely of short roots, for some i in $[k]$. By a direct check we obtain that then Δ_i has rank 1 and Δ_{i-1} is of type B_3 or G_2 ; in particular $i = k$. Now $\theta_{k-1}^\vee + \theta_k^\vee$ must be dominant for Δ_{k-1} , but this is not the case. In fact, θ_k is the short root of Π_{k-1} , while θ_{k-1} is long. If α is the (unique) root in Π'_{k-1} , then α is long, thus $(\theta_{k-1}^\vee, \alpha) = 1$ and $(\theta_k^\vee, \alpha) < -1$, hence $(\theta_{k-1}^\vee + \theta_k^\vee, \alpha) < 0$.

Finally we prove (2). Assume $\alpha \in \Pi'_i$. Clearly $(\alpha, \theta_j) = 0$ for $1 \leq j < i$. Moreover, as remarked above, we have $(\alpha, \theta_{i+1}) < 0$ and, since $\alpha \notin \Delta_j$ for $j > i$, $(\alpha, \theta_j) \leq 0$ for $j \geq i + 2$. Since θ_i and θ_{i+1} are long we obtain $(\tau, \alpha) = (\theta_i^\vee, \alpha) + (\theta_{i+1}^\vee, \alpha) + \sum_{j \geq i+2} (\theta_j^\vee, \alpha) = 1 - 1 + \sum_{j \geq i+2} (\theta_j^\vee, \alpha)$. Since τ is dominant it follows that $(\theta_j^\vee, \alpha) = 0$ for all $j \geq i + 2$. \square

Remark 5.7. The proof of Theorem 5.6 makes clear the following facts.

- (1) Assume $\tau \in X^+$ and $\tau = \theta_1^\vee + \dots + \theta_k^\vee$, as in 5.6. Then $\frac{(\theta, \theta)}{2} \tau = \theta_1 + \dots + \theta_k$, and $\frac{(\theta, \theta)}{2}(\tau, \tau) = 2k$.
- (2) Assume $\tau, \tilde{\tau} \in X^+$, $\tau = \theta_1^\vee + \dots + \theta_k^\vee$, and $\tilde{\tau} = \tilde{\theta}_1^\vee + \dots + \tilde{\theta}_k^\vee$ as in Theorem 5.6. Let h be the maximal index such that $\theta_h = \tilde{\theta}_h$. Then $\theta_i = \tilde{\theta}_i$ for $1 \leq i \leq h$; moreover, $\theta_i \perp \tilde{\theta}_j$ for all $i, j > h$. It follows that $\frac{(\theta, \theta)}{2}(\tau, \tilde{\tau}) = 2h$. In particular the orthogonal decomposition of $\tau \in X^+$ provided by Theorem 5.6 is unique.
- (3) Let $\alpha \in \Delta_\ell^+$. Then there exists a unique maximal good chain $\Delta = \{\Delta_1 \supset \dots \supset \Delta_k\}$ such that $\alpha \in \Delta_i$ for $1 \leq i \leq k$. Moreover, if $\{\Delta = \Delta'_1 \supset \dots \supset \Delta'_h\}$ is any good chain such that $\alpha \in \Delta'_h$, then $h \leq k$ and $\Delta'_i = \Delta_i$ for $1 \leq i \leq h$. This is easily seen by induction; indeed, if $\{\Delta = \Delta_1 \supset \dots \supset \Delta_i\}$

is a good chain such that $\alpha \in \Delta_i$, then α belongs to at most one connected component of $\Delta_i \cap \theta_i^\perp$, hence there is at most one good chain $\{\Delta = \Delta_1 \supset \cdots \supset \Delta_{i+1}\}$ which extends $\{\Delta = \Delta_1 \supset \cdots \supset \Delta_i\}$ and such that $\alpha \in \Delta_{i+1}$.

Examples.

$\Delta \cong A_n$. Set $\Delta_i = \langle \alpha_i, \dots, \alpha_{2n-i+1} \rangle$ $1 \leq i \leq \lfloor \frac{n+1}{2} \rfloor$. Then the possible good chains are $\mathcal{S}_j = \{\Delta_1, \dots, \Delta_j\}$, $1 \leq j \leq \lfloor \frac{n+1}{2} \rfloor$. The corresponding elements in X are $\tau_j = \omega_j^\vee + \omega_{n-j+1}^\vee$ (here ω_i^\vee denotes the i th fundamental coweight).

$\Delta \cong E_6$. Due to condition (2), there are only two good chains:

$$\{\Delta\}, \quad \{\Delta, \langle \alpha_1, \alpha_3, \alpha_4, \alpha_5, \alpha_6 \rangle\}.$$

Indeed $X^+ = \{\theta^\vee, \omega_1^\vee + \omega_6^\vee\}$.

$\Delta \cong B_{2n+1}$. Due to conditions (2) and (3), there are the following good chains:

$$\begin{aligned} & \{\Delta\}, \quad \{\Delta, \langle \alpha_3, \alpha_4, \dots, \alpha_{2n+1} \rangle\}, \dots \\ & \{\Delta, \langle \alpha_3, \alpha_4, \dots, \alpha_{2n+1} \rangle\}, \dots, \langle \alpha_{2n-1}, \alpha_{2n}, \dots, \alpha_{2n+1} \rangle\}, \quad \{\Delta, \langle \alpha_1 \rangle\}. \end{aligned}$$

§6 THE POSETS $\mathcal{I}_{ab}(\tau)$

In this section we begin to study the poset of abelian ideals encoded by elements in \widehat{W} of the form $t_\tau v$, for a fixed $\tau \in X$ and v varying in W . We prove that this poset has minimum, obtained for a special element $t_\tau v_\tau$ which will be analysed in detail. Then we introduce the posets $\mathcal{I}_{ab}(\alpha, \tau)$, which will be extensively studied in the next section.

Notation. Let $\tau \in X^+$. According to Proposition 5.6, and Remark 5.7 (2) there exists a unique good chain of irreducible subsystems $\{\Delta = \Delta_1 \supset \cdots \supset \Delta_k\}$ such that $\tau = \theta_1^\vee + \cdots + \theta_k^\vee$, where θ_i is the highest root of Δ_i . We shall call $\tau = \theta_1^\vee + \cdots + \theta_k^\vee$ *the standard orthogonal decomposition of τ* , and $\{\Delta = \Delta_1 \supset \cdots \supset \Delta_k\}$ *the good chain of τ* . Moreover we set

$$\Delta(\tau) = \Delta_k, \quad \Pi(\tau) = \Pi \cap \Delta(\tau), \quad \theta(\tau) = \theta_k, \quad \Pi'(\tau) = \{\alpha \in \Pi(\tau) \mid (\alpha, \theta(\tau)) \neq 0\}.$$

We set $\Delta(0) = \Pi(0) = \emptyset$. Then, by 5.6, given $\alpha \in \Pi$ we have $(\alpha, \tau) \neq 0$ if and only if $\alpha \in \Pi'(\tau)$.

We set, for $\tau \in X$,

$$\mathcal{W}_{ab}(\tau) = \{w \in \mathcal{W}_{ab} \mid w = t_\tau v, \ v \in W\},$$

$$\mathcal{I}_{ab}(\tau) = \{i \in \mathcal{I}_{ab} \mid w_i \in \mathcal{W}_{ab}(\tau)\}.$$

6.1. Let $w \in \widehat{W}$. By Proposition 3.1 (2), $w \in \mathcal{W}_{ab}$ if and only if $w(C_1) \subset C_2$. Let $x \in C_1$, and $w = t_\tau v$, with $\tau \in Q^\vee$ and $v \in W$. Then $w(x) \in C_2$ if and only if $(t_\tau v(x), \beta) > 0$ for all $\beta \in \Pi$, and $(t_\tau v(x), \theta) < 2$. Equivalently,

$w(x) \in C_2$ if and only if $0 < (t_\tau v(x), \beta) < 2$ for all $\beta \in \Delta^+$. For $\beta \in \Delta$ have $(t_\tau v(x), \beta) = (\tau + v(x), \beta) = (\tau, \beta) + (x, v^{-1}(\beta))$; moreover $0 < |(x, v^{-1}(\beta))| < 1$ and $(x, v^{-1}(\beta)) > 0$ if and only if $v^{-1}(\beta) > 0$. Set

$$\Delta_\tau^i = \{\alpha \in \Delta \mid (\alpha, \tau) = i\}.$$

Then we obtain:

- (1) $t_\tau v \in \mathcal{W}_{ab}$ if and only if the following conditions hold: $\tau \in X$; for all $\beta \in \Delta_\tau^0 \cap \Delta^+$, $v^{-1}(\beta) > 0$; for all $\beta \in \Delta_\tau^2$, $v^{-1}(\beta) < 0$.

By 4.1, for $\tau \in X$ we have $(\tau, \theta) = 2$ unless $\tau = 0$. In this case, by 3.1 (2), we have that $t_\tau v \in \mathcal{W}_{ab}$ if and only if $t_\tau v = 1$, the identity of \widehat{W} . So we have, equivalently,

- (2) $t_\tau v \in \mathcal{W}_{ab}$ if and only if the following conditions hold: $\tau \in X$; for all $\beta \in \Pi$, if $(\tau, \beta) = 0$, then $v^{-1}(\beta) > 0$; if $t_\tau v \neq 1$, then $v^{-1}(\theta) < 0$.

Using Theorem 5.6, we also obtain:

- (3) $t_\tau v \in \mathcal{W}_{ab}$ if and only if $\tau \in X$, $L_v \subseteq \Pi'(\tau)$, and, if $t_\tau v \neq 1$, $\theta \in N(v)$.

Moreover, by Proposition 3.3, $w \in \mathcal{W}_{ab}$ if and only if $N(w) \subseteq \Delta^- + \delta$. We have $w^{-1}(-\beta + \delta) = v^{-1}t_{-\tau}(-\beta + \delta) = v^{-1}(-\beta + (1 - (\tau, \beta))\delta) = v^{-1}(-\beta) + (1 - (\tau, \beta))\delta$. Hence $w^{-1}(-\beta + \delta) < 0$ if and only if $(\tau, \beta) > 1$ or $(\tau, \beta) = 1$ and $v^{-1}(\beta) > 0$. Thus if $w \in \mathcal{W}_{ab}$, since $(\tau, \beta) \leq 2$ for all $\tau \in X$ and $\beta \in \Delta^+$, we obtain (cfr. [CMP, Lemma C]):

- (4) $N(t_\tau v) = \{-\beta + \delta \mid \beta \in \Delta_\tau^2 \cup (\Delta_\tau^1 \setminus N(v))\}$,

hence:

- (5) $\mathfrak{g}_\beta \subseteq \mathfrak{i}(t_\tau v)$ if and only if $\beta \in \Delta_\tau^2 \cup (\Delta_\tau^1 \setminus N(v))$.

By (1), if $w \in \mathcal{W}_{ab}$, then $\Delta_\tau^2 \subseteq N(v)$. Therefore by (5), if $t_\tau v$ and $t_\tau v'$ belong to \mathcal{W}_{ab} ($\tau \in X$, $v, v' \in W$), then the inclusion of the corresponding ideals corresponds to the reverse weak order of v, v' :

- (6) $\mathfrak{i}(t_\tau v) \subseteq \mathfrak{i}(t_\tau v')$ if and only if $v' \leq v$.

Consider

$$\Delta_\tau^{even} = \{\alpha \in \Delta \mid (\alpha, \tau) \text{ is even}\}.$$

It is clear that for $\tau \in X$

- (7) $\Delta_\tau^2 = (\Delta_\tau^{even} \setminus \Delta_\tau^0) \cap \Delta^+$.

We remark that Δ_τ^0 is the parabolic standard subsystem of $\widehat{\Delta}$ generated by $\Pi \Pi'(\tau)$, and, by [CMP, Lemma 2.6 (2)], Δ_τ^{even} is isomorphic to the standard parabolic subsystem of $\widehat{\Delta}$ generated by $\widehat{\Pi} \setminus \Pi'(\tau)$.

Proposition 6.2. *Let $w \in \mathcal{W}_{ab}$, $\tau \in X^+$, and let $\tau = \theta_1^\vee + \cdots + \theta_k^\vee$ be the standard orthogonal decomposition of τ . Then $\mathbf{i}(w) \in \mathcal{I}_{ab}(\tau)$ if and only if $-\theta_i + \delta \in N(w)$ for all $i \in \{1, \dots, k\}$ and $-\beta + \delta \notin N(w)$ for all $\beta \in \Delta_k \cap \theta_k^\perp$.*

Proof. Assume $w = t_\tau v$ with $v \in W$. Since $(\tau, \theta_i) = 2$ for all $i \in \{1, \dots, k\}$, by 6.1 (4) we have that $w^{-1}(-\theta_i + \delta) < 0$, for all $i \in \{1, \dots, k\}$. Moreover, if $\beta \in \Delta_k \cap \theta_k^\perp$, we have $(\tau, \beta) = 0$, hence by 6.1 (4) $-\beta + \delta \notin N(w)$. Conversely, let $w = t_{\tilde{\tau}} v$ with $\tilde{\tau} \neq \tau$ and $\tilde{\tau} = \tilde{\theta}_1^\vee + \cdots + \tilde{\theta}_{\tilde{k}}^\vee$ be the standard orthogonal decomposition of $\tilde{\tau}$. Let $h \in \{1, \dots, \min\{k, \tilde{k}\}\}$ be the maximal index such that $\theta_h = \tilde{\theta}_h$. Then, by Remark 5.7 (2) at least one of k, \tilde{k} is strictly greater than h . Moreover, if $h < k$, then $\theta_{h+1} \perp \tilde{\tau}$ hence, by 6.1 (4), $-\theta_{h+1} + \delta \notin N(w)$. If $h = k$, then we have that $\tilde{\theta}_{h+1} \in \Delta_k \cap \theta_k^\perp$ and $-\tilde{\theta}_{h+1} + \delta \in N(w)$. \square

As recalled in 2.9, for each $I \subseteq \Pi$ there is a unique $v \in W$ with $L_v \subseteq I$ of maximal length, and maximal, indeed, with respect to the weak order: just take $v = w_0^I w_0$, where w_0^I is the longest element in $\langle s_\alpha \mid \alpha \notin I \rangle$. Then $N(v) = \Delta^+ \setminus \Delta^I$, where Δ^I is the subsystem of Δ generated by $\Pi \setminus I$.

Notation. We fix $\tau \in X$ and set

$$v_\tau = w_0^I w_0, \quad \text{with} \quad I = \Pi'(\tau).$$

By 6.1 (3) $t_\tau v_\tau \in \mathcal{W}_{ab}$. Moreover we have the following result.

Proposition 6.3.

- (1) $\mathbf{i}(t_\tau v_\tau) = \min \mathcal{I}_{ab}(\tau)$;
- (2) $\dim \mathbf{i}(t_\tau v_\tau) = \frac{1}{2}(|\Delta_\tau^{\text{even}}| - |\Delta_\tau^0|)$.

Proof. (1) From 6.1 (6) we directly obtain that $\mathbf{i}(t_\tau v_\tau)$ is minimal in $\mathcal{I}_{ab}(\tau)$.
 (2) We have $N(v_\tau) = \Delta^+ \setminus \Delta^I = \Delta^+ \setminus \Delta_\tau^0 = \Delta_\tau^2 \cup \Delta_\tau^1$, hence, by 6.1 (5), $\mathbf{i}(t_\tau v_\tau) = \bigoplus_{\alpha \in \Delta_\tau^2} \mathfrak{g}_\alpha$. In particular $\dim \mathbf{i}(t_\tau v_\tau) = |\Delta_\tau^2|$, and by 6.1 (7) we obtain formula (2). \square

Lemma 6.4. *The element v_τ is an involution.*

Proof. We have, for any $I \subseteq \Pi$, that $(w_0^I w_0)^{-1} = w_0^J w_0$ with $J = -w_0(I)$. So, in our case, we have to prove that $-w_0(I) = I$, for $I = \Pi'(\tau)$. By Lemma 5.3 (2) w_0 preserves all irreducible components of Δ_θ and acts on them as their longest element. Since the longest element maps the highest root to its opposite, by recursion we obtain that $-w_0$ fixes τ , for each $\tau \in X$, and therefore stabilizes $\Pi'(\tau)$. \square

Lemma 6.5. *Let $I \subseteq \Pi$, $v = w_0^I w_0$, and $y \in W$. If v is an involution, then $L_{vy} \subseteq I$ if and only if $L_y \subseteq I$. In this case $-vN(y) \subseteq N(v)$ and $N(vy) = N(v) \setminus -vN(y)$. In particular $l(vy) = l(v) - l(y)$.*

Proof. For any $u \in W$ we have $L_u \subseteq I$ if and only if $\overline{N}(u) \subseteq \Delta \setminus \Delta^I$, where Δ^I is the subsystem of Δ generated by $\Pi \setminus I$. Recall that $\overline{N}(vy) = \overline{N}(v) + v\overline{N}(y)$. If $L_y \not\subseteq I$, let $\alpha \in L_y \setminus I$. Then $\alpha \in N(y)$, whence $v(\alpha) \in vN(y)$, but $v(\alpha) = v^{-1}(\alpha) > 0$,

hence $v(\alpha) \notin \overline{N}(v)$; therefore $v(\alpha) \in \overline{N}(vy)$. Since $\overline{N}(v) = \Delta \setminus \Delta^I$, we obtain that $\overline{N}(vy) \not\subseteq \Delta \setminus \Delta^I$, hence $L_{vy} \not\subseteq I$. Conversely, if $\overline{N}(y) \subseteq \Delta \setminus \Delta^I$, then $vN(y) = v^{-1}N(y) < 0$. By 2.7, it follows that $N(vy) \subseteq N(v)$, hence $L_{vy} \subseteq L_v$. Moreover, $-v(N(y)) \subseteq N(v)$ and $l(vy) = l(v) - l(y)$. \square

Proposition 6.6. *Assume $\tau \in X$, $v, v' \in W$. Then $t_\tau v \in \mathcal{W}_{ab}$ if and only if $v = v_\tau y$ with $L_y \subseteq \Pi'(\tau)$.*

Assume $t_\tau v, t_\tau v' \in \mathcal{W}_{ab}$, $v = v_\tau y$, and $v' = v_\tau y'$. Then $\mathfrak{i}(t_\tau v) \subseteq \mathfrak{i}(t_\tau v')$ if and only if $N(y) \subseteq N(y')$.

Proof. The first assertion follows directly from 6.1 (3), 6.4, and 6.5.

Under the assumptions of the second assertion, by 6.5 we have $-v_\tau N(y) \subseteq N(v_\tau)$, $-v_\tau N(y') \subseteq N(v_\tau)$, $N(v_\tau y) = N(v_\tau) \setminus -v_\tau N(y)$, and $N(v_\tau y') = N(v_\tau) \setminus -v_\tau N(y')$. Therefore, $v' \leq v$ if and only if $y \leq y'$. This proves the claim. \square

It is clear that the arguments of section 4 can be specialized to $\mathcal{I}_{ab}(\tau)$. We obtain the following result.

Proposition 6.7. *Let $\mathfrak{i} \in \mathcal{I}_{ab}(\tau)$. Then \mathfrak{i} is maximal in $\mathcal{I}_{ab}(\tau)$ if and only if $w_{\mathfrak{i}}(\Pi) \cap (\Delta^- + \delta) = \emptyset$. In this case, if $\tau \neq 0$, we have $w_{\mathfrak{i}}^{-1}(-\theta + 2\delta) \in \Pi$, hence if $w_{\mathfrak{i}} = t_\tau v$, then $v^{-1}(-\theta) \in \Pi_\ell$.*

Proof. We notice that if $\mathfrak{j} \in \mathcal{I}_{ab}(\tau)$, then $w_{\mathfrak{j}} = w_{\mathfrak{i}}x$ with $x \in W$. So, we can just mimic the proof of 4.4 considering Π in place of $\widehat{\Pi}$. \square

The above result leads us to analyse $v^{-1}(\theta)$ for all $v \in W$ such that $t_\tau v \in \mathcal{I}_{ab}(\tau)$. We first study what $v_\tau(\theta)$ is.

For any root β and any simple root α let $n_\alpha(\beta)$ denote the coefficient of α in β and set $m_\alpha = n_\alpha(\theta)$:

$$\beta = \sum_{\alpha \in \Pi} n_\alpha(\beta)\alpha, \quad \theta = \sum_{\alpha \in \Pi} m_\alpha\alpha.$$

Proposition 6.8. *Fix $\tau \in X^+$.*

- (1) *We have $v_\tau(\Pi \setminus \Pi'(\tau)) = \Pi \setminus \Pi'(\tau)$.*
- (2) *Set $B_\tau = v_\tau(\Pi'(\tau))$. For each $\beta \in B_\tau$, there exists a unique $\alpha \in \Pi'(\tau)$ such that $n_\alpha(\beta) \neq 0$ and, for such an α , we have $n_\alpha(\beta) = -1$. In particular $B_\tau \subseteq \Delta^-$.*
- (3) *Let $\beta \in B_\tau$ and $\alpha \in \Pi'(\tau)$ such that $n_\alpha(\beta) = -1$. Then $-\beta$ is the maximal element in $\{\gamma \in \Delta^+ \mid \text{Supp}(\gamma) \cap \Pi'(\tau) = \{\alpha\}, n_\alpha(\gamma) = 1\}$, with respect to the standard partial order of Q .*
- (4) *We have $v_\tau(\theta) = -\theta(\tau)$. Moreover, $n_\alpha(\theta(\tau)) = m_\alpha$ for all $\alpha \in \Pi'(\tau)$ and $\theta(\tau)$ is the minimal root with this property, with respect to the standard partial order of Q .*

Proof.

(1) The claim follows since v_τ maps $\Pi \setminus \Pi'(\tau)$ to $-w_0(\Pi \setminus \Pi'(\tau))$, which is equal to $\Pi \setminus \Pi'(\tau)$, as seen in the proof of Lemma 6.4.

(2) Recall that $v_\tau = v_\tau^{-1}$; thus $B_\tau = v_\tau(\Pi'(\tau))$ is a set of negative roots. By (1) v_τ preserves $\Pi \setminus \Pi'(\tau)$, and since $v_\tau(\Pi)$ is a basis of Δ , we obtain that $B_\tau \cup (\Pi \setminus \Pi'(\tau))$ is a basis of Δ . In particular, each $\alpha \in \Pi'(\tau)$, is a nonnegative or nonpositive linear combination of $B_\tau \cup (\Pi \setminus \Pi'(\tau))$. This implies that for each $\alpha \in \Pi'(\tau)$, some $\beta \in B_\tau$ is supported by α and, moreover, α has coefficient -1 in such a β . Moreover, each $\beta \in B_\tau$ is supported by a unique $\alpha \in \Pi'(\tau)$.

(3) Let $\alpha \in \Pi'(\tau)$ and β_α be the unique root in B_τ supported by α . If $\gamma \in \Delta^+$, $\text{Supp}(\gamma) \cap \Pi'(\tau) = \{\alpha\}$, and $n_\alpha(\gamma) = 1$ then, arguing as in (2) we find that γ equals $-\beta_\alpha$ minus a sum of roots in $\Pi \setminus \Pi'(\tau)$. This proves the claim.

(4) We have $v_\tau(\theta) = \sum_{\alpha \in \Pi} m_\alpha v_\tau(\alpha)$. For each $\alpha \in \Pi'(\tau)$, set $\gamma_\alpha = v_\tau(\alpha)$, and let α' be the unique root of $\Pi'(\tau)$ which supports γ_α . Then, by (2), $n_{\alpha'}(v_\tau(\theta)) = -m_\alpha$, hence $m_\alpha \leq m_{\alpha'}$. Moreover $\{\alpha' \mid \alpha \in \Pi'(\tau)\} = \Pi'(\tau)$, hence $\{m_{\alpha'} \mid \alpha \in \Pi'(\tau)\} = \{m_\alpha \mid \alpha \in \Pi'(\tau)\}$. It follows that $\sum_{\alpha \in \Pi'(\tau)} m_\alpha = \sum_{\alpha \in \Pi'(\tau)} m_{\alpha'}$, hence $m_\alpha = m_{\alpha'}$.

Thus each $\alpha \in \Pi'(\tau)$ occurs in $-v_\tau(\theta)$ with coefficient m_α . We show that $-v_\tau(\theta)$ is the minimal root with this property. As seen in (1), v_τ permutes $\Pi \setminus \Pi'(\tau)$; since $\theta + \eta$ is not a root for each $\eta \in Q^+$, we obtain that $v_\tau(\theta) + \eta$ is not a root for all $\eta \in Q^+(\Pi \setminus \Pi'(\tau))$. Equivalently $-v_\tau(\theta) - \eta$ is not a root for all $\eta \in Q^+(\Pi \setminus \Pi'(\tau))$, hence $-v_\tau(\theta)$ is minimal among the roots in which α occurs with coefficient m_α for all $\alpha \in \Pi'(\tau)$. Now we know from the orthogonal decomposition of $\tau = \theta_1^\vee + \dots + \theta_k^\vee$ that $(\tau, \theta_i) = 2$ for $i = 1, \dots, k$. Since $n_\alpha(\theta_i) \leq n_\alpha(\theta_1) = m_\alpha$ for all $\alpha \in \Pi'(\tau)$ and since $\Pi'(\tau)$ is the set of all simple roots non orthogonal to τ , we obtain $n_\alpha(\theta_i) = m_\alpha$ for $i = 1, \dots, k$. In particular we obtain that $-v_\tau(\theta) \leq \theta(\tau)$. But $\Pi'(\tau)$ is the set of all simple roots which belong to $\Delta(\tau)$ and are not orthogonal to $\theta(\tau)$, hence we have $(-v_\tau(\theta), \theta^\vee(\tau)) = (\theta(\tau), \theta^\vee(\tau)) = 2$. Since $\theta(\tau)$ is long this implies that $-v_\tau(\theta) = \theta(\tau)$. \square

Lemma 6.9. *Let $\alpha \in \Delta_\ell^+$, $v \in W$ and assume that $\alpha \notin \Delta(\tau)$ and $v(\alpha) = \theta(\tau)$. Then $L_v \not\subseteq \Pi'(\tau)$.*

Proof. By 2.3 (5), $L_v \not\subseteq \Pi'(\tau)$, if and only if there exists some $\gamma \in N(v)$ such that $\text{Supp}(\gamma) \cap \Pi'(\tau) = \emptyset$. We shall find such a γ . Since $\alpha \notin \Delta(\tau)$, we have that $v \notin W(\Delta(\tau))$, hence $N(v) \not\subseteq \Delta(\tau)$. By 2.8, $N(v) \cap \Delta(\tau) = N(v')$ for some $v' \in W(\Delta(\tau))$, hence there exists $v'' \in W$ such that $v = v'v''$, and $v'N(v'') = N(v) \setminus \Delta(\tau)$. Set $N(v'') = \{\beta_1, \dots, \beta_h\}$, and $\gamma_i = v'(\beta_i)$, for $i = 1, \dots, h$, so that $N(v) \setminus \Delta(\tau) = \{\gamma_1, \dots, \gamma_h\}$. By 2.4 (1), we may assume that $v = s_{\gamma_h} \dots s_{\gamma_1} v'$. Conditions $\alpha \notin \Delta(\tau)$ and $v' \in W(\Delta(\tau))$ imply $v'(\alpha) \neq \theta(\tau)$, hence $s_{\gamma_h} \dots s_{\gamma_1}(\theta(\tau)) \neq \theta(\tau)$. This implies that there exists $i \in \{1, \dots, h\}$ such that $\gamma_i \not\leq \theta(\tau)$. Since $\theta(\tau) \notin N(v)$, we have that $\theta(\tau) \neq \gamma_i$, hence $(\gamma_i, \theta^\vee(\tau)) = \pm 1$. Assume first $(\gamma_i, \theta^\vee(\tau)) = -1$, so that $\gamma_i + \theta(\tau) \in \Delta^+$. By 6.8 (4), for any $\beta \in \Pi'(\tau)$ we have that $n_\beta(\theta(\tau)) = m_\beta$, hence that $n_\beta(\gamma_i) = 0$, and we are done. So assume $(\gamma_i, \theta^\vee(\tau)) = 1$, so that $\gamma_i - \theta(\tau) \in \Delta$. Since $\gamma_i \notin \Delta(\tau)$, we have that $\gamma_i \not\leq \theta(\tau)$, hence $\gamma_i = \theta(\tau) + \gamma'_i$, for some $\gamma'_i \in \Delta^+$. Since $\gamma_i \in N(v)$, and $\theta(\tau) \notin N(v)$, by 2.3 (3) we obtain that $\gamma'_i \in N(v)$. But $(\gamma'_i, \theta^\vee(\tau)) = -1$, so we are reduced to the previous case. \square

For $\tau \in X$ and $\alpha \in \Delta_\ell^+$, we set

$$\mathcal{W}_{ab}(\tau, \alpha) = \{w \in \mathcal{W}_{ab}(\tau) \mid w(\alpha) = -\theta + 2\delta\};$$

$$\mathcal{I}_{ab}(\tau, \alpha) = \{i \in \mathcal{I}_{ab} \mid w_i \in \mathcal{W}_{ab}(\tau, \alpha)\}.$$

Let $\alpha \in \Delta_\ell^+$ and $w \in \mathcal{W}_{ab}$ be such that $w(\alpha) = -\theta + 2\delta$. Then, if $w = t_\tau v$ with $\tau \in Q^\vee$ and $v \in W$, we have $v(\alpha) = -\theta$. Conversely, if $v(\alpha) = -\theta$, then $v \neq 1$, hence by 3.1 (2) $\tau \neq 0$. By 4.1 it follows that $(\tau, \theta^\vee) = 2$, hence $t_\tau v(\alpha) = t_\tau(-\theta) = -\theta - (\tau, -\theta)\delta = -\theta + 2\delta$. Therefore

$$\mathcal{W}_{ab}(\tau, \alpha) = \{t_\tau v \in \mathcal{W}_{ab}(\tau) \mid v(\alpha) = -\theta\}.$$

It is clear that $\mathcal{W}_{ab}(\tau, \alpha)$ is empty for $\tau = 0$. If $\tau \in X^+$ and $t_\tau v \in \mathcal{W}_{ab}$, then by 6.1 (2) we have that $v^{-1}(-\theta) \in \Delta_\ell^+$, hence

$$\mathcal{W}_{ab}(\tau) = \bigcup_{\alpha \in \Delta_\ell^+} \mathcal{W}_{ab}(\tau, \alpha), \quad \text{and} \quad \mathcal{I}_{ab}(\tau) = \bigcup_{\alpha \in \Delta_\ell^+} \mathcal{I}_{ab}(\tau, \alpha)$$

Proposition 6.10. *Assume $\tau \in X^+$ and $\alpha \in \Delta_\ell^+$. If $\mathcal{I}_{ab}(\tau, \alpha) \neq \emptyset$, then $\alpha \in \Delta(\tau)$.*

Proof. Assume $t_\tau v \in \mathcal{I}_{ab}(\tau, \alpha)$. Then by 6.6 $v = v_\tau y$ with $L_y \subseteq \Pi'(\tau)$; moreover, using 6.4, and 6.8 (4), $y(\alpha) = v_\tau^{-1}v(\alpha) = v_\tau(-\theta) = \theta(\tau)$. By 6.9, this implies that $\alpha \in \Delta(\tau)$. \square

§7 THE POSETS $\mathcal{I}_{ab}(\tau, \alpha)$

In this section we prove that, for $\alpha \in \Delta_\ell^+$, $\mathcal{I}_{ab}(\tau, \alpha)$ is not empty if and only if $\alpha \in \Delta(\tau)$, and in this case $\mathcal{I}_{ab}(\tau, \alpha)$ has a unique minimal element and a unique maximal element.

Let

$$\theta^\vee = \sum_{\alpha \in \Pi} m_\alpha^\vee \alpha^\vee.$$

We have $\theta^\vee = \frac{2}{(\theta, \theta)} \theta = \frac{2}{(\theta, \theta)} \sum_{\alpha \in \Pi} m_\alpha \frac{(\alpha, \alpha)}{2} \alpha^\vee$, hence $m_\alpha^\vee = \frac{(\alpha, \alpha)}{(\theta, \theta)} m_\alpha$.

The dual Coxeter number of \mathfrak{g} is

$$g = 1 + \sum_{\alpha \in \Pi} m_\alpha^\vee.$$

Part of next proposition is also proved in [S] and [P].

Proposition 7.1. *Let α be any long root. Then:*

- (1) $\{v \in W \mid v(\alpha) = \theta\}$ has a unique element y_α of minimal length.
- (2) $L_{y_\alpha} \subseteq \Pi'_\theta$; conversely, if $v' \in W$, $v'(\alpha) = \theta$, and $L_{v'} \subseteq \Pi'_\theta$, then $v' = y_\alpha$.
- (3) If $\alpha \in \Pi_\ell$, then $\ell(y_\alpha) = g - 2$;
- (4) y_α^{-1} is the unique minimal element, with respect to the weak order, in $\{v \in W \mid v(\theta) = \alpha\}$;
- (5) $N(y_\alpha^{-1}) = \{\beta \in \Delta^+ \mid (\beta, \alpha^\vee) = -1\}$.

Proof.

(1), (4). We start with proving that there exists a unique minimal element, with respect to the weak order, in $\{v \in W \mid v(\theta) = \alpha\}$. This will imply (1) and (4). Let $m = \min_{v \in W} \{\ell(v) \mid v(\theta) = \alpha\}$ and let $v \in W$ be any element such that $v(\theta) = \alpha$ and $\ell(v) = m$. Let $v' \in W$ be such that $v'(\theta) = \alpha$. Then $v' = vx$ with $x \in \text{Stab}_W(\theta)$. We shall prove that $\ell(v') = \ell(v) + \ell(x)$: by 2.6, this implies in particular that $N(v) \subseteq N(v')$. Let $v = s_{i_1} \cdots s_{i_k}$, $x = s_{j_1} \cdots s_{j_h}$ be reduced expressions. If by contradiction $s_{i_1} \cdots s_{i_k} s_{j_1} \cdots s_{j_h}$ is not reduced, then there exist $u <_B v$, $y <_B x$, $<_B$ being the Bruhat order, such that $uy = vx$. But $\text{Stab}_W(\theta) = W(\Delta_\theta) = \langle s_\beta \mid \beta \in \Pi \setminus \Pi'_\theta \rangle$, therefore, since $x \in \text{Stab}_W(\theta)$, we have $y \in \text{Stab}_W(\theta)$, too. It follows that $u(\alpha) = \theta$, against the assumption that $\ell(v)$ is minimal. We set $y_\alpha = v^{-1}$; then it is clear that y_α is the unique element of minimal length in W which maps α to θ .

(2) For all $y \in W$, if $y(\alpha) = \theta$, then $y = xy_\alpha$ with $x \in \text{Stab}_W(\theta)$ and $\ell(y) = \ell(x) + \ell(y_\alpha)$. In particular, $\ell(s_\beta y_\alpha) > \ell(y_\alpha)$ for all $\beta \in \Pi \setminus \Pi'_\theta$, hence $L_{y_\alpha} \subseteq \Pi'_\theta$. Moreover, $L_x \subseteq \Pi \setminus \Pi'_\theta$ and $N(x) \subseteq N(y)$, hence, if $y \neq y_\alpha$, $L_y \not\subseteq \Pi'_\theta$.

(5) Assume $y_\alpha(\beta) < 0$. Then $\alpha \neq \beta$ and $-y_\alpha(\beta) \in N(y_\alpha)$. By item (2), $\text{Supp}(-y_\alpha(\beta)) \cap \Pi'_\theta \neq \emptyset$, hence $(\beta, \alpha^\vee) = -(-y_\alpha(\beta), \theta^\vee) = -1$. Conversely, assume $(\beta, \alpha^\vee) = -1$. Then $(y_\alpha(\beta), \theta^\vee) = -1$, hence $y_\alpha(\beta) < 0$.

(3) Now we assume $\alpha \in \Pi$ and prove that $m = g - 2$. Let $\alpha \in \Pi_\ell$ and $v \in W$ be such that $v(\alpha) = \theta$. We first show that $\ell(v) \geq g - 2$. Let $v = s_{\gamma_1} \cdots s_{\gamma_k}$ be a reduced expression. Set $v_0 = 1$, and $v_h = \prod_{i=1}^h s_{\gamma_i}$, for $1 \leq h \leq k$. Since $N(v_h) \subseteq N(v)$, $v_h^{-1}(\theta)$ is a long positive root for $0 \leq h \leq k$; in particular, $v_{h-1}^{-1}(\theta) \neq \gamma_h$, for $0 < h \leq k$. For any positive root β and any $\gamma \in \Pi$, we have $s_\gamma(\beta) = \beta - (\beta, \gamma^\vee)\gamma$. Moreover, if β is long and $\beta \neq \gamma$, then $|(\beta^\vee, \gamma)| \leq 1$, hence $|(\beta, \gamma^\vee)| = \left| (\beta^\vee, \gamma) \frac{(\beta, \beta)}{(\gamma, \gamma)} \right| \leq \left| \frac{(\beta, \beta)}{(\gamma, \gamma)} \right| = \left| \frac{(\theta, \theta)}{(\gamma, \gamma)} \right|$. Therefore $v_{h-1}^{-1}(\theta) - v_h^{-1}(\theta) \leq \frac{(\theta, \theta)}{(\gamma_h, \gamma_h)} \gamma_h$, for $1 \leq h \leq k$, hence

$\theta - \alpha = v_0^{-1}(\theta) - v_k^{-1}(\theta) \leq \sum_{i=1}^k \frac{(\theta, \theta)}{(\gamma_i, \gamma_i)} \gamma_i$. Since $\theta = \sum_{\gamma \in \Pi} m_\gamma \gamma = \sum_{\gamma \in \Pi} m_\gamma^\vee \frac{(\theta, \theta)}{(\gamma, \gamma)} \gamma$, we obtain that each $\gamma \neq \alpha$ occurs in the sequence $(\gamma_1, \dots, \gamma_k)$ at least m_γ^\vee times, and α at least $m_\alpha^\vee - 1$ times. It follows that $\ell(v) = k \geq \sum_{\gamma \in \Pi} m_\gamma^\vee - 1 = g - 2$. Next,

we show that there exists a $w \in W$ such that $w(\alpha) = \theta$ and $\ell(w) \leq g - 2$. If β is a long root and $\beta \neq \theta$, then there exists a simple root γ such that $(\beta, \gamma^\vee) < 0$ and hence $s_\gamma(\beta) = \beta + \frac{(\theta, \theta)}{(\gamma, \gamma)} \gamma$. Therefore, if α is a simple long root, we can find a sequence of simple roots $(\gamma_1, \dots, \gamma_k)$ such that $(s_{\gamma_{i-1}} \cdots s_{\gamma_1}(\alpha), \gamma_i^\vee) < 0$, $s_{\gamma_i} \cdots s_{\gamma_1}(\alpha) = s_{\gamma_{i-1}} \cdots s_{\gamma_1}(\alpha) + \frac{(\theta, \theta)}{(\gamma_i, \gamma_i)} \gamma_i$, for $i \leq k$, and $s_{\gamma_k} \cdots s_{\gamma_1}(\alpha) = \theta$. Then clearly $k = \sum_{\gamma \in \Pi} m_\gamma^\vee - 1 = g - 2$ and therefore $\ell(s_{\gamma_k} \cdots s_{\gamma_1}) \leq g - 2$. This concludes the proof. \square

Proposition 7.2. *Let $\alpha \in \Delta_\ell^+$. Then for any pair β_1, β_2 in Δ^+ such that $\theta = \beta_1 + \beta_2$, at most one of β_1, β_2 lies in $N(y_\alpha)$; moreover if $\beta \perp \theta$, then $\beta \notin N(y_\alpha)$.*

If α is simple, exactly one of β_1, β_2 lies in $N(y_\alpha)$; in particular

$$g - 2 = \frac{1}{4}(|\Delta| - |\Delta_\theta| - 2).$$

Conversely, assume that $y \in W$ is such that for any pair β_1, β_2 in Δ^+ such that $\theta = \beta_1 + \beta_2$, exactly one of β_1, β_2 lies in $N(y)$, and, moreover, $\theta \notin N(y)$. Then there exists $\alpha \in \Pi_\ell$ such that $y(\alpha) = \theta$.

Proof. We have $\theta \notin N(y_\alpha)$, therefore, by 2.3 (3), for any pair β_1, β_2 in Δ^+ such that $\theta = \beta_1 + \beta_2$, at most one of β_1, β_2 lies in $N(y_\alpha)$. Moreover, by 7.1 (2) and 2.3 (5), no root orthogonal to θ may belong to $N(y_\alpha)$. If $\alpha \in \Pi$, since $y_\alpha(\alpha) > 0$, we have $N(y_\alpha s_\alpha) = N(y_\alpha) \cup \{\theta\}$; therefore at least one of β_1, β_2 lies in $N(y_\alpha s_\alpha)$ and hence in $N(y_\alpha)$. By 7.1 (2) we have that, moreover, no root in Δ_θ belongs to $N(y_\alpha)$, therefore, using 7.1 (3), we obtain $g - 2 = \ell(y_\alpha) = \frac{1}{2}(|\Delta^+ \setminus \Delta_\theta^+| - 1) = \frac{1}{4}(|\Delta| - |\Delta_\theta| - 2)$.

Now let $y \in W$ be such that for any pair β_1, β_2 in Δ^+ such that $\theta = \beta_1 + \beta_2$, exactly one of β_1, β_2 lies in $N(y)$, and, moreover, $\theta \notin N(y)$. Then $N(y) \cup \{\theta\}$ still verifies conditions (a) and (b) of 2.3 (4), therefore there exists $y' \in W$ such that $N(y') = N(y) \cup \{\theta\}$. By 2.6, $y' = yx$ with $\ell(y') = \ell(y) + \ell(x)$, therefore $\ell(x) = 1$; equivalently, $x = s_\alpha$ for some $\alpha \in \Pi$. Then by 2.4 we obtain that $\theta = y(\alpha)$, hence that $\alpha \in \Pi_\ell$. \square

Notation. For $\alpha \in \Delta_\ell^+$, we set $\check{w}_\alpha = s_0 y_\alpha$. Recall that $y_\alpha s_\alpha = s_\theta y_\alpha$ and $t_{\theta^\vee} = s_0 s_\theta$, therefore we have

$$\check{w}_\alpha = s_0 y_\alpha = t_{\theta^\vee} y_\alpha s_\alpha.$$

Proposition 7.3. *Let $\alpha \in \Delta_\ell^+$.*

- (1) *We have $\mathcal{I}_{ab}(\theta^\vee, \alpha) = \{i(\check{w}_\alpha)\}$.*
- (2) *For any pair β_1, β_2 in Δ^+ such that $\theta = \beta_1 + \beta_2$, at most one of $-\beta_1 + \delta, -\beta_2 + \delta$ lies in $N(\check{w}_\alpha)$, and if $\alpha \in \Pi$ exactly one. Moreover, $\alpha_0 \in N(\check{w}_\alpha)$, and $\beta \notin N(\check{w}_\alpha)$ for all $\beta \in \widehat{\Delta}^+ \cap \alpha_0^\perp$.*
- (3) *$N(\check{w}_\alpha^{-1}) = \{\beta \in \Delta^+ \mid (\beta, \alpha) = -1\} \cup \{-\alpha + \delta\}$.*

Proof.

(1) follows from 6.1 (3) and 7.1 (2).

(2) If $\theta = \beta_1 + \beta_2$ with $\beta_1, \beta_2 \in \Delta^+$, then $(\beta_i, \theta^\vee) = 1$ for $i = 1, 2$, so that $s_0(\beta_1) = -\beta_2 + \delta$ and $s_0(\beta_2) = -\beta_1 + \delta$. Since $N(\check{w}_\alpha) = N(s_0) + s_0 N(y_\alpha)$, we get the claim by Proposition 7.2.

(3) follows from 7.1 (5), since $N(\check{w}_\alpha^{-1}) = N(y_\alpha^{-1}) \cup y_\alpha^{-1} N(s_0)$. \square

Remark. It is clear that for all $\tau \in X^+$ we have that $\mathcal{I}_{ab}(\tau) = \bigcup_{\alpha \in \Delta_\ell^+} \mathcal{I}_{ab}(\tau, \alpha)$.

Therefore by 7.3 (1) we obtain that $|\mathcal{I}_{ab}(\theta^\vee)| = |\Delta_\ell^+|$. By [CMP, Theorem 2.4] we obtain that $|\Delta_\ell^+| = |W|/|W(\widehat{\Pi} \setminus \Pi'_\theta)|$.

We can apply Proposition 7.1, in particular, to the irreducible parabolic subsystems $\Delta(\tau)$, $\tau \in X^+$. We set

$$\begin{aligned}\Pi_\ell(\tau) &= \Pi_\ell \cap \Pi(\tau) \\ \Delta_\ell^+(\tau) &= \Delta_\ell^+ \cap \Delta(\tau).\end{aligned}$$

For $\alpha \in \Pi_\ell(\tau)$ we denote by $y_{\tau,\alpha}$ be the unique element of minimal length in $W(\Delta(\tau))$ such that $y_{\tau,\alpha}(\alpha) = \theta(\tau)$.

Proposition 7.4. *Let $\tau \in X^+$ and $\alpha \in \Delta_\ell^+(\tau)$. Then $t_\tau v_\tau y_{\tau,\alpha} \in \mathcal{W}_{ab}(\tau, \alpha)$. In particular, $\mathcal{I}_{ab}(\tau, \alpha)$ is not empty if and only if $\alpha \in \Delta_\ell^+(\tau)$.*

Proof. The “only if” part was stated in 6.10. Conversely, if $\alpha \in \Delta_\ell^+(\tau)$, then $y_{\tau,\alpha}(\alpha) = \theta(\tau)$ and $L_{y_{\tau,\alpha}} \subseteq \Pi'(\tau)$. Moreover, since $v_\tau y_{\tau,\alpha}(\alpha) = -\theta$, we have $\theta \in N(v_\tau y_{\tau,\alpha})$. By Lemma 6.5 and 6.1 (3), it follows that $t_\tau v_\tau y_{\tau,\alpha} \in \mathcal{I}_{ab}(\tau, \alpha)$. \square

Remark. The element y_α defined in Proposition 7.1 equals $y_{\theta^\vee, \alpha}$, in the above notation. Comparing Propositions 7.3 (1) and 7.4 we obtain that, indeed, $v_{\theta^\vee} y_\alpha = y_\alpha s_\alpha$.

Proposition 7.5. *Let $\tau \in X^+$, $\alpha \in \Delta_\ell^+(\tau)$, and $y \in W$ be such that $y(\alpha) = \theta(\tau)$ and $L_y \subseteq \Pi'(\tau)$. Then $y_{\tau,\alpha} \leq y$ and $y_{\tau,\alpha}^{-1} y \in W(\Pi \cap \alpha^\perp)$. In particular*

$$i(t_\tau v_\tau y_{\tau,\alpha}) = \min \mathcal{I}_{ab}(\tau, \alpha).$$

Proof. By 2.8, there exists $y' \in W(\Delta(\tau))$ such that $N(y') = N(y) \cap \Delta(\tau)$ and $y = y'x$ with $\ell(y) = \ell(y') + \ell(x)$. Set $\alpha' = y'^{-1}(\theta(\tau))$. Clearly $\alpha' \in \Delta^+(\tau)$, and since $L_{y'} \subseteq \Pi'(\tau)$, by 7.1 (2) we obtain that $y' = y_{\tau,\alpha'}$. Then $x = y_{\tau,\alpha'}^{-1}y$; hence $y = y_{\tau,\alpha'}x$ and $N(y) = N(y_{\tau,\alpha'}) \cup y_{\tau,\alpha'}N(x)$. We prove that x is a product of reflections with respect to simple roots orthogonal to α' . Let $x = x_1 \cdots x_k$ with x_i simple reflection for $1 \leq i \leq k$. Assume by contradiction that $x_j = s_\beta$ with $\beta \not\perp \alpha'$ for some $j \in \{1, \dots, k\}$, and $x_i(\alpha') = \alpha'$ for $i < j$. Set $x' = x_1 \cdots x_{j-1}$, and $y' = y_{\tau,\alpha'}x'$. Then $y'(\beta) \in N(y) \setminus N(y_{\tau,\alpha'})$, hence $y'(\beta) \notin \Delta(\tau)$; in particular $y'(\beta) \not\leq \theta(\tau)$. We have $y'(\alpha') = \theta(\tau)$ and, since $(\beta, \alpha') \neq 0$, $(y'(\beta), \theta(\tau)) \neq 0$, hence either $y'(\beta) - \theta(\tau)$, or $y'(\beta) + \theta(\tau)$ is a root. In the second case, by 6.8 (4) we obtain that $n_\gamma(y'(\beta)) = 0$ for all $\gamma \in \Pi'(\tau)$: this is impossible, since $y'(\beta) \in N(y)$ and $L_y \subseteq \Pi'(\tau)$. In the first case, since $y'(\beta) \not\leq \theta(\tau)$, we obtain that $y'(\beta) - \theta(\tau) > 0$; moreover, since $y'(\beta) \in N(y)$ and $\theta(\tau) \notin N(y)$, by 2.3 (3) $y'(\beta) - \theta(\tau) \in N(y)$. As above this leads to a contradiction, since $n_\gamma(y'(\beta) - \theta(\tau)) = 0$ for all $\gamma \in \Pi'(\tau)$. Thus we have in particular that $x(\alpha') = \alpha'$, hence $y(\alpha') = \theta(\tau)$; it follows that $\alpha' = \alpha$. By Proposition 6.6 we obtain that $i(t_\tau v_\tau y_{\tau,\alpha}) = \min \mathcal{I}_{ab}(\tau, \alpha)$. \square

We set, for $\tau \in X^+$, $\alpha \in \Delta_\ell^+(\tau)$

$$\begin{aligned}W_\alpha &= W(\Pi \cap \alpha^\perp), \\ P_{\tau,\alpha} &= \{\beta \in \Pi \setminus \Pi(\tau) \mid \beta \not\leq \Delta(\tau), \beta \perp \alpha\}.\end{aligned}$$

Lemma 7.6. *Let $\tau \in X^+$, $\alpha \in \Delta_\ell^+(\tau)$, and $y \in W$ be such that $y(\alpha) = \theta(\tau)$ and $L_y \subseteq \Pi'(\tau)$. Set $x = y_{\tau,\alpha}^{-1}y$. Then $x \in W_\alpha$ and $L_x \subseteq P_{\tau,\alpha}$.*

Conversely, if $x \in W_\alpha$ and $L_x \subseteq P_{\tau,\alpha}$, then $y_{\tau,\alpha}x(\alpha) = \theta(\tau)$ and $L_{y_{\tau,\alpha}x} \subseteq \Pi'(\tau)$. Moreover, $\ell(y_{\tau,\alpha}x) = \ell(y_{\tau,\alpha}) + \ell(x)$.

Proof. By Proposition 7.5, we have that $x \in W_\alpha$, hence $N(x) \subseteq \Delta(\Pi \cap \alpha^\perp)$; moreover $N(y) = N(y_{\tau,\alpha}) \cup y_{\tau,\alpha}N(x)$. Therefore it suffices to prove that for all $\beta \in \Pi \cap \alpha^\perp \setminus P_{\tau,\alpha}$ we have that $\beta \notin N(x)$. Assume $\beta \in \Pi \cap \alpha^\perp \setminus P_{\tau,\alpha}$; this means that either $\beta \in \Pi(\tau)$, or $\beta \notin \Pi(\tau)$ and $\beta \perp \Delta(\tau)$. First assume $\beta \notin \Pi(\tau)$ and $\beta \perp \Delta(\tau)$. Then, since $y_{\tau,\alpha} \in W(\Delta(\tau))$, we obtain that $y_{\tau,\alpha}(\beta) = \beta$. By assumption, $\beta \notin N(y)$, hence $\beta \notin N(x)$. Next assume $\beta \in \Pi(\tau)$. Then $y_{\tau,\alpha}(\beta) \in \Delta(\tau)$ and, since $\beta \perp \alpha$, $y_{\tau,\alpha}(\beta) \perp \theta(\tau)$. It follows that $\text{Supp}(y_{\tau,\alpha}(\beta)) \cap \Pi'(\tau) = \emptyset$; therefore $y_{\tau,\alpha}(\beta) \notin N(y)$, whence $\beta \notin N(x)$.

Conversely, let $x \in W_\alpha$ be such that $L_x \subseteq P_{\tau,\alpha}$. It is clear that $y_{\tau,\alpha}x(\alpha) = \theta(\tau)$. We use 2.3 (5): since $N(y_{\tau,\alpha}) \subseteq \Pi'(\tau)$, it suffices to prove that, for all $\beta \in N(x)$, $\text{Supp}(y_{\tau,\alpha}(\beta)) \cap \Pi'(\tau) \neq \emptyset$, and $y_{\tau,\alpha}(\beta) > 0$. By assumption, for each $\beta \in N(x)$, $\text{Supp}(\beta) \cap P_{\tau,\alpha} \neq \emptyset$. Since $y_{\tau,\alpha} \in W(\Delta(\tau))$, $y_{\tau,\alpha}$ acts on any root by summing linear combinations of roots in $\Pi(\tau)$. Since $P(\tau, \alpha) \cap \Pi(\tau) = \emptyset$, we obtain that $\text{Supp}(y_{\tau,\alpha}(\beta)) \cap P_{\tau,\alpha} \neq \emptyset$; moreover $y_{\tau,\alpha}(\beta) > 0$. But $\beta \perp \alpha$, hence $y_{\tau,\alpha}(\beta) \perp \theta(\tau)$. Since $\theta(\tau)$ has negative scalar product with all roots in $P_{\tau,\alpha}$, we obtain that $\text{Supp}(y_{\tau,\alpha}(\beta))$ also contains some root having positive scalar product with $\theta(\tau)$. By Lemma 5.5 (3) it follows that $\text{Supp}(y_{\tau,\alpha}(\beta)) \cap \Pi'(\tau) \neq \emptyset$. This implies the thesis. \square

Notice that if $\beta \in \Pi \setminus \Pi(\tau)$ and $\beta \not\perp \Delta(\tau)$, then $(\beta, \theta(\tau)) < 0$. Let $\{\Delta = \Delta_1 \supset \dots \supset \Delta_k\}$ be the good chain of τ . By Lemma 5.5, for all $\beta \in \Pi$ we have $(\beta, \theta(\tau)) < 0$ if and only if $\beta \in \Pi'_{k-1}$, where $\Pi'_{k-1} = \{\beta \in \Pi \mid (\beta, \theta_{k-1}) \neq 0\}$, θ_{k-1} being the highest root of Π_{k-1} . Thus we obtain that $P_{\tau,\alpha} = \Pi'_{k-1} \cap \alpha^\perp$. We set, for $\alpha \in \Delta_\ell^+$ and $\tau \in X^+$

$$W_{\alpha,\tau} = W((\Pi \cap \alpha^\perp) \setminus \Pi'_{k-1}).$$

As seen in 2.9, $W_{\alpha,\tau} \setminus W_\alpha$ is a poset with minimum and maximum. If w_0^α is the longest element of W_α , and $w_0^{\alpha,\tau}$ is the longest element of $W_{\alpha,\tau}$, then $w_0^{\alpha,\tau}w_0^\alpha$ corresponds to the maximal element of $W_{\alpha,\tau} \setminus W_\alpha$. We set

$$\check{y}_{\alpha,\tau} = y_{\alpha,\tau}w_0^{\alpha,\tau}w_0^\alpha.$$

We can now describe the poset structure of $\mathcal{I}_{ab}(\tau, \alpha)$.

Proposition 7.7. *Let $\tau \in X^+$, and $\alpha \in \Delta_\ell^+(\tau)$. Then the poset $\mathcal{I}_{ab}(\tau, \alpha)$ is isomorphic to $W_{\alpha,\tau} \setminus W_\alpha$. In particular $\mathcal{I}_{ab}(\tau, \alpha)$ has a unique maximal element and a unique minimal element. Moreover,*

$$\max \mathcal{I}_{ab}(\tau, \alpha) = i(t_\tau v_\tau \check{y}_{\alpha,\tau}).$$

Proof. Assume $i, j \in \mathcal{I}_{ab}(\tau, \alpha)$. Then by Proposition 6.6, $w_i = t_\tau v_\tau y$, $w_j = t_\tau v_\tau y'$, with $L_y, L_{y'} \subseteq \Pi'(\tau)$. Moreover $i \subseteq j$ if and only if $y \leq y'$. By Lemma 7.6, there exist $x, x' \in W_\alpha$ such that $L_x, L_{x'} \subseteq P_{\tau,\alpha}$, $y = y_{\tau,\alpha}x$, and $y' = y_{\tau,\alpha}x'$. Moreover, $y \leq y'$ if and only if $x \leq x'$. By 2.9, this implies the claim. \square

§8 MAXIMAL ELEMENTS IN $\mathcal{I}_{ab}(\tau, \alpha)$

In the previous section we have shown that $\mathcal{I}_{ab}(\tau, \alpha)$ is not empty if and only if $\alpha \in \Delta_\ell^+(\tau)$ and in this case it has unique minimal and maximal elements, but at this point we have no information about the relationships between $\mathcal{I}_{ab}(\tau, \alpha)$ and $\mathcal{I}_{ab}(\tau', \alpha)$. In this section we shall prove that $\max \mathcal{I}_{ab}(\tau, \alpha) \subseteq \max \mathcal{I}_{ab}(\tau', \alpha)$ if and only if $\tau \leq \tau'$. We remark that if $\tau \leq \tau'$, then $\max \mathcal{I}_{ab}(\tau, \alpha)$ and $\min \mathcal{I}_{ab}(\tau', \alpha)$ may be not comparable.

Let $\alpha \in \Delta_\ell^+$. We set

$$\text{Adj}(\alpha) = \{\alpha' \in \Pi \mid (\alpha, \alpha') \neq 0\};$$

$$\Phi_*(\alpha) = \{\beta \in \Delta \mid n_{\alpha'}(\beta) = m_{\alpha'}, \text{ for all } \alpha' \in \text{Adj}(\alpha)\};$$

$$\mathbf{i}_*(\alpha) = \sum_{\beta \in \Phi_*(\alpha)} \mathfrak{g}_\beta.$$

Notice that $\text{Adj}(\alpha) \cap \text{Supp}(\alpha) \neq \emptyset$, since $(\alpha, \alpha) \neq 0$.

Consider the parabolic subsystems of $\widehat{\Delta}$, Δ , generated by all simple roots which are orthogonal to α , $\Delta(\widehat{\Pi} \cap \alpha^\perp)$ and $\Delta(\Pi \cap \alpha^\perp)$, respectively. Recall that, for $\alpha \in \Delta_\ell^+$, we have set $W_\alpha = W(\Pi \cap \alpha^\perp)$. We also set:

$$\widehat{W}_\alpha = W(\widehat{\Pi} \cap \alpha^\perp), \quad \widehat{\Delta}_\alpha = \Delta(\widehat{\Pi} \cap \alpha^\perp), \quad \Delta_\alpha = \Delta(\Pi \cap \alpha^\perp).$$

It is clear that $\Delta_\alpha \subseteq \widehat{\Delta}_\alpha$ and that equality holds if and only if $\alpha \not\perp \theta$.

Lemma 8.1. (1) For all $\alpha \in \Delta_\ell^+$, $\mathbf{i}_*(\alpha)$ is an abelian ideal of \mathfrak{b} .
(2) Assume $\alpha \perp \theta$, and set $w_{*,\alpha} = w_{\mathbf{i}_*(\alpha)}$. Then

$$N(w_{*,\alpha}) = -\Phi_*(\alpha) + \delta = \widehat{\Delta}_\alpha^+ \setminus \Delta_\alpha^+;$$

in particular $w_{*,\alpha} \in \mathcal{W}_{ab} \cap \widehat{W}_\alpha$.

(3) If $x \in \widehat{W}_\alpha$ and $L_x \subseteq \{\alpha_0\}$, then $x \in \mathcal{W}_{ab}$ and $x \leq w_{*,\alpha}$. In particular, for all $x \in \mathcal{W}_{ab} \cap \widehat{W}_\alpha$, we have $x \leq w_{*,\alpha}$.

Proof.

(1) It is clear that $\Phi_*(\alpha) \subseteq \Delta^+$ and is an abelian set, i.e., for all $\beta, \beta' \in \Phi_*(\alpha)$, $\beta + \beta'$ is not a root. It is also clear that if $\beta \in \Phi_*(\alpha)$ and $\beta' \in \Delta^+$ is such that $\beta' \geq \beta$, then $\beta' \in \Phi_*(\alpha)$. This proves that $\mathbf{i}_*(\alpha)$ is an abelian ideal of \mathfrak{b} .

(2) The first equality follows from section 3 and holds even if $\alpha \not\perp \theta$. We prove the second equality.

Assume that $\alpha_0 \perp \alpha$ and $\beta \in \widehat{\Delta}_\alpha^+ \setminus \Delta_\alpha^+$. Then $\beta = c\alpha_0 + \sum_{\gamma \in \Pi \cap \alpha^\perp} c_\gamma \gamma$, with $c, c_\gamma \in \mathbb{N}$ and $c > 0$, hence $\beta = \overline{\beta} + c\delta$ with $\overline{\beta} = -c\theta + \sum_{\gamma \in \Pi \cap \alpha^\perp} c_\gamma \gamma$. Now $n_{\alpha'}(\overline{\beta}) = -cm_{\alpha'}$ for all $\alpha' \in \text{Adj}(\alpha)$. It follows that $c = 1$, hence that $-\overline{\beta} \in \Phi_*(\alpha)$, and $\beta \in -\Phi_*(\alpha) + \delta$. Conversely, if $\overline{\beta} \in \Delta^+$ is such that $n_{\alpha'} = m_{\alpha'}$ for all $\alpha' \in \text{Adj}(\alpha)$, then $-\beta + \delta - \alpha_0$ is either 0, or a non negative linear combination of roots in $\Pi \cap \alpha^\perp$, hence $-\beta + \delta \in \widehat{\Delta}_\alpha^+ \setminus \Delta_\alpha^+$.

(3) Assume $x \in \widehat{W}_\alpha$ and $L_x \subseteq \{\alpha_0\}$. Then $N(x) \subseteq \widehat{\Delta}_\alpha^+$; moreover, by 2.3 (3) we have that $N(x) \cap \Delta = \emptyset$, hence $N(x) \subseteq \widehat{\Delta}_\alpha^+ \setminus \Delta_\alpha^+$. By (2) and 2.6 we obtain that $x \leq w_{*,\alpha}$, and hence that $x \in \mathcal{W}_{ab}$. \square

Remark. Since for all $w \in \mathcal{W}_{ab}$ we have $L_w \subseteq \{\alpha_0\}$, by Lemma 8.1 (3) and 2.9 we obtain that the poset $\mathcal{W}_{ab} \cap \widehat{W}_\alpha$ is isomorphic to $W_\alpha \setminus \widehat{W}_\alpha$.

Lemma 8.2. *Let $\{\Delta_1 = \Delta \supset \dots \supset \Delta_k\}$ be a good chain, and $\Pi_i = \Delta_i \cap \Pi$ for $i = 1, \dots, k$. Assume that $k \geq 2$ and $\alpha \in \Pi_k$. Then $n_\alpha(\theta_{k-1}) = m_\alpha$.*

Proof. We assume by contradiction that $n_\alpha(\theta_{k-1}) < m_\alpha$. We shall prove that then for all $\beta \in \Pi_k$ we have that $n_\beta(\theta_{k-1}) < m_\beta$: this is impossible since, by 6.8 (4), for $\beta \in \Pi'_k$ we have $m_\beta = n_\beta(\theta_k) \leq n_\beta(\theta_{k-1})$.

Assume $\beta \in \Pi_k$. We first prove that, if $n_{\beta'}(\theta_{k-1}) < m_{\beta'}$ for some $\beta' \in \text{Adj}(\beta)$, then $n_\beta(\theta_{k-1}) < m_\beta$. Since $\beta \in \Delta_k$, we have $(\beta, \theta_i) = 0$ for all $i \in [k-1]$, hence $0 = (\beta, \theta) = \sum_{\beta' \in \text{Adj}(\beta)} m_{\beta'}(\beta, \beta')$, and $0 = (\beta, \theta_{k-1}) = \sum_{\beta' \in \text{Adj}(\beta)} n_{\beta'}(\theta_{k-1})(\beta, \beta')$.

Since $(\beta, \beta) > 0$ and $(\beta, \beta') < 0$ for all $\beta' \neq \beta$ in $\text{Adj}(\beta)$, we obtain that, if $n_{\beta'}(\theta_{k-1}) < m_{\beta'}$ for some $\beta' \neq \beta$ in $\text{Adj}(\beta)$, then $n_\beta < m_\beta$.

Now, for all $\beta \in \Pi_k$, there exists a sequence, $\beta_1 = \alpha, \dots, \beta_h = \beta$, in Π_k , such that $\beta_{i+1} \in \text{Adj}(\beta_i)$ for $i = 1, \dots, h-1$. If $n_\alpha(\theta_{k-1}) < m_\alpha$, we obtain inductively that $n_\beta(\theta_{k-1}) < m_\beta$. \square

Let $\alpha \in \Delta_\ell^+$. We recall that, by Remark 5.7 (3), there exists a unique maximal good chain $\{\Delta = \Delta_1 \supset \dots \supset \Delta_k\}$ such that $\alpha \in \Delta_i$ for $1 \leq i \leq k$: we shall call this chain the *maximal good chain of α* . Any good chain the elements of which contain α is an initial section of the maximal one.

Notation. Let $\alpha \in \Delta_\ell^+$. We denote by $\{\Delta = \Delta_1(\alpha) \supset \dots \supset \Delta_{k(\alpha)}(\alpha)\}$ the maximal good chain of α , and for $1 \leq h \leq k(\alpha)$, we denote by $\theta_h(\alpha)$ the highest root of $\Delta_h(\alpha)$. Moreover we set $\tau_0(\alpha) = 0$, and

$$\tau_h(\alpha) = \sum_{i=1}^h \theta_i(\alpha)^\vee, \quad \text{for } 1 \leq h \leq k(\alpha), \quad X_\alpha = \{\tau_h(\alpha) \mid 1 \leq h \leq k(\alpha)\}.$$

It is clear that if $\alpha \in \Delta_\ell^+$, then $\tau \in X_\alpha$ if and only if $\alpha \in \Delta_\ell^+(\tau)$. Thus by Remark 5.7 and Proposition 7.4 we have that $\mathcal{I}_{ab}(\tau, \alpha) \neq \emptyset$ if and only if $\tau \in X_\alpha$.

Proposition 8.3. *Let $\alpha \in \Delta_\ell^+$ and assume that $k(\alpha) \geq 2$. Then $-\theta_h(\alpha) + \delta \in \widehat{\Delta}_\alpha$, for all $h \in \{1, \dots, k(\alpha) - 1\}$. Equivalently, for $1 \leq h < k(\alpha)$, we have $n_\beta(\theta_h(\alpha)) = m_\beta$, for all $\beta \in \text{Adj}(\alpha)$.*

Proof. Set $k = k(\alpha)$. We shall prove that for all $\beta \in \text{Adj}(\alpha)$ we have $n_\beta(\theta_{k-1}) = m_\beta$; this clearly implies that $n_\beta(\theta_h) = m_\beta$ for all $h \in \{1, \dots, k-1\}$, which is equivalent to the thesis. Let $\beta \in \text{Adj}(\alpha)$. If $\beta \in \Pi_k$, then the claim follows directly from Lemma 8.2. Assume $\beta \notin \Pi_k$. We first show that $\beta \in \Pi_{k-1}$. If not, consider the maximal $i \in \{1, \dots, k-2\}$ such that $\beta \in \Pi_i$. Since $\alpha \in \Pi_{i+1}$, β cannot belong to any irreducible component of $\Delta_i \cap \theta_i^\perp$ different from Δ_{i+1} ; it follows that $\beta \in \Pi'_i$.

But then, by property (2) of Definition 5.4, $\beta \perp \Delta_{i+2}$: this is impossible, since $\alpha \in \Delta_{i+2}$ and $\beta \in \text{Adj}(\alpha)$. Thus we have $\beta \in \Pi_{k-1} \setminus \Pi_k$ and $\beta \not\perp \Delta_k$: this implies that $\beta \in \Pi'_{k-1}$, hence, by 6.8 (4), $n_\beta(\theta_{k-1}) = m_\beta$. \square

In the next Lemma we consider the element \check{w}_α defined in section 7.

Lemma 8.4. (1) *Let $\alpha \in \Delta_\ell^+$ and $x \in \mathcal{W}_{ab} \cap \widehat{W}_\alpha$. Then $\check{w}_\alpha x \in \mathcal{W}_{ab}$ and $\mathbf{i}(\check{w}_\alpha x) \supseteq \mathbf{i}(\check{w}_\alpha)$. Moreover, $\dim \mathbf{i}(\check{w}_\alpha x) = \ell(\check{w}_\alpha) + \ell(x)$.*
 (2) *Let $\alpha \in \Delta_\ell^+$ and $x \in \widehat{W}_\alpha$ be such that $\check{w}_\alpha x \in \mathcal{W}_{ab}$. Then $x \in \mathcal{W}_{ab} \cap \widehat{W}_\alpha$.*

Proof. (1) If $x \in \mathcal{W}_{ab} \cap \widehat{W}_\alpha$, then we have that $N(x) \subseteq \Delta^- + \delta$, and $N(x) \perp \alpha$. Consider $z \in N(x)$, $z = -\beta + \delta$, $\beta \in \Delta^+$. We have $\beta \perp \alpha$, so, by 7.1 (5), $\beta \notin N(y_\alpha^{-1})$; it follows that $y_\alpha N(x) \subseteq \Delta^- + \delta$. Moreover, $y_\alpha N(x) \perp \theta$, therefore $\check{w}_\alpha N(x) = s_0 y_\alpha N(x) = y_\alpha N(x)$. In particular, $\check{w}_\alpha N(x) \in \widehat{\Delta}^+$, hence by 2.7 $\check{w}_\alpha \leq \check{w}_\alpha x$. Moreover, $N(\check{w}_\alpha x) = N(\check{w}_\alpha) \cup \check{w}_\alpha N(x) \subseteq \Delta^- + \delta$, hence $\check{w}_\alpha x \in \mathcal{W}_{ab}$. By 3.2 we obtain that $\mathbf{i}(\check{w}_\alpha) \subseteq \mathbf{i}(\check{w}_\alpha x)$. Moreover, we have $\dim \mathbf{i}(\check{w}_\alpha x) = \ell(\check{w}_\alpha) + \ell(x)$.

(2) By 8.1 (3) it suffices to prove that $L_x \subseteq \{\alpha_0\}$. If $L_x \not\subseteq \{\alpha_0\}$, then there exists $\beta \in \Pi \cap \alpha^\perp$ such that $\beta \in N(x)$. By Proposition 7.3 (3) we have $\check{w}_\alpha(\beta) > 0$, hence $\check{w}_\alpha(\beta) \in N(\check{w}_\alpha x) \subseteq \Delta^- + \delta$. It follows that $\check{w}_\alpha s_\beta \in \mathcal{W}_{ab}$ and $\mathbf{i}(\check{w}_\alpha s_\beta) \supsetneq \mathbf{i}(\check{w}_\alpha)$. But $\mathbf{i}(\check{w}_\alpha s_\beta) \in \mathcal{I}_{ab}(\theta^\vee, \alpha)$, and, by 7.3 (1), $\mathcal{I}_{ab}(\theta^\vee, \alpha) = \{\mathbf{i}(\check{w}_\alpha)\}$, thus we get a contradiction. \square

Lemma 8.5. *Let $\alpha \in \Delta_\ell^+$. Assume $\tau \in X$, $x \in \mathcal{W}_{ab}(\tau) \cap \widehat{W}_\alpha$ and set $2h = \frac{(\theta, \theta)}{2}(\tau, \tau)$. Then:*

- (1) $h \in \{0, \dots, k(\alpha) - 1\}$ and $\check{w}_\alpha x \in \mathcal{W}_{ab}(\tau_{h+1}(\alpha), \alpha)$;
- (2) $y_\alpha(\tau) = \theta_2(\alpha)^\vee + \dots + \theta_{h+1}(\alpha)^\vee$.

Proof. By 8.4 (1), $\check{w}_\alpha x \in \mathcal{W}_{ab}$; moreover, it is clear that $\check{w}_\alpha x(\alpha) = -\theta + 2\delta$. Let $x = t_\tau v$, with $v \in W$. Then $\check{w}_\alpha x \in \mathcal{I}_{ab}(\theta^\vee + y_\alpha(\tau), \alpha)$. By Proposition 7.4 and Remark 5.7 (3), we obtain that $\theta^\vee + y_\alpha(\tau) = \tau_j(\alpha)$ for some $j \in \{1, \dots, k(\alpha)\}$. Since $\alpha \perp \tau$, we have that $\theta \perp y_\alpha(\tau)$, hence $\frac{(\theta, \theta)}{2}(\theta^\vee + y_\alpha(\tau), \theta^\vee + y_\alpha(\tau)) = 2(h+1)$. By Remark 5.7 (1), it follows that $\theta^\vee + y_\alpha(\tau) = \tau_{h+1}(\alpha)$ and hence that $1 \leq h+1 \leq k(\alpha)$, so that $h \in \{0, \dots, k(\alpha) - 1\}$. Moreover $y_\alpha(\tau) = \theta_2(\alpha)^\vee + \dots + \theta_{h+1}(\alpha)^\vee$. \square

Notation. Let $\alpha \in \Delta_\ell^+$. If $k(\alpha) > 1$ and $h \in [k(\alpha) - 1]$, we set

$$\Psi_h(\alpha) = \{\beta \in \Delta_h(\alpha) \mid \beta = \theta_h(\alpha), \text{ or } \theta_h(\alpha) - \beta \in \Delta_\alpha\};$$

$$\Phi_h(\alpha) = \bigcup_{i=1}^h \Psi_i(\alpha); \quad \mathbf{i}_h(\alpha) = \sum_{\beta \in \Phi_h(\alpha)} \mathfrak{g}_\beta.$$

Moreover, we set $\Phi_0(\alpha) = \emptyset$, and $\mathbf{i}_0(\alpha)$ to be the zero ideal. By Proposition 8.3 we directly obtain that $\Phi_h(\alpha) \subseteq \Phi_*(\alpha)$ for all $h \in [k(\alpha) - 1]$. We remark that the subsets $\Psi_h(\alpha)$, $h \in [k(\alpha) - 1]$, are pairwise disjoint.

Lemma 8.6. *Let $\alpha \in \Delta_\ell^+$. Then $\mathfrak{i}_h(\alpha)$ is an abelian ideal of \mathfrak{b} , for $0 \leq h < k(\alpha)$. Set $\check{w}_{h,\alpha} = w_{\mathfrak{i}_h(\alpha)}$. Then $\check{w}_{h,\alpha} \in \mathcal{W}_{ab}(\tau_h(\alpha)) \cap \widehat{W}_\alpha$. In particular $\check{w}_\alpha \check{w}_{h,\alpha} \in \mathcal{W}_{ab}(\tau_{h+1}(\alpha), \alpha)$ and $\dim \mathfrak{i}(\check{w}_\alpha \check{w}_{h,\alpha}) = \ell(\check{w}_\alpha) + \ell(\check{w}_{h,\alpha})$.*

Proof. We set $k = k(\alpha)$; $\Delta_h = \Delta_h(\alpha)$, $\theta_h = \theta_h(\alpha)$, $\tau_h = \tau_h(\alpha)$, and $\Phi_h = \Phi_h(\alpha)$, for $0 \leq h < k$.

If $h = 0$, the claim is trivial. Assume $k \geq 2$ and $1 \leq h \leq k - 1$. Since $\Phi_h(\alpha) \subseteq \Phi_*(\alpha)$, by Lemma 8.1 (1) we obtain that $\mathfrak{i}_h(\alpha)$ is an abelian subalgebra.

Then let $\beta \in \Phi_h$ and $\beta' \in \Delta^+$ be such that $\beta' \geq \beta$. We shall prove that $\beta' \in \Phi_h$; this will imply that $\mathfrak{i}_h(\alpha)$ is an abelian ideal. It is clear that $n_\gamma(\beta') = m_\gamma$ for all $\gamma \in \text{Adj}(\alpha)$. Let $i \in [h]$ be such that $\beta \in \Delta_i$ and $\theta_i - \beta \in \Delta_\alpha \cup \{0\}$. Assume first $\beta' \in \Delta_i$. Then, since θ_i is dominant in Δ_i , we have $(\theta_i, \beta') \geq (\theta_i, \beta) > 0$. Hence either $\beta' = \theta_i$, or $\theta_i - \beta' \in \Delta$. In the latter case, since $\theta_i - \beta' \leq \theta_i - \beta$, we have that $\theta_i - \beta' \in \Delta_\alpha$; therefore $\beta' \in \Phi_h$. Next assume $\beta' \notin \Delta_i$ and let j , $1 \leq j \leq i - 1$, be such that $\beta' \in \Delta_j$ and $\beta' \notin \Delta_{j+1}$. If $\beta' \perp \theta_j$, then β' is included in some connected component of $\Delta_j \cap \theta_j^\perp$. But, for all $\alpha' \in \text{Adj}(\alpha)$, we have $n_{\alpha'}(\beta') = m_{\alpha'} > 0$; moreover $\text{Adj}(\alpha) \cap \text{Supp}(\alpha) \neq \emptyset$, and $\text{Supp}(\alpha) \in \Delta_{j+1}$, hence we obtain that $\beta' \in \Delta_{j+1}$, a contradiction. Hence $\beta' \not\perp \theta_j$. It follows that either $\beta' = \theta_j$, or $\theta_j - \beta' \in \Delta$, and, indeed, $\theta_j - \beta' \in \Delta_\alpha$. Thus $\beta' \in \Phi_h$. This concludes the proof that $\mathfrak{i}_h(\alpha) \in \mathcal{I}_{ab}$.

By definition, $\theta_i \in \Phi_h$ for all $i \in [h]$; moreover $\beta \notin \Phi_h$ for all $\beta \in \Delta_h \cap \theta_h^\perp$. It follows that $-\theta_i + \delta \in N(\check{w}_{h,\alpha})$ for all $i \in [h]$, and $-\beta + \delta \notin N(\check{w}_{h,\alpha})$ for all $\beta \in \Delta_h \cap \theta_h^\perp$. By Proposition 6.2, it follows that $\check{w}_{h,\alpha} \in \mathcal{W}_{ab}(\tau_h)$.

If $\beta \in \Psi_j(\alpha)$, for some $j \in [h]$, then, by definition, $-\beta + \delta - (-\theta_j + \delta) \in \Delta_\alpha$, and since, by 8.3, $-\theta_j + \delta \in \widehat{\Delta}_\alpha$, we obtain that $-\beta + \delta \in \widehat{\Delta}_\alpha$. Since $N(\check{w}_{h,\alpha}) = \{-\beta + \delta \mid \beta \in \Phi_h\}$, we obtain that $N(\check{w}_{h,\alpha}) \subseteq \widehat{\Delta}_\alpha$, hence that $\check{w}_{h,\alpha} \in \widehat{W}_\alpha$.

By 8.5 and 8.4 (1) we finally obtain that $\check{w}_\alpha \check{w}_{h,\alpha} \in \mathcal{W}_{ab}(\tau_{h+1}(\alpha), \alpha)$, and $\dim \mathfrak{i}(\check{w}_\alpha \check{w}_{h,\alpha}) = \ell(\check{w}_\alpha) + \ell(\check{w}_{h,\alpha})$. \square

Remark. For $\alpha \in \Delta_\ell^+$ and $0 \leq h < k(\alpha)$, let $\check{w}_{h,\alpha} = w_{\mathfrak{i}_h(\alpha)}$, as in the statement of 8.6. Notice that $\mathfrak{i}_0(\alpha)$ is the zero ideal, hence $\check{w}_{0,\alpha} = 1$. Notice, moreover, that, for $h > 0$

$$N(\check{w}_{h,\alpha}) = -\Phi_h(\alpha) + \delta.$$

Proposition 8.7. *Let $\alpha \in \Delta_\ell^+$, $0 \leq h < k(\alpha)$, and $\mathfrak{i} \in \mathcal{I}_{ab}(\tau_{h+1}(\alpha), \alpha)$. Then $\mathfrak{i} \supseteq \mathfrak{i}(\check{w}_\alpha)$, and $\check{w}_\alpha^{-1} \mathfrak{i} \in \mathcal{W}_{ab}(\tau_h(\alpha)) \cap \widehat{W}_\alpha$.*

Proof. Set $\tau = \tau_{h+1}(\alpha)$. By Lemma 8.6 we have that $\check{w}_\alpha \check{w}_{h,\alpha} \in \mathcal{W}_{ab}(\tau, \alpha)$. Set $j = \min \mathcal{I}_{ab}(\tau, \alpha)$. By Proposition 7.5, for each $w \in \mathcal{W}_{ab}(\tau, \alpha)$ we have that $w = w_j y$ with $y \in W_\alpha$. In particular, for all $w, w' \in \mathcal{W}_{ab}(\tau, \alpha)$ we have that $w^{-1} w' \in W_\alpha$. Thus $w_{\mathfrak{i}} = \check{w}_\alpha \check{w}_{h,\alpha} y$, with $y \in W_\alpha$, and therefore $\check{w}_{h,\alpha} y \in \widehat{W}_\alpha$. Then by Lemma 8.4 (2) we obtain that $\check{w}_{h,\alpha} y \in \mathcal{W}_{ab} \cap \widehat{W}_\alpha$, and it is clear that $\check{w}_{h,\alpha} y \in \mathcal{W}_{ab}(\tau_h(\alpha), \alpha)$. Moreover, by 8.4 (1), $\mathfrak{i} \supseteq \mathfrak{i}(\check{w}_\alpha)$. \square

For $\alpha \in \Delta_\ell^+$ we set

$$\mathcal{I}_{ab}(\alpha) = \bigcup_{\tau \in X_\alpha} \mathcal{I}_{ab}(\tau, \alpha).$$

These posets have been studied also in [P], [S]. If $\mathbf{i} \in \mathcal{I}_{ab}$, then $\mathbf{i} \in \mathcal{I}_{ab}(\alpha)$ if and only if $w_{\mathbf{i}}(\alpha) = -\theta + 2\delta$. We can now obtain the following results.

Corollary 8.8. *(Suter) The poset $\mathcal{I}_{ab}(\alpha)$ is isomorphic to $W_\alpha \setminus \widehat{W}_\alpha$. In particular, it has a unique minimal element and a unique maximal element. Moreover,*

$$\min \mathcal{I}_{ab}(\alpha) = \mathbf{i}(\check{w}_\alpha), \quad \max \mathcal{I}_{ab}(\alpha) = \mathbf{i}(\check{w}_\alpha w_{*,\alpha}).$$

Proof. By Proposition 8.7 and Lemma 8.4, $\mathbf{i} \in \mathcal{I}_{ab}(\alpha)$ if and only if $w_{\mathbf{i}} = \check{w}_\alpha x$ with $x \in \mathcal{W}_{ab} \cap \widehat{W}_\alpha$; moreover, in this case, $N(\check{w}_\alpha x) = N(\check{w}_\alpha) \cup \check{w}_\alpha N(x)$. Therefore, if $\mathbf{i}, \mathbf{j} \in \mathcal{I}_{ab}(\alpha)$, $w_{\mathbf{i}} = \check{w}_\alpha x$, and $w_{\mathbf{j}} = \check{w}_\alpha y$, with $x, y \in \mathcal{W}_{ab} \cap \widehat{W}_\alpha$, then $\mathbf{i} \subseteq \mathbf{j}$ if and only if $x \leq y$. By Lemma 8.1 and the subsequent remark we get the claim. \square

Corollary 8.9. *Assume $\mathbf{i} \in \mathcal{I}_{ab}$, $\alpha \in \Delta_\ell^+$ and $\tau \in X_\alpha$.*

- (1) $\mathbf{i} \in \mathcal{I}_{ab}(\alpha)$ if and only if $\min \mathcal{I}_{ab}(\alpha) \subseteq \mathbf{i} \subseteq \max \mathcal{I}_{ab}(\alpha)$.
- (2) $\mathbf{i} \in \mathcal{I}_{ab}(\tau, \alpha)$ if and only if $\min \mathcal{I}_{ab}(\tau, \alpha) \subseteq \mathbf{i} \subseteq \max \mathcal{I}_{ab}(\tau, \alpha)$.

Proof. (1) The “only if” part is a trivial consequence of Corollary 8.8. Conversely, assume that $\min \mathcal{I}_{ab}(\alpha) \subseteq \mathbf{i} \subseteq \max \mathcal{I}_{ab}(\alpha)$. Then there exists $x \in W$ such that $w_{\mathbf{i}} = \check{w}_\alpha x$ and $N(w_{\mathbf{i}}) = N(\check{w}_\alpha) \cup \check{w}_\alpha N(x)$. Since $N(w_{\mathbf{i}}) \subseteq N(\check{w}_\alpha w_{*,\alpha})$, it follows that $N(x) \subseteq N(w_{*,\alpha}) \subseteq \widehat{\Delta}_\alpha$; therefore $x(\alpha) = \alpha$ and $w_{\mathbf{i}}(\alpha) = \check{w}_\alpha(\alpha) = -\theta + 2\delta$.

(2) The “only if” part follows from Propositions 7.5 and 7.7. Conversely, assume $\min \mathcal{I}_{ab}(\tau, \alpha) \subseteq \mathbf{i} \subseteq \max \mathcal{I}_{ab}(\tau, \alpha)$, and let $\tau = \tau_h(\alpha)$, $h \in [k(\alpha)]$. By item (1), $\mathbf{i} \in \mathcal{I}_{ab}(\alpha)$, hence $\mathbf{i} \in \mathcal{I}_{ab}(\tau_{h'}(\alpha), \alpha)$, for some $h' \in [k(\alpha)]$. By Proposition 6.2, $\mathfrak{g}_{\theta_{\mathbf{i}}(\alpha)} \subset \mathbf{i}$ if and only if $i \in [h']$; similarly, $\mathfrak{g}_{\theta_{\mathbf{i}}(\alpha)}$ is contained in $\min \mathcal{I}_{ab}(\tau, \alpha)$ and $\max \mathcal{I}_{ab}(\tau, \alpha)$, hence in \mathbf{i} , if and only if $i \in [h]$. Hence $h = h'$. \square

We set

$$\mathbf{m}(\alpha) = \max \mathcal{I}_{ab}(\alpha),$$

thus $\mathbf{m}(\alpha) = \mathbf{i}(\check{w}_\alpha w_{*,\alpha})$.

Corollary 8.10. *(Suter). For all $\alpha \in \Delta_\ell^+$*

$$\dim \mathbf{m}(\alpha) = g - 1 + \frac{1}{2} \left(|\widehat{\Delta}_\alpha| - |\Delta_\alpha| \right).$$

If, moreover, $\alpha \perp \theta$, denote by $\widehat{\Delta}_1(\alpha)$ the connected component of $\widehat{\Delta}_\alpha$ which includes α_0 and set $\widehat{\Delta}_1^0(\alpha) = \widehat{\Delta}_1(\alpha) \cap \Delta$. Then

$$\dim \mathbf{m}(\alpha) = g - 1 + \frac{1}{2} \left(|\widehat{\Delta}_1(\alpha)| - |\widehat{\Delta}_1^0(\alpha)| \right).$$

By Proposition 7.7, if $\mathcal{I}_{ab}(\tau, \alpha)$ is not empty, then it has a unique maximal element. For all $1 \leq h \leq k(\alpha)$, we set

$$\mathbf{m}_h(\alpha) = \max \mathcal{I}_{ab}(\tau_h(\alpha), \alpha).$$

We can now explicitly describe $\mathbf{m}_h(\alpha)$.

Proposition 8.11. *For all $\alpha \in \Delta_\ell^+$, and $0 \leq h < k(\alpha)$, we have*

$$\mathfrak{m}_{h+1}(\alpha) = \mathfrak{i}(\check{w}_\alpha \check{w}_{h,\alpha}).$$

Proof. Set $\tau^- = \tau_h(\alpha)$. By Proposition 8.7, $w_{\mathfrak{m}_{h+1}(\alpha)} = \check{w}_\alpha x$, with $x \in \mathcal{W}_{ab}(\tau^-) \cap \widehat{W}_\alpha$. By Lemma 8.6 it suffices to prove that $\check{w}_\alpha x \leq \check{w}_\alpha \check{w}_{h,\alpha}$, and this is equivalent to prove that $N(x) \subseteq N(\check{w}_{h,\alpha})$. Recall that $N(\check{w}_{h,\alpha}) = \{-\beta + \delta \mid \beta \in \Phi_h(\alpha)\}$. Assume $\beta \in N(x)$ and $\beta = -\bar{\beta} + \delta$, with $\bar{\beta} \in \Delta^+$. We shall prove that $\bar{\beta} \in \Phi_h(\alpha)$.

We set $\Delta_i = \Delta_i(\alpha)$, and $\theta_i = \theta_i(\alpha)$, for $1 \leq i \leq h$. Since $\Delta_h = \Delta(\tau^-)$, by Proposition 6.2 we have that $\bar{\beta} \notin \Delta_h \cap \theta_h^\perp$, in particular $\bar{\beta} \notin \Delta_{h+1}$. Let $i \in [h]$ be such that $\bar{\beta} \in \Delta_i$ and $\bar{\beta} \notin \Delta_{i+1}$; then $\bar{\beta} \in \Delta_j$ for $1 \leq j \leq i$. We first prove that there exists $j \in \{1, \dots, i\}$ such that $(\bar{\beta}, \theta_j) > 0$. We assume, by contradiction, that $(\bar{\beta}, \theta_j) = 0$ for $1 \leq j \leq i$. Since $\bar{\beta} \notin \Delta_{i+1}$, we obtain that $\bar{\beta}$ belongs to a connected component of $\Delta_i \cap \theta_i^\perp$ distinct from Δ_{i+1} , in particular $\bar{\beta} \perp \Delta_{i+1}$, and hence $\bar{\beta} \perp \theta_j$ also for $j \geq i+1$. But this implies that $\bar{\beta} \perp \tau^-$, which is impossible, by 6.1 (4). Therefore there exists $j \in \{1, \dots, i\}$ such that $(\bar{\beta}, \theta_j) > 0$. Then either $\bar{\beta} = \theta_j$, or $\theta_j - \bar{\beta} \in \Delta^+$. Moreover, since $-\theta_j + \delta$ and β belong to $\widehat{\Delta}_\alpha$, we have that $\theta_j - \bar{\beta} = \beta - (-\theta_j + \delta) \in \widehat{\Delta}_\alpha \cup \{0\}$. But $\theta_j - \bar{\beta} \in \Delta \cup \{0\}$, therefore $\theta_j - \bar{\beta} \in \Delta_\alpha \cup \{0\}$. This means that $\bar{\beta} \in \Phi_h(\alpha)$. \square

For any good chain $\{\Delta = \Delta_1 \subset \dots \subset \Delta_k\}$, we have that $\Pi_i \cup \{-\theta_i + \delta\}$ is a basis for a subsystem $\widehat{\Delta}_i$ of $\widehat{\Delta}$ which can be identified with the affine root system of Δ_i .

Let $\alpha \in \Delta_\ell^+$, $k = k(\alpha)$, and $\Delta_i = \Delta_i(\alpha)$, for $i \in [k]$. We consider the parabolic subsystem of $\widehat{\Delta}_i$ generated by $(\Pi_i \cup \{-\theta_i + \delta\}) \cap \alpha^\perp$: this is a finite and not necessarily irreducible system. If $1 \leq i < k$, then $-\theta_i + \delta \perp \alpha$, and we denote by $\widehat{\Delta}_i(\alpha)$ the irreducible component of $\Delta((\Pi_i \cup \{-\theta_i + \delta\}) \cap \alpha^\perp)$ which includes $-\theta_i + \delta$. By 8.3, $\widehat{\Delta}_i(\alpha) \subseteq \widehat{\Delta}_\alpha$. Notice that if $\alpha \perp \theta$, then $\widehat{\Delta}_1(\alpha)$ is equal to the system $\widehat{\Delta}_1$ which occurs in 8.10.

If $k(\alpha) > 1$ and $1 \leq h \leq k(\alpha) - 1$, we set

$$\overline{\Psi}_h(\alpha) = \{\beta \in \widehat{\Delta}_h(\alpha)^+ \mid (\beta, (-\theta_h + \delta)^\vee) = -1\} \cup \{0\}.$$

Lemma 8.12. *Let $1 \leq h < k(\alpha)$. We have*

$$\overline{\Psi}_h(\alpha) = \{\theta_h - \beta \mid \beta \in \Psi_h(\alpha)\}.$$

In particular,

$$|\Psi_h(\alpha)| = |\overline{\Psi}_h(\alpha)| = g_h(\alpha) - 1,$$

where $g_h(\alpha)$ is the dual Coxeter number of $\widehat{\Delta}_h(\alpha)$.

Proof. Since $\widehat{\Delta}_h(\alpha)^+ \subseteq \widehat{\Delta}_\alpha^+$, by Lemma 8.1 we obtain that $\widehat{\Delta}_h(\alpha)^+ \setminus \Delta \subseteq -\Phi_*(\alpha) + \delta$; in particular $\widehat{\Delta}_h(\alpha)^+ \setminus \Delta$ is an abelian set. If $\beta \in \widehat{\Delta}_h(\alpha)^+$ and $(\beta, (-\theta_h + \delta)^\vee) = -1$, then $\beta + (-\theta_h + \delta)$ is a root, and therefore $\beta \in \Delta$. Then it is clear that $\theta_h - \beta \in \Delta_h$, hence, by definition, $\theta_h - \beta \in \Psi_h(\alpha)$.

Conversely, assume $\beta \in \Psi_h(\alpha)$ and $\beta \neq \theta_h$. Then it is clear that $(\theta_h - \beta, (-\theta_h + \delta)^\vee) = -1$; moreover, $\theta_h - \beta$ is positive and belongs, by definition, to Δ_α . It follows that $\theta_h - \beta \in \overline{\Psi}_h(\alpha)$.

By Proposition 7.1 (3) and (5) we have that $|\overline{\Psi}_h(\alpha)| = g_h(\alpha) - 1$. \square

Theorem 8.13. *Let $\alpha \in \Delta_\ell^+$. Then $\mathfrak{m}_h(\alpha) \subset \mathfrak{m}_{h+1}(\alpha)$, for $1 \leq h < k(\alpha)$, and $\mathfrak{m}_{k(\alpha)}(\alpha) = \mathfrak{m}(\alpha)$. Moreover,*

$$\dim \mathfrak{m}_h(\alpha) = g - 1 + \sum_{i=1}^{h-1} (g_i(\alpha) - 1).$$

Proof. By Proposition 8.11, $w_{\mathfrak{m}_{h+1}(\alpha)} = \check{w}_\alpha \check{w}_{h,\alpha}$, for $0 \leq h < k(\alpha)$; moreover, by Lemma 8.6, $N(\check{w}_\alpha \check{w}_{h,\alpha}) = N(\check{w}_\alpha) \cup \check{w}_\alpha N(\check{w}_{h,\alpha})$. We have $N(w_{0,\alpha}) = \emptyset$, and $N(\check{w}_{h,\alpha}) = -\Phi_h(\alpha) + \delta$ for $1 \leq h < k(\alpha)$; since $\Phi_{h-1}(\alpha) \subset \Phi_h(\alpha)$, we obtain that $\check{w}_\alpha w_{h-1,\alpha} < \check{w}_\alpha \check{w}_{h,\alpha}$, for $1 \leq h < k(\alpha)$. Therefore $\mathfrak{m}_h(\alpha) \subset \mathfrak{m}_{h+1}(\alpha)$. It follows that $\mathfrak{m}_{k(\alpha)}(\alpha) = \max \mathcal{I}_{ab}(\alpha)$, hence, by Corollary 8.8, $\mathfrak{m}_{k(\alpha)}(\alpha) = \mathfrak{m}(\alpha)$.

By Lemma 8.6, we have that $\dim \mathfrak{m}_h(\alpha) = \ell(\check{w}_\alpha) + \ell(w_{h-1,\alpha})$. By 7.1 (3), $\ell(\check{w}_\alpha) = g - 1$. Moreover, $|N(w_{h-1,\alpha})| = |\Phi_{h-1}(\alpha)| = \left| \bigcup_{i=1}^{h-1} \Psi_i \right|$, and since the sets $\Psi_i(\alpha)$ are pairwise disjoint, by 8.12, we obtain that $\ell(w_{h-1,\alpha}) = \sum_{i=1}^{h-1} (g_i(\alpha) - 1)$. \square

Lemma 8.14. *Assume $\alpha \in \Pi_\ell(\tau)$, $\tau \in X_\alpha$, and $\mathfrak{j} \in \mathcal{I}_{ab}$. If there exists $\mathfrak{i} \in \mathcal{I}_{ab}(\tau, \alpha)$, such that $\mathfrak{j} \supseteq \mathfrak{i}$, then $w_{\mathfrak{i}}^{-1} w_{\mathfrak{j}} \in \widehat{W}_\alpha$. In particular $w_{\mathfrak{j}}(\alpha) = -\theta + 2\delta$.*

Proof. By assumption $w_{\mathfrak{j}} \geq w_{\mathfrak{i}}$, in the weak order, hence there exists $x \in \widehat{W}$ such that $w_{\mathfrak{j}} = w_{\mathfrak{i}}x$ and $\ell(w_{\mathfrak{j}}) = \ell(w_{\mathfrak{i}}) + \ell(x)$. Moreover $N(w_{\mathfrak{j}}) = N(w_{\mathfrak{i}}) \cup w_{\mathfrak{i}}N(x)$, with $w_{\mathfrak{i}}N(x) \subseteq \widehat{\Delta}^+$, and indeed, since $w_{\mathfrak{j}} \in \mathcal{W}_{ab}$, $w_{\mathfrak{i}}N(x) \subseteq \Delta^- + \delta$. Assume by contradiction that $x \notin \widehat{W}_\alpha$ and set $x = x's_\beta x''$, with $x' \in \widehat{W}_\alpha$, $\beta \in \widehat{\Pi} \setminus \alpha^\perp$, and $\ell(x) = \ell(x') + 1 + \ell(x'')$, so that $x'(\beta) \in N(x)$. Notice that then $\beta \neq \alpha$, otherwise we have $\alpha = x'(\alpha) \in N(x)$ and hence $-\theta + 2\delta = w_{\mathfrak{i}}(\alpha) \in N(w_{\mathfrak{j}})$, which is impossible. Thus, since α is simple and long, we have $(\beta, \alpha^\vee) = -1$, hence $(w_{\mathfrak{i}}x'(\beta), w_{\mathfrak{i}}x'(\alpha)) = (w_{\mathfrak{i}}x'(\beta), -\theta^\vee + 2\delta) = -1$. But this implies that $w_{\mathfrak{i}}x'(\beta) \in \Delta^+ + \mathbb{Z}\delta$, which is impossible since $w_{\mathfrak{i}}x'(\beta) \in N(w_{\mathfrak{j}}) \subseteq \Delta^- + \delta$. We have therefore that $x \in \widehat{W}_\alpha$. \square

Notice that if $\alpha \in \Delta_\ell(\tau)$ and $\tau = \theta_1^\vee + \dots + \theta_k^\vee$ is the standard orthogonal decomposition of τ , then $\tau = \tau_k(\alpha)$.

Corollary 8.15. *Assume $\tau \in X^+$, $\alpha, \beta \in \Pi_\ell(\tau)$, $\alpha \neq \beta$, $\mathfrak{i} \in \mathcal{I}_{ab}(\tau, \alpha)$, and $\mathfrak{j} \in \mathcal{I}_{ab}(\tau, \beta)$. Then $\mathfrak{i}, \mathfrak{j}$ are not comparable. In particular, if $\tau = \theta_1^\vee + \dots + \theta_k^\vee$ is the standard orthogonal decomposition of τ , the maximal elements of $\mathcal{I}_{ab}(\tau)$ are exactly the ideals $m_k(\alpha)$ with $\alpha \in \Pi_\ell(\tau)$.*

Proof. Assume by contradiction $\mathfrak{i} \subseteq \mathfrak{j}$. Then by 8.14 $\mathfrak{j} \in \mathcal{I}_{ab}(\tau, \alpha)$, which is absurd. The last assertion follows from Proposition 6.7. \square

Remark. It would be interesting to investigate the inclusion relationships between ideals in $\mathcal{I}_{ab}(\tau, \alpha)$, $\mathcal{I}_{ab}(\tau, \beta)$ for $\alpha, \beta \in \Delta_\ell^+(\tau)$. Related results can be found in [P, Corollary 3.3] and in [St2]; we hope to deal with this problem in a subsequent paper.

Proposition 8.16. *If $\alpha \in \Pi_\ell$, then $\mathfrak{m}(\alpha)$ is a maximal abelian ideal of \mathfrak{b} .*

Proof. If $\mathfrak{i} \supseteq \mathfrak{m}(\alpha)$, then by Lemma 8.14 $w_{\mathfrak{i}} \in \mathcal{I}_{ab}(\tau, \alpha)$ for some $\tau \in X_\alpha$, hence by Theorem 8.13 $\mathfrak{i} \subseteq \mathfrak{m}(\alpha)$. \square

Remark. Notice that if $\tau = \tau_{k(\alpha)}(\alpha)$, then $\alpha \not\perp \tau$; conversely, if $\tau \in X_\alpha$ and $\tau = \tau_h(\alpha)$ with $1 \leq h < k(\alpha)$, then $\alpha \perp \tau$. Hence the maximal ideals of \mathfrak{b} which belong to $\mathcal{I}_{ab}(\tau)$ are exactly the $\mathfrak{m}(\alpha)$ with $\alpha \in \Pi_\ell(\tau)$ and $\alpha \not\perp \tau$.

Corollary 8.17. *(Panyushev). The map $\alpha \mapsto \mathfrak{m}(\alpha)$ is a bijection between long simple roots and maximal abelian ideals of \mathfrak{b} .*

Proof. The claim follows from Proposition 8.16, Corollary 8.15, and Theorems 4.4, 8.13. \square

§9 DIMENSION FORMULAS FOR $\mathfrak{m}_h(\alpha)$

In this section we determine the dimension of $\mathfrak{m}_h(\alpha)$, for all $\alpha \in \Pi_\ell$ and $1 \leq h \leq k(\alpha)$. This calculation has the following interesting consequence. It is easy to see that the maximal dimension d of a maximal commutative subalgebra of \mathfrak{g} coincides with the maximal dimension of an abelian ideal of \mathfrak{b} (for a proof see [R, §7]). Hence

$$d = \max_{\alpha \in \Pi_\ell} \{ \dim \mathfrak{m}(\alpha) \}.$$

As expected from Theorem 8.13 we shall see that the maximum is obtained when α is a long simple root having maximal distance from the affine node in the extended Dynkin diagram of Δ . The dimensions $\dim \mathfrak{m}_h(\alpha)$ will be uniformly calculated using Theorem 8.13: these calculations afford, as a special case, a conceptual explanation of the case by case calculations in [PR, Tables I, II], and in turn, a proof of Malcev's classical result on the dimension of a maximal commutative subalgebra [M]. We use for Dynkin diagrams the numbering of [B].

Before looking at the ideals $\mathfrak{m}_h(\alpha)$ for each type of root system, we note that, in general, for any long simple root α we have $\mathfrak{m}_1(\alpha) = \max \mathcal{I}_{ab}(\theta^\vee, \alpha)$ and $\dim \mathfrak{m}_1(\alpha) = g - 1$.

Type A_n

In this case the element of X^+ are linearly ordered and are $\lfloor \frac{n+1}{2} \rfloor$ in number. Indeed, there exists a unique maximal good chain $\{\Delta_1 = \Delta \supset \cdots \supset \Delta_{\lfloor \frac{n+1}{2} \rfloor}\}$; moreover, any good chain is an initial section of the maximal one. We have $\hat{\Delta}_i = \Delta(\alpha_i, \dots, \alpha_{n-i+1}) \cong A_{2n-i+2}$; if θ_i^\vee is the highest root of Δ_i and $\tau_i = \theta_1^\vee + \cdots + \theta_i^\vee$ we have $\tau_i = \omega_i^\vee + \omega_{n-i+1}^\vee$. Clearly, $\tau_i < \tau_j$ when $i < j$. If $\alpha_h \in \Delta_i$, then $\hat{\Delta}_k(\alpha_h) \cong A_{n-2k}$ for $1 \leq k \leq i-1$. Hence for $1 \leq i \leq \lfloor \frac{n+1}{2} \rfloor$ and $i \leq h \leq n-i+1$

$$\dim (\mathfrak{m}_i(\alpha_h)) = n + \sum_{k=1}^{i-1} (n-2k) = i(n-i+1).$$

We have $d = \lfloor \frac{(n+1)^2}{4} \rfloor$; this value is obtained for $i = \lfloor \frac{n+1}{2} \rfloor$ and $h = \lfloor \frac{n+1}{2} \rfloor$ if n is odd and $h = \frac{n}{2}, \frac{n}{2} + 1$ if n is even.

Type C_n

Again X^+ is linearly ordered and consists of n elements $\tau_1 = \theta^\vee < \dots < \tau_n = 2\omega_n^\vee$; moreover $\Pi_\ell = \{\alpha_n\}$ and, for $1 \leq i \leq n-1$, $\widehat{\Delta}_i(\alpha_n) \cong C_{n-i}$, so that

$$\dim (\mathfrak{m}_i(\alpha_n)) = n + \sum_{k=1}^{i-1} (n-k) = \frac{(2n-i+1)i}{2}.$$

We have $d = \frac{n(n+1)}{2} = \dim (\mathfrak{m}_n(\alpha_n))$.

Type B_n

In this case there are exactly two maximal good chains. The first is $\{\Delta_1 = \Delta \supset \dots \supset \Delta_{\lfloor \frac{n}{2} \rfloor}\}$ with $\Delta_i = \Delta(\alpha_{2i-1}, \dots, \alpha_n)$ and the other is $\{\Delta_1 = \Delta \supset \tilde{\Delta}_2\}$ where $\tilde{\Delta}_2$ is the rank one system generated by α_1 ; this the good chain of α_1 . The good chain of α_2 consists only of Δ . We have

$$\begin{aligned} \dim (\mathfrak{m}_1(\alpha_h)) &= 2n-2, \quad 1 \leq h \leq n-1, \\ \dim (\mathfrak{m}_2(\alpha_1)) &= 2n-1. \end{aligned}$$

For $h \geq 3$

$$\widehat{\Delta}_k(\alpha_h) \cong \begin{cases} A_1 & h = 2k+1, \\ A_3 & h = 2k+2, \\ D_{h-2k+1} & 2k+3 \leq h \leq n-1. \end{cases}$$

Hence, for $3 \leq h \leq n-1, i \geq 2$

$$\dim (\mathfrak{m}_i(\alpha_h)) = 2n-2 + \begin{cases} \sum_{k=1}^{i-2} (2h-4k-1) + 1 & h = 2i-1, \\ \sum_{k=1}^{i-2} (2h-4k-1) + 3 & h = 2i, \\ \sum_{k=1}^{i-1} (2h-4k-1) & h > 2i-1. \end{cases}$$

Note that the second summand in the r.h.s. (i.e., $\ell(\check{w}_{i,\alpha_h})$) is independent from n . The previous calculation can be expressed as follows

$$\dim (\mathfrak{m}_i(\alpha_h)) = \begin{cases} \frac{4n+h^2-3h-2}{2} & \text{if the ideal is maximal,} \\ 2(n-i^2) + (i-1)(2h+1) & \text{otherwise.} \end{cases}$$

For $n=3$ we have $d=5$, obtained for $h=1, i=2$. For $n \geq 4$ we have $d = \frac{n(n-1)}{2} + 1$ and this value is obtained for $h=n-1$ and $i = \lfloor \frac{n}{2} \rfloor$ (for $n=4$ it is also obtained for $h=1, i=2$).

Type D_n

The structure of X^+ is exactly the same as for B_n . The only difference is that any good chain starts with a system of type D_n , hence the only difference in the

calculations lies in the Coxeter number of the whole system. The final results are as follows.

$$\begin{aligned} \dim (\mathfrak{m}_1(\alpha_h)) &= 2n - 3, \quad 1 \leq h \leq n - 1, \\ \dim (\mathfrak{m}_2(\alpha_1)) &= 2n - 2, \end{aligned}$$

and for $3 \leq h \leq n - 1, i \geq 2$

$$\dim (\mathfrak{m}_i(\alpha_h)) = \begin{cases} \frac{4n+h^2-3h-4}{2} & \text{if the ideal is maximal,} \\ 2(n-i^2) + (i-1)(2h+1) - 1 & \text{otherwise.} \end{cases}$$

Type E_6

X^+ consists of two elements $\tau_1 = \theta^\vee < \tau_2 = \theta^\vee + \theta^\vee(A_5)$ and $\Pi(\tau_2) = \Pi \setminus \{\alpha_2\}$. We have

$$\widehat{\Delta}_1(\alpha_i) \cong \begin{cases} A_5 & i = 1, 6, \\ A_2 & i = 3, 5, \\ A_1 & i = 4. \end{cases}$$

therefore

$$\begin{aligned} \dim (\mathfrak{m}_1(\alpha_h)) &= 11, \quad 1 \leq h \leq 6 \\ \dim (\mathfrak{m}_2(\alpha_1)) &= \dim (\mathfrak{m}_2(\alpha_6)) = 16, \\ \dim (\mathfrak{m}_2(\alpha_3)) &= \dim (\mathfrak{m}_2(\alpha_5)) = 13, \\ \dim (\mathfrak{m}_2(\alpha_4)) &= 12. \end{aligned}$$

Type E_7

X^+ consists of three elements $\tau_1 = \theta^\vee < \tau_2 = \theta^\vee + \theta^\vee(D_6) < \tau_3 = \theta^\vee + \theta^\vee(D_6) + \theta^\vee(A_1)$ and $\Pi(\tau_2) = \Pi \setminus \{\alpha_1\}$, $\Pi(\tau_3) = \{\alpha_7\}$. We have

$$\widehat{\Delta}_1(\alpha_i) \cong \begin{cases} A_1 & i = 2, \\ A_3 & i = 3, 5, \\ A_2 & i = 4, \\ A_5 & i = 6, \\ D_6 & i = 7, \end{cases} \quad \widehat{\Delta}_2(\alpha_7) \cong A_1.$$

therefore

$$\begin{aligned} \dim (\mathfrak{m}_1(\alpha_h)) &= 17, \quad 1 \leq h \leq 7 \\ \dim (\mathfrak{m}_2(\alpha_2)) &= 18, \\ \dim (\mathfrak{m}_2(\alpha_3)) &= \dim (\mathfrak{m}_2(\alpha_5)) = 20, \\ \dim (\mathfrak{m}_2(\alpha_4)) &= 19, \\ \dim (\mathfrak{m}_2(\alpha_6)) &= 22, \\ \dim (\mathfrak{m}_2(\alpha_7)) &= 26, \\ \dim (\mathfrak{m}_3(\alpha_7)) &= 27. \end{aligned}$$

Type E_8

X^+ consists of two elements $\tau_1 = \theta^\vee < \tau_2 = \theta^\vee + \theta^\vee(E_7)$ and $\Pi(\tau_2) = \Pi \setminus \{\alpha_8\}$. We have

$$\widehat{\Delta}_1(\alpha_i) \cong \begin{cases} A_7 & i = 1, \\ A_5 & i = 2, 3, \\ A_4 & i = 4, \\ A_3 & i = 5, \\ A_2 & i = 6, \\ A_1 & i = 7, \end{cases}$$

therefore

$$\begin{aligned} \dim(\mathfrak{m}_1(\alpha_h)) &= 29, & 1 \leq h \leq 8 \\ \dim(\mathfrak{m}_2(\alpha_1)) &= 36, \\ \dim(\mathfrak{m}_2(\alpha_2)) &= \dim(\mathfrak{m}_2(\alpha_3)) = 34, \\ \dim(\mathfrak{m}_2(\alpha_4)) &= 33, \\ \dim(\mathfrak{m}_2(\alpha_5)) &= 32, \\ \dim(\mathfrak{m}_2(\alpha_6)) &= 31, \\ \dim(\mathfrak{m}_2(\alpha_7)) &= 30. \end{aligned}$$

Type F_4

X^+ consists of two elements $\tau_1 = \theta^\vee < \tau_2 = \theta^\vee + \theta^\vee(C_3)$ and $\Pi(\tau_2) = \{\alpha_2, \alpha_3, \alpha_4\}$. Since $\widehat{\Delta}(\alpha_2) \cong A_1$ we have

$$\begin{aligned} \dim(\mathfrak{m}_1(\alpha_h)) &= 8, & h = 1, 2 \\ \dim(\mathfrak{m}_2(\alpha_2)) &= 9. \end{aligned}$$

Type G_2

We have $X^+ = \{\theta^\vee\}$. There is only one maximal ideal corresponding to the unique simple long root; it has dimension 3.

Remark. It is also possible to obtain the previous results using Proposition 7.7. Indeed

$$\dim(\max \mathcal{I}_{ab}(\alpha, \tau)) = \ell(v_\tau) + \ell(y_{\tau, \alpha}) + \ell(w_0^\alpha) - \ell(w_0^{\alpha, \tau})$$

where w_0^α is the longest element in W_α and $w_0^{\alpha, \tau}$ is the longest element in $W((\pi \cap \alpha^\perp) \setminus P_{\tau, \alpha})$.

Now $\ell(v_\tau) = |\Delta_\tau^2| = |\Delta_\tau^{\text{even}}| - |\Delta_\tau^0|$, $\ell(y_{\tau, \alpha}) = g(\Delta(\tau)) - 2$ and the last two summands in the r.h.s. of the previous formula can be easily calculated determining the number of positive roots of the corresponding subsystems.

As an example, we will write down the computations for types A_n and E_6 for the maximal ideals, displaying the results as follows. Set $\tau_\alpha = \tau_{k(\alpha)}(\alpha)$.

long simple root α	τ_α	$\Delta_{\tau_\alpha}^{\text{even}}$	$\Delta(\tau_\alpha)$	$\ell(w_0^\alpha)$
$\dim(\mathfrak{i}_\alpha)$		$\Delta_{\tau_\alpha}^0$		$\ell(w_0^{\alpha, \tau_\alpha})$

We use the following notational conventions: if in the rightmost column appears a root system of negative rank, then the whole column must not be considered (e.g., type A_n , $k = 1$). Moreover a root system of rank zero should be regarded as the empty set.

Type A_n

For $1 \leq k \leq n$ we have

$$\begin{array}{ccccc} \alpha_k & \omega_k^\vee + \omega_{n-k+1}^\vee & A_{2k-1} \times A_{n-2k} & A_{n-2k+2} & A_{n-k-1} \\ k(n-k+1) & & A_{k-1} \times A_{n-2k} \times A_{k-1} & & A_{n-2k} \times A_{k-2} \end{array}$$

Type E_6

$$\begin{array}{ccccc} \alpha_1, \alpha_6 & \omega_1^\vee + \omega_6^\vee & D_5 & A_5 & A_4 \\ 16 & & D_4 & & A_3 \\ \alpha_3, \alpha_5 & \omega_1^\vee + \omega_6^\vee & D_5 & A_5 & A_1 \times A_2 \\ 13 & & D_4 & & A_2 \\ \alpha_4 & \omega_1^\vee + \omega_6^\vee & D_5 & A_5 & \\ 12 & & D_4 & & \\ \alpha_2 & \omega_2^\vee & A_1 \times A_5 & E_6 & \\ 11 & & A_5 & & \end{array}$$

ACKNOWLEDGMENT

The authors wish to thank Corrado De Concini for several useful discussions.

REFERENCES

- [B] N. Bourbaki, *Groupes et algèbres de Lie*, Hermann, Paris, 1968.
- [CMP] P. Cellini, P. Möseneder Frajria and P. Papi, *Compatible discrete series*, Pacific J. of Math. (to appear), [math.RT/0207275](#).
- [CP1] P. Cellini and P. Papi, *The structure of total reflection orders in affine root systems*, J. Algebra **205** (1998), 207–226.
- [CP2] P. Cellini and P. Papi, *ad-nilpotent ideals of a Borel subalgebra*, J. Algebra **225** (2000), 130–141.
- [CP3] P. Cellini and P. Papi, *ad-nilpotent ideals of a Borel subalgebra II*, J. Algebra **258** (2002), 112–121.
- [D] M. Dyer, *Hecke algebras and shellings of Bruhat intervals*, Comp. Math. **89** (1993), 91–115.
- [H] J.E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge University Press, 1990.
- [IM] N. Iwahori, H. Matsumoto, *On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups*, Inst. Hautes Études Sci. Publ. Math. **25** (1965), 5–48.

- [Kac] V.G. Kac, *Infinite Dimensional Lie Algebras*, Cambridge University Press, 1990.
- [K1] B. Kostant, *Eigenvalues of a Laplacian and commutative Lie subalgebras*, *Topology* **3**, suppl. **2** (1965), 147–159.
- [K2] B. Kostant, *The Set of Abelian ideals of a Borel Subalgebra, Cartan Decompositions, and Discrete Series Representations*, *Internat. Math. Res. Notices* **5** (1998), 225–252.
- [M] A. Mal'cev, *Commutative subalgebras of semi-simple Lie algebras*, *Translations of the Amer. Math. Soc. Series 1* **9** (1951), 214–227.
- [P] D. Panyushev, *Abelian ideals of a Borel subalgebra and long positive roots*, *IMRN* **35** (2003), 1889–1913.
- [PR] D. Panyushev, G. Röhrle, *Spherical orbits and abelian ideals*, *Adv. Math.* **159** (2001), 229–246.
- [R] G. Röhrle, *On normal abelian subgroups in parabolic groups*, *Ann. Inst. Fourier, Grenoble* **48** (1998), 1455–1482.
- [St1] J. Stembridge, *The partial order of dominant weights*, *Adv. Math.* **136** (1998), 340–364.
- [St2] J. Stembridge, *Quasi-minuscule quotients and reduced words for reflections*, *J. Alg. Comb.* **13** (2001), 275–293.
- [S] R. Suter, *Abelian ideals in a Borel subalgebra of a semisimple Lie algebra* (2002), [math.RT/0210463](#).

PAOLA CELLINI
 DIPARTIMENTO DI SCIENZE
 UNIVERSITÀ “G. D’ANNUNZIO”
 VIALE PINDARO 87
 65127 PESCARA — ITALY
 E-MAIL: cellini@sci.unich.it

PAOLO PAPI
 DIPARTIMENTO DI MATEMATICA ISTITUTO G. CASTELNUOVO
 UNIVERSITÀ DI ROMA “LA SAPIENZA”
 PIAZZALE ALDO MORO 5
 00185 ROMA — ITALY
 E-MAIL: papi@mat.uniroma1.it