

# The Generalized Witt Algebras using Additive Maps

Ki-Bong Nam and Mook Ok Wang

September 25, 2017

## Abstract

Kawamoto generalized the Witt algebra using  $F[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$  instead of  $F[x_1, \dots, x_n]$ . We construct the generalized Witt algebra  $W(g_p, n)$  by using an additive map  $g_p$  from a set of integers into a field of characteristic zero where  $1 \leq p \leq n$ . We show that the Lie algebra  $W(g_p, n)$  is simple if  $g_p$  is injective, and also the Lie algebra  $W(g_p, n)$  has no ad-diagonalizable elements.

## 1 Introduction

Let  $F$  be a field of characterisitic zero. The Witt algebra is called the general algebra by Rudakov [8]. Kac [2] studied the generalized Witt algebra on the  $F$ -algebra in the formal power series  $F[[x_1, \dots, x_n]]$  for a positive integer  $n$ . Nam [5] constructs the Lie algebra on the  $F$ -subalgebra  $F[e^{\pm x_1}, \dots, e^{\pm x_n}, x_1, \dots, x_{n+m}]$  in the formal power series  $F[[x_1, \dots, x_{n+m}]]$  for the positive integers  $n$  and  $m$ .

The Witt algebra  $W(n)$  has a basis

$$\{x_1^{a_1} \cdots x_n^{a_n} \partial_i | a_1, \dots, a_n \in N, 1 \leq i \leq n\}$$

with Lie bracket on basis elements

$$\begin{aligned} & [x_1^{a_1} \cdots x_n^{a_n} \partial_i, x_1^{b_1} \cdots x_n^{b_n} \partial_j] \\ &= b_i x_1^{a_1+b_1} \cdots x_n^{a_n+b_n} x_i^{-1} \partial_j - a_j x_1^{a_1+b_1} \cdots x_n^{a_n+b_n} x_j^{-1} \partial_i \end{aligned}$$

where  $N$  is the set of non-negative integers.

Consider the generalized Witt algebra  $W(n, m)$  having a basis

$$\{e^{a_1 x_1} \cdots e^{a_n x_n} x_1^{u_1} \cdots x_{n+m}^{u_{n+m}} \partial_k | a_1, \dots, a_n, u_1, \dots, u_{n+m} \in Z, 1 \leq k \leq n+m\}$$

with Lie bracket on basis elements given by

$$\begin{aligned} & [e^{a_1 x_1} \cdots e^{a_n x_n} x_1^{u_1} \cdots x_{n+m}^{u_{n+m}} \partial_i, e^{b_1 x_1} \cdots e^{b_n x_n} x_1^{t_1} \cdots x_{n+m}^{t_{n+m}} \partial_j] \\ &= b_i e^{a_1 x_1 + b_1 x_1} \cdots e^{a_n x_n + b_n x_n} x_1^{u_1+t_1} \cdots x_{n+m}^{u_{n+m}+t_{n+m}} \partial_j \end{aligned}$$

$$\begin{aligned}
& +t_i e^{a_1 x_1 + b_1 x_1} \dots e^{a_n x_n + b_n x_n} x_1^{u_1 + t_1} \dots x_{n+m}^{u_{n+m} + t_{n+m}} x_i^{-1} \partial_j \\
& -a_i e^{a_1 x_1 + b_1 x_1} \dots e^{a_n x_n + b_n x_n} x_1^{u_1 + t_1} \dots x_{n+m}^{u_{n+m} + t_{n+m}} \partial_i \\
& -u_i e^{a_1 x_1 + b_1 x_1} \dots e^{a_n x_n + b_n x_n} x_1^{u_1 + t_1} \dots x_{n+m}^{u_{n+m} + t_{n+m}} x_j^{-1} \partial_i
\end{aligned}$$

where  $b_i = 0$  if  $n + 1 \leq i \leq n + m$ , and  $a_j = 0$  if  $n + 1 \leq j \leq n + m$  (see [4, 5, 7]). In [5, 8], it is noted that the Lie subalgebra of  $W(n, m)$  is the Witt algebra  $W(n)$  on  $F[x_1, \dots, x_m]$ . Let  $g_p$  be an additive map from  $Z$  into  $F$  where  $1 \leq p \leq n$ . Let us define as  $W(g_p, n)$  the Lie algebra with basis

$$B := \left\{ \binom{a_1}{i_1} \dots \binom{a_n}{i_n}_k \mid a_1, \dots, a_n, i_1, \dots, i_n \in Z, 1 \leq k \leq n \right\} \quad (1)$$

and a Lie bracket on basis elements given by

$$\begin{aligned}
& \left[ \binom{a_1}{i_1} \dots \binom{a_n}{i_n}_k, \binom{b_1}{j_1} \dots \binom{b_n}{j_n}_l \right] \\
& = g_k(b_k) \binom{a_1 + b_1}{i_1 + j_1} \dots \binom{a_n + b_n}{i_n + j_n}_l \\
& + j_k \binom{a_1 + b_1}{i_1 + j_1} \dots \binom{a_k + b_k}{i_k + j_k - 1} \binom{a_{k+1} + b_{k+1}}{i_{k+1} + j_{k+1}} \dots \binom{a_n + b_n}{i_n + j_n}_l \\
& - g_l(a_l) \binom{a_1 + b_1}{i_1 + j_1} \dots \binom{a_n + b_n}{i_n + j_n}_l \\
& - i_l \binom{a_1 + b_1}{i_1 + j_1} \dots \binom{a_l + b_l}{i_l + j_l - 1} \binom{a_{l+1} + b_{l+1}}{i_{l+1} + j_{l+1}} \dots \binom{a_n + b_n}{i_n + j_n}_k.
\end{aligned} \quad (2)$$

It follows from [1, 4, 5, 6] that the above bracket is extended linearly to the given basis  $B$ . Also, it is not hard to show that the above bracket satisfies the Jacobi identity. In section 2, we will prove the following main theorems. Throughout the paper, a given map  $g_p$  means an additive and injective map from a set of integers into a field of characteristic zero.

**Theorem 1** The Lie algebra  $W(g_p, n)$  is simple.

**Theorem 2** For any automorphism of  $\theta \in \text{Aut}(W(g, 1))$

$$\theta \left( \binom{0}{1}_1 \right) = \sum_j C_j \left( \binom{0}{j}_1 \right)$$

where  $C_j \in F$ .

**Theorem 3** Each derivation of  $W(g, 1)_+$  can be written as a sum of an inner derivation and a scalar derivation [6].

## 2 Simplicities of $W(g_p, n)$

The Lie algebra  $W(g_p, n)$  has a  $Z^n$ -gradation as mentioned in [3] : that is,

$$W(g_p, n) = \bigoplus_{(a_1, \dots, a_n) \in Z^n} W_{(a_1, \dots, a_n)} \quad (3)$$

where each is the subspace of  $W(g_p, n)$  with a basis

$$\bar{B} = \left\{ \binom{a_1}{i_1} \cdots \binom{a_n}{i_n} \mid a_1, \dots, a_n, i_1, \dots, i_n \in Z, 1 \leq l \leq n \right\}$$

Let  $W_{(a_1, \dots, a_n)}$  denote the  $(a_1, \dots, a_n)$ -homogeneous component of  $W(g_p, n)$  and elements in the  $W_{(a_1, \dots, a_n)}$ -homogeneous  $(a_1, \dots, a_n)$ -homogeneous elements. Note that the  $(0, \dots, 0)$  - homogeneous component is isomorphic to the Witt algebra  $W(n)$  [8]. From now on let  $(0, \dots, 0)$ -homogeneous component denote the 0-homogeneous component. For the simplicity of  $W(g_p, n)$ , we define the map as an additive and injective map. Now we introduce a lexicographic ordering of two basis elements of  $W(g_p, n)$  as follows : for any two elements  $\binom{a_1}{i_1} \cdots \binom{a_n}{i_n}_l, \binom{b_1}{j_1} \cdots \binom{b_n}{j_n}_k$ , we have

$$\binom{a_1}{i_1} \cdots \binom{a_n}{i_n}_l > \binom{b_1}{j_1} \cdots \binom{b_n}{j_n}_k$$

if  $(a_1, \dots, a_n, i_1, \dots, i_n, l) > (b_1, \dots, b_n, j_1, \dots, j_n, k)$  by the natural lexicographic ordering in  $Z^{2n+1}$ .

For any element  $l \in W(g_p, n)$ ,  $l$  can be written as follows using the ordering and gradation :

$$\begin{aligned} l = & \sum_{i_1, \dots, i_n, p} C(i_1, \dots, i_n, p) \binom{a_{11}}{i_1} \cdots \binom{a_{1n}}{i_n}_l + \cdots \\ & + \sum_{j_1, \dots, j_n, q} C(j_1, \dots, j_n, q) \binom{a_{t1}}{j_1} \cdots \binom{a_{tn}}{j_n}_q \end{aligned}$$

where  $1 \leq p, \dots, q \leq n$ ,  $(a_{11}, \dots, a_{1n}) > \dots > (a_{t1}, \dots, a_{tn})$  and

$$C(i_1, \dots, i_n, p), \dots, C(j_1, \dots, j_n, q) \in F.$$

Next, define the string number  $st(l) = t$  for  $l$  (see [5, 6]), and  $l_p(l)$  as  $\max\{i_1, \dots, i_n, \dots, j_1, \dots, j_n\}$ . For any basis element  $\binom{a_1}{i_1} \cdots \binom{a_n}{i_n}_l$  in  $\bar{B}$ , let us refer to  $a_1, \dots, a_n$  as upper indices and  $i_1, \dots, i_n$  as lower indices.

**Remark 2.1** If  $g_p$  is an inclusion, then  $W(g_p, n) = W(n, 0)$  is the generalized Witt algebra which is studied by Nam [5, 6].

**Lemma 1** *If  $l$  is any non-zero element, then the ideal  $\langle l \rangle$  generated by  $l$  contains an element whose lower indices are positive.*

*Proof.* Let  $l$  be a nonzero element of  $W(g_p, n)$ . Take an element  $M = \binom{0}{j_1} \cdots \binom{0}{j_n}_t$  such that  $j_1 \gg \cdots \gg j_n$  and  $t$  such that either  $a_t \neq 0$  or  $i_t \neq 0$  in  $l$ , where  $a \gg b$  means  $a$  is sufficiently larger than  $b$ . Then  $0 \neq [M, l]$  is the required element.  $\square$

**Lemma 2** *If an ideal  $I$  of  $W(g_p, n)$  contains  $\binom{0}{0} \cdots \binom{0}{0}_l$  for  $1 \leq l \leq n$ , then  $I = W(g_p, n)$ .*

*Proof.* Since  $W(n)$  is a simple subalgebra of  $W(g_p, n)$  with the basis  $\{\binom{0}{i_1} \cdots \binom{0}{i_n}_l | i_1, \dots, i_n \in Z, 1 \leq l \leq n\}$ , the ideal  $\langle \binom{0}{0} \cdots \binom{0}{0}_i \rangle$  contains  $W(n)$ , where  $\langle \binom{0}{0} \cdots \binom{0}{0} \rangle$  is the ideal of which is generated by  $\binom{0}{0} \cdots \binom{0}{0}_l$  for a fixed  $t \in \{1, \dots, n\}$  (see [8]). It follows from Lemma 2 of [5] and Lemma 3 of [6] that the basis elements of  $W(g_p, n)$  are contained in  $I$  by using the injectiveness of  $g_p$ .  $\square$

**Theorem 1** *The Lie algebra  $W(g_p, n)$  is simple.*

*Proof.* It is not difficult to prove this theorem by induction on  $st(l)$  for any element  $l$  in any ideal  $I$  of  $W(g_p, n)$  and by considering the one to one properties of  $g_p$ .

**Corollary 1** *The Lie algebra  $W(n, 0)$  is simple.*

*Proof.* If we take an additive embedding  $g_p : Z \rightarrow F$ ,  $1 \leq p \leq n$ , then we get the required result (see [5, 6]).  $\square$

It is an interesting problem to find all the automorphisms of the subalgebra  $W(g, 1)$  of  $W(g_p, n)$ .

**Theorem 2** *For any automorphism  $\theta \in \text{Aut}(g_p, n)$*

$$\theta\left(\binom{0}{1}_1\right) = \sum_j C_j \binom{0}{j}_1$$

where  $C_j \in F$ .

*Proof.* It is not difficult to prove this theorem using the gradation of (2.1) and the  $W(g, 1)$  acting of  $\binom{0}{0}_1$  on the zero homogeneous component  $W_0$  whose vector basis is  $\{\binom{0}{i}_1 | i \in Z\}$  as an ad-map.  $\square$

If we consider the Lie subalgebra of  $W(g, 1)$  such that all the lower indices are zero, then this subalgebra is a Block algebra  $W(1)$  which is also called the centerless Virasoro algebra [1]. Thus all the automorphisms of this Lie algebra can be determined by Theorem 3 in [1].

Consider the Lie subalgebra  $W(1, 1)^+$  of  $W(n, m)$  in [6] with basis

$$\{e^{ax} x^i y^j \partial_x, e^{bx} x^l y^m \partial_y | a, b \in Z, i, j, l, m \in N\}$$

where  $N$  is the set of non-negative integers.

**Conjecture** For any  $\theta \in \text{Aut}(W(1, 1)^+)$   $\theta(y\partial_y) = (\alpha y + \beta)\partial_y$  for some  $0 \neq \alpha, \beta \in F$ .

The element  $l \in W(g_p, n)$  is an ad-diagonalizable element if  $[l, m] = \alpha(m)m$  for any  $m \in B$  given in (1) and for some  $\alpha(m) \in F$ .

**Proposition 1** *The Lie algebra  $W(g, n)$  has no ad-diagonalizable element with respect to the basis given in (1).*

*Proof.* Since  $W(g, n)$  is  $Z^n$ -graded Lie algebra all the ad-diagonalizable elements are in the  $(0, \dots, 0)$ -homogeneous component.  $W_{(0, \dots, 0)}$  is isomorphic to  $W(n)$  as Lie algebras where  $W(n)$  is the well known Witt algebra [8]. Thus all the ad-semisimple elements of  $W(g_p, n)$  are of the form  $\sum_{i=1}^n C_i \binom{0}{1}_i$ , where  $C_i \in F$ . But  $[\sum_{i=1}^n C_i \binom{0}{1}_i, \binom{a}{0}_j] \neq \alpha \binom{a}{0}_j$  for any  $\alpha \in F$ . Therefore, we proved the proposition.  $\square$

**Remark 2.2** For another proof of Proposition 2.6, see Corollary 1 of [5].

### 3 Derivation of $W(g, 1)_+$

Consider the subalgebra  $W(g, 1)_+$  of  $W(g_p, n)$  with basis

$$B_+ := \left\{ \binom{a}{i}_1 \mid a \in Z, i \in N \right\}$$

where  $N$  is the set of non-negative integers.

In this section we determine all the derivations of the Lie algebra  $W(g, 1)_+$ . Ikeda and Kawamoto found all the derivations of the Kawamoto algebra  $W(G, I)$  [4] in their paper [2]. It is very important to find all the derivations of a given Lie algebra to compare with other Lie algebras. Let  $L$  be a Lie algebra over any field  $F$ . An  $F$ -linear map  $D$  from  $L$  to  $L$  is a derivation if  $D([l_1, l_2]) = [D(l_1), l_2] + [l_1, D(l_2)]$  for any  $l_1, l_2 \in L$ .

Let  $L$  be a Lie algebra over any field  $F$ . Define the derivation  $D$  of  $L$  to be a scalar derivation if for all basis elements  $l$  of  $L$ ,  $D(l) = f_l l$  for some scalar  $f_l \in F$  [1], [7].

We need the following lemma.

**Lemma 3** *Let  $D$  be a derivation of  $W(g, 1)_+$ . If  $D(\partial) = 0$ , then  $D = f \text{ad}_\partial + S$  where  $f \in F$  and  $S$  is a scalar derivation.*

*Proof.* It is not difficult to prove this lemma using the gradation and the ordering of the Lie algebra  $W(g, 1)_+$ . [6].  $\square$

**Theorem 3** *Each derivation of  $W(g, 1)_+$  can be written as a sum of an inner derivation and a scalar derivation.*

*Proof.* Let  $D$  be any derivation of  $W(g, 1)_+$ . Then  $D(\partial) = f\partial$  for some  $f \in F[e^{\pm x}, x]$ . Since  $\partial : F[e^{\pm x}, x] \rightarrow F[e^{\pm x}, x]$  is onto, there is a function  $g \in F[e^{\pm x}, x]$  such that  $\partial(g) = f$ . Then  $ad_{g\partial}(\partial) = [\partial, g\partial] = f\partial = D(\partial)$ . We have  $(D - ad_{g\partial})(\partial) = 0$ . By Lemma 3 we have  $D = ad_{g\partial} + cad_{\partial} + S$  [6]. Therefore, we have proved the theorem.  $\square$

**Remark 3.1** Let  $Z^+$  and  $F^+$  be the additive group of  $Z$  and  $F$  respectively. For any  $g \in Hom(Z^+, F^+)$ , we have  $g(1) = m$  for some fixed  $m \in F$ , thus the Witt algebra  $W(n)$  can not be changed by using the idea of the additive map in this paper [5].

## References

- [1] R. Block, *On torsion-free abelian groups and Lie algebras*, Proc. Amer. Math. Soc. 9(1958), 613-620.
- [2] T. Ikeda and N. Kawamoto, *On the derivations of generalized Witt algebras*, Hiroshima Math. J., 20 (1990), 47-55.
- [3] V. G. Kac, *Description of filtered Lie algebra with which graded Lie algebras of Cartan type are associated*, Izv. Akad. Nauk SSSR, Ser. Mat. Tom. 38(1974), 832-834.
- [4] N. Kawamoto, *On G-Graded Automorphisms of generalized Witt algebras*, Contem. Math. A.M.S., 184(1995), 225-230.
- [5] N. Kawamoto, *Generalized Witt algebras over a field of characteristic zero*, Hiroshima. Math. J., 16(1986), 417-426.
- [6] Ki-Bong Nam, *Generalized Witt algebras over a field of characteristic zero*, UW-Madison, thesis, (1998), 1-40.
- [7] Ki-Bong Nam, *Simple Lie algebras which generalize Witt algebras I*, Kyungpook Math. J., to appear, 1-9.
- [8] D. S. Passman, *Simple Lie Algebras of Witt-Type*, to appear, 1-9.
- [9] A. N. Rudakov, *Groups of Automorphisms of Infinite-Dimensional Simple Lie Algebras*, Math. USSR-Izv. 3(1969), 707-722.

Department of Mathematics, UW-Madison, WI 53706

e-mail :nam@math.wisc.edu

Department of Mathematics, Hanyang University-Ansan, Ansan, Korea