

Determining a Semisimple Group from its Representation Degrees

BY

MICHAEL LARSEN*

*Department of Mathematics, Indiana University
Bloomington, IN 47405, USA*

ABSTRACT

The Lie algebra of a compact semisimple Lie group G is determined by the degrees of the irreducible representations of G . However, two different groups can have the same representation degrees.

0. Introduction

A compact semisimple Lie group G has a finite number of representations of any given degree. In this paper we investigate to what extent G is determined by the multi-set of degrees of its representations or equivalently, the multi-set of degrees of its irreducible representations. This problem arises naturally in at least two different ways. On the one hand, following Witten [7], we define the zeta-function $\zeta_G(s)$ of a compact semisimple Lie group as the Dirichlet series

$$\zeta_G(s) = \sum_{V \text{ irreducible}} (\dim V)^{-s}.$$

The question in this paper then belongs to the familiar class of problems asking when two different mathematical objects can have the same zeta-function. On the other hand, the multi-set of degrees of the irreducible representations of G comprises a (small) part of the data encoded in the representation category $\text{Rep}_{\mathbb{C}}G$. We know [2] that the full structure of neutral Tannakian category on $\text{Rep}_{\mathbb{C}}G$ enables us to recover the complexification $G_{\mathbb{C}}$ of G and therefore G itself (as a maximal compact subgroup of $G_{\mathbb{C}}$). For semisimple groups, though, there are results (see for instance, [5]) showing that $\text{Rep}_{\mathbb{C}}G$ has more data than we need to determine G . In this paper we are asking whether the \mathbb{C} -linear abelian category with fiber functor underlying $\text{Rep}_{\mathbb{C}}G$ already determines G .

* Partially supported by NSF Grant DMS-0100537

The main theorem of this paper is that $\zeta_G(s)$ determines the Lie algebra of G up to isomorphism. Our basic tool is the Weyl dimension formula, which expresses the dimension of each irreducible representation of a group G as a product over the set of positive roots of G . We try to recover the geometry of the root system of G from the factorizations of the representation degrees, the basic difficulty being that there are many ways to factor a given degree. Our strategy is to choose values of n which have few factorizations or at least few factorizations which could possibly arise from Weyl's formula.

The simplest idea is to consider prime powers $n = p^k$. It is an easy consequence of Weyl's formula that $\dim V_{p\lambda + (p-1)\rho} = p^{|R^+|} \dim V_\lambda$, where V_λ is the irreducible representation of G of highest weight λ , R^+ is the set of positive roots of G , and ρ is the half-sum of the elements of R^+ . An elaboration of this argument, given in §1, shows that the prime power coefficients determine the number of irreducible factors of R and the number of roots in each factor. Unfortunately, there are infinitely many pairs of distinct irreducible root systems with the same number of roots, and prime power coefficients alone are in general insufficient to disambiguate further. For example, $\mathrm{SO}(15)$ and $\mathrm{PSp}(14)$ both have the property that the number of their irreducible representations of degree p^k is 1 if $k \in 49\mathbb{Z}$ and 0 otherwise (for $p \gg 0$, this follows from the discussion in §1, but a computer search is needed to confirm it for small primes.)

We therefore broaden our search to encompass degrees n in which the largest power of some prime p dividing n is not too much smaller than n . To illustrate this idea, we consider the case that R is irreducible, and exclude weights λ with $\lambda + \rho$ divisible by p (such weights are not “allowable at p ” in the terminology of this paper). We compare the size of the logarithm of the largest power of p dividing $\dim V_\lambda$ to that of $\dim V_\lambda$ itself and prove that

$$(0.0.1) \quad \lim_{p \rightarrow \infty} \sup_{\lambda \neq 0 \text{ allowable}} \frac{-\log |\dim V_\lambda|_p}{\log \dim V_\lambda} = \lim_{p \rightarrow \infty} \sup_{\lambda \neq 0 \text{ allowable}} \frac{\log p \operatorname{ord}_p \dim V_\lambda}{\log \dim V_\lambda}$$

approaches a rational limit strictly between 0 and 1 which depends on R . This limit is the **efficiency** of R (defined in §3 directly in terms of the geometry of R). A more efficient root system always has allowable representation degrees which cannot be achieved by an allowable weight of a less efficient system. Once the efficiency is known, we can begin studying the families of representations (indexed by p) whose degrees achieve it. It turns out the optimal ones, in a sense made explicit below, lie in finitely many one parameter families of the form

$p \mapsto p\mu + \nu$. For each such family, the degree is given as a polynomial in p , the “Weyl polynomial” associated to μ and ν . The Weyl polynomials of a simple root system encode enough information, ultimately, to extract the root system giving rise to them.

The process of excluding non-allowable representations is carried out in §1 and is somewhat delicate; because our groups may not be products of almost simple factors, we must restrict attention to primes which are congruent to 1 modulo a sufficiently divisible integer. Weyl polynomials are introduced in §2 and the basic pairs (μ, ν) used in the proof of the main theorem are given. The computation of efficiencies occupies much of §3 and is rather involved. There does not seem to be a simple formula for the answer: for instance, the efficiency of E_8 turns out to be $\frac{63}{117}$. The proof of the main theorem, by means of a rather complicated induction on the number of factors, is given in §4.

To show that our result is in some sense sharp, in §5 we give a construction of pairs of non-isomorphic compact semisimple Lie groups with the same Witten zeta-function. This construction is modelled on a well known theorem of F. Gassmann [3] asserting that the Dedekind zeta-function $\zeta_K(s)$ of a number field K does not determine K up to isomorphism. The proof of Gassmann’s theorem is purely group-theoretic. Namely, there exists a finite group G with subgroups H_1 and H_2 , not conjugate in G , whose elements can be put into one-to-one correspondence in such a way that corresponding elements *are* conjugate in G . The fixed fields of H_1 and H_2 in a Galois extension with group G then have the same zeta-function. T. Sunada [6] used the same trick to construct pairs of non-isometric isospectral manifolds; here π_1 plays the role of the Galois group. Like Sunada, we exploit π_1 , but our fundamental groups are abelian, so there are no inner automorphisms to work with. Instead we make use of outer automorphisms. We construct subgroups H_1 and H_2 of a suitable $\pi_1(G)$, whose elements can be put into one-to-one correspondence in such a way that corresponding elements always lie in the same orbit of $\text{Aut}(\pi_1(G))$ but such that the groups H_1 and H_2 as a whole do not. The corresponding covering groups can then be shown to have the same representation degrees.

This work was initiated during a visit to the Hebrew University in Jerusalem in the summer of 2002. I would like to thank the Einstein Institute of Mathematics for its hospitality during this visit.

1. Modifying the zeta-function

Let G be a compact semisimple group, \tilde{G} its universal cover, and G_1, \dots, G_m the almost simple direct factors of G . Let Λ_G and $\Lambda_{\tilde{G}}$ denote the weight lattices of G and \tilde{G} respectively, so

$$[\Lambda_{\tilde{G}} : \Lambda_G] = |\ker(\tilde{G} \rightarrow G)|.$$

We write V_λ for the irreducible representation of \tilde{G} with highest weight λ ; we can regard V_λ as a G -representation if and only if $\lambda \in \Lambda_G$, and all irreducible representations of G arise in this way. We have a direct sum decomposition

$$\Lambda_{\tilde{G}} = \bigoplus_{i=1}^m \Lambda_{G_i},$$

and we write $\lambda = \lambda_1 + \dots + \lambda_m$. Thus λ is a dominant weight of \tilde{G} if and only if λ_i is a dominant weight of G_i for all i . There is a tensor decomposition

$$V_\lambda = V_{\lambda_1} \otimes \dots \otimes V_{\lambda_m}.$$

Let R (resp. R_i) denote the root system of G (resp. G_i). We fix a positive Weyl chamber of R (equivalently, positive Weyl chambers for each R_i .) For each root α , we denote the dual root $\alpha^\vee = \frac{2\alpha}{\|\alpha\|^2}$, and we write ρ (resp. ρ_i) for the half-sum of positive roots in R (resp. R_i). The Weyl dimension formula [1] VIII §9 Th. 2 asserts

$$\dim V_\lambda = \prod_{\alpha \in R^+} \frac{\alpha^\vee(\lambda + \rho)}{\alpha^\vee(\rho)} = \prod_{i=1}^m \prod_{\alpha \in R_i^+} \frac{\alpha^\vee(\lambda_i + \rho_i)}{\alpha^\vee(\rho_i)} = \prod_{i=1}^m \dim V_{\lambda_i}.$$

It follows that

$$(1.0.1) \quad \dim V_{p\lambda_i + (p-1)\rho_i} = p^{|R_i^+|} \dim V_{\lambda_i}.$$

Thus,

$$\dim V_{(p^{a_i}-1)\rho_1 + \dots + (p^{a_m}-1)\rho_m} = p^{\sum a_i |R_i^+|}.$$

Note that when p is odd, $(p^{a_i} - 1)\rho_i$ lies in the root lattice of G_i , so

$$\sum_{i=1}^m (p^{a_i} - 1)\rho_i \in \Lambda_G.$$

Therefore, every coefficient of $\zeta_G(s)$ is greater than equal to the corresponding coefficient of

$$\prod_{i=1}^m (1 - p^{-|R_i^+|s})^{-1}.$$

We will see that for fixed G , when $p \gg 0$ is a sufficiently large prime, the latter expression gives precisely the part of the Dirichlet series $\zeta_G(s)$ consisting of p -power terms.

LEMMA 1.1: *An irreducible root system R cannot be contained in the union of two proper subspaces of $\text{Span } R$.*

Proof: Let $E = \text{Span } R$ and V and W proper subspaces such that $R \subset V \cup W$. We use induction on $\dim E$, the lemma being trivial when $\dim E = 1$. Replacing V and W by $\text{Span}(V \cap R)$ and $\text{Span}(W \cap R)$ respectively, we may assume without loss of generality that $V \cap R$ and $W \cap R$ are root systems with span V and W respectively. If $W \subset V$, then $R \subset V \subsetneq E = \text{Span } R$, which is absurd; likewise, V is not a subspace of W . If $W \subset V^\perp$, then $R = (R \cap V) \cup (R \cap V^\perp)$, contrary to the hypothesis of irreducibility. By the induction hypothesis, there exists a root α in W which is neither in V nor in V^\perp . Thus $\alpha^\perp \cap V$ is a proper subspace of V , and there exists a root $\beta \in V \cap R$ which is neither in W nor in α^\perp . As α and β are neither scalar multiples of one another nor mutually orthogonal, either $\alpha + \beta$ or $\alpha - \beta$ belongs to R (depending on whether the inner product of α and β is positive or negative.) Either way, $\alpha + \beta \notin V \cup W$, and the lemma follows by induction. \square

LEMMA 1.2: *If $p \gg 0$, then*

$$\dim V_{\lambda_i} \in p^{\mathbb{N}} \Leftrightarrow \lambda_i + \rho_i \in p^{\mathbb{N}} \rho_i.$$

Proof: One direction is trivial. By (1.0.1), it suffices to show $\dim V_{\lambda_i} \in p^{\mathbb{N}}$ implies $\lambda_i = 0$ or $\lambda_i + \rho_i \in p\Lambda_{G_i}$. If p does not divide $\lambda_i + \rho_i$, then

$$S = \{\alpha^\vee \in R_i^\vee \mid \alpha^\vee(\lambda_i + \rho_i) \in p\mathbb{Z}\}$$

is a proper root subsystem of R_i^\vee . For $p \gg 0$, every root in $\alpha^\vee \in R \cap \text{Span } S$ such that $p\alpha^\vee \in S$ lies in S itself. (In fact, an examination of all equal-rank

subsystems of all irreducible root systems show that it suffices to take $p \geq 5$.) Therefore, S is closed. By Lemma 1.1, the complement of S spans R_i^\vee , so there are finitely many elements $\mu \in \Lambda_{G_i}$ such that

$$\beta^\vee(\mu + \rho_i) \leq \prod_{\alpha^\vee \in R_i^\vee} \alpha^\vee(\rho_i)$$

for all $\beta^\vee \in R_i^\vee \setminus S$. If $p \gg 0$, then $p \nmid \dim V_\mu$ for all μ in this set. Thus $\dim V_\mu \in p^\mathbb{N}$ implies $\mu = 0$. \square

COROLLARY 1.3: *The zeta-function of G determines, for each integer n , the number of irreducible factors of the root system of G of cardinality n .*

This is not enough to determine the root system of G . For one thing, the root systems B_i and C_i each have $2i^2$ roots. We also have coincidences with $|R_i| = 12$ (A_3 and G_2) $|R_i| = 72$ (B_6/C_6 , and E_6), $|R_i| = 240$ (A_{15} and E_8), and a sparse but infinite set of cases in which a root system of type A has the same number of roots of one of type B, C, or D. So we need to refine the method. We note, however, that fixing $\zeta_G(s)$ determines the rank of G . We may therefore define N_G to be the factorial of the number of roots in the root system of G . Thus N_G depends on the one hand only on the Lie algebra of G and on the other only on the zeta-function of G . It also has the property that $N_G \Lambda_G$ is contained in the root lattice of G . We define

$$\zeta_G^*(s) = \zeta_G(s) \prod_{p \equiv 1 \pmod{N_G}} \prod_{i=1}^m (1 - p^{-|R_i^+|s}).$$

Note that for $p \equiv 1 \pmod{N_G}$,

$$\lambda_1 + \cdots + \lambda_m \in \Lambda_G \Leftrightarrow \lambda_1 + \cdots + \lambda_{i-1} + (p\lambda_i + (p-1)\rho_i) + \lambda_{i+1} + \cdots + \lambda_m \in \Lambda_G.$$

DEFINITION 1.4: *We say that a weight λ_i is **allowable at a prime p** if $p \nmid \lambda_i + \rho_i$.*

DEFINITION 1.5: *We say that a weight $\lambda_1 + \cdots + \lambda_m \in \Lambda_G$ is **allowable** if for every prime $p \equiv 1 \pmod{N_G}$ and every positive integer $i \leq m$, λ_i is allowable at p . We write Λ_G^* for the set of allowable dominant weights of G . Thus,*

$$\zeta_G^*(s) = \sum_{\lambda \in \Lambda_G^*} \dim(V_\lambda)^{-s}.$$

It is easier to work with $\zeta_G^*(s)$ than with $\zeta_G(s)$ because for $\lambda \in \Lambda_G^*$, the condition that $\dim V_\lambda$ is highly divisible by $p \equiv 1 \pmod{N_G}$ gives substantial information about the structure of the Lie algebra of G , whereas for non-allowable λ , the same divisibility condition could hold for essentially trivial reasons.

2. Weyl polynomials

In this section we consider a fixed simply connected almost simple group G , i.e., G is connected and simply connected, and $G/Z(G)$ is simple. We are interested in the dimensions of representations of G belonging to one-parameter families, specifically the families $p \mapsto p\mu + \nu$. We let R denote the root system of G ; by the weight lattice of R , we mean the weight lattice of G .

DEFINITION 2.1: *Given an irreducible root system R and elements μ and ν of the weight lattice of R , the **Weyl polynomial** $P_{\mu,\nu}^R(x) \in \mathbb{Q}[x]$ is given by the formula*

$$P_{\mu,\nu}^R(x) = \frac{\prod_{\alpha \in R^+} (\alpha^\vee(\mu)x + \alpha^\vee(\nu + \rho))}{\prod_{\alpha \in R^+} \alpha^\vee(\rho)}.$$

When $\lambda = n\mu + \nu$ is dominant,

$$(2.1.1) \quad \dim V_\lambda = P_{\mu,\nu}^R(n)$$

We will always assume that the following two conditions hold:

$$(2.1.2) \quad \alpha^\vee(\mu) \geq 0 \quad \forall \alpha \in R^+$$

and

$$(2.1.3) \quad \alpha^\vee(\mu) = 0 \Rightarrow \alpha^\vee(\nu) \geq 0 \quad \forall \alpha \in R^+.$$

This implies that (2.1.1) holds for all $n \gg 0$. If p is a prime greater than the maximum of $|\alpha^\vee(\mu)|$, $|\alpha^\vee(\nu + \rho)|$, and $|\alpha^\vee(\rho)|$ for any root α , then $\text{ord}_p(\dim V_{p\mu+\nu})$ is equal to $|(\nu + \rho)^\perp \cap R^+|$, the multiplicity of 0 as a root of $P_{\mu,\nu}^R(x)$. In particular, this multiplicity is $|R^+|$ if and only if $\nu + \rho = 0$, which corresponds to the case that for all p , $p\mu + \nu$ is not allowable at p .

PROPOSITION 2.2: *For every irreducible root system R , there exist weights μ and ν satisfying (2.1.2) and (2.1.3) and such that $(\nu + \rho)^\perp \cap R$ is a proper closed root subsystem of R of maximal order. Moreover, we can choose μ and ν such that $\mu + \nu$ lies in the root lattice.*

Proof: Let ϖ_i denote the i th fundamental weight in the Bourbaki ordering [1] Planches. By definition, $\alpha_j^\vee(\varpi_i) = \delta_{ij}$, so the Dynkin diagram of the root system $\varpi_i \perp \cap R$ is obtained from that of R by deleting the i th vertex.

All claims made below about Weyl orbits of weights ϖ_i for exceptional groups were checked with [4]. Note that [4] uses the Dynkin root ordering, which differs from the Bourbaki ordering for E_n .

For $R = A_n$, we may take

$$\mu = -\nu = \varpi_2 + \varpi_3 + \cdots + \varpi_n.$$

As $\nu + \rho = \varpi_1$ is the fundamental weight associated to vertex 1, $(\nu + \rho)^\perp \cap R$ is of type A_{n-1} .

For $R = B_n$, we may take

$$\mu = -\nu = \varpi_2 + \varpi_3 + \cdots + \varpi_n.$$

As $\nu + \rho = \varpi_1$, $(\nu + \rho)^\perp \cap R$ is of type B_{n-1} .

For $R = C_n$, we may take

$$\mu = -\nu = \varpi_1 + \varpi_2 + \cdots + \varpi_{n-2} + 2\varpi_{n-1}.$$

As $\nu + \rho = \varpi_n - \varpi_{n-1}$ lies in the Weyl orbit of ϖ_1 , $(\nu + \rho)^\perp \cap R$ is of type C_{n-1} .

For $R = D_n$, we may take

$$\mu = -\nu = \varpi_1 + \varpi_2 + \cdots + \varpi_{n-3} + 2\varpi_{n-2}.$$

As $\nu + \rho = \varpi_n + \varpi_{n-1} - \varpi_{n-2}$ lies in the Weyl orbit of ϖ_1 , $(\nu + \rho)^\perp \cap R$ is of type D_{n-1} .

For $R = E_6$, we may take

$$\mu = \varpi_2 + \varpi_3 + \varpi_5 + \varpi_6, \quad \nu = -\varpi_2 - 2\varpi_3 - 2\varpi_5 - \varpi_6.$$

As $\nu + \rho = \varpi_1 - \varpi_3 + \varpi_4 - \varpi_5$ lies in the Weyl orbit of ϖ_1 , $(\nu + \rho)^\perp \cap R$ is of type D_5 .

For $R = E_7$, we may take

$$\mu = \varpi_1 + \varpi_3 + \varpi_4 + \varpi_6, \quad \nu = -\varpi_1 - \varpi_3 - 2\varpi_4 - 2\varpi_6.$$

As $\nu + \rho = \varpi_2 - \varpi_4 + \varpi_5 - \varpi_6 + \varpi_7$ lies in the Weyl orbit of ϖ_7 , $(\nu + \rho)^\perp \cap R$ is of type E_6 .

For $R = E_8$, we may take

$$\mu = \varpi_1 + \varpi_3 + \varpi_4 + \varpi_6 + \varpi_8, \quad \nu = -\varpi_1 - \varpi_3 - 2\varpi_4 - 2\varpi_6 - 2\varpi_8.$$

As $\nu + \rho = \varpi_2 - \varpi_4 + \varpi_5 - \varpi_6 + \varpi_7 - \varpi_8$ lies in the Weyl orbit of ϖ_8 , $(\nu + \rho)^\perp \cap R$ is of type E_7 .

For $R = F_4$, we may take

$$\mu = \varpi_1 + \varpi_2, \quad \nu = -\varpi_1 - 2\varpi_2.$$

As $\nu + \rho = -\varpi_2 + \varpi_3 + \varpi_4$ lies in the Weyl orbit of ϖ_4 , $(\nu + \rho)^\perp \cap R$ is of type B_3 .

For $R = G_2$, we may take

$$\mu = \varpi_2, \quad \nu = -2\varpi_2.$$

As $\nu + \rho = \varpi_1 - \varpi_2$ lies in the Weyl orbit of ϖ_2 , $(\nu + \rho)^\perp \cap R$ is of type A_1 .

Except in the exceptional cases, $\mu + \nu = 0$, so the root lattice condition is trivial. It is likewise trivial for E_8 , F_4 , and G_2 . This leaves E_6 and E_7 , where the claim can be checked by [4]. \square

We recall that the **height** of a rational number m/n is $\max(|m|, |n|)$, where $\gcd(m, n) = 1$.

LEMMA 2.3: *If R_1 and R_2 are irreducible root systems and μ_i, ν_i are roots of R_i for $i = 1, 2$, either $P_{\mu_1, \nu_1}^{R_1}(x)$ is a constant multiple of $P_{\mu_2, \nu_2}^{R_2}(x)$ or the height*

$$\text{ht}\left(\frac{P_{\mu_1, \nu_1}^{R_1}(p)}{P_{\mu_2, \nu_2}^{R_2}(p)}\right)$$

goes to ∞ as p increases without bound.

Proof: More generally, if $f(x)$ is any non-constant rational function over \mathbb{Q} , the height of $f(n)$ goes to ∞ as $n \rightarrow \infty$. Indeed, writing $f(x)$ as $\frac{aQ(x)}{bR(x)}$, where a and b are relatively prime integers and $Q(x)$ and $R(x)$ are monic polynomials with integer coefficients, the greatest common divisor of $Q(n)$ and $R(n)$ divides the (non-zero) resultant of the two polynomials. Therefore, if either Q or R is different from 1, the height of $Q(n)/R(n)$ (and therefore the height of $f(n)$) goes to infinity. \square

3. Efficiency of root systems

In this section we give a definition of efficiency purely in terms of root systems and compute the efficiency of each simple root system. It is not obvious that this coincides with the characterization (0.0.1) given in the introduction, though the equivalence can be deduced easily from Proposition 4.3 below.

DEFINITION 3.1: *The **efficiency** $\text{eff}(R)$ of an irreducible root system R is the supremum of $\frac{|R'|}{|R|-|R''|}$, where R' and R'' are closed root subsystems of R such that $R' \cap R'' = \emptyset$. The **level** $\text{lev}(R)$ is the smallest value of $|R'^+|$ for which $\frac{|R'|}{|R|-|R''|}$ achieves $\text{eff}(R)$.*

This definition is connected to the study of Weyl polynomials by the following proposition:

PROPOSITION 3.2: *Let R be an irreducible root system with half-sum of roots ρ , and consider all pairs of weights μ and ν of R satisfying (2.1.2), (2.1.3), and $\nu + \rho \neq 0$. The maximum of*

$$\frac{\text{ord}_{x=0} P_{\mu,\nu}^R(x)}{\deg P_{\mu,\nu}^R(x)}$$

is given by $\text{eff}(R^\vee)$, and this maximum is achieved only when $\text{ord}_{x=0} P_{\mu,\nu}^R(x) \geq \text{lev}(R^\vee)$.

Note that $R = R^\vee$ except for the case B_n/C_n when $n \geq 3$, and even here, as we will see, $\text{eff}(R) = \text{eff}(R^\vee)$ and $\text{lev}(R) = \text{lev}(R^\vee)$, so in fact Proposition 3.2 remains true when all the occurrences of R^\vee are replaced with R .

Proof: Let $(R^\vee)' = (\nu + \rho)^\perp \cap R^\vee$, and $(R^\vee)'' = \mu^\perp \cap R^\vee$. As $\nu + \rho \neq 0$, $(R^\vee)''$ is a proper subsystem of R^\vee . If $\mu = 0$, $P_{\mu,\nu}^R(x)$ is a non-zero constant, so the maximum will not be achieved by such a choice. Therefore, $(R^\vee)'$ and $(R^\vee)''$ are proper closed root subsystems of R^\vee , disjoint by (2.1.3). A factor $\alpha^\vee(\mu)x + \alpha^\vee(\nu + \rho)$ of $P_{\mu,\nu}^R(x)$ has degree 0 if $\alpha \in (R^\vee)''$ and 1 otherwise. The order of this polynomial at 0 is 1 if $\alpha \in (R^\vee)'$ and 0 otherwise. Taking the product over R^+ , we see that

$$\frac{\text{ord}_{x=0} P_{\mu,\nu}^R(x)}{\deg P_{\mu,\nu}^R(x)} \leq \text{eff}(R^\vee)$$

for all μ and ν satisfying the hypotheses of the proposition. All that remains to be shown is that a pair $((R^\vee)', (R^\vee)'')$ achieving the bound $\text{eff}(R)$ can always

be found by taking a suitable choice of μ and ν . We claim that the choices in Proposition 2.2 do the trick. This follows from the explicit computation of efficiencies of all irreducible root systems given by the following proposition.

PROPOSITION 3.3: *The efficiencies of root systems of type A_n , B_n , C_n , D_n , E_6 , E_7 , E_8 , F_4 , and G_2 are $\frac{n}{n+2}$, $\frac{n-1}{n+1}$, $\frac{n-1}{n+1}$, $\frac{n-1}{n+1}$, $\frac{10}{17}$, $\frac{3}{5}$, $\frac{63}{117}$, $\frac{3}{7}$, and $\frac{1}{5}$ respectively. In each case the efficiency is achieved only when R' is a proper closed root subsystem of maximal order in R . Explicitly, the levels of these root systems are $\frac{n^2-n}{2}$, $(n-1)^2$, $(n-1)^2$, $(n-1)(n-2)$, 20, 36, 63, 9, and 1 respectively. Two inequivalent irreducible root systems with the same efficiency and the same level must be of the form B_n and C_n for the same $n \geq 3$.*

Proof: We prove that in each case the efficiency is achieved when R' is the largest proper root subsystem and R'' is as large as possible given $R' \cap R'' = \emptyset$. It suffices to determine R'_0 and R''_0 satisfying these conditions and show that if R' is not of maximal order, then $\frac{|R'|}{|R|-|R''|} < \frac{|R'_0|}{|R|-|R''_0|}$. In particular, it suffices to prove $|R'| + |R''| \leq |R'_0| + |R''_0|$.

For every closed root subsystem $S \subset R$, there exist choices of basis for R and S such that every positive root of S is positive in R and every simple root of S is simple in R ([1] VI, §1, Prop. 24). Thus we can regard root subsystems as arising from subgraphs of the Dynkin diagram of R . This is convenient, among other things, for computing the largest subsystems of the different root systems (which are always obtained by removing a single vertex from the Dynkin diagram). Explicitly, the proper closed root systems of maximal order for A_n , B_n , C_n , D_n , E_6 , E_7 , E_8 , F_4 , and G_2 are A_{n-1} , B_{n-1} , C_{n-1} , D_{n-1} , D_5 , E_6 , E_7 , B_3/C_3 , and A_1 respectively. These give the values for $\text{lev}(R)$ stated in the proposition.

It is also worth noting that if $\text{rank } R'$ is $n-1$, and f is a non-zero linear form vanishing on the hyperplane $\text{Span } R'$, then the Coxeter number h of every simple factor of R'' disjoint from R' satisfies

$$(3.3.1) \quad h \leq 1 + \sup_{s_1, s_2 \in R \setminus R'} \frac{f(s_1)}{f(s_2)}.$$

Indeed, a choice of half-plane with boundary $\text{Span } R'$ determines a basis on R'' . If $\alpha_1, \dots, \alpha_t$ are the simple roots and $\alpha = \sum n_i \alpha_i$ the highest root of a simple factor of R'' with Coxeter number h , then

$$\sup_{s_1, s_2 \in R \setminus R'} \frac{f(s_1)}{f(s_2)} \geq \sup_j \sum_{i=1}^t n_i \frac{f(\alpha_i)}{f(\alpha_j)} \geq \sum_{i=1}^t n_i = h - 1.$$

The value on the right hand side of (3.3.1) can be computed easily by restricting the adjoint representation of a group of type R to $U(1)$ times a group of type R' and looking at the $U(1)$ -components of the resulting direct sum.

With these general facts in hand, we now examine the cases.

For A_n , we use ε -coordinates, in which the roots are the vectors of length $\sqrt{2}$ in $\ker \Sigma: \mathbb{Z}^{n+1} \rightarrow \mathbb{Z}$, Σ denoting the sum of coordinates. Let $R'_0 = \varepsilon_1^\perp \cap A_n = A_{n-1}$ and $R''_0 = \{\pm(\varepsilon_1 - \varepsilon_2)\}$. We define an equivalence relation on $\{1, \dots, n+1\}$ by setting $i \sim j$ if and only if $\varepsilon_i - \varepsilon_j \in R' \cup \{0\}$. The equivalence classes I_1, \dots, I_c have cardinality a_1, \dots, a_c where $\sum a_k = n+1$ and $R' = \prod_{a_k > 1} A_{a_k-1}$. Given $k \neq l$, the set $\{\varepsilon_i - \varepsilon_j \in R'' \mid i \in I_k, j \in I_l\}$ has at most $\min(a_i, a_j) \leq (a_i - 1)(a_j - 1) + 1$ elements since R' and R'' are disjoint. Thus

$$\begin{aligned} |R'| + |R''| &\leq \sum_k a_k^2 - a_k + \sum_{k \neq l} (a_k a_l - a_k - a_l + 2) \\ &= (n+1)^2 - (n+1) - 2(c-1)(n+1) + 2c(c-1) \\ &= (n+2-c)^2 + c^2 - (n+2). \end{aligned}$$

Thus $|R'| + |R''| < n^2 - n + 2$ unless $c \leq 2$ or $c \geq n$. For $c \leq 2$, $|R''| \leq 2$, and for $c \geq n$, $|R'| \leq 2$, so either way $|R'| + |R''|$ is bounded above by $|R'_0| + |R''_0|$.

For B_n (resp. C_n) we use the ε -coordinate system, in which the short (resp. long) vectors are ε_i (resp. $2\varepsilon_i$). Let $R'_0 = \varepsilon_1^\perp \cap B_n = B_{n-1}$ (resp. $\varepsilon_1^\perp \cap C_n = C_{n-1}$) and $R''_0 = \{\pm\varepsilon_1\}$ (resp. $\{\pm 2\varepsilon_1\}$). Let I' be the set of $i \in \{1, \dots, n\}$ such that ε_i (resp. $2\varepsilon_i$) lies in R' and I'' the corresponding set for R'' . As R' and R'' are disjoint, $I' \cap I'' = \emptyset$, and we let I denote the complement of $I' \cup I''$. By disjointness, $\pm\varepsilon_i \pm \varepsilon_j \notin R''$ for $i, j \in I'$ and $\pm\varepsilon_i \pm \varepsilon_j \notin R'$ for $i, j \in I''$; the set $\{\pm\varepsilon_i \pm \varepsilon_j \mid i \in I, j \in I'\}$, which is disjoint from R' , meets R'' in at most $2 \min(|I|, |I'|)$ elements, and likewise $\{\pm\varepsilon_i \pm \varepsilon_j \mid i \in I, j \in I''\}$, which is disjoint from R'' , meets R' in at most $2 \min(|I|, |I''|)$ elements. Finally, for each $i, j \in I$, $i \neq j$, the set $\{\pm\varepsilon_i \pm \varepsilon_j\}$ meets R' in at most 2 vectors and R'' in at most 2 vectors. Combining these facts, we get

$$\begin{aligned} (3.3.2) \quad |R'| + |R''| &\leq (2|I'|^2 + |I|^2 - |I| + 2 \min(|I|, |I''|)) \\ &\quad + (2|I''|^2 + |I|^2 - |I| + 2 \min(|I|, |I'|)) \\ &\leq (2|I'|^2 + |I|^2 - |I| + |I| + |I''|) + (2|I''|^2 + |I|^2 - |I| + |I| + |I'|) \\ &= 2|I|^2 + 2(|I'| + \frac{1}{4})^2 + 2(|I''| + \frac{1}{4})^2 - \frac{1}{4}. \end{aligned}$$

As $|I| + |I'| + |I''| = n$, for $n \geq 4$, this quantity is $\leq |R'_0| + |R''_0| = 2n^2 - 4n + 4$ unless one of $|I|$, $|I'|$, and $|I''|$ is at least $n - 1$. If $|I| = n - 1$, then the first inequality of (3.3.2) reads

$$|R'| + |R''| \leq 2n^2 - 6n + 8 \leq 2n^2 - 4n + 4 = |R'_0| + |R''_0|.$$

If $|I| = n$, then $|R'| \leq n^2 - n$ and $|R''| \leq n^2 - n$, with equality only if the systems are of type A_{n-1} . If R' is of type A_{n-1} , the right hand side of (3.3.1) is 3 (resp. 2), so assuming $n \geq 4$, R'' cannot also be of type A_{n-1} . Thus,

$$\frac{|R'|}{|R| - |R''|} < \frac{n(n-1)}{2n^2 - n(n-1)} = \frac{n-1}{n+1} = \frac{|R'_0|}{|R| - |R''_0|}.$$

Thus $|R'| + |R''|$ is indeed maximized when R' is B_{n-1} (resp. C_{n-1}). The same is true for $n = 2$ and $n = 3$ by a simple case analysis.

For D_n , we again use the ε -coordinate system, in which the roots are the length $\sqrt{2}$ vectors in \mathbb{Z}^n . Let $R'_0 = \varepsilon_1^\perp \cap D_n = D_{n-1}$ and $R''_0 = \{\pm\varepsilon_1 \pm \varepsilon_2\} = D_2$. The subsystem R' defines an equivalence relation \sim on $\{1, 2, \dots, n\}$, where $i \sim j$ if $i = j$ or $\{\pm\varepsilon_i \pm \varepsilon_j\} \subset R'$. As R' is a closed root subsystem, there can be at most one \sim equivalence class with 2 or more elements; indeed,

$$\{\pm\varepsilon_i \pm \varepsilon_k\} \subset \text{Span}\{\pm\varepsilon_i \pm \varepsilon_j, \pm\varepsilon_k \pm \varepsilon_l\}.$$

Let I' denote this class (or the empty set) and I'' the corresponding set for R'' . As $R' \cap R'' = \emptyset$, $|I' \cap I''| \leq 1$. Let I denote the complement of $I' \cup I''$. Reasoning as in the case B_n/C_n , $\pm\varepsilon_i \pm \varepsilon_j \notin R''$ for $i, j \in I'$ and $\pm\varepsilon_i \pm \varepsilon_j \notin R'$ for $i, j \in I''$; the set $\{\pm\varepsilon_i \pm \varepsilon_j \mid i \in I, j \in I'\}$, which is disjoint from R' , meets R'' in at most $2 \min(|I|, |I'|)$ elements, and likewise $\{\pm\varepsilon_i \pm \varepsilon_j \mid i \in I, j \in I''\}$, which is disjoint from R'' , meets R' in at most $2 \min(|I|, |I''|)$ elements. Again for each $i, j \in I$, $i \neq j$, the set $\{\pm\varepsilon_i \pm \varepsilon_j\}$ meets R' in at most 2 vectors and R'' in at most 2 vectors, so

$$\begin{aligned} |R'| + |R''| &\leq (2|I'|^2 - 2|I'| + |I|^2 - |I| + 2 \min(|I|, |I''|)) \\ &\quad + (2|I''|^2 - 2|I''| + |I|^2 - |I| + 2 \min(|I|, |I'|)) \\ &\leq 2|I|^2 + 2|I'|^2 - |I'| + 2|I''|^2 - |I''| \\ &= 2|I|^2 + 2(|I'| - \frac{1}{4})^2 + 2(|I''| - \frac{1}{4})^2 - \frac{1}{4}. \end{aligned}$$

Moreover, $|I'|, |I''| \leq n-1$, $|I|+|I'| \leq n$, $|I|+|I''| \leq n$, and $|I|+|I'|+|I''| \leq n+1$.
If

$$|R'| + |R''| > |R'_0| + |R''_0| = 2n^2 - 6n + 8,$$

either $|I|$, $|I'|$, or $|I''|$ is at least $n-1$. Now $|I| \geq n-1$ implies $I' = I'' = \emptyset$, so either $R' = A_{n-1}$ or $|R'| \leq (n-1)(n-2)$, and likewise for R'' . If $|R'| \leq (n-1)(n-2)$ and $|R''| \leq (n-1)(n-2)$, then $\frac{|R'|}{|R|+|R''|} \leq \frac{n-2}{n+1} < \frac{|R'_0|}{|R|+|R''_0|}$. If $R' = A_{n-1}$, the upper bound (3.3.1) is 2, so R'' is of the form A_1^r (and vice versa). If R' or R'' has r mutually orthogonal roots, since they are closed root subsystems, the Dynkin diagram of D_n must have r vertices, no two of which share an edge, so $r \leq n/2 + 1$, and

$$|R'| + |R''| \leq n(n-1) + n + 2 \leq 2n^2 - 6n + 8 = |R'_0| + |R''_0|,$$

as long as $n \geq 5$. For $n = 4$, we check by hand that there do not exist 3 mutually orthogonal roots in $D_4 \setminus A_3$, so (D_3, D_2) is still optimal.

For E_8 , $|R| = 240$. The largest closed subsystem, E_7 , has 126 roots, and it can be realized as the set of roots orthogonal to a given root α ; D_7 is next with 84. As $\frac{126}{240-6} > \frac{84}{240-84}$, it suffices to prove that the largest closed root subsystem of E_8 disjoint from $E_7 = \alpha^\perp \cap E_8$ has order 6. The orthogonal projection of any root onto $\mathbb{R}\alpha$ is in $\{-\alpha, -\frac{1}{2}\alpha, 0, \frac{1}{2}\alpha, \alpha\}$, so the right hand side of (3.3.1) is 3. Thus all factors of R'' are of type A_1 or A_2 . Moreover, a factor of type A_2 must contain α itself (which is the only root whose projection onto $\mathbb{R}\alpha$ equals α), so if R'' contains any factor of the form A_2 , then $R'' = A_2$. Otherwise, $R'' = A_1^n$. We look for the largest value such that A_1^n can be embedded as a closed subsystem in E_8 with no root orthogonal to α . Equivalently, we fix an embedding of A_1^n in E_8 and ask whether any root of E_8 fails to be orthogonal to every root of A_1^n . Equivalently, we ask whether the restriction of the adjoint representation of E_8 to A_1^n has an irreducible summand which is non-trivial on each factor A_1 .

To construct mutually orthogonal roots in E_8 , we start with a root β_1 , choose $\beta_2 \in \beta_1^\perp \cap E_8 = E_7$, choose $\beta_3 \in \beta_2^\perp \cap E_7 = D_6$, choose $\beta_4 \in \beta_3^\perp \cap D_6 = A_1 \times D_4$, and so on. All choices are equivalent until β_4 , which can be chosen in A_1 or in D_4 . In the first case, $A_1^4 \subset D_4^\perp \cap E_8 = D_4$, so the system is not closed. We use branching rules [4], starting with the adjoint representation of E_8 , restricting to $A_1 \times E_7$, taking only representations of E_7 which appear opposite a non-trivial factor in A_1 , and so on. We get the (56-dimensional) representation V_{ϖ_7} of E_7 ,

and then the (12-dimensional) representation V_{ϖ_1} of D_6 , then a representation of $A_1 \times D_4$ which is 2-dimensional and therefore trivial on the D_4 factor. We conclude that A_1^3 can be realized (and it is indeed a closed subsystem of E_8 since no simply laced rank 3 system contains A_1^3 as a proper subsystem), but A_1^4 cannot.

For E_7 , $|R| = 126$. The largest subsystem, E_6 , has 72 roots; D_6 comes next, with 60, and then A_6 , with 42. As $\frac{72}{126-6} > \frac{42}{126-42}$, we need only consider the cases $R' = E_6$ and $R' = D_6$. Writing E_6 as the orthogonal complement of the fundamental weight ϖ_7 the right hand side of (3.3.1) is 2, so we need only consider $R'' = A_1^r$. We proceed as before, with $\beta_1 \in E_7$, $\beta_2 \in \beta_1^\perp \cap E_7 = D_6$, and $\beta_3 \in \beta_2^\perp \cap D_6 = A_1 \times D_4$. Applying the branching rules for V_{ϖ_7} , which we have already described, we see that if we choose β_3 to be in the factor A_1 , the restriction of V_{ϖ_7} to A_1^3 has a summand non-trivial on every factor, so we can take $R'' = A_1^3$. Any fourth root β_4 mutually orthogonal to the first three must lie in the D_4 factor, and we know from the E_8 analysis that ϖ_7 cannot be non-orthogonal to β_1 , β_2 , and such a β_4 . When $R' = D_6 = \alpha^\perp \cap E_7$, the argument for E_8 shows that either $R'' = A_2$ or $R'' = A_1^n$ where $n \leq 7$ for dimension reasons. As $\frac{72}{126-6} > \frac{60}{126-14}$, we see that (E_6, A_1^3) is optimal.

For E_6 , $|R| = 72$. The largest subsystem, D_5 , has 40 roots, the next largest, A_5 , has 30, and the third largest, D_4 , has 24. As $\frac{40}{72-4} > \frac{24}{72-24}$, we need only consider $D_5 = \varpi_1^\perp \cap E_6$ and $A_5 = \alpha^\perp \cap E_6$, where α is a root. For $R' = D_5$, the right hand side of (3.3.1) is 2, so $R'' = A_1^n$. Choose $\beta_1 \in E_6$, $\beta_2 \in \beta_1^\perp \cap E_6 = A_5$, and $\beta_3 \in \beta_2^\perp \cap A_5 = A_3$. In the restriction of V_{ϖ_1} to $A_1 \times A_5$, the only summand non-trivial on A_1 is 6-dimensional on A_5 ; its restriction to $A_1 \times A_3$ is the a direct sum of a representation trivial on A_1 and a representation trivial on A_3 . It follows that the restriction of V_{ϖ_1} to A_1^2 has a summand non-trivial on both factors but the restriction to A_1^3 has no such summand. For $R' = A_5$, the right hand side of (3.3.1) is 3, so as in the E_8 case, $R'' = A_2$ or $R'' = A_1^n$, and of course $n \leq 5$. As $\frac{40}{72-4} > \frac{30}{72-10}$, $(R', R'') = (D_5, A_1^2)$ is optimal.

For F_4 , $|R| = 48$, and the largest proper closed root subsystems are obtained by deleting an endpoint vertex from the Dynkin diagram. They are of type B_3 and C_3 , and are both of order 9; in each case, $\text{Span } R' = \alpha^\perp \cap R$ for a root α , short or long respectively. Either way, the bound on the Coxeter number of components of R'' is 3, so we need only consider A_1 , A_2 , $A_1 \times A_1$, and $A_2 \times A_1$ as possibilities for R'' . In each case, A_2 can be realized; for example, if $\alpha = \varpi_4$

with respect to the standard coordinates on F_4 , R'' can be taken to have basis $\{2\varpi_4 - \varpi_3, \varpi_3 - \varpi_4\}$. In each case, α or $-\alpha$ is the highest vector of the A_2 , so $A_2 \times A_1$ cannot be disjoint from $\text{Span } R'$ (since $\pm\alpha$ belong to both). Finally, if R' is not of type B_3 or C_3 , it has at most 8 roots, and $8 + 8 < 18 + 6$.

The case G_2 is trivial since any proper root subsystem has rank 1 and therefore order 2.

□

DEFINITION 3.4: *We say an irreducible root system S is **greater than or equal to** an irreducible root system T (and write $S \geq T$) if and only if $\text{eff}(S) > \text{eff}(T)$ or $\text{eff}(S) = \text{eff}(T)$ and $\text{lev}(S) \leq \text{lev}(T)$. If $S \geq T$ but $T \not\geq S$, we say S is **strictly greater than** T .*

An immediate consequence of Proposition 3.3 is that $S \geq T$ and $T \geq S$ implies $S = T$ or $S = T^\vee$.

4. The main theorem

LEMMA 4.1: *Let R be an irreducible root system spanning a Euclidean space E and Q a quadratic form on E such that $Q(\alpha) = 0$ for all $\alpha \in R$. Then $Q = 0$.*

Proof: Let $B(x, y) = Q(x + y) - Q(x) - Q(y)$ denote the associated bilinear form. If α and β are roots of R making an obtuse angle, then $\alpha + \beta$ is again a root, so $B(\alpha, \beta) = 0$. Likewise, if α and β make an acute angle, $\alpha - \beta$ is a root, so again $B(\alpha, \beta) = 0$. The roots in $R \setminus \alpha^\perp$ span E by Lemma 1.1. Therefore, $B(\alpha, E) = 0$. As every $\alpha \in R$ lies in the null space of B , it follows that B and therefore Q must be 0. □

PROPOSITION 4.2: *If R_1 and R_2 are irreducible root systems with $R_1 \geq R_2$, then there exist weights μ_1 and ν_1 and an integer N such that:*

- 1) μ_1 and ν_1 satisfy (2.1.2) and (2.1.3);
- 2) $\nu_1 + \rho_1 \neq 0$;
- 3) $\mu_1 + \nu_1$ lies in the root lattice of R_1 ;
- 4) $\frac{\text{ord}_{x=0} P_{\mu_1, \nu_1}^{R_1}(x)}{\deg P_{\mu_1, \nu_1}^{R_1}(x)} = \text{eff}(R_1)$;
- 5) If μ_2 and ν_2 are weights of R_2 and c is a positive constant such that

$$(4.2.1) \quad P_{\mu_2, \nu_2}^{R_2}(x) = c P_{\mu_1, \nu_1}^{R_1}(x),$$

then

- a) $\text{eff}(R_1) = \text{eff}(R_2)$ and $\text{lev}(R_1) = \text{lev}(R_2)$.
- b) $c \geq 1$, with equality only if $R_1 = R_2$;
- c) If $p > N$ and $\text{ord}_p c > 0$, then p divides $\nu_2 + \rho_2$, i.e., ν_2 is not allowable at p .

Proof:

First we observe that condition (4) implies that $P_{\mu_1, \nu_1}^{R_1}(x)$ is not a monomial in x , so that (4.2.1) implies the same for $P_{\mu_2, \nu_2}^{R_2}(x)$. This implies that $\nu_2 + \rho_2 \neq 0$, so

$$\frac{\text{ord}_{x=0} P_{\mu_2, \nu_2}^{R_2}(x)}{\deg P_{\mu_2, \nu_2}^{R_2}(x)} \leq \text{eff}(R_2).$$

The hypothesis $\text{eff}(R_1) \geq \text{eff}(R_2)$ then implies $\text{eff}(R_1) = \text{eff}(R_2)$. Proposition 3.2 and equation (4.2.1) then imply

$$\text{lev}(R_2) \leq \text{ord}_{x=0} P_{\mu_2, \nu_2}^{R_2}(x) = \text{ord}_{x=0} P_{\mu_1, \nu_1}^{R_1}(x) = \text{lev}(R_1),$$

so $\text{lev}(R_2) = \text{lev}(R_1)$.

For (b), we choose μ_1 and ν_1 as in Proposition 2.2. We note that for any integer n , whether or not $\lambda_2 = n\mu_2 + \nu$ is dominant, $P_{\mu_2, \nu_2}^{R_2}(n)$ is either zero or ± 1 times the dimension of an irreducible representation of a Lie group of type R_2 . Indeed, suppose $\alpha^\vee(\lambda_2 + \rho_2) \neq 0$ for all α . Then defining w in the Weyl group of R_2 as the unique element such that $w(\lambda_2 + \rho_2)$ is dominant, and setting $\lambda'_2 = w(\lambda_2 + \rho_2) - \rho_2$, the weight λ'_2 is dominant. Thus,

$$\begin{aligned} \dim V_{\lambda'_2}^2 &= (-1)^{|R_2^+|} \frac{\prod_{\alpha \in R} \alpha^\vee(\lambda'_2 + \rho_2)}{\prod_{\alpha \in R} \alpha^\vee(\rho_2)} \\ &= (-1)^{|R_2^+|} \frac{\prod_{\alpha \in R} \alpha^\vee(\lambda_2 + \rho_2)}{\prod_{\alpha \in R} \alpha^\vee(\rho_2)} = P_{\mu_2, \nu_2}^{R_2}(n)^2. \end{aligned}$$

For R_1 of type A_n , B_n , C_n , or D_n , we have $P_{\mu_1, \nu_1}^{R_1}(1) = 1$, so $P_{\mu_2, \nu_2}^{R_2}(1) = c$ implies $c \in \mathbb{Z}$ and therefore $c \geq 1$. The same is true for G_2 , since $P_{\mu_1, \nu_1}^{R_1}(2) = 1$. For F_4 ,

$$\gcd(P_{\mu_1, \nu_1}^{R_1}(2), P_{\mu_1, \nu_1}^{R_1}(3)) = \gcd(52, 340119) = 1,$$

so again c must be an integer. For E_6 , we have

$$\gcd(P_{\mu_1, \nu_1}^{R_1}(2), P_{\mu_1, \nu_1}^{R_1}(3), P_{\mu_1, \nu_1}^{R_1}(4)) = \gcd(1728, 3171108447, 71292900343808) = 1,$$

so again $c \in \mathbb{Z}$. For E_7 , we have

$$\gcd(P_{\mu_1, \nu_1}^{R_1}(2), P_{\mu_1, \nu_1}^{R_1}(3)) = \gcd(573440, 33940969546604175) = 5,$$

so $5c \in \mathbb{Z}$. Thus, $|P_{\mu_2, \nu_2}^{R_2}(2)|$ is a positive integer multiple of $\frac{573440}{5} = 114688$. However, the lowest degree of any irreducible representation of a Lie group of type E_7 which is divisible by 114688 is 573440 itself. For E_8 ,

$$\begin{aligned} \gcd(P_{\mu_1, \nu_1}^{R_1}(2), P_{\mu_1, \nu_1}^{R_1}(3)) &= \gcd(4096000, 2665014302693985712862760000) \\ &= 8000, \end{aligned}$$

and the smallest dimension of any irreducible representation of a Lie group of type E_8 which is divisible by $512 = \frac{4096000}{8000}$ is 4096000.

To finish proving (b), it suffices to show that if R_1 is B_n or C_n , $n \geq 3$, $R_2 = R_1^\vee$, μ_1 and ν_1 are defined as in Proposition 2.2, and μ_2 and ν_2 are any weights of R_2 , then $P_{\mu_1, \nu_1}^{R_1} \neq \pm P_{\mu_2, \nu_2}^{R_2}$. If μ and ν are weights of B_n such that $\text{ord}_{x=0} P_{\mu, \nu}^{B_n}(x)$ achieves the maximum $(n-1)^2$, then $\nu + \rho$ must be a scalar multiple $d\varepsilon_i$ of an ε -basis vector, and writing $\mu = (a_1, \dots, a_n)$, either $a_i = \pm a_j$ for some $j \neq i$ or some $a_j = 0$. We can always replace ν by any linked weight $w(\nu + \rho) - \rho$ with the effect of changing $P_{\mu, \nu}^{B_n}$ by a factor of ± 1 , so without loss of generality we may assume μ is dominant, which means we may assume either $a_1 > a_2 > \dots > a_{n-1} > a_n = 0$ (and $i = n$) or, renumbering, $a_1 > \dots > a_i = a_i > \dots > a_{n-1} > 0$. In the first case,

$$(4.2.2) \quad x^{-(n-1)^2} P_{\mu, \nu}^{B_n}(x) = \frac{d^{2n} F(a_1, \dots, a_{n-1}) \prod_{i=1}^{n-1} (a_i x - d)(a_i x + d)}{\Delta_n}$$

where

$$\Delta_n = \prod_{\alpha \in B_n} \alpha^\vee(\rho_{B_n}) = \prod_{\alpha \in C_n} \alpha^\vee(\rho_{C_n})$$

and

$$F(a_1, \dots, a_{n-1}) = \prod_{i=1}^{n-1} a_i \prod_{i=1}^{n-2} \prod_{j=i+1}^{n-1} (a_i^2 - a_j^2).$$

In the second case, $x^{-(n-1)^2} P_{\mu, \nu}^{B_n}(x)$ is

$$(4.2.3) \quad \frac{d^{2n} F(a_1, \dots, a_{n-1}) (a_i x + d)(2a_i x + d) \prod_{j \neq i} ((a_i + a_j)x + d)((a_j - a_i)x + d)}{\Delta_n}$$

If A is the number of positive roots of $x^{-(n-1)^2} P_{\mu, \nu}^{B_n}(x)$ and B is the number of negative roots, then (4.2.2) applies if and only if $A = B$, and otherwise (4.2.3) applies with $i = \sup(A, B) - (n-1)$. In either case, the roots determine the

vector (a_1, \dots, a_{n-1}, d) up to scalar multiplication by a positive rational number and therefore the polynomial itself up to multiplication by an element of $(\mathbb{Q}^*)^{n^2}$.

The case analysis for $P_{\mu, \nu}^{C_n}(x)$ is exactly the same, and the formulas (4.2.2) and (4.2.3) the same except without the factor of 2^n . As $2^n \notin (\mathbb{Q}^*)^{n^2}$, $P_{\mu_1, \nu_1}^{R_1} \neq P_{\mu_2, \nu_2}^{R_2}$.

Given two pairs (μ_1, ν_1) and (μ_2, ν_2) of weights of an irreducible root system R , $i = 1, 2$, we have maps $f_i: R^+ \rightarrow \mathbb{Q}^2$ defined by

$$f_i(\alpha) = (\alpha^\vee(\mu_i), \alpha^\vee(\nu_i)).$$

Suppose that $P_{\mu_2, \nu_2}^R(x) = cP_{\mu_1, \nu_1}^R(x) \neq 0$. Then $(0, 0) \notin f_i(R^+)$ and for some permutation $\sigma: R^+ \rightarrow R^+$, $f_2(\alpha)$ is a scalar multiple of $f_1(\alpha)$ for all α . As there are only finitely many possibilities for σ , by making N large enough we may assume without loss of generality that σ is the identity. Thus,

$$\alpha^\vee(\mu_1)\alpha^\vee(\nu_2 + \rho_2) - \alpha^\vee(\mu_2)\alpha^\vee(\nu_1 + \rho_1) = 0$$

for all $\alpha^\vee \in R^\vee$. Regarding the above expression as a quadratic form on R^\vee , by Lemma 4.1, it is trivial, so $(\mu_2, \nu_2 + \rho_2) = C(\mu_1, \nu_1 + \rho_1)$ for some non-zero rational number C . Therefore $c = C^{|R^+|}$, and claim (c) follows. \square

PROPOSITION 4.3: *Let $l \in \mathbb{N}$, $e \in \mathbb{Q}$, and G a simply connected almost simple Lie group such that for every almost simple factor G_i of G with root system R_i , either $\text{eff}(R_i) < e$ or $\text{eff}(R_i) = e$ and $\text{lev}(R_i) \geq l$. Let $A(G)$ denote the set of pairs (p, λ) , where p is a prime and λ is a dominant weight of G allowable at p . Then*

$$(4.3.1) \quad \inf_{(p, \lambda) \in A(G)} e \log \dim V_\lambda - (\log p) \text{ord}_p \dim V_\lambda > -\infty.$$

Moreover, for every $C \in \mathbb{R}$, there exists a finite set

$$L_{G, e, l, C} = \{(\mu, \nu) \mid \mu, \nu \in \Lambda_G\}$$

such that for all $(p, \lambda) \in A(G)$ such that $\text{ord}_p \dim V_\lambda \leq l$ and

$$(4.3.2) \quad e \log \dim V_\lambda - (\log p) \text{ord}_p \dim V_\lambda \leq C,$$

there exists $(\mu, \nu) \in L_{G, e, l, C}$ such that $\lambda = p\mu + \nu$ and either $\mu = 0$, or μ is a non-zero weight of an almost simple factor G_i of G with $\text{eff}(R_i) = e$ and $\text{lev}(R_i) = l$.

Proof: Suppose first that G is almost simple with root system R and Coxeter number h . We take advantage of the equalities $|R| = |R^\vee|$, $\text{eff}(R) = \text{eff}(R^\vee)$, and $\text{lev}(R) = \text{lev}(R^\vee)$ to avoid burdening our notation unnecessarily with $^\vee$ superscripts. Define

$$(4.3.3) \quad S_i = \{\alpha^\vee \in R^\vee : p^i | \alpha^\vee(\lambda + \rho)\}, \quad R'_i = R^\vee \cap \text{Span } S_i$$

Thus,

$$R^\vee \supset R'_1 \supset R'_2 \supset \cdots \supset R'_k \supsetneq R'_{k+1} = \emptyset.$$

An examination of the list of all equal-rank root subsystems of all irreducible root systems shows that $\alpha \in R'_i$ implies either 2α or 3α lies in the abelian group generated by S_i . As λ is allowable at p , R'_i is a proper subsystem of R^\vee except possibly when $i = 1$ and $p \leq 3$. Let R''_i denote the closed root subsystem of R^\vee generated by the simple roots α^\vee such that

$$(4.3.4) \quad 0 < \alpha^\vee(\lambda + \rho) < \frac{p^i}{3h}.$$

For any root in R''_i , $0 < \alpha^\vee(\lambda + \rho) < \frac{p^i}{3}$, so neither $2\alpha^\vee$ nor $3\alpha^\vee$ lies in the abelian group generated by S_i , which means $R'_i \cap R''_i = \emptyset$. We have

$$(4.3.5) \quad \text{ord}_p \dim V_\lambda \leq \sum_{i=1}^k |(R'_i)^+|$$

with equality if $p \geq 5$, and

$$(4.3.6) \quad \log \dim V_\lambda \geq -|R^+| \log 3h + \log p \sum_{i=1}^k (|R^+| - |(R''_i)^+|).$$

Except possibly when $i = 1$ and $p \leq 3$,

$$(4.3.7) \quad \text{eff}(R)(|R^+| - |(R''_i)^+|) \geq |(R'_i)^+|,$$

so

$$(4.3.8) \quad \text{eff}(R) \log p \sum_{i=1}^k (|R^+| - |(R''_i)^+|) \geq -\text{eff}(R)|R^+| \log 3 + \log p \sum_{i=1}^k |(R'_i)^+|.$$

By (4.3.6), (4.3.8), and (4.3.5),

$$\begin{aligned} e \log \dim V_\lambda - \log p \text{ord}_p \dim V_\lambda &\geq \text{eff}(R) \log \dim V_\lambda - \log p \text{ord}_p \dim V_\lambda \\ &\geq -\text{eff}(R)|R^+| \log 9h. \end{aligned}$$

This gives (4.3.1).

If $e > \text{eff}(R)$, then

$$e \log \dim V_\lambda - \log p \text{ord}_p \dim V_\lambda \geq (e - \text{eff}(R)) \log \dim V_\lambda - \text{eff}(R) |R^+| \log 9h$$

implies that the left hand side is $\geq C$ for all but finitely many weights λ , independent of p . We can therefore define $L_{G,e,l,C}$ to be the set of pairs $(0, \nu)$, as ν ranges over all such λ . If $l < \text{lev}(R)$, then for $p \geq 5$, (4.3.5) is an equality, and so $|(R'_i)^+| < \text{lev}(R)$ for $1 \leq i \leq k$. It follows that (4.3.7) is strict and that

$$\begin{aligned} & \text{eff}(R) \log \dim V_\lambda - \log p \text{ord}_p \dim V_\lambda \\ & \geq \{ \text{eff}(R)(|R^+| - |(R''_i)^+|) - |(R'_i)^+| \} \log p - \text{eff}(R) |R^+| \log 9h > C \end{aligned}$$

for $p \gg 0$. We therefore obtain an upper bound on p , which then gives an upper bound on $\dim V_\lambda$ and consequently a finite set of possible λ , and we can set $L_{G,e,l,C}$ to be a set of pairs $(0, \lambda)$ as before. The only way in which $\dim V_\lambda$ could fail to be bounded by the condition (4.3.2) is if $\text{eff}(R) = e$ and $\text{lev}(R) = l$ and moreover $|(R'_i)^+| = \text{lev}(R)$ for all $i \leq k$. For $p \geq 5$, equality holds in (4.3.5), and this last condition implies $k = 1$.

To prove that λ can be written $p\mu + \nu$, where μ and ν are drawn from finite sets independent of p , we may assume that we have fixed $S = R'_1$ and $T = R''_1$. The Weyl formula gives

$$\begin{aligned} \log \dim V_\lambda & \geq (|S^+|) \log p + \sum_{\alpha \in S^+} \log \frac{\alpha^\vee(\lambda + \rho)}{p} + \sum_{\alpha \in T^+} \log \alpha^\vee(\lambda + \rho) \\ & \quad + (|R^+| - |S^+| - |T^+|) \log \frac{p}{3h} - \sum_{\alpha \in R^+} \log \alpha^\vee(\rho) \\ & = (|R^+| - |T^+|) \log p + \sum_{\alpha \in S^+} \log \frac{\alpha^\vee(\lambda + \rho)}{p} + \sum_{\alpha \in T^+} \log \alpha^\vee(\lambda + \rho) \\ & \quad - (|R^+| - |S^+| - |T^+|) \log 3h - \sum_{\alpha \in R^+} \log \alpha^\vee, \end{aligned}$$

so there are only finitely many possible values for the integers $p^{-1} \alpha^\vee(\lambda + \rho)$ for $\alpha \in S^+$ and $\alpha^\vee(\lambda + \rho)$ for $\alpha \in T^+$ which will bound

$$\log \dim V_\lambda - (|R^+| - |T^+|) \log p = \frac{\text{eff}(R) \log \dim V_\lambda - \log p \text{ord}_p \dim V_\lambda}{\text{eff}(R)}$$

above. For every $\beta^\vee \in \mathbb{Z}R^\vee \cap \text{Span } S \cap \text{Span } T$, $\beta^\vee(\lambda + \rho)$ is both divisible by p and bounded absolutely. Fixing a basis of $\text{Span } S \cap \text{Span } T$ in the root lattice of R^\vee , we conclude that for $p \gg 0$, $\beta^\vee(\lambda + \rho) = 0$ for all $\beta^\vee \in \text{Span } S \cap \text{Span } T$. As S^\vee and T^\vee generate the root lattice, there exists a unique weight ν such that $\alpha^\vee(\nu + \rho) = 0$ for all $\alpha^\vee \in S$ and $\alpha^\vee(\nu + \rho) = \alpha^\vee(\lambda + \rho)$ for all $\alpha^\vee \in T$. Letting $\mu = (\lambda - \nu)/p$, we see that $\alpha^\vee(\mu)$ is an integer for all $\alpha^\vee \in S \cup T$ and therefore for all roots in R^\vee ; it follows that μ is also a weight, and as μ and ν are determined independently of p by the values $p^{-1}\alpha^\vee(\lambda + \rho)$ for $\alpha^\vee \in S^+$ and $\alpha^\vee(\lambda + \rho)$ for $\alpha^\vee \in T$, there are only finitely many possibilities for (μ, ν) . Thus the proposition is true when G is almost simple. Note that in every case where $\mu \neq 0$ arises, $\text{ord}_p \dim V_\lambda = l$, the maximum allowable value.

Now consider a general $G = G_1 \times \cdots \times G_n$. Let

$$B_i = \inf_{(p, \lambda) \in A(G_i)} e \log \dim V_\lambda - (\log p) \text{ord}_p \dim V_\lambda$$

Then letting λ_i denote the i th component of λ ,

$$\begin{aligned} e \log \dim V_\lambda - (\log p) \text{ord}_p \dim V_\lambda &= \sum_i e \log \dim V_{\lambda_i} - (\log p) \text{ord}_p \dim V_{\lambda_i} \\ &\geq \sum B_i, \end{aligned}$$

so $B = \sum_i B_i$ is a lower bound for (4.3.1). On the other hand,

$$e \log \dim V_\lambda - (\log p) \text{ord}_p \dim V_\lambda \leq C$$

implies

$$e \log \dim V_{\lambda_i} - (\log p) \text{ord}_p \dim V_{\lambda_i} \leq C - \sum_{j \neq i} B_j = C - B + B_i$$

so there exist sets $L_{G_i, e, l, C - B + B_i}$ such that each $\lambda_i = p\mu_i + \nu_i$ for some $(\mu_i, \nu_i) \in L_{G_i, e, l, C - B + B_i}$. Letting

$$L_{G, e, l, C} = \left\{ \left(\sum_i \mu_i, \sum_i \nu_i \right) \mid (\mu_i, \nu_i) \in L_{G_i, e, l, C - B + B_i} \right\},$$

we obtain a finite set of pairs (μ, ν) of the desired kind. In particular, for each λ satisfying the condition $\text{ord}_p \dim V_\lambda \leq l$, there can be at most one coordinate i with $\mu_i \neq 0$ because

$$\text{ord}_p \dim V_{p\mu_i + \nu_i} = l$$

for each non-zero μ_i and each $p \gg 0$.

□

THEOREM 4.4: *The zeta-function of a compact semisimple Lie group G determines the Lie algebra of the group up to isomorphism.*

Proof: Let G^1 and G^2 be compact semisimple Lie groups with $\zeta_{G^1}(s) = \zeta_{G^2}(s)$ but with non-isomorphic Lie algebras. By Corollary 1.3, the number of factors of type A_1 in G^1 and G^2 are the same. Let R_1^i, R_2^i, \dots be the irreducible factors of the root system R^i of G^i . If G^1 and G^2 have different Lie algebras, by a suitable renumbering, we may assume that for some $k \geq 0$

$$R_1^1 \geq R_2^1 \geq \dots, R_1^2 \geq R_2^2 \geq \dots, R_i^1 = R_i^2 \quad \forall i \leq k,$$

and there is no factor in $R_{k+1}^2, R_{k+2}^2, \dots$ which equals R_{k+1}^1 or is strictly greater than it. In particular, R^1 has at least $k+1$ irreducible factors (though R^2 might have only k .) Moreover, R_{k+1}^1 is not of type A_1 . Under these conditions, we will find an integer n such that the n th coefficient of $\zeta_{G^1}^*(s)$ is positive and the n th coefficient of $\zeta_{G^2}^*(s)$ is zero, which by Corollary 1.3 implies $\zeta_{G^1}(s) \neq \zeta_{G^2}(s)$. Without loss of generality we may assume that G^1 is adjoint and G^2 is simply connected, so both are products of almost simple factors: $G^i = \prod_j G_j^i$.

In fact, we will prove the following stronger statement: under these conditions, if r is a fixed positive integer, then there exist primes $p_1 \gg \dots \gg p_k$, all congruent to 1 (mod $N_{G^1} = N_{G^2}$) and an allowable weight λ^1 of G^1 such that

$$(4.4.1) \quad \text{ord}_{p_i} \dim V_{\lambda^1} = \text{lev}(R_i^1), \quad 1 \leq i \leq k+1$$

and there is no allowable weight λ^2 of G^2 such that the following conditions hold:

$$(4.4.2) \quad \text{ord}_{p_i} \dim V_{\lambda^2} = \text{lev}(R_i^1), \quad 1 \leq i \leq k+1;$$

$$(4.4.3) \quad \dim V_{\lambda^2} \mid r \dim V_{\lambda^1};$$

$$(4.4.4) \quad \dim V_{\lambda^2} \leq \dim V_{\lambda^1}.$$

Explicitly, we may take $\lambda^1 = \lambda_1^1 + \dots + \lambda_{k+1}^1$, where $\lambda_i^1 = p_i \mu_i^1 + \nu_i^1$, and μ_i^1 and ν_i^1 are defined by Proposition 2.2 as weights of the factor \tilde{G}_i^1 of \tilde{G}^1 . Note that the congruence condition on p_i guarantees that each λ_i^1 lies in the root lattice of \tilde{G}_i^1 and therefore defines a representation of G_i^1 . Note also that we take all primes p_i in the construction to be larger than the bound N in Proposition 4.2 (which

depends only on the root systems of G^1 and G^2 .) We also choose each p_i large enough that for each $j \leq k+1$, $p_i > P_{\mu_j^1, \nu_j^1}^{R_j^1}(2)$; again, this condition depends only on G^1 . By choosing each p_j to be congruent to 2 modulo a sufficiently high power of $p_{j+1} \cdots p_k p_{k+1}$, we can guarantee that $p_i \nmid \dim V_{\lambda_j^1}$ for $i > j$, and therefore, we can guarantee (4.4.1).

Suppose the claim does not hold for $k = 0$. If $R_1^1 = A_1$, we are done. Otherwise, $P_{\mu_1^1, \nu_1^1}^{R_1^1}(x)$ is not constant. As

$$\text{eff}(R_1^1) = \frac{\text{ord}_{x=0} P_{\mu_1^1, \nu_1^1}^{R_1^1}(x)}{\deg P_{\mu_1^1, \nu_1^1}^{R_1^1}(x)},$$

we have

$$(4.4.5) \quad \text{eff}(R_1^1) \log \dim V_{\lambda_1^1} \leq \text{lev}(R_1^1) \log p_1 + C,$$

for some constant C depending only on R^1 . By (4.4.3),

$$\text{eff}(R_1^1) \log \dim V_{\lambda^2} \leq \text{lev}(R_1^1) \log p_1 + \text{eff}(R_1^1) \log r + C.$$

Applying Proposition 4.3 with $e = \text{eff}(R_1^1)$, $l = \text{lev}(R_1^1)$, $G = G^2$, $p = p_1$, and λ^2 an allowable weight of G^2 satisfying (4.4.2)–(4.4.4), we deduce that λ^2 must be of the form $p_1 \mu + \nu$, where μ and ν belong to finite sets independent of p_1 . As $P_{\mu_1^1, \nu_1^1}^{R_1^1}(x)$ is not constant, $\dim V_{\lambda_1^1} \rightarrow \infty$ as $p_1 \rightarrow \infty$, so for $p_1 \gg 0$, $\mu \neq 0$. Again by Proposition 4.3, μ must be a weight of an almost simple factor G_1^2 of G^2 , and R_1^2 must have the same efficiency and level as R_1^1 . (By hypothesis, R_1^2 and R_1^1 cannot be equal, so at this point we must be in the B_n/C_n case, though we do not use this fact explicitly.) Writing $G^2 = G_1^2 \times H^2$, we can decompose $\lambda^2 = \lambda_1^2 + (\lambda^2 - \lambda_1^2)$, where the second summand is a weight of H^2 , and $\lambda_1^2 = p_1 \mu_1^2 + \nu_1^2$, where μ_1^2 and ν_1^2 belong to finite sets independent of p_1 . By (4.3.1) and (4.4.5), the fraction $\frac{\dim V_{\lambda_1^1}}{\dim V_{\lambda_1^2}}$ is bounded above, and by (4.4.3), its denominator is bounded above by r , so the height of the fraction is bounded. By Lemma 2.3, therefore, $\dim V_{\lambda^2}$ is the value at p_1 of a Weyl polynomial which is a scalar multiple of $P_{\mu_1^1, \nu_1^1}^{R_1^1}(x)$. By Proposition 4.2, $\dim V_{\lambda_1^2} > \dim V_{\lambda_1^1}$, contrary to (4.4.4).

The induction step is similar, but we must exercise some additional care. We write $G^1 = G_1^1 \times H^1$. We assume p_1 is large compared to the degree of the representation of $\dim V_{\lambda^1 - \lambda_1^1}$. In other words, the upper bound on

$$\text{eff}(R_1^1) \log \dim V_{\lambda^1} - \text{lev}(R_1^1) \log p_1$$

depends on p_2, p_3, \dots, p_{k+1} , which we regard as constant, but not on p_1 , which we take as large as we need. Applying Proposition 4.3 and Lemma 2.3 as before, there exists a simple factor G_1^2 of G^2 , weights μ_1^2 and ν_1^2 of G_1^2 , and a rational number $c \geq 1$, such that the root system R_1^2 has the same efficiency and level as R_1^1 , $\lambda_1^2 = p\mu_1^2 + \nu_1^2$, and

$$cP_{\mu_1^1, \nu_1^1}^{R_1^1}(x) = P_{\mu_1^2, \nu_1^2}^{R_1^2}(x).$$

Moreover $c = 1$ implies $R_1^1 = R_1^2$. As λ_1^1 and λ_1^2 are allowable, Proposition 4.2 guarantees that $\text{ord}_{p_i} c = 0$ for all i . Writing $G^2 = G_1^2 \times H^2$, $\lambda^2 - \lambda_1^2$ is an allowable dominant weight of H^2 , and for $i \geq 2$,

$$\begin{aligned} \text{ord}_{p_i} \dim V_{\lambda^2 - \lambda_1^2} &= \text{ord}_{p_i} \dim V_{\lambda^2} - \text{ord}_{p_i} \dim V_{\lambda_1^2} \\ &= \text{ord}_{p_i} \dim V_{\lambda^1} - \text{ord}_{p_i} P_{\mu_1^2, \nu_1^2}^{R_1^2}(p_1) \\ &= \text{ord}_{p_i} \dim V_{\lambda^1} - \text{ord}_{p_i} P_{\mu_1^1, \nu_1^1}^{R_1^1}(p_1) \\ &= \text{ord}_{p_i} \dim V_{\lambda^2 - \lambda_1^2} = \text{lev}(R_i^1). \end{aligned}$$

As $\dim V_{\lambda^2 - \lambda_1^2}$ divides $\text{ht}(c)r \dim V_{\lambda^1 - \lambda_1^1}$, and $\text{ht}(c)r$ can be bounded independent of the choice of primes p_i , we can apply the induction hypothesis to conclude that

$$\dim V_{\lambda^2 - \lambda_1^2} > \dim V_{\lambda^1 - \lambda_1^1}.$$

Therefore

$$\begin{aligned} \dim V_{\lambda^2} &= \dim V_{\lambda_1^2} \dim V_{\lambda^2 - \lambda_1^2} > \dim V_{\lambda_1^2} \dim V_{\lambda^1 - \lambda_1^1} \\ &= P_{\mu_1^2, \nu_1^2}^{R_1^2}(p_1) \dim V_{\lambda^1 - \lambda_1^1} \geq P_{\mu_1^1, \nu_1^1}^{R_1^1}(p_1) \dim V_{\lambda^1 - \lambda_1^1} \\ &= \dim V_{\lambda_1^1} \dim V_{\lambda^1 - \lambda_1^1} = \dim V_{\lambda^1}, \end{aligned}$$

contrary to (4.4.4). □

5. The Gassmann Phenomenon

Let $G = \text{SU}(2)^n$ and $Z = Z(G)$. Irreducible representations of G are indexed by ordered n -tuples of non-negative integers: $\lambda = (a_1, \dots, a_n)$. The representation associated to λ is

$$V_\lambda = V_{(a_1, \dots, 0)} \otimes \cdots \otimes V_{(0, \dots, a_n)} = W_{a_1} \boxtimes \cdots \boxtimes W_{a_n},$$

where W_i denotes the i th symmetric power of the natural representation of $\mathrm{SU}(2)$. The restriction of V_λ to $Z \cong \mathbb{F}_2^n$ is isotypic with character χ_λ . In coordinates,

$$\chi_\lambda(a_1, \dots, a_n) = (-1)^{a_1 z_1 + \dots + a_n z_n}.$$

For any character $\chi \in Z^*$, we write

$$X_\chi = \{\lambda \in \mathbb{N}^n \mid \chi_\lambda = \chi\}.$$

Thus, X_χ consists of n -tuples of non-negative integers such that each coordinate has prescribed parity.

Let Z_i ($i = 1, 2$) denote any subgroup of Z , and let $G^i = G/Z_i$. Then,

$$\zeta_{G^i}(s) = \sum_{\chi \in Z_i^\perp} \sum_{\lambda \in X_\chi} (\dim V_\lambda)^{-s}.$$

Now,

$$\dim V_\lambda = \prod_{i=1}^n (a_i + 1),$$

so

$$\sum_{\lambda \in X_\chi} (\dim V_\lambda)^{-s} = (2^{-s} \zeta(s))^O (\zeta(s) - 2^{-s} \zeta(s))^E,$$

where O (resp. E) is the number of coordinates which are constrained to be odd (resp. even) in X_χ . Thus $O + E = n$ and $E - O$ is the trace of $\sigma(\chi)$, where $\sigma: Z^* = \mathbb{F}_2^n \rightarrow \mathrm{GL}_n(\mathbb{C})$ is defined by

$$\sigma(z_1^*, \dots, z_n^*) = \begin{pmatrix} (-1)^{z_1^*} & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & (-1)^{z_n^*} \end{pmatrix}.$$

As long as

$$|\{z^* \in Z_1^\perp \mid \sigma(z^*) = c\}| = |\{z^* \in Z_2^\perp \mid \sigma(z^*) = c\}|$$

for all $c \in \mathbb{Z}$, we have $\zeta_{G^1}(s) = \zeta_{G^2}(s)$.

LEMMA 5.1: *There exists an integer n and two injective homomorphisms*

$$\phi_1, \phi_2: \mathbb{F}_2^3 \rightarrow \mathbb{F}_2^n$$

such that $\sigma(\phi_1(\mathbb{F}_2^3))$ and $\sigma(\phi_2(\mathbb{F}_2^3))$ are not conjugate in $\mathrm{GL}_n(\mathbb{C})$ but for all $c \in \mathbb{Z}$,

$$|\{x \in \mathbb{F}_2^3 \mid \mathrm{tr}(\sigma(\phi_1(x))) = c\}| = |\{x \in \mathbb{F}_2^3 \mid \mathrm{tr}(\sigma(\phi_2(x))) = c\}|.$$

Proof: To guarantee non-conjugacy of images it suffices to arrange that the character of $\sigma \circ \phi_1$ precomposed by any automorphism of \mathbb{F}_2^3 is distinct from the character of $\sigma \circ \phi_2$. To achieve this, we fix an injective function

$$f_1: \mathbb{F}_2^3 \rightarrow 8\mathbb{Z}$$

such that

$$f_1(0) \geq \sum_{x \neq 0} |f(x)|.$$

By Fourier inversion, f_1 is the character of an effective representation ψ_1 . Regarding it as a homomorphism to the diagonal matrices in $\mathrm{GL}_n(\mathbb{C})$, we deduce that there exists ϕ_1 such that $\psi_1 = \sigma \circ \phi_1$. Composing any non-linear automorphism of the pointed set $(\mathbb{F}_2^3, 0)$ with f_1 we obtain a new function f_2 , therefore a new representation ψ_2 , and therefore a new homomorphism ϕ_2 satisfying the conditions of the lemma. \square

Setting $Z_i = \phi_i(\mathbb{F}_2^3)^\perp$ and $G^i = G/Z_i$, the lemma implies $\zeta_{G^1}(s) = \zeta_{G^2}(s)$. Moreover, G^1 and G^2 cannot be isomorphic; indeed an isomorphism ι would give rise to an isomorphism $\tilde{\iota}$ between universal covers, and therefore a diagram

$$\begin{array}{ccc} G = \tilde{G}^1 & \xrightarrow{\tilde{\iota}} & \tilde{G}^2 = G \\ \downarrow & & \downarrow \\ G^1 & \xrightarrow{\iota} & G^2. \end{array}$$

An inner automorphism is trivial on Z , so this means an element of $\mathrm{Out}(G) = S_n$ sends Z_1 to Z_2 , contrary to Lemma 5.1. We conclude:

THEOREM 5.2: *There exist non-isomorphic compact semisimple Lie groups with the same Witten zeta-function.*

REFERENCES

- [1] Bourbaki, N.: Groupes et algèbres de Lie, Chap. 5–6 (1968), Chap. 7–8 (1975), Hermann, Paris.
- [2] Deligne, Pierre; Milne, James S.; Ogus, Arthur; Shih, Kuang-yen: Hodge cycles, motives, and Shimura varieties. Lecture Notes in Mathematics, 900. Springer-Verlag, Berlin-New York, 1982.
- [3] Gassmann, F.: Bemerkungen zu vorstehenden Arbeit von Hurwitz. *Math. Z.* **25** (1926) 124–143.
- [4] Moody, R. V.; Patera, J.; Rand, D. W.: simpLie Version 2.1, Macintosh Software for Representations of Simple Lie Algebras.
- [5] Larsen, M.; Pink, R.: Determining representations from invariant dimensions. *Invent. Math.* **102** (1990), no. 2, 377–398.
- [6] Sunada, T.: Riemannian coverings and isospectral manifolds. *Annals of Math.* **121** (1985) 169–186.
- [7] Witten, E.: On quantum gauge theories in two dimensions. *Comm. Math. Phys.* **141** (1991), no. 1, 153–209.