

# Properties of the Solomon algebra homomorphism

**Christophe Hohlweg**

Institut de Recherche Mathématique Avancée,  
Université Louis Pasteur et CNRS,  
7 rue René Descartes,  
67084 Strasbourg Cedex, France  
E-mail: hohlweg@math.u-strasbg.fr

February 1, 2008

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The Solomon homomorphism</b>	<b>3</b>
2.1	The Solomon descent algebra . . . . .	3
2.2	Coxeter classes . . . . .	4
2.3	The Solomon homomorphism . . . . .	7
2.4	Image of the Solomon homomorphism . . . . .	8
<b>3</b>	<b>Character formula and enumeration</b>	<b>9</b>
3.1	The Solomon homomorphism is an isometry . . . . .	9
3.2	Idempotents in the group algebra . . . . .	10
<b>4</b>	<b>Conjectures and results</b>	<b>12</b>
4.1	The symmetry conjecture . . . . .	12
4.2	Orthogonal idempotents and character formulas . . . . .	15
4.3	The double coset conjecture . . . . .	17
4.4	Results . . . . .	19
4.5	The case of a single generator . . . . .	20
4.6	Proof of Theorem 4.2 . . . . .	24

## 1 Introduction

Let  $(W, S)$  be a finite Coxeter system. Let  $I \subseteq S$  and  $W_I$  be the *parabolic subgroup* of  $W$  generated by  $I$ . We denote  $X_I$  the cross section of  $W/W_I$  consisting of the unique coset representatives of minimal length. Its elements are called *the minimal coset representatives* (see [3], [12], 5.12).

The *Solomon descent algebra*  $\Sigma W$  (first defined in [19]) is the subalgebra of  $\mathbb{Q}W$  generated by  $\{x_I\}_{I \subset S}$ , where  $x_I = \sum_{w \in X_I} w$ .

Solomon has also shown that the linear function  $\Phi$  which maps  $x_I$  to  $1_{W_I}^W$ , the induced character of the trivial representation on  $W_I$  to  $W$ , is an algebra homomorphism called *the Solomon homomorphism*.

For the symmetric groups (which are Coxeter groups of type  $A$ ), properties of  $\Phi$  are related to enumerative results: certain joint statistics on the symmetric group may be enumerated by the scalar products of appropriate characters.

For example, Gessel has shown ([9]) that the number of element in  $X_I \cap X_J^{-1}$  is the scalar product of the characters  $1_{W_I}^W$  and  $1_{W_J}^W$ . We show that this result is equivalent to the following:  $\Phi$  is an isometry for a particular scalar product on  $\mathbb{Q}S_n$  (defined by  $\langle w, v \rangle = 1$  if  $w = v^{-1}$ ,  $= 0$  otherwise). In Section 3, we extend this result of Gessel on symmetric groups to all finite Coxeter groups.

As another example, there is a set of orthogonal idempotents  $E_\lambda \in \Sigma S_n$ , indexed by the partitions  $\lambda$  of  $n$ , which decompose  $\mathbb{Q}S_n$  (see [6]). Gessel and Reutenauer have shown ([10]) that the scalar product of  $1_{W_I}^W$  and  $\chi_\lambda$  - the character of the left action of  $S_n$  on  $\mathbb{Q}S_n E_\lambda$  - is the number of elements in  $X_I$  of cycle type  $\lambda$  (equivalently, in the conjugation class corresponding to  $\lambda$ ). By this way, Jöllenbeck and Reutenauer in [13] have shown that  $\Phi$  has the following symmetry property:

$$(\star) \quad \Phi(x)(y) = \Phi(y)(x)$$

for all  $x, y \in \Sigma S_n$  ( $\Phi(x)$  is linearly extended to  $\mathbb{Q}S_n$ ). In other words

$$\sum_{w \in X_I} 1_{W_J}^W(w) = \sum_{g \in X_J} 1_{W_I}^W(g), \text{ for all } I, J \subset S.$$

Conversely, the symmetry property implies the previous enumerative result.

In our attempt to extend to all finite Coxeter groups the symmetry property  $(\star)$ , we use the work of Bergeron, Bergeron, Howlett and Taylor ([1]): they have found for each finite Coxeter group a set of orthogonal idempotents  $E_\lambda$ , indexed by Coxeter classes which decompose  $\mathbb{Q}W$  (a Coxeter class is an equivalence class of subsets of  $S$  conjugated under  $W$ ). In Section 4, we prove the scalar product of  $1_{W_I}^W$  and  $\chi_\lambda$  - the character of the right action of  $W$  on  $E_\lambda \mathbb{Q}W$  - is equal to the number of elements in  $X_I$  of Coxeter type  $\lambda$  if and only if  $\Phi$  has the symmetry property  $(\star)$  (the Coxeter type may be viewed as the cycle type of a permutation). Thus, again, properties of  $\Phi$  are strongly related to enumerative results.

We show that  $\Phi$  has property  $(\star)$  for all finite Coxeter groups whose irreducible components are of type  $A_n, E_6, E_7, E_8, F_4, H_3, H_4$  or  $I_2(m)$ . Whether this is true for all finite Coxeter groups is an open conjecture. Moreover, we state a stronger conjecture (obtained jointly with M. Schöcker) that depends on two subsets  $I, J$  of  $S$ ; we prove this new conjecture in the following cases:  $I = \emptyset, I = S$  and  $I$  is a singleton. In particular, the new conjecture is true for dihedral groups. The general case remains open.

As a byproduct of our techniques, in Section 2, we give a quick proof of the following result: For each conjugacy class  $C$ , if  $I, J$  are conjugated under  $W$ , then  $|C \cap X_I| = |C \cap X_J|$ . This result has been obtained independently for all Coxeter groups by Fleischmann in [4]. We introduce the image of  $\Phi$  in the algebra of class functions and we describe it using some results of Bergeron, Bergeron, Howlett and Taylor ([1]). Moreover, we show that  $\Phi$  is surjective if and only if  $W$  is a direct product of symmetric groups.

## 2 The Solomon homomorphism

### 2.1 The Solomon descent algebra

Let  $w \in W$ . We denote by  $\ell(w)$  the length of  $w$  as a word in the elements of  $S$ . The *ascent set* of  $w$  is the set

$$S(w) = \{s \in S \mid \ell(ws) > \ell(w)\}.$$

The *descent set* of  $w$  is the set

$$D(w) = \{s \in S \mid \ell(ws) < \ell(w)\}.$$

Observe that  $D(w) \amalg S(w) = S$ . Let  $I \subset S$ , then the set of minimal coset representatives  $X_I$  can also be viewed as

$$\begin{aligned} X_I &= \{x \in W \mid \ell(xs) > \ell(x), \forall s \in I\} \\ &= \{w \in W \mid S(w) \supset I\} \\ &= \{w \in W \mid D(w) \subset S - I\}. \end{aligned}$$

**Parabolic components:** Any  $w \in W$  has a unique factorization  $w = w^I w_I$  such that  $w^I \in X_I$  and  $w_I \in W_I$ . Moreover,  $\ell(w) = \ell(w^I) + \ell(w_I)$  ([12], 5.12). We call the couple  $(w^I, w_I)$  the *parabolic components* of  $w$ .

Let  $I, J \subseteq S$  and  $X_I^{-1} = \{x^{-1} \mid x \in X_I\}$ . Then the set

$$X_{IJ} = X_I^{-1} \cap X_J$$

is a cross section of the double cosets  $W_I \backslash W / W_J$ , consisting again of the unique representatives of minimal length ([19], Lemma 1).

Following [1], Section 2, for  $I, K \subset S$ , we denote  $X_{I \cap K}^I = W_I \cap X_K$ . Then  $X_{I \cap K}^I$  is a cross section of  $W_I / W_{I \cap K}$  which is constituted by the minimal coset representatives of  $W_{K \cap I}$  in  $W_I$ . Finally, we have the following properties:

1. for  $K \subset I \subset S$ ,  $X_I X_K^I = X_K$ ;
2. for  $I, J \subset S$ ,  $X_J$  is the disjoint union of  $X_{I \cap b J b^{-1}}^I b$  for all  $b \in X_{IJ}$  and  $W_I \cap b W_J b^{-1} = W_{I \cap b J b^{-1}}$ . In particular, for all  $w \in W$ , there is a unique  $b \in X_{IJ}$  such that  $w^J \in X_{I \cap b J b^{-1}}^I b$ ;

3. for  $K, L \subset S$  and  $b \in X_{KL}$  such that  $bLb^{-1} = K$ , we have  $X_K b = X_L$ .

**Double parabolic components:** We have an analogue decomposition in parabolic components for double cosets: any  $w \in W$  has a unique factorization  $w = abc$  such that  $b \in X_{IJ}$ ,  $a \in X_{I \cap bJb^{-1}}^I$  and  $c \in W_J$ . Moreover,  $\ell(w) = \ell(a) + \ell(b) + \ell(c)$  ([19], Lemma 1). We call the triplet  $(a, b, c)$  the *double parabolic components* of  $w$ .

For  $I \subset S$ , we consider the element  $x_I = \sum_{w \in X_I} w$  in  $\mathbb{Q}W$ . Solomon has shown in [19] that the vector subspace  $\Sigma W$  of  $\mathbb{Q}W$  generated by these elements is a subalgebra. Moreover  $\{x_I\}_{I \subset S}$  is a basis that satisfies

$$x_I x_J = \sum_{K \subset S} a_{IJK} x_K,$$

where  $a_{IJK} = |\{w \in X_{IJ} \mid w^{-1}W_I w \cap W_J = W_K\}|$ .

Following the case of  $A_n$  and  $B_n$  (symmetric and hyperoctahedral groups, see [16], [2]), we call  $\Sigma W$  the *Solomon descent algebra*. Note that in these two case, the basis  $\tilde{x}_I = x_{S-I} = \sum_{D(w) \subset I} w$  is more commonly used. Note also that  $\dim \Sigma W = 2^{|S|}$ .

We use a basis of  $\Sigma W$  which is useful to enumerate the elements of  $W$  which have a given descent or ascent set. Let  $I, J \subset S$ . The set of elements of  $W$ , which have as ascent set  $I$ , is  $Y_I = \{w \in W \mid S(w) = I\}$ . We consider the following element in  $\mathbb{Q}W$ :

$$y_I = \sum_{w \in Y_I} w.$$

Then

$$x_J = \sum_{I \supset J} y_I$$

and, by inclusion-exclusion,

$$y_I = \sum_{J \supset I} (-1)^{|I-J|} x_J.$$

Therefore  $\{y_I\}_{I \subset S}$  is a basis of  $\Sigma W$ .

## 2.2 Coxeter classes

Let  $I, J \subset S$ . We say that  $I$  and  $J$  are *W-conjugate* if there is  $w \in W$  such that  $wIw^{-1} = J$ . In this case, we write  $I \sim_W J$ .

The relation  $\sim_W$ , on the set of subsets of  $S$ , is an equivalence relation. For this relation, an equivalence class is called a *Coxeter class*. The set of Coxeter classes of  $W$  is denoted  $\Lambda(W)$ . For  $I \subset S$ , write  $\lambda(I)$  its Coxeter class.

An element  $c \in W$  with  $c = s_1 \dots s_n$  ( $s_i \in S$ ) is a *Coxeter element* if the  $s_i$  are all distinct and if  $S = \{s_1, \dots, s_n\}$  ( $c$  is the product of all elements of

$S$ , without repetition). We write  $c_I$  for a Coxeter element of  $W_I$ . It is well-known that the following statements give three equivalent characterisations of the  $W$ -conjugation ( See [1], Proposition 4.4 and [8], Proposition 3.1.15):

1.  $I \sim_W J$ ,
2. the subgroups  $W_I$  and  $W_J$  are conjugated,
3. the Coxeter elements  $c_I$  of  $W_I$  and  $c_J$  of  $W_J$  are conjugated.

Denote by  $C(w)$  the conjugacy class of an element  $w \in W$  and  $Cl(W)$  the set of conjugacy classes of  $W$ . A *cuspidal class* is a conjugacy class  $C$  such that  $C \cap W_I = \emptyset$  for all  $I \subsetneq S$ . A  *$I$ -cuspidal class* is a cuspidal class of the parabolic subgroup  $W_I$ . As example, the Coxeter elements are all conjugated and their conjugacy class is cuspidal (see [8], Proposition 3.1.6).

An element  $w \in W$  is an  *$I$ -element* ( $I \subset S$ ) if  $C(w) \cap W_I$  is a  $I$ -cuspidal class. If  $J \in \lambda(I)$ , then  $w$  is also a  $J$ -element (since  $W_J$  is conjugated to  $W_I$ ,  $C(w) \cap W_I$  is conjugated to  $C(w) \cap W_J$ . As subgroups of  $W_J$  are conjugated to subgroups of  $W_I$ ,  $C(w) \cap W_J$  is a  $J$ -cuspidal class).

More generally, for any  $\lambda \in \Lambda$ , we say that  $w \in W$  is a  *$\lambda$ -element* (or is of *Coxeter type  $\lambda$* ) if  $w$  is a  $I$ -element for some  $I \in \lambda$ . Denote by  $C(\lambda)$  the set of  $\lambda$ -elements. We call it *the  $\lambda$ -set*.

It is well-known that conjugacy classes and Coxeter classes coincide in the symmetric group. The subclass of finite Coxeter groups that have this property is given by the following proposition. We will see that this is also the subclass of finite Coxeter groups for which the Solomon homomorphism is surjective.

**Proposition 2.1.** *Let  $(W, S)$  be a finite Coxeter system then the following propositions are equivalent:*

1.  $W$  is a direct product of symmetric groups;
2.  $|\Lambda(W)| = |Cl(W)|$ ;
3. Coxeter classes coincide with conjugacy classes.

We need two lemmas to prove Proposition 2.1. The first one implies that there are more conjugacy classes than Coxeter classes.

**Lemma 2.2.** *Let  $\lambda \in \Lambda$ , then  $C(\lambda) = \coprod C$ , where the disjoint union is taken over all  $C \in Cl(W)$  such that  $C \subset C(\lambda)$ . Moreover,  $W$  is the disjoint union of its  $\lambda$ -sets, for  $\lambda \in \Lambda$ .*

*Proof.* If  $w \in C(\lambda)$ , then  $C(w) \cap W_I$  is a cuspidal class of  $W_I$  (for some  $I \in \lambda$ ). Thus all  $g \in C(w)$  are  $\lambda$ -elements. Therefore  $C(w) \subset C(\lambda)$ . Finally, for all  $w \in C(\lambda)$ , there is a conjugacy class  $C$  such that  $w \in C \subset C(\lambda)$  and the lemma is proved.  $\square$

It is well-known that the conjugacy classes of  $W$  are the direct product of conjugacy classes of its irreducible components (as  $W$  is isomorphic to the direct product of them). This second lemma gives an analogue for Coxeter classes.

**Lemma 2.3.** *Let  $(W, S)$  be a finite Coxeter system. For  $i = 1, \dots, n$ , let  $(W_i, K_i)$  be the irreducible components of  $(W, S)$ . For  $\lambda \in \Lambda(W)$  and  $I \in \lambda$ , denote  $I_p = I \cap K_p$  and  $\pi_p(\lambda) = \lambda(I_p) \in \Lambda(W_p)$ . Then the following mapping*

$$\begin{aligned} \pi : \Lambda(W) &\longrightarrow \Lambda(W_1) \times \dots \times \Lambda(W_n) \\ \lambda &\longmapsto (\pi_1(\lambda), \dots, \pi_n(\lambda)) \end{aligned}$$

*is a bijection. Moreover, for any  $\lambda \in \Lambda(W)$*

$$|C(\lambda)| = \prod_{p=1}^n |C(\pi_p(\lambda))|.$$

*Proof.* It is readily seen that one just has to prove the lemma in the case  $n = 2$  (and conclude by induction on  $n$ ). As  $W_p$ ,  $p = 1, 2$ , are the irreducible components of  $W$ , they can be viewed as normal parabolic subgroup of  $W$  such that  $W = W_1 W_2$ . Therefore, for any  $w \in W$ ,  $w = w_1 w_2 = w_2 w_1$  with  $w_p \in W_p$ . Observe that  $w^{K_1} = w_{K_2} = w_2$  and  $w_{K_1} = w^{K_2} = w_1$ . For  $I \subset S$  and  $p = 1, 2$ ,  $W_p \cap I \subset W_p \cap S = K_p$ . Thus  $I_p = I \cap K_p = I \cap W_p$  and  $I$  is the disjoint union of  $I_1$  and  $I_2$ . Finally, since each element of  $W_1$  commutes with each element of  $W_2$ ,  $w I_p w^{-1} = w_p I_p w_p^{-1}$ .

One first shows that  $\pi$  is well defined. Let  $I, J \in \lambda$ , there is  $w \in W$  such that  $w I w^{-1} = J$ . As  $W_p$  is a normal subgroup, for  $p = 1, 2$ ,  $w I_p w^{-1} = J_p$ . Moreover, since  $w I_p w^{-1} = w_p I_p w_p^{-1}$ , one has  $w_p \in W_p$  such that  $w_p I_p w_p^{-1} = J_p$ , hence  $\lambda(I_p) = \lambda(J_p) \in \Lambda(W_p)$ . Then  $\pi_p$ , and therefore  $\pi$ , are well defined.

Let  $(\lambda_1, \lambda_2) \in \Lambda(W_1) \times \Lambda(W_2)$  and  $I_p \in \lambda_p$ . Then  $I = I_1 \cup I_2 \subset S$  and  $\lambda(I) \in \Lambda(W)$ . It is readily seen that  $\pi_p(\lambda(I)) = \lambda_p$  hence  $\pi(\lambda(I)) = (\lambda_1, \lambda_2)$ . Therefore,  $\pi$  is surjective.

Let  $\lambda, \mu \in \Lambda(W)$  such that  $\pi(\lambda) = \pi(\mu)$ . So  $\pi_p(\lambda) = \pi_p(\mu)$ . Let  $I \in \lambda$  and  $J \in \mu$ , then  $I_p$  and  $J_p$  are  $W_p$ -conjugate by an element  $w_p \in W_p$ . Write  $w = w_1 w_2 = w_2 w_1$ , then one has  $w I w^{-1} = J$ . Therefore,  $I \sim_W J$  so  $\lambda = \mu$ . One concludes that  $\pi$  is injective, hence bijective.

Let  $\lambda \in \Lambda(W)$  and  $I \in \lambda$ . If  $w \in C(\lambda)$ , then  $C(w) \cap W_I$  is a  $I$ -cuspidal class. It is well-known that  $C^p(w_p)$  is a conjugacy class of  $W_p$  and  $C^p(w_p) \cap W_{I_p}$  is an  $I_p$ -cuspidal class (see [8], exercise 3.10). Therefore, the mapping

$$\begin{aligned} \eta : C(\lambda) &\longrightarrow C(\pi_1(\lambda)) \times C(\pi_2(\lambda)) \\ w &\longmapsto (w_1, w_2) \end{aligned}$$

is well defined and injective since the parabolic components  $(w_1, w_2)$  of  $w$  are uniquely determined. Moreover,  $\eta$  is easily seen to be surjective, hence the lemma is proved.  $\square$

*Proof of Proposition 2.1.* One may assume that  $W$  is irreducible, by Lemma 2.3.

1.  $|\Lambda(W)| \leq |Cl(W)|$ , and  $|\Lambda(W)| < |Cl(W)|$  if  $W$  contains a parabolic subgroup  $W_I$  that has at least two cuspidal classes. Indeed, denote by  $C_1^I$  and  $C_2^I$  two distinct cuspidal classes of  $W_I$ . Let  $w_1 \in C_1^I$  and  $w_2 \in C_2^I$ .

Recall a useful result about cuspidal classes: if  $I \subset S$ , if  $C^I \in Cl(W_I)$  is a  $I$ -cuspidal class and if  $w \in C^I$ , then  $C^I = C(w) \cap W_I$  (see [8], Theorem 3.2.11). This implies that the conjugacy classes  $C(w_1)$  and  $C(w_2)$  of  $W$  are distinct and contained in  $C(\lambda(I))$ . Therefore  $|\Lambda(W)| < |Cl(W)|$ .

2. The Coxeter group of type  $I_2(m)$ ,  $m \geq 4$  has at least two cuspidal classes. Indeed, let  $S = \{s, t\}$ , then the conjugacy class  $C_1$  of the Coxeter element  $st$  is cuspidal. Moreover,  $stst$  is an element of minimal length in its conjugacy class  $C_2$ . As  $\ell(st) = 2 < 4 = \ell(stst)$ ,  $C_2$  is distinct of  $C_1$ . Observe that  $C_2$  is cuspidal, hence  $C_1, C_2$  are two distinct cuspidal classes.
3. The Coxeter group of type  $D_4$  has at least two cuspidal classes. Indeed,  $S = \{1, 2, 3, 4\}$  with  $m(1, 2) = m(2, 3) = m(2, 4) = 3$ . The conjugacy class  $C_1$  of the Coxeter element 1234 is cuspidal. Moreover 12342 is an element of minimal length in its conjugacy class  $C_2$  thus  $C_2$  is distinct of  $C_1$ . Observe that  $C_2$  is cuspidal.
4. The only irreducible Coxeter group that has no parabolic subgroup of type  $D_4$  and  $I_2(m)$  is the symmetric group (type  $A$ , see classification of finite Coxeter groups in [12]).

□

### 2.3 The Solomon homomorphism

Denote by  $\mathbb{Q}_{class}W$  the set of class functions on  $W$ . In his article [19], Solomon shows that the following linear homomorphism

$$\begin{aligned} \Phi : \Sigma W &\longrightarrow \mathbb{Q}_{class}W \\ x_I &\longmapsto 1_{W_I}^W \end{aligned}$$

is an algebra homomorphism. Moreover he shows that

$$\ker \Phi = Rad \Sigma W$$

is generated by the elements  $x_I - x_J$  with  $I \sim_W J$ .

**Remark:** Note that the theorem implies that if  $I \sim_W J$ , then  $1_{W_I}^W = 1_{W_J}^W$ ; this may also be deduced from general results on induction of characters, since  $W_I, W_J$  are conjugated in  $W$ .

The following proposition has been obtained independently for all Coxeter groups by Fleischmann ([4]).

**Proposition 2.4.** *Let  $I, J \subset S$  such that  $I \sim_W J$  then*

1. *for any conjugacy class  $C$ ,  $|X_I \cap C| = |X_J \cap C|$ ;*
2. *for any Coxeter class  $\lambda$ ,  $|X_I \cap C(\lambda)| = |X_J \cap C(\lambda)|$ .*

**Lemma 2.5.** *Extend linearly class functions to  $\mathbb{Q}W$ . Let  $x \in \ker \Phi$ , then*

$$f(x) = 0, \quad \forall f \in \mathbb{Q}_{class}W.$$

*Proof.* Let  $f \in \mathbb{Q}_{class}W$ . As  $f$  is a linear combination of characters, one can assume that  $f$  is a character, associated to some representation  $\mu$  of  $W$ , of degree  $n$ . As  $x \in \ker \Phi = \text{Rad } \Sigma W$ ,  $x$  is a nilpotent element of  $\Sigma W$ . Thus, the characteristic polynomial of  $\mu x$  is  $X^n$ . As  $f(x) = \text{tr}(\mu x)$  is the coefficient of  $\pm X^{n-1}$  in this polynomial, one has  $f(x) = 0$ .  $\square$

*Proof of Proposition 2.4.* Let  $f_C$  be the characteristic function of the conjugacy class  $C$ . It is well-known that  $f_C \in \mathbb{Q}_{class}W$ . Then  $f_C(x_I - x_J) = 0$ , by Lemma 2.5. Hence

$$|X_I \cap C| = f_C(x_I) = f_C(x_J) = |X_J \cap C|.$$

The second statement is immediate from 1 and Lemma 2.2.  $\square$

## 2.4 Image of the Solomon homomorphism

Let  $\lambda$  be a Coxeter class. As  $1_{W_I}^W = 1_{W_J}^W$  for all  $I, J \in \lambda$ , we denote  $\varphi_\lambda$  instead of  $1_{W_I}^W$  for all  $I \in \lambda$ . Following [1] (Theorem 4.5), we consider the matrix

$$A = (\varphi_\lambda(c_\mu))_{\lambda, \mu \in \Lambda},$$

where  $c_\mu$  is an arbitrary  $\mu$ -element (for example a Coxeter  $\mu$ -element). In [1], the authors have shown that:

1.  $A$  is an invertible matrix;
2.  $\dim \text{Rad } \Sigma = \dim \ker \Phi = 2^{|S|} - |\Lambda(W)|$ .

They have also introduced the characteristic class function of  $\lambda$ -elements

$$\xi_\lambda = \begin{cases} 1 & \text{if } w \in C(\lambda); \\ 0 & \text{otherwise.} \end{cases}$$

Following [1], we denote by  $\nu_{\lambda\mu}$  the coefficients of  $A^{-1}$ . Observe that if  $w \in C(\mu)$ , then  $\xi_\lambda(w) = \xi_\lambda(c_\mu)$ , for a Coxeter  $\mu$ -element  $c_\mu$ . Then we have

$$\xi_\lambda = \sum_{\beta \in \Lambda} \nu_{\lambda\beta} \varphi_\beta$$

since  $\sum_{\beta \in \Lambda} \nu_{\lambda\beta} \varphi_\beta(c_\mu) = \delta_{\lambda,\mu}$ .

We denote  $\Gamma W$  the image of  $\Phi$  in  $\mathbb{Q}_{class}W$  (spanned by  $\{1_{W_I}^W\}_{I \in \Lambda}$ ). As  $\Sigma W$  is a  $\mathbb{Q}$ -algebra,  $\Gamma W$  is also a  $\mathbb{Q}$ -algebra. The results of Bergeron, Howlett and Taylor can be rewritten as follow:  $\{\varphi_\lambda\}_{\lambda \in \Lambda(W)}$ ,  $(\xi_\lambda)_{\lambda \in \Lambda}$  are two basis of  $\Gamma W$  and  $\dim \Gamma W = |\Lambda(W)|$ . Therefore, the result of Bergeron, Bergeron, Howlett and Taylor about the basis  $(\xi_\lambda)_{\lambda \in \Lambda}$  implies the following proposition, which states that,  $\Gamma W$  is the set of functions on  $W$  stable on Coxeter classes.



**Proposition 2.6.** *A class function  $f$  is in  $\Gamma W$  if and only if  $f$  is a  $\mathbb{Q}$  valued function and is stable on  $\lambda$ -sets, for all  $\lambda \in \Lambda$ .*

*Proof.* By Lemma 2.2,  $\xi_\lambda = \sum_{C \subset C(\lambda)} f_C$  where  $f_C$  is the characteristic function of the conjugacy class  $C$ . If  $f$  is stable on  $\lambda$ -set, for any  $\lambda$ , the coefficients of  $f$  in the basis  $\{f_C\}_{C \in Cl(W)}$  are equal if the conjugacy classes are in the same  $\lambda$ -set. Therefore  $f$  may be written in the basis  $\{\xi_\lambda\}_{\lambda \in \Lambda W}$  with coefficients in  $\mathbb{Q}$  if  $f$  is a  $\mathbb{Q}$  valued function. On the other side, if  $f \in \Gamma W$ , it is readily seen that  $f$  is  $\mathbb{Q}$ -valued and is stable on all  $\lambda$ -sets.  $\square$

We conclude the study of the image of  $\Phi$  with the following corollary.

**Corollary 2.7 (of Proposition 2.1).** *Let  $(W, S)$  be a finite Coxeter system. Then  $\Phi$  is surjective if and only if  $W$  is a direct product of symmetric groups.*

### 3 Character formula and enumeration

#### 3.1 The Solomon homomorphism is an isometry

It is well known that irreducible characters of  $W$  are  $\mathbb{R}$ -valued. More precisely, there is a finite field extension  $\mathbb{K}$  of  $\mathbb{Q}$  such that the characters of  $W$  can be realized over  $\mathbb{K}$  (see [8], Chapter 5). For example, for type  $A_n$ ,  $B_n$  and  $D_n$ ,  $\mathbb{K} = \mathbb{Q}$ . As a well-known consequence, each element of  $W$  is conjugated to its inverse and the classical scalar product on  $\mathbb{Q}_{class}W$  is given by

$$\langle f, g \rangle_W = \frac{1}{|W|} \sum_{w \in W} f(w)g(w).$$

We consider the following scalar product on  $\mathbb{Q}W$ . For  $w, g \in W$  let

$$\langle w, g \rangle = \begin{cases} 1 & \text{if } w = g^{-1}; \\ 0 & \text{otherwise.} \end{cases}$$

This scalar product is of great interest concerning the problems of enumeration: Let  $M \subset W$  and  $z_M = \sum_{w \in M} w \in \mathbb{Q}W$ . Then the number of elements of  $W$  which are in  $N$  and whose inverse is in  $N \subset W$  is equal to  $\langle z_M, z_N \rangle$ .

For example, let  $I, J \subset S$ ; then

1.  $\langle x_I, x_J \rangle = |X_{IJ}|$  is the number of  $w$  in  $W$  such that the ascent set of  $w$  contains  $I$  and that the ascent set of  $w^{-1}$  contains  $J$ ;
2.  $\langle y_I, y_J \rangle$  is the number of  $w$  in  $W$  such that the ascent set of  $w$  equals  $I$  and that the ascent set of  $w^{-1}$  equals  $J$ , that is, the number of  $w$  in  $W$  such that the descent set of  $w$  equals  $S - I$  and that the descent set of  $w^{-1}$  equals  $S - J$ .

The following theorem gives an isometry property of  $\Phi$  for this scalar product and the classical scalar product on  $\mathbb{Q}_{class}W$ . This is a generalization of Gessel's result for symmetric groups (see [9]).

**Theorem 3.1.** *Let  $x, y \in \Sigma W$ , then  $\langle \Phi(x), \Phi(y) \rangle_W = \langle x, y \rangle$ .*

*Proof.* As  $\{x_I\}_{I \subset S}$  is a basis of  $\Sigma W$ , one just has to prove the theorem for these elements. Let  $I, J \subset S$ , let  $\chi_I$  (resp.  $\chi_J$ ) be a character on  $W_I$  (resp.  $W_J$ ). Denote  $\chi_I^W$  (resp.  $\chi_J^W$ ) be the character induced by  $\chi_I$  (resp.  $\chi_J$ ) on  $W$ . As  $X_{IJ}$  is a set of representatives of double coset  $W_I w W_J$ , Mackey's Theorem (see [14], Theorem 3.3, p.86) implies

$$\chi_I^W \chi_J^W = \sum_{x \in X_{IJ}} \beta_x^W,$$

where  $\beta_x^W$  denote the induced character on  $W$  of the character  $\beta_x$  on the subgroup  $W_I \cap x W_J x^{-1} = W_{I \cap x J x^{-1}}$  defined by

$$\beta_x(w) = \chi_I(w) \chi_J(x^{-1} w x),$$

for  $w \in W_{I \cap x J x^{-1}}$ . Applying this to  $\chi_I = 1_{W_I}$  and  $\chi_J = 1_{W_J}$ , one has

$$1_{W_I}^W 1_{W_J}^W = \sum_{x \in X_{IJ}} 1_{W_{I \cap x J x^{-1}}}^W.$$

Therefore

$$\begin{aligned} \langle \Phi(x_I), \Phi(x_J) \rangle_W &= \langle 1_{W_I}^W 1_{W_J}^W, 1 \rangle_W \\ &= \sum_{x \in X_{IJ}} \langle 1_{W_{I \cap x J x^{-1}}}^W, 1 \rangle_W \\ &= \sum_{x \in X_{IJ}} \langle 1, 1 \rangle_{W_{I \cap x J x^{-1}}}, \text{ by Frobenius reciprocity} \\ &= |X_{IJ}| \\ &= \langle x_I, x_J \rangle. \end{aligned}$$

□

**Corollary 3.2 (of proof).** *Let  $I, J \subset S$ , then*

$$|X_{IJ}| = \langle 1_{W_I}^W, 1_{W_J}^W \rangle_W.$$

*Equivalently, the number of  $w$  in  $W$  such that the ascent set of  $w$  contains  $I$  and that the ascent set of  $w^{-1}$  contains  $J$  is equal to  $\langle 1_{W_I}^W, 1_{W_J}^W \rangle_W$ .*

### 3.2 Idempotents in the group algebra

Let  $e = \sum_{w \in W} e_w w \in \mathbb{Q}W$  be an idempotent. Then  $e\mathbb{Q}W$  (resp.  $\mathbb{Q}We$ ) is a right (resp. left)  $W$ -module by right (resp. left) multiplication. Denote by  $\chi_R$  (resp.  $\chi_L$ ) the character of the representation  $e\mathbb{Q}W$  (resp.  $\mathbb{Q}We$ ).

**Proposition 3.3.** *The right  $W$ -module  $e\mathbb{Q}W$  and the left  $W$ -module  $\mathbb{Q}We$  have the same character  $\chi_e(w) = \chi_L(w) = \chi_R(w)$ . In particular  $\dim \mathbb{Q}We = \dim e\mathbb{Q}W$ .*

The proposition is an immediate corollary of the following well-known lemma. For a proof, see [16], Lemma 8.4, where it is given for any finite group and for the classical scalar product on  $\mathbb{Q}W$  (for which  $W$  is an orthonormal basis).

**Lemma 3.4.** *For any idempotent  $e \in \mathbb{Q}W$  and any  $w \in W$  we have*

$$\chi_L(w) = \chi_R(w) = \sum_{x \in W} \langle x^{-1}ex, w \rangle.$$

The next well-known lemma is a important tool for the following.

**Lemma 3.5.** *For any idempotents  $e$  and  $e'$  in  $\mathbb{Q}W$  we have:*

$$\langle \chi_e, \chi_{e'} \rangle_W = \chi_{e'}(e).$$

*Proof.*

$$\begin{aligned} \langle \chi_e, \chi_{e'} \rangle_W &= \frac{1}{|W|} \sum_{w \in W} \chi_e(w) \chi_{e'}(w) \\ &= \frac{1}{|W|} \sum_{w \in W} \sum_{x \in W} \langle x^{-1}ex, w \rangle \chi_{e'}(w), \quad \text{by Lemma 3.4} \\ &= \frac{1}{|W|} \sum_{x \in W} \sum_{w \in W} e_{xw^{-1}x^{-1}} \chi_{e'}(w) \\ &= \frac{1}{|W|} \sum_{x \in W} \sum_{w \in W} e_{xw^{-1}x^{-1}} \chi_{e'}(w^{-1}), \quad \text{since } w \sim w^{-1} \\ &= \frac{1}{|W|} \sum_{x \in W} \sum_{w \in W} e_{xw^{-1}x^{-1}} \chi_{e'}(xw^{-1}x^{-1}) \\ &= \frac{1}{|W|} \sum_{x \in W} \sum_{g \in W} e_g \chi_{e'}(g) \\ &= \chi_{e'}(e). \end{aligned}$$

□

For  $I \subset S$ ,  $e_I = \frac{1}{|W_I|} \sum_{w \in W_I} w$  is an idempotent of  $\mathbb{Q}W$ . Observe that  $\chi_{e_I} = 1_{W_I}^W$ . Let

$$\text{Fix}_{W_I}(e\mathbb{Q}W) = \{x \in e\mathbb{Q}W \mid xu = x, \forall u \in W_I\}.$$

Observe that  $e\mathbb{Q}W$  is a right  $W_I$ -module and that  $W_I$  acts trivially on  $e\mathbb{Q}We_I$ . Therefore

$$\text{Fix}_{W_I}(e\mathbb{Q}W) = e\mathbb{Q}We_I$$

and

$$\dim \text{Fix}_{W_I}(e\mathbb{Q}W) = \langle \chi_e, 1_{W_I}^W \rangle_W,$$

see [5], p.15 – 16. Hence

$$\dim e\mathbb{Q}We_I = \langle \chi_e, 1_{W_I}^W \rangle_W = \chi_e(e_I).$$

In [18], the author has looked at the following idempotents: Let  $I \subset S$  and  $\epsilon$  be the sign character, denote

$$\tilde{e}_I = \frac{1}{|W_I|} \sum_{w \in W_I} \epsilon(w)w.$$

Solomon has shown in [18] that the character  $\tilde{\chi}_I$  of the action of  $W$  on  $\tilde{e}_{S-I}e_I\mathbb{Q}W$  is  $\Phi(y_I)$ . We deduce, from Theorem 3.1, the following corollary which generalize to finite Coxeter groups a Gessel's result for symmetric groups ([9]) and a Poirier's result for hyperoctahedral groups ([15]).

**Corollary 3.6.** *Let  $I, J \subset S$ , then*

1. *the number of  $w$  in  $X_J$  such that the descent set of  $w^{-1}$  equals  $S - I$  is  $\langle \tilde{\chi}_I, 1_{W_J}^W \rangle_W = \dim \text{Fix}_{W_J}(\tilde{e}_{S-I}e_I\mathbb{Q}W)$ ;*
2. *the number of  $w$  in  $W$  such that the descent set of  $w$  equals  $S - I$  and that the descent set of  $w^{-1}$  equals  $S - J$  is  $\langle \tilde{\chi}_I, \tilde{\chi}_J \rangle_W$ .*

*Proof.*

$$\begin{aligned} \langle \tilde{\chi}_I, \tilde{\chi}_J \rangle_W &= \langle \Phi(y_I), \Phi(y_J) \rangle_W \\ &= \langle y_I, y_J \rangle, \quad \text{by Theorem 3.1.} \end{aligned}$$

As  $\langle y_I, y_J \rangle$  is the number of  $w$  in  $W$  such that the descent set of  $w$  equals  $S - I$  and that the descent set of  $w^{-1}$  equals  $S - J$ , the second statement is proved. As  $\Phi$  is an algebra homomorphism, the first one follows from

$$x_J = \sum_{K \supset J} y_K.$$

□

## 4 Conjectures and results

### 4.1 The symmetry conjecture

First, as for each  $x \in \Sigma W$ ,  $\Phi(x)$  is a class function on  $W$ , we can linearly extend it to  $\mathbb{Q}W$ .

In [13], Jöllenbeck and Reutenauer prove that if  $W$  is a Coxeter group of type  $A_n$ , then  $\Phi(x)(y) = \Phi(y)(x)$ , for all  $x, y \in \Sigma W$ . They sketch the following conjecture.

**Conjecture 4.1 (Symmetry conjecture).** *Let  $W$  be a finite Coxeter group. Then for all  $x, y \in \Sigma W$  we have*

$$\Phi(x)(y) = \Phi(y)(x).$$

The purpose of this section is to show the following theorem, which solves this conjecture for a subclass of finite Coxeter groups.

**Theorem 4.2.** *Let  $W$  be a finite Coxeter group whose irreducible components are of type  $A_n$ ,  $E_6$ ,  $E_7$ ,  $E_8$ ,  $F_4$ ,  $H_3$ ,  $H_4$  or  $I_2(m)$ . Then for all  $x, y \in \Sigma W$  we have*

$$\Phi(x)(y) = \Phi(y)(x).$$

**Corollary 4.3.** *With the same hypothesis than Theorem 4.2 one has for any  $I, J \subset S$ :*

$$\sum_{w \in X_I} 1_{W_J}^W(w) = \sum_{g \in X_J} 1_{W_I}^W(g).$$

**Restriction to a basis.** In the next proposition, one readily sees that our investigation can be restricted to  $x_I, x_J \in \Sigma W$ , for  $I, J \subset S$ . It also proves Corollary 4.3.

**Proposition 4.4.** *Let  $W$  be a finite Coxeter group. Then the following propositions are equivalent:*

1.  $\Phi(x)(y) = \Phi(y)(x)$ ,  $\forall x, y \in \Sigma W$ ;
2.  $\Phi(x_I)(x_J) = \Phi(x_J)(x_I)$ ,  $\forall I, J \subset S$ ;
3.  $\sum_{w \in X_I} 1_{W_J}^W(w) = \sum_{g \in X_J} 1_{W_I}^W(g)$ ,  $\forall I, J \subset S$ .

**Restriction to irreducible Coxeter systems.** Now, we show that it is enough to look at irreducible finite Coxeter groups to prove the conjecture. Let  $(W, S)$  be a finite Coxeter system with  $(W_1, I_1), \dots, (W_n, I_n)$  its irreducible components. Thus  $W$  is isomorphic to  $W_1 \times \dots \times W_n$ .

It is well-known that if  $\chi_i \in \mathbb{Q}_{class} W_i$  and  $\chi_j \in \mathbb{Q}_{class} W_j$  then the tensor product

$$\chi_i \otimes \chi_j(w_1, w_2) = \chi_i(w_1)\chi_j(w_2), \quad \forall (w_1, w_2) \in W_i \times W_j.$$

is a class function in  $\mathbb{Q}_{class}(W_i \times W_j)$ . For any  $w \in W$ , there is a unique  $w_i \in W_i$  such that  $w = w_1 \dots w_n$ . Then

$$\chi_1 \otimes \dots \otimes \chi_n(w) = \chi_1(w_1) \dots \chi_n(w_n).$$

Denote by  $\Phi_p$  the Solomon algebra homomorphism from  $\Sigma W_p$  to  $\mathbb{Q}_{class} W_p$ .

**Proposition 4.5.** *If for  $p = 1, \dots, n$ ,*

$$\Phi_p(x_p)(y_p) = \Phi_p(y_p)(x_p), \quad \forall x_p, y_p \in \Sigma W_p$$

*then*

$$\Phi(x)(y) = \Phi(y)(x), \quad \forall x, y \in \Sigma W.$$

*Proof.* Proposition 4.4 implies that one just has to show  $\Phi(x_K)(x_L) = \Phi(x_L)(x_K)$ , for all  $K, L \in S$ . Following the proof of Lemma 2.3, one just shows the proposition for  $p = 1, 2$  and for  $M \subset S$ , one denotes  $M_p = M \cap I_p \subset I_p$ ,  $p = 1, 2$ . Remember that  $X_{M_p}^{I_p} = W_p \cap X_p$ . By [1], Lemma 3.1, one has

$$X_M = X_{M_1}^{I_1} X_{M_2}^{I_2}$$

hence in  $\mathbb{Q}W$ ,

$$x_M = x_{M_1}^{I_1} x_{M_2}^{I_2}.$$

It is well-known that

$$1_{W_M}^W = 1_{W_{M_1}}^{W_1} \otimes 1_{W_{M_2}}^{W_2},$$

therefore, for  $K, L \subset S$  one has

$$\begin{aligned} \Phi(x_K)(x_L) &= \sum_{w \in X_L} 1_{W_K}^W(w) \\ &= \sum_{(w_1, w_2) \in X_{L_1}^{I_1} \times X_{L_2}^{I_2}} 1_{W_{K_1}}^{W_1} \otimes 1_{W_{K_2}}^{W_2}(w_1 w_2) \\ &= \sum_{(w_1, w_2) \in X_{L_1}^{I_1} \times X_{L_2}^{I_2}} 1_{W_{K_1}}^{W_1}(w_1) 1_{W_{K_2}}^{W_2}(w_2) \\ &= \sum_{w_1 \in X_{L_1}^{I_1}} 1_{W_{K_1}}^{W_1}(w_1) \sum_{w_2 \in X_{L_2}^{I_2}} 1_{W_{K_2}}^{W_2}(w_2) \\ &= 1_{W_{K_1}}^{W_1} \left( \sum_{w_1 \in X_{L_1}^{I_1}} w_1 \right) 1_{W_{K_2}}^{W_2} \left( \sum_{w_2 \in X_{L_2}^{I_2}} w_2 \right) \\ &= 1_{W_{K_1}}^{W_1}(x_{L_1}^{I_1}) 1_{W_{K_2}}^{W_2}(x_{L_2}^{I_2}) \\ &= 1_{W_{L_1}}^{W_1}(x_{K_1}^{I_1}) 1_{W_{L_2}}^{W_2}(x_{K_2}^{I_2}), \text{ as } x_{K_p}^{I_p}, x_{L_p}^{I_p} \in \Sigma W_p \\ &= \Phi(x_L)(x_K). \end{aligned}$$

□

#### Restriction to Coxeter classes.

**Proposition 4.6.** *Let  $I, J \subset S$ . If  $I \sim_W J$  then  $\Phi(x_I)(x_J) = \Phi(x_J)(x_I)$ .*

*Proof.* As  $x_I - x_J \in \ker \Phi$ , one has  $\Phi(x_I) = \Phi(x_J)$ . By Lemma 2.5, one has  $\Phi(x_J)(x_J - x_I) = 0$ . Finally,

$$\Phi(x_I)(x_J) = \Phi(x_J)(x_J) = \Phi(x_J)(x_I).$$

□

## 4.2 Othogonal idempotents and character formulas

In [1], the authors have shown that for any finite Coxeter group  $W$ , there is a family of idempotents  $E_\lambda \in \Sigma W$  indexed by the Coxeter classes  $\lambda \in \Lambda(W)$  satisfying  $\Phi(E_\lambda) = \xi_\lambda$ . We call a family of *good idempotents* such a family. We denote  $\chi_\lambda$  the character of the right action of  $W$  on  $E_\lambda \mathbb{Q}W$ .

Recall that if  $e, e'$  are idempotents in  $\mathbb{Q}W$ , with associated characters  $\chi, \chi'$ , then  $\dim(e\mathbb{Q}We') = \langle \chi, \chi' \rangle_W$ . In particular, if  $e' = e_I = \frac{1}{|W_I|} \sum_{w \in W_I} w$ , then  $\dim(e\mathbb{Q}We_I) = \dim(\text{Fix}_{W_I}(e\mathbb{Q}W)) = \langle \chi, 1_{W_I}^W \rangle_W$ . The purpose of this section is to show the following theorem and corollary. The theorem states that the symmetry conjecture (Conjecture 4.1) is equivalent to the calculation of the cardinality of  $C(\lambda) \cap X_I$  as the scalar product of  $1_{W_I}^W$  with  $\chi_\lambda$ .

**Theorem 4.7.** *Let  $W$  be a finite Coxeter group. Then the following items are equivalent:*

1. *The symmetry conjecture holds;*
2. *For any family of good idempotents  $(E_\lambda)_{\lambda \in \Lambda(W)} \subset \Sigma W$ , we have*

$$|C(\lambda) \cap X_I| = \langle \chi_\lambda, 1_{W_I}^W \rangle_W,$$

*for all  $I \subset S$  and for all  $\lambda \in \Lambda(W)$ .*

**Remark.** In the case  $A_{n-1} = S_n$ , Coxeter classes and conjugacy classes are in bijection. Moreover, they are indexed by the partitions of  $n$ .

In [6] (see also [16], Section 9.2), the authors have shown the existence of a good family in the case  $A_{n-1}$ . They have also shown that each indecomposable projective right module generated by  $E_\lambda$ ,  $\lambda$  partition of  $n$ , is the multilinear part of the subspaces generated by symmetrized Lie polynomials of type  $\lambda$ .

Finally, in [13], the authors found an analog of Theorem 4.7 for the case  $A_{n-1}$  and they proved Theorem 4.2 in the case of symmetric group using [16], Corollary 9.44 that is they proved that the second part of Theorem 4.7 appear if  $W = A_{n-1}$ . This is how Jöllenbeck and Reutenauer proved Theorem 4.2 in this case.

The following corollary, which generalizes a result by Gessel and Reutenauer on the symmetric groups ([10]) and Poirier for hyperoctahedral groups ([15]), is an immediate consequence of Theorem 4.7 and Theorem 4.2.

**Corollary 4.8.** *Let  $W$  be a finite Coxeter group such that its irreducible components are of type  $A_n, E_6, E_7, E_8, F_4, H_3, H_4$  or  $I_2(m)$ . Then the number of  $\lambda$ -elements of  $W$  which have  $I$  as ascent set is  $\langle \chi_\lambda, \tilde{\chi}_I \rangle_W$ , where  $\tilde{\chi}_I = \Phi(y_I)$ .*

We need the following lemma to prove Theorem 4.7. It is the consequence of the results on  $\ker \Phi$  ([19]) and  $\text{Im} \Phi$  ([1]).

**Lemma 4.9.** *Let  $(E_\lambda)_{\lambda \in \Lambda(W)} \subset \Sigma W$  be a family of good idempotents and denote  $\Sigma_\lambda W$  the subspace of  $\Sigma W$  generated by  $\{E_\lambda\}_{\lambda \in \Lambda(W)}$  then*

1.  $\Sigma_\Lambda W$  is a subalgebra of  $\Sigma W$  and  $(E_\lambda)_{\lambda \in \Lambda(W)}$  is one of its bases;
2.  $\Sigma W = \ker \Phi \oplus \Sigma_\Lambda W$ ;
3.  $\Phi$  is an isomorphism from  $\Sigma_\Lambda W$  to  $\Gamma W$ .

*Proof.* As the  $E_\lambda$  are orthogonal idempotents,  $\Sigma_\Lambda W$  is an algebra. To prove that  $(E_\lambda)_{\lambda \in \Lambda(W)}$  is a basis of  $\Sigma_\Lambda W$ , one just has to show that  $(E_\lambda)_{\lambda \in \Lambda(W)}$  is free. Let  $a_\lambda \in \mathbb{Q}$  such that

$$\sum_{\lambda \in \Lambda(W)} a_\lambda E_\lambda = 0.$$

As  $E_\lambda \in \Sigma W$ , this implies

$$\begin{aligned} \Phi\left(\sum_{\lambda \in \Lambda(W)} a_\lambda E_\lambda\right) &= \sum_{\lambda \in \Lambda(W)} a_\lambda \Phi(E_\lambda) \\ &= \sum_{\lambda \in \Lambda(W)} a_\lambda \xi_\lambda \\ &= 0. \end{aligned}$$

As  $(\xi_\lambda)_{\lambda \in \Lambda(W)}$  is a basis of  $\Gamma W$ ,  $a_\lambda = 0$  for all  $\lambda \in \Lambda(W)$ . Therefore,  $(E_\lambda)_{\lambda \in \Lambda(W)}$  is a basis of  $\Sigma_\Lambda W$  and  $\dim \Sigma_\Lambda W = |\Lambda(W)|$ .

Let  $x \in \ker \Phi \cap \Sigma_\Lambda W$ , then  $\Phi(x) = \sum_{\lambda \in \Lambda(W)} a_\lambda \xi_\lambda = 0$  where  $a_\lambda \in \mathbb{Q}$ . For any  $\lambda \in \Lambda(W)$  and  $w$  a  $\lambda$ -element,  $\Phi(x)(w) = a_\lambda = 0$ . Therefore  $x = 0$ . This implies  $\ker \Phi \cap \Sigma_\Lambda W = \{0\}$  thus  $\ker \Phi \oplus \Sigma_\Lambda W$ . As  $\dim \Sigma_\Lambda W + \dim \ker \Phi = \dim \Sigma W$ ,  $\Sigma W = \ker \Phi \oplus \Sigma_\Lambda W$ .

Therefore, the restriction of  $\Phi$  to  $\Sigma_\Lambda W$  is an isomorphism from  $\Sigma_\Lambda W$  to  $\Gamma W$ .  $\square$

**Corollary 4.10.** *Any family of good idempotents is a family of othogonal idempotents.*

*Proof.* Let  $\lambda, \mu$  be two distincts Coxeter classes. Then  $\Phi(E_\lambda E_\mu) = \xi_\lambda \xi_\mu = 0$ , hence  $E_\lambda E_\mu \in \ker \Phi$ . As  $E_\lambda E_\mu \in \Sigma_\Lambda W$ , one has  $E_\lambda E_\mu = 0$  by Lemma 4.9 (2).  $\square$

**Remark.** In [1], the authors have obtained a set of good idempotents  $(E_\lambda)_{\lambda \in \Lambda(W)} \subset \Sigma W$  such that their sum for  $\lambda \in \Lambda(W)$  equals the identity of  $\mathbb{Q}W$ . In this case,  $\mathbb{Q}W = \bigoplus_{\lambda \in \Lambda(W)} E_\lambda \mathbb{Q}W$ .

*Proof of Theorem 4.7.* One may assume that  $x = x_I$  and  $y = x_J$  for some  $I, J \subset S$ , by Proposition 4.4.

One has  $|C(\lambda) \cap X_I| = \xi_\lambda(x_I) = \Phi(E_\lambda)(x_I)$ . By Lemma 3.5,  $\Phi(x_I)(E_\lambda) = 1_{W_I}^W(E_\lambda) = \langle \chi_\lambda, 1_{W_I}^W \rangle_W$ , hence (1) implies (2).



(2) implies (1): As  $\{E_\lambda\}_{\lambda \in \Lambda W}$  is a basis of  $\Sigma_\Lambda W$  (Lemma 4.9) and  $\{x_I\}_{I \subset S}$  is a basis of  $\Sigma W$ , one just has to show that  $1_{W_I}^W(E_\lambda) = \Phi(E_\lambda)(x_I)$ . One has

$$\begin{aligned} \xi_\lambda(x_I) &= \Phi(E_\lambda)(x_I) \\ &= |X_I \cap C(\lambda)| \\ &= \langle \chi_\lambda, 1_{W_I}^W \rangle_W \\ &= 1_{W_I}^W(E_\lambda), \quad \text{by Lemma 3.5.} \end{aligned}$$

□

### 4.3 The double coset conjecture

Let  $I, J \subset S$  and  $b \in X_{IJ}$ . We consider the set

$$W(I, J, b) = \{w \in W \mid w^J b^{-1} = (wb^{-1})_I\},$$

where  $(w^J, w_J)$  (resp.  $((wb^{-1})^I, (wb^{-1})_I)$ ) are the parabolic components of  $w$  (resp.  $wb^{-1}$ ).

In [11], the authors have shown that  $W(I, J, e) = W(J, I, e)$ , for all  $I, J \subset S$ . For  $b \neq e$ ,  $W(I, J, b)$  is not generally equal to  $W(J, I, b^{-1})$ . They stated the following conjecture.

**Conjecture 4.11.** *If  $I, J \subset S$  then for all  $b \in X_{IJ}$ ,  $|W(I, J, b)| = |W(J, I, b^{-1})|$ .*

The following proposition shows that Conjecture 4.11 implies Conjecture 4.1.

**Proposition 4.12.** *Let  $I, J \subset S$  then*

$$\Phi(x_J)(x_I) = \sum_{b \in X_{IJ}} |W(I, J, b)|.$$

**Lemma 4.13.** *Let  $I, J \subset S$  and  $b \in X_{IJ}$ . Let  $w \in W$  then*

1. *if  $w \in X_{J \cap b^{-1}Ib}$ ,  $w_J \in X_{J \cap b^{-1}Ib}^J$ ;*
2. *If  $w \in W(I, J, b)$ ,  $w \in W_I b W_J$  and  $w^J \in X_{I \cap b J b^{-1}}^I b$ ;*
3.  *$w \in W(I, J, b)$  if and only if there are  $a \in X_{I \cap b J b^{-1}}^I$ ,  $c \in X_{J \cap b^{-1}Ib}^J$  and  $u \in X_I$  such that  $w = abc = uab$ . Moreover,  $a, c, u$  are unique.*

*Proof.* 1. As  $X_J X_{J \cap b^{-1}Ib}^J = X_{J \cap b^{-1}Ib}$ ,  $w_J \in X_{J \cap b^{-1}Ib}^J$ .

2. One has

$$W_I w W_J = W_I w^J w_J W_J = W_I w^J W_J = W_I (wb^{-1})_I b W_J = W_I b W_J.$$

As seen in Section 2.1,  $w^J \in X_{I \cap b J b^{-1}}^I b$ .

3. One has  $w^J = ab$  with a unique  $a \in X_{I \cap bJb^{-1}}^I$ , by Lemma 2. Let  $u = w^I \in X_I$  and  $c = w_J$  uniquely determined as parabolic components. Then

$$\begin{aligned} w = uab &\in X_I X_{I \cap bJb^{-1}}^I b \\ &= X_{I \cap bJb^{-1}} b, && \text{see Section 2.1} \\ &= X_{J \cap b^{-1}Jb}, && \text{see Section 2.1.} \end{aligned}$$

By (1),  $c \in X_{J \cap b^{-1}Ib}^J$ .

If they are  $a \in X_{I \cap bJb^{-1}}^I$ ,  $c \in X_{J \cap b^{-1}Ib}^J$  and  $u \in X_I$  such that  $w = abc = uab$ , then  $w^J b^{-1} = abb^{-1} = a = (wb^{-1})_I$ . Thus  $w \in W(I, J, b)$ .  $\square$

*Proof of Proposition 4.12.* As  $1_{W_J}^W(u) = |\{g \in X_J \mid ug \in gW_J\}|$  (see [12], Section 3.14),

$$\begin{aligned} \Phi(x_J)(x_I) &= \sum_{u \in X_I} 1_{W_J}^W(u) \\ &= |\{(u, g) \in X_I \times X_J \mid ug \in gW_J\}|. \end{aligned}$$

As  $X_J$  is the disjoint union of  $X_{I \cap bJb^{-1}}^I b$  for all  $b \in X_{IJ}$ , the double parabolic components of  $g \in X_{I \cap bJb^{-1}}^I b$  are  $(a, b, e)$ . Hence,

$$\begin{aligned} \Phi(x_J)(x_I) &= \sum_{b \in X_{IJ}} |\{(u, a) \in X_I \times X_{I \cap bJb^{-1}}^I \mid uab \in abW_J\}| \\ &= \sum_{b \in X_{IJ}} |W'(I, J, b)|. \end{aligned}$$

One just has to show that  $|W(I, J, b)| = |W'(I, J, b)|$ .

By Lemma 4.13, (3), for any  $w \in W$  one has unique  $a \in X_{I \cap bJb^{-1}}^I$ ,  $c \in X_{J \cap b^{-1}Ib}^J$  and  $u \in X_I$  such that  $w = abc = uab$ . Therefore, the following mapping

$$\begin{aligned} f : W(I, J, b) &\longrightarrow W'(I, J, b) \\ w &\longmapsto (u, a) \end{aligned}$$

is well defined and is injective (parabolic components are unique and determine uniquely each  $w \in W$ ).

Let  $(w, a) \in W'(I, J, b)$ , then

$$(wab)^J b^{-1} = (ab)b^{-1} = a = ((wab)b^{-1})_I.$$

Then  $wab \in W(I, J, b)$  and  $f(wab) = (w, a)$ . Therefore  $f$  is a bijection and the proposition is proved.  $\square$

#### 4.4 Results

The following proposition proves Conjecture 4.11 for any  $I \subset S$  and  $J = \emptyset, S$ .

**Proposition 4.14.** *If  $(W, S)$  is a finite Coxeter system and if  $I \subset S$ , then*

1.  $|W(I, \emptyset, b)| = |W(\emptyset, I, b^{-1})| = |W_I|$ , for all  $b \in X_{I\emptyset}$ ;
2.  $X_{IS} = \{e\}$  and  $W(I, S, e) = W(S, I, e) = X_I$ .

*Proof.* 1. Observe that  $X_{\emptyset I} = X_I$ . If  $w \in W$  then  $(w^\emptyset, w_\emptyset) = (w, e)$ . Let  $b \in X_I^{-1}$ . Then, on one hand,

$$\begin{aligned} W(I, \emptyset, b) &= \{w \in W \mid w^\emptyset b^{-1} = (wb^{-1})_I\} \\ &= \{w \in W \mid w = (wb^{-1})_I b\} \\ &= \{w \in W \mid w \in W_I b\} \\ &= W_I b \end{aligned}$$

On the other hand,

$$\begin{aligned} W(\emptyset, I, b^{-1}) &= \{w \in W \mid w^I b = (wb)_\emptyset\} \\ &= \{w \in W \mid w^I b = e\} \\ &= \{w \in W \mid w^I = b^{-1}\} \\ &= \{w \in W \mid w \in b^{-1} W_I\} \\ &= b^{-1} W_I \end{aligned}$$

2. Observe that  $X_{IS} = \{e\}$  then  $(w^S, w_S) = (e, w)$ . On one hand,

$$\begin{aligned} W(I, S, b) &= W(I, S, e) \\ &= \{w \in W \mid w^S = w_I\} \\ &= \{w \in W \mid e = w_I\} \\ &= \{w \in W \mid w^I \in X_I\} \\ &= X_I \end{aligned}$$

On the other hand,

$$\begin{aligned} W(S, I, b^{-1}) &= W(S, I, e) \\ &= \{w \in W \mid w^I = w_S\} \\ &= \{w \in W \mid W^I = w\} \\ &= X_I \end{aligned}$$

□

The following corollary, that proves Conjecture 4.1 for any  $I \subset S$  and  $J = \emptyset, S$ , may be proved directly using Proposition 4.4.

**Corollary 4.15.** *Let  $(W, S)$  be a finite Coxeter system then for any  $I \subset S$*

$$1. \Phi(x_\emptyset)(x_I) = \Phi(x_I)(x_\emptyset),$$

$$2. \Phi(x_S)(x_I) = \Phi(x_I)(x_S).$$

*Proof.* It is an immediate consequence of Proposition 4.14 and Propostion 4.12.  $\square$

**Proposition 4.16.** *Let  $(W, S)$  be a finite Coxeter system,  $I, J \subset S$  and  $b \in X_{IJ}$  such that  $bJb^{-1} \subset I$ . Then*

$$|W(I, J, b)| = |W(J, I, b^{-1})| = \frac{|W_I|}{|W_J|}.$$

*Proof.* On one hand, observe that  $X_{J \cap b^{-1}Ib}^J = X_J^J = \{e\}$ . Thus  $W(I, J, b) = X_{I \cap bJb^{-1}}^I b$ , by Lemma 4.13. Therefore,

$$\begin{aligned} |W(I, J, b)| &= |X_{I \cap bJb^{-1}}^I| \\ &= \frac{|W_I|}{|W_{I \cap bJb^{-1}}|} \\ &= \frac{|W_I|}{|W_I \cap bW_Jb^{-1}|} \\ &= \frac{|W_I|}{|bW_Jb^{-1}|} \\ &= \frac{|W_I|}{|W_J|}, \quad \text{as } bW_Jb^{-1} \subset W_I. \end{aligned}$$

On the other hand,  $W(J, I, b^{-1}) \subset b^{-1}X_{I \cap bJb^{-1}}^I$  by Lemma 4.13. Let  $a \in X_{I \cap bJb^{-1}}^I$ , then

$$\begin{aligned} b^{-1}ab &\in X_I X_{I \cap bJb^{-1}}^I b \\ &= X_{J \cap b^{-1}Ib} \\ &= X_J, \text{ as } J \subset b^{-1}Ib. \end{aligned}$$

Hence  $b^{-1}a = uab \in W(J, I, b^{-1})$  by Lemma 4.13. Therefore  $W(J, I, b^{-1}) = b^{-1}X_{I \cap bJb^{-1}}^I$  and the lemma is proved.  $\square$

## 4.5 The case of a single generator

We denote  $t$  for  $\{t\} \subset S$  to simplify the notations.

**Theorem 4.17.** *Let  $(W, S)$  be a finite Coxeter system,  $t \in S$  and  $I \subset S$  then for all  $b \in X_{It}$*

$$|W(I, t, b)| = |W(t, I, b^{-1})|.$$

**Corollary 4.18.** *The double coset conjecture holds for dihedral groups.*

*Proof.* As for  $I_2(m)$ ,  $S = \{s, t\}$ , the subsets of  $S$  are  $\emptyset$ ,  $s$ ,  $t$ ,  $S$ , one concludes with Corollary 4.15 and Theorem 4.17.  $\square$

**Corollary 4.19.** *Let  $(W, S)$  be a finite Coxeter system. Then*

$$\Phi(x_I)(x_t) = \Phi(x_t)(x_I),$$

for any  $t \in S$  and any  $I \subset S$ . On other words,

$$\sum_{w \in X_I} 1_{W_t}^W(w) = \sum_{g \in X_t} 1_{W_I}^W(g),$$

for any  $t \in S$  and any  $I \subset S$ .

*Proof of the theorem.* 1. If  $btb^{-1} \in I$ , one concludes by Proposition 4.16.

2. From now on, one supposes that  $btb^{-1} \notin I$ . Thus

$$X_{I \cap btb^{-1}}^I = W_I \implies W_I b \subset X_t$$

and

$$X_{t \cap b^{-1}Ib}^t = W_t \implies W_t b^{-1} \subset X_I.$$

3. Let  $d \in X_{II}$  such that  $btb \in W_I d W_I$ . Then  $d^2 = e$ . Indeed, as  $d \in X_{II}$  is the unique element of minimal length in  $W_I d W_I$  and as

$$(btb^{-1})^{-1} = btb^{-1} \in W_I d^{-1} W_I = W_I d W_I,$$

one has  $d = d^{-1}$ .

4. There is a unique  $a_0 \in X_{I \cap dId}^I$  such that  $btb^{-1} = a_0 d a_0^{-1}$ . Indeed, there is a unique  $(a_0, a_1) \in X_{I \cap dId}^I \times W_I$  such that  $btb^{-1} = a_0 d a_1$ . Thus

$$tb^{-1} a_1 = b^{-1} a_0 d \in X_I X_{I \cap dId}^I d = X_{I \cap dId} d = X_{I \cap dId}.$$

Then  $a_1 \in X_{I \cap dId}^I$ , by Lemma 4.13 (1).

Finally one has

$$btb^{-1} = (btb^{-1})^{-1} = a_0 d a_1^{-1} = a_1 d a_0^{-1}.$$

By unicity of double parabolic components,  $a_0 = a_1$ .

5. One defines a mapping:

$$\begin{aligned} \sigma : W_I &\longrightarrow btb^{-1} W_I btb^{-1} \\ a &\longmapsto btb^{-1} a btb^{-1} \end{aligned}$$

As  $\sigma$  is the restriction of the conjugation by  $btb^{-1}$  on  $W_I$ , it is a isomorphism. Denote

$$Fix_\sigma W_I = \{a \in W_I \mid \sigma(a) = a\}.$$

As  $e \in W_I$  and  $\sigma(e) = e$ , this set is non empty. One will show that

$$|W(I, t, b)| = |W_I| + |Fix_\sigma W_I|.$$

In fact, one will prove that

$$(\star) \quad W(I, t, b) = W_I b \amalg a_0^{-1} Fix_\sigma W_I b t.$$

It is immediate that  $W_I b \subset W(I, t, b)$ .

Let  $w = a_0^{-1} a b t \in Fix_\sigma W_I$ . As  $a_0^{-1} a \in W_I$ ,  $a_0^{-1} a b \in X_t$ , by 2. Moreover,

$$\begin{aligned} a_0^{-1} a b t b^{-1} a^{-1} a_0 &= a_0^{-1} (a a_0 d a_0^{-1}) a^{-1} a_0, \text{ as } b t b^{-1} = a_0 d a_0^{-1} \\ &= a_0^{-1} (a_0 d a_0^{-1} a) a^{-1} a_0, \text{ as } a \in Fix_\sigma W_I \\ &= d \in X_{II} \subset X_I. \end{aligned}$$

Therefore  $w \in W(I, t, b)$  that implies  $a_0^{-1} Fix_\sigma W_I b t \subset W(I, t, b)$ .

Conversely, let  $w \in W(I, t, b)$ , then there is  $a \in W_I$  such that  $w = ab \in W_I b$  or  $w = abt$ . In the second case  $abt b^{-1} a^{-1} = u \in X_I$ , by Lemma 4.13. As  $u$  is an involution,  $u \in X_{II}$ . But  $u \in a W_I d W_I a^{-1} = W_I d W_I$  therefore  $u = d$ .

One writes  $w = a_0^{-1} (a_0 a) b t$ . Denote  $a' = a_0 a \in W_I$  thus

$$a_0^{-1} a' b t b^{-1} a'^{-1} a_0 = a b t b^{-1} a^{-1} = d.$$

therefore

$$a' b t b^{-1} a'^{-1} = a_0 d a_0^{-1} = b t b^{-1}.$$

This implies  $a' \in Fix_\sigma W_I$  thus  $w \in a_0^{-1} Fix_\sigma W_I b t$  and  $(\star)$  is proved.

6. If  $a \in W_I$  and  $d_a \in X_{t\{t\}}$  such that  $b^{-1} a b \in W_t d_a W_t$ . Then the following propositions are equivalent:

- i)  $t d_a = d_a t$ ,
- ii)  $b^{-1} a b = t b^{-1} a b t = d_a$ ,
- iii)  $\{b^{-1} a, t b^{-1} a\} \subset W(t, I, b^{-1})$ ,
- iv)  $b^{-1} a b = t b^{-1} a b t$ .

$i) \Rightarrow ii)$ : If  $t d_a = d_a t$  then  $X_{t \cap d_a t d_a^{-1}}^t = \{e\}$  and  $W_t d_a W_t = \{d_a, d_a t\}$ . As  $b^{-1} a b \in W_t d_a W_t$ ,  $b^{-1} a b = d_a$  or  $b^{-1} a b = d_a t$ . In the second case, that is  $ab = b d_a t$ , one has  $b d_a \in X_t d_a = X_{d_a^{-1} t d_a} = X_t$ . Thus  $ab = b d_a t \notin X_t$ , which is a contradiction with  $ab \in W_I b \subset X_t$ . Therefore  $b^{-1} a b = d_a = t d_a t = t b^{-1} a b t$ .

$ii) \Rightarrow iii)$ : If  $b^{-1} a b = t b^{-1} a b t = d_a$ , then  $b^{-1} a = d_a b$  and  $t b^{-1} a = d_a t b$ . Therefore,  $\{b^{-1} a, t b^{-1} a\} \subset W(t, I, b^{-1})$ , by Lemma 4.13 3.

$iii) \Rightarrow i)$ : If  $\{b^{-1} a, t b^{-1} a\} \subset W(t, I, b^{-1})$ , thus  $b^{-1} a b$  and  $t b^{-1} a b t$  are in  $X_t$ . One supposes that  $t d_a \neq d_a t$ . Then  $X_{t \cap d_a t d_a^{-1}}^t = W_t$ , hence  $W_t d_a \subset$

$X_t$ .

Therefore,

$$b^{-1}ab = \begin{cases} d_a \in X_t & \text{or} \\ td_a \in X_t \end{cases}.$$

Thus

$$tb^{-1}abt = \begin{cases} td_at \notin X_t & \text{or} \\ d_at \notin X_t \end{cases}.$$

Which is a contradiction with *iii*). Hence  $b^{-1}ab = tb^{-1}abt \in X_t$ . Therefore,

$$b^{-1}ab = tb^{-1}abt = \begin{cases} d_a \in X_t & \text{or} \\ td_a \in X_t \end{cases}.$$

Finally, either  $b^{-1}ab = d_a$ , then  $tb^{-1}abt = td_at = d_a$  hence  $d_at = td_a$ ; or  $b^{-1}ab = td_a$ , then  $tb^{-1}abt = d_at = td_a$  hence  $d_at = td_a$ .

*i*)  $\iff$  *iv*): As *i*)  $\Rightarrow$  *ii*), one has in particular  $b^{-1}ab = tb^{-1}abt$  thus *i*)  $\Rightarrow$  *iv*).

If  $b^{-1}ab = tb^{-1}abt$ , so

$$b^{-1}ab = tb^{-1}abt = \begin{cases} d_a = td_at \in X_I & \text{or} \\ td_a = d_at \in X_I & \text{or} \\ d_at = td_a \notin X_I & \text{or} \\ td_at = d_a \notin X_I \end{cases}.$$

Observe that  $td_a = d_at$  in all the cases.

7. One has:

$$|W(I, t, b^{-1})| = |W_I| + |\{a \in W_I \mid d_at = td_a\}|.$$

Indeed, let  $a \in W_I$  such that  $d_at \neq td_a$  thus  $X_{t \cap d_atd_a^{-1}}^t = W_t$ . As  $b^{-1}ab \in W_t d_a W_t$ , one has

$$b^{-1}ab = \begin{cases} d_a \in X_I & \text{or} \\ td_a \in X_I & \text{or} \\ d_at \notin X_I & \text{or} \\ td_at \notin X_I \end{cases}.$$

Observe that either

$$b^{-1}a \in W(t, I, b^{-1}) \quad \text{and} \quad tb^{-1}a \notin W(t, I, b^{-1})$$

or

$$b^{-1}a \notin W(t, I, b^{-1}) \quad \text{and} \quad tb^{-1}a \in W(t, I, b^{-1}).$$

As  $W_I$  is the disjoint union of  $\{a \in W_I \mid d_at = td_a\}$  and  $\{a \in W_I \mid d_at \neq td_a\}$ , one has:

$$\begin{aligned} |W(t, I, b^{-1})| &= \sum_{a \in W_I \mid d_at = td_a} 2 + \sum_{a \in W_I \mid d_at \neq td_a} 1, \quad \text{by (6)} \\ &= 2|\{a \in W_I \mid d_at = td_a\}| + |\{a \in W_I \mid d_at \neq td_a\}| \\ &= |W_I| + |\{a \in W_I \mid d_at = td_a\}|. \end{aligned}$$

8. By 5 and 7, one just has to show that  $|Fix_\sigma W_I| = |\{w \in W_I \mid d_a t = t d_a\}|$  to finish the proof.

$$\begin{aligned}
a \in Fix_\sigma W_I &\iff abtb^{-1} = btb^{-1}a \\
&\iff b^{-1}ab = tb^{-1}abt \\
&\iff d_a t = t d_a, \quad \text{by 6} \\
&\iff a \in \{w \in W_I \mid d_a t = t d_a\}
\end{aligned}$$

□

#### 4.6 Proof of Theorem 4.2

By Proposition 4.4 and Proposition 4.5, one shows the theorem by using the classification of irreducible finite Coxeter groups (see [12]).

**Case  $A_n$ :** This case has been shown by Jöllenbeck and Reutenauer in [13]. An idea of their proof is to be found in the remark following Theorem 4.7.

**Case  $I_2(m)$ :** It is given by Proposition 4.12 and Corollary 4.18.

**Cases  $E_6, E_7, E_8, F_4, H_3, H_4$ :** We begin to give a lemma.

**Lemma 4.20.** *Let  $I, J \subset S$ , then*

$$\sum_{w \in X_I} 1_{W_J}^W(w) = \sum_{C \in Cl(W)} |X_I \cap C| 1_{W_J}^W(w_C),$$

where  $w_C$  is an arbitrary element in  $C$ .

*Proof.* One has

$$\sum_{w \in X_I} 1_{W_J}^W(w) = \sum_{C \in Cl(W)} \sum_{w \in X_I \cap C} 1_{W_J}^W(w).$$

As  $1_{W_J}^W$  is constant in conjugacy classes, one chooses an arbitrary element  $w_C \in C$ . Hence

$$\begin{aligned}
\sum_{w \in X_I} 1_{W_J}^W(w) &= \sum_{C \in Cl(W)} \sum_{w \in X_I \cap C} 1_{W_J}^W(w_C) \\
&= \sum_{C \in Cl(W)} 1_{W_J}^W(w_C) \sum_{w \in X_I \cap C} 1 \\
&= \sum_{C \in Cl(W)} |X_I \cap C| 1_{W_J}^W(w_C).
\end{aligned}$$

□



As, if  $J \sim J'$ ,  $1_{W_J}^W = 1_{W_{J'}}^W$  and  $|X_I \cap C| = |X_J \cap C|$  for some conjugacy class  $C$  (see Proposition 2.4), it is sufficient to consider Coxeter Classes of  $I$  and  $J$  in Lemma 4.20. Therefore, we construct two matrices

$$A = (a_{\lambda C})_{\lambda \in \Lambda(W), C \in Cl(W)} \text{ and } B = (b_{C'\mu})_{C' \in Cl(W), \mu \in \Lambda(W)},$$

where  $a_{\lambda C} = |X_I \cap C|$ , for some  $I \in \lambda$  and  $b_{C'\mu} = \varphi_\mu(w'_C)$ . Then  $AB = D = (d_{\lambda\mu})_{\lambda, \mu \in \Lambda(W)}$  where

$$d_{\lambda\mu} = \sum_{C \in Cl(W)} |X_I \cap C| 1_{W_J}^W(w_C) = \sum_{w \in X_I} 1_{W_J}^W(w), \text{ for some } I \in \lambda, J \in \mu.$$

Hence to prove Conjecture 4.1 amounts to prove that  $D$  is a symmetric matrix.

For these exceptional cases, we used the GAP part of CHEVIE ([7] and [17]) to obtain  $D$ . This program contains some useful functions to compute the matrix  $A$  and the matrix  $B$ , therefore the matrix  $D$ .

1. the GAP functions `CharTable`, `PositionId`, `StoreFusion` and `Induced` are used to compute the coefficients of  $A$ ;
2. The functions `ReflectionSubgroup`, `ReducedRightCosetRepresentatives` and `PositionClass` are used to compute the coefficients of  $B$ .

We just have to give a set of representants of Coxeter classes. Following the notation in [8], the following algorithm determines a set of representants for each Coxeter Class.

**Algorithm 4.21.** *[Construction of Representant of Coxeter Classes] Given a finite Coxeter System  $(W, S)$   $\Lambda(W)_{rep}$  - a set of representant of  $\Lambda(W)$  - is determinated. As  $I \sim J$  iff  $|I| = |J|$ , one just has to consider the subset of  $S$  of cardinality  $i$ , for  $i = 1, \dots, |S|$  to construct  $\Lambda(W)_{rep}$ .*

1. **[Initialize]** Let  $PS_i = \{I_1, \dots, I_m\}$  be the set of subsets of  $S$  of cardinality  $i$ . Set  $\Lambda(W)_{rep}^i \leftarrow I_1$  and

$$E_1 \leftarrow \{I_p \mid c_{I_p} \sim c_{I_1}\}.$$

2. **[Next  $k$ ]** If  $I_k \in E_{k-1}$ , then set  $k \leftarrow k + 1$ ,  $E_{k+1} \leftarrow E_k$ . Otherwise set

$$\Lambda(W)_{rep}^i \leftarrow \Lambda(W)_{rep}^i \cup I_k$$

and

$$E_k \leftarrow E_{k-1} \cup \{I_p \mid c_{I_p} \sim c_{I_k}\}$$

and  $k \leftarrow k + 1$ .

At this step, each subset of  $S$  in  $\Lambda(W)_{rep}^i$  is not conjugated to each of the others and each subset of  $S$  in  $E_k$  has a representant of it Coxeter class in  $\Lambda(W)_{rep}^i$ .

3. **[Loop]** If  $k < m + 1$  repeat step 2 else stop.

Then

$$\Lambda(W)_{rep} = \bigcup_{i=0}^{|S|} \Lambda(W)_{rep}^i.$$

By Corollary 4.19, one just has to take the submatrix  $A'$  of  $A$  and  $B'$  of  $B$  indexed by Coxeter Classes  $\lambda, \mu$  such that  $|I|, |J| \geq 2$  for some  $I \in \lambda$  and  $J \in \mu$ . The above algorithm can be initialize to  $i = 2$ .

As  $|E_8| > 6.10^8$ , using Corollary 4.19 is necessary to restrain the cardinality of cosets in this very large case. Finally, still in this case, we cannot directly construct and stock the set of minimal coset representatives  $X_I$  with GAP. Indeed for a set of two commuting generators in  $E_8$ ,  $|X_I| > 150.10^6$ . So we used the fact that for  $I \subset S$  and  $s \in S - I$  then

$$X_I = X_K X_I^K,$$

where  $K = S - \{s\}$ . If we keep in memory  $X_I$ , it take the memory of  $X_K$  times  $X_I^K$ , rather than if we keep in memory  $X_K$  and  $X_I^K$ , they take the memory for  $X_K$  plus  $X_I^K$ .

Finally, we just have to compute the submatrix  $D'$  of  $D$ , indexed by Coxeter Classes  $\lambda, \mu$  such that  $|I|, |J| \geq 2$ . Observe in the following that  $D'$  is a symmetric matrix. (The following matrix and tables have been obtained by one week of computation with GAP-CHEVIE; the original program also verified that  $D'$  is symmetric). This complete our proof.

**Remark about GAP-CHEVIE notations:** The set  $S$  of generators are given in CHEVIE by integers from 1 to  $n$ , see [7] for more informations. The set of representants  $\{I\}$  of Coxeter classes is given as a set of Coxeter  $I$ -elements.

**Coxeter group of type  $H_3$**  The set of representants of Coxeter classes that we consider is:

$$\{ 12, 123, 13, 23 \}.$$

Therefore the submatrix of  $D'$  indexed by them is

$$\begin{pmatrix} 24 & 12 & 40 & 32 \\ 12 & 1 & 30 & 20 \\ 40 & 30 & 52 & 46 \\ 32 & 20 & 46 & 38 \end{pmatrix}$$

**Coxeter group of type  $H_4$**  The set of representants of Coxeter classes that we consider is

$$\{ 12, 123, 1234, 124, 13, 134, 23, 234 \}.$$

Therefore the submatrix of  $D'$  indexed by them is

$$\begin{pmatrix} 4080 & 2040 & 1440 & 2840 & 5760 & 3280 & 4800 & 2640 \\ 2040 & 404 & 120 & 1116 & 4112 & 1602 & 2952 & 958 \\ 1440 & 120 & 1 & 720 & 3600 & 1200 & 2400 & 600 \\ 2840 & 1116 & 720 & 1828 & 4744 & 2288 & 3672 & 1664 \\ 5760 & 4112 & 3600 & 4744 & 7144 & 5104 & 6360 & 4592 \\ 3280 & 1602 & 1200 & 2288 & 5104 & 2730 & 4080 & 2132 \\ 4800 & 2952 & 2400 & 3672 & 6360 & 4080 & 5484 & 3504 \\ 2640 & 958 & 600 & 1664 & 4592 & 2132 & 3504 & 1510 \end{pmatrix}$$

**Coxeter group of type  $F_4$**  The set of representants of Coxeter classes that we consider is:

$$\{ 12, 123, 1234, 124, 13, 134, 23, 234, 34 \}.$$

Therefore the submatrix of  $D'$  indexed by them is:

$$\begin{pmatrix} 396 & 252 & 192 & 324 & 552 & 396 & 456 & 312 & 600 \\ 252 & 70 & 24 & 148 & 364 & 170 & 232 & 84 & 312 \\ 192 & 24 & 1 & 96 & 288 & 96 & 144 & 24 & 192 \\ 324 & 148 & 96 & 224 & 432 & 250 & 312 & 170 & 396 \\ 552 & 364 & 288 & 432 & 600 & 432 & 512 & 364 & 552 \\ 396 & 170 & 96 & 250 & 432 & 224 & 312 & 148 & 324 \\ 456 & 232 & 144 & 312 & 512 & 312 & 416 & 232 & 456 \\ 312 & 84 & 24 & 170 & 364 & 148 & 232 & 70 & 252 \\ 600 & 312 & 192 & 396 & 552 & 324 & 456 & 252 & 396 \end{pmatrix}.$$

**Coxeter group of type  $E_6$**  The set of representants of Coxeter classes that we consider is

$$\{12, 123, 1234, 12345, 123456, 12346, 1235, 12356, 1245, 125, 13, 134, 13456, 1356, 2345\}.$$

Therefore the submatrix of  $D'$  indexed by them is:

$$\begin{pmatrix} 29136 & 21744 & 16128 & 13728 & 12960 & 14544 & 18144 & 15456 & 17088 & 23040 & 26784 & 19776 & 14448 & 17856 & 15408 \\ 21744 & 12858 & 6882 & 4890 & 4320 & 5616 & 9252 & 6564 & 7950 & 14712 & 18432 & 10488 & 5412 & 8640 & 6300 \\ 16128 & 6882 & 1704 & 642 & 432 & 1088 & 3976 & 1864 & 2648 & 9108 & 12024 & 4432 & 840 & 3144 & 1380 \\ 13728 & 4890 & 642 & 87 & 27 & 341 & 2540 & 926 & 1378 & 7092 & 9432 & 2632 & 162 & 1776 & 444 \\ 12960 & 4320 & 432 & 27 & 1 & 216 & 2160 & 720 & 1080 & 6480 & 8640 & 2160 & 72 & 1440 & 270 \\ 14544 & 5616 & 1088 & 341 & 216 & 661 & 3080 & 1304 & 1882 & 7800 & 10368 & 3336 & 464 & 2304 & 852 \\ 18144 & 9252 & 3976 & 2540 & 2160 & 3080 & 6152 & 3894 & 4908 & 11256 & 14472 & 6976 & 2878 & 5460 & 3576 \\ 15456 & 6564 & 1864 & 926 & 720 & 1304 & 3894 & 1996 & 2690 & 8700 & 11448 & 4320 & 1104 & 3132 & 1602 \\ 17088 & 7950 & 2648 & 1378 & 1080 & 1882 & 4908 & 2690 & 3604 & 10080 & 13176 & 5552 & 1652 & 4140 & 2274 \\ 23040 & 14712 & 9108 & 7092 & 6480 & 7800 & 11256 & 8700 & 10080 & 16392 & 20016 & 12576 & 7644 & 10752 & 8544 \\ 26784 & 18432 & 12024 & 9432 & 8640 & 10368 & 14472 & 11448 & 13176 & 20016 & 23832 & 15984 & 10224 & 14112 & 11160 \\ 19776 & 10488 & 4432 & 2632 & 2160 & 3336 & 6976 & 4320 & 5552 & 12576 & 15984 & 7888 & 3088 & 6264 & 3864 \\ 14448 & 5412 & 840 & 162 & 72 & 464 & 2878 & 1104 & 1652 & 7644 & 10224 & 3088 & 250 & 2052 & 624 \\ 17856 & 8640 & 3144 & 1776 & 1440 & 2304 & 5460 & 3132 & 4140 & 10752 & 14112 & 6264 & 2052 & 4644 & 2784 \\ 15408 & 6300 & 1380 & 444 & 270 & 852 & 3576 & 1602 & 2274 & 8544 & 11160 & 3864 & 624 & 2784 & 1074 \end{pmatrix}.$$

**Coxeter group of type  $E_7$**  The set of representants of Coxeter classes that we consider is

$\{\lambda_1 = 12, \lambda_2 = 123, \lambda_3 = 1234, \lambda_4 = 12345, \lambda_5 = 123456, \lambda_6 = 1234567, \lambda_7 = 123457, \lambda_8 = 12346, \lambda_9 = 123467, \lambda_{10} = 1235, \lambda_{11} = 12356, \lambda_{12} = 123567, \lambda_{13} = 12357, \lambda_{14} = 1245, \lambda_{15} = 124567, \lambda_{16} = 12457, \lambda_{17} = 125, \lambda_{18} = 1257, \lambda_{19} = 13, \lambda_{20} = 134, \lambda_{21} = 13456, \lambda_{22} = 134567, \lambda_{23} = 13467, \lambda_{24} = 1356, \lambda_{25} = 2345, \lambda_{26} = 234567, \lambda_{27} = 23457, \lambda_{28} = 24567, \lambda_{29} = 2457, \lambda_{30} = 257\}.$

We give here the submatrix of  $D'$  indexed by this Coxeter classes. We have decomposed this matrix into three tables.

	$\lambda_1$	$\lambda_2$	$\lambda_3$	$\lambda_4$	$\lambda_5$	$\lambda_6$	$\lambda_7$	$\lambda_8$	$\lambda_9$	$\lambda_{10}$
$\lambda_1$	1733376	1320192	981504	823296	749568	725760	774912	877824	800256	1111296
$\lambda_2$	1320192	818544	452496	314592	258816	241920	278976	365568	300480	601968
$\lambda_3$	981504	452496	129972	50868	29412	24192	38160	84180	50220	271872
$\lambda_4$	823296	314592	50868	8896	2508	1512	5432	28056	11804	169328
$\lambda_5$	749568	258816	29412	2508	222	56	1330	15060	5294	131864
$\lambda_6$	725760	241920	24192	1512	56	1	756	12096	4032	120960
$\lambda_7$	774912	278976	38160	5432	1330	756	3198	20466	8122	145800
$\lambda_8$	877824	365568	84180	28056	15060	12096	20466	52152	28862	208448
$\lambda_9$	800256	300480	50220	11804	5294	4032	8122	28862	13762	161392
$\lambda_{10}$	1111296	601968	271872	169328	131864	120960	145800	208448	161392	410472
$\lambda_{11}$	936960	427152	134928	66140	45756	40320	53716	93912	63908	261352
$\lambda_{12}$	827904	326304	67260	22032	12334	10080	16286	41510	23056	181688
$\lambda_{13}$	969984	463968	167688	92004	67416	60480	76680	121980	88332	295176
$\lambda_{14}$	1044480	522096	191568	98680	68648	60480	80272	135896	94472	333824
$\lambda_{15}$	798720	297600	46452	8768	3026	2016	5490	25720	11126	158296
$\lambda_{16}$	933888	419808	123504	54780	35364	30240	42828	83056	53392	252776
$\lambda_{17}$	1398528	924672	577632	440352	381312	362880	402240	489408	423648	716544
$\lambda_{18}$	1177344	678720	347808	237840	194688	181440	210000	278736	227424	485568
$\lambda_{19}$	1612800	1147680	761760	586080	508320	483840	535680	649440	564480	914400
$\lambda_{20}$	1217280	689184	312000	181920	134496	120960	152448	232896	173088	472032
$\lambda_{21}$	8478672	335136	61764	14212	5636	4032	9396	35372	16456	183680
$\lambda_{22}$	773376	276576	36156	4580	1012	576	2674	19092	7298	143496
$\lambda_{23}$	903936	389328	100488	39404	23928	20160	30192	64996	39184	227136
$\lambda_{24}$	1072128	551424	218304	122088	89688	80640	101976	159696	116328	359904
$\lambda_{25}$	956160	427680	111312	37824	19344	15120	27168	69696	38688	252720
$\lambda_{26}$	772608	274896	34200	3424	402	126	1806	17714	6392	141928
$\lambda_{27}$	876288	360672	76800	21824	10096	7560	14992	45960	23512	203488
$\lambda_{28}$	870912	351792	67284	15480	5892	4032	9900	38460	17760	194280
$\lambda_{29}$	1067520	539280	198096	100752	69264	60480	80976	139728	96288	345264
$\lambda_{30}$	1421568	942336	585216	443520	382464	362880	403200	494208	426240	729216

	$\lambda_{11}$	$\lambda_{12}$	$\lambda_{13}$	$\lambda_{14}$	$\lambda_{15}$	$\lambda_{16}$	$\lambda_{17}$	$\lambda_{18}$	$\lambda_{19}$	$\lambda_{20}$
$\lambda_1$	936960	827904	969984	1044480	798720	933888	1398528	1177344	1612800	1217280
$\lambda_2$	427152	326304	463968	522096	297600	419808	924672	678720	1147680	689184
$\lambda_3$	134928	67260	167688	191568	46452	123504	577632	347808	761760	312000
$\lambda_4$	66140	22032	92004	98680	8768	54780	440352	237840	586080	181920
$\lambda_5$	45756	12334	67416	68648	3026	35364	381312	194688	508320	134496
$\lambda_6$	40320	10080	60480	60480	2016	30240	362880	181440	483840	120960
$\lambda_7$	53716	16286	76680	80272	5490	42828	402240	210000	535680	152448
$\lambda_8$	93912	41510	121980	135896	25720	83056	489408	278736	649440	232896
$\lambda_9$	63908	23056	88332	94472	11126	53392	423648	227424	564480	173088
$\lambda_{10}$	261352	181688	295176	333824	158296	252776	716544	485568	914400	472032
$\lambda_{11}$	138040	78624	167784	188688	60868	128288	547776	332976	718560	297216
$\lambda_{12}$	78624	33508	104076	113584	20118	67948	448992	248496	596160	198912
$\lambda_{13}$	167784	104076	197748	222384	84948	157728	583104	366336	756000	335808
$\lambda_{14}$	188688	113584	222384	254328	91056	178592	642624	410064	835200	385344
$\lambda_{15}$	60868	20118	84948	91056	8150	49644	421152	223440	563040	169824
$\lambda_{16}$	128288	67948	157728	178592	49644	117004	542112	323568	714240	288096
$\lambda_{17}$	547776	448992	583104	642624	421152	542112	1023360	790080	1238400	806112
$\lambda_{18}$	332976	248496	366336	410064	223440	323568	790080	558336	990720	554400
$\lambda_{19}$	718560	596160	756000	835200	563040	714240	1238400	990720	1470240	1023840
$\lambda_{20}$	297216	198912	335808	385344	169824	288096	806112	554400	1023840	546624
$\lambda_{21}$	75116	27576	102048	111768	13228	63656	460512	252768	614880	201792
$\lambda_{22}$	51936	15250	74904	78088	4752	41176	399936	207744	534240	149952
$\lambda_{23}$	108044	52608	137052	153784	35912	97240	512640	298032	679680	256704
$\lambda_{24}$	212328	135576	246192	281328	112752	202368	669792	434880	869760	417408
$\lambda_{25}$	120048	55344	152688	171888	35088	108672	555360	330480	728640	285120
$\lambda_{26}$	50604	14118	73320	76240	3694	39412	398688	205776	532800	147648
$\lambda_{27}$	88616	36032	116664	129440	19904	76672	486528	273984	645120	225792
$\lambda_{28}$	80400	29436	108072	120144	13788	67728	478176	263520	640800	216576
$\lambda_{29}$	194832	115728	228672	263712	91584	182640	660864	420240	862560	402048
$\lambda_{30}$	555264	451584	589824	653184	421632	546048	1042560	799488	1267200	824832

	$\lambda_{21}$	$\lambda_{22}$	$\lambda_{23}$	$\lambda_{24}$	$\lambda_{25}$	$\lambda_{26}$	$\lambda_{27}$	$\lambda_{28}$	$\lambda_{29}$	$\lambda_{30}$
$\lambda_1$	847872	773376	903936	1072128	956160	772608	876288	870912	1067520	1421568
$\lambda_2$	335136	276576	389328	551424	427680	274896	360672	351792	539280	942336
$\lambda_3$	61764	36156	100488	218304	111312	34200	76800	67284	198096	585216
$\lambda_4$	14212	4580	39404	122088	37824	3424	21824	15480	100752	443520
$\lambda_5$	5636	1012	23928	89688	19344	402	10096	5892	69264	382464
$\lambda_6$	4032	576	20160	80640	15120	126	7560	4032	60480	362880
$\lambda_7$	9396	2674	30192	101976	27168	1806	14992	9900	80976	403200
$\lambda_8$	35372	19092	64996	159696	69696	17714	45960	38460	139728	494208
$\lambda_9$	16456	7298	39184	116328	38688	6392	23512	17760	96288	426240
$\lambda_{10}$	183680	143496	227136	359904	252720	141928	203488	194280	345264	729216
$\lambda_{11}$	75116	51936	108044	212328	120048	50604	88616	80400	194832	555264
$\lambda_{12}$	27576	15250	52608	135576	55344	14118	36032	29436	115728	451584
$\lambda_{13}$	102048	74904	137052	246192	152688	73320	116664	108072	228672	589824
$\lambda_{14}$	111768	78088	153784	281328	171888	76240	129440	120144	263712	653184
$\lambda_{15}$	13228	4752	35912	112752	35088	3694	19904	13788	91584	421632
$\lambda_{16}$	63656	41176	97240	202368	108672	39412	76672	67728	182640	546048
$\lambda_{17}$	460512	399936	512640	669792	555360	398688	486528	478176	660864	1042560
$\lambda_{18}$	252768	207744	298032	434880	330480	205776	273984	263520	420240	799488
$\lambda_{19}$	614880	534240	679680	869760	728640	532800	645120	640800	862560	1267200
$\lambda_{20}$	201792	149952	256704	417408	285120	147648	225792	216576	402048	824832
$\lambda_{21}$	20172	8284	47200	135144	48192	7024	28968	21756	113808	463104
$\lambda_{22}$	8284	2150	28592	99576	25344	1392	13744	8688	78720	400896
$\lambda_{23}$	47200	28592	78284	177432	85920	27228	58968	50748	157776	517248
$\lambda_{24}$	135144	99576	177432	307128	199536	97872	153744	143544	289920	678528
$\lambda_{25}$	48192	25344	85920	199536	92688	23376	62304	53568	179424	565632
$\lambda_{26}$	7024	1392	27228	97872	23376	596	12120	7068	76224	398592
$\lambda_{27}$	28968	13744	58968	153744	62304	12120	39088	31224	132096	489600
$\lambda_{28}$	21756	8688	50748	143544	53568	7068	31224	22656	120672	478080
$\lambda_{29}$	113808	78720	157776	289920	179424	76224	132096	120672	269952	665856
$\lambda_{30}$	463104	400896	517248	678528	565632	398592	489600	478080	665856	1051776

**Coxeter group of type  $E_8$**  The set of representants of Coxeter classes that we consider is:

( $\lambda_1 = 12$ ,  $\lambda_2 = 123$ ,  $\lambda_3 = 1234$ ,  $\lambda_4 = 12345$ ,  $\lambda_5 = 123456$ ,  $\lambda_6 = 1234567$ ,  
 $\lambda_7 = 12345678$ ,  $\lambda_8 = 1234568$ ,  $\lambda_9 = 123457$ ,  $\lambda_{10} = 1234578$ ,  $\lambda_{11} = 12346$ ,  
 $\lambda_{12} = 123467$ ,  $\lambda_{13} = 1234678$ ,  $\lambda_{14} = 123468$ ,  $\lambda_{15} = 1235$ ,  $\lambda_{16} = 12356$ ,  
 $\lambda_{17} = 123567$ ,  $\lambda_{18} = 1235678$ ,  $\lambda_{19} = 123568$ ,  $\lambda_{20} = 12357$ ,  $\lambda_{21} = 1245$ ,  
 $\lambda_{22} = 124567$ ,  $\lambda_{23} = 1245678$ ,  $\lambda_{24} = 12457$ ,  $\lambda_{25} = 125$ ,  $\lambda_{26} = 1257$ ,  
 $\lambda_{27} = 13$ ,  $\lambda_{28} = 134$ ,  $\lambda_{29} = 13456$ ,  $\lambda_{30} = 134567$ ,  $\lambda_{31} = 1345678$ ,  
 $\lambda_{32} = 13467$ ,  $\lambda_{33} = 134678$ ,  $\lambda_{34} = 1356$ ,  $\lambda_{35} = 2345$ ,  $\lambda_{36} = 234567$ ,  
 $\lambda_{37} = 2345678$ ,  $\lambda_{38} = 23457$ ,  $\lambda_{39} = 234578$ ).

We give here the submatrix of  $D'$  indexed by this Coxeter classes. We have decomposed this matrix into four tables.

	$\lambda_1$	$\lambda_2$	$\lambda_3$	$\lambda_4$	$\lambda_5$	$\lambda_6$	$\lambda_7$	$\lambda_8$	$\lambda_9$	$\lambda_{10}$
$\lambda_1$	437160960	339793920	255744000	214179840	192015360	180034560	174182400	183121920	198650880	186301440
$\lambda_2$	339793920	219699360	126612000	88485120	70796160	62082720	58060800	64477440	76479840	66998880
$\lambda_3$	255744000	126612000	41733360	17606160	9810000	6918480	5806080	7849680	12730800	8969280
$\lambda_4$	214179840	88485120	17606160	3848832	1146672	530880	362880	767760	2308128	1171296
$\lambda_5$	192015360	70796160	9810000	1146672	143460	29772	13440	80592	611292	236028
$\lambda_6$	180034560	62082720	6918480	530880	29772	1596	240	15242	270192	93484
$\lambda_7$	174182400	58060800	5806080	362880	13440	240	1	6720	181440	60480
$\lambda_8$	183121920	64477440	7849680	767760	80592	15242	6720	44802	403548	151210
$\lambda_9$	198650880	76479840	12730800	2308128	611292	270192	181440	403548	1353996	658812
$\lambda_{10}$	186301440	66998880	8969280	1171296	236028	93484	60480	151210	658812	293848
$\lambda_{11}$	228648960	102617280	27612240	10012704	5163984	3517500	2903040	4058736	7034268	4761696
$\lambda_{12}$	205424640	82549440	16471440	4418016	1919700	1213572	967680	1455716	2934284	1824168
$\lambda_{13}$	189527040	69662880	10325040	1800768	575808	320262	241920	413784	1105044	598522
$\lambda_{14}$	209064960	86107680	18873840	5760384	2723772	1788716	1451520	2102964	3929460	2548040
$\lambda_{15}$	289059840	165227040	78967200	49372800	37176768	31559040	29030400	33144576	41255520	34897344
$\lambda_{16}$	244039680	118974240	41486880	20691600	13721040	10883304	9676800	11728152	16206312	12737136
$\lambda_{17}$	212613120	89501760	21350160	7495728	4031292	2868280	2419200	3241876	5366368	3749372
$\lambda_{18}$	192890880	72646560	12023040	2620992	1017696	617378	483840	758036	1687240	997262
$\lambda_{19}$	220078080	97133760	27315840	11716608	7220784	5533552	4838400	6049880	8880696	6707120
$\lambda_{20}$	252288000	128331360	50086080	27667872	19535400	16043784	14515200	17056944	22359768	18235608
$\lambda_{21}$	272010240	144900000	58440480	30997632	20717568	16357344	14515200	17662080	24329280	19130976
$\lambda_{22}$	202014720	79449120	14485680	3241824	1157268	644524	483840	830428	2034528	1140620
$\lambda_{23}$	186301440	67004640	8979360	1183392	248676	103974	69120	161932	670552	304712
$\lambda_{24}$	240215040	114829920	37707360	17456208	10890768	8315728	7257600	9104720	13278240	10052528
$\lambda_{25}$	339009280	245329920	156637440	119022336	100806912	91517184	87091200	93984768	106502400	96609024
$\lambda_{26}$	302561280	182062080	96975360	66309888	52937856	46522368	43545600	48278784	57293184	50215296
$\lambda_{27}$	401526800	301190400	205804800	159148800	134887680	122135040	116121600	125556480	142352640	128977920
$\lambda_{28}$	316477440	189504000	92640000	55407360	39456000	32179200	29030400	34310400	44862720	36564480
$\lambda_{29}$	217912320	92171520	20264400	5483088	2181720	1266912	967680	1591872	3538656	2079576
$\lambda_{30}$	195425280	73762560	11420880	1850592	459240	204064	138240	304136	1076372	515792
$\lambda_{31}$	183121920	64434240	7825680	785184	101592	29822	17280	60852	420692	166772
$\lambda_{32}$	232657920	106981920	31490400	13092576	7648560	5640672	4838400	6269376	9679568	7059296
$\lambda_{33}$	205655040	82964160	16991040	4849920	2253600	1486784	1209600	1744056	3295576	2127960
$\lambda_{34}$	276480000	150465600	64491840	36772128	26098848	21407328	19353600	22766976	29768256	24300000
$\lambda_{35}$	251596800	121875840	37313280	14054400	6943104	4503168	3628800	5326848	9694080	6350976
$\lambda_{36}$	192199680	71055360	10046880	1273920	198048	55904	30240	117144	694728	284088
$\lambda_{37}$	180126720	62232480	7044960	581664	43704	5694	2160	23510	300392	107758
$\lambda_{38}$	225054720	99004320	24805440	8013312	3656160	2289504	1814400	2759136	5386368	3393840
$\lambda_{39}$	202245120	79833600	14928480	3561984	1361136	789948	604800	993696	2276868	1320864

	$\lambda_{11}$	$\lambda_{12}$	$\lambda_{13}$	$\lambda_{14}$	$\lambda_{15}$	$\lambda_{16}$	$\lambda_{17}$	$\lambda_{18}$	$\lambda_{19}$	$\lambda_{20}$
$\lambda_1$	228648960	205424640	189527040	209064960	289059840	244039680	212613120	192890880	220078080	252288000
$\lambda_2$	102617280	82549440	69662880	86107680	165227040	118974240	89501760	72646560	97133760	128331360
$\lambda_3$	27612240	16471440	10325040	18873840	78967200	41486880	21350160	12023040	27315840	50086080
$\lambda_4$	10012704	4418016	1800768	5760384	49372800	20691600	7495728	2620992	11716608	27667872
$\lambda_5$	5163984	1919700	575808	2723772	37176768	13721040	4031292	1017696	7220784	19535400
$\lambda_6$	3517500	1213572	320262	1788716	31559040	10883304	2868280	617378	5533552	16043784
$\lambda_7$	2903040	967680	241920	1451520	29030400	9676800	2419200	483840	45315200	14515200
$\lambda_8$	4058736	1455716	413784	2102964	60756288	11728152	3241876	758036	6049880	17056944
$\lambda_9$	7034284	2934284	1105044	3929460	41255520	16206312	5366368	1687240	8880696	22359768
$\lambda_{10}$	4761696	1824168	598522	2548040	34897344	12737136	3749372	997262	6707120	18235608
$\lambda_{11}$	17498496	9734340	5677848	11465916	60756288	29139360	13445220	6837720	18242544	36592632
$\lambda_{12}$	9734340	4762868	2383000	5924412	45895296	19414512	7478296	3101068	11268400	25837632
$\lambda_{13}$	5677848	2383000	948816	3182820	36815232	13934304	4443900	1404342	7543112	19588272
$\lambda_{14}$	11465916	5924412	3182820	7197180	48708288	21425424	8819584	3992964	12776752	27992592
$\lambda_{15}$	60756288	45895296	36815232	48708288	115584864	75212064	51515040	39026592	57994176	83879424
$\lambda_{16}$	29139360	19414512	13934304	21425424	75212064	41410752	23615184	15378384	28820496	49126992
$\lambda_{17}$	13445220	7478296	4443900	8819584	51515040	23615184	10498956	5317000	14573384	30287568
$\lambda_{18}$	6837720	3101068	1404342	3992964	39026592	15738384	5317000	1930534	8569664	21186744
$\lambda_{19}$	18242544	11268400	7543112	12776752	57994176	28820496	14573384	8569664	18901640	35693808
$\lambda_{20}$	36592632	25837632	19588272	27992592	83879424	49126992	30287568	21186744	35693808	56943600
$\lambda_{21}$	41920320	28605120	20803008	31287360	95738400	56280480	33963456	22816416	40300608	65026656
$\lambda_{22}$	8273472	3741876	1643036	4821348	43491840	17712384	6318552	2293132	9973912	24004224
$\lambda_{23}$	4773192	1836244	609716	2559460	34903776	12747216	3760768	1008216	6716768	18242856
$\lambda_{24}$	25772256	16401888	11196690	18379008	71406624	37965408	20531616	12598624	25688512	45662208
$\lambda_{25}$	132655104	112520448	99330048	115941888	192948480	148253184	119280384	102279168	126604800	157102848
$\lambda_{26}$	77854848	62131968	52286976	64998528	132725760	92219520	67873152	54608640	74390016	100761600
$\lambda_{27}$	175737600	149921280	132503040	154172160	244166400	193363200	158112000	136339200	166717440	202901760
$\lambda_{28}$	69719040	50734080	39014400	54366720	133651200	86676480	57722880	41913600	65579520	96606720
$\lambda_{29}$	11999184	5813304	2791248	7258320	52382496	22940424	9063408	3704664	13435968	30030528
$\lambda_{30}$	6229488	2529564	915192	3432572	39367392	15106488	4810340	1432848	8168868	21089976
$\lambda_{31}$	4057860	1468100	428652	2111844	33130848	11729184	3251596	771600	6059208	17058936
$\lambda_{32}$	20798256	12513072	8054880	14307584	64718688	32587152	16345568	9287520	21236736	40084896
$\lambda_{33}$	10187232	5144832	2701152	6313696	46364160	19874784	7876864	3430720	11676352	26291712
$\lambda_{34}$	47684160	34102080	26024832	36765504	101408256	61969248	39464928	28053504	45759168	70604352
$\lambda_{35}$	23837184	13286016	7615872	15634944	74620800	37613952	18058752	9239040	23968512	46244736
$\lambda_{36}$	5339088	2029112	636624	2847288	37450368	13938816	4168160	1092280	7383336	19768752
$\lambda_{37}$	3601860	1258932	341066	1843456	31700928	10988856	2929372	645916	5608648	16158480
$\lambda_{38}$	15251184	7964928	4245936	9629280	57751776	26730960	11561520	5340288	16268256	34126176
$\lambda_{39}$	8629560	4008432	1838640	5103384	43913952	18091512	6611700	2505792	10290600	24394272

	$\lambda_{21}$	$\lambda_{22}$	$\lambda_{23}$	$\lambda_{24}$	$\lambda_{25}$	$\lambda_{26}$	$\lambda_{27}$	$\lambda_{28}$	$\lambda_{29}$	$\lambda_{30}$
$\lambda_1$	272010240	202014720	186301440	240215040	359009280	302561280	410572800	316477440	217912320	195425280
$\lambda_2$	144900000	79449120	67004640	114829920	245329920	182062080	301190400	189504000	92171520	73762560
$\lambda_3$	58440480	14485680	8979360	37707360	156637440	96975360	205804800	92640000	20264400	11420880
$\lambda_4$	30997632	3241824	1183392	17456208	119022336	66309888	159148800	55407360	5483088	1850592
$\lambda_5$	20717568	1157268	248676	10890768	100806912	52937856	134887680	39456000	2181720	459240
$\lambda_6$	16357344	644524	103974	8315728	91517184	46522368	122135040	32179200	1266912	204064
$\lambda_7$	14515200	483840	69120	7257600	87091200	43545600	116121600	29030400	967680	138240
$\lambda_8$	17662080	830428	161932	9104720	93984768	48278784	125556480	34310400	1591872	304136
$\lambda_9$	24329280	2034528	670552	13278240	106502400	57293184	142352640	44862720	3538656	1076372
$\lambda_{10}$	19130976	1140620	304712	10052528	96609024	50215296	128977920	36564480	2079576	515792
$\lambda_{11}$	41920320	8273472	4773192	25772256	132655104	77854848	175737600	69719040	11999184	6229488
$\lambda_{12}$	28605120	3741876	1836244	16401888	112520448	62131968	149921280	50734080	5813304	2529564
$\lambda_{13}$	20803008	1643036	609716	11196960	99330048	52286976	132503040	39014400	2791248	915192
$\lambda_{14}$	31287360	4821348	2559460	18379008	115941888	64998528	154172160	54366720	7258320	3432572
$\lambda_{15}$	95738400	43491840	34903776	71406624	192948480	132725760	244166400	133651200	52382496	39367392
$\lambda_{16}$	56280480	17712384	12747216	37965408	148253184	92219520	193363200	86676480	22940424	15106488
$\lambda_{17}$	338363456	6318552	3760768	20531616	119280384	67873152	158112000	57722880	9063408	4810340
$\lambda_{18}$	22816416	2293132	1008216	12598624	102279168	54608640	136339200	41913600	3704664	1432848
$\lambda_{19}$	40300608	9973912	6716768	25688512	126604800	74390016	166717440	65579520	13435968	8168688
$\lambda_{20}$	65026656	24004224	18242856	45662208	157102848	100761600	202901760	96606720	30030528	21089976
$\lambda_{21}$	75382176	26365632	19138656	52452096	173882880	113431680	224570880	111989760	33873792	22662144
$\lambda_{22}$	26365632	2781016	1153396	14736768	109458432	59635968	146085120	47696640	4562088	1696248
$\lambda_{23}$	19138656	1153396	316042	10062240	96609024	50216832	128977920	36564480	2094384	529212
$\lambda_{24}$	52452096	14736768	10062240	34527840	144317952	88473216	189008640	82394880	19656384	12217728
$\lambda_{25}$	173882880	109458432	96609024	144317952	269266176	208760832	323066880	217232640	122595840	103799808
$\lambda_{26}$	113431680	59635968	50216832	88473216	208760832	149380608	259614720	151741440	69357312	55273728
$\lambda_{27}$	224570880	146085120	128977920	189008640	323066880	259614720	380816640	274648320	163296000	138620160
$\lambda_{28}$	111989760	47696640	36564480	82394880	217232640	151741440	274648320	157002240	59082240	42251520
$\lambda_{29}$	33873792	4562088	2094384	19656384	122595840	69357312	163296000	59082240	7227744	2992704
$\lambda_{30}$	22662144	1696248	529212	12217728	103799808	55273728	138620160	42251520	2992704	858964
$\lambda_{31}$	17628864	845324	177668	9098832	93987072	48285696	125452800	34218240	1604760	322976
$\lambda_{32}$	45871008	10992384	7072288	29190720	136832256	81780096	180195840	74215680	15152928	8804144
$\lambda_{33}$	29095296	4115744	2140864	16857120	112909824	62584320	150128640	51194880	6258240	2881408
$\lambda_{34}$	81255744	31848480	24312672	58146048	179071488	118874880	229547520	117849600	39694752	28099584
$\lambda_{35}$	54028800	11355264	6352128	33866496	152239104	92799360	201139200	87736320	16608384	8449920
$\lambda_{36}$	20966976	1254496	297368	11094080	101090304	53230848	134991360	39705600	2326656	531624
$\lambda_{37}$	16495872	682704	118388	8417920	91657728	46665984	122238720	32348160	1329144	282012
$\lambda_{38}$	38966976	6556128	3402288	23387376	129238272	74884992	171590400	66055680	9916080	4642992
$\lambda_{39}$	26796672	3034584	1332756	15110736	109840896	60064128	146292480	48142080	4887600	1918056

	$\lambda_{31}$	$\lambda_{32}$	$\lambda_{33}$	$\lambda_{34}$	$\lambda_{35}$	$\lambda_{36}$	$\lambda_{37}$	$\lambda_{38}$	$\lambda_{39}$
$\lambda_1$	183121920	232657920	205655040	276480000	251596800	192199680	180126720	225054720	202245120
$\lambda_2$	64434240	106981920	82964160	150465600	121875840	71055360	62232480	99004320	79833600
$\lambda_3$	7825680	31490400	16991040	64491840	37313280	10046880	7044960	24805440	14928480
$\lambda_4$	785184	13092576	4849920	36772128	14054400	1273920	581664	8013312	3561984
$\lambda_5$	101592	7648560	2253600	26098848	6943104	198048	43704	3656160	3611136
$\lambda_6$	29822	5640672	1486784	21407328	4503168	55904	5694	2289504	789948
$\lambda_7$	17280	4838400	1209600	19353600	3628800	30240	2160	1814400	604800
$\lambda_8$	60852	6269376	1744056	22766976	5326848	117144	23510	2759136	993696
$\lambda_9$	420692	9679568	3295576	29768256	9694080	694728	300392	5386368	2276868
$\lambda_{10}$	166672	7059296	2127960	24300000	6350976	284088	107758	3393840	1320864
$\lambda_{11}$	4057860	20798256	10187232	47684160	23837184	5339088	3601860	15251184	8629560
$\lambda_{12}$	1468100	12513072	5144832	34102080	13286016	2029112	1258932	7964928	4008432
$\lambda_{13}$	428652	8054880	2701152	26024832	7615872	636624	341066	4245936	1838640
$\lambda_{14}$	2111844	14307584	6313696	36765504	15634944	2847288	1843456	9629280	5103384
$\lambda_{15}$	33130848	64718688	46364160	101408256	74620800	37450368	31700928	57751776	43913952
$\lambda_{16}$	11729184	32587152	19874784	61969248	37613952	13938816	10988856	26730960	18091512
$\lambda_{17}$	3251596	16345568	7876864	39464928	18058752	4168160	2929372	11561520	6611700
$\lambda_{18}$	771600	9287520	3430720	28053504	9239040	1092280	645916	5340288	2505792
$\lambda_{19}$	6059208	21236736	11676352	45759168	23968512	7383336	5608648	16268256	10290600
$\lambda_{20}$	17058936	40084896	26291712	70604352	46244736	19768752	16158480	34126176	24394272
$\lambda_{21}$	17628864	45871008	29095296	81255744	54028800	20966976	16495872	38966976	26796672
$\lambda_{22}$	845324	10992384	4115744	31848480	11355264	1254496	682704	6556128	3034584
$\lambda_{23}$	177668	7072288	2140864	24312672	6352128	297368	118388	3402288	1332756
$\lambda_{24}$	9098832	29190720	16857120	58146048	33866496	11094080	8417920	23387376	15110736
$\lambda_{25}$	93987072	136832256	112909824	179071488	152239104	101090304	91657728	129238272	109840896
$\lambda_{26}$	48285696	81780096	62584320	118874880	92799360	53230848	46665984	74884992	60064128
$\lambda_{27}$	125452800	180195840	150128640	229547520	201139200	134991360	122238720	171590400	146292480
$\lambda_{28}$	34218240	74215680	51194880	117849600	87736320	39705600	32348160	66055680	48142080
$\lambda_{29}$	1604760	15152928	6258240	39694752	16608384	2326656	1329144	9916080	4887600
$\lambda_{30}$	322976	8804144	2881408	28099584	8449920	531624	228012	4642992	1918056
$\lambda_{31}$	76360	6271344	1757552	22752288	5300352	138008	38334	2758608	1007208
$\lambda_{32}$	6271344	24146240	12973696	51653952	27668736	7839456	5731696	18505056	11352048
$\lambda_{33}$	1757552	12973696	5530880	34602912	13800960	2368040	1534416	8413056	4383360
$\lambda_{34}$	22752288	51653952	34602912	87158304	60018048	26369280	21546720	44713152	32272992
$\lambda_{35}$	5300352	27668736	13800960	60018048	33041664	7168896	4625280	21125376	11822976
$\lambda_{36}$	138008	7839456	2368040	26369280	7168896	256784	72328	3819456	1460880
$\lambda_{37}$	38334	5731696	1534416	4625280	72328	11002	2368224	830520	830520
$\lambda_{38}$	2758608	18505056	8413056	44713152	21125376	3819456	2368224	13699968	6915312
$\lambda_{39}$	1007208	11352048	4383360	32272992	11822976	1460880	830520	6915312	3292116

## Acknowledgments

The author is indebted to C. Reutenauer for his invaluable contribution during this work. He also thanks M. Schocker for the many hours passed on the double coset conjecture; and R.B Howlett for our useful discussions. Finally, he is grateful to F. Lübeck for his helpful remarks about CHEVIE and GAP.

## References

- [1] F. Bergeron, N. Bergeron, R. B. Howlett and D. E. Taylor: *A decomposition of the descent algebra of a finite Coxeter group.*, J. Algebraic Combin., 1992, **1**, p.23-44.
- [2] N. Bergeron: *A Decomposition of the Descent Algebra of the Hyperoctahedral Group II*, J. Algebra, 1992, **148**, p.98-122.
- [3] N. Bourbaki: *Groupes et algèbres de Lie*, Chap. 4-6, Hermann, 1968.
- [4] P. Fleischmann: *On pointwise conjugacy of distinguished coset representatives in Coxeter groups*, J. Group Theory, 2002, **5**, p.269 -283.
- [5] W. Fulton, J. Harris: *Representation theory, a first course*, Springer, 1999, **129**.
- [6] A. M. Garsia, C. Reutenauer: *A decomposition of Solomon's descent algebra*, Adv. in Math., 1989, **77**, p.189-262.
- [7] M. Geck, G. Hiss, F. Lübeck, G. Malle, and G. Pfeiffer. *CHEVIE – A system for computing and processing generic character tables for finite groups of Lie type, Weyl groups and Hecke algebras.* *Appl. Algebra Engrg. Comm. Comput.*, 7:175–210, 1996.
- [8] M. Geck, G. Pfeiffer: *Characters of Finite Coxeter Groups and Iwahori-Hecke Algebras*, London Math. Soc. Mon. New Series, 2000, **21**.
- [9] I. Gessel: *Multipartite P-partitions and inner product of skew Schur functions*, 1984, Contemp. Math., **34**, p.289-301.
- [10] I. Gessel, C. Reutenauer: *Counting permutations with given cycle structure and descent set*, 1993, J. Combin. Theory Ser. A, **64**, p.189-215.
- [11] C. Hohlweg, M. Schocker: *On Deodhar's Lemma and a parabolic symmetry of finite Coxeter groups*, Preprint Institut de Recherche Mathématique avancée (IRMA), 2002.
- [12] J.E. Humphreys: *Reflection groups and Coxeter groups*, 1990, Cambridge university press, **29**.
- [13] A. Jöllenbeck, C. Reutenauer: *Eine Symmetrieeigenschaft von Solomons Algebra und der höheren Lie-Charaktere*, Abh. Math. Sem. Univ. Hamburg, **71**, 2001, P.105-111.



- [14] W. Ledermann: *Introduction to group characters*, 1977, Cambridge university press.
- [15] S. Poirier: *Cycle type and descent set in wreath products*, 1998, Discrete Math., **180**, p.315-343.
- [16] C. Reutenauer: *Free Lie algebras*, 1993, London Math. Soc. Mon. New Series, **7**.
- [17] Martin Schönert et al. *GAP – Groups, Algorithms, and Programming*. Lehrstuhl D für Mathematik, Rheinisch Westfälische Technische Hochschule, Aachen, Germany, fifth edition, 1995.
- [18] L. Solomon: *A Decomposition of the Group Algebra of a finite Coxeter group*, 1968, J. Algebra, **9**, No 2, p.220-239.
- [19] L. Solomon: *A Mackey formula in the group ring of a Coxeter group*, 1976, J. Algebra, **41**, p.255-268.