

# Ramified Partition Algebras

P P Martin and A Elgamal\*

Mathematics Department, City University,  
Northampton Square, London EC1V 0HB, UK.

## Abstract

For each natural number  $n$ , poset  $T$ , and  $|T|$ -tuple of scalars  $\mathbf{Q}$ , we introduce the ramified partition algebra  $P_n^{(T)}(\mathbf{Q})$ , which is a physically motivated and natural generalization of the partition algebra [24, 25] (the partition algebra coincides with case  $|T| = 1$ ). For fixed  $n$  and  $T$  these algebras, like the partition algebra, have a basis independent of  $\mathbf{Q}$ . We investigate their representation theory in case  $T = \underline{2} := (\{1, 2\}, \leq)$ . We show that  $P_n^{(2)}(\mathbf{Q})$  is quasi-hereditary over field  $k$  when  $Q_1 Q_2$  is invertible in  $k$  and  $k$  is such that certain finite group algebras over  $k$  are semisimple (e.g. when  $k$  is algebraically closed, characteristic zero). Under these conditions we determine an index set for simple modules of  $P_n^{(2)}(\mathbf{Q})$ , and construct standard modules with this index set. We show that there are unboundedly many choices of  $\mathbf{Q}$  such that  $P_n^{(2)}(\mathbf{Q})$  is *not* semisimple for sufficiently large  $n$ , but that it is generically semisimple for all  $n$ .

We construct tensor space representations of certain non-semisimple specializations of  $P_n^{(2)}(\mathbf{Q})$ , and show how to use these to build clock model transfer matrices [24] in arbitrary physical dimensions.

## 1 Introduction

For  $k$  a ring and  $Q \in k$  the partition algebras  $P_n(Q)$  ( $n = 1, 2, \dots$ ) are a tower of finite dimensional unital  $k$ -algebras. These algebras first arose in the context of transfer matrix algebras in Statistical Mechanics [24, 25], but are also known to play roles in invariant theory [21, 28] analogous to that of the Brauer algebra [4, 35, 19], and in the study of Schur algebras (see [20, 30] and cf. [11, §4]). They have a rich and (for  $k = \mathbb{C}$ ) tractable representation theory [26, 36]. The ramified partition algebras  $P_n^{(T)} = P_n^{(T)}(\mathbf{Q})$  are a generalisation depending on a poset  $T$ , which is again physically motivated (see below), but also natural as an abstract extension, as we shall see. In this paper we construct these new algebras (in §2), and describe the representation theory of the simplest non-trivial cases (§3).

An important tool in our approach is *quasi-heredity* [8, 9, 10]. When applicable this provides a crucial organizational scheme for an algebra, via constructions for a set of *standard modules* (the set of *heads* of these modules constituting a complete set of simple modules [10, §A1]). When, as here, quasi-heredity ‘commutes’ with a

---

\*Sadly Ahmed died before this work was completed. His memory lives on.

globalisation functor [16, §6.2] [5] on a tower of algebras we have a singularly powerful representation theoretic tool, even when the algebras are semisimple (indeed in semisimple cases it typically *reduces* to a glorified Jones basic construction [15, 18]). Let  $P_n^{(T)}\text{-mod}$  denote the category of left  $P_n^{(T)}$ -modules. We may suppose without loss of generality, as we shall see, that the underlying set of the poset  $T$  is  $\{1, 2, \dots, d\}$ . Write  $\mathbf{Q} = (Q_1, Q_2, \dots) \in k^d$  and set  $Q^\pi = \prod_{t \in T} Q_t \in k$ . We shall show that for  $Q^\pi$  invertible in  $k$  there exists an idempotent  $e^T \in P_n^{(T)}$  such that  $e^T P_n^{(T)} e^T \cong P_{n-1}^{(T)}$ . It follows that there are functors  $\mathcal{F} : P_n^{(T)}\text{-mod} \rightarrow P_{n-1}^{(T)}\text{-mod}$  and  $\mathcal{G} : P_{n-1}^{(T)}\text{-mod} \rightarrow P_n^{(T)}\text{-mod}$ , called *localisation* and *globalisation* respectively, which port information between layers in the tower. Taken in conjunction with quasi-heredity this provides the framework for an iterative mechanism of analysis (with the base case  $n = 1$  being amenable to brute force).<sup>1</sup>

This paper is concerned with the *representation theory* of the ramified partition algebras. This *requires* no reference to, nor interest in, the physical applications, but a remark on the authors' guiding motivation in constructing and studying these algebras is in order (see §4.1 for a more detailed exposition). The guiding motivation for this work is the study of Statistical Mechanical lattice models [2, 32] of three dimensional physical systems by exact computation [3, 33]. There has been some progress in this study using the ordinary partition algebra  $P_n = P_n(Q)$  [29], which underlies the transfer matrix algebra formalism (see §4.1) for *Potts models* in high dimensions [24, 2]. However, as may be observed from [25, 7] and references therein, there is a problem with the ordinary partition algebra approach in descending in dimension to a physical lattice capturing 3d (or indeed any finite  $d > 2$ ) spatial geometry. We suggest that a better foundation would be provided by a more general algebra of *clock models* [13, 12] or similar models [1, 14]. That is, one in which the set of *spin configurations* is equipped, though the form of the interaction Hamiltonian (see §4.1), with a richer structure than the discrete topology (which is the structure effectively provided by the Potts Hamiltonian). Our reasoning is as follows. The defining basis of the partition algebra is built from the set  $N$  of spatial lattice sites (with  $|N| = n$ ) by forming the set of partitions of  $N$ . A partition forms a base for a rather crude topology on  $N$  (a *partition topology*). In the partition algebra this constrains the spatial geometry *and* the spin interactions to be very crude — and in this sense the two limitations are linked. Accordingly we look here at a generalization of the partition algebra suitable for the interactions in clock models. The idea is that the *set* partitions of  $P_n(Q)$  should be glorified with a structure interpolating between the discrete topology and the metric topology of 3d Euclidean space. The challenge is that this structure not destroy the property of associative algebra.

The ordinary partition algebra is the case  $T = \{1\}$ , however the representation theory of  $P_n^{(T)}$  is vastly richer than that of  $P_n$  in general, as we will show. The simplest interesting new case is  $T = \underline{2} := (\{1, 2\}, 1 \leq 2)$ . In this paper we concentrate mainly on this case. We determine conditions for it to be quasi-hereditary, and construct its standard modules. We show that  $P_n^{(2)}(\mathbf{Q})$  is generically semisimple, but (in §4) we also exhibit, for certain values of  $\mathbf{Q}$ , *special* representations  $R$  which may be used to show non-semisimplicity at these values of  $\mathbf{Q}$ . It is also these

---

<sup>1</sup>Both globalisation and quasi-heredity apply to the ordinary partition algebra, and are key tools in its analysis [26]. The utility of globalisation is a natural consequence of the physical context of these algebras (cf. [27]), but it *is* intriguing to see how often quasi-heredity also occurs in this setting [6, 26, 36] (in as much as a physical interpretation is so far lacking).

representations which may be used in clock model transfer matrix computations (cf. [24]). Results of these computations for clock models will be presented in a separate paper, but to underline the motivation we include a small sample of such results here — zeros of the partition function for  $Z_Q$ -symmetric models (recall that zeros converge on the real axis at a phase transition point [24, 31]).

The representation  $R$  is also the obvious generalization of the ‘Potts’ representation of  $P_n$  which arises in invariant theory [24, 21, 30]. An analysis of this, and of the generic representation theory of  $P_n^{(T)}$ , is the first step in seeking for  $P_n^{(T)}$  a role in invariant theory analogous to that of  $P_n$ .

## 2 Preliminaries

Let  $(T, \leq)$  be a poset of degree  $d$ ,  $k$  a field,  $Q_t \in k$  for each  $t \in T$ , and  $k$ -algebra  $P_n(Q_t)$  the ordinary partition algebra. The  $T$ -ramified partition algebra will be defined to be a certain subalgebra of the tensor product  $\bigotimes_{t \in T} P_n(Q_t)$ , depending on  $\leq$ .

### 2.1 Construction of $P_n^{(T)}$

First we recall the definition of  $P_n(Q)$  [26, §1]. Given a set  $U$  we write  $\mathbf{E}_U$  for the set of partitions of  $U$  (we identify this with the set of equivalence relations on  $U$ , writing  $\sim^a$  for the relation corresponding to  $a \in \mathbf{E}_U$ ). Note that any bijection  $f : U \rightarrow V$  naturally underlies a corresponding bijection  $\mathbf{E}^f : \mathbf{E}_U \rightarrow \mathbf{E}_V$ .

For  $a \in \mathbf{E}_U$  we write  $q \in a$  for  $q \subseteq U$  a part of  $a$  as a partition. For  $S \subset U$  write  $a|_S$  for the restriction of  $a$  to  $S$ . Given  $n \in \mathbb{N}$  we write  $\underline{n} = \{1, 2, \dots, n\}$ ,  $\underline{n}' = \{1', 2', \dots, n'\}$ , and so on. We set  $p_n = \mathbf{E}_{\underline{n} \cup \underline{n}'}$ , and  $kp_n$  the  $k$ -space with basis  $p_n$ .

Let  $f : \underline{n} \cup \underline{n}' \rightarrow \underline{n}' \cup \underline{n}''$  be the map  $i \mapsto i'$ . Then given  $a, b \in p_n$  we define  $\mathbf{d}_u(ab) \in \mathbf{E}_{\underline{n} \cup \underline{n}' \cup \underline{n}''}$  by identifying with the equivalence relation generated by  $a$  and  $\mathbf{E}^f(b)$ . Let  $g : \underline{n} \cup \underline{n}'' \rightarrow \underline{n} \cup \underline{n}'$  be the map  $i \mapsto i$  for  $i \in \underline{n}$  and  $i'' \mapsto i'$  for  $i'' \in \underline{n}''$ . Then define  $\mathbf{d}(ab) = \mathbf{E}^g(\mathbf{d}_u(ab)|_{\underline{n} \cup \underline{n}''}) \in p_n$ . Fixing  $Q \in k$  we now define the product  $ab \in kp_n$  by

$$ab = Q^{c(a,b)} \mathbf{d}(ab) \tag{1}$$

where  $c(a, b) = \#\{q \subseteq \underline{n}' : q \in \mathbf{d}_u(ab)\}$ . In [26, §1] it is shown that this product ( $k$ -linearly extended) endows  $kp_n$  with the structure of a unital associative  $k$ -algebra, which we call the partition algebra, and denote by  $P_n(Q)$ .

Now fix a poset  $(T, \leq)$  with  $T = \{1, 2, \dots, d\}$  but  $\leq$  arbitrary, and an element  $\mathbf{Q} = (Q_1, Q_2, \dots, Q_d) \in k^d$ . Our aim is to define the  $T$ -ramified partition algebra as a subalgebra of  $P_n^{(d)} = \bigotimes_{t \in T} P_n(Q_t)$ , depending on the partial order  $\leq$ . We will often concentrate on the special case  $T = \{1, 2\}$  with  $1 \leq 2$  (and where a poset is intended,  $\underline{2}$  will denote this one, and  $\underline{d}$  the natural ordered generalization), although our construction is general.

Given  $a, b \in \mathbf{E}_U$  we say that  $a$  is a *refinement* of  $b$  (see e.g. [22, ch.4 §7]), denoted  $a \leq b$ , if each part of  $b$  is a union of one or more parts of  $a$ ; i.e. if  $i \sim^a j$  implies  $i \sim^b j$ . It is easy to show that

**Proposition 1** *Let  $a, b \in \mathbf{E}_U$  and  $U' \subseteq U$ . Then  $a \leq b$  implies  $a|_{U'} \leq b|_{U'}$ .*

For  $S = \mathbf{E}_U$  we define  $S^{(T)}$  to be the subset of  $\prod_T S$  given by the elements  $a = (a_t : t \in T)$  for which  $t \leq t'$  implies  $a_t$  is a refinement of  $a_{t'}$ . We call any such  $a \in S^{(T)}$  a  $T$ -ramified partition. For example  $(\{\{1\}, \{2\}\}, \{\{1, 2\}\}) \in \mathbf{E}_{\{1,2\}}^{(2)}$ .

**Proposition 2** *For any  $d$ -tuple  $\mathbf{Q} = (Q_1, \dots, Q_d) \in k^d$ , the set  $p_n^{(T)}$  is a basis for a subalgebra of  $\bigotimes_{t \in T} P_n(Q_t)$ .*

*Proof:* It is enough to show that if  $a, b \in p_n^{(T)}$  then  $\mathbf{d}(a_t b_t)$  is a refinement of  $\mathbf{d}(a_{t'} b_{t'})$  if  $t \leq t'$ . But if  $t \leq t'$  then  $(a_t, a_{t'})$  and  $(b_t, b_{t'})$  both appear in  $p_n^{(2)}$  (where we identify  $\underline{2} = \{1, 2\}$  with  $\{t, t'\}$ ) so we are done if we can show for  $\{(a, b), (c, d)\} \subseteq p_n^{(2)}$  that  $\mathbf{d}(ac)$  a refinement of  $\mathbf{d}(bd)$ . Now by proposition 1 it is enough to show that  $\mathbf{d}_u(ac) \leq \mathbf{d}_u(bd)$ . Suppose  $i \sim^{\mathbf{d}_u(ac)} j$  and that the connection is established through some sequence of connections alternately within  $a$  and  $c$ . Each connection within  $a$  is within  $b$ , each connection within  $c$  is within  $d$ , so  $i \sim^{\mathbf{d}_u(bd)} j$ .  $\square$

Fixing  $\mathbf{Q}$  as above we denote the  $k$ -algebra with basis  $p_n^{(T)}$  by  $P_n^{(T)} = P_n^{(T)}(\mathbf{Q})$ . This is the  $T$ -ramified partition algebra.

Two posets are isomorphic if there is a bijective homomorphism between the underlying sets which preserves the partial order. It will be evident that the construction of  $P_n^{(T)}$  depends on the partial order, but not on the names of the elements of  $T$ . Thus  $P_n^{(T)}$  is defined for poset  $T$  with *any* underlying set, and  $T \cong T'$  implies  $P_n^{(T)} \cong P_n^{(T')}$ .

Not all choices of  $T$  are *necessarily* physically interesting, in the sense of the motivation discussed in the introduction. From an abstract point of view, as we will see later, the choice of  $T$  as the poset of divisors of a natural number gives rise to some interesting representations. But we will concentrate largely on  $T = \underline{2}$ , which is among the main motivating examples, captures much of the flavour of the general case (consider the proof of proposition 2), and already has a very rich structure.

Refinement equips any  $\mathbf{E}_U$ , and in particular  $p_n$ , with the property of lattice. The bottom element  $\{U\}$  is called the *trivial* partition.

## 2.2 Examples and notations

Putting  $T = \{1\}$  we recover the partition algebra itself. Recall from [26, §1] the special elements in basis  $p_n$  denoted  $1$ ,  $A^i$ , and  $A^{ij}$  as follows:

$$1 = \{\{1, 1'\}, \{2, 2'\}, \dots, \{i, i'\}, \dots, \{n, n'\}\}$$

$A^i$  differs from  $1$  by  $\{i, i'\} \rightsquigarrow \{i\}, \{i'\}$ ; and  $A^{ij}$  differs from  $1$  by  $\{i, i'\}, \{j, j'\} \rightsquigarrow \{i, i', j, j'\}$  (see also the inner components of figure 2 later). Thus  $1 \in p_1$  is  $\{\{1, 1'\}\}$  and  $A^1 \in p_1$  is  $\{\{1\}, \{1'\}\}$ . See [26, §1] also for the partition algebra *diagram calculus*.

The next simplest case is  $T = \underline{2}$ . An example of the  $\underline{2}$ -ramified partition algebra product is illustrated in figure 1 (the formal interpretation of this diagram is given shortly). The basis  $p_1^{(2)}$  consists of three elements:  $(1, 1) = (\{\{1, 1'\}\}, \{\{1, 1'\}\})$ ;  $(A^1, 1) = (\{\{1\}, \{1'\}\}, \{\{1, 1'\}\})$ ; and  $(A^1, A^1) = (\{\{1\}, \{1'\}\}, \{\{1\}, \{1'\}\})$ . Note that  $\{(A^1, A^1)\}$  is a basis for an ideal, as is  $\{(A^1, A^1), (A^1, 1)\}$  — see later.

It is convenient to introduce a ‘serial’ shorthand notation and also a diagram calculus for  $P_n^{(d)}$ . The shorthand for a  $\underline{2}$ -ramified partition is to write out the more refined (‘inner’) partition and then to group the parts to indicate the less




Figure 1: The algebra product in  $P_3^{(2)}$  in an example:  $ab = Q_1 c$ . Diagrams for  $a$  and  $b$  (drawn one above the other on the left) are juxtaposed to overlay and identify the vertices indicated by dotted lines, and these vertices are then removed in exchange for an appropriate scalar factor (cf. [26, §1]).

refined ('outer') partition by using another nest of brackets. Thus in  $p_1^{(2)}$ ,  $(1, 1) = \{\{\{1, 1'\}\}\}$ ,  $(A^1, 1) = \{\{\{1\}, \{1'\}\}\}$ , and  $(A^1, A^1) = \{\{\{1\}\}, \{\{1'\}\}\}$ . There is an obvious generalization for  $T$  any chain.

The diagram calculus applies the same principle to partition algebra diagrams (in these the elements of  $\underline{n}, \underline{n}'$  are represented as two rows of vertices arranged with  $i'$  vertically below  $i$ , so composition involves vertical juxtaposition). Inner parts are indicated by 'hard wiring' vertices together as in ordinary partition algebra diagrams. Outer parts are indicated by drawing 'islands' around clusters of connected vertices. See figure 1 for some examples. Note by equation (1) that inner (outer) clusters *isolated from the exterior in composition* are replaced by a factor  $Q_1$  (resp.  $Q_2$ ). As for  $P_n(Q)$ , there will not be a strictly planar representation in general. Consider also the elements in figure 2, and note that

$$(A^i, A^i)(A^{ij}, A^{ij})(A^i, A^i) = (A^i, A^i) \quad (A^{ij}, A^{ij})(A^i, A^i)(A^{ij}, A^{ij}) = (A^{ij}, A^{ij}). \quad (2)$$

It will be evident from the diagrams that  $P_n^{(2)}$  is isomorphic to its opposite (indeed this is clearly true for any  $T$ ). We denote the image of an element  $a$  under this isomorphism by  $a^\circ$  (as a diagram it is reflected top to bottom).

### 2.3 Canonical inclusions

Let  $S_n$  denote the symmetric group on  $\underline{n}$ . By  $S_n \hookrightarrow P_n$  we denote the natural monomorphism (and here, as usual, we write  $\sigma_i$  for the 2-cycle  $(i \ i+1) \in S_n$ , and write  $\sigma_{ij}$  for the 2-cycle  $(ij)$ ).

For  $t \in T$  define

$$\begin{aligned} \text{pr}_t : p_n^{(T)} &\rightarrow p_n \\ (a_1, a_2, \dots, a_d) &\mapsto a_t. \end{aligned}$$

Since  $a \in p_n$  is a refinement of itself we have




Figure 2: The multiplicative unit in  $P_4^{(2)}$  together with various other elements. In these examples  $i = 2$  and  $j = 3$ . Note that these ‘diagrams’ do not necessarily embed in the plane in any natural way, and that care must be taken with interpretation of ‘crossing’ lines. Over/under information is not relevant, but may appear in diagrams as a guide to the eye.

**Proposition 3** *The diagonal inclusion*

$$\begin{aligned} D^{(T)} : p_n &\rightarrow p_n^{(T)} \\ a &\mapsto (a, a, \dots, a) \end{aligned}$$

(or simply  $D$ , where no ambiguity arises) extends to an algebra morphism

$$D^{(T)} : P_n(Q^\pi) \hookrightarrow P_n^{(T)}(\mathbf{Q}).$$

For example, set  $e^T = D^{(T)}(A^n)$ , then  $e^T e^T = Q^\pi e^T$ .

There are various ways to inject  $P_{n-1}^{(T)}$  into  $P_n^{(T)}$ . We treat as inclusion that which appends to every basis element a part which is  $\{n, n'\}$  at every level of  $T$ . For example, in case  $T = \underline{2}$  the map on basis elements is given explicitly by

$$a \mapsto a \cup \{\{\{n, n'\}\}\}.$$

NB, including  $P_{n-1}^{(T)}$  in  $P_n^{(T)}$  in this way means that  $e^T P_{n-1}^{(T)} = P_{n-1}^{(T)} e^T$ , and hence that  $e^T P_n^{(T)}$  is a left  $P_{n-1}^{(T)}$ -module.

By  $p_n \xrightarrow{I_n} p_n^{(\underline{2})}$  we denote the *inner* inclusion of  $p_n$ , i.e. where the inner partition in  $In(p)$  is  $p$  and the outer partition is trivial.

Note that the serial notation for a  $\underline{2}$ -ramified partition is unique if written out so that at each step the numerically lowest possible element not yet written is written next (for which purpose take  $i' = n + i$ ).

Let us now fix the case  $T = \underline{2}$  until further notice.

### 3 Structure of $P_n^{(\underline{2})}$

### 3.1 Filtration by ideals

For a partition  $a \in p_n$  (or any restriction thereof) let  $\pi(a)$  be the set of *propagating* parts of  $a$ , i.e. those intersecting both  $\underline{n}$  and  $\underline{n}'$ . Then the *propagating number*  $\#^P(a) := |\pi(a)|$  (see [26]). For  $a \in p_n^{(2)}$  let  $\#^P(a) \in \mathbb{N}_0 \times \mathbb{N}_0$  denote  $(\#^P(a_1), \#^P(a_2))$ , and similarly for any  $T$ .

For  $a \in p_n^{(2)}$  let the *propagating index*  $\#^-(a)$  be the list of  $\#^P(a_1|_q)$  over all  $q \in \pi(a_2)$  arranged so that  $\#^-(a)_i \geq \#^-(a)_{i+1}$ . For example, the three  $\underline{2}$ -ramified partitions illustrated in figure 1 have  $\#^-(a) = (1, 1)$ ,  $\#^-(b) = (1, 0, 0)$ ,  $\#^-(c) = (1, 0)$ . Note that the values taken by the propagating index differ from integer partitions only in that there may be zero parts.

A generalization of the propagating index may be formulated for other choices of  $T$ , but we will not pursue the point here.

For  $\lambda$  a propagating index we denote by  $p_n^{(2)}(\lambda)$  the subset of elements  $a \in p_n^{(2)}$  such that  $\#^-(a) = \lambda$ .

The *propagating envelope*  $\#^{env}(a)$  is  $\sum_i f(\#^-(a)_i)$  where  $f(0) = 1$  and  $f(j) = j$  otherwise. This number is the minimum  $n$  required so that there is some  $b \in p_n^{(2)}$  with  $\#^-(b) = \#^-(a)$ . For example  $\#^-(\{\{1, 2\}, \{1', 2'\}\}) = (0)$ , so  $\#^{env}(\{\{1, 2\}, \{1', 2'\}\}) = 1$ ; and  $a, b, c$  from figure 1 have  $\#^{env}(a) = 2$ ,  $\#^{env}(b) = 3$ ,  $\#^{env}(c) = 2$ .

NB,  $\#^P$  and  $\#^{env}$  are determined by  $\#^-$  (specifically  $\#^P(a_1) = \sum_i \#^-(a)_i$  and  $\#^P(a_2)$  is the number of (integer) parts of  $\#^-(a)$ ). The basis  $p_n^{(2)}$  is fixed, and  $\#^-$  is invariant, under the opposite isomorphism.

We denote by  $p_n(i)$  the subset of elements  $a \in p_n$  such that  $\#^P(a) = i$ . Recall

**Proposition 4** [26] *Let  $a \in p_n(i)$  and  $b \in p_n(i')$ . Then there exist  $c, d \in p_n$  such that  $a = cbd$  iff  $i \leq i'$ .*

This is a powerful tool in the analysis of  $P_n(Q)$ . The next few propositions build up to a  $T = \underline{2}$  generalization (proposition 8).

**Proposition 5** *The subset  $G_n$  of elements of the types illustrated in figure 2 (or equivalently  $\{(A^1, 1), (A^1, A^1), (1, A^{12}), (A^{12}, A^{12})\} \cup \{(\sigma_i, \sigma_i); i = 1, 2, \dots, n-1\}$ ) generates  $P_n^{(2)}$ .*

*Proof:* Let  $a$  be an element of the defining basis. We construct it as a product of elements of  $G_n$  as follows. Step 1 is to construct an element of  $p_n^{(T)}$  with ‘enough’ propagating connections to act as a base for the construction of  $a$ . For non-empty  $q \subseteq \underline{n}$ ,  $q' \subseteq \underline{n}'$  let  $l(q \cup q')$  (resp.  $l'(q \cup q')$ ) be the numerically lowest unprimed (resp. primed) element of the argument. For  $a \in p_n^{(2)}$  let  $\pi_1(a) = \pi(a_1)$  and  $\pi_2(a) = \{q \in \pi(a_2) \mid q \cap r = \emptyset \ \forall r \in \pi(a_1)\}$ . Let  $\mathcal{X}_t(a) = \cup_{q \in \pi_t(a)} \{l(q), l'(q)\}$ , and  $\mathcal{X}^o(a) = \mathcal{X}_1(a) \cup \mathcal{X}_2(a)$  — the union of the set of low numbered pairs for inner propagating parts of  $a$  and the set of low numbered pairs for outer propagating parts containing no inner propagating parts. Now consider the set of basis elements of the form  $w \in S_n \hookrightarrow P_n \xrightarrow{D} P_n^{(T)}$ , i.e. those generated by the  $(\sigma_i, \sigma_i)$ s. Note that there is always at least one such  $w$  which restricts, on  $\mathcal{X}^o(a)$ , to the same pairings as  $a$  does (albeit diagonally). Choose any one such  $w$ . Step 2: Apply  $(A^i, A^i)$ s so as to cut all the connections *not* on  $\mathcal{X}^o(a)$ . Step 3: Take the resultant basis element and make all the additional outer connections required for  $a$ , i.e. those *within*  $\underline{n}$  and those within  $\underline{n}'$ , using  $(1, A^{ij})$ s from the left and right as appropriate. Step 4: Make the additional inner connections for  $a$  using  $(A^{ij}, A^{ij})$ s as appropriate (NB, these will




Figure 3: The special element  $\mathcal{I}_\lambda$  of  $p_{14}^{(2)}((3, 2^2, 1^3, 0))$ .

not make any further *outer* connections, cf. step 3 and the definition of  $p_n^{(2)}$ . Finally make the required inner disconnections, i.e. between those pairs in  $\mathcal{X}^o(a)$  referring to outer but not inner propagating parts, using  $(A^i, 1)$ s.  $\square$

**Proposition 6** For  $a, b \in p_n^{(T)}$

(i)  $\#^P(\mathbf{d}(ab))_t \leq \min(\#^P(a)_t, \#^P(b)_t) \forall t \in T$ .

(ii) If  $\#^P(a)_t > 0$  and  $t \leq t'$  then  $\#^P(a)_{t'} > 0$ .

And for  $T = \underline{2}$

(iii) Let  $a, b$  be such that  $\#^P(\mathbf{d}(ab))_2 = \#^P(a)_2$ . Then  $\#^-(\mathbf{d}(ab))_i \leq \#^-(a)_i$  for all suitable  $i$ .

(iv)  $\#^{env}(\mathbf{d}(ab)) \leq \#^{env}(a)$ .

*Proof:* The first claim is elementary in as much as it holds in  $P_n^{(d)}$ , and  $P_n^{(d)} \supseteq P_n^{(T)}$ . The second and third follow from the fact that each propagating part of  $a_1$  is a subset of a propagating part of  $a_2$ . To demonstrate (iv) let  $\#_0^{env}(a)$  be the number of trouser legs (outer propagating parts) containing no legs (inner propagating parts), then  $\#^{env}(a) = \#^P(a_1) + \#_0^{env}(a)$ . The first summand is non-increasing in any product by (i). The second summand may increase, but by (i) case  $t = 2$  if it does so then for each new empty trouser (increase of one) there must be at least one non-empty trouser lost, so the first summand must decrease by at least one.  $\square$

It follows that  $\#^-$  provides a means of filtering ideals of  $P_n^{(2)}$ .

Let us pick out a special element  $\mathcal{I}_\lambda$  of  $p_n^{(2)}(\lambda)$  for each  $\lambda$  as follows. Define  $x_0 = \{\{\{1\}, \{1'\}\}\}$  and

$$x_m = \{\{\{1, 1'\}, \{2, 2'\}, \dots, \{m, m'\}\}\}.$$

Recall the usual Young subgroup construction whereby  $S_n \times S_m \hookrightarrow S_{n+m}$ . We denote by  $a \times b$  the analogous combination of elements  $a \in P_n^{(T)}$  and  $b \in P_m^{(T)}$  into an element of  $P_{n+m}^{(T)}$ . Then

$$\mathcal{I}_\lambda = (\times_i x_{\lambda_i}) \times \left( \times_{i=1}^{n-\#^{env}(\lambda)} \{\{\{1\}\}, \{\{1'\}\}\} \right)$$

where  $\#^{env}(\lambda)$ , resp.  $\#^P(\lambda)$ , denotes the  $\#^{env}$ , resp.  $\#^P$ , of any  $a$  with  $\#^-(a) = \lambda$ . An example is given in figure 3.

Let  $\Lambda$  denote the set of possible propagating indices  $\lambda$ . We write  $\bar{\lambda} = (\bar{\lambda}_1^1, \bar{\lambda}_2^2, \dots)$  for  $\lambda \in \Lambda$  expressed in the exponent notation (i.e. a repeated component is replaced by an exponent recording the number of repeats, thus  $(1, 1, 1) = (1^3)$  and so on).

Note that for each  $\lambda \in \Lambda$  there is an  $m \in \mathbb{N}_0$  such that  $\lambda \in \mathbb{N}_0^m$  ( $\mathbb{N}_0^0$  is to be understood as the empty list). Each  $\mathbb{N}_0^m$  has a natural action of  $S_m$  upon it (permuting the indices), and each  $S_m$  orbit in  $\mathbb{N}_0^m$  has a unique element which is a






Figure 4: The bottom of  $(\Lambda, \leq)$ .

$\lambda$ . This is called the *dominant* element. If we equate a  $\lambda$  with an element  $\mu$  of  $\mathbb{N}_0^m$  we will always intend the dominant element in the orbit of  $\mu$ .

We define a partial order  $\leq$  on  $\Lambda$  as follows. First write a relation  $\lambda' \rho \lambda$  if

(0)  $\lambda' = \lambda$ ; or

(i)  $\lambda' = (\lambda_1, \lambda_2, \dots, \lambda_{i-1}, \lambda_i - 1, \lambda_{i+1}, \dots)$ ; or

(ii)  $\lambda' = (\lambda_1, \lambda_2, \dots, \lambda_{i-1}, \lambda_i + \lambda_j, \lambda_{i+1}, \dots, \lambda_{j-1}, \lambda_{j+1}, \dots)$ .

Now write  $\lambda' \leq \lambda$  for the transitive completion of this relation (it is manifestly antisymmetric). Note that in case (i)  $\#^P(\lambda) - \#^P(\lambda') = (1, 0)$ ; and in case (ii)  $\#^P(\lambda) - \#^P(\lambda') = (0, 1)$ .

Examples:  $() \leq \lambda$  for all  $\lambda$  and, fixing  $n$ ,  $(1^n) \geq \lambda$  for all  $\lambda$ . The poset is *not* in general a lattice. The Hasse diagram starts as in figure 4.

**Proposition 7** For  $a, b \in p_n^{(2)}$ ,  $\#^-(\mathbf{d}(ab)) \leq \#^-(a)$ .

*Proof:* It is enough to show the result true for  $a$  or  $b$  a generator (as in proposition 5). But multiplication by a generator only implements one of (0)–(ii) above.  $\square$

**Proposition 8** Put  $P = P_n^{(2)}$  and  $p = p_n^{(2)}$ .

(i) Let  $a \in p(\lambda)$  and  $b \in p(\lambda')$ . Then there exist  $c, d \in p$  such that  $a = cbd$  iff  $\lambda \leq \lambda'$ .

(ii)  $\cup_{\lambda' \leq \lambda} p(\lambda')$  is a basis for  $PT_\lambda P$ .

(iii)  $\lambda' \geq \lambda$  if and only if  $PT_{\lambda'} P \supseteq PT_\lambda P$ .

*Proof:* (i) The *only if* part follows from proposition 7.

For each suitable  $i$ , there is an outer part of  $b$  with  $\lambda'_i$  inner propagating parts. For each such  $i$ , consider acting on  $b$  with  $(A^j, A^j)$ s so as to break up inner parts, until only the lowest numbered primed and unprimed elements in each inner propagating part remain connected (i.e., in a total of  $\lambda'_i$  primed/unprimed pairs). Note by proposition 4 that this process is *reversible* (for example if  $j$  was in the same inner part as  $k$  then  $(A^{jk}, A^{jk})$  acting on the left would restore the connection). Then act with  $(\sigma_{ij}, \sigma_{ij})$ s until these lowest numbered elements coincide with the corresponding data for  $\mathcal{I}_{\lambda'}$ . This step is reversible since  $\sigma_{ij}$  is an involution. If  $\lambda = \lambda'$  then  $a$  may also be taken reversibly to  $\mathcal{I}_{\lambda'}$ . To obtain  $c$  and  $d$  simply compose the series of generators taking  $b$  to  $\mathcal{I}_\lambda$  then the reverses which would take  $\mathcal{I}_\lambda$  to  $a$ .

Finally, by definition, if  $\lambda < \lambda'$  then we can move from  $\lambda'$  to  $\lambda$  by a sequence of relations  $\rho$  of types (i) and (ii). Note that  $\mathcal{I}_{\lambda'}$  may be taken to an element of  $p_n(\lambda')$  by a sequence of multiplications by generators which tracks this.

It will be evident that (i) implies (ii) implies (iii).  $\square$

An element  $a$  of a  $k$ -algebra is a *pre-idempotent* if there exists  $\alpha \in k$  such that  $\alpha a$  is idempotent.

For  $\lambda \in \Lambda \setminus \{()\}$  define  $\mathcal{I}'_\lambda \in p_n^{(2)}$  whose inner partition is given by  $\text{pr}_{inner}(\mathcal{I}'_\lambda) = \text{pr}_{inner}(\mathcal{I}_\lambda)$ ; and whose outer partition agrees with  $\text{pr}_{outer}(\mathcal{I}_\lambda)$  except on the tail of non-propagating parts (if any), whose elements are instead included in the last (i.e. rightmost as in figure 3) propagating part.

**Proposition 9** *Again put  $P = P_n^{(2)}$  and suppose  $Q_1$  invertible in  $k$ . Then each  $\mathcal{I}'_\lambda$  is pre-idempotent, and these pre-idempotents commute with each other, and obey*

$$P\mathcal{I}'_\lambda P = P\mathcal{I}_\lambda P.$$

*Proof:* Note that each  $\text{pr}_{inner}(\mathcal{I}'_\lambda)$  may be expressed in the form  $\prod_{j \in J} A^j$  for some  $J$ , and hence is pre-idempotent in  $P_n(Q_1)$  provided that  $Q_1$  invertible. Further, the elements of  $P_n$  of this form commute. Meanwhile  $\text{pr}_{outer}(\mathcal{I}'_\lambda)$  may be expressed in the form  $\prod_{(i,j) \in J} A^{ij}$  for some  $J$ , and hence is idempotent in  $P_n(Q_2)$ . Again, all such elements of  $P_n$  commute with each other. Thus the set of objects  $\mathcal{I}'_\lambda$  are commuting pre-idempotents in  $P_n^{(d)}$ , and hence also in  $P_n^{(2)}$ .

Now  $\mathcal{I}'_\lambda$  differs from  $\mathcal{I}_\lambda$  only in the tail (if any) of pairs of parts of the outer partition of the form  $\{j\}\{j'\}$ . If  $\lambda \neq ()$  then this is preceded by at least one propagating part. Let  $\{\dots (j-1)(j-1)'\}$  be the last such. Then  $A^{(j-1)j}\{\dots (j-1)(j-1)'\}\{j\}\{j'\}A^{(j-1)j} = \{\dots (j-1)(j-1)'jj'\}$  and  $A^j\{\dots (j-1)(j-1)'jj'\}A^j = \{\dots (j-1)(j-1)'\}\{j\}\{j'\}$  establishing the equality of ideals.  $\square$

## 3.2 On heredity

Recall that propagating index determines propagating number. Define  $\Lambda^m \subset \Lambda$  by  $\lambda \in \Lambda^m$  implies  $\#^P(\text{pr}_{inner}(a)) + \#^P(\text{pr}_{outer}(a)) = m$  for any  $a \in p_n^{(2)}(\lambda)$ . For example,  $(0,0), (1) \in \Lambda^2$ .

**Proposition 10** *There exist idempotents  $\mathcal{I}_m$  such that*

- (i)  $P_n^{(2)}\mathcal{I}_m P_n^{(2)} = \sum_{\lambda \in \Lambda^m} P_n^{(2)}\mathcal{I}'_\lambda P_n^{(2)}$  for all  $m$ ; and
- (ii)  $P_n^{(2)}\mathcal{I}_{m+1} P_n^{(2)} \supset P_n^{(2)}\mathcal{I}_m P_n^{(2)}$  for all  $m = 1, \dots, 2n-1$ .

*Proof:* Given a pair  $(e_x, e_y)$ , say, of commuting idempotents in an algebra  $A$  one may construct an orthogonal pair  $(e_x(1-e_y), e_y)$  and hence an idempotent  $e_x - e_x e_y + e_y$ . Clearly  $A(e_x - e_x e_y + e_y)A \supseteq A e_x (e_x - e_x e_y + e_y)A + A e_y (e_x - e_x e_y + e_y)A = A e_x A + A e_y A$  (saturating the inequality) — NB, sums inside and outside of brackets have different meanings here. Applying this process as many times as necessary to the idempotents with  $\lambda \in \Lambda^m$  (noting proposition 9) we obtain our candidate for  $\mathcal{I}_m$  satisfying (i). For (ii), observe that every  $\lambda \in \Lambda^m$  ( $m < 2n$ ) obeys  $\lambda < \lambda'$  for some  $\lambda' \in \Lambda^{m+1}$ .  $\square$

Suppose that  $Q_1 Q_2$  is invertible in  $k$ . Then  $\mathcal{I}_\lambda$  and  $\mathcal{I}'_\lambda$  are pre-idempotent (this being so we will not further labour the distinction between  $\mathcal{I}_\lambda$  and its idempotent form). It is clear in this case that  $\mathcal{I}_\emptyset$  is a heredity idempotent in the sense of [8] (or see [26]).

**Proposition 11**  $P_n^{(2)}$  is quasi-hereditary with heredity chain  $(\mathcal{I}_{2n}, \mathcal{I}_{2n-1}, \dots, \mathcal{I}_1, \mathcal{I}_\emptyset)$ .

*Proof:* Write  $P_i$  for  $P_n^{(2)}/P_n^{(2)}\mathcal{I}_{i-1}P_n^{(2)}$ . Given proposition 10 it remains to show: (i) that subalgebra  $\mathcal{I}_i P_n^{(2)} \mathcal{I}_i$  is semi-simple in  $P_i$ ; and (ii) that  $P_n^{(2)}\mathcal{I}_i \otimes_{\mathcal{I}_i P_n^{(2)} \mathcal{I}_i} \mathcal{I}_i P_n^{(2)} \cong P_n^{(2)}\mathcal{I}_i P_n^{(2)}$  in  $P_i$ . A basis for  $P_i$  is  $\cup_{\lambda \in \Lambda^i} p_n^{(2)}(\lambda)$ . It follows from proposition 7 that




Figure 5: Constructing an element of  $I_\lambda PI_\lambda$  in case  $\lambda = (3, 2^2, 1^3, 0)$  (the middle two rows of points are here to exhibit membership of  $I_\lambda PI_\lambda$  — on ignoring these the diagram becomes an element of the basis  $p_{14}^{(2)}((3, 2^2, 1^3, 0)) \cap I_\lambda PI_\lambda$ ). The shading here is a guide to the eye *only*.

the product of any two basis elements not from the same  $p_n^{(2)}(\lambda)$  is zero in  $P_i$ . For (i), one then notes that the subalgebra  $\mathcal{I}_i P_n^{(2)} \mathcal{I}_i$  will break up as a direct sum with summands of form  $\mathcal{I}_\lambda P_n^{(2)} \mathcal{I}_\lambda$ . The basis of *this* algebra given by its intersection with the given basis of  $P_i$  (illustrated in figure 5) is closed under multiplication, with multiplicative unit  $\mathcal{I}_\lambda$ , and invertible elements. Thus this algebra is isomorphic to a group algebra. (We will identify the group in section 3.3.)

For (ii), one considers the obvious map

$$(aI_i, I_i b) \mapsto aI_i b$$

in terms of basis elements. Again, since the  $\{I_\lambda\}$  are orthogonal (in  $P_i$ ) it is enough to consider  $(aI_\lambda, I_\lambda b) \mapsto aI_\lambda b$ . As written, this requires that the kernel coincides with elements of the form  $(ac, b) - (a, cb)$  where  $c \in \mathcal{I}_i P_n^{(2)} \mathcal{I}_i$ . The kernel clearly includes this set, so it is enough to show that it is no larger. The map is realizable at the level of basis elements, from which the extent of the kernel is immediate, as follows.

Let  $q$  be an element of  $p_n^{(2)}(\lambda) \cap PI_\lambda$  (which set is a basis for  $PI_\lambda$  in  $P_i$ ), and  $a$  an element of  $p_n^{(2)}(\lambda) \cap I_\lambda PI_\lambda$  (similarly a basis for  $I_\lambda PI_\lambda$ ). Up to a scalar  $qa$  is in  $p_n^{(2)}(\lambda) \cap PI_\lambda$  and the map

$$p_n^{(2)}(\lambda) \cap I_\lambda PI_\lambda \rightarrow p_n^{(2)}(\lambda) \cap PI_\lambda$$

given in this way by  $a \mapsto qa$  is injective for any  $q$  (again consider figure 5, for example). Define an equivalence relation on  $p_n^{(2)}(\lambda) \cap PI_\lambda$  by  $q \sim q'$  if  $qa = q'$  for some  $a$  as above. Let  $q^\sim$  be a set of representatives of equivalence classes. Then

each element of  $p_n^{(2)}(\lambda) \cap PI_\lambda$  may be written uniquely in the form  $qa$  where  $q \in q^\sim$  (i.e.  $(q, a) \mapsto qa$  is a bijection).

The same argument works on the right.  $\square$

### 3.3 Left ideals, sections and special modules

Let  $P_n^{(2)}[\lambda]$  denote the  $\lambda$  section in the filtration given by proposition 8(ii) (i.e. with basis  $p_n^{(2)}(\lambda)$ ). Let  $P_n^{(2)}[\lambda]'$  denote the image of  $P_n^{(2)}\mathcal{I}_\lambda$  in this section (cf. [26, §3]). The algebra which is the image of  $\mathcal{I}_\lambda P_n^{(2)}\mathcal{I}_\lambda$  in this section acts on  $P_n^{(2)}[\lambda]'$  from the right. So in addition to being a left  $P_n^{(2)}$ -module  $P_n^{(2)}[\lambda]'$  is also a right module for the group algebra of a permutation group  $S[\lambda]$  whose action is to permute the primed sides of outer propagating parts with the same number of inner propagating parts; and to permute the primed elements of inner propagating parts within each outer propagating part — thus

$$S[\lambda] = \prod_i S_{\bar{\lambda}_i} \wr S_{\bar{\lambda}^i}$$

(consider the action from below of  $S_1 \times S_2 \times S_3 \times S_1$  on outer and  $S_3 \times (S_2 \times S_2) \times (S_1 \times S_1 \times S_1) \times S_0$  on inner parts in figure 3). Indeed,  $P_n^{(2)}[\lambda]'$  is a sum of copies of the right regular module. Thus if  $L_\Theta$  is a simple left  $kS[\lambda]$ -module we may define a left  $P_n^{(2)}$ -module

$$\mathcal{S}_\Theta = \mathcal{S}_\Theta(n) = P_n^{(2)}[\lambda]' \otimes_{kS[\lambda]} L_\Theta.$$

This is an indecomposable summand of  $P_n^{(2)}[\lambda]'$  and hence a standard module of  $P_n^{(2)}$  (in the quasi-hereditary sense).

Evidently, every simple  $P_n^{(2)}$ -module appears as a subquotient of some  $P_n^{(2)}[\lambda]'$ , and each  $P_n^{(2)}[\lambda]'$  is a sum of modules of form  $\mathcal{S}_\Theta$ , with  $\mathcal{S}_\Theta \cong \mathcal{S}_{\Theta'}$  iff  $L_\Theta \cong L_{\Theta'}$ . Thus every simple  $P_n^{(2)}$ -module appears as a subquotient of some  $\mathcal{S}_\Theta$ . The following result is well known.

**Proposition 12** [23, §I.Appendix B] *Let  $G$  be a group with  $C_G$  conjugacy classes. Then equivalence classes of simple  $G \wr S_n$ -modules are indexed by  $C_G$ -tuples of integer partitions of total degree  $n$ .  $\square$*

It follows that for each  $\lambda$  there is a distinct inequivalent  $\mathcal{S}_\Theta$  for each *tuple* of multipartitions (tuples of integer partitions are called multipartitions) in which the  $i^{\text{th}}$  multipartition is as specified in the proposition in case  $S_{\bar{\lambda}_i} \wr S_{\bar{\lambda}^i}$ .

Let us denote by  $\Gamma_n$  the set of all such indices  $\Theta$  unioned over all  $\lambda$  such that  $\#^{\text{env}}(\lambda) = n$ . For example,  $\Gamma_0$  consists of a single label  $((), \text{say})$  corresponding to the nominal irreducible representation of a product over an empty set of wreaths (it will be evident from the construction that the correct interpretation is as a single copy of the ground ring);  $\Gamma_1$  consists of labels,  $(0), (1)$ , say, corresponding to the single irreducible representations of each of  $S_1 \wr S_0$  and  $S_1 \wr S_1$  respectively;  $\Gamma_2$  consists of labels,  $(0^{(2)}), (0^{(1^2)})$  (the irreducibles of  $S_2 \wr S_0$ ),  $(1, 0)$  (the irreducible of  $(S_1 \wr S_1) \times (S_1 \wr S_0)$ ),  $(1^{(2)}), (1^{(1^2)})$  (irreducibles of  $S_2 \wr S_1$ ),  $(1, 1)^{(2)}$  and  $(1, 1)^{(1^2)}$  (irreducibles of  $S_1 \wr S_2$ ), say. See figure 6 for more examples.













Note that the  $kS[\lambda]$  action may be realized as a subalgebra of  $P_n^{(T)}$  in a similar way to  $kS_n$ . The extra structure in  $kS[\lambda]$  is just so that it commutes with  $\mathcal{I}_\lambda$ . When we confuse elements of  $kS[\lambda]$  with elements of  $P_n^{(T)}$  it will be through this realization.

Since  $kS[\lambda]$  is a group algebra we may take it semisimple for  $k$  a suitable field of characteristic 0 (such as  $\mathbb{C}$ ). Let  $e_\Theta$  then be a primitive idempotent of  $kS[\lambda]$  such that  $kS[\lambda]e_\Theta = L_\Theta$  and  $e_\Theta kS[\lambda]e_\Theta = ke_\Theta$ . We may define an inner product  $\langle | \rangle$  on  $\mathcal{S}_\Theta$  by

$$\langle a|b \rangle = e_\Theta \mathcal{I}_\lambda = a^\circ b$$

(to be understood in the  $\lambda$  section, which makes the opposite right module to  $\mathcal{S}_\Theta$  also a dual).

For example,  $\mathcal{I}_\emptyset = \prod_{i=1}^n (A^n, A^n)$  and  $S[\emptyset] = S_0$  which here means the trivial group (so there is only one irreducible representation and  $e_\emptyset = 1$ ). A basis for  $P_n^{(T)} \mathcal{I}_\emptyset$  is  $\{wx \mid w \in \mathbf{E}_n^{(T)}, x = \{\{\{1'\}\}, \{\{2'\}\}, \dots, \{\{n'\}\}\}\}$ . Thus the gram matrix  $G_\emptyset$  of the inner product with respect to this basis has rows and columns conveniently indexed by  $\mathbf{E}_n^{(T)}$ . The  $ab^{th}$  entry of  $G_\emptyset$  is  $\prod_i Q_i^{h_i}$  where  $h_i$  is the number of parts in  $a_i b_i$  (the normal composition of partitions). In particular for  $n = 2$  the  $\lambda = \emptyset$  module has gram matrix computed as follows:

			
			
	$Q_1^2 Q_2^2$	$Q_1 Q_2^2$	$Q_1 Q_2$
			
	$Q_1 Q_2^2$	$Q_1 Q_2^2$	$Q_1 Q_2$
			
	$Q_1 Q_2$	$Q_1 Q_2$	$Q_1 Q_2$
			

whereupon

$$|G_\emptyset^{m=2}| = Q_1^3 Q_2^4 (Q_1 - 1)(Q_2 - 1).$$

Similarly we have (for the dimension twelve module at  $n = 3$ )

$$|G_\emptyset^{n=3}| = Q_1^{12} Q_2^{20} (Q_1 - 2)(Q_2 - 2)(Q_1 - 1)^7 (Q_2 - 1)^7$$

and (for the dimension 60 module at  $n = 4$ )

$$|G_\emptyset^{m=4}| = Q_1^{60} Q_2^{119} (Q_1 - 3)(Q_2 - 3)(Q_1 - 2)^{13} (Q_2 - 2)^{11} (Q_1 - 1)^{45} (Q_2 - 1)^{48}.$$

What is striking about these results is that the determinants factorise over  $\mathbb{Z}[Q_1] \cup \mathbb{Z}[Q_2]$  rather than  $\mathbb{C}[Q_1, Q_2]$ <sup>2</sup>.

<sup>2</sup>Indeed, computing the determinant of a common or garden 60x60 matrix with entries polynomial in 2 variables is probably beyond the reach of modern computers. We used the GAP package[17], some lucky guesses and some saturated bounds to reduce to a few tens of minutes computation on a powerful laptop.

**Proposition 13** *For all  $n$  the module  $\mathcal{S}_\emptyset$  is generically simple.*

*Proof:* Consider  $G_\emptyset$  for  $n$  arbitrary. It is enough to show that this is non-singular on an open subset of parameter space. Recall that the composition of partitions required to compute the inner product involves a transitive closure. It follows that the diagonal entry in each row of  $G_\emptyset$  is always of (possibly equal) highest total degree. Consider the matrix obtained by deleting the rows (and columns) where it is uniquely of highest degree. Iterating this procedure (cf. [29]) we arrive at the empty matrix. It follows that the determinant of the gram matrix is a finite polynomial in  $Q_1, Q_2$ .  $\square$

### 3.4 Modules $\mathcal{S}_\Theta(n)$ with $\#^{env}(\Theta) = n$

Let us extend the definition of  $\#^{env}$  so  $\#^{env}(\Theta) = \#^{env}(\lambda)$ , where  $\lambda$  is the propagating index underlying  $\Theta$ . Then ‘modules  $\mathcal{S}_\Theta$  with  $\#^{env}(\Theta) = n$ ’ are the  $P_n^{(2)}$ -modules  $\mathcal{S}_\Theta$  whose canonical basis elements have propagating index which cannot be realized in  $P_{n-1}^{(2)}$ . They also obey

$$D^{(T)}(A^i)\mathcal{S}_\Theta = D^{(T)}(A^{ij})\mathcal{S}_\Theta = 0$$

for all  $i, j$ , whereas no  $\mathcal{S}_\Theta$  with  $\#^{env}(\Theta) < n$  has this property (for, in this case we may construct a basis element from a basis element for the equivalent module at level  $n - 1$  by  $a \mapsto a \cup \{\{\{n\}\}, \{\{n'\}\}\}$  — and this is not killed by  $D^{(T)}(A^n)$ ).

**Proposition 14** *If  $\#^{env}(\Theta) = n$  and  $Q_{inner}$  invertible in  $k$  then  $\mathcal{S}_\Theta(n)$  is simple.*

*Proof:* Consider the gram matrix as constructed in the previous section. Because of the quotient (we are in the  $\lambda$  section) we may apply the pigeonhole principle to see that the inner product  $\langle a|b \rangle$  between any two basis elements is zero unless each outer part of  $a$  meets only one outer part of  $b^\circ$  in the composition. Thus the gram matrix breaks up into blocks indexed by the possible arrangements of outer parts in the restriction of  $a$  to  $\underline{n}'$  (or equivalently the restriction of  $b^\circ$  to  $\underline{n}$ ), and this arrangement characterizes all the products in the block. Since  $\#^{env}(\Theta) = n$  this implies that, in a block, each outer part from  $a$  meeting one from  $b^\circ$  carries the same number of inner propagating lines. Within the block, then, up to overall factors of  $Q_{inner}$ , the inner product coincides with that of the corresponding simple  $kS[\lambda]$ -module. Each block determinant is therefore the same, and non-zero. Hence the whole gram matrix is non-singular.  $\square$

### 3.5 Generic representation theory

For any  $T$  recall that  $e^T$  commutes with  $P_{n-1}^{(T)}$ , and that  $e^T e^T = Q^\pi e^T$ . Thus

**Proposition 15** *For  $\prod_{t \in T} Q_t$  invertible in  $k$ ,*

$$e^T P_n^{(T)} e^T = P_{n-1}^{(T)} e^T \cong P_{n-1}^{(T)} \quad (3)$$

*is an algebra isomorphism; and*

$$\begin{aligned} \mathcal{G} : P_{n-1}^{(T)}\text{-mod} &\rightarrow P_n^{(T)}\text{-mod} \\ M &\mapsto P_n^{(T)} e^T M \end{aligned}$$

*is a full embedding of  $P_{n-1}^{(T)}\text{-mod}$  in  $P_n^{(T)}\text{-mod}$  [16, §6.2] (cf. [26]).*

Note that  $\mathcal{G}(P_{n-1}^{(2)}\mathcal{I}_\lambda) = P_n^{(2)}\mathcal{I}_\lambda$  (NB, strictly speaking  $\mathcal{I}_\lambda$  is a different object on each side of this equation, i.e. depending on the level  $n$ ; it is this ‘globalization’ identity which justifies omission of the  $n$  label).

By equation (3), it is the quotient

$$P[n] := P_n^{(T)} / P_n^{(T)} e^T P_n^{(T)}$$

whose representations will be missed in the embedding (or equivalently, those left modules  $M$  for which  $e^T.M = 0$ ). Thus

**Proposition 16** *If  $\Lambda_n^{(T)}$  is an index set for the equivalence classes of simple modules of  $P_n^{(T)}$  then*

(i)  $|\Lambda_0^{(T)}| = 1$ ;

and if  $Q_1 Q_2$  invertible in  $k$  then

(ii)  $\Lambda_n^{(2)} = \Lambda_{n-1}^{(2)} \cup \Gamma_n = \cup_{0 \leq i \leq n} \Gamma_i$ , and

(iii)  $\{\text{head } \mathcal{S}_\Theta \mid \Theta \in \Lambda_n^{(2)}\}$  is a complete set of representatives of the equivalence classes of simple modules of  $P_n^{(2)}$ .

In other words, the set of equivalence classes of irreducible representations is indexed by the possible values of  $\#^-$  together with the corresponding set of tuples of multipartitions described in section 3.3.

Recall [27, §2.2] that  $P_n^m$  is the subalgebra of  $P_{n+m}$  in which each of the nodes labelled  $n+1, \dots, n+m$  is *identified* with its primed partner. The algebras  $P_n^m$  generalize in a natural way to algebras  $P_n^{(T)m}$  (any  $T$ ), whereupon  $P_n^{(T)} \subset P_n^{(T)1} \subset P_{n+1}^{(T)}$ . The chain of inclusions from  $P_0^{(T)}$  upward obtained from this sequence is denoted  $P_*^{(T)}$ .

As for restriction rules, it will be clear that the module  $\mathcal{S}_()$  in  $P_{2n}^{(2)}$  restricts to the regular representation of  $P_n^{(2)}$ . En route we have the restriction

$$\text{Res}_{P_n^{(2)}}^{P_{n-1}^{(2)}} \mathcal{S}_() \cong \mathcal{S}_() \oplus \mathcal{S}_{(0)} \oplus \mathcal{S}_{(1)}$$

and so on. The first few layers are depicted in figure 6. We do not need the general result here, but see proposition 20.

Given a module  $M \in P_{n-1}^{(T)}\text{-mod}$  there is a globalization functor  $\mathcal{G}$  and, via the injection of  $P_{n-1}^{(T)}$  into  $P_n^{(T)}$ , an induction functor  $\text{Ind}$  acting on it, so that  $\mathcal{G}M$  and  $\text{Ind}M$  are in  $P_n^{(T)}\text{-mod}$  for any  $n$ . In writing down such modules, we will not generally give an explicit indication of the level  $n$  unless it seems helpful to do so.

We now assemble the tools for proving generic semisimplicity. By construction (or using [10]) we have

**Proposition 17**

$$\mathcal{G}\mathcal{S}_\lambda(n) \cong \mathcal{S}_\lambda(n+1)$$

Recall [9]

**Proposition 18** *Projective modules of quasi-hereditary algebras have filtrations by standard modules.*

We denote the multiplicity in this filtration by  $[P_\lambda : \mathcal{S}_\mu]$ . The following is readily proved (cf. [26]):




Figure 6: Generic restriction rules for the generic irreducible  $\#^P = (0, 0)$  representation of  $P_5^{(2)}$  through the chain of restrictions  $P_*^{(2)}$ . The top layer is  $P_0^{(2)}$  and the bottom is  $P_5^{(2)}$ . In each layer up to  $P_2^{(2)1}$  a complete list of irreducible representations is given. Thereafter only irreducibles occurring in the restriction are recorded. The irreducibles are labelled by their  $\#^P$  index (i.e. ambiguously) and by their dimension.



**Proposition 19** For  $M \in P_n^{(T)} - \text{mod}$ ,

$$\text{Res}_{n+1}^{n+2} \mathcal{G} \mathcal{G} M \cong \text{Ind}_n^{n+1} M$$

$$\text{Ind } \mathcal{G} M \cong \mathcal{G} \text{Ind } M. \quad (4)$$

Thus if  $\mathcal{S}_\lambda(n)$  is projective for all  $n$  then so is  $\text{Res } \mathcal{S}_\lambda(n)$ . Whereupon

**Proposition 20**  $[\text{Res } \mathcal{S}_\lambda : \mathcal{S}_\mu] \neq 0$  implies  $\#^{\text{env}}(\lambda) - \#^{\text{env}}(\mu) \in \{-1, 0, +1\}$ .

*Proof:* Consider the restriction  $\text{Res}_{P_{n-1}^{(2)}}^{P_n^{(2)}} P_n^{(2)}[\hat{\lambda}]'$  where  $\hat{\lambda}$  is such that  $\mathcal{S}_\lambda$  is a summand.

Since  $P_n^{(2)}[\hat{\lambda}]'$  is a direct sum of  $\mathcal{S}_\lambda$ s all with the same  $\#^{\text{env}}$ , it is enough to show an equivalent result for this module.

In our basis of  $P_n^{(2)}[\hat{\lambda}]'$  the elements conform to one of the following possibilities for the connectivity of node  $n$  (the distinguished node, as it were, on restriction).

(1a) Node  $n$  is the sole unprimed element in a propagating part (inner not outer) — this contributes to a basis for a module isomorphic to  $P_{n-1}^{(2)}[\lambda']'$  with  $\lambda' \rho \hat{\lambda}$  via a  $\rho$ -step of type (i) (i.e. with  $\#^{\text{env}}(\lambda') = \#^{\text{env}}(\lambda) - 1$ ).

(1b) Node  $n$  is the sole unprimed element in a propagating part (inner and outer) — this means that  $\#^P(\hat{\lambda})$  includes a singleton, and the situation is as above except that this singleton has disappeared in  $\lambda'$ . Again  $\#^{\text{env}}(\lambda') = \#^{\text{env}}(\lambda) - 1$ .

(1c) Node  $n$  is the sole unprimed element in a propagating part (outer not inner) — this time  $\#^P(\hat{\lambda})$  includes a zero element and the situation is as above except that this has disappeared in  $\lambda'$ . Again  $\#^{\text{env}}(\lambda') = \#^{\text{env}}(\lambda) - 1$ .

(2a) Node  $n$  is a singleton both with respect to inner and outer partitions — the set of such objects forms a basis for a module isomorphic to  $P_{n-1}^{(2)}[\hat{\lambda}]'$ .

(2b) Node  $n$  is connected, but in such a way that regarding it as primed (for the purpose of determining the propagating index) does not change the propagating index (i.e. both the inner and outer part in which it lies have other nodes of both primed and unprimed type). Again sets of such objects form bases for modules isomorphic to  $P_{n-1}^{(2)}[\hat{\lambda}]'$ , provided we quotient by the modules produced at (1abc) above.

(3a) Node  $n$  is in a part (inner not outer) containing only unprimed elements — in the restriction there is no action on this node, so, quotienting by all the modules discussed so far, it may be regarded much as if it were primed. Thus we can build a basis for a copy of a module  $P_{n-1}^{(2)}[\lambda']'$  where  $\lambda'$  is such that one element of  $\#^P(\lambda')$  is incremented by 1 compared to  $\#^P(\lambda)$ , and hence  $\#^{\text{env}}(\lambda') = \#^{\text{env}}(\lambda) + 1$ .

(3b) Node  $n$  is in a part (inner and outer) containing only unprimed elements — as above, except that  $\#^P(\lambda')$  has a new singleton element. Again  $\#^{\text{env}}(\lambda') = \#^{\text{env}}(\lambda) + 1$ .

(3c) Node  $n$  is in a part (outer not inner) containing only unprimed elements — as above, except that  $\#^P(\lambda')$  has a new zero element. Again  $\#^{\text{env}}(\lambda') = \#^{\text{env}}(\lambda) + 1$ .

□

**Theorem 1**  $P_n^{(2)}(\mathbf{Q})$  is generically semisimple.

*Proof:* Let  $\text{Prop}_\lambda$  be that  $\mathcal{S}_\lambda = P_\lambda = S_\lambda$ , the simple module, for all  $n$  (i.e. this module lies in a singleton block). Then we are done if  $\text{Prop}_\lambda$  true for all  $\lambda$ . Suppose it is true for all  $\{\lambda \mid \#^{\text{env}}(\lambda) \leq m\}$ . Work by induction on  $m$ . The base case is

$m = 0$ , which is true generically by proposition 13 using quasi-heredity and Brauer–Humphreys reciprocity [9].

Applying equation (4) to case  $M = 1$ ,  $n = 0$  we see that every standard module with  $\#^{env}(\nu) \leq m$  appears in the  $m^{th}$  induction (and hence also the  $m^{th}$  restriction) of  $\mathcal{S}_0$ . Thus for each  $\nu$  with  $\#^{env}(\nu) = m + 1$  there is a  $\lambda$  with  $\#^{env}(\lambda) = m$  such that  $[\text{Res } \mathcal{S}_\lambda : \mathcal{S}_\nu] \neq 0$ . We now use this  $\lambda$  and the inductive assumptions to establish  $\text{Prop}_\nu$ .

Consider the Frobenius reciprocity

$$\text{Hom}_{n+1}(\text{Ind } \mathcal{S}_\kappa, \mathcal{S}_\lambda) \cong \text{Hom}_n(\mathcal{S}_\kappa, \text{Res } \mathcal{S}_\lambda) \quad (5)$$

By proposition 20, if  $\#^{env}(\kappa) \geq m + 2$  then this space is empty (since it is empty on the left by the inductive assumption). On the other hand, by assumption, all the indecomposable summands  $P_\chi$  of  $\text{Res } \mathcal{S}_\lambda$  are simple except possibly those with  $\#^{env}(\chi) = m + 1$ . But suppose some  $\mathcal{S}_\phi$  a submodule of  $P_\chi$ . Then it is simple by equation (5) and quasi-heredity (the only possible maps into it would show up in equation (5) for some  $\kappa$ ), and so  $\phi = \chi$ , and indeed  $P_\chi$  is simple by Brauer–Humphreys reciprocity and quasi-heredity. Thus in particular our  $\mathcal{S}_\nu = P_\nu = S_\nu$ .  $\square$

## 4 Exceptional representations

Fix  $Q \in \mathbb{N}$  and let  $V_Q$  denote the  $k$ -space with basis  $\{e_1, \dots, e_Q\}$ . Then for each ordinary partition algebra  $P_n(Q)$  there a  $Q^n$ -dimensional *Potts representation*  $R$  acting on  $V = V_Q^{\otimes n}$ . This is described, for example, in [28] (see also below). The underlying idea is that each basis element of  $V$  is a colouring of the  $n$  lattice sites ( $\{1, \dots, Q\}$  being the colours), and that such a colouring is ‘consistent’ with a given partition of the sites only if the elements within each part in the partition have the same colour. Fixing  $\mathbf{Q}$  a function from  $T$  to  $\mathbb{N}$ , the tensor product algebra  $\bigotimes_t P_n(Q_t)$  may be equipped with a corresponding product representation. It will be evident that this representation is  $(\prod_t Q_t)^n$ -dimensional. This is to say, it acts on a space isomorphic to that of the Potts representation of  $P_n(\prod_t Q_t)$  (recall the ‘diagonal’ isomorphism of  $P_n(\prod_t Q_t)$  to a subalgebra of our algebra). We will again write  $R$  both for this representation of  $\bigotimes_t P_n(Q_t)$  and for its restriction to  $P_n^{(T)}$ .

**Theorem 2** *For any  $\mathbf{Q} \in \mathbb{N}^d$  there exists an  $m \in \mathbb{N}$  such that  $P_n^{(T)}(\mathbf{Q})$  is non-semisimple for all  $n > m$ .*

*Proof:* The proof is a straightforward generalization of the corresponding result for the ordinary partition algebra as in [29]. In short, one compares the growth rate (as  $n$  increases) of dimension of standard module  $\mathcal{S}_0$  with that of the Potts representation  $R$ , having established that, if semisimple, the former is a summand of the latter (consider  $R(\mathcal{I}_0)$ ).  $\square$

### 4.1 Application to physical models

For  $H$  a graph let  $\mathfrak{E}_H$  (resp.  $\mathfrak{V}_H$ ) denote its edge (resp. vertex) set. Define graph  $A_l$  by  $\mathfrak{V}_{A_l} = \{1, 2, \dots, l\}$  and  $\mathfrak{E}_{A_l} = \{\{i, j\} | i - j = \pm 1\}$ , and graph  $\hat{A}_l$  by  $\mathfrak{V}_{\hat{A}_l} = \{1, 2, \dots, l\}$  and  $\mathfrak{E}_{\hat{A}_l} = \{\{i, j\} | i - j \equiv \pm 1 \pmod{l}\}$ , as usual.

Fix  $Q \in \mathbb{N}$ . A *spin configuration* on graph  $H$  is an element of  $S_H = \text{hom}(\mathfrak{V}_H, \{1, 2, \dots, Q\})$  (i.e. a colouring of the vertices of  $H$  from  $Q$  colours). For  $s \in S_H$  let  $s_i$  denote the ‘colour’ of vertex  $i$ .

Given a graph  $H$  let graph  $G = H \times A_l$  be given by  $\mathfrak{V}_{H \times A_l} = \mathfrak{V}_H \times \mathfrak{V}_{A_l}$  and  $\{(a, b), (c, d)\} \in \mathfrak{E}_{H \times A_l}$  if either  $a = c$  and  $\{b, d\} \in \mathfrak{E}_{A_l}$  or  $b = d$  and  $\{a, c\} \in \mathfrak{E}_H$ . We may break graph  $G$  up into  $l$  ‘lateral’ layers  $H_k$  ( $k = 1, 2, \dots, l$ ) each isomorphic to  $H$ , and  $l - 1$  ‘transverse’ layers  $H_{k,k+1}$  each consisting of the edges between a certain pair of lateral layers.

A  $Q$ -state model on  $G$  may be constructed as follows [2, 24]. An *edge Hamiltonian*  $\mathcal{H}$  is a map from  $\text{hom}(\mathfrak{V}_{A_2}, \{1, 2, \dots, Q\})$  to the real numbers (here  $A_2$  stands for any of the 2 vertex subgraphs of  $G$  containing precisely one edge). From this we form *graph Hamiltonian*

$$\mathcal{H}_G = \sum_{\{i,j\} \in \mathfrak{E}_G} \mathcal{H}(s_i, s_j).$$

Given such a function and a complex variable  $\beta$  define the *partition function*

$$Z = Z(\beta) = \sum_{s \in S_G} \exp(\beta \mathcal{H}_G(s)).$$

In Statistical Mechanics one wants to know the analytic structure of  $Z(\beta)$ .<sup>3</sup> If the Hamiltonian  $\mathcal{H}(s_i, s_j)$  is integer valued then, regarding  $Z$  as a function of  $\exp(\beta)$ , this structure consists of roots. (More generally of limits of distributions of roots as  $l$  and  $H$  grow large in some way.)

For  $s, t \in S_H$  let  $T_{st}^{(l)}$  denote the partition function in case the colours of vertices in  $H_1$  (resp.  $H_l$ ) have been *fixed* to  $s$  (resp.  $t$ ). Note that, given a bijection  $\mathfrak{V}_H \leftrightarrow \{1, 2, \dots, n\}$ ,  $S_H$  is a basis for  $V$  in a natural way. Thus  $T^{(l)}$  is a matrix in  $\text{End}(V)$ . In case  $l = 1$  the first and last layers are identified, and  $T^{(1)}$  is a diagonal matrix. We may similarly introduce a matrix  $T^{(-)}$  of partition functions associated to a single transverse layer  $H_{k,k+1}$ . Here  $s$  determines the colour of vertices in layer  $k$ , and  $t$  in layer  $k + 1$ . These layer matrices are called *transfer matrices*. Note that we may use them to build up  $T^{(l)}$  one layer at a time:

$$T^{(l)} = (T^{(1)} T^{(-)})^l T^{(1)}.$$

Similarly, we can introduce matrices which will build up  $T^{(1)}$  and  $T^{(-)}$  one *edge* at a time.

For certain choices of  $\mathcal{H}(s_i, s_j)$  the representation  $R$  can be used to build these transfer matrices. The representation  $R$  of the full tensor product algebra may be used to build transfer matrices for models with a significantly greater variety of Hamiltonian than can the restriction of  $R$  to  $P_n(\prod_t Q_t)$ . The point about  $P_n^{(T)}$  is that  $R$  may be restricted to it *without* losing the greater variety. A suitable concrete example will illustrate these points. Let  $\mathbf{Q} = (Q_1, Q_2) = (2, 3)$ . Let  $D_j$  be the  $j \times j$  matrix with all entries 1, and  $I_j$  be the  $j \times j$  unit matrix. Then

$$R((A^i, A^i)) = I_{6^{i-1}} \otimes D_{Q_1} \otimes D_{Q_2} \otimes I_{6^{n-i-2}}$$

$$R((A^i, 1)) = I_{6^{i-1}} \otimes D_{Q_1} \otimes I_{Q_2} \otimes I_{6^{n-i-2}}$$

---

<sup>3</sup>Actually  $Z$  will only be physically interesting, within the framework described, for certain carefully chosen  $G$  and  $\mathcal{H}$ ; and interesting constructions are possible outside of this framework. Neither point need concern us here.

(cf. [2, §12.4]).

The colour-valued variables  $s_i$  associated to the vertices  $i$  of  $G$  are called *spins*. Note that  $R$  above acts on a space appropriate for a  $Q_1 Q_2 = 6$ -state model. Considering spins  $s_i$  as taking values from the integers mod. 6, then a  $Z_6$ -symmetric model [2, 12] is one in which the edge Hamiltonian takes the form  $\mathcal{H}(s_i, s_j) = f(s_i - s_j)$ , where  $f$  may be any real valued even function <sup>4</sup>. Suppose we want to build the transfer matrix for such a model with  $f(0) = 2$ ,  $f(1) = f(2) = 1$ ,  $f(3) = 0$  (such as the model examined in [24, p.305]). Then the transfer matrix factor building in the contribution of that edge to the Hamiltonian is

$$\begin{aligned} R((1 + (e^\gamma - 1)(A^{ij}, A^{ij}))(1 + (e^\beta - 1))(1, A^{ij})) \\ = R((1 + (e^\beta - 1))(1, A^{ij}) + e^\beta(e^\gamma - 1)(A^{ij}, A^{ij})) \end{aligned}$$

if lattice edge  $(i, j)$  is in the transfer matrix layer (i.e. it is in some  $H_k$ ), and

$$R((e^\beta(e^\gamma - 1)1 + (e^\beta - 1)(A^i, 1) + (A^i, A^i)))$$

if  $(i, j)$  is perpendicular to the transfer matrix layer; where  $\gamma = 2$ ,  $\beta = -1$ .

The analytic structure of the partition function  $Z$  for such models reveals physical *phase transitions* as points on the positive  $\exp(\beta)$ -axis which are approached by distributions of roots as the system size grows large (see [24] for details). In computing this structure we are, at base, using the algebra as nothing more than a formal aid to computation (the globalization functor  $\mathcal{G}$  gives explicit control of the thermodynamic, i.e. large  $n$ , limit as a global algebra limit, but there is no way yet to use this in specific calculations). None the less, the results are of considerable interest both in the way they model phase transitions and in what they tell us about specific physical systems. We will postpone such interpretation to a separate paper but, in the renaissance spirit, include a couple of results here (in the form of distributions of roots), for illustration. (There has been renewed interest in results of this kind recently, although mostly obtained by brute force calculation, see [31].) Figures 7 and 8 show the roots of  $Z$  in  $\exp(\beta)$  for  $Q_1 = Q_2 = 2$  and  $f(0) = 3$ ,  $f(1) = 1$  and  $f(2) = 0$ , for various graphs. The results for various different graphs are shown, as a rough indication of the sense in which the distribution of roots may approach a limit as  $G$  becomes large. <sup>5</sup> The graphs concerned are given by  $(H = \hat{A}_7, l = 9)$  and  $(H = \hat{A}_8, l = 9)$  and  $(H = \hat{A}_8, l = 13)$  and  $(H = \hat{A}_{10}, l = 13)$  (i.e. these examples are models of two-dimensional physical systems).

## 5 Discussion

There remain a number of interesting open problems on the representation theory side. We have not determined the exceptional structure of  $P_n^{(2)}$  (cf.  $P_n$ , where complete results are known), nor addressed its representation theory in positive characteristic (nothing is known here for  $P_n$  either, although it *is* known to be a

---

<sup>4</sup>Directed graphs and non-even functions produce *chiral* models, which have also been studied, but need not concern us here.

<sup>5</sup>A better indication is perhaps given by the *animated* combined version of these figures available online from [www.city.ac.uk/mathematics/algebra/statinfo/StatMechIntro2.html](http://www.city.ac.uk/mathematics/algebra/statinfo/StatMechIntro2.html) or from PPM's homepage.




Figure 7: Zeros of the partition function in  $e^\beta$  for  $7 \times 9$  (dots) and  $8 \times 9$  (crosses) square lattice systems with  $Q_1 = Q_2 = 2$ .




Figure 8: Zeros of the partition function in  $e^\beta$  for  $8 \times 13$  (crosses) and  $10 \times 13$  (dots) square lattice systems with  $Q_1 = Q_2 = 2$ .

hard problem [30]). As for more exotic posets  $T$ , we do not know even the generic structure of  $P_n^{(T)}$  (indeed we do not even know if it *has* a generic structure).

As a quasi-hereditary algebra, it is natural to ask if  $P_n^{(2)}$  has tilting modules [9]. These have yet to be constructed even for  $P_n$ .

The assumptions of this paper restrict us to finite  $T$ , however there is no serious obstruction to relaxing this constraint. Infinite  $T$  lacks, for the moment, physical motivation, however it offers some intriguing possibilities cf. [34] (and references therein) for example.

**Acknowledgements.** PPM would like to thank Anton Cox for invaluable discussions, and EPSRC for partial financial support.

## Appendix: Rank of $P_n^{(2)}$

The degree of  $p_n^{(2)}$ , or the set  $\mathbf{E}_N^{(2)}$  of  $\underline{2}$ -ramified partitions of any finite set  $N$ , may be computed as follows. The *shape* of a partition (of  $N$  say) is the list of degrees of its parts, expressed as an integer partition. For  $\mu$  such a shape, let  $g_\mu$  be the number of partitions with that shape. Let  $b_n$  be the number of partitions of  $\underline{n}$ . Then

$$|p_n^{(2)}| = \sum_{\mu} g_{\mu} \prod_i b_{\mu_i}$$

The first few of these are

$$p_1^{(T)} : \begin{array}{r} \frac{g_{\mu} \times \prod}{1 \times 1} \quad \mu \\ \frac{1 \times 2}{\sum = 3} \quad (1, 1) \\ \quad \quad \quad (2) \end{array}$$

$$\begin{array}{r} \frac{g_{\mu} \times \prod}{1 \times 1} \quad \mu \\ \frac{3 \times 2}{\sum = 12} \quad (1, 1, 1) \\ \quad \quad \quad (2, 1) \\ \quad \quad \quad (3) \end{array}$$

$$p_2^{(T)} : \begin{array}{r} \frac{g_{\mu} \times \prod}{1 \times 1} \quad \mu \\ \frac{6 \times 2}{\sum = 60} \quad (1, 1, 1, 1) \\ \quad \quad \quad (2, 1, 1) \\ \quad \quad \quad (3, 1) \\ \quad \quad \quad (2, 2) \\ \quad \quad \quad (4) \end{array}$$

$g_\mu \times \Pi$	$\mu$
$1 \times 1$	$(1, 1, 1, 1, 1)$
$10 \times 2$	$(2, 1, 1, 1)$
$10 \times 5$	$(3, 1, 1)$
$15 \times 2 \times 2$	$(2, 2, 1)$
$5 \times 15$	$(4, 1)$
$10 \times 2 \times 5$	$(3, 2)$
$1 \times 52$	$(5)$
<hr/>	
$\Sigma = 358$	

These figures may be compared with the summed squares of dimensions of generic irreducible representations of the corresponding algebras in figure 6.

## References

- [1] J Ashkin and E Teller, Phys Rev **64** (1943), 178–184.
- [2] R J Baxter, *Exactly solved models in statistical mechanics*, Academic Press, New York, 1982.
- [3] V V Bazhanov and R J Baxter, *Star triangle relation for a 3-dimensional model*, J Stat Phys **71** (1994), 839–864.
- [4] R Brauer, *On algebras which are connected with the semi-simple continuous groups*, Annals of Mathematics **38** (1937), 854–872.
- [5] E Cline, B Parshall, and L Scott, *Finite-dimensional algebras and highest weight categories*, J. reine angew. Math. **391** (1988), 85–99.
- [6] ———, *Generic and  $q$ -rational representation theory*, Publ RIMS **35** (1999), 31–90.
- [7] S Dasmahapatra and P P Martin, *On the algebraic approach to cubic lattice Potts models*, Journal of Physics A **29** (1996), 263–278.
- [8] V. Dlab and C.M. Ringel, *A construction for quasi-hereditary algebras*, Compos. Math. **70** (1989), 155–175.
- [9] S. Donkin, *On tilting modules for algebraic groups*, Math. Z. **212** (1993), 39–60.
- [10] ———, *The  $q$ -Schur algebra*, LMS Lecture Notes Series, vol. 253, Cambridge University Press, 1998.
- [11] ———, *Symmetric and exterior powers, linear source modules and representations of Schur superalgebras*, Proc LMS **83** (2001), 647–680.
- [12] M B Einhorn, R Savit, and E Rabinovici, *A physical picture for the phase transition in  $Z_N$  symmetric models*, Nucl Phys B **170** [FS1] (1980), 16.
- [13] S Elitzur, R B Pearson, and J Shigemitsu, *The phase structure of discrete abelian spin and gauge systems*, Phys Rev D **19** (1979), 3698.
- [14] C Fan, Phys Lett **39A** (1972), 136–136.



- [15] F M Goodman, P de la Harpe, and V F R Jones, *Coxeter–Dynkin diagrams and towers of algebras*, Math Sci Research Inst Publ, Springer–Verlag, Berlin, 1989.
- [16] J A Green, *Polynomial representations of  $GL_n$* , Springer–Verlag, Berlin, 1980.
- [17] The GAP Group, *GAP — Groups, Algorithms, and Programming, version 4.2*, Aachen, St. Andrews, 1999.
- [18] T Halverson and A Ram, *Characters of algebras containing a Jones basic construction...*, Adv Math **116** (1995), 263–321.
- [19] P Hanlon and D Wales, *A tower construction for the radical in Brauer’s centralizer algebras*, J Algebra **164** (1994), 773–830.
- [20] A Henke and S Koenig, *Relating polynomial  $gl(n)$ –representations in different degrees*, J reine angew Math (to appear).
- [21] V F R Jones, in Subfactors (Singapore), World Scientific, 1994.
- [22] H F Mattson Jr., *Discrete mathematics with applications*, Wiley, Singapore, 1993.
- [23] I Macdonald, *Symmetric functions and Hall polynomials*, 2 ed., Oxford, 1995.
- [24] P P Martin, *Potts models and related problems in statistical mechanics*, World Scientific, Singapore, 1991.
- [25] ———, *Temperley–Lieb algebras for non–planar statistical mechanics — the partition algebra construction*, Journal Of Knot Theory and its Ramifications **3** (1994), no. 1, 51–82.
- [26] ———, *The structure of the partition algebras*, J Algebra **183** (1996), 319–358.
- [27] ———, *The partition algebra and the Potts model transfer matrix spectrum in high dimensions*, J Phys A **32** (2000), 3669–3695.
- [28] P P Martin and G Rollet, *The Potts model representation and a Robinson–Schensted correspondence for the partition algebra*, Compositio Math **112** (1998), 237–254. Corrigendum: multiple substitutions of  $\Rightarrow$  for  $-$  symbol!
- [29] P P Martin and H Saleur, *Algebras in higher dimensional statistical mechanics — the exceptional partition algebras*, Lett. Math. Phys. (1994), no. 30, 179–185.
- [30] P P Martin and D Woodcock, *The partition algebras and a new deformation of the Schur algebras*, J Algebra **203** (1998), 91–124.
- [31] V Matveev and R Shrock, *Complex-temperature phase diagram of the 1D  $Z(6)$  clock model and its connection with higher-dimensional models*, PHYSICS LETTERS A **221** (1996), 343–349.
- [32] R K Pathria, *Statistical Mechanics*, Pergamon, 1972.

- [33] T Regge and R Zecchina, *Combinatorial and topological approach to the 3D Ising model*, JOURNAL OF PHYSICS A-MATHEMATICAL AND GENERAL **33** (2000), 741–761.
- [34] A M Vershik and Yu V Yakubovich, *On continuous partition lattice*, preprint (St. Petersburg) (1998).
- [35] H Weyl, *Classical groups*, Princeton, Princeton, 1946, ch.III-V.
- [36] C. Xi, *Partition algebras are cellular*, Compositio Mathematica **119** (1999), 107–118.