

COHOMOLOGY AT INFINITY AND THE WELL-ROUNDED RETRACT FOR GENERAL LINEAR GROUPS

AVNER ASH AND MARK MCCONNELL

INTRODUCTION

(0.1). Let \mathbf{G} be a reductive algebraic group defined over \mathbb{Q} , and let Γ be an arithmetic subgroup of $\mathbf{G}(\mathbb{Q})$. Let X be the symmetric space for $\mathbf{G}(\mathbb{R})$, and assume X is contractible. Then the cohomology (mod torsion) of the space X/Γ is the same as the cohomology of Γ . In turn, X/Γ will have the same cohomology as W/Γ , if W is a “spine” in X . This means that W (if it exists) is a deformation retract of X by a Γ -equivariant deformation retraction, that W/Γ is compact, and that $\dim W$ equals the virtual cohomological dimension (vcd) of Γ . Then W can be given the structure of a cell complex on which Γ acts cellularly, and the cohomology of W/Γ can be found combinatorially.

Spines have been found for many groups \mathbf{G} (see (2.6) below, and [A1] [M-M0] [M-M1]). This paper concerns the case where \mathbf{G} is the restriction of scalars of a general linear group over a number field k with ring of integers \mathcal{O} . This means $\mathbf{G}(\mathbb{Q}) = \mathrm{GL}_n(k)$. We shall take Γ to be a subgroup of finite index in $\mathrm{GL}_n(\mathcal{O})$, or more generally in $\mathrm{GL}(\mathcal{P})$, where \mathcal{P} is a projective \mathcal{O} -module of rank n . In [A3], Ash found a spine W for these \mathbf{G} , calling W the *well-rounded retract*. (This generalized [Sou2].) The retract has been used in computations: see [Sou1] [A-G-G] [A-M1] [A-M2] [vG-T]. We remark that $k = \mathbb{Q}$, $\Gamma \subseteq \mathrm{GL}_n(\mathbb{Z})$ for $n = 2, 3, 4$ still provide our main cases of computational interest. For imaginary quadratic fields k and $n = 2$, see [Men] [S-V] [V]. For real quadratic k and $n = 2$, see [B].

(0.2). The present paper extends [A3] to deal with the “cusps” of X/Γ , in a way which we now describe.

Borel and Serre [B-S] introduced a bordification \bar{X} of X such that \bar{X}/Γ is a compactification of X/Γ . The space \bar{X}/Γ has the same homotopy type as X/Γ , and therefore has the same homotopy type as W/Γ . The boundary of \bar{X}/Γ is a union of finitely many “faces”, one for each equivalence class mod Γ of parabolic \mathbb{Q} -subgroups of \mathbf{G} .

Research partially supported by NSA grant MDA-904-94-H-2030 (first author) and NSF grants DMS-9206659 and 9401460 (second author). This manuscript is submitted for publication with the understanding that the United States government is authorized to reproduce and distribute reprints.

The Borel-Serre boundary gives a geometric approach for distinguishing the cuspidal cohomology within the whole cohomology of Γ . A cohomology class on \bar{X}/Γ with coefficients in \mathbb{C} can be viewed, via the de Rham theorem, as coming from a Γ -invariant automorphic form on $\mathbf{G}(\mathbb{R})$. If the class has non-zero restriction to the boundary, it is certainly not a cusp form. The converse is not completely true, but the difference between the “interior cohomology” (the span of classes that restrict to zero on the boundary) and the “cuspidal cohomology” (the span of classes whose associated automorphic forms are cuspidal) is fairly well understood, thanks to the theory of Eisenstein series. In many practical examples, like the $\mathrm{GL}_4(\mathbb{Z})$ example mentioned below, there is no difference; to compute the cuspidal cohomology in certain degrees, we simply compute the restriction of the cohomology to the boundary.

A related question is the existence of “ghost classes”—those classes which restrict to zero on each face of the boundary, but do not restrict to zero on the whole boundary. Harder has given examples of ghost classes [H]; it is of interest to know how widespread the phenomenon is.

In specific cases, then, one may need to compute the cohomology of the boundary of \bar{X}/Γ , the cohomology of individual boundary faces, and the restriction map from $H^*(\bar{X}/\Gamma)$ to these groups.

This paper builds a technology for these computations, when \mathbf{G} is the general linear group in (0.1) and W is the well-rounded retract of [A3]. To each Borel-Serre boundary face in \bar{X} , we associate a closed subcomplex W' of W . Let Γ' be the stabilizer of W' in Γ . We prove that the inclusion $W'/\Gamma' \hookrightarrow W/\Gamma$ induces the same map on cohomology as the inclusion of the corresponding boundary face into \bar{X}/Γ . Our main result, Theorem 9.5, is that there is a spectral sequence converging to $H^*(\partial\bar{X}/\Gamma)$ whose E_1 term is the direct sum of the cohomology of the various W'/Γ' . There are only finitely many terms in this direct sum. There is a map of spectral sequences, starting with a natural map from $H^*(W/\Gamma)$ to the E_1 term, that induces the canonical restriction map $H^*(\bar{X}/\Gamma) \rightarrow H^*(\partial\bar{X}/\Gamma)$ up to canonical isomorphism. There is a dual statement in homology (10.2). We do not assume that Γ is torsion-free. We use (co)homology with coefficients in any fixed abelian group.

We can also find the maps on (co)homology induced by the inclusion of a single boundary face into \bar{X}/Γ , without any spectral sequences (10.3).

Each W'/Γ' is a finite cell complex, so its cohomology can be worked out combinatorially. See (6.3) for an example. We emphasize that each W' is a subcomplex of W , so that we can work with it in terms already available once we have computed W in any given case.

In a future paper, we will use this method to study the congruence subgroups $\Gamma_0(p)$ of prime level p in $\mathrm{GL}_4(\mathbb{Z})$. For a range of p , we will compute $H^5(\Gamma_0(p), \mathbb{C})$, and will isolate the subspace of classes that restrict to zero on the boundary. We will compute the Hecke eigenvalues of these classes, using either the new techniques of Paul Gunnells, or the new techniques of R. MacPherson and the second author. We will try to determine whether any of the cusp forms we find are functorial lifts from smaller groups such as Sp_4 or O_4 .

(0.3). We have long wished to see spelled out in detail the relationship between the well-rounded retract and the Borel-Serre compactification. The retraction $r : X \rightarrow W$ carries any $x \in X$ into W by a combination of geodesic actions depending on x —see (4.2). Each

Borel-Serre boundary component e in \bar{X} has a tubular neighborhood \bar{N} , homeomorphic to $e \times [0, 1]^l$ for appropriate l , where the fibers $[0, 1]^l$ are given by the geodesic action. The retraction r is constant on the interior of each fiber, and carries the interior of \bar{N} onto the corresponding cell complex W' .

The main idea of the paper is to extend the retraction $r : X \rightarrow W$ to a continuous Γ -equivariant map $\bar{r} : \bar{X} \rightarrow W$. It is relatively easy to see that the continuous extension exists and is Γ -equivariant. However, $r = r_1$ really comes from a deformation retraction $r_t : X \times [0, 1] \rightarrow X$, and it is harder to prove that \bar{r} comes from a deformation retraction of \bar{X} onto W . A simple example will illustrate the problem. The map $f_t : \mathbb{R} \times [0, 1] \rightarrow \mathbb{R}$ given by $(x, t) \mapsto (1 - t) \cdot x$ is a deformation retraction of \mathbb{R} onto $\{0\}$. The compactification $[-\infty, \infty]$ of \mathbb{R} certainly has a deformation retraction onto $\{0\}$. But f_t does not extend to the deformation retraction for $[-\infty, \infty]$.

To fix the problem with $[-\infty, \infty]$, we can first use any deformation retraction onto $[-x_0, x_0]$ for some large x_0 , and then use the old f_t to retract $[-x_0, x_0]$ to $\{0\}$. We do something similar to prove \bar{r} is a deformation retraction. We use Saper's recent construction [S] of a "central tile" $X_0 \subset X$ and a deformation retraction $r_{t,S}$ of \bar{X} onto X_0 . This X_0 is a large codimension-zero subset of X , which we may take arbitrarily close to $\partial\bar{X}$ in all directions. We show that composing our r_t with Saper's $r_{t,S}$ gives a deformation retraction $R_t : \bar{X} \times [0, 1] \rightarrow \bar{X}$ with $\bar{r} = R_1$.

By what we have said, \bar{r} gives a homotopy equivalence between \bar{X}/Γ and W/Γ . We show it also gives a homotopy equivalence between any Borel-Serre boundary component \bar{e}/Γ' and the corresponding cell complex W'/Γ' . It is then routine to show that \bar{r} gives an isomorphism between a standard spectral sequence for $H^*(\partial\bar{X}/\Gamma)$ and the spectral sequence described in (0.2). This proves our main theorem.

The sections of the paper follow this outline. In Sections 1–4 we summarize basic material on algebraic groups, lattices, the well-rounded retract and retraction, and the Borel-Serre compactification. None of this material is new, but we must recall it to establish our notation, and because we emphasize some less familiar aspects of it—marked lattices, the flag of successive minima, the orthogonal scaling group (which is the geodesic action from a lattice's point of view), and the connections between r_t and the geodesic action. Section 5 summarizes our notation. In Section 6 we define the sets $W_{\mathcal{F}}$ (called W' in this introduction). In Section 7 we define the tubular neighborhoods $\bar{N}_{\mathcal{F}}$ (called \bar{N} here); this is the technical heart of the paper, where we explore the connections between the well-rounded retraction and the geodesic action. Section 8 summarizes the material from [S] that we need, and contains our results on R_t and \bar{r} . In Section 9, we use \bar{r} to prove our main theorem on spectral sequences.

(0.4). The results in this paper could be extended in several ways. First, [A3] defines W for GL_n of any finite dimensional division algebra over \mathbb{Q} , not necessarily commutative. Our results could be generalized to that case, presumably with only technical changes. Second, our arguments work for SL as well as GL , with minor changes (10.5). See (10.4) for comments on Γ -equivariant cohomology, and (10.6)–(10.7) for other connections with reduction theory.

Parts (1.6), (2.6), and (6.3) present the case $k = \mathbb{Q}$ in more detail. We hope they help

guide the reader through the technicalities that involve structure over k versus structure over \mathbb{Q} .

We are grateful to J. R. Myers and Steve Zucker for helpful conversations. We especially thank Chun-Nip Lee and Leslie Saper. We are grateful to the referee for several comments. We thank Bob MacPherson and Dan Grayson for introducing us to some of these ideas, and Connie Chen for help with the illustration. We thank the NSA and the NSF for support.

SECTION 1—BACKGROUND ON ALGEBRAIC GROUPS

(1.1). We fix a number field k of degree d over \mathbb{Q} , with ring of integers \mathcal{O} . Let v run through the archimedean places of k ; the completion k_v is either \mathbb{R} or \mathbb{C} . Let \mathbb{R}_+ be the positive real numbers.

(1.2). Fix an integer $n \geq 1$. The algebraic group GL_n is defined over \mathbb{Q} , and hence over k . Let \mathbf{G} be the restriction of scalars of GL_n from k down to \mathbb{Q} .

To see this algebraic group concretely, fix a \mathbb{Q} -basis of k . Let k_1 be any commutative k -algebra. A point of \mathbf{G} with entries in k_1 is an $n \times n$ matrix with non-zero determinant whose (i, j) -th entry is a d -tuple of coordinates $(x_{ij}^{(1)}, \dots, x_{ij}^{(d)})$ with $x_{ij}^{(l)} \in k_1$. The d -tuple gives an element of $k \otimes_{\mathbb{Q}} k_1$ with respect to the fixed \mathbb{Q} -basis of k . Addition and multiplication of d -tuples are defined using the corresponding operations in k . We fix such a realization of \mathbf{G} throughout the paper.

(1.3). Let $S = k \otimes_{\mathbb{Q}} \mathbb{R}$. The real and complex embeddings of k induce a canonical isomorphism of \mathbb{R} -algebras $S \cong \prod_v k_v$. Clearly $k \hookrightarrow S$ by the canonical diagonal embedding.

Set $G = \mathbf{G}(\mathbb{R}) = \mathrm{GL}_n(S)$. Each entry in a matrix of G breaks up as a product over the v 's, so that $G \cong \prod_v \mathrm{GL}_n(k_v)$.

(1.4). Let ${}_{\mathbb{Q}}\mathbf{T}$ be the following maximal \mathbb{Q} -split torus of \mathbf{G} : if the fixed basis of k over \mathbb{Q} has 1 as its first element, then ${}_{\mathbb{Q}}\mathbf{T}$ is the subgroup where all the $x_{ij}^{(m)}$ are zero except for $x_{ii}^{(1)}$, $i = 1, \dots, n$. Write ${}_{\mathbb{Q}}\mathbf{T} = \{\mathrm{diag}(a_1, \dots, a_n)\}$ with $a_i = x_{ii}^{(1)}$.

The split radical $R_d \mathbf{G}$ of \mathbf{G} is the subgroup of ${}_{\mathbb{Q}}\mathbf{T}$ in which all the a_i are equal to each other. Let H be the identity component of $R_d \mathbf{G}(\mathbb{R})$. These are the *positive real homotheties*: $h \in H$ acts by a common positive real scalar on each factor $\mathrm{GL}_n(k_v)$ of G . We identify $H \cong \mathbb{R}_+$.

(1.5). Let $\mathbf{a}, \mathbf{b} \in S^n$. These break up as a product of vectors $\mathbf{a}_v, \mathbf{b}_v \in (k_v)^n$. Write $\mathbf{a}_v = (a_1, \dots, a_n)$ with $a_i \in k_v$, and write \mathbf{b}_v similarly. For each v , let $\langle (a_1, \dots, a_n), (b_1, \dots, b_n) \rangle_v = \sum_{i=1}^n \mathrm{trace}_{\mathbb{R}}^{k_v}(a_i \bar{b}_i)$, where the bar denotes complex conjugation when v is complex and the identity when v is real. Let $\langle \mathbf{a}, \mathbf{b} \rangle = \sum_v \langle \mathbf{a}_v, \mathbf{b}_v \rangle_v$, a positive-definite product on S^n . Let K_v be the subgroup of $\mathrm{GL}_n(k_v)$ preserving \langle, \rangle_v . Let $K = \prod_v K_v$; this is the largest subgroup of G preserving \langle, \rangle , and is a maximal compact subgroup of G . Let $\|\mathbf{a}\| = \langle \mathbf{a}, \mathbf{a} \rangle^{1/2}$.

Definition. Let $X = (HK) \backslash G$.

In the language of [B-S], X is a homogeneous space for G of type S - \mathbb{Q} . It is a symmetric Riemannian space for the Lie group $H \backslash G$.

(1.6). *Example of $\mathrm{GL}_n(\mathbb{Z})$.* Let $k = \mathbb{Q}$. Then $G = \mathrm{GL}_n(\mathbb{R})$, the group of $n \times n$ invertible matrices with real entries. The group ${}_{\mathbb{Q}}\mathbf{T}$ consists of the diagonal matrices, and $H = \{\mathrm{diag}(h, \dots, h) \mid h > 0\}$. The space X is the symmetric space consisting of the real $n \times n$ symmetric positive-definite matrices mod homotheties. When $n = 2$, X is isomorphic to the upper half-plane (see (2.6)).

SECTION 2—MARKED LATTICES. THE WELL-ROUNDED RETRACT

We recall the definition and basic properties of the well-rounded retract for GL_n . Except for the emphasis on marked lattices, the material comes from [A3]; we specialize this to the number field case, and omit the proofs. For the $\mathrm{GL}_n(\mathbb{Z})$ case, see (2.6).

(2.1). The space S^n of column vectors is an \mathbb{R} -vector space. S (and hence $k \subset S$) acts on it by coordinate-wise multiplication, and $G = \mathrm{GL}_n(S)$ acts on it on the left. These module structures commute with each other. If A is a subset of S^n , the S -module it spans in S^n is denoted $S \cdot A$.

(2.2). A \mathbb{Z} -lattice in a vector space V over \mathbb{Q} is a finitely generated subgroup of V which contains a \mathbb{Q} -basis of V . A \mathbb{Z} -lattice in a vector space over \mathbb{R} is a finitely generated subgroup which contains an \mathbb{R} -basis that is also a \mathbb{Z} -basis.

Fix once and for all a \mathbb{Z} -lattice L_0 in k^n . Let \mathcal{O}_0 be the subset of k which stabilizes L_0 . Then \mathcal{O}_0 is an order in k . Let Γ_0 be the arithmetic group consisting of the $\gamma \in G$ such that $\gamma \cdot L_0 = L_0$. Let Γ be an arithmetic subgroup of finite index in Γ_0 .

We shall identify lattices that differ only by a homothety, namely L and hL for $h \in H$. Let Y be the set of all \mathbb{Z} -lattices in S^n that are stable under \mathcal{O}_0 and are isomorphic to L_0 as \mathcal{O}_0 -module. Then we may identify Y with $H \backslash G / \Gamma_0$, where for $g \in G$ the lattice $L = gL_0$ corresponds to the coset $g\Gamma_0$. Give Y the topology coming from $H \backslash G / \Gamma_0$.

For any subset $A \subseteq L = gL_0$, the S -span $S \cdot A$ is a free S -module whose S -rank equals the dimension over k of the k -vector space spanned by $g^{-1}A$.

(2.3). The group K acts on Y , and in fact

$$K \backslash Y = X / \Gamma_0,$$

since both are $(HK) \backslash G / \Gamma_0$. However, the points of view on the two sides of this equation are different: $K \backslash Y$ is a space of lattices modulo rotations, while X / Γ_0 is a homogeneous space modulo an arithmetic group. We now make explicit how to go back and forth between the two sides.

Definition. A *marked lattice* in S^n is a function $f : L_0 \rightarrow S^n$ of the form $f(\mathbf{x}) = g\mathbf{x}$ for $g \in G$. We shall identify marked lattices that differ only by a homothety, i.e. f and hf for $h \in H$.

Let $Y' = H \backslash G$ be the set of marked lattices. A marked lattice f gives rise to an ordinary lattice $L \in Y$ by setting $L = f(L_0)$. This realizes the projection $Y' = H \backslash G \rightarrow Y = H \backslash G / \Gamma_0$. On the other hand, $f(\mathbf{x}) = g\mathbf{x}$ gives a point $(HK)g \in X$, which realizes the projection $Y' = H \backslash G \rightarrow X = (HK) \backslash G$. The right action of $g_1 \in G$ on Y' sends $f : \mathbf{x} \mapsto g\mathbf{x}$

to $\mathbf{x} \mapsto gg_1\mathbf{x}$. The left action of $k_1 \in K$ on Y' sends f to $k_1 \cdot f$. Diagram (2.3.1) shows these spaces.

$$(2.3.1) \quad \begin{array}{ccc} & Y' = H \backslash G & \\ & \text{(marked lattices)} & \\ \swarrow & & \searrow \\ X = HK \backslash G & & Y = H \backslash G / \Gamma_0 \\ \text{(homogeneous space)} & & \text{(lattices)} \\ \searrow & & \swarrow \\ & X/\Gamma_0 = K \backslash Y & \\ & \text{(arithmetic quotient)} & \end{array}$$

(2.4). Let f be a marked lattice, fixed within its homothety class. Let $L = f(L_0)$. The *arithmetic minimum* of f is defined to be $m(f) = \min\{\|\mathbf{a}\| \mid \mathbf{a} \in L - \{0\}\}$. This number is positive. The set of *minimal vectors* of f is defined to be $M(f) = \{\mathbf{a} \in L \mid \|\mathbf{a}\| = m(L)\}$.

The definitions of $m(f)$ and $M(f)$ use only the image L of f , so they descend to functions $m(L)$ and $M(L)$ on Y . On the other hand, $m(f)$ and $M(f)$ are K -equivariant, since K preserves $\|\dots\|$.

Unless otherwise specified, we normalize f (resp. L) in its homothety class so that $m(f)$ (resp. $m(L)$) equals 1.

(2.5). Definition. A marked lattice f is *well-rounded* if $M(f)$ spans S^n as S -module.

By the Γ_0 -invariance and K -equivariance noted in (2.4), we get a definition of well-rounded elements in all the spaces in (2.3.1). For instance, a point in X is well-rounded if and only if, for the class of marked lattices $f \bmod K$ that represents it, one (and hence each) of the $M(f)$ spans S^n as S -module. The definitions in X/Γ_0 and Y are similar.

Definition. W is the set of well-rounded elements in X . We call this the *well-rounded retract* in X . For any arithmetic group $\Gamma \subseteq \Gamma_0$, W/Γ is called the *well-rounded retract* in X/Γ .

In Section 4, we will explain why these are deformation retracts.

(2.6). Example of $\mathrm{GL}_n(\mathbb{Z})$. As in (1.6), take $k = \mathbb{Q}$ and $G = \mathrm{GL}_n(\mathbb{R})$. If L_0 is the standard lattice $\mathbb{Z}^n \subset \mathbb{R}^n$, then $\Gamma_0 = \mathrm{GL}_n(\mathbb{Z})$, and Y is the set of all rank- n lattices in $\mathbb{R}^n \bmod$ homotheties. The product \langle, \rangle is the standard dot product.

Let's work out the $\mathrm{GL}_2(\mathbb{Z})$ case in detail. For each marked lattice $f : L_0 \rightarrow \mathbb{R}^2$, rotate the image until $f((1,0))$ lies on the positive x -axis, and use a homothety to make $\|f((1,0))\| = 1$. This identifies X with the upper half-plane \mathfrak{h} by sending $x = HK \begin{pmatrix} 1 & a \\ 0 & b \end{pmatrix}$ (with $b > 0$) to the point $a + bi \in \mathfrak{h}$. The well-rounded retract W is the tree shown in the figure below. The lower branches break in two infinitely many times as they approach the

horizontal axis $\{b = 0\}$, which is not pictured.

The group G is generated by $\mathrm{SL}_2(\mathbb{R})$, H , and $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Under our identification of X with \mathfrak{h} , $\mathrm{SL}_2(\mathbb{R})$ acts on \mathfrak{h} by linear fractional transformations as usual, H acts trivially, and $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ acts by $a + bi \mapsto -a + bi$. The well-known fundamental domain for the action of $\mathrm{SL}_2(\mathbb{Z})$ on \mathfrak{h} is $F = \{a + bi \in \mathfrak{h} \mid |a| \leq \frac{1}{2}, a^2 + b^2 \geq 1\}$, shown with hatching. A fundamental domain for $\mathrm{GL}_2(\mathbb{Z})$ is the right-hand half of F , namely $F^* = \{a + bi \in F \mid a \geq 0\}$, shown with double hatching. The space X/Γ_0 is homeomorphic to F^* . The retract W/Γ_0 is identified with the bottom arc of F^* .

SECTION 3—THE BOREL-SERRE COMPACTIFICATION

In this section we recall the construction of the Borel-Serre manifold with corners \bar{X} , and we summarize facts about flags and parabolic subgroups. The main goal of the section is Proposition 3.6, which relates the geodesic action on X to the idea of rescaling a lattice in directions given by a fixed flag \mathcal{F} .

(3.1). Recall that ${}_{\mathbb{Q}}\mathbf{T} = \{\mathrm{diag}(a_1, \dots, a_n)\}$ is a maximal \mathbb{Q} -split torus in \mathbf{G} . The set of \mathbb{Q} -roots of \mathbf{G} with respect to ${}_{\mathbb{Q}}\mathbf{T}$ is $\{a_i a_j^{-1} \mid 1 \leq i \neq j \leq n\}$. Fix the fundamental system of simple roots $\Delta = \{a_i a_{i+1}^{-1} \mid i = 1, \dots, n-1\}$.

(3.2). We recall some facts about parabolic \mathbb{Q} -subgroups and flags. Let $\{e_1, \dots, e_n\}$ be the standard k -basis of k^n . Let $J \subseteq \Delta$. Form a graph with vertices in $\{1, \dots, n\}$ and with an edge joining vertices i and $i+1$ if and only if $a_i a_{i+1}^{-1} \in J$. The graph has $l = \#(\Delta - J) + 1$ connected components. Say the j -th component in order from left to right has ν_j vertices. Consider the flag $\mathcal{F}_J = \{0 \subsetneq V_1 \subsetneq \dots \subsetneq V_l = k^n\}$ in which V_j is spanned by $\{e_1, \dots, e_{\nu_1 + \dots + \nu_j}\}$. The standard parabolic \mathbb{Q} -subgroup \mathbf{P}_J is the subgroup of \mathbf{G} such that $\mathbf{P}_J(\mathbb{Q})$ is the stabilizer of \mathcal{F}_J in $\mathbf{G}(\mathbb{Q})$. In the coordinates of (1.2), \mathbf{P}_J has the usual block upper-triangular form, with diagonal blocks of sizes $\nu_1 \times \nu_1, \dots, \nu_l \times \nu_l$. Every parabolic \mathbb{Q} -subgroup \mathbf{P} of \mathbf{G} is conjugate by some $g \in \mathbf{G}(\mathbb{Q})$ to a unique \mathbf{P}_J .

Let $P_J = \mathbf{P}_J(\mathbb{R})$, and $P = \mathbf{P}(\mathbb{R})$ in general.

For $g \in \mathbf{G}(\mathbb{Q})$, the group of \mathbb{Q} -points of $g^{-1}\mathbf{P}_Jg$ is the stabilizer of the flag $\mathcal{F} = g^{-1}\mathcal{F}_J$. Any flag of the latter form is called a \mathbb{Q} -flag. There is a one-to-one correspondence between the parabolic \mathbb{Q} -subgroups and the \mathbb{Q} -flags they stabilize. We write $\mathcal{F}' \supseteq \mathcal{F}$ if every V_j belonging to \mathcal{F} is also a member of \mathcal{F}' .

For the \mathbb{Q} -flag $\mathcal{F} = \{0 \subsetneq V_1 \subsetneq \cdots \subsetneq V_l = k^n\}$, we set $\#\mathcal{F} = l$. Thus $\#\mathcal{F} = n$ when P is a Borel subgroup, $\#\mathcal{F} = 2$ for maximal proper P , and $\#\mathcal{F} = 1$ when $P = G$.

(3.3). Let ${}_{\mathbb{Q}}\mathbf{T}_J$ be the subgroup of ${}_{\mathbb{Q}}\mathbf{T}$ in which the i -th and $(i+1)$ -st diagonal entries are equal whenever $t_i t_{i+1}^{-1} \in J$. Let T_J be the identity component of ${}_{\mathbb{Q}}\mathbf{T}_J(\mathbb{R})$, and let $A_J = H \backslash T_J$.

Let $P = g^{-1}P_Jg$ for $g \in \mathbf{G}(\mathbb{Q})$. There is a semi-direct product decomposition $P = M_P T_P U_P$. Here U_P is the unipotent radical of P , $T_P = g^{-1}T_Jg$, and $M_P T_P$ is the unique Levi subgroup of P which is stable under the Cartan involution of G which fixes $g^{-1}Kg$. We set $A_P = H \backslash T_P$.

When $P \subseteq P'$, $A_{P'}$ is naturally identified with a subgroup of A_P by taking the kernel of appropriate roots in Δ .

(3.4). We recall the definition of the geodesic action [B-S, §3]. Let $P = g^{-1}P_Jg$ for $g \in \mathbf{G}(\mathbb{Q})$. Let Z be the identity component of the center of $M_P T_P$. Let $y \in X$. There is a $p \in P$ with $y = HKgp$. For any $z \in Z$, define

$$y \circ z = HKgzp.$$

The right side is independent of the choices of g and p . This action of Z is the *geodesic action* on X associated to P . It is a proper and free action, and it commutes with the ordinary action of P . The group A_P is contained in $H \backslash Z$, so it operates on X by the geodesic action.

(3.5). We recall the construction of \bar{X} [B-S, §§5–7]. The roots in $\Delta - J$ determine an isomorphism between A_J and $(0, \infty)^{l-1}$, sending the point

$$\text{diag}(\underbrace{a_1, \dots, a_1}_{\nu_1}, \dots, \underbrace{a_l, \dots, a_l}_{\nu_l}) \quad (a_j > 0, \text{ mod } H)$$

to $(a_1 a_2^{-1}, \dots, a_{l-1} a_l^{-1}) \in (0, \infty)^{l-1}$. Let \bar{A}_J be the partial compactification of A_J corresponding to $[0, \infty)^{l-1}$ under this isomorphism. The *corner* $X(P_J)$ associated to P_J is the fiber product $X(P_J) = X \times^{A_J} \bar{A}_J$, where A_J acts on X by the geodesic action. Conjugating by an element of $\mathbf{G}(\mathbb{Q})$, we define \bar{A}_P and a corner $X(P) = X \times^{A_P} \bar{A}_P$ for any P .

Whenever $P \subseteq P'$, there is a map $X(P') \hookrightarrow X(P)$ induced by the inclusion on X ; we identify $X(P')$ with its image under this map. As a set, \bar{X} is the union of the $X(P)$ over all P . One puts on \bar{X} the topology such that the original topology on each corner is preserved, \bar{X} is Hausdorff, and Γ acts properly on \bar{X} with a compact quotient.

Let $e(P)$ be the set corresponding to $X \times^{A_P} \{0\}^{l-1}$ in the corner $X(P) = X \times^{A_P} \bar{A}_P$. As a set, \bar{X} is the disjoint union of X and the $e(P)$ for all proper parabolic \mathbb{Q} -subgroups P . We call $e(P)$ the *boundary face corresponding to P* .

Let $\bar{e}(P)$ be the closure of $e(P)$ in \bar{X} . We have $\bar{e}(P) = \coprod_{P' \subseteq P} e(P')$. The sets $\bar{e}(P_1)$ and $\bar{e}(P_2)$ are disjoint unless $P_1 \cap P_2$ is a parabolic \mathbb{Q} -subgroup P_3 , in which case $\bar{e}(P_1) \cap \bar{e}(P_2) = \bar{e}(P_3)$.

Notice that the largest subgroup of Γ which stabilizes $X(P)$, $e(P)$, and $\bar{e}(P)$ is $\Gamma \cap P$.

Modding out by the geodesic action of A_P gives a $(\Gamma \cap P)$ -equivariant bundle map $q_P : X(P) \rightarrow e(P)$. The orbits in $X(P)$ under $M_P U_P$ are sections of this bundle called the *canonical cross-sections*.

(3.6). We now present a central tool of the paper, an interpretation of the geodesic action in the setting of marked lattices.

Let f be a marked lattice. Let $\mathcal{F} = \{0 = V_0 \subsetneq \cdots \subsetneq V_l = k^n\}$ be a \mathbb{Q} -flag, with $P = g^{-1}P_J g$ the parabolic \mathbb{Q} -subgroup determined by \mathcal{F} . Throughout the paper, we will abuse notation by writing $f(V_j \cap L_0) \otimes_{\mathbb{Q}} \mathbb{R} \subseteq S^n$ as $f(V_j)$; this is an S -linear subspace of S^n , of S -rank equal to $\dim_k V_j$. Set \check{V}_j equal to the orthogonal complement of $f(V_{j-1})$ in $f(V_j)$ with respect to \langle, \rangle . We have $S^n = \check{V}_1 \oplus \cdots \oplus \check{V}_l$ as an orthogonal direct sum.

Definition. Let $\mathfrak{A}_{\mathcal{F},f}^*$ be the group of \mathbb{R} -linear maps $S^n \rightarrow S^n$ which act by positive real homotheties on each summand \check{V}_j separately. Let $\mathfrak{A}_{\mathcal{F},f} = \mathfrak{A}_{\mathcal{F},f}^*/H$. We call $\mathfrak{A}_{\mathcal{F},f}$ the *orthogonal scaling group with respect to \mathcal{F} and f* .

Note that every $\alpha \in \mathfrak{A}_{\mathcal{F},f}^*$ is S -linear. The group $\mathfrak{A}_{\mathcal{F},f}$ acts on Y' via $f_1 \mapsto \alpha \cdot f_1$. However, it does not act on X . (The next proposition will show that $\mathfrak{A}_{\mathcal{F},f}$ acts on $Y' = H \backslash G$ as a lift of the action of A_P on $X = HK \backslash G$, but this lift is only determined by an arbitrary choice of an element of K . The choice is encoded in f .)

Let $\Psi : Y' = H \backslash G \rightarrow X = HK \backslash G$ be the projection. Let $y = \Psi(f)$. Let $\iota : A_P \rightarrow \mathfrak{A}_{\mathcal{F},f}$ be the following map: if $z \in A_P$ with

$$(3.6.1) \quad z = g^{-1} \operatorname{diag}(\underbrace{a_1, \dots, a_1}_{\nu_1}, \dots, \underbrace{a_l, \dots, a_l}_{\nu_l})g \quad (a_j > 0),$$

let $\iota(z)$ be the element of $\mathfrak{A}_{\mathcal{F},f}$ which (for each $j = 1, \dots, l$) acts on \check{V}_j by the scalar a_j . It is clear that ι is an isomorphism of groups.

Proposition. For any $z \in A_P$, with ι as above,

$$(3.6.2) \quad y \circ z = \Psi(\iota(z) \cdot f).$$

Remark. The proposition is stated in [G, (2.4)], without proof.

Proof. We use the notation of (3.4). Write $y = HKgp$ and $p = g^{-1}p_J g$ for $p_J \in P_J$. Consider the marked lattice $f_1 : \mathbf{x} \mapsto gp\mathbf{x} = p_J g\mathbf{x}$. Since $\Psi(f_1) = y = \Psi(f)$, we may choose a $k_1 \in K$ such that $f_1 = k_1 \cdot f$.

Write $z \in A_P$ as $z = g^{-1}z_J g$ for $z_J = \operatorname{diag}(a_1, \dots, a_1, \dots, a_l, \dots, a_l) \in A_J$. Then

$$y \circ z = HKgzp = HKz_J p_J g.$$

The marked lattice $f_z : \mathbf{x} \rightarrow z_J p_J g \mathbf{x}$ descends mod K to $y \circ z$.

Let \mathcal{I}_J be the flag in S^n whose j -th member is the coordinate subspace spanned by $\{e_1, \dots, e_{\nu_1 + \dots + \nu_j}\}$. (This \mathcal{I}_J is essentially $\mathcal{F}_J \otimes_{\mathbb{Q}} \mathbb{R}$.) The map $\mathbf{x} \mapsto g \mathbf{x}$ carries \mathcal{F} to \mathcal{I}_J , and the further multiplication by p_J and z_J preserves \mathcal{I}_J . Thus for any $z \in A_P$, $f_z(V_j)$ is exactly the j -th member of \mathcal{I}_J . (The idea is that, out of the whole K -equivalence class of marked lattices representing $y \circ z$, f_z is a representative for which the $f_z(V_j)$ will be in standard position.) The orthocomplement of $f_z(V_{j-1})$ in $f_z(V_j)$ is the coordinate subspace E_j spanned by $\{e_{\nu_1 + \dots + \nu_{j-1} + 1}, \dots, e_{\nu_1 + \dots + \nu_j}\}$. The E_j are orthogonal, and z_J acts by the map $\alpha_1 : S^n \rightarrow S^n$ characterized by the property that it acts on each E_j by the scalar a_j . In short, the left-hand side of (3.6.2) is the projection under Ψ of the marked lattice given by the composition $L_0 \xrightarrow{f_1} S^n \xrightarrow{\alpha_1} S^n$.

On the right-hand side of (3.6.2), observe that $\Psi(\alpha \cdot f) = \Psi(k_1 \alpha \cdot f) = \Psi((k_1 \alpha k_1^{-1}) f_1)$. Since $f_1 = k_1 \cdot f$, the orthogonal complement of $k_1 \cdot f(V_{j-1})$ in $k_1 \cdot f(V)$ equals that of $f_1(V_{j-1})$ in $f_1(V_j)$; this says exactly that $k_1 \cdot \check{V}_j = E_j$ for all j . It is immediate that $k_1 \alpha k_1^{-1}$ is the same map as α_1 . Thus the right-hand side of (3.6.2) is also Ψ applied to the composition $L_0 \xrightarrow{f_1} S^n \xrightarrow{\alpha_1} S^n$. \square

Note. We will always normalize elements of A_P and $\mathfrak{A}_{\mathcal{F},f}$ in their classes modulo H so that $a_1 = 1$.

(3.7). While A_P and $\mathfrak{A}_{\mathcal{F},f}$ act on different spaces, we use similar notation for the two actions. We write

$$\boldsymbol{\rho} = (\rho_1, \dots, \rho_{l-1}) \in (0, \infty)^{l-1}$$

for a point in either group. The ρ_i are the coordinates coming from the simple roots in J : that is, $\rho_j = a_j/a_{j+1}$ for $j = 1, \dots, l-1$. By Proposition 3.6, the point of $\mathfrak{A}_{\mathcal{F},f}$ corresponding to $(\rho_1, \dots, \rho_{l-1})$ acts on \check{V}_j by the scalar

$$(3.7.1) \quad a_j = (\rho_1 \cdots \rho_{j-1})^{-1} \quad (j = 2, \dots, l).$$

(Because of our normalizations, the action on \check{V}_1 is by 1.)

Remark. The ρ_j 's are good coordinates for rescaling sublattices within lattices, while the a_j are the natural coordinates for the geodesic action. Though (3.7.1) is messy, we cannot avoid using it, since part of our goal is to relate lattices and the geodesic action. The inverses in (3.7.1) arise because the well-rounded retraction goes away from the Borel-Serre boundary.

Multiplication in A_P corresponds to coordinate-wise multiplication of the ρ_i . As in (3.5), the ρ_j extend to coordinates on \bar{A}_P with $(\rho_1, \dots, \rho_{l-1}) \in [0, \infty)^{l-1}$. Going to infinity in $X(P)$ means going to zero in the ρ_j variables. We introduce a partial ordering on A_P and \bar{A}_P , where $(\rho_1, \dots, \rho_{l-1}) \leq (\rho'_1, \dots, \rho'_{l-1})$ if and only if $\rho_j \leq \rho'_j$ for each j .

The action of $\boldsymbol{\rho} = (\rho_1, \dots, \rho_{l-1})$ on Y' is denoted $f \mapsto \boldsymbol{\rho} \cdot f$. This descends mod Γ_0 to an action $L \mapsto \boldsymbol{\rho} \cdot L$ on Y . Descending mod K as in Proposition 3.6, $\boldsymbol{\rho} \cdot x$ denotes the geodesic action on points x of X or $X/(\Gamma \cap P)$.

(3.8). We will need to consider \mathbb{Q} -flags $\mathcal{M} = \{0 \subsetneq M^{(1)} \subseteq M^{(2)} \subseteq \dots \subseteq M^{(n)} = k^n\}$ in which some of the inclusions can be equalities. If \mathcal{F} is obtained from \mathcal{M} by replacing each string of equalities with a single space V_j , we call \mathcal{F} the *irredundant version* of \mathcal{M} . If \mathfrak{A} is the orthogonal scaling group with respect to this flag and some f , we always write $\rho = (\rho_1, \dots, \rho_{l-1}) \in \mathfrak{A}$ with reference to \mathcal{F} , not \mathcal{M} .

SECTION 4—THE WELL-ROUNDED RETRACTION

In this section, we define the well-rounded retraction r_t . The material comes from [A3], though we emphasize different things (marked lattices, the flag of successive minima).

(4.1). Let f be a marked lattice, with $L = f(L_0)$. We normalize f up to homothety so that $m(f) = 1$. If $V \subseteq k^n$ is a k -subspace, $f(V) \otimes_{\mathbb{Q}} \mathbb{R}$ is abbreviated $f(V)$.

Recall that $Y' = H \backslash G$ is the space of marked lattices (2.3). For $i = 1, \dots, n$, let

$$Y'_i = \{f \in Y' \mid \text{rank}_S(S \cdot M(f)) \geq i\}.$$

Observe that $Y' = Y'_1 \supset Y'_2 \supset \dots$, and that Y'_n is the set of well-rounded elements of Y' . For $i = 1, \dots, n-1$, we will define a deformation retraction $r_t^{(i)} : Y'_i \times [0, 1] \rightarrow Y'_i$ with image Y'_{i+1} . The deformation retraction will be equivariant for both the K and Γ_0 actions. For any $f \in Y'$, we will also define a \mathbb{Q} -flag $\mathcal{M} = \{0 \subseteq M^{(1)} \subseteq \dots \subseteq M^{(n)} = k^n\}$ depending on f . (This \mathcal{M} could have equalities, as in (3.8).) The definitions proceed simultaneously by induction on i .

Let $f_1 = f$, and for $i > 1$ let $f_i = r_1^{(i-1)}(f_{i-1})$ inductively. Let $M^{(i)}$ be the k -span of $f_i^{-1}(M(f_i))$. This is a subspace of k^n of dimension $\geq i$. If f_i is in Y'_{i+1} already, set $r_t^{(i)}(f_i) = f_i$ for all $t \in [0, 1]$, and set $\mu_i = 1$. Otherwise, consider the S -linear maps φ_μ obtained by multiplying in $f_i(M^{(i)})$ by 1 and in the orthocomplement $f_i(M^{(i)})^\perp$ by a scalar $\mu > 0$. By [A3, p. 463], there is a unique smallest $\mu_i \in (0, 1)$ such that

$$(4.1.1) \quad \varphi_{\mu_i} \cdot f_i \text{ still has arithmetic minimum equal to 1.}$$

This $\varphi_{\mu_i} \cdot f_i$ lies in Y'_{i+1} . We set $r_t^{(i)}(f_i) = \varphi_{1+(\mu_i-1)t} \cdot f_i$. This is continuous in both the f_i and t variables, and defines a homotopy between $r_0^{(i)} = (\text{identity on } Y'_i)$ and $r_1^{(i)} : Y'_i \rightarrow Y'_{i+1}$.

Definition. The *well-rounded retraction* r_t on Y' is obtained by composing the deformation retractions $r_t^{(i)}$ in the order $i = 1, \dots, n-1$. More precisely, $r_t : Y' \times [0, 1] \rightarrow Y'$ is given by

$$r_t(f) = r_{t(n-1)-(i-1)}^{(i)} \circ r_1^{(i-1)} \circ \dots \circ r_1^{(1)}(f),$$

where i is determined by the rule $t \in [\frac{i-1}{n-1}, \frac{i}{n-1}]$ for $i \in \{1, \dots, n-1\}$.

Let $r = r_1$. We will also call this the *well-rounded retraction*.

Notice that $M^{(i+1)} \supseteq M^{(i)}$ by construction. This gives the

Definition. The *flag of successive minima* for f is the \mathbb{Q} -flag $\mathcal{M} = \{0 \subseteq M^{(1)} \subseteq \cdots \subseteq \cdots M^{(n)} = k^n\}$.

This \mathcal{M} is an invariant of f 's class mod K , and is equivariant for Γ_0 . It is a “lower semi-continuous” function of f in the sense that for any $f \in Y'$, there is a neighborhood U of f such that for all $f' \in U$, the flag \mathcal{M}' of successive minima for f' satisfies $\mathcal{M}' \subseteq \mathcal{M}$.

(4.2). Remark. Once the inductive construction of r is completed, we obtain the following interpretation of $r(f)$ as the result of applying a geodesic action to f . Let \mathcal{M} be the flag of successive minima for f , with irredundant version \mathcal{F} . Let $\boldsymbol{\mu} = (\mu_1^{-1}, \dots, \mu_{n-1}^{-1}) \in \mathfrak{A}_{\mathcal{F}, f}$ with the μ_i depending on f as in (4.1). (As in (3.8), μ_i^{-1} is omitted when $M^{(i)} = M^{(i+1)}$.) Then

$$r(f) = \boldsymbol{\mu} \cdot f.$$

(4.3). We now adapt the definition of the well-rounded retraction to the other spaces in (2.3.1). In each case, the retraction will still be denoted r_t for $t \in [0, 1]$, with $r = r_1$, and will be called the “well-rounded retraction”.

The whole construction is Γ_0 - and K -equivariant. Hence r_t descends to deformation retractions of Y onto the set of well-rounded lattices, of X onto W , and of X/Γ onto W/Γ for any $\Gamma \subseteq \Gamma_0$ of finite index.

We recall the main theorem of [A3].

Theorem. *The map r_t is a Γ_0 -equivariant deformation retraction of X onto W . The quotient W/Γ_0 is compact. The dimension of W and W/Γ_0 is the virtual cohomological dimension of Γ_0 , which equals $\dim X - (n - 1)$. The space W has a natural structure as a cell complex on which Γ_0 acts cell-wise with finite stabilizers of cells. The first barycentric subdivision of the cell structure descends to a finite cell complex structure on W/Γ_0 .*

The example of $\mathrm{GL}_2(\mathbb{Z})$ is discussed in (6.3).

Remark. When the class number h of k is > 1 , [A3] defines a whole $(h - 1)$ -dimensional family of retracts, depending on a parameter called a *set of weights*. For simplicity, we will use only the trivial set of weights (identically equal to 1). Readers may easily make the needed changes if they wish to use a non-trivial set of weights.

SECTION 5—SUMMARY OF NOTATIONAL CONVENTIONS

The letter f always stands for a marked lattice $f : \mathbf{x} \mapsto g\mathbf{x}$ for $g \in G$, modulo the homotheties H . We let $L = f(L_0)$, sometimes without explicit mention. We let L' be the lattice for f' , etc. As in (3.6), $f(V_j \cap L_0) \otimes_{\mathbb{Q}} \mathbb{R} \subseteq S^n$ will be written $f(V_j)$. Let \check{V}_j be the orthogonal complement of $f(V_{j-1})$ in $f(V_j)$, so that $S^n = \check{V}_1 \oplus \cdots \oplus \check{V}_l$ as an orthogonal direct sum. As in (2.4), lattices are always scaled by a homothety so that their shortest non-zero vector has length 1.

For the rest of the paper, P denotes the real points of a parabolic \mathbb{Q} -subgroup. Let $\mathcal{F} = \{0 \subsetneq V_1 \subsetneq \cdots \subsetneq \cdots V_l = k^n\}$ be the \mathbb{Q} -flag for P as in (3.2), and $\mathfrak{A} = \mathfrak{A}_{\mathcal{F}, f}$ the orthogonal scaling group with respect to \mathcal{F} and f (see (3.6)–(3.7)).

Whenever we mention P , the letters \mathcal{F} and \mathfrak{A} will refer to the objects corresponding to P , and vice versa. The symbol P' is associated to \mathcal{F}' and \mathfrak{A}' , etc.

From Section 6 on, we will often write f for the point that f determines in X or X/Γ . We will speak of marked lattices f “in” X or W , or write L for the point that L determines in X/Γ . These abuses are justified because our constructions involving marked lattices are K -equivariant.

SECTION 6—THE SETS $W_{\mathcal{F}}$

(6.1). Throughout this section, $\mathcal{F} = \{0 = V_0 \subsetneq V_1 \subsetneq \cdots V_l = k^n\}$ will be a fixed \mathbb{Q} -flag.

Definition. Let $W_{\mathcal{F}}$ be the set of all marked lattices f in W such that for all $j = 1, \dots, l$,

$$S \cdot (M(L) \cap f(V_j)) = f(V_j).$$

($W_{\mathcal{F}}$ was called W' in the introduction.) Geometrically, $W_{\mathcal{F}}$ consists of the lattices $L = f(L_0)$ whose intersection with each subspace $f(V_j)$ is well-rounded in that subspace. By the definition of the cell structure on W [A3, pp. 466–7], $W_{\mathcal{F}}$ is a closed subcomplex of W and is stable under $\Gamma \cap P$. A cell in $W_{\mathcal{F}}$ is said to *respect* \mathcal{F} . If $\mathcal{F} = \gamma\mathcal{F}'$ for $\gamma \in \Gamma$, then the images of $W_{\mathcal{F}}$ and $W_{\mathcal{F}'}$ in W/Γ are equal. The \mathbb{Q} -flags fall into only finitely many equivalence classes mod Γ , so only finitely many distinct subcomplexes of W/Γ arise in this way.

If \mathcal{M} is a \mathbb{Q} -flag with possible equalities (3.8), and \mathcal{F} is the irredundant version of \mathcal{M} , let $W_{\mathcal{M}} = W_{\mathcal{F}}$.

(6.2). The following is immediate from the definitions in (4.1).

Lemma. *If f has flag of successive minima \mathcal{M} , then $r(f) \in W_{\mathcal{M}}$.*

(6.3). *Example of $\mathrm{GL}_2(\mathbb{Z})$.* Refer again to the picture in (2.6). For the standard flag $\mathcal{F} = \{0 \subsetneq \mathbb{Q} \cdot (1, 0) \subsetneq \mathbb{Q}^2\}$ with associated P , the space $W_{\mathcal{F}}$ is the horizontal sequence of arcs forming the top of W . The boundary face $e(P)$ in \bar{X} (not shown) is the horizontal line at $b = \infty$. The geodesic action by A_P flows straight up, pushing X upward to converge to $e(P)$. The well-rounded retraction pulls the region above W straight down onto $W_{\mathcal{F}}$. For torsion-free Γ , $e(P)/(\Gamma \cap P)$ and $W_{\mathcal{F}}/(\Gamma \cap P)$ are homeomorphic to circles, and the well-rounded retraction induces the obvious isomorphism between the cohomology of these spaces. (The situation is more complicated for $n > 2$, when the boundary faces are no longer disjoint.)

Remark. For small n , the $W_{\mathcal{F}}/(\Gamma \cap P)$ are all readily computable. See (10.6).

SECTION 7—NEIGHBORHOODS OF INFINITY IN \bar{X}

We now prove our main results explaining the connection between the well-rounded retraction r , the geodesic action, and the topology of the Borel-Serre boundary. Parts (7.1)–(7.4) give a chain of propositions that allow us to define a tubular neighborhood $\bar{N}_{\mathcal{F}}$ of $e(P)$ (called \bar{N} and e in the introduction). In (7.5)–(7.6) we prove that $r : X \rightarrow W$ has the continuous extension $\bar{r} : \bar{X} \rightarrow W$, where (for all \mathcal{F}) \bar{r} is constant on the fibers of $\bar{N}_{\mathcal{F}}$ and carries $\bar{N}_{\mathcal{F}}$ onto $W_{\mathcal{F}}$.

(7.1). Fix a marked lattice f , with $L = f(L_0)$. As in (4.1), let $\mathcal{M} = \{0 \subseteq M^{(1)} \subseteq \dots \subseteq M^{(n)} = k^n\}$ be the flag of successive minima for f . Let \mathcal{F} be the irredundant version of \mathcal{M} , with associated P . Let $\mathfrak{A} = \mathfrak{A}_{\mathcal{F}, f}$. Let $f'' = r(f)$ be the image of f under the well-rounded retraction.

The idea of the following lemma is that if f is close enough to $e(P)$, then the well-rounded retraction carries the whole orthant $\{\boldsymbol{\rho} \in \mathfrak{A} \mid \boldsymbol{\rho} \leq 1\} \cdot f$ to a single point of W . The lemma says that, to know that f is “close enough” to $e(P)$, it suffices to know that f ’s flag of successive minima is the flag \mathcal{F} corresponding to P (or at least contains this \mathcal{F} —see (7.2)).

Lemma. *Let f , \mathcal{M} , \mathcal{F} , P , and \mathfrak{A} be as above. For any $\boldsymbol{\rho} = (\rho_1, \dots, \rho_{l-1}) \in \mathfrak{A}$ with $\boldsymbol{\rho} \leq (1, \dots, 1)$, $f' = \boldsymbol{\rho} \cdot f$ satisfies $r(f') = r(f) = f''$. The common image f'' will lie in $W_{\mathcal{M}} = W_{\mathcal{F}}$.*

Proof. It suffices to prove the lemma when $\boldsymbol{\rho} = (1, \dots, 1, \rho_j, 1, \dots, 1)$ for a fixed $\rho_j \leq 1$; the general case follows by applying this for each j . Let i be the unique index such that $V_j = M^{(i)}$ and $M^{(i)} \subsetneq M^{(i+1)}$.

The marked lattices f and f' agree on $L_0 \cap M^{(i)}$, since $\boldsymbol{\rho}$ acts by 1 in that region. For $\mathbf{x} \in L_0$ with $\mathbf{x} \notin M^{(i)}$, no vector $f'(\mathbf{x})$ is shorter than the corresponding vector $f(\mathbf{x})$, since $\boldsymbol{\rho}$ magnifies lengths in the directions perpendicular to $f(M^{(i)})$ by $\rho_j^{-1} \geq 1$. And $f(\mathbf{x})$ cannot be a minimal vector of f , since the minimal vectors lie in $f(M^{(1)})$, which is contained in $f(M^{(i)})$. It follows that the well-rounded retractions for f and f' will proceed exactly the same up through the end of the $r^{(i-1)}$ step—that is, if μ_ι and μ'_ι (for f and f' resp.) are as in (4.1), then

$$(7.1.1) \quad \mu_\iota = \mu'_\iota \text{ for all } \iota < i.$$

Set $g = r_1^{(i-1)} \circ \dots \circ r_1^{(1)}(f)$ and $g' = r_1^{(i-1)} \circ \dots \circ r_1^{(1)}(f')$. It follows from (3.7.1) and (7.1.1) that $g' = (1, \dots, 1, \rho_j^{-1}, 1, \dots, 1) \cdot g$. The $r_t^{(i)}$ stage of the retraction will, by definition, carry g to

$$(7.1.2) \quad (1, \dots, 1, \mu_i^{-1}, 1, \dots, 1) \cdot g,$$

and carry g' to

$$(7.1.3) \quad (1, \dots, 1, \mu_i'^{-1}, 1, \dots, 1) \cdot g'.$$

However, by the uniqueness statement in (4.1.1), μ_i^{-1} must equal $\mu_i'^{-1} \rho_i$. Thus the marked lattices (7.1.2) and (7.1.3) will coincide. The $r^{(i+1)}$ and later stages of the retraction will agree on this common lattice, so the final values $r(f)$ and $r(f')$ are equal. By (6.2), $f'' \in W_{\mathcal{M}}$. \square

(7.2). We record a corollary of the previous lemma. The idea is that if f ’s flag of successive minima refines the flag \mathcal{F}' for some P' , then f is “close enough” to $e(P')$. After all, if f is close enough to $e(P)$, it should be close enough to any larger Borel-Serre boundary component $e(P')$ that has $e(P)$ in its closure.

Corollary. *Let the notation be as in (7.1). Let \mathcal{F}' be any \mathbb{Q} -flag with $\mathcal{F}' \subseteq \mathcal{F}$, and let \mathfrak{A}' be the orthogonal scaling group for \mathcal{F}' and f . Then for any $\boldsymbol{\rho} \in \mathfrak{A}'$ with $\boldsymbol{\rho} \leq (1, \dots, 1)$, $\boldsymbol{\rho} \cdot f$ and f map to the same point under the well-rounded retraction. This point lies in $W_{\mathcal{F}'}$.*

Proof. This is trivial, since \mathfrak{A}' is naturally a subgroup of \mathfrak{A} , and $W_{\mathcal{F}} \subseteq W_{\mathcal{F}'}$. \square

(7.3). In (7.1) we took an f close to $e(P)$ and studied the orthant $\{\boldsymbol{\rho} \leq 1\} \cdot f$. Now, instead, we take an arbitrary f . We show in (7.3) (actually in the corollary) that there is a $\mathbf{t} \in \mathfrak{A}$ which pushes f so close to $e(P)$ that the orthant $\{\boldsymbol{\rho} \leq \mathbf{t}\} \cdot f$ still has the property of (7.1)—namely, r maps it to a single point in $W_{\mathcal{F}}$.

Lemma. *Choose any marked lattice f . Fix a \mathbb{Q} -flag $\mathcal{F} = \{0 \subsetneq V_1 \subsetneq \dots \subsetneq V_l = k^n\}$, and let $\mathfrak{A} = \mathfrak{A}_{\mathcal{F}, f}$. Then there exists $\mathbf{t} = (t_1, \dots, t_{l-1}) \in \mathfrak{A}$, depending on f , such that for any $\boldsymbol{\rho} \in \mathfrak{A}$ with $\boldsymbol{\rho} \leq \mathbf{t}$, the flag \mathcal{M}' of successive minima for $f' = \boldsymbol{\rho} \cdot f$ satisfies $\mathcal{M}' \supseteq \mathcal{F}$.*

Proof. Let $j \in \{1, \dots, l-1\}$. Let $d_j = \dim_k V_j$. Let $\overset{\circ}{L} = f(V_j \cap L_0)$. This is a \mathbb{Z} -lattice in the S -submodule $\overset{\circ}{V} = f(V_j \otimes_{\mathbb{Q}} \mathbb{R}) \subseteq S^n$. Perform the well-rounded retraction on $\overset{\circ}{L}$ within $\overset{\circ}{V}$. As in (4.1), this involves multiplying by constants $\mu_1, \dots, \mu_{d_j-1} \leq 1$ in various subspaces of $\overset{\circ}{V}$. (We must first rescale f by a homothety so that $m(\overset{\circ}{L}) = 1$; this does not affect the value of the μ_i .) Let $\beta_j = \mu_1 \cdots \mu_{d_j-1}$.

The orthogonal projection of $L = f(L_0)$ onto $f(V_j)^\perp$ is a lattice L^\dagger under the restriction of the inner product to $f(V_j)^\perp$. Let $\alpha_j > 0$ be the length of the shortest non-zero vector of L^\dagger .

Let $t_1 = \min(1, \frac{1}{2}\alpha_1\beta_1)$, and inductively let $t_j = \min(1, \frac{1}{2}(t_1 \cdots t_{j-1})^{-1}\alpha_j\beta_j)$. This means $(t_1 \cdots t_j)^{-1} \geq 2\alpha_j^{-1}\beta_j^{-1}$ for $j = 1, \dots, l-1$.

Now let $f' = \boldsymbol{\rho} \cdot f$ for $\boldsymbol{\rho} \leq \mathbf{t}$. Fix j . Let \mathbf{a} be any vector of L lying outside $f(V_j \cap L_0)$. Let $\mathbf{b} = \boldsymbol{\rho} \cdot \mathbf{a}$ be the corresponding vector for f' . Notice that $\boldsymbol{\rho}$ multiplies \check{V}_{j+1} by $(\rho_1 \cdots \rho_j)^{-1}$, by (3.7.1). Since $\rho_j \leq 1$ for all j , $\boldsymbol{\rho}$ multiplies by factors at least as large as $(\rho_1 \cdots \rho_j)^{-1}$ in each summand of $\check{V}_{j+1} \oplus \dots \oplus \check{V}_l = f(V_j)^\perp$.

Imagine performing the well-rounded retraction on $\overset{\circ}{L}$, but carrying out all the retractions on $L' = f'(L_0)$ itself; more precisely, consider the lattice

$$(7.3.1) \quad (1, \dots, 1, \mu_{d_j-1}^{-1}, 1, \dots, 1) \cdot (1, \dots, 1, \mu_{d_j-2}^{-1}, 1, \dots, 1) \cdots (\mu_1^{-1}, 1, \dots, 1) \cdot L'$$

where the μ_i are the constants for $\overset{\circ}{L}$. (The μ_i^{-1} is in the i -th position when the minimal vectors of $\overset{\circ}{L}$ at the i -th step of the retraction span the i -th member of the irredundant version of the flag of successive minima for $\overset{\circ}{L}$.) Let $\mathbf{b}^{(j)}$ be the vector corresponding to \mathbf{b}

in (7.3.1). Observe, using (3.7.1), that

$$\begin{aligned}
\|\mathbf{b}^{(j)}\| &\geq \|(\text{orthogonal projection of } \mathbf{b}^{(j)} \text{ onto } f(V_j)^\perp)\| \\
&= \mu_1 \cdots \mu_{d_j-1} \cdot \|(\text{orthogonal projection of } \mathbf{b} \text{ onto } f(V_j)^\perp)\| \\
&= \beta_j \cdot \|(\text{orthogonal projection of } \mathbf{b} \text{ onto } f(V_j)^\perp)\| \\
&\geq \beta_j \cdot (\rho_1 \cdots \rho_j)^{-1} \cdot \|(\text{orthogonal projection of } \mathbf{a} \text{ onto } f(V_j)^\perp)\| \\
&\geq \beta_j \cdot (t_1 \cdots t_j)^{-1} \cdot \alpha_j \\
&\geq 2 \\
&> 1.
\end{aligned}$$

This says that the image of L' under the well-rounded retraction on L up through the $r^{(d_j-1)}$ stage is equal to (7.3.1): no vector outside \mathring{L} will affect the stopping times of the retraction on L' , since each will have length > 1 during the entire process leading to (7.3.1). In particular, $f(V_j)$ is a member of \mathcal{M}' , the flag of successive minima of L' , which is what we wanted to prove. \square

Corollary. *With notation as in the lemma, the points $\boldsymbol{\rho} \cdot f$ with $\boldsymbol{\rho} \leq \mathbf{t}$ map to a common point of $W_{\mathcal{F}}$ under the well-rounded retraction r .*

Proof. Apply Corollary 7.2 to the marked lattice $\mathbf{t} \cdot f$ of the lemma. \square

(7.4). We can now construct neighborhoods of the Borel-Serre boundary component $e(P)$ that are well behaved with respect to the well-rounded retraction. Fix a \mathbb{Q} -flag \mathcal{F} , with its associated P .

Recall that the corner $X(P)$ is a fiber bundle with base $e(P)$ and fiber \bar{A}_P . By a *neighborhood in X near $e(P)$* , we mean an open set in X whose intersection with every fiber contains $\{a \in A_P \mid a < a_0\}$ for some $a_0 \in A_P$ depending on the fiber.

Proposition. *There is a neighborhood $N_{\mathcal{F}}$ in X near $e(P)$ such that, on the intersection of $N_{\mathcal{F}}$ with a fiber of $X(P)$, the well-rounded retraction r takes a constant value. Furthermore, r carries $N_{\mathcal{F}}$ surjectively to $W_{\mathcal{F}}$.*

Proof. As in (3.5), let C_1 be the canonical cross-section $HK \cdot 1 \cdot M_P U_P$. Let f_1 be a marked lattice corresponding to a point in C_1 . For this f_1 , let $\mathfrak{A} = \mathfrak{A}_{\mathcal{F}, f_1}$, and let $\mathbf{t}_1 \in \mathfrak{A}$ be the point constructed in Lemma 7.3. We define

$$(7.4.1) \quad N_{\mathcal{F}} = \{\boldsymbol{\rho}_1 \cdot f_1 \mid \boldsymbol{\rho}_1 < \mathbf{t}_1\},$$

where f_1 ranges over all marked lattices representing points of C_1 , and \mathbf{t}_1 depends on f_1 . For any given f_1 , the map r is constant on the set $\{\boldsymbol{\rho}_1 \cdot f_1\} = N_{\mathcal{F}} \cap (\text{fiber})$, by Corollary 7.3.

To prove $N_{\mathcal{F}}$ is an open set in X , we must show the t_j vary continuously with f_1 . The μ_i of (4.1) vary continuously as a function of f , by [A3, §3(i)]. Hence the β_j of (7.3) vary continuously with f . For any continuous family of lattices (not necessarily normalized

by $m(L) = 1$), the function $m(L)$ is also continuous in L ; applying this to L^\dagger (defined in (7.3)), we see the α_j of (7.3) are continuous in f . Hence the t_j are continuous in f .

By Corollary 7.2, r carries $N_{\mathcal{F}}$ into $W_{\mathcal{F}}$. In the rest of the proof, we prove the map is surjective. Let f be a marked lattice representing a point in $W_{\mathcal{F}}$. Let $d_j = \dim_k V_j$. Let f_1 be the marked lattice corresponding to the point in C_1 in the A_P -fiber through f ; let $\rho \cdot f = f_1$. Choose \mathbf{t} for f_1 as in Lemma 7.3. For a sufficiently small $\tau \in (0, 1)$, $\tau = (\tau, \dots, \tau)$ satisfies $\tau < \mathbf{t} \cdot \rho$, implying $\tau \cdot f \in N_{\mathcal{F}}$. We will show r carries $\tau \cdot f$ back to f .

The lattice for $\tau \cdot f$ is the lattice for f scaled by a factor of $1/\tau^{j-1}$ on \check{V}_j . The minimal vectors of f span their intersection with $f(V_1)$, by the definition of $W_{\mathcal{F}}$. Since $1/\tau > 1$, the minimal vectors of $\tau \cdot f$ are exactly the minimal vectors of f that lie in $f(V_1)$. Hence the minimal vectors of $\tau \cdot f$ lie in $f(V_1)$ and span $f(V_1)$. This means the steps $r^{(1)}, \dots, r^{(d_1-1)}$ of the well-rounded retraction are trivial.

The retraction $r^{(d_1)}$ fixes $f(V_1)$ and acts by a scalar $\mu_{d_1} \leq 1$ on all orthogonal directions. Consider what happens as a scalar μ descends from 1 to μ_{d_1} . The marked lattice $\tau \cdot f$ becomes

$$(7.4.2) \quad (\mu^{-1}\tau, \tau, \dots, \tau) \cdot f.$$

As long as $1/(\mu^{-1}\tau) > 1$, there are no new minimal vectors. But when $\mu^{-1}\tau = 1$, the minimal vectors of (7.4.2) are exactly the minimal vectors of f that lie in $f(V_2)$. These span $f(V_2)$ (again by the definition of $W_{\mathcal{F}}$), which has rank bigger than the rank of $f(V_1)$. So by (4.1.1) for $i = d_1$, we must have $\mu_{d_1} = \tau$, and at the end of the $r^{(d_1)}$ stage we have produced $(1, \tau, \dots, \tau) \cdot f$. The well-rounded retraction goes on similarly to produce $(1, 1, \tau, \dots, \tau) \cdot f, \dots$, and finally $(1, \dots, 1) \cdot f = f$. \square

(7.5). The neighborhood $N_{\mathcal{F}}$ defined in Proposition 7.4 lies in X , the interior of \bar{X} . We now extend it to a neighborhood $\bar{N}_{\mathcal{F}}$ of $e(P)$ in the corner $X(P)$. As in (3.7), the identification $\mathfrak{A} = \mathfrak{A}_{\mathcal{F}, f_1} \cong A_P$ extends to an identification $\bar{\mathfrak{A}} = \bar{\mathfrak{A}}_{\mathcal{F}, f_1} \cong \bar{A}_P$, where $\bar{\mathfrak{A}}$ is obtained from \mathfrak{A} by allowing one or more of the ρ_j in $(\rho_1, \dots, \rho_{j-1}) \in \mathfrak{A}$ to become 0. With the notation of (7.4.1),

Definition. $\bar{N}_{\mathcal{F}} = \{f = \rho_1 \cdot f_1 \mid \rho_1 \in \bar{\mathfrak{A}}_{\mathcal{F}, f_1}, \rho_1 < \mathbf{t}_1\}$, where f_1 ranges over all marked lattices representing points of C_1 , and \mathbf{t}_1 depends on f_1 as in Lemma 7.3.

It is clear that $\bar{N}_{\mathcal{F}}$ is a tubular neighborhood of $e(P)$ in $X(P)$, in the sense that $\bar{N}_{\mathcal{F}}$ is homeomorphic to $e(P) \times [0, 1)^{l-1}$ with $e(P)$ included as $e(P) \times (0, \dots, 0)$. Also, the set $\bar{N}_{\mathcal{F}}$ is $(\Gamma_0 \cap P)$ -invariant.

(7.6). We now define a map $\bar{r} : \bar{X} \rightarrow W$ that continuously extends the well-rounded retraction r . Choose $x \in \bar{X}$. If $x \in X$, let $\bar{r}(x) = r(x)$. Otherwise, $x \in e(P)$ for a unique P with associated \mathcal{F} . Hence $x \in \bar{N}_{\mathcal{F}}$. Consider the fiber φ' of $\bar{N}_{\mathcal{F}}$ over x . By Proposition 7.4, r takes a constant value on $\varphi' \cap X$. Define $\bar{r}(x)$ to be this common value $r(\varphi' \cap X)$.

Proposition. *The map \bar{r} is a continuous Γ -equivariant extension of r to \bar{X} . For any \mathbb{Q} -flag \mathcal{F} , it is constant on the \bar{A}_P -fibers of $\bar{N}_{\mathcal{F}}$. It is a retraction $\bar{X} \rightarrow W$ (i.e. it fixes W pointwise).*

Proof. It suffices to prove \bar{r} is constant on φ' . For then, the continuity of \bar{r} on $\bar{N}_{\mathcal{F}}$ will follow from the tubular neighborhood structure and the continuity of r on $N_{\mathcal{F}}$. The continuity of \bar{r} on \bar{X} will follow from its continuity on X and on each $\bar{N}_{\mathcal{F}}$, because these form an open cover of \bar{X} . The Γ -equivariance of \bar{r} will follow from that of r .

Let y lie on the fiber φ' of $\bar{N}_{\mathcal{F}}$ over x , with $y \neq x$. If $y \in X$, then $\bar{r}(x) = r(x) = \bar{r}(y)$ by construction. Next, suppose $y \in e(P_1)$ for some P_1 with associated \mathcal{F}_1 . Then $P_1 \supseteq P$. Now A_{P_1} is naturally a subgroup of A_P by (3.3), and $\bar{A}_{P_1} \subset \bar{A}_P$. So φ' fibers over $\bar{e}(P_1)$ with fibers given by the action of \bar{A}_{P_1} . Let φ'_1 be the intersection of φ' with the \bar{A}_{P_1} -fiber of $\bar{N}_{\mathcal{F}_1}$ over y . Then

$$\begin{aligned} \bar{r}(y) &= r(\varphi'_1 \cap X) && \text{by def. of } \bar{r}(y) \\ &= r(\varphi' \cap X) && \text{since } \varphi'_1 \cap X \subseteq \varphi' \cap X, \text{ and} \\ & && r \text{ is constant on the latter} \\ &= \bar{r}(x) && \text{by def. of } \bar{r}(x). \end{aligned}$$

This proves \bar{r} is constant on the \bar{A}_P -fibers of $\bar{N}_{\mathcal{F}}$.

Finally, \bar{r} fixes W pointwise because r does. \square

SECTION 8—EXTENDING THE DEFORMATION RETRACTION TO THE BOREL-SERRE BOUNDARY

(8.1). We recall from [S] some facts about Saper's tiling of \bar{X} . Saper constructs a Γ -equivariant subspace X_0 of X called the *central tile*. This is a codimension-zero submanifold with corners in X . The closed boundary faces $\partial^P X_0$ are in one-to-one correspondence with the proper parabolic \mathbb{Q} -subgroups P ; inclusions $P \subseteq P'$ correspond to inclusions $\partial^P X_0 \subseteq \partial^{P'} X_0$, and conversely. Every $y \in \partial X_0$ lies on $\partial^P X_0$ for a unique smallest P . There is a Γ -equivariant piecewise-analytic deformation retraction $r_S : \bar{X} \rightarrow X_0$; more precisely, r_S fits into a piecewise-analytic family $r_{t,S}$ for $t \in [0, 1]$ with $r_{0,S} = (\text{identity on } \bar{X})$ and $r_{1,S} = r_S$. The retraction r_S is uniquely characterized by

$$(8.1.1) \quad r_S(y \circ a) = y \quad \text{whenever } y \in \partial^P X_0 \text{ and } a \in \bar{A}_P, a \leq 1.$$

The converse of (8.1.1) also holds: if $y \in \partial^P X_0$, and P is the smallest with this property, then

$$(8.1.2) \quad r_S(y') = y \quad \text{implies} \quad \exists a \in \bar{A}_P, a \leq 1, \text{ such that } y' = y \circ a.$$

For any arithmetic subgroup Γ , the quotient X_0/Γ is compact. Also, for any P and given any open neighborhood U of $\bar{e}(P)/(\Gamma \cap P)$ in \bar{X}/Γ , we may choose X_0 so that

$$(8.1.3) \quad \partial^P X_0/(\Gamma \cap P) \subseteq U.$$

In fact, given any central tile, we may choose enlargements of it whose complements in \bar{X} , modulo Γ , form a fundamental system of neighborhoods of $\partial \bar{X}/\Gamma$ in \bar{X}/Γ .

(8.2). We will now fix a certain Γ -invariant open neighborhood B of $\partial\bar{X}$ in \bar{X} . There are only finitely many classes of \mathbb{Q} -flags mod Γ . Let \mathcal{F} represent such a class. Let $\bar{N}_{\mathcal{F}}$ be the tubular neighborhood of $e(P)$ in $X(P)$ as in (7.6), and let

$$B = \bigcup_{\mathcal{F}} \bigcup_{\gamma \in \Gamma} \gamma \cdot \bar{N}_{\mathcal{F}}.$$

Clearly B/Γ is an open neighborhood of $\partial\bar{X}/\Gamma$ in \bar{X}/Γ .

From now on, we fix a central tile X_0 with the following properties:

- (i) $\partial^P X_0 \subseteq \bigcup_{\mathcal{F}'} \bigcup_{\gamma \in \Gamma \cap P} (\gamma \cdot \bar{N}_{\mathcal{F}'})$ for any proper parabolic \mathbb{Q} -subgroup P with associated \mathcal{F} , where \mathcal{F}' runs through a set of $(\Gamma \cap P)$ -representatives of \mathbb{Q} -flags containing \mathcal{F} (the number of such representatives is finite);
- (ii) $\partial X_0 \subseteq B$, and $X_0 \cup B = \bar{X}$;
- (iii) the well-rounded retract W lies in the interior of X_0 .

By (8.1.3), it is possible to arrange for (i). Part (ii) follows from (i), since there are only finitely many \mathcal{F} mod Γ . To arrange for (iii), recall that W/Γ is compact, so that there is a Γ -invariant open neighborhood of $\partial\bar{X}$ in \bar{X} whose closure does not meet W ; enlarge X_0 if necessary so that its boundary lies in this neighborhood.

(8.3). We now construct the map R_t described in (0.3).

Definition. The map $R_t : \bar{X} \times [0, 1] \rightarrow \bar{X}$ is given as follows. If $t \in [0, \frac{1}{2}]$, let $R_t(x) = r_{2t,S}(x)$. If $t \in [\frac{1}{2}, 1]$, let $R_t(x) = r_{2t-1}(r_S(x))$.

The map R_t is well-defined after $t = \frac{1}{2}$ because the image of r_S lies in X (not $\partial\bar{X}$), and r is defined on X .

Proposition. The map R_t is a Γ -equivariant deformation retraction of \bar{X} onto W .

Proof. R_t is Γ -equivariant, since both $r_{t,S}$ and r_t have this property. Since $W \subseteq X_0$ by (iii) in (8.2), W is fixed pointwise by r_S . We know W is fixed pointwise by r . Hence the image of R_1 is at least W . On the other hand, the image is contained in W because the image of r is W . \square

(8.4). We now show that our global retraction \bar{r} of (7.6) is really a deformation retraction.

Theorem. $\bar{r} = R_1$.

Proof. If $x \in X_0$, then r_S does not move x , and $R_1(x) = r(x) = \bar{r}(x)$. Next, assume $x \in \bar{X}$ with $x \notin X_0$. The retraction r_S carries x to a point y , and there is a unique smallest P such that y lies on the face $\partial^P X_0$. However, by (8.1.2), x is of the form $x = y \circ a$ for some $a \in \bar{A}_P$, $a \leq 1$. By (8.2) part (i), $y \in \bar{N}_{\mathcal{F}''}$ for some $\mathcal{F}'' \supseteq \mathcal{F}$. Since $\bar{A}_P \subseteq \bar{A}_{P''}$, we have $x \in \bar{N}_{\mathcal{F}''}$, and x lies on the same $\bar{A}_{P''}$ -fiber of $\bar{N}_{\mathcal{F}''}$ as y . By (7.6) applied to \mathcal{F}'' , we have $\bar{r}(x) = \bar{r}(y)$. But $\bar{r}(y) = r(y) = R_1(x)$ by the definition of R_t . \square

(8.5). Here is a simple consequence of Theorem 8.4.

Corollary. *The map $\bar{r} : \bar{X} \rightarrow W$ is a Γ -equivariant homotopy equivalence. \bar{X}/Γ and W/Γ have the same homotopy type via the map $\bar{r} \bmod \Gamma$.*

We now generalize this corollary to all the boundary components.

Theorem. *Let P be a proper parabolic \mathbb{Q} -subgroup, with associated \mathcal{F} . Then \bar{r} induces a $(\Gamma \cap P)$ -equivariant homotopy equivalence of $\bar{e}(P)$ and $W_{\mathcal{F}}$.*

Proof. We must define a $(\Gamma \cap P)$ -equivariant homotopy inverse \bar{s} for $\bar{r}|_{\bar{e}(P)}$. Choose a point $x \in W_{\mathcal{F}}$, represented by a marked lattice f . Let $\boldsymbol{\tau} = (\tau, \dots, \tau) \in \mathfrak{A} = \mathfrak{A}_{\mathcal{F}, f}$. We define $\bar{s}(x)$ to be the point of \bar{X} corresponding to

$$(8.5.1) \quad \lim_{\tau \rightarrow 0+} \boldsymbol{\tau} \cdot f.$$

For all $\tau > 0$, $\boldsymbol{\tau} \cdot f$ lies in a single A_P -fiber of X . Let q_P be as in (3.5). It is clear that the limit in (8.5.1) is $q_P(x)$. Thus an equivalent definition is

$$(8.5.2) \quad \bar{s} \text{ is the restriction of } q_P \text{ to } W_{\mathcal{F}}.$$

By (8.5.2), \bar{s} is continuous. Since q_P commutes with the action of P , \bar{s} is $(\Gamma \cap P)$ -equivariant.

We now show that $\bar{r} \circ \bar{s}$ is the identity on $W_{\mathcal{F}}$. For a sufficiently small $\tau_0 > 0$, $(\tau_0, \dots, \tau_0) \cdot f$ will lie in $N_{\mathcal{F}}$. By (7.6), the value of \bar{r} on the point $(\tau, \dots, \tau) \cdot f$ is the same for all $\tau \in [0, \tau_0]$. Using (8.5.1), $\bar{r} \circ \bar{s}(x) = \bar{r}((\tau_0, \dots, \tau_0) \cdot f)$. However, the proof of Proposition 7.4 shows $\bar{r}((\tau_0, \dots, \tau_0) \cdot f) = f$.

Next, we show $\bar{s} \circ \bar{r}|_{\bar{e}(P)}$ is homotopic to the identity on $\bar{e}(P)$. Consider the family of maps

$$(8.5.3) \quad q_P \circ R_t \quad \text{restricted to } \bar{e}(P).$$

Here R_t is from (8.3), and q_P is extended continuously to a neighborhood of $\bar{e}(P)$ in \bar{X} in a natural way [B-S, (5.5)]. (A dangerous mistake would be to assume that the canonical cross-sections in $X(P)$ extend over $\bar{e}(P)$ as cross-sections of q_P .) The function in (8.5.3) is continuous because the image of $\bar{e}(P)$ under R_t lies in $\bar{e}(P) \cup X$, where q_P is continuous. At $t = 0$, (8.5.3) is the identity $\bar{e}(P) \rightarrow \bar{e}(P)$. At $t = 1$, for any $x \in \bar{e}(P)$,

$$\begin{aligned} q_P \circ R_1(x) &= q_P(\bar{r}(x)) && \text{by (8.4)} \\ &= \bar{s}(\bar{r}(x)) && \text{by (8.5.2).} \end{aligned}$$

Thus $q_P \circ R_t$ affords a $(\Gamma \cap P)$ -equivariant homotopy between $\bar{s} \circ \bar{r}|_{\bar{e}(P)}$ and the identity. \square

Remark. Formula (8.5.3) defines a well-rounded retraction within $\bar{e}(P)$ itself, carrying $\bar{e}(P)$ onto a subset of $e(P)$ homeomorphic to $W_{\mathcal{F}}$. We know that $\bar{e}(P)/(\Gamma \cap P)$ is a compactification of a bundle whose base is a locally symmetric space (perhaps with finite quotient singularities) for a group of lower rank, and whose fiber is an arithmetic quotient of U_P . Accordingly, there is a cellular quotient map from $W_{\mathcal{F}}/(\Gamma \cap P)$ to the well-rounded retract in the lower-rank locally symmetric space. This map may be constructed by dualizing the techniques of [M2]; see (10.7).

SECTION 9—THE MAIN RESULTS

In (9.5) we establish our main result, as we construct the spectral sequence described in the introduction. The bulk of the proofs resides in (9.2)–(9.4). We set up two spectral sequences, one (9.2.1) computing the cohomology of the Borel-Serre boundary $\partial\bar{X}/\Gamma$, and the other (9.3.1) based on the subcomplexes $W_{\mathcal{F}}/(\Gamma \cap P)$ in the well-rounded retract W/Γ . Because the map \bar{r} gives homotopy equivalences on \bar{X}/Γ and all its faces, it induces an isomorphism between these spectral sequences, and between their abutments. We set up canonical maps $H^*(X/\Gamma) \rightarrow (9.2.1)$ and $H^*(W/\Gamma) \rightarrow (9.3.1)$. Finally, we show that the map $H^*(W/\Gamma) \rightarrow (9.3.1)$ computes the canonical restriction map $H^*(\bar{X}/\Gamma) \rightarrow H^*(\partial\bar{X}/\Gamma)$.

(9.1). Let $A^{p,q}$ be any first-quadrant double complex. There are two standard filtrations of the double complex, $\{A^{p,q} \mid q \geq q_0\}$ and $\{A^{p,q} \mid p \geq p_0\}$. The spectral sequences these give are called respectively the *type I* and *type II sequences* for $A^{p,q}$. Our main results involve type II sequences. The cohomology of the single complex associated to $A^{p,q}$ is called the *abutment* of (either) spectral sequence.

Throughout the paper, (co)homology groups have coefficients in any fixed abelian group.

(9.2). We set up a Mayer-Vietoris spectral sequence for the cohomology of $\partial\bar{X}/\Gamma$. Let Φ_l be a set of representatives of the Γ -equivalence classes of the \mathbb{Q} -flags $\mathcal{F} = \{0 \subsetneq V_1 \subsetneq \cdots \subsetneq V_l = k^n\}$ (see (3.2)). There are only finitely many such equivalence classes. Let $\mathcal{E} = \{\bar{e}(P)/(\Gamma \cap P) \mid \mathcal{F} \in \Phi_2\}$ (corresponding to maximal proper parabolic \mathbb{Q} -subgroups). This \mathcal{E} is a cover of $\partial\bar{X}/\Gamma$. By (3.5), the non-empty $(p+1)$ -fold intersections of distinct members of \mathcal{E} are exactly the sets in $\{\bar{e}(P)/(\Gamma \cap P) \mid \mathcal{F} \in \Phi_{p+2}\}$. This means we are in the correct setting to use the Čech cohomology techniques of [B-T, §15].

For $0 \leq p \leq n-2$, define

$$\mathcal{X}^{p,q} = \bigoplus_{\mathcal{F} \in \Phi_{p+2}} C^q(\bar{e}(P)/(\Gamma \cap P)),$$

where C^q denotes the singular q -cochains. We make $\mathcal{X}^{p,q}$ into a double complex, where the vertical maps are $(-1)^p$ times the coboundary maps, and the horizontal maps are induced as in Čech cohomology from the inclusions $\bar{e}(P) \hookleftarrow \bar{e}(P')$ for $\mathcal{F}' \supseteq \mathcal{F}$, with the alternating sign convention of [B-T, (15.7.1)]. Exactly as in [B-T, (15.7)], we see that the q -th row of $\mathcal{X}^{p,q}$ is exact, except at the $p=0$ position where the kernel equals $C^q(\partial\bar{X}/\Gamma)$. Thus the type I spectral sequence for $\mathcal{X}^{p,q}$ collapses at E_2 to $H^*(\partial\bar{X}/\Gamma)$. Therefore, the total complex of $\mathcal{X}^{p,q}$ computes $H^*(\partial\bar{X}/\Gamma)$.

Let $E_{r,\mathcal{X}}^{p,q}$ be the type II spectral sequence for $\mathcal{X}^{p,q}$. We have

$$(9.2.1) \quad E_{1,\mathcal{X}}^{p,q} = \bigoplus_{\mathcal{F} \in \Phi_{p+2}} H^q(\bar{e}(P)/(\Gamma \cap P)) \Rightarrow H^{p+q}(\partial\bar{X}/\Gamma).$$

(9.3). For $0 \leq p \leq n-2$, define

$$\mathcal{W}^{p,q} = \bigoplus_{\mathcal{F} \in \Phi_{p+2}} C^q(W_{\mathcal{F}}/(\Gamma \cap P)),$$

made into a double complex in the same way as $\mathcal{X}^{p,q}$. (Recall that $W_{\mathcal{F}} \hookrightarrow W_{\mathcal{F}'}$ whenever $\mathcal{F}' \supseteq \mathcal{F}$.) Let $E_{r,\mathcal{W}}^{p,q}$ be the type II spectral sequence for $\mathcal{W}^{p,q}$. As above,

$$(9.3.1) \quad E_{1,\mathcal{W}}^{p,q} = \bigoplus_{\mathcal{F} \in \Phi_{p+2}} H^q(W_{\mathcal{F}}/(\Gamma \cap P)).$$

Let \mathcal{W}^* denote the single complex associated to $\mathcal{W}^{p,q}$. Thus the abutment of (9.3.1) is $H^{p+q}(\mathcal{W}^*)$.

(9.4). We now define a map of double complexes. Consider the single-column double complex

$$\hat{\mathcal{X}}^{p,q} = \begin{cases} C^q(\bar{X}/\Gamma) & \text{if } p = 0 \\ 0 & \text{if } p > 0. \end{cases}$$

There is an obvious map $\hat{\mathcal{X}}^{p,q} \rightarrow \mathcal{X}^{p,q}$, where for $p = 0$ we map $\omega \in C^q(\bar{X}/\Gamma)$ to the direct sum over $\mathcal{F} \in \Phi_2$ of the restriction of ω to $C^q(\bar{e}(P)/(\Gamma \cap P))$.

Lemma. *The map $\hat{\mathcal{X}}^{p,q} \rightarrow \mathcal{X}^{p,q}$ is a chain map of double complexes.*

Proof. Let $\omega \in C^q(\bar{X}/\Gamma)$, and let τ be its image in $\mathcal{X}^{0,q}$. The only thing to check is that the horizontal arrow $\delta : \mathcal{X}^{0,q} \rightarrow \mathcal{X}^{1,q}$ carries τ to 0. But as we have said, the kernel of δ is precisely the set of elements that come from a global cochain on $\partial\bar{X}/\Gamma$. And τ does come from a global cochain, namely the restriction of ω to $\partial\bar{X}/\Gamma$. \square

Considering the type I spectral sequences, we see that $\hat{\mathcal{X}}^{p,q} \rightarrow \mathcal{X}^{p,q}$ induces on the abutments the canonical restriction map $H^q(\bar{X}/\Gamma) \rightarrow H^q(\partial\bar{X}/\Gamma)$.

Next, consider

$$\hat{\mathcal{W}}^{p,q} = \begin{cases} C^q(W/\Gamma) & \text{if } p = 0 \\ 0 & \text{if } p > 0, \end{cases}$$

made into a double complex the same way $\hat{\mathcal{X}}$ was. There is an obvious map $\psi : \hat{\mathcal{W}}^{p,q} \rightarrow \mathcal{W}^{p,q}$ carrying $\omega \in C^q(W/\Gamma)$ to the direct sum over $\mathcal{F} \in \Phi_2$ of the restrictions of ω to $C^q(W_{\mathcal{F}}/(\Gamma \cap P))$. Again, $\psi : \hat{\mathcal{W}}^{p,q} \rightarrow \mathcal{W}^{p,q}$ is a chain map of double complexes.

(9.5). We now give the main result of the paper.

Theorem. (i) *We have a commutative diagram*

$$\begin{array}{ccc} H^*(\bar{X}/\Gamma) & \xrightarrow{\text{restriction}} & H^*(\partial\bar{X}/\Gamma) \\ \cong \downarrow & & \downarrow \cong \\ H^*(W/\Gamma) & \xrightarrow{\psi^*} & H^*(\mathcal{W}^*) \end{array}$$

where the vertical maps are natural isomorphisms and the top map is the canonical restriction map.

(ii) *There is a spectral sequence*

$$(9.5.1) \quad E_{1,\mathcal{W}}^{p,q} = \bigoplus_{\mathcal{F} \in \Phi_{p+2}} H^q(W_{\mathcal{F}}/(\Gamma \cap P)) \Rightarrow H^{p+q}(\mathcal{W}^*),$$

and ψ induces a map of spectral sequences given on the E_1 page by the natural restriction map $H^q(W/\Gamma) \rightarrow E_{1,\mathcal{W}}^{0,q}$.

Here X is as in (1.5), Γ is any arithmetic subgroup of Γ_0 as in (2.2), Φ_l is as in (9.2), and P is the \mathbb{R} -points of the parabolic \mathbb{Q} -subgroup associated to \mathcal{F} as in (3.2). The space $W_{\mathcal{F}}$, a closed subcomplex of the well-rounded retract W , is defined in (6.1). We use any fixed abelian group of coefficients.

Proof. First we observe that the global deformation retraction $\bar{r} : \bar{X} \rightarrow W$ induces a map of double complexes $\mathcal{X}^{p,q} \rightarrow \mathcal{W}^{p,q}$. The map \bar{r} commutes with the vertical maps of these double complexes, because any continuous map commutes with the coboundary maps in singular cohomology. More importantly, \bar{r} commutes with the horizontal maps because it is defined *globally* on \bar{X} , and because the horizontal maps are induced from inclusion maps on various subspaces of \bar{X} .

It follows that \bar{r} induces a homomorphism of spectral sequences $E_{r,\mathcal{X}}^{p,q} \rightarrow E_{r,\mathcal{W}}^{p,q}$. The map on E_1 terms comes from the maps $H^q(\bar{e}(P)/(\Gamma \cap P)) \rightarrow H^q(W_{\mathcal{F}}/(\Gamma \cap P))$ induced for all \mathcal{F} by \bar{r} . By Theorem 8.5, the latter maps are canonical isomorphisms. By [Br, VII.2.6], the right-hand arrow in (i) is an isomorphism.

Similarly, \bar{r} induces a map $\hat{\mathcal{X}}^{p,q} \rightarrow \hat{\mathcal{W}}^{p,q}$. By Corollary 8.4, it gives a canonical isomorphism in cohomology. This is the left vertical arrow in (i).

Part (9.4) defined the maps of double complexes that induce the horizontal arrows in (i), and it explained why the top arrow is the canonical restriction. This proves (i).

The spectral sequence in (ii) is (9.3.1). The assertion about ψ in (ii) is immediate from the definition of ψ . \square

SECTION 10—PRACTICAL COMMENTS AND RELATED RESULTS

(10.1). We make some remarks about how to use the spectral sequence (9.5.1) in practice. When we work with the finite cell complex W/Γ , we represent each cell by some piece of data. (See (10.6), [M-M0], or [A-G-G].) Since $W_{\mathcal{F}}$ is a subcomplex of W , it is tempting to imagine representing each cell of $W_{\mathcal{F}}/(\Gamma \cap P)$ by some piece of the W/Γ data. However, two cells of $W_{\mathcal{F}}$ may be equivalent mod Γ and yet inequivalent mod $(\Gamma \cap P)$. An example when $\Gamma = \mathrm{GL}_3(\mathbb{Z})$, $\mathcal{F} = \{0 \subseteq (*, *, 0) \subseteq \mathbb{Q}^3\}$, is provided by the two 0-cells of W whose minimal vectors are given by the columns of the following matrices (together with their negatives):

$$\begin{pmatrix} 1 & 0 & 1 & 0 & 0 & -1 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}.$$

The cells are Γ -equivalent. (In the language of [M-M0], their \mathcal{C} -configurations are both complete quadrilaterals.) But any element of Γ preserving \mathcal{F} must carry the left-hand matrix to a matrix with three 0's in its bottom row.

In general, each Γ -class of cells of $W_{\mathcal{F}}$ will break up into finitely many $(\Gamma \cap P)$ -classes. So a cell in $W_{\mathcal{F}}/(\Gamma \cap P)$ is represented by a piece of data for W/Γ , plus a finite amount of extra structure depending on \mathcal{F} . Say one has worked out all the classes in (9.5.1) as explicit cocycles on the cell complexes $W_{\mathcal{F}}/(\Gamma \cap P)$. To compute the d_1 and higher differentials, one has to work out how the data-with-extra-structure in $W_{\mathcal{F}}/(\Gamma \cap P)$ is connected with the data-with-extra-structure in $W_{\mathcal{F}'}/(\Gamma \cap P)$ for $\mathcal{F}' \supseteq \mathcal{F}$. SHEAFHOM, a suite of object-oriented programs currently under development by the second author, may help to automate such computations.

Say that Γ is *small enough* if, whenever $\mathcal{F}, \mathcal{F}'$ are distinct \mathbb{Q} -flags and $W_{\mathcal{F}} \cap W_{\mathcal{F}'} \neq \emptyset$, there is no element of Γ carrying \mathcal{F} to \mathcal{F}' . When Γ is small enough in this sense, we need never worry about the extra structure just described. To see why, assume there were two cells $C_a, C_b \in W_{\mathcal{F}}$ which were not $(\Gamma \cap P)$ -equivalent but such that $C_b = \gamma \cdot C_a$ for some $\gamma \in \Gamma$. Set $\mathcal{F}' = \gamma^{-1}\mathcal{F}$; we have $\mathcal{F}' \neq \mathcal{F}$ since $\gamma \notin P$. Because C_b respects \mathcal{F} , C_a must respect \mathcal{F}' . Hence $C_a \in W_{\mathcal{F}} \cap W_{\mathcal{F}'}$. Since Γ was assumed small enough, the existence of such a γ is a contradiction.

When $k = \mathbb{Q}$ and $n \leq 4$, one can use the methods of [M1] to show that the principal congruence subgroups $\Gamma(N) \subset \mathrm{GL}_n(\mathbb{Z})$ of prime level $N \geq 3$ are small enough in the above sense. We do not know whether all neat Γ are small enough.

(10.2). For convenience, we record the homology version of our main theorem. We use the obvious analogues of the notation of (9.5).

Theorem. (i) *We have a commutative diagram*

$$\begin{array}{ccc} H_*(\partial\bar{X}/\Gamma) & \xrightarrow{\text{inclusion}} & H_*(\bar{X}/\Gamma) \\ \cong \downarrow & & \downarrow \cong \\ H_*(\mathcal{W}_*) & \xrightarrow{\psi_*} & H_*(W/\Gamma) \end{array}$$

where the vertical maps are natural isomorphisms and the top map is the canonical inclusion map.

(ii) *There is a spectral sequence*

$$E_{p,q}^1 = \bigoplus_{\mathcal{F} \in \Phi_{p+2}} H_q(W_{\mathcal{F}}/(\Gamma \cap P)) \Rightarrow H_{p+q}(\mathcal{W}_*),$$

and ψ induces a map of spectral sequences given on the E_1 page by the natural inclusion map $E_{0,q}^1 \rightarrow H_q(W/\Gamma)$.

This is proved by dualizing the constructions of Section 9.

(10.3). Consider one Borel-Serre boundary face $\bar{e}(P)/(\Gamma \cap P)$. As we mentioned in the introduction, our methods allow us to compute the canonical maps $H^*(\bar{X}/\Gamma) \rightarrow H^*(\bar{e}(P)/(\Gamma \cap P))$ and $H_*(\bar{e}(P)/(\Gamma \cap P)) \rightarrow H_*(\bar{X}/\Gamma)$ without any spectral sequences. We simply use the cellular map $W_{\mathcal{F}}/(\Gamma \cap P) \rightarrow W/\Gamma$.

In the application to $\mathrm{GL}_4(\mathbb{Z})$ described near the end of (0.2), we are primarily interested in the degree-five homology. We believe that in degree five, all the homology of the boundary comes from the maximal boundary faces. We could therefore find the image of

$$\left(\bigoplus_{\substack{\mathcal{F} \text{ for maximal} \\ \text{faces mod } \Gamma}} H_5(W_{\mathcal{F}}/(\Gamma \cap P)) \right) \rightarrow H_5(W/\Gamma)$$

without computing any spectral sequences.

It would very interesting to find the image of the map $H_*(\partial \bar{X}/\Gamma) \rightarrow H_*(\bar{X}/\Gamma)$ without having to find $H_*(\partial \bar{X}/\Gamma)$ directly, and without having to work out any spectral sequences.

(10.4). All our results hold for Γ -equivariant cohomology. That is, one may replace $H^*((...)/\Gamma)$ with $H_{\Gamma}^*((...))$ in the theorems of Section 9 and throughout the paper. Of course, if we use coefficients in a field of characteristic 0 or of characteristic prime to the order of any torsion element of Γ , then the two kinds of cohomology are canonically isomorphic. All these results hold for homology.

In the Γ -equivariant setting, the summands in each column of our spectral sequences would be replaced by $H_{(\Gamma \cap P)}^*(W_{\mathcal{F}})$, and each of these would have to be computed by a spectral sequence in its own right. We would get a spectral sequence of spectral sequences.

Both $H_{\Gamma}^*(X)$ and $H^*(X/\Gamma)$ are interesting objects from the point of view of number theory and automorphic forms. Both have Hecke operators acting on them. In [A4], the first author has conjectured that any Hecke eigenclass in $H_{\Gamma}^*(X; \mathbb{F}_p)$, for any p , should have an attached mod p representation of the absolute Galois group of \mathbb{Q} . In [A-M2], we have conjectured the same for the Hecke module $H^*(X/\Gamma; \mathbb{F}_p)$. Both papers provide examples.

(10.5). The method of this paper can easily be extended to SL_n . If G were $\mathrm{SL}_n(S)$, then $K \backslash G$ would be the product of the irreducible symmetric spaces for the $\mathrm{SL}_n(k_v)$. In the case $G = \mathrm{GL}_n(S)$ of this paper, $HK \backslash G$ is the product of the $\mathrm{SL}_n(k_v)$ symmetric spaces and a Euclidean factor $H \backslash (\prod_v \mathbb{R}_+) \cong \mathbb{R}_+^{r_1+r_2-1}$, where r_1, r_2 are the number of real and complex places of k , respectively.

(10.6). Let $k = \mathbb{Q}$, so that $\Gamma \subseteq \mathrm{GL}_n(\mathbb{Z})$ or $\mathrm{SL}_n(\mathbb{Z})$, and let $n \leq 4$. In [M1] and the survey article [M-M0], an interpretation of the cell structure on the well-rounded retract W is given. There is a one-to-one correspondence in which each cell corresponds to a configuration of points and lines (called a \mathcal{C} -configuration) in the projective space $\mathbb{P}^{n-1}(\mathbb{Q})$. It is easy to interpret our $W_{\mathcal{F}}$ in this language.

(10.7). Again let $k = \mathbb{Q}$ and $n \leq 4$. Let \check{X}/Γ be any Satake compactification of X/Γ . In [M2], one constructs a cell structure on \check{X}/Γ . Intersecting with X/Γ gives a “locally-compact cell structure” on X/Γ that is dual to the well-rounded retract W/Γ inside X/Γ .

(see [M1, §2] for definitions and precise statements). There is a cellular map from the barycentric subdivision of the boundary of \check{X}/Γ to the barycentric subdivision of W/Γ . This offers an interesting contrast to the present paper.

REFERENCES

- [A1] A. Ash, *Deformation retracts with lowest possible dimension of arithmetic quotients of self-adjoint homogeneous cones*, Math. Annalen **225** (1977), 69–76.
- [A2] ———, *Cohomology of congruence subgroups of $SL(n, \mathbb{Z})$* , Math. Annalen **249** (1980), 55–73.
- [A3] ———, *Small-dimensional classifying spaces for arithmetic subgroups of general linear groups*, Duke Jour. **51** (1984), 459–468.
- [A4] ———, *Galois representations attached to mod p cohomology of $GL(n, \mathbb{Z})$* , Duke Jour. **65** (1992), 235–255.
- [A-G-G] A. Ash, D. Grayson, P. Green, *Computations of cuspidal cohomology of congruence subgroups of $SL(3, \mathbb{Z})$* , Jour. Number Theory **19** (1984), 412–436.
- [A-M1] A. Ash, M. McConnell, *Doubly cuspidal cohomology for principal congruence subgroups of $GL(3, \mathbb{Z})$* , Math. of Computation **59** (1992), 673–688.
- [A-M2] ———, *Experimental indications of three-dimensional Galois representations from the cohomology of $SL(3, \mathbb{Z})$* , Jour. Exper. Math. **1** (1992), 209–223.
- [B-S] A. Borel, J.-P. Serre, *Corners and arithmetic groups*, Comm. Math. Helv. **48** (1973), 436–491.
- [B-T] R. Bott, L. W. Tu, *Differential forms in algebraic topology*, Springer, New York, 1982.
- [Br] K. Brown, *Cohomology of Groups*, Springer, New York, 1982.
- [B] A. Brownstein, thesis (unpublished).
- [G] D. Grayson, *Reduction theory using semistability*, Comm. Math. Helv. **59** (1984), 600–634.
- [H] G. Harder, *Some results on the Eisenstein cohomology of arithmetic subgroups of GL_n* , Cohomology of Arithmetic Groups and Automorphic Forms (J.-P. Labesse and J. Schwermer, eds.), Springer Lecture Notes 1447, Springer, New York, 1990, pp. 85–153.
- [L-S] R. Lee, R. H. Szczarba, *On the torsion in $K_4(\mathbb{Z})$ and $K_5(\mathbb{Z})$* , Duke Jour. **45** (1978), 101–129.
- [M-M0] R. MacPherson, M. McConnell, *Classical Projective Geometry and Modular Varieties*, Algebraic Analysis, Geometry, and Number Theory: Proceedings of the JAMI Inaugural Conference (J.-I. Igusa, ed.), Johns Hopkins University Press, 1989, pp. 237–290.
- [M-M1] ———, *Explicit reduction theory for Siegel modular threefolds*, Invent. Math. **111** (1993), 575–625.
- [M1] M. McConnell, *Classical projective geometry and arithmetic groups*, Math. Annalen **290** (1991), 441–462.
- [M2] ———, *Cell decompositions of Satake compactifications for $SL(n, \mathbb{R})$* , preprint.
- [Men] E. Mendoza, *Cohomology of PGL_2 over imaginary quadratic integers*, Bonner Math. Schriften, no. 128 (1979), Universität Bonn, Mathematisches Institut, Bonn.
- [S] L. Saper, *Tilings and finite energy retractions of locally symmetric spaces*, preprint.
- [S-V] J. Schwermer, K. Vogtmann, *The integral homology of SL_2 and PSL_2 of euclidean imaginary quadratic integers*, Comment. Math. Helv. **58** (1983), 573–598.
- [Sou1] C. Soulé, *The cohomology of $SL(3, \mathbb{Z})$* , Topology **17** (1978), 1–22.
- [Sou2] ———, *Thèse*, Univ. Paris VII.
- [Što] M. I. Štogrin, *Locally quasi-densest lattice packings of spheres*, Soviet Math. Doklady **15** (1974), 1288–1292.
- [vG-T] B. van Geemen, J. Top, *A non-selfdual automorphic representation of GL_3 and a Galois representation*, Invent. Math. **117** (1994), 391–401.
- [V] K. Vogtmann, *Rational homology of Bianchi groups*, Math. Annalen **272** (1985), 399–419.

E-mail address: `ash@math.ohio-state.edu`

OKLAHOMA STATE UNIVERSITY, MATH, 401 MATH SCIENCES, STILLWATER, OK 74078-1058

E-mail address: `mmcconn@math.okstate.edu`