

## ON UNITARY REPRESENTATIONS OF GROUPS OF ISOMETRIES

V. V. USPENSKIJ

ABSTRACT. Is the topological group of all motions (including translations) of an infinite-dimensional Hilbert space  $H$  isomorphic to a subgroup of the unitary group  $U(H)$ ? This question was asked by Su Gao. We answer the question in the affirmative.

## 1. INTRODUCTION

For a Hilbert space  $H$  we denote by  $U_s(H)$  the group of all unitary operators on  $H$ , equipped with the strong operator topology. This topology is the same as the topology of pointwise convergence, it is induced on  $U(H)$  by the product topology on  $H^H$ . The group  $U_s(H)$  is a topological group, in other words, its topology is compatible with its group structure.

Given a topological group  $G$ , a *unitary representation* of  $G$  is a continuous homomorphism  $f : G \rightarrow U_s(H)$ . The representation  $f$  is *faithful* if it is injective, and *topologically faithful* if  $f$  is a homeomorphic embedding of  $G$  into  $U_s(H)$ . Thus a group admits a topologically faithful unitary representation if and only if it is isomorphic to a topological subgroup of a group of the form  $U_s(H)$ .

Every locally compact group  $G$  (in particular, every Lie group) admits a topologically faithful unitary representation. For example, the regular representation on  $L^2(G)$  is such. Beyond the class of locally compact groups the situation is less clear. There are topological groups with a countable base which do not admit any faithful unitary representation or even any faithful representation by isometries on a reflexive Banach space (Megrelishvili). For example, the group  $H(I)$  of all self-homeomorphisms of the interval  $I = [0, 1]$  has this property [2]. It follows that  $\text{Is}(U)$ , the group of isometries of the Urysohn universal metric space  $U$ , also has this property, since it contains a topological copy of  $H(I)$ . See [4], [5], [6] for the proof and a discussion of the space  $U$  and its group of isometries.

The question arises: what are metric spaces  $M$  for which the group  $\text{Is}(M)$  of all isometries of  $M$  admits a (topologically) faithful unitary representation? We consider the topology of pointwise convergence on  $\text{Is}(M)$  (or the compact-open topology, which is the same on  $\text{Is}(M)$ ). If  $M$  is locally compact and connected, the group  $\text{Is}(M)$  is locally compact [1, Ch.1, Thm.4.7] and hence embeds in a unitary group. On the other hand, if  $M$  is the Urysohn space  $U$  (which topologically is the same as a Hilbert space [7]), then  $\text{Is}(M)$  has no faithful unitary representations, as we noted above.

Let  $H$  be a (real or complex) Hilbert space. Consider the topological group  $\text{Is}(H)$  of all (not necessarily linear) isometries of  $H$ . This group is a topological semidirect product of the subgroup of translations (which is isomorphic to the additive group of

---

*Date:* 10 June 2004.

*Key words and phrases.* Hilbert space, motion, representation, unitary groups, isometry.

*2000 Mathematics Subject Classification:* Primary: 43A65. Secondary: 43A35, 22D10, 54E35.

$H$ ) and the unitary group  $U(H_{\mathbf{R}})$  of the real Hilbert space  $H_{\mathbf{R}}$  underlying  $H$ . Su Gao asked the author whether the topological group  $\text{Is}(H)$  is isomorphic to a subgroup of a unitary group. The aim of the present note is to answer this question in the affirmative. It follows that the additive group of  $H$  admits a topologically faithful unitary representation. Even this corollary does not look self-evident.

**Theorem 1.1.** *Let  $H$  be a Hilbert space, and let  $G = \text{Is}(H)$  be the topological group of all (not necessarily linear) isometries of  $H$ . Then  $G$  is isomorphic to a subgroup of a unitary group.*

In Section 2 we remind some basic facts concerning positive-definite functions on groups, and prove (a generalization of) Theorem 1.1 in Section 3.

## 2. POSITIVE-DEFINITE FUNCTIONS

For a complex matrix  $A = (a_{ij})$  we denote by  $A^*$  the matrix  $(\bar{a}_{ji})$ .  $A$  is *Hermitian* if  $A = A^*$ . A Hermitian matrix  $A$  is *positive* if all the eigenvalues of  $A$  are  $\geq 0$  or, equivalently, if  $A = B^2$  for some Hermitian  $B$ . A complex function  $p$  on a group  $G$  is *positive-definite* if for every  $g_1, \dots, g_n \in G$  the  $n \times n$ -matrix  $p(g_i^{-1}g_j)$  is Hermitian and positive. If  $f : G \rightarrow U(H)$  is a unitary representation of  $G$ , then for every vector  $v \in H$  the function  $p = p_v$  on  $G$  defined by  $p(g) = (gv, v)$  is positive-definite. (We denote by  $(x, y)$  the scalar product of  $x, y \in H$ .) Conversely, let  $p$  be a positive-definite function on  $G$ . Then  $p = p_v$  for some unitary representation  $f : G \rightarrow U(H)$  and some  $v \in H$ . Indeed, consider the group algebra  $\mathbf{C}[G]$ , equip it with the scalar product defined by  $(g, h) = p(h^{-1}g)$ , quotient out the kernel, and take the completion to get  $H$ . The regular representation of  $G$  on  $\mathbf{C}[G]$  gives rise to a unitary representation on  $H$  with the required property. If  $G$  is a topological group and the function  $p$  is continuous, the resulting unitary representation is continuous as well. These arguments yield the following criterion:

**Proposition 2.1.** *A topological group  $G$  admits a topologically faithful unitary representation (in other words, is isomorphic to a subgroup of  $U_s(H)$  for some Hilbert space  $H$ ) if and only if for every neighborhood  $U$  of the neutral element  $e$  of  $G$  there exist a continuous positive-definite function  $p : G \rightarrow \mathbf{C}$  and  $a > 0$  such that  $p(e) = 1$  and  $|1 - p(g)| > a$  for every  $g \in G \setminus U$ .*

*Proof.* The necessity is easy: consider convex linear combinations of the functions  $p_v$  defined above, where  $v \in H$  is a unit vector. We prove that the condition is sufficient. According to the remarks preceding the proposition, for every neighborhood  $U$  of  $e$  there exists a unitary representation  $f_U : G \rightarrow U_s(H)$  such that  $f_U^{-1}(V) \subset U$  for some neighborhood  $V$  of 1 in  $U_s(H)$ . Indeed, pick a positive-definite function  $p$  on  $G$  such that  $p(e) = 1$  and

$$\inf\{|1 - p(g)| : g \in G \setminus U\} = a > 0.$$

Let  $f_U$  be a representation such that  $p(g) = (f_U(g)v, v)$  for every  $g \in G$  and some unit vector  $v \in H$ . Then the neighborhood

$$V = \{A \in U_s(H) : |1 - (Av, v)| < a\}$$

of 1 in  $U_s(H)$  is as required.

It follows that the Hilbert direct sum of the representations  $f_U$ , taken over all neighborhoods  $U$  of  $e$ , is topologically faithful.  $\square$

Let  $G$  be a locally compact Abelian group,  $\hat{G}$  its Pontryagin dual. Recall that continuous positive-definite functions on  $\hat{G}$  are precisely the Fourier transforms of positive measures on  $G$  (Bochner's theorem). In particular, every positive-definite function  $p : \mathbf{R}^k \rightarrow \mathbf{C}$  has the form

$$p(x) = \int_{\mathbf{R}^k} \exp(-2\pi i(x, y)) d\mu(y)$$

for some positive measure  $\mu$  on  $\mathbf{R}^k$ . Here  $(x, y) = \sum_{i=1}^k x_i y_i$  for  $x = (x_1, \dots, x_k)$  and  $y = (y_1, \dots, y_k)$ . Let  $p(x) = \exp(-\pi(x, x))$ . Then  $p$  is the Fourier transform of the Gaussian measure  $p(y)dy$  and hence is positive-definite. Thus for any  $x_1, \dots, x_n \in \mathbf{R}^k$  the matrix  $(\exp(-\|x_i - x_j\|^2))$  is positive. Here  $\|\cdot\|$  is the Hilbert space norm on  $\mathbf{R}^k$  defined by  $\|y\| = \sqrt{(y, y)}$ . If  $H$  is an infinite-dimensional Hilbert space and  $x_1, \dots, x_n \in H$ , then the finite-dimensional linear subspace of  $H$  spanned by  $x_1, \dots, x_n$  is isometric to  $\mathbf{R}^k$  for some  $k$ . Thus we arrive at the following:

**Lemma 2.2.** *Let  $H$  be a Hilbert space, and let  $x_1, \dots, x_n$  be points in  $H$ . Then the symmetric  $n \times n$ -matrix  $(\exp(-\|x_i - x_j\|^2))$  is positive.*

### 3. PROOF OF THE MAIN THEOREM

We prove a theorem which is more general than Theorem 1.1.

**Theorem 3.1.** *Let  $(M, d)$  be a metric space, and let  $G = \text{Is}(M)$  be its group of isometries. Suppose that there exists a real-valued positive-definite function  $p : \mathbf{R} \rightarrow \mathbf{R}$  such that:*

- (1)  $p(0) = 1$ , and for every  $\epsilon > 0$  we have  $\sup\{p(x) : |x| \geq \epsilon\} < 1$ ;
- (2) for every points  $a_1, \dots, a_n \in M$  the symmetric real  $n \times n$ -matrix  $(p(d(a_i, a_j)))$  is positive.

*Then the topological group  $G$  is isomorphic to a subgroup of a unitary group.*

In virtue of Lemma 2.2, this theorem is indeed stronger than Theorem 1.1: if  $M = H$ , we can take for  $p$  the function defined by  $p(x) = \exp(-x^2)$ .

*Proof.* According to Proposition 2.1, we must show that there are sufficiently many positive-definite functions on  $G$ . For every  $a \in M$  consider the function  $p_a : G \rightarrow \mathbf{R}$  defined by  $p_a(g) = p(d(ga, a))$ . This function is positive-definite. Indeed, let  $g_1, \dots, g_n \in G$ . Since each  $g_i$  is an isometry of  $M$ , we have  $d(g_i^{-1}g_j a, a) = d(g_i a, g_j a)$ , and hence the matrix  $(p_a(g_i^{-1}g_j)) = (p(d(g_i^{-1}g_j a, a))) = (p(d(g_i a, g_j a)))$  is positive by our assumption.

Let  $U$  be a neighborhood of 1 (= the identity map of  $M$ ) in  $G$ . Without loss of generality we may assume that

$$U = \{g \in G : d(ga_i, a_i) < \epsilon, i = 1, \dots, n\}$$

for some  $a_1, \dots, a_n \in M$  and some  $\epsilon > 0$ . Let  $t$  be the average of the functions  $p_{a_1}, \dots, p_{a_n}$ . Then  $t$  is a positive-definite function on  $G$  such that

$$\sup\{t(g) : g \in G \setminus U\} < 1.$$

Indeed, let  $\delta > 0$  be such that  $p(x) \leq 1 - \delta$  for every  $x$  such that  $|x| \geq \epsilon$ . If  $g \in G \setminus U$ , then  $d(ga_i, a_i) \geq \epsilon$  for some  $i$ , hence  $p_{a_i}(g) = p(d(ga_i, a_i)) \leq 1 - \delta$  and

$$t(g) = \frac{1}{n} \sum_{k=1}^n p_{a_k}(g) \leq 1 - \frac{\delta}{n}.$$

We invoke Proposition 2.1 to conclude that  $G$  has a topologically faithful representation.  $\square$

## REFERENCES

- [1] Sh. Kobayashi, K. Nomizu, *Foundations of differential geometry. Vol. 1*, Interscience Publishers, New York–London, 1963.
- [2] M.G. Megrelishvili, *Every semitopological semigroup compactification of the group  $H_+[0, 1]$  is trivial*, Semigroup Forum **63** (2001), no. 3, 357–370.
- [3] V. Pestov, *Topological groups: Where to from here?*, Topology Proc., **24** (1999), 421–502. E-print: math.GN/9910144
- [4] V. Uspenskij, *On the group of isometries of the Urysohn universal metric space*, Comment. Math. Univ. Carolinae **31** (1990), no. 1, 181–182.
- [5] V. Uspenskij, *On subgroups of minimal topological groups*, E-print: math.GN/0004119
- [6] V.V. Uspenskij, *Compactifications of topological groups*, Proceedings of the Ninth Prague Topological Symposium (2001), 331–346 (electronic), Topol. Atlas, North Bay, ON, 2002. E-print: math.GN/0204144
- [7] V.V. Uspenskij, *The Urysohn universal metric space is homeomorphic to a Hilbert space*, Topology Appl. **139** (2004), 145–149. E-print: math.GN/0309101

DEPARTMENT OF MATHEMATICS, 321 MORTON HALL, OHIO UNIVERSITY, ATHENS, OHIO 45701, USA

*E-mail address:* uspensk@math.ohiou.edu