

A REMARK ON CUSPIDAL LOCAL SYSTEMS

VICTOR OSTRIK

ABSTRACT. In this note we show that all reductive groups are clean in characteristic ≥ 3 . In characteristic 2 there are two cuspidal local systems (one for F_4 and one for E_8) which can not be handled by our method.

1. INTRODUCTION

1.1. Let G be a connected reductive group over an algebraically closed field of characteristic $p \geq 0$. In the study of the character sheaves on the group G ([5]) an important technical assumption is that the group G is clean. This means that all cuspidal local systems (see [4]) on all Levi subgroups of G are clean (that is the IC-extension of these local systems coincides with the extension by zero). It is expected that any group G is clean; equivalently any cuspidal local system is clean for any reductive group G . It is known [5] that the last assertion is equivalent to the similar assertion with G assumed to be almost simple. It was shown in [5] that any cuspidal local system is clean if

- 1) G is of classical type and p is arbitrary;
- 2) G is of type E_6 and $p = 0$ or $p > 2$;
- 3) G is of type G_2 , F_4 , E_7 and $p = 0$ or $p > 3$;
- 4) G is of type E_8 and $p = 0$ or $p > 5$.

In [7, 8] T. Shoji used the Shintani descent theory to improve the bounds for p above for G of types G_2 , F_4 , E_8 .

The aim of this note is to present a simple argument which proves

Theorem 1. *Let G be an almost simple group. Then any cuspidal local system on G is clean except, possibly, two cases: $p = 2$ and G is of type F_4 or E_8 .*

Moreover, in each unsettled case there is exactly one cuspidal local system which is not known to be clean. In view of results of Lusztig [5] and Shoji [7, 8] Theorem 1 is new only for G of type E_6 and $p = 2$ and G of type E_7 and $p = 3$.

1.2. **Applications.** Recall that Lusztig defined a class of perverse sheaves on group G called the admissible complexes, see [4]. It is known that the character sheaves form a subset of admissible sheaves, see [5]. It is known in many cases that actually character sheaves coincide with admissible complexes, see [5, 7, 8].

Theorem 2. *The class of character sheaves coincides with the class of admissible complexes for any $p \geq 0$.*

In view of Lusztig's results in [5] 7.1 Theorem 2 is an immediate consequence of Theorem 2.12 (a) below.

Recall that Lusztig defined generalized Springer correspondence in [4] which is a bijection between the irreducible $Ad(G)$ -equivariant local systems supported on the unipotent orbits and the irreducible representations of some collection of Coxeter groups. The generalized Springer correspondence is known explicitly [6, 9,

5] in all cases with two very small gaps: in the case when the Coxeter group is of type G_2 there is an ambiguity in attaching the local systems to two-dimensional representations of this group for G of type E_6 when $p = 2$ and E_8 when $p = 3$. This ambiguity can be now removed using the method used by Lusztig in [5] 24.10 to handle a similar problem for E_6 , $p > 3$. We use below the notations from [9].

Proposition 1. *Let G be of type E_6 and $p = 2$ (respectively, of type E_8 and $p = 3$). Under the generalized Springer correspondence the reflection representation of G_2 corresponds to the local system supported on the orbit of type A_5 (respectively E_7).*

We omit the proof since it coincides with [5] 24.10 (note that the calculation of the corresponding generalized Green functions are almost identical for E_6 and E_8).

1.3. Acknowledgment. I learned the definition of the automorphism $\Theta_{\mathcal{F}}$ which is crucial for this paper from Roman Bezrukavnikov who in turn learned it from Vladimir Drinfeld. I am happy to thank both of them. I am deeply grateful to George Lusztig for very useful conversations. Thanks are also due to Toshiaki Shoji for interesting comments. This work was supported in part by NSF grants DMS-0098830 and DMS-0111298.

2. PROOFS

2.1. Let $a, p : G \times G$ be the adjoint action and the second projection respectively: $a(g, x) = gxg^{-1}$, $p(g, x) = x$. Let \mathcal{F} be a complex of (constructible) sheaves on G which is $Ad(G)$ -equivariant in the naive sense: we are given an isomorphism $\xi : a^*\mathcal{F} \rightarrow p^*\mathcal{F}$ satisfying the cocycle relation, see e.g. [7]. Let $\Delta : G \rightarrow G \times G$ be the diagonal embedding. Obviously $p\Delta = a\Delta = \text{Id}$. The following definition is crucial for this note:

Definition 2.1. We define a canonical automorphism $\Theta_{\mathcal{F}}$ of \mathcal{F} as the composition:

$$\mathcal{F} = (a\Delta)^*\mathcal{F} = \Delta^*a^*\mathcal{F} \xrightarrow{\Delta^*\xi} \Delta^*p^*\mathcal{F} = (p\Delta)^*\mathcal{F} = \mathcal{F}.$$

Remark 2.2. (i) The definition above makes sense in the case when G is a finite group. In this case $\Theta_{\mathcal{F}}$ is well known in the conformal field theory under the name of T -matrix, see e.g. [1].

(ii) The automorphism $\Theta_{\mathcal{F}}$ was used in [2, 7] to prove that the characteristic functions of character sheaves are eigenvectors for Shintani descent.

We are going to apply this definition for two kinds of complexes: the usual constructible sheaves and the perverse sheaves.

Lemma 2.3. *Let \mathcal{F} be a simple $Ad(G)$ -equivariant perverse sheaf. Then $\Theta_{\mathcal{F}} = \theta_{\mathcal{F}}\text{Id}_{\mathcal{F}}$ for some scalar $\theta_{\mathcal{F}}$.*

Proof. This is an immediate consequence of the Schur's Lemma. \square

Let \mathcal{O} be an adjoint orbit in G and let $x \in \mathcal{O}$. Recall that the functor $\mathcal{L} \mapsto \mathcal{L}_x$ defines an equivalence $\{G\text{-equivariant local systems } \mathcal{L} \text{ on } \mathcal{O}\} \rightarrow \{\text{Representations of } A_G(x) := Z_G(x)/Z_G(x)^0\}$. Let \bar{x} be the class of $x \in Z_G(x)$ in the group $A_G(x)$. Observe that $\bar{x} \in A_G(x)$ is central.

Lemma 2.4. *Under the equivalence above we have $\Theta_{\mathcal{L}} = \bar{x}|_{\mathcal{L}_x}$.*

Proof. This is a direct consequence of definition. \square

Combining Lemmas 2.3 and 2.4 one can calculate the number $\theta_{\mathcal{F}}$ for an irreducible $Ad(G)$ -equivariant perverse sheaf \mathcal{F} in the following way: take any point $x \in G$ such that $\mathcal{F}_x \neq 0$, then $\bar{x} \in A_G(x)$ acts on \mathcal{F}_x via the scalar $\theta_{\mathcal{F}}$.

Now let P be a parabolic subgroup of G with Levi quotient L . Recall (see [5]) that the induction functors $\text{Ind}_P : \{ Ad(L)\text{-equivariant sheaves on } L \} \rightarrow \{ Ad(G)\text{-equivariant sheaves on } G \}$ is defined as follows: consider the variety $\tilde{G}_P = \{(x, gP) \in G \times G/P \mid g^{-1}xg \in P\}$. The group G acts on \tilde{G}_P in the following way: $h \cdot (x, gP) = (h x h^{-1}, h g P)$. It is easy to see that we have an equivalence: $\nu : \{ G\text{-equivariant sheaves on } \tilde{G}_P \} \simeq \{ Ad(P)\text{-equivariant sheaves on } P \}$. Let $m : P \rightarrow L$ be the canonical projection and let $n : \tilde{G}_P \rightarrow G$ be defined by $n(x, gP) = x$. Then the functor $\text{Ind}_P := n_! \nu^{-1} m^*$ (we use here just naive notion of the equivariance but a similar construction holds for example in the equivariant derived category).

Lemma 2.5. *The automorphism $\Theta_{\mathcal{F}}$ commutes with the induction functor:*

$$\text{Ind}_P(\Theta_{\mathcal{F}}) = \Theta_{\text{Ind}_P(\mathcal{F})}.$$

Proof. Easy. □

Remark 2.6. A statement similar to Lemma 2.5 involving the Shintani descent twisting operator instead of $\Theta_{\mathcal{F}}$ is contained in [2, 7].

2.2. Calculation of $\theta_{\mathcal{F}}$ for some unipotent cuspidal pairs. We refer the reader to [4] for the definition of the cuspidal pair (C, \mathcal{L}) for the group G (recall only that here C is some inverse image of conjugacy class under projection $G \rightarrow G/Z_G^0$ and \mathcal{L} is some $Ad(G)$ -equivariant local system on G). In this section we consider the case when G is semisimple and C is an unipotent class (such cuspidal pairs are called unipotent) and calculate $\theta_{\mathcal{F}}$ for $\mathcal{F} = \mathcal{L}$ (extended by zero to G). First note that

Lemma 2.7. *Assume that the characteristic of k is good for G . Then $\theta_{\mathcal{L}} = 1$ for any unipotent cuspidal pair (C, \mathcal{L}) .*

Proof. Obviously it is enough to prove the Lemma for simply connected almost simple groups. For groups of type A the order of $A_G(u)$ for any unipotent element u is relatively prime to the characteristic of k and hence $\bar{u} = 1$ (since the order of u is some power of the characteristic); the result follows from Lemma 2.4. For other classical groups $A_G(u)$ is a 2-group; thus $\bar{u} \in A_G(u)$ is trivial. Similarly, for groups of type E_6, E_7 the order of $A_G(u)$ is always relatively prime with the characteristic (assumed to be good). If the group G is of type G_2, F_4, E_8 then the unipotent cuspidal local system (C, \mathcal{L}) is unique. In these cases for $u \in C$ one has $A_G(u) = S_3, S_4, S_5$ respectively. Thus $A_G(u)$ has trivial center, hence $\bar{u} = 1$. □

Now we are going to calculate $\theta_{\mathcal{L}}$ for unipotent cuspidal pairs in the exceptional groups. We are going to use the following fact due to T. Springer and B. Lou [10, 3]:

Theorem 2.8. *Let $u \in G$ be a regular unipotent element and let $U \subset G$ be the maximal unipotent subgroup containing u . Then $Z_G(u) = Z_G \times Z_U(u)$; moreover the group $Z_U(u)/Z_U^0(u)$ is cyclic and is generated by \bar{u} .*

The list of unipotent cuspidal pairs for exceptional groups is given in [9]. We give the values of $\theta_{\mathcal{L}}$ in all bad characteristic cases using the notations of *loc. cit.* In the table below ζ (respectively ξ) is a fixed primitive root of unity of degree 3 (respectively 5).

	G	$\text{char}(k)$	class of u	$A_G(u)$	ϕ	$\theta_{\mathcal{L}}$
1	G_2	2	G_2	\mathbb{Z}_2	-1	-1
2			$G_2(a_1)$	S_3	ε	1
3	G_2	3	G_2	\mathbb{Z}_3	$\zeta^{\pm 1}$	$\zeta^{\pm 1}$
4			$G_2(a_1)$	\mathbb{Z}_2	-1	1
5	F_4	2	F_4	\mathbb{Z}_4	$\pm i$	$\pm i$
6			$F_4(a_1)$	\mathbb{Z}_2	-1	-1
7			$F_4(a_2)$	D_8	ε	1
8			$F_4(a_3)$	S_3	ε	1
9	F_4	3	F_4	\mathbb{Z}_3	$\zeta^{\pm 1}$	$\zeta^{\pm 1}$
10			$F_4(a_3)$	S_4	ε	1
11	E_6	2	E_6	\mathbb{Z}_6	$-\zeta^{\pm 1}$	-1
12			$A_5 + A_1$	\mathbb{Z}_6	$-\zeta^{\pm 1}$	1
13	E_6	3	E_6	\mathbb{Z}_3	$\zeta^{\pm 1}$	$\zeta^{\pm 1}$
14	E_7	2	E_7	\mathbb{Z}_4	$\pm i$	$\pm i$
15	E_7	3	E_7	\mathbb{Z}_6	$-\zeta^{\pm 1}$	$\zeta^{\pm 1}$
16			$D_6(a_2) + A_1$	$S_3 \times \mathbb{Z}_2$	$-\varepsilon$	-1
17	E_8	2	$E_8(a_1)$	\mathbb{Z}_4	$\pm i$	$\pm i$
18			$E_7(a_2) + A_1$	$S_3 \times \mathbb{Z}_2$	$-\varepsilon$	-1
19			$D_8(a_1)$	D_8	ε	1
20			$2A_4$	S_5	ε	1
21	E_8	3	$E_7 + A_1$	\mathbb{Z}_6	$-\zeta^{\pm 1}$	$\zeta^{\pm 1}$
22			$2A_4$	S_5	ε	1
23	E_8	5	E_8	\mathbb{Z}_5	$\zeta^{\pm 1}, \zeta^{\pm 2}$	$\zeta^{\pm 1}, \zeta^{\pm 2}$
24			$2A_4$	S_5	ε	1

Comments on the calculation. In cases 1,3,5,9,11,13,14,15,23 the calculation is immediate from 2.8; in cases 2,4,7,8,10,19,20,22,24 the calculation is immediate from the fact that \bar{u} is central in $A_G(u)$. The calculation in case 12 is as follows: assume that $\bar{u} \neq 1$ in this case, then for the representation $\phi' = -1$ of $A_G(u)$ we will have $\phi'(\bar{u}) = -1$; this is a contradiction since the local system \mathcal{L}' corresponding to ϕ' appears in the principal series (see [9]) and thus we have $\theta_{\mathcal{L}'} = 1$ by Lemma 2.5; thus $\bar{u} = 1$ and the result follows. The similar method applies to cases 16,17,18,21. Finally, for the case 6 see [7] 7.2 (it is stated there that \bar{u} is nontrivial in this case; one way to see this is an explicit calculation; one can also use the results on Shintani descent, see *loc. cit.*).

Remark 2.9. For types G_2 , F_4 , E_8 the numbers in the table were computed by T. Shoji [7, 8] as the eigenvalues of the twisting operator.

The important consequence of the calculation above is the following

Corollary 2.10. *Let $\mathcal{L}_1 \neq \mathcal{L}_2$ be two cuspidal local systems on a simple group G . Then $\theta_{\mathcal{L}_1} \neq \theta_{\mathcal{L}_2}$ except for $\text{char}(k) = 2$ and G is of type F_4 or E_8 .*

Remark 2.11. Note that the results above together with the explicit knowledge of the generalized Springer correspondence for exceptional groups [9] allow to determine \bar{u} for all unipotent elements u in these groups.

2.3. Cleanness of cuspidal local systems. Recall that a local system \mathcal{L} on a locally closed subset U of a variety X is called clean if $j_!\mathcal{L} = j_{!*}\mathcal{L}$ where $j : U \subset X$ is the obvious embedding. It is expected that all cuspidal local systems are clean. Here is a main result of this note:

Theorem 2.12. (a) *Any cuspidal sheaf is a character sheaf.*

(b) *Assume that G is an almost simple exceptional group. Let (C, \mathcal{L}) be a cuspidal pair for G . Then \mathcal{L} is clean except possibly two cases: k is of characteristic 2 and*

1) G is of type F_4 and C is unipotent orbit of type $F_4(a_2)$ (there is a unique such cuspidal pair);

2) G is of type E_8 and C is unipotent orbit of type $D_8(a_1)$ (there is a unique such cuspidal pair).

Proof. The theorem is known to be true for classical groups and for exceptional groups in good characteristic [5]. The proof for exceptional groups in bad characteristic is quite similar to proofs in [5]. First one shows that any cuspidal character sheaf is clean (except, possibly, two cases in (b)) using Proposition 7.9 of [5] III with the action of the center replaced by the automorphism $\Theta_{\mathcal{F}}$ and using Corollary 2.10 (recall that it is enough to consider the cuspidal local systems supported on the unipotent orbits, see [5] 7.11). Then the results of [5] provide a classification of character sheaves. Finally one compares the list of cuspidal character sheaves with the list of cuspidal local systems (known from [4]) and deduces (a). The case when $p = 2$ and G is of type F_4 or E_8 requires additional arguments, see [7] 7.3 and [8] 5.3.

□

REFERENCES

- [1] B. Bakalov, A. Kirillov, Jr., *Lectures on tensor categories and modular functor*, Amer. Math. Soc., 2000.
- [2] M. Eftekhari, *Descente de Shintani des faisceaux caractères*, C. R. Acad. Sci. Paris Sér. I Math. **318** (1994), no. 4, 305-308.
- [3] B. Lou, *The centralizer of a regular unipotent element in a semi-simple algebraic group*, Bull. Amer. Math. Soc. **74** (1968), 1144-1146.
- [4] G. Lusztig, *Intersection cohomology complexes on a reductive group*, Inv. Math. **75** (1984), 205-272.
- [5] G. Lusztig, *Character sheaves* I, Adv. in Math. **56** (1985), 193-237; II, Adv. in Math. **57** (1985), 226-265; III, Adv. in Math. **57** (1985), 266-315; IV, Adv. in Math. **59** (1986), 1-63; V, Adv. in Math. **61** (1986), 103-155.
- [6] G. Lusztig, N. Spaltenstein, *On the generalized Springer correspondence for classical groups*, Algebraic groups and related topics (Kyoto/Nagoya, 1983), 289-316, Adv. Stud. Pure Math., 6, North-Holland, Amsterdam, 1985.
- [7] T. Shoji, *Character sheaves and almost characters on reductive groups* I, Adv. in Math. **111** (1995), 244-313.
- [8] T. Shoji, *Character sheaves and almost characters on reductive groups* II, Adv. in Math. **111** (1995), 314-354.
- [9] N. Spaltenstein, *On the generalized Springer correspondence for exceptional groups*, Algebraic groups and related topics (Kyoto/Nagoya, 1983), 317-339, Adv. Stud. Pure Math., 6, North-Holland, Amsterdam, 1985.
- [10] T. Springer, *Some arithmetical results on semi-simple Lie algebras*, Inst. Hautes Études Sci. Publ. Math. **30** (1966), 115-141.