# arXiv:math/9905005v1 [math.RT] 2 May 1999

# Raising operators for the Whittaker wave functions of the Toda chain and intertwining operators.

A.Chervov <sup>1</sup>

Moscow State University

and

Institute of Theoretical and Experimental Physics 117259 Moscow, Russia

### Abstract.

Intertwiners between representations of Lie groups can be used to obtain relations for matrix elements. We apply this technique to obtain different identities for the wave functions of the open Toda chain, in particular raising operators and bilinear relations for the wave functions at different energy levels. We also recall the group theory approach to the Toda chain: treating the wave functions as matrix elements in irreducible representations between the so-called Whittaker vectors, integral representations of the wave functions, etc.

### 1. Introduction

Interrelations between the theory of integrable systems and representation group theory proved to be fruitful for the both fields. The subject has become so vast and diverse that it looks impossible to pick up a few of surveys which would cover all the important questions. Some topics are covered in reviews [1, 2, 3] and books [4, 5]. Review [6] is closely related to our consideration. Usually it is possible to find a group theory interpretation for all the important structures of integrable systems. Phase space, symplectic structure, Hamiltonians, Lax pairs, etc. can be interpreted in terms of an underlying group symmetry for both classical and quantum integrable systems. However we will not discuss these questions here.

The main goal of this paper is to demonstrate how representation theory, and, more concretely, intertwining operators, can be used to obtain relations between wave-functions at different energy levels. We will work out only the simplest case of the open Toda chain with two and three particles, but the method can be applied to other integrable systems. The paper mainly follows the ideas of [7, 8].

Let us recall one of the main ideas of the group theory approach to integrable systems: solutions of integrable equations usually are matrix elements or traces of the groups elements in irreducible representations. Main examples we keep in mind are: the tau-function of the Kadomtsev-Petviashvili which is a matrix element in fundamental representations of the group  $GL_{\infty}$  (see [1, 6, 8]), and the wave functions of many-body problems like Toda or Calogero-Sutherland-Moser, which are matrix elements in infinite-dimensional irreducible representations of the semisimple groups (see for example [7]).

The idea to obtain different relations for wave functions is quite simple and goes as follows: let functions  $f_i$  ( $g_i$ ) be solutions of some integrable system corresponding to some representation  $V_i$  ( $W_i$  respectively) of some group, (it can be the trace or matrix element, can be a wave function or a tau function, etc.) If one has an intertwinning operator  $V_1 \otimes V_2 \otimes ... \otimes V_n \to W_1 \otimes W_2 \otimes ... \otimes W_m$  then there 'usually' exist the following relation:  $D(f_1 f_2 ... f_n) = \tilde{D}(g_1 g_2 ... g_m)$ , where  $D, \tilde{D}$  are some differential operators. Miraculously enough this simple idea leads to non-trivial relations. It was demonstrated in [8, 9] that in such way one can obtain famous bilinear identities for KP,KdV tau-functions, which corresponds to intertwiners between fundamental representations of the  $GL_{\infty}$ ,  $\hat{SL}(n)$ . In [10], it was shown, that, in this way, one can obtain the Baxter relation for transfer matrices of integrable lattice models. The above idea is also used in representation theory approach to Knizhnik-Zamolodchikov equation [12, 11, 5], and in the theory of special functions (see, for example [13, 14]).

<sup>&</sup>lt;sup>1</sup>E-mail: chervov at vitep1.itep.ru, alex at lchervova.home.bio.msu.ru

<sup>2&#</sup>x27;usually' means that there are no such relations for generic matrix elements, while they typically emerge for solutions of integrable systems.

In this paper, we show how to obtain different relations for the Toda wave-functions applying the above idea. The group theory approach to the classical Toda chain has been first proposed in [15], group theory approach to the quantum Toda chain has been proposed by Kostant and Kazhdan at late seventies (unpublished) and further elaborated in [16], [7]. Let us recall it following [7]. Consider the group  $SL(n+1,\mathbb{R})$  with the standard Chevalley generators  $e_i$ ,  $h_i$ ,  $f_i$ , and let  $V_{\lambda}$  be irreducible representations with weight  $\lambda$  of the principal series with weight  $\lambda$ . Let us introduce the so called Whittaker vector  $|w\rangle \in V_{\lambda}$ , i.e. the eigenvector of generators  $e_i$ . Then, one considers the Whittaker function, i.e. matrix element  $W_{\lambda}(\phi_1,...,\phi_n) = \langle w|exp(\phi_1h_1+\phi_2h_2+...+\phi_nh_n)|w\rangle$ . The main observation is that function  $W_{\lambda}(\phi_1,...,\phi_n)$  is a common eigenfunction of Toda Hamiltonians, which turn out to be Casimirs of the group  $SL(n+1,\mathbb{R})$ , while the weight  $\lambda$  marks the corresponding energy level. (Existence and other issues on Whittaker vectors were deeply explored by Kostant [17]).

The relations between Whittaker functions  $W_{\lambda}(\phi)$  for different  $V_{\lambda}$  will be obtained as follows: let us find the intertwinning operators  $V_{\lambda_1} \otimes V_{fin} \to V_{\lambda_2}$  then we will show that there exists a relations  $\mathcal{D}\left(W_{\lambda_1}(\phi_1,...,\phi_n)\right) = \tilde{\mathcal{D}}\left(W_{\lambda_2}(\phi_1,...,\phi_n)\right)$  (see formulas 29,71) where  $\mathcal{D},\tilde{\mathcal{D}}$  are some differential operators, and  $V_{fin}$  is any finite-dimensional representation of the group  $SL(n+1,\mathbb{R})$ . This relation is an example of the so-called raising operators [18]. This idea was suggested to us by S. Kharchev to whom we are deeply indebted. Analogously we will construct bilinear relations from the intertwining operators  $V_{\lambda_1} \otimes V_{\lambda_2} \to V_{\lambda_3} \otimes V_{\lambda_4}$ . One can proceed this way to construct other relations like above, for example  $V_{\lambda_1} \otimes V_{\lambda_2} = \bigoplus_i V_{\lambda_i}$  then one can obtain relation like  $W_{\lambda_1} \otimes W_{\lambda_2} = \sum_i D_i(W_{\lambda_i})$  (see formula 39).

Let us also note that the existence of the bilinear identities in this situation is quite surprising, because it means that wave-function, which are the eigen-functions of Toda Hamiltonians, i.e. eigenfunctions of some *linear* differential operators, satisfy some system of *nonlinear* differential equations. One cannot separate some finite number of wave-functions from this system, only infinite set of them is closed nonlinear system. Nevertheless it is some system of nonlinear equations, but we do not know its sense, its hamiltonian description (if it is) or anything else, it seems to be interesting to clarify it.

Let us describe what kind of technical problems arise in proceeding to the concrete formulas. Let  $|w\rangle_{\lambda_1}$  be the Whittaker vector in representation  $V_{\lambda_1}$ , let us consider the intertwiner  $\Phi: V_{\lambda_1} \to V_{\lambda_2} \otimes V_{fin}$  then  $\Phi|w\rangle_{\lambda_1} = \sum_i |w\rangle_i \otimes |i\rangle$  where  $|i\rangle$  is some basis in finite dimensional representation  $V_{fin}$ . to construct the explicit relations one needs to find such polynomials  $P_i$  that  $|w\rangle_i = P_i(f_i, h_i)|w\rangle_{\lambda_2}$  where  $P_i$  are some polynomials of generators  $f_i, h_i$ , i.e. one needs to express the components of the image of Whittaker vector in spaces  $V_{\lambda_1}$  as application of some operators to Whittaker in space  $V_{\lambda_2}$ . This is the main technical problem we treat in this paper, we solve it for the cases  $SL(2,\mathbb{R})$  and  $SL(3,\mathbb{R})$ , but we were unable to find the general formula for the case of  $SL(n,\mathbb{R})$ .

Let us mention that the SL(2) Toda equation reduces to the well-known textbook equation, whose solutions are Bessel and Mcdonald functions (depends on necessary asymptotics, we need Mcdonald one), and the relation from theorem 2.11 one can find, for example in [19].

Let us argue that the scheme like above can be applied to other integrable models. One knows that if instead of the Whittaker vector  $|w\rangle$  one considers a spherical vector  $|s\rangle$  i.e. a vector such that it is remains fixed under the action of the maximal compact subgroup then one can obtain wave function for Calodgero-Sutherland integrable system zonal spherical function, obviously if one considers different intertwiners one can obtain different relations between wave functions. Also instead of finite-dimensional Lie group G one can takes affine group G, which corresponds to periodic Toda chain, or quantum group  $U_q(G)$ , which leads to difference equations [20, 21]. Obviously, our scheme of obtaining different relations will work in this situations also. But of course explicit constructions of these relations seems to be not obvious.

We should also mention that raising operators considered here were widely explored recently [18], but all approaches are completely different from our's and very simple idea, that the origin of this relations is presence of the intertwining operators, was not present in the literature to our knowledge. Let us repeat that from our point of view one can consider all models in the same setup. Also we see that the group theory approach shows that such at first sight not related subjects such as Baxter relation important in lattice models [10] and conformal field theory [22], raising operators in integrable quantum many body problems [18] and bilinear relations in the theory of classical nonlinear equations [1, 6] have the same explanations from the point of view of intertwining operators.

Exposition is organized as follows: section 2 is devoted to the case of SL(2) and the demonstration of the main ideas. In the section 3 we consider the case of SL(3), which is mainly analogous to the SL(2) case, but that is the case, where first arises the problems in writing the explicit formulas for the action of the intertwining operators on the Whittaker vectors. And we show how these problems can be solved for SL(3). Section 4 is devoted to the SL(N) case. We recall representation theory approach to the SL(N) Toda chain and discuss the problem of obtaining explicit raising operators. In section 5 we make some concluding remarks.

# 2. SL(2) Toda Chain.

In this section we will show how to obtain the raising operators, bilinear, etc. identities for the wave functions of SL(2) Toda. We will do it with the help of the intertwining operators between the tensor products of representations

of SL(2). Before doing this in subsections 1 and 2 we will recall representation theory interpretation of the Toda wave functions as matrix element in irreducible representation, here we closely follow [7]. Let us sketch the content: we consider Whittaker vector  $|w\rangle$  in irreducible representation  $V_{\lambda}$  of SL(2), i.e. such vector that it is eigenvector for the generator e, consider Whittaker function  $W(\phi) = \langle w|exp(\phi h)|w \rangle$ , we show that it is eigenfunction for Toda hamiltonian  $\frac{1}{2}\partial_{\phi}^2 + exp(-2\phi) + \partial_{\phi}$ , the reason is that this Hamiltonian is simply the Casimir of SL(2), then we consider intertwining operator  $\Phi: V_{\lambda+1} \to V_{\lambda} \otimes V_1$ , where  $V_1$  is standard two-dimensional representation of SL(2), and the main technical point is to show that  $\Phi(|w>_{\lambda+1}) = \frac{1}{\lambda+1} \frac{1}{2} (h+\lambda+2) |w>_{\lambda} \otimes |0> +\mu_R|w>_{\lambda} \otimes |1>$ , it is quite obvious for the case of SL(2), because we know the representation of the Whittaker vector in terms of the action of the generator f on the vacuum vector in module  $V_{\lambda}$ , the fact that such representation for the case of SL(N) is unreasonable is the main problem for writing the explicit formulas for the SL(N) case. Then we proceed to the bilinear relations, to find them one should find the intertwining operator between some representations:  $V_{\lambda_1} \otimes V_{\lambda_2} \to V_{\lambda_3} \otimes V_{\lambda_4}$ . To find such intertwiners is not an easy problem, but there is a trick which is due to [8] that works as follows: if we have intertwiners :  $V_{\lambda+1} \to V_{\lambda} \otimes V_1$  and  $V_{\mu+1} \to V_{\mu} \otimes V_1$  we can consider their tensor product :  $V_{\lambda+1} \otimes V_{\mu+1} \to V_{\lambda} \otimes V_1 \otimes V_{\mu} \otimes V_1$  then consider intertwiner:  $V_1 \otimes V_1 \to \mathbb{C}$ , then consider the composition:  $V_{\lambda_1+1} \otimes V_{\mu_1+1} \to V_{\lambda_1} \otimes V_1 \otimes V_{\mu_1} \otimes V_1 \to V_{\lambda+1} \otimes V_{\mu+1}$ , hence we get desirable intertwiner (consideration instead of  $V_1$  other finite dimensional representations leads to various kinds of necessary intertwiners).

2.1. **Notations:** Let us denote by  $(\lambda, n)$  the product  $(\lambda)(\lambda-1)(\lambda-2)...(\lambda-n+1)$ , we will often use this abbreviation. The Lie algebra SL(2) is defined by the generators e, f, h satisfying the following relations:

$$[e, f] = h, [h, e] = 2e, [h, f] = -2f.$$

The fundamental representation:

$$h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \ e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \ f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \tag{1}$$

The quadratic Casimir operator:

$$C = (ef + fe) + \frac{1}{2}h^2 = 2fe + h + \frac{1}{2}h^2.$$
 (2)

Highest weight representation:

We consider the principal (spherical) series of representations, induced by the one-dimensional representations of the Borel subgroup. The space of representation  $V_{\lambda}$  is functions of one real variable x and matrix elements are defined by integrals with the flat measure. The action of the algebra is given by differential operators:

$$e = \frac{\partial}{\partial x}, \ h = -2x\frac{\partial}{\partial x} + \lambda, \ f = -x^2\frac{\partial}{\partial x} + \lambda x.$$
 (3)

The action of the group is given by the formula:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} : x^k \to (a+cx)^{\lambda-k} (b+dx)^k \tag{4}$$

Obviously that Constants are vacuum (i.e. the highest weight) vectors for such representation i.e. e(Constant) = $0, h(Constant) = \lambda(Constant)$ 

2.2. Whittaker function. Definition 2.1: The vector  $|w\rangle_{\lambda}^{\mu}$  is called Whittaker vector in representation  $V_{\lambda}$ , if it is eigenvector for generator e i.e.  $e|w>^{\mu}_{\lambda}=\mu|w>^{\mu}_{\lambda}$  for some constant  $\mu$ .

Lemma 2.2: For given  $\mu$  and irreducible representation  $V_{\lambda}$  ( $\lambda \neq 1, 2, 3, ...$ ) there exists a unique (up to a scalar) Whittaker vector  $|w\rangle_{\lambda}^{\mu} \in V_{\lambda}$ , which can be expressed as follows:

$$|w\rangle_{\lambda}^{\mu} = exp(\mu x) = \sum_{i=0}^{\infty} \frac{\mu^n f^n}{n!(\lambda, n)} |vac\rangle$$
(5)

where we denote  $(\lambda, n) = \lambda(\lambda - 1)...(\lambda - n + 1)$  and  $|vac\rangle$  - vacuum (the highest weight) vector in  $V_{\lambda}$ .

The proof is obvious.

The dual Whittaker vector:  ${}^{\mu}_{\lambda} < w|$  is given  $\sum_{n=0}^{\infty} {}_{\lambda} < 0| \frac{\mu^n e^n}{n!(\lambda,n)} = x^{-\lambda-2} exp(-\mu x)$  (see [7]). Definition 2.3: The Whittaker function  $W^{\mu_L,\mu_R}_{\lambda}(\phi)$  is the function given by:

$$W_{\lambda}^{\mu_L,\mu_R}(\phi) = {}^{\mu_L}_{\lambda} < w|exp(\phi h)|w>_{\lambda}^{\mu_R}$$
 (6)

Remark: we will sometimes omit indexes  $\lambda, \mu$  of Whittaker vectors and Whittaker functions, if they are not important.

Proposition 2.4:  $W_{\lambda}(\phi) = exp(\lambda\phi) \sum_{i=0}^{\infty} \frac{(\mu_R)^i (\mu_L)^n exp(-2i\phi)}{i!(\lambda,i)}$ The proof is obvious recalling (6) and  $\langle vac|e^n|f^n|vac \rangle = n!(\lambda,n)$ 

Remark: for the case  $SL(2,\mathbb{R})$  Whittaker function coincides with the textbook Macdonald function (up to some exponents and scalars), this can be easy seen from the integral representation below, see also [7].

Proposition 2.5: The Whittaker function satisfies the equation:

$$\left[\frac{1}{2}\frac{\partial^2}{\partial\phi^2} + \frac{\partial}{\partial\phi} - 2\mu_R\mu_L e^{-2\phi}\right]W_\lambda(\phi) = \left(\frac{1}{2}\lambda^2 + \lambda\right)W_\lambda(\phi) \tag{7}$$

Hence, the function  $\Psi_{\lambda}(\phi) = exp(\phi)W_{\lambda}(\phi)$  satisfies two particle open Toda equation:

$$\left[\frac{1}{2}\frac{\partial^2}{\partial\phi^2} - 2\mu_R\mu_L e^{-2\phi}\right]\Psi_\lambda(\phi) = \left(\frac{1}{2}\lambda^2 + \lambda\right)\Psi_\lambda(\phi),\tag{8}$$

The proof is very simple, but we will reproduce it, since it is the same for all kinds of integrable many body problems. The main idea is that the Hamiltonian is a Casimir operators, hence it acts by scalar on every matrix element, that's how RHS of 7 appears, on the other hand it is some concrete operator - that's how LHS of 7 appears.

Proof: The Casimir operator  $C=2fe+h+\frac{1}{2}h^2$  acts as scalar  $\frac{1}{2}\lambda^2+\lambda$  on the representation  $V_{\lambda}$ . Hence:  $< w|exp(\phi h)C|w>=(\frac{1}{2}\lambda^2+\lambda)W_{\lambda}(\phi)$  On the other hand  $< w|exp(\phi h)C|w>=< w|exp(\phi h)(\frac{1}{2}h^2+h+2fe)|w>=(\frac{1}{2}\frac{\partial^2}{\partial\phi^2}+\frac{\partial}{\partial\phi})W_{\lambda}(\phi)+2< w|f|exp(\phi(h-2))e|w>=(\frac{1}{2}\frac{\partial^2}{\partial\phi^2}+\frac{\partial}{\partial\phi}+2\mu_R\mu_lexp(-2\phi))W_{\lambda}(\phi)$  hence we obtain RHS of 7. Q.E.D.

Remark: any matrix element of the irreducible representation is eigenfunction if all the Casimir operators of any group. It may seem that hence we obtain integrable system. Of course, it's not true, because usually the dimension of a group is something like  $n^2$  where the number of independent Casimirs is something like n. So one does not have enough Hamiltonians in involution. So one should reduce the phase space from the hole group to some submanifold of it, but the problem is that Casimir operator acting on the function on this submanifold can give a function beyond that class. In our case we saw that  $\langle w|exp(\phi h)C|w \rangle = (\frac{1}{2}\frac{\partial^2}{\partial \phi^2} + \frac{\partial}{\partial \phi} + 2\mu_R\mu_lexp(-2\phi))W_\lambda(\phi)$ . This is due to the clever choice: we considered matrix element of the Cartan group element between Whittaker vectors, the idea to do so is due to Kazhdan and Kostant. For example if one takes some arbitrary vectors  $\langle u|, |v\rangle$  then  $\langle u|exp(\phi h)C|v\rangle = (\frac{1}{2}\frac{\partial^2}{\partial \phi^2} + \frac{\partial}{\partial \phi})W_\lambda(\phi) + 2\langle u|exp(\phi h)fe|v\rangle$ , and one can do nothing with the  $\langle u|exp(\phi h)fe|v\rangle$  one cannot in general express it via  $\langle u|exp(\phi h)|v\rangle$ . So one cannot obtain the LHS of (7), despite one has RHS. The general conception of the consistent hamiltonian reductions is due to Drinfeld and Sokolov [3].

Proposition 2.6 (Integral representation of the Whittaker function):

$$W_{\lambda}(\phi) = \left(\frac{1}{\mu_R}\right)^{-(\lambda+1)} \int_0^{\infty} x^{-(\lambda+2)} e^{-\frac{\mu_L \mu_R e^{-2\phi}}{x} - x} dx = 2e^{(\lambda+1)\phi} \left(\sqrt{\frac{\mu_L}{\mu_R}}\right)^{-(\lambda+1)} K_{\lambda+1} (2\sqrt{\mu_L \mu_R} e^{-\phi}) dx. \tag{9}$$

- where  $K_{\lambda}(z)$  is the Macdonald function.

Remark: for the convergence we should require  $Re\mu_L\mu_R > 0$ .

The theorem is tautology after recalling that the invariant pairing  $\langle v|u\rangle$  is realized as integral  $\int_0^\infty$ , and dual Whittaker vector  $\langle w|$  is realized as  $x^{-2(\lambda+1)}e^{-\frac{\mu_L}{x}}$ . The second equality is the textbook integral formula for the Macdonald function.

2.3. Intertwining operators. Denote by  $V_1$  the fundamental representation of the  $SL(2,\mathbb{R})$ , denote |0> the highest weight vector, |1> the lowest weight vector, i.e.

$$|0\rangle = \begin{pmatrix} 1\\0 \end{pmatrix}, |1\rangle = \begin{pmatrix} 0\\1 \end{pmatrix}, \tag{10}$$

hence e|0>=0, f|0>=|1>, h|0>=|0>; e|1>=|0>, f|1>=0, h|1>=-|0>.

Proposition 2.7: The isomorphism  $\Phi_{\lambda} = \Phi_{\lambda,+} \oplus \Phi_{\lambda,-} : V_{\lambda+1} \oplus V_{\lambda-1} \to V_{\lambda} \otimes V_1$  is given by the formulas:

$$\Phi_{\lambda,+}: f^n|vac\rangle_{\lambda+1} \to f^n|vac\rangle_{\lambda} \otimes |0\rangle + nf^{n-1}|vac\rangle_{\lambda} \otimes |1\rangle, \tag{11}$$

$$\Phi_{\lambda,+}^{-1}: f^{n}|vac>_{\lambda} \otimes |0> \to \frac{(\lambda+1-n)}{\lambda+1} f^{n}|vac>_{\lambda+1}, \ \Phi_{\lambda,+}^{-1}: f^{n}|vac>_{\lambda} \otimes |1> \to \frac{1}{\lambda+1} f^{n+1}|vac>_{\lambda+1}, \ \ (12)$$

$$\Phi_{\lambda,-}: f^n|vac>_{\lambda-1} \to f^{n+1}|vac>_{\lambda} \otimes |0> + (n-\lambda)f^n|vac>_{\lambda} \otimes |1>, \tag{13}$$

$$\Phi_{\lambda,-}^{-1}: f^n|vac>_{\lambda}\otimes|0> \to \frac{n}{\lambda+1}f^{n-1}|vac>_{\lambda-1}, \ \Phi_{\lambda,-}^{-1}: f^n|vac>_{\lambda}\otimes|1> \to \frac{-1}{\lambda+1}f^n|vac>_{\lambda-1}.$$
 (14)

The action of dual operators  $\Phi^*$ ,  $\Phi_1^*$ ,  $\Phi_{-1}^*$  on the dual anti-representations are given by the same formulas (up to scalars) with the change of f to e, which follows from the Chevalley antiinvolution:

$$\Phi_1^*:_{\lambda+1} < vac|e^n \to_{\lambda} < vac|e^n \otimes <0| + n_{\lambda} < vac|e^{n-1} \otimes <1|$$

$$\tag{15}$$

$$\Phi_1^{-1*}:_{\lambda} < vac|e^n \otimes <0| \rightarrow \frac{(\lambda+1-n)}{\lambda+1} |_{\lambda+1} < vac|e^n, |\Phi_1^{-1*}:_{\lambda} < vac|e^n \otimes <1| \rightarrow \frac{1}{\lambda+1} |_{\lambda+1} < vac|e^{n+1}, (16)$$

$$\Phi_{-1}^*: {}_{\lambda-1} < vac|e^n \to \frac{1}{\lambda+1} ({}_{\lambda} < vac|e^{n+1} \otimes <0| + (n-\lambda)_{\lambda} < vac|e^n \otimes <1|)$$

$$\tag{17}$$

$$\Phi_{-1}^{-1*}:_{\lambda} < vac|e^{n} \otimes <0| \to (n\lambda)_{\lambda-1} < vac|e^{n-1}, \ \Phi_{-1}^{-1*}:_{\lambda} < vac|e^{n} \otimes <1| \to (-\lambda)_{\lambda-1} < vac|e^{n}. \tag{18}$$

The invariant pairing is given by: <0|0>=1, <0|1>=0, <1|0>=0, <1|1>=1.

The proof is obvious, let us only note that in order to find the intertwiners  $\Phi_{\pm 1}^{-1}$  from  $V_{\lambda\pm 1}\otimes V_1\to V_{\lambda}$  it is more convinient to express them via the dual intertwiners:  $V_{\lambda\pm 1}\to V_1^*\otimes V_{\lambda}$ , which can be easily found.

Proposition 2.8: Intertwiners act on Whittaker vectors as follows:

$$|w\rangle_{\lambda+1} \to \frac{(h+\lambda+2)}{2(\lambda+1)}|w\rangle\otimes|0\rangle + \frac{\mu_R}{\lambda+1}|w\rangle\otimes|1\rangle,\tag{19}$$

$$|w>_{\lambda-1} \to \frac{\lambda}{\mu_R} \frac{1}{2} (-h+\lambda)|w> \otimes |0> -\lambda|w> \otimes |1>, \tag{20}$$

$$|w>_{\lambda}\otimes|0> \to |w>_{\lambda+1} + \frac{\mu_R}{\lambda(\lambda+1)}|w>_{\lambda-1}, \tag{21}$$

$$|w\rangle_{\lambda}\otimes|1\rangle \to \frac{(\lambda+1-h)}{2\mu_R}|w\rangle_{\lambda+1} - \frac{\lambda+1+h}{2\lambda}|w\rangle_{\lambda-1}, \tag{22}$$

$$_{\lambda} < w | \otimes < 0 | \rightarrow \mu_{L \lambda - 1} < w | + _{\lambda + 1} < w |, \tag{23}$$

$$_{\lambda} < w | \otimes < 1 | \rightarrow -\frac{h + \lambda + 1}{2\lambda} |_{\lambda - 1} < w | \otimes < 1 | + \frac{\lambda + 1 - h}{2\mu_L} |_{\lambda + 1} < w |. \tag{24}$$

Proof: The proof is simple, let us consider only the first equality. We will use the following trivial fact:

 $(\lambda - n + 1)f^n|vac\rangle_{\lambda} = \frac{1}{2}(\lambda + 2 + h)f^n|vac\rangle_{\lambda}.$ 

$$\Phi_{1}(|w>_{\lambda+1}) = \Phi_{1}(\sum_{n=0}^{\infty} \frac{\mu_{R}^{n}}{n!(\lambda+1,n)} f^{n} |vac>_{\lambda} + 1) = \sum_{n=0}^{\infty} \frac{\mu_{R}^{n}}{n!(\lambda+1,n)} (f^{n} |vac>_{\lambda} \otimes |0> + nf^{n-1} |vac>_{\lambda} \otimes |1>) = \frac{1}{\lambda+1} \sum_{n=0}^{\infty} \frac{\mu_{R}^{n}}{(n)!(\lambda,n)} (\lambda-n+1) f^{n} |vac>_{\lambda} \otimes |0> + \frac{\mu_{R}}{\lambda+1} |w>_{\lambda} \otimes |1> = \frac{(\lambda+2+h)}{2(\lambda+1)} |w>_{\lambda} \otimes |0> + \frac{\mu_{R}}{\lambda+1} |w>_{\lambda} \otimes |1>.$$
O.E.D.

Remark: let us note that such simple formulas like above cannot be true for an arbitrary vector  $|v\rangle_{\lambda} \in V_{\lambda}$  i.e. of course from the irredicibility of repsentations follows that there exist series O(f,h,e) such that  $\Phi_1(|v\rangle_{\lambda+1}) = O(f,h,e)|v\rangle_{\lambda}$ , but in general such expression will be complicated.

Let us find the intertwiners between tensor product of infinite-dimensional representations.

Proposition 2.9: the isomorphism  $\Phi = \bigoplus_{k=0}^{\infty} \Phi_k : V_{\lambda} \otimes V_{\nu} \to \bigoplus_{k=0}^{\infty} V_{\lambda+\nu-k}$  in realization (3) is given by the formulas:

$$\Phi_k: x^n \otimes x^m \to C_k \sum_{i=0}^k (-1)^i \binom{n}{k-i} \binom{m}{i} \frac{(\nu)_k}{(\nu)_i} \frac{(\lambda)_k}{(\lambda)_{k-i}} x^{n+m-k}, \tag{25}$$

where 
$$C_k = \frac{\left(\prod_{i=1}^k (i(\nu^2(\lambda - i + 1) + \lambda^2(\nu - i + 1))\right)^{\frac{1}{2}}}{(\nu\lambda)^k (\sum_{i=0}^k \frac{(\nu)_k}{(\nu)_i} \frac{(\lambda)_k}{(\lambda)_{k-i}})}$$
 (26)

Or one can rewrite it as follows. Denote by  $mul_k$  operator  $V_{\lambda} \otimes V_{\nu} \to V_{\lambda+\nu-k}$  which acts:  $f(x) \otimes g(x) \to f(x)g(x)$ . Then:

$$\Phi_k: f(x) \otimes g(x) \to C_k(mul_k) * (\sum_{i=0}^k \frac{(-1)^i}{(k-i)!i!} \frac{(\nu)_k}{(\nu)_i} \frac{(\lambda)_k}{(\lambda)_{k-i}} \partial_x^{k-i} f(x) \otimes \partial_x^i g(x)).$$
 (27)

The proof can be found in the literature, or proceeded as follows: from the equality:  $\Phi_k * h = h * \Phi_k$  follows  $\Phi_k(x^n \otimes x^m) = c_{n,m}x^{n+m-k}$ , using  $\Phi_k * e = e * \Phi_k$  one obtains the recurrence relation for the coefficients  $c_{n,m}$ . Solution of this relation can be guessed after considering several examples of small k = 0, 1, 2.

Proposition 2.10: The action of  $\Phi_k$  on Whittaker vectors is given by:

$$\Phi_k: |w>_{\lambda}^{\mu_1} \otimes |w>_{\nu}^{\mu_2} \to C_k \sum_{i=0}^k \frac{(-1)^i}{(k-i)!i!} \frac{(\nu)_k}{(\nu)_i} \frac{(\lambda)_k}{(\lambda)_{k-i}} (\mu_1)^{k-i} (\mu_2)^i) |w>_{\lambda+\nu-k}^{\mu_1+\mu_2}. \tag{28}$$

This easily follows from 27.

### 2.4. Relations for the Whittaker wave functions.

Theorem (Raising operators) 2.11 The following relations holds:

$$W_{\lambda+1}(\phi) = exp(\phi) \frac{\partial_{\phi} + \lambda + 2}{2(\lambda+1)} W_{\lambda}(\phi), \ \mu_L W_{\lambda-1}(\phi) = \frac{\lambda}{\mu_R} exp(\phi) \frac{-\partial_{\phi} + \lambda}{2} W_{\lambda}(\phi). \tag{29}$$

Remark: easy to see that these relations are consistent: application of one then another gives that  $W_{\lambda}$  satisfy Toda equation.

Proof: We will calculate  $_{\lambda} < w | \otimes < 0 | \Phi_1 exp(\phi h) | w >_{\lambda+1}$  in two ways, first we will apply  $\Phi_1$  to  $exp(\phi h) | w >_{\lambda+1}$  and then take pairing, hence we will obtain the RHS of (29); to obtain the LHS of (29) we will apply  $\Phi_1^*$  to  $_{\lambda} < w | \otimes < 0 |$  and consider the pairing after.

 $_{\lambda} < w | \otimes < 0 | \ \Phi_1 exp(\phi h) | w>_{\lambda+1} = _{\lambda} < w | \otimes < 0 | \ exp(\phi h) \Phi_1 | w>_{\lambda+1} = _{\lambda} < w | \otimes < 0 | \ exp(\phi h) (\frac{(h+\lambda+2)}{2(\lambda+1)} | w>_{\lambda} \otimes | 0> + \frac{\mu_R}{\lambda+1} | w> \otimes | 1>) = _{\lambda} < w | \otimes < 0 | \ \frac{(h+\lambda+2)}{2(\lambda+1)} exp(\phi h) | w>_{\lambda} \otimes exp(\phi h) | 0> = exp(\phi) \frac{(\partial_{\phi}+\lambda+2)}{2(\lambda+1)} _{\lambda} < w | \otimes < 0 | \ exp(\phi h) | w>_{\lambda} \otimes exp(\phi h) | 0> = exp(\phi) \frac{(\partial_{\phi}+\lambda+2)}{2(\lambda+1)} _{\lambda} < w | \otimes < 0 | \ exp(\phi h) | w>_{\lambda} \otimes exp(\phi h) | 0> = exp(\phi) \frac{(\partial_{\phi}+\lambda+2)}{2(\lambda+1)} _{\lambda} < w |$ 

 $_{\lambda} < w | \otimes < 0 | \Phi_1 exp(\phi h) | w>_{\lambda+1} = (_{\lambda} < w | \otimes < 0 | \Phi_1^*) | exp(\phi h) | w>_{\lambda+1} = _{\lambda+1} < w | exp(\phi h) | w>_{\lambda+1} = W_{\lambda+1}(\phi).$  Hence we get the LHS of (29). Q.E.D.The proof of the second equality in (29) is the same.

One can deduce directly from the (29) or deduce representation theoretically (as we will do for the demonstration of the idea) the following relations:

Theorem (Baxter like relations) 2.12:

$$exp(\phi)W_{\lambda}(\phi) = \frac{\mu_{L}\mu_{R}}{\lambda(\lambda+1)}W_{\lambda-1}(\phi) + W_{\lambda+1}(\phi),$$

$$exp(-\phi)W_{\lambda}(\phi) = \left(\frac{\partial_{\phi} + \lambda + 1}{2}\right)^{2}W_{\lambda-1}(\phi) + \frac{1}{\mu_{L}\mu_{R}}\left(\frac{\lambda + 1 - \partial_{\phi}}{2}\right)^{2}W_{\lambda+1}(\phi).$$
(30)

Remark: analogous considerations for the infinite-dimensional algebras leads to Baxter relations for the quantum transfer matrices in lattice models, see [10].

Proof: We will use isomorphism  $\Phi: V_{\lambda+1} \oplus V_{\lambda-1} \to V_{\lambda} \otimes V_1$  and calculate the pairing  $< 0 | < w | exp(\phi h) | w > | 0 >$  first directly, obtaining LHS, then applying isomorphism  $\Phi^{-1}$  to |w| > | 0 > and  $\Phi^{-1}$  to |w| < 0 |, and obtain RHS.

$$\exp(\phi)W_{\lambda}(\phi) = {}_{\lambda} < w | \otimes < 0 | \exp(\phi h) | 0 > \otimes | w >_{\lambda} = 
(\mu_{L | \lambda - 1} < w | + |_{\lambda + 1} < w |) \exp(\phi h) (\frac{\mu_{R}}{\lambda(\lambda + 1)} | w >_{\lambda - 1} + | w >_{\lambda + 1}) = 
= \frac{\mu_{L} \mu_{R}}{\lambda(\lambda + 1)} |_{\lambda - 1} < w | \exp(\phi h) | w >_{\lambda - 1} + |_{\lambda + 1} < w | \exp(\phi h) | w >_{\lambda + 1} = \frac{\mu_{L} \mu_{R}}{\lambda(\lambda + 1)} W_{\lambda - 1}(\phi) + W_{\lambda + 1}(\phi).$$
(31)

Q.E.D.The proof of the second relation is the same, but one should consider  $_{\lambda} < w | \otimes < 1 | exp(\phi h) | 1 > \otimes | w >_{\lambda}$  instead of  $_{\lambda} < w | \otimes < 0 | exp(\phi h) | 0 > \otimes | w >_{\lambda}$ . Obviously  $0 = _{\lambda} < w | \otimes < 0 | exp(\phi h) | 1 > \otimes | w >_{\lambda}$ , so one cannot obtain any more relations like above.

Theorem (Bilinear relations) 2.13:

$$\left(-\frac{\mu_2^R exp(\phi_1 - \phi_2)}{2(\lambda + 1)(\nu + 1)}(\partial_{\phi_1} + \lambda + 2) + \frac{exp(\phi_2 - \phi_1)\mu_1^R}{2(\lambda + 1)(\nu + 1)}(\partial_{\phi_2} + \nu + 2)\right)W_{\lambda}^{\mu_1^L, \mu_1^R}(\phi_1)W_{\nu}^{\mu_2^L, \mu_2^R}(\phi_2) =$$
(32)

$$= \left(-\frac{\nu + 1 - \partial_{\phi_2}}{2\mu_2^L} + \frac{\lambda + 1 - \partial_{\phi_1}}{2\mu_1^L}\right) W_{\lambda+1}^{\mu_1^L, \mu_1^R}(\phi_1) W_{\nu+1}^{\mu_2^L, \mu_2^R}(\phi_2)$$
(33)

Proof: let us denote by  $\Phi_{\lambda,\nu}$  the intertwiner:  $V_{\lambda+1} \otimes V_{\nu+1} \to V_{\lambda} \otimes V_{\nu}$ . we will construct that intertwiner using a trick due to [8]: let us denote by S the intertwiner  $V_1 \otimes V_1 \to \mathbb{C}$ , than  $\Phi_{\lambda,\nu}$  is given by composition:  $(id \otimes S \otimes id)(\Phi_{\lambda,+} \otimes \Phi_{\nu,+})$ . Lemma 2.14: The action of the  $\Phi_{\lambda,\nu}$  on Whittaker vectors is given by the formulas:

$$\Phi_{\lambda,\nu} : exp(\phi_1 h)|w>_{\lambda+1}^{\mu_1^R} \otimes exp(\phi_2 h)|w>_{\nu+1}^{\mu_2^R} \to \\
-exp(\phi_1 - \phi_2) \frac{h + \lambda + 2}{2(\lambda + 1)} exp(\phi_1 h)|w>_{\lambda}^{\mu_1^R} \otimes \frac{\mu_2^R exp(\phi_2 h)}{\nu + 1} exp(\phi_2 h)|w>_{\nu}^{\mu_2^R} +$$
(34)

$$+exp(\phi_{2}-\phi_{1})\frac{\mu_{1}^{R}exp(\phi_{1}h)}{(\lambda+1)}|w>_{\lambda}^{\mu_{1}^{R}}\otimes\frac{(h+\nu+2)}{2(\nu+1)}exp(\phi_{2}h)|w>_{\nu}^{\mu_{2}^{R}},$$
(35)

$$\Phi_{\lambda,\nu}^*: \ ^{\mu_1^L}{}_{\lambda} < w | \otimes \ ^{\mu_2^L}{}_{\nu} < w | \rightarrow - \frac{\mu_1^L}{\lambda + 1} < w | \otimes \ ^{\mu_2^L}{}_{\nu + 1} < w | \frac{\nu + 1 - h}{2\mu_2^L} + \frac{\mu_1^L}{\lambda + 1} < w | \frac{\lambda + 1 - h}{2\mu_1^L} \otimes \ ^{\mu_2^L}{}_{\nu + 1} < w |. \ \ (36)$$

The proof of the lemma follows from (11) and recalling the fact that action of S is given by: S(|0>|1>)=0, S(|1>|0>)=0, S(|1>|0>)=1, S(|0>|1>)=-1 and respectively:  $S^*:\mathbb{C}\to\mathbb{V}\otimes\mathbb{V}$  acts as follows:  $1S^*=-<0|<1|+<1|<0|$ .

The proof of the theorem easily follows from the lemma:  $_{\lambda} < w|_{\otimes_{\nu}} < w|$   $(\Phi_{\lambda,\nu}exp(\phi_1h)|w>_{\lambda+1}\otimes exp(\phi_2h)|w>_{\nu+1})$  equals to LHS of 32 and  $(_{\lambda} < w|_{\otimes_{\nu}} < w|\Phi^*_{\lambda,\nu})$   $exp(\phi_1h)|w>_{\lambda+1}\otimes exp(\phi_2h)|w>_{\nu+1}$  equals to RHS of 32.

And obviously  $_{\lambda} < w | \otimes_{\nu} < w | (\Phi_{\lambda,\nu} exp(\phi_1 h) | w >_{\lambda+1} \otimes exp(\phi_2 h) | w >_{\nu+1}) = (_{\lambda} < w | \otimes_{\nu} < w | \Phi^*_{\lambda,\nu}) exp(\phi_1 h) | w >_{\lambda+1} \otimes exp(\phi_2 h) | w >_{\nu+1}.$  Q.E.D.

Theorem 2.15 (Nonlinear equations):

$$\frac{(\partial_{\phi} + \lambda + 2)}{2(\lambda + 1)} W_{\lambda}(\phi) \frac{\mu_2^R (1 - \partial_{\phi})}{\nu + 1} W_{\nu}(\phi) + \frac{\mu_1^R}{\lambda + 1} W_{\lambda}(\phi) \frac{(\partial_{\phi} + \lambda + 2)}{\nu + 1} (1 + \partial_{\phi}) W_{\nu}(\phi) = \tag{37}$$

$$= -W_{\lambda+1}(\phi) \frac{\nu + 1 - \partial_{\phi}}{2\mu_{1}^{L}} \partial_{\phi} W_{\nu+1}(\phi) + \frac{\lambda + 1 - \partial_{\phi}}{2\mu_{1}^{L}} W_{\lambda+1}(\phi) \partial_{\phi} W_{\nu+1}(\phi). \tag{38}$$

The proof is the standard procedure of deducing nonlinear equations from the bilinear ones: one should substitute  $\phi_2 = \phi_1 + \delta$  and consider the Taylor expansion of the  $W(\phi_2)$  in  $\delta$  and necessary relations are obtained as equalities in different powers  $\delta^k$ , the above one is for k = 1. In this way one obtains Kadomtsev-Petviashvili hierarchy from the Hirota bilinear relations.

Theorem 2.16 (Product formula):

$$W_{\lambda}^{\mu_L,\mu_R}(\phi)W_{\nu}^{\mu'_L,\mu'_R}(\phi) =$$
 (39)

$$=\sum_{k=0}^{\infty}C_{k}W_{\lambda+\nu-k}^{\mu_{L}+\mu'_{L},\mu_{R}+\mu'_{R}}(\phi)\sum_{i=0}^{k}\frac{(-1)^{i}\mu_{R}^{i}(\mu'_{R})^{(k-i)}}{(k-i)!i!}\binom{n}{k-i}\binom{m}{i}\sum_{i=0}^{k}\frac{(-1)^{i}\mu_{L}^{i}(\mu'_{L})^{(k-i)}}{(k-i)!i!}\binom{n}{k-i}\binom{m}{i}$$

To prove the theorem above one should note that LHS in 39 equals to  $_{\lambda} < w | \otimes_{\nu} < w | exp(\phi h) | w >_{\lambda} \otimes | w >_{\nu}$ , on the other hand one can compute this expression by use of isomorphism from proposition 2.9 and obtain RHS of 39. The calculation is very similar to the one in proof of the theorem 2.12 (Baxter relation).

So in this section we demonstrated how to deduce different relations for the Whittaker wave function of Toda by use of intertwiners.

3. 
$$SL(3)$$
 Toda Chain.

In this section we will show how to obtain raising operators and the bilinear identities for the wave functions of SL(3) Toda. The idea is completely analogous to the case of SL(2): the wave functions are matrix elements of SL(3) in irreducible representations, so considerations of the different intertwiners:  $V_{\lambda} \otimes V_{finite-dim} \to V_{\lambda\pm 1}$ ,  $V_{\lambda_1} \otimes V_{\lambda_2} \to V_{\nu_1} \otimes V_{\nu_2}$ , etc. gives different relations for the wave functions. The main technical problem here is to find the explicit expression for the image of the Whittaker vector  $|w>_{\lambda}$  through the Whittaker vector  $|w>_{\lambda\pm 1}$ : find P(f,h), such that  $\Phi(|w>_{\lambda}) = P(f,h)(|w>_{\lambda+1}$ , pay attention that we need expression of the operator P in terms of generators of the SL(3), (expression in operators  $x_{i,j}, \partial_{x_{i,j}}$  in Borel-Weil realization one can easily find). In the case of SL(2) such problem was easily solved by the fact that there was an explicit realization of the vector  $|w>_{\lambda}$ :  $|w>_{\lambda}=\sum_{i=0}\frac{\mu^n f^n}{n!(\lambda,n)}|vac>_{\lambda}$ , but in case of SL(3) such formula is not reasonable or useable. So one need to find the main formula:  $\Phi(|w>_{\lambda}) = P(f,h)(|w>_{\lambda+1}$  using more or less the only information about vector  $|w>_{\lambda}$  that it is eigenvector of the generators  $e_i$ . So it is the main difference with the case of SL(2) and we will mostly pay attention to this question and we will be brief in questions analogous to the case of SL(2).

3.1. **Notations:** The Lie algebra SL(3) is defined by the generators  $e_1, e_2, e_{12}, h_1, h_2, f_1, f_2, f_{12}$ . Sometimes we will refer  $e_{12}$  as  $e_3$ , and  $f_{12}$  as  $f_3$ . Generators satisfy the following relations:

$$[e_1, f_1] = h_1, [h_1, e_1] = 2e_1, [h_1, f_1] = -2f_1,$$
 (40)

$$[e_2, f_2] = h_2, \quad [h_2, e_2] = 2e_2, \quad [h_2, f_2] = -2f_2,$$
 (41)

$$[e_{12}, f_{12}] = h_1 + h_2, \quad [h_1 + h_2, e_{12}] = 2e_{12}, \quad [h_1 + h_2, f_{12}] = -2f_{12},$$
 (42)

$$[h_1, e_2] = -e_2, \quad [h_1, f_2] = f_2, \quad [h_2, f_1] = f_1, \quad [h_2, e_1] = e_1,$$
 (43)

$$[e_1, e_2] = e_{12}, [e_1, e_{12}] = 0, [e_2, e_{12}] = 0,$$
 (44)

$$-[f_1, f_2] = f_{12}, \quad [f_1, f_{12}] = 0, \quad [f_2, f_{12}] = 0, \tag{45}$$

$$[e_1, f_2] = 0, \quad [e_2, f_1] = 0.$$
 (46)

(47)

The first fundamental representation:

$$h_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, h_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \tag{48}$$

$$e_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, e_{12} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \tag{49}$$

$$f_1 = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, f_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, f_{12} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$
 (50)

Quadratic Casimir operator:

$$C_2 = \sum_{i=1}^{3} (e_i f_i + f_i e_i) + \frac{2}{3} \left( \sum_{i=1}^{2} h_i^2 + h_1 h_2 \right) =$$
(51)

$$=2\sum_{i=1}^{3}f_{i}e_{i}+2\sum_{i=1}^{2}h_{i}+\frac{2}{3}\left(\sum_{i=1}^{2}h_{i}^{2}+h_{1}h_{2}\right).$$

$$(52)$$

Highest weight representation:

We consider the principal (spherical) series of representations, induced by the one-dimensional representations of the Borel subgroup. Let us describe Borel-Weil realization. The space of representation  $V_{\lambda}$   $\lambda = (\lambda_1, \lambda_1)$  is functions of three real variable  $x_1, x_2, x_{12}$ , which are matrix elements of the 3\*3 upper triangular matrices, i.e. coordinates on the biggest Schubert cell of the SL(3) flag variety G/B. Matrix elements are defined by integrals with the flat measure. The action of the algebra is given by the differential operators:

$$e_{1} = \frac{\partial}{\partial x_{1}}, \quad e_{2} = \frac{\partial}{\partial x_{2}} + x_{1} \frac{\partial}{\partial x_{12}}, \quad e_{12} = \frac{\partial}{\partial x_{12}},$$

$$h_{1} = -2x_{1} \frac{\partial}{\partial x_{1}} + x_{2} \frac{\partial}{\partial x_{2}} - x_{12} \frac{\partial}{\partial x_{12}} + \lambda_{1},$$

$$h_{2} = x_{1} \frac{\partial}{\partial x_{1}} - 2x_{2} \frac{\partial}{\partial x_{2}} - x_{12} \frac{\partial}{\partial x_{12}} + \lambda_{2},$$

$$f_{1} = \lambda_{1}x_{1} - x_{1}^{2} \frac{\partial}{\partial x_{1}} - x_{1}x_{12} \frac{\partial}{\partial x_{12}} - (x_{12} - x_{1}x_{2}) \frac{\partial}{\partial x_{2}},$$

$$f_{2} = \lambda_{2}x_{2} + x_{12} \frac{\partial}{\partial x_{1}} - x_{2}^{2} \frac{\partial}{\partial x_{2}},$$

$$f_{12} = \lambda_{1}x_{12} - \lambda_{2}(x_{1}x_{2} - x_{12}) - x_{1}x_{12} \frac{\partial}{\partial x_{1}} + x_{2}(x_{1}x_{2} - x_{12}) \frac{\partial}{\partial x_{2}} - x_{12}^{2} \frac{\partial}{\partial x_{12}}.$$

$$(53)$$

Obviously that Constants are vacuum (i.e. the highest weight) vectors in such representation, i.e.  $e_i(Constant) = 0$ ,  $h_i(Constant) = \lambda_i(Constant)$ .

3.2. Whittaker function. Definition 3.1: The vector  $|w>_{\lambda}^{\mu_1,\mu_2}$  is called Whittaker vector in representation  $V_{\lambda}$ , if it is eigenvector for generators  $e_1,e_2$  i.e.  $e_1|w>_{\lambda}^{\mu_1,\mu_2}=\mu_1|w>_{\lambda}^{\mu_1,\mu_2}$  and  $e_2|w>_{\lambda}^{\mu_1,\mu_2}=\mu_2|w>_{\lambda}^{\mu_1,\mu_2}$  for some constants  $\mu_i$ . Obviously  $e_3|w>_{\lambda}^{\mu_1,\mu_2}=0$ . We will sometimes omit indexes  $\lambda,\mu$ , if they remains unchanged during the calculations.

Lemma 3.2: For given  $\mu_i$  and the irreducible representation  $V_{\lambda}$  ( $\lambda_i \neq 1, 2, 3, ...$ ) there exist a unique (up to scalar) Whittaker vector  $|w\rangle \in V_{\lambda}$ , which can be expressed as follows:

$$|w\rangle = exp(\mu_1 x_1 + \mu_2 x_2)$$
 (56)

The proof is obvious.

Remark: in case of sl(2) we also gave an expression for the Whittaker vector  $|w\rangle$  in terms of the generator f, i.e.  $|w\rangle = \sum_{i=0}^{u} \frac{\mu^n f^n}{n!(\lambda,n)} |vac\rangle$  one can see that such expression in case of sl(3) is rather complicated and unusable.

Lemma 3.3: The dual Whittaker vector:  $\langle w |$  is given by the formula:

$$\langle w| = (x_{12} - x_1 x_2)^{-(\lambda+2)} x_{12}^{-(\lambda+2)} e^{-\mu_2^L \frac{x_1}{x_{12}} - \mu_1^L \frac{x_2}{x_1 x_2 - x_{12}}}.$$
 (57)

see [7].

Definition 3.4: The Whittaker function  $W_{\lambda}^{\mu_L,\mu_R}(\phi_1,\phi_2)$  is the function given by:

$$W_{\lambda}^{\mu_L,\mu_R}(\phi_1,\phi_2) = \langle w|exp(\phi_1h_1 + \phi_2h_2)|w \rangle$$
 (58)

We will sometimes omit indexes  $\lambda, \mu$ .

Proposition 3.5: The Whittaker function satisfies the equation:

$$\left(\frac{2}{3}\left(\frac{\partial^2}{\partial\phi_1^2} + \frac{\partial^2}{\partial\phi_2^2} + \frac{\partial^2}{\partial\phi_1\partial\phi_2}\right) + \frac{\partial}{\partial\phi_1} + \frac{\partial}{\partial\phi_2} - \mu_1^L \mu_1^R e^{\phi_2 - 2\phi_1} - \mu_2^L \mu_2^R e^{\phi_1 - 2\phi_2}\right) W_{\lambda}(\phi_1, \phi_2) =$$

$$(2\lambda_1 + 2\lambda_2 + \frac{2}{3}(\lambda_1^2 + \lambda_2^2 + \lambda_1\lambda_2) W_{\lambda}(\phi_1, \phi_2). \tag{59}$$

Remark: the proof is based on idea that this hamiltonian is the second order Casimir operator for the SL(3), and completely analogous to the SL(2) case. The function  $W_{\lambda}(\phi_1, \phi_2)$  is also eigenfunction for the third order hamiltonian, which is the image of the third order Casimir for the SL(3). So we see that Whittaker function is the wave function for the SL(3) quantum Toda chain. These facts are standard so we are very brief.

Proposition 3.6 (Integral representation of the Whittaker function) ([7]):

$$W_{\lambda}(\phi_1, \phi_2) = e^{(\lambda_1 + 1)\phi_1 + (\lambda_2 + 1)\phi_2} \int dx_1 dx_2 dx_{12} (x_{12} - x_1 x_2)^{-\lambda_2 - 2} x_{12}^{-\lambda_1 - 2}) \times$$
(60)

$$\times e^{-\mu_2^L \frac{x_1}{x_{12}} - \mu_1^L \frac{x_2}{x_1 x_2 - x_{12}} - \mu_1^R x_1 e^{\phi_2 - 2\phi_1} - \mu_2^R x_2 e^{\phi_1 - 2\phi_2}}.$$
(61)

The theorem is simple corollary of the facts that the invariant pairing  $\langle v|u \rangle$  is realized as integral  $\int$  with the flat measure, and dual Whittaker vector  $\langle w|$  is realized as  $(x_{12}-x_1x_2)^{(-\lambda_2-2)}x_{12}^{(-\lambda_1-2)}exp(-\mu_2^L\frac{x_1}{x_{12}}-\mu_1^L\frac{x_2}{x_1x_2-x_{12}})$ .

3.3. **Intertwining operators.** Denote by  $V_{(1,0)}$  the first fundamental representation of the  $SL(3,\mathbb{R})$ , denote by |0> the highest weight vector, by |1> the vector  $f_1|0>$ , by |2> the vector  $f_2f_1|0>=f_3|0>$ , i.e.:

$$|0\rangle = \begin{pmatrix} 1\\0\\0 \end{pmatrix}, |1\rangle = \begin{pmatrix} 0\\1\\0 \end{pmatrix}, |2\rangle = \begin{pmatrix} 0\\0\\1 \end{pmatrix}, \tag{62}$$

hence  $e_1|0>=0, e_2|0>=0, h_1|0>=|0>, h_2|0>=0.$ 

Proposition 3.7: Let us denote by  $\Phi_{\lambda} = \Phi_{\lambda,(+1,0)} \oplus \Phi_{\lambda,(-1,1)} \oplus \Phi_{\lambda,(0,-1)}$  the ismorphism  $V_{\lambda+(1,0)} \oplus V_{\lambda+(-1,1)} \oplus V_{\lambda+(0,-1)} \to V_{\lambda} \otimes V_1$ . The operators  $\Phi_{\lambda,(+1,0)}$  and  $\Phi_{\lambda,(+1,0)}^{-1}$  in Borel-Weil realization are given by the formulas:

$$\Phi_{\lambda,(+1,0)}: x_1^{n_1} x_2^{n_2} x_3^{n_3} | 0 >_{\lambda} \otimes | 0 > \to x_1^{n_1} x_2^{n_2} x_3^{n_3} | 0 >_{\lambda+(1,0)}, \tag{63}$$

$$\Phi_{\lambda,(+1,0)}: x_1^{n_1} x_2^{n_2} x_3^{n_3} | 0 >_{\lambda} \otimes | 1 > \to x_1^{n_1+1} x_2^{n_2} x_3^{n_3} | 0 >_{\lambda+(1,0)}, \tag{64}$$

$$\Phi_{\lambda,(+1,0)}: x_1^{n_1} x_2^{n_2} x_3^{n_3} |0>_{\lambda} \otimes |2> \to x_1^{n_1} x_2^{n_2} x_3^{n_3+1} |0>_{\lambda+(1,0)}.$$

$$\tag{65}$$

$$(\Phi_{\lambda,(+1,0)})^{-1}: \\ x_1^{n_1}x_2^{n_2}x_3^{n_3}|0>_{\lambda+(1,0)} \to \frac{1}{(\lambda_1+1)(\lambda_1+\lambda_2+2)}((n_1n_2x_3+((\lambda_1+2)n_3+n_2n_3)x_1x_2)x_1^{n_1-1}x_2^{n_2-1}x_3^{n_3-1}|0>_{\lambda}\otimes|2> + \\ (n_1(\lambda_1+\lambda_2+2-n_2)x_3+n_3(\lambda_2-n_2)n_3x_2)x_1^{n_1-1}x_2^{n_2}x_3^{n_3-1})|0>_{\lambda}\otimes|1> + \\ (-n_1n_2x_3^2+n_3(-\lambda_2+n_2)x_1^2x_2^2+ \\ +((\lambda_1+1)(\lambda_1+\lambda_2+2)-n_1(\lambda_1+\lambda_2+2)+n_1n_2-n_3(\lambda_1+2)-n_2n_3)x_1x_2x_3)x_1^{n_1-1}x_2n_2-1x_3^{n_3-1}|0>_{\lambda}\otimes|0> 0$$

This proposition is auxiliary for us. The main formulas we need to obtain the relations between the Whittaker wave functions are the formulas from the next proposition. But to obtain them we need this statement. Actually we use it rather small, only to fix the constants (see below).

To prove the formula 65 one should only note that the representation  $V_{(1,0)}$  also have Borel-Weil realization and the above intertwiner is given by the multiplication of functions from the two representations. The proof of the formula 66 is more complicated, but it is standard representation theory reasoning, and should be available in the literature, but we were not able to find it. So we sketch the proof. Let us denote the three components of the operator  $(\Phi_{\lambda,(+1,0)})^{-1}$  by  $\Psi_0, \Psi_1, \Psi_2$ , i.e.  $(\Phi_{\lambda,(+1,0)})^{-1}|v>=\Psi_0(|v>)\otimes |0>+\Psi_1(|v>)\otimes |1>+\Psi_2(|v>)\otimes |2>$ , first simple, but useful step in proving 66 is that all the three operators  $\Psi_i$  can be easily expressed one throw another, so it's enough to find only one of them. Actually it is easy to see that, for example:  $\Psi_1=[\Psi_2,f_2], \Psi_0=[\Psi_2,f_3]$ . Looking at the formula 66 it is clear that operator  $\Psi_2$  is much simpler than the other two, so it is rather useful that we can express them through the operator  $\Psi_2$ . Second step is to find  $\Psi_2$ . It goes as follows, first let us note that:  $\Psi_2(x_1^{k_1}x_2^{k_2}x_3^{k_3})=\alpha_{k_1,k_2,k_3}x_1^{k_1-1}x_2^{k_2-1}x_3^{k_3}+\beta_{k_1,k_2,k_3}x_1^{k_1}x_2^{k_2}x_3^{k_3-1}$ . That easily follows from the equalities:  $\Psi_2h_1=h_1\Psi_2,\Psi_2h_2=h_1\Psi_2$  and  $e_1^{k_1+1}\Psi_2(x_1^{k_1}x_2^{k_2}x_3^{k_3})=0, e_3^{k_1+1}\Psi_2(x_1^{k_1}x_2^{k_2}x_3^{k_3})=0$ . the second point is to use the commutation relation of  $e_2$  and  $\Psi_2$  and to find recurrence relations for  $\alpha_{k_1,k_2,k_3},\beta_{k_1,k_2,k_3}$ , they can be solved directly after some work. Hence we find  $\Psi_2$  and as explained above it follows that we find  $\Phi_1,\Phi_0$ .

Remark: Some other useful formulas for the intertwiners we put to the Appendix. Let us note that it is rather easy to find the formulas for the intertwiners in Verma realization (see Appendix), not Borel-Weil, but the problem is that expression of the vector  $x_1^{k_1}x_2^{k_2}x_3^{k_3}$  through the vectors  $f_1^{n_1}f_2^{n_2}f_3^{n_3}$  is rather complicated and untreatable.

Proposition 3.8: The action of the intertwiner  $(\Phi_{\lambda,(+1,0)})$  on the Whittaker vector can be expressed as follows:

$$(\Phi_{\lambda,(+1,0)})|w>_{\lambda}^{\mu_1,\mu_2}\otimes|0>\to|w>_{\lambda+(1,0)}^{\mu_1,\mu_2}=exp(\mu_1x_1+\mu_2x_2)|0>_{\lambda+(1,0)},$$
(67)

$$(\Phi_{\lambda,(+1,0)})|w>_{\lambda}^{\mu_1,\mu_2}\otimes|1> \to \bar{\Lambda}_1|w>_{\lambda+(1,0)}^{\mu_1,\mu_2} = x_1 exp(\mu_1 x_1 + \mu_2 x_2)|0>_{\lambda+(1,0)},$$
(68)

$$(\Phi_{\lambda,(+1,0)})|w>_{\lambda}^{\mu_1,\mu_2}\otimes|2> \to \frac{1}{\mu_1}(f_2+\bar{\Lambda}_2\mu_2-\lambda_2\bar{\Lambda}_2|w>_{\lambda+(1,0)}^{\mu_1,\mu_2}=x_{12}exp(\mu_1x_1+\mu_2x_2)|0>_{\lambda+(1,0)}.$$
(69)

where 
$$\bar{\Lambda}_1 = \frac{2(\lambda_1+1)+\lambda_2-2h_1-h_2}{3\mu_1}$$
 and  $\bar{\Lambda}_2 = \frac{(\lambda_1+1)+2\lambda_2-h_1-2h_1}{3\mu_2}$ 

where  $\bar{\Lambda}_1 = \frac{2(\lambda_1+1)+\lambda_2-2h_1-h_2}{3\mu_1}$  and  $\bar{\Lambda}_2 = \frac{(\lambda_1+1)+2\lambda_2-h_1-2h_2}{3\mu_2}$ The action of the intertwiner  $(\Phi_{\lambda,(+1,0)})^{-1}$  on the Whittaker vector can be expressed as follows:

$$(\Phi_{\lambda,(+1,0)})^{-1}|w>_{\lambda+1}^{\mu_1,\mu_2} \to \frac{1}{(\lambda_1+1)(\lambda_1+\lambda_2+2)}(\mu_1\mu_2|w>_{\lambda}^{\mu_1,\mu_2}\otimes|2> + \frac{1}{\mu_2}(\tilde{\Lambda}_2+2)|w>_{\lambda}^{\mu_1,\mu_2}\otimes|1> + \frac{1}{\mu_2}(\tilde{\Lambda}_1^2-(\lambda_2-3)\tilde{\Lambda}_1))|w>_{\lambda}^{\mu_1,\mu_2}\otimes|0>).$$
 (70)

where  $\tilde{\Lambda}_1 = \frac{2h_1 + h_2 + \lambda_1 + 2\lambda_2}{3}$  and  $\tilde{\Lambda}_2 = \frac{2h_2 + h_1 + \lambda_2 + 2\lambda_1}{3}$ This proposition is crucial in this section. The relations for the Whittaker wave functions (71,72) easily follows from it. Let us sketch the proof and explain the difference with SL(2) case, which is much more simple. We discuss the analogous reasonings with more details in the next section in the case of SL(n). Let us denote by  $P_0, P_1, P_2$  the operators such that:  $(Phi_{\lambda,(+1,0)})^{-1}|w>_{\lambda+1}=\Psi_0|w>_{\lambda}\otimes|0>+\Psi_1|w>_{\lambda}\otimes|1>+\Psi_2|w>_{\lambda}\otimes|2>$ . We will look for them as some polynomials  $P_i(h_i, f_i)$  from the generators  $e_i$ . From the commutation relations with the operators  $e_i$ and uniqueness of the Whittaker vector it's easy to see that the operator  $P_2$  is some constant depending only on  $\mu_i$ . To fix this constant we look on the formulas 66, 56 and see that  $P_2 = \mu_1 \mu_2$ . Also from the commutation relations with operators  $e_i$  follows the following relations:  $[P_1, e_1] = 0$ ,  $[P_1, e_2] = e_2 + Constant$  so  $A_1$  equals  $\frac{1}{\mu_2}\Lambda_2 + C_1$ , where  $\Lambda_i$ are fundamental coroots, i.e. elements of Cartan subalgebra such that  $[\Lambda_i, e_j] = \delta_{i,j}$ , the constant  $C_1$  can also be fixed by the formulas 66,56. Analogously  $A_0$  satisfy the identities:  $[P_0, e_2] = 0$ ,  $[P_0, e_1] = \frac{\alpha}{\mu_1} A_1 e_1 + \beta P_1$ , where  $\alpha + \beta = 1$ . the solution of this equation is given by the ansatz  $P_0 = \frac{1}{\mu_2} (f_1 + \frac{1}{\mu_1} (\Lambda_1^2 + C_0 \Lambda_1))$  the constants  $c_0, \alpha, \beta$  also can be fixed by the formulas 66,56. So the necessary formulas for  $P_i$  are obtained.

Let us one more time emphasize the difference with the SL(2) case, where we had the formula for the Whittaker vector  $|w\rangle = \sum_{i=0}^{n} \frac{\mu^n f^n}{n!(\lambda,n)} |vac\rangle$ . Hence the proposition 2.8, which is analogous to the proposition 3.8 above immediately ately follows from the formula for the action of the intertwiner on the vector  $f^n|vac>$ . In the present case the formula for |w> in terms of the generators  $f_i$  is too complicated to extract something from it, so the proof of the proposition above was based mostly on the property of the Whittaker vector:  $e_i|w>=\mu_i|w>$ .

## 3.4. Relations for the Whittaker wave functions.

Theorem (Raising operators) 3.9: The following relations holds:

$$W_{\lambda+1}(\phi_1, \phi_2) = \frac{exp(\phi_1)}{\mu_1^R(\lambda_1 + 1)(\lambda_1 + \lambda_2 + 2)} (\mu_1^R exp(-2\phi_1 + \phi_2) + \frac{1}{\mu_2^R} (\left(\frac{2\partial_{\phi_1} + \partial_{\phi_2} + \lambda_1 + 2\lambda_2}{3}\right)^2 - (\lambda_2 - 3) \frac{2\partial_{\phi_1} + \partial_{\phi_2} + \lambda_1 + 2\lambda_2}{3} - 2(\lambda_2 - 1)) W_{\lambda}(\phi_1, \phi_2).$$

$$(71)$$

This proposition follows from the 67.70 in a way completely analogous to the SL(2) case (proposition 2.11), so we omit the details.

Theorem (Bilinear relations) 3.10:

$$((\frac{1}{\mu_{2}^{\prime L}}(\mu_{2}^{\prime R}exp(-\phi_{2}+2\phi_{1})+(\frac{-\partial_{\phi_{2}^{\prime}}-2\partial_{\phi_{1}^{\prime}}+(\nu_{2}+1)+2\nu_{1}}{3\mu_{1}^{\prime L}})^{2} - (\nu_{1})\frac{-\partial_{\phi_{2}^{\prime}}-2\partial_{\phi_{1}^{\prime}}+(\nu_{2}+1)+2\nu_{1}}{3\mu_{1}^{\prime L}}) - \frac{-2\partial_{\phi_{1}}-\partial_{\phi_{2}}+2(\lambda_{1}+1)+\lambda_{2}}{3\mu_{1}^{\prime L}} \frac{-2\partial_{\phi_{2}^{\prime}}-\partial_{\phi_{1}^{\prime}}+2(\nu_{2}+1)+\nu_{1}}{3\mu_{2}^{\prime L}} + (\frac{1}{\mu_{1}^{\prime L}}(\mu_{1}^{R}exp(-\phi_{1}+2\phi_{2})+(\frac{-\partial_{\phi_{1}}-2\partial_{\phi_{2}}+(\nu_{1}+1)+2\nu_{2}}{3\mu_{2}^{\prime L}}))^{2} - (\nu_{2})\frac{-\partial_{\phi_{1}}-2\partial_{\phi_{2}}+(\nu_{1}+1)+2\nu_{2}}{3\mu_{2}^{\prime L}}))W_{\lambda+1}^{\mu_{1}^{\prime L},\mu_{2}^{\prime L},\mu_{1}^{\prime R},\mu_{2}^{\prime R}}(\phi_{1},\phi_{2})W_{\nu+1}^{\mu_{1}^{\prime L},\mu_{2}^{\prime L},\mu_{1}^{\prime R},\mu_{2}^{\prime R}}(\phi_{1}^{\prime},\phi_{2}^{\prime}) = \frac{1}{(\lambda_{1}+1)(\lambda_{1}+\lambda_{2}+2)}\frac{1}{(\nu_{2}+1)(\nu_{1}+\nu_{2}+2)}((\frac{1}{\mu_{1}^{\prime R}}(\mu_{1}^{\prime L}exp(-2\phi_{1}+\phi_{2})+\frac{1}{\mu_{2}^{\prime R}}((\frac{2\partial_{\phi_{1}}+\partial_{\phi_{2}}+\lambda_{1}+2\lambda_{2}}{3})^{2} - (\lambda_{2}-3)\frac{2\partial_{\phi_{1}}+\partial_{\phi_{2}}+\lambda_{1}+2\lambda_{2}}{3} - 2(\lambda_{2}-1))))\mu_{1}^{\prime R}\mu_{2}^{\prime R} + \frac{(\partial_{\phi_{1}}+2\partial_{\phi_{2}}+2\lambda_{1}+\lambda_{2})}{3\mu_{2}}\frac{(\partial_{\phi_{1}^{\prime}}+2\partial_{\phi_{1}}+2\nu_{1}+\nu_{1})}{3\mu_{1}} - -\mu_{1}\mu_{2}(\frac{1}{\mu_{2}^{\prime R}}(\mu_{2}^{\prime L}exp(-2\phi_{2}^{\prime}+\phi_{1}^{\prime})+\frac{1}{\mu_{1}^{\prime R}}((\frac{2\partial_{\phi_{2}^{\prime}}+\partial_{\phi_{1}^{\prime}}+\nu_{2}+2\nu_{1}}{3})^{2} - (\nu_{1}-3)\frac{2\partial_{\phi_{2}^{\prime}}+\partial_{\phi_{1}^{\prime}}+\nu_{2}+2\nu_{1}}{3} - 2(\nu_{1}-1))))W_{\lambda}^{\mu_{1}^{\prime L},\mu_{2}^{\prime L},\mu_{1}^{\prime R},\mu_{2}^{\prime R}}(\phi_{1},\phi_{2})W_{\nu}^{\mu_{1}^{\prime L},\mu_{2}^{\prime L},\mu_{1}^{\prime R},\mu_{2}^{\prime R}}(\phi_{1}^{\prime},\phi_{2}^{\prime}).$$
 (72)

The prooof is also completely analogous to the SL(2) case, so we refer the reader to the previous section.

One can obtain the nonlinear equations from this bilinear one, as it was done in the SL(2) case, but we omit these considerations due to their length.

4. 
$$SL(n)$$
 Toda Chain.

4.1. Notations. Algebra sl(n) is completely given by the generators  $e_i, h_i, f_i, i = 1, ..., n-1$  and commutation relations:

$$[e_i, h_j] = -A_{ij}e_{\pm,i}, \quad [f_i, h_j] = A_{ij}e_{\pm,i}, \quad [e_i, f_j] = \delta_{ij}h_j, \quad i, j = 1, \dots, N - 1,$$
(73)

and the Serre relations

$$\operatorname{ad}_{T_{+i}}^{1-A_{ij}}(T_{\pm j}) = 0, (74)$$

where  $ad_x^k(y) \equiv [x, [x, ..., [x, y]..]]$ 

Where  $A_{ij}$  is Cartan matrix, which is for the algebra SL(N) is equal to

$$\begin{pmatrix} 2 & -1 & 0 & 0 & \dots & 0 \\ -1 & 2 & -1 & 0 & \dots & 0 \\ 0 & -1 & 2 & -1 & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}$$

Quadratic Casimir operator is

$$C_2 = \sum_{\alpha \in \Delta} e_{\alpha} f_{\alpha} + \sum_{ij}^{N-1} A_{ij}^{-1} h_i h_j,$$
 (75)

where the first sum goes over all (positive and negative) roots.

In the case of generic SL(N) group, one can define the (right) regular representation only in general terms of the group acting on the space of the algebra of functions:

$$\pi_{reg}(h)f(g) = f(gh). \tag{76}$$

Therefore, we use from now on mostly group (not algebra) terms. Still, we can restrict the space of functions to the irreducible representations in the generic situation. For doing this, we consider the representation denoted by  $V_{\lambda}$ induced by one-dimensional representations of the Borel subgroup. That is, we reduce the space of all functions to the functions satisfying the following covariance property:

$$f_{\lambda}(bg) = \chi_{\lambda}(b)f_{\lambda}(g), \tag{77}$$

where b is an element of the Borel subgroup of lower-triangle matrices and  $\chi_{\lambda}$  is the character of the Borel subgroup of the form:

$$\chi_{\lambda}(b) = \prod_{i=1}^{N-1} |b_{ii}|^{(\lambda-\rho)e_{-i}} (\operatorname{sign}b_{ii})^{\epsilon_i}, \tag{78}$$

where  $\epsilon_i$  are equal to either 0 or 1. For the sake of simplicity, we consider the representations with all these sign factors to be zero although other cases can be also easily treated. The representation constructed belongs to the principal (spherical) series.

Thus, our representation is given by restricting the space of functions to the functions defined on the coset  $B \setminus G$ , the biggest cell of which, in turn, may be identified with the strictly upper-triangular matrices  $N_+$ . We will denote by  $x_{i,j}$  the matrix elements of n\*n matrices, and discussion above gives us realization of the representation of the SL(n) in the space of functions of  $x_{i,j}$ ,  $1 \le i < j \le n$ . The explicit formulas can be found for example in [23, 24].

4.2. Whittaker function. Definition 4.1: The vector  $|w\rangle_{\lambda}^{\mu_1,...,\mu_{n-1}}$  is called Whittaker vector in representation  $V_{\lambda}$ , if it is eigenvector for generators  $e_1,...,e_{n-1}$  i.e.  $e_i|w\rangle_{\lambda}=\mu_i|w\rangle_{\lambda}$  for some constants  $\mu_i$ . Obviously  $[e_i,e_j]|w\rangle_{\lambda}^{\mu_1,\mu_2}=0$ . We will sometimes omit indexes  $\lambda,\mu$ , if they remains unchanged during the calculations.

Lemma 4.2: For given  $\mu_i$  and irreducible representation  $V_{\lambda}$  there exist a unique (up to scalar) Whittaker vector  $|w\rangle \in V_{\lambda}$ , which can be expressed as follows:

$$|w\rangle = exp(\mu_1 x_{1,2} + \mu_2 x_{2,3} + \dots + \mu_{n-1} x_{n-1,n})$$
(79)

The proof is obvious due to explicit formulas for generators  $e_i$  in Borel-Weil realization.  $e_i$  acts on such vector as  $\partial_{x_{i,i+1}}$ , see [7] for details.

Definition 4.3: The Whittaker function  $W_{\lambda}^{\mu_{i}^{L},\mu_{j}^{R}}(\phi_{1},\phi_{2},...,\phi_{n-1})$  is the function given by:

$$W_{\lambda}^{\mu_i^L, \mu_j^R}(\phi_1, \phi_2) = \langle w | exp(\phi_1 h_1 + \phi_2 h_2 + \dots + \phi_{n-1} h_{n-1}) | w \rangle$$
(80)

We will sometimes omit indexes  $\lambda, \mu$ .

Proposition 4.4: The Whittaker function satisfies the equation:

$$\left(\sum_{ij} A_{ij}^{-1} \partial_{\phi_i} \partial_{\phi_j} + 2\sum_{ij} A_{ij}^{-1} \partial_{\phi_j} - 2\sum_{i} \mu_i^L \mu_i^R exp(\sum_j A_{i,j} \phi_j)\right) W_{\lambda}^{\mu_i^L, \mu_j^R}(\phi_i) = (\lambda^2 - \rho^2) W_{\lambda}^{\mu_i^L, \mu_j^R}(\phi_i). \tag{81}$$

where  $\rho$  is one half of sum of simple roots.

Remark: the proof is based on the idea, that this hamiltonian is the second order Casimir operator for the SL(n), and completely analogous to the SL(2) case. The function  $W_{\lambda}(\phi_i)$  is also eigenfunction for the higher order hamiltonians, which are the images of the higher order Casimirs for the SL(n). Hence, we see that Toda is completely integrable system and the Whittaker function is the wave function for all Hamiltonians of the SL(n) quantum Toda chain. These facts are standard so we are very brief (see [7]).

Proposition 4.5 (Integral representation of the Whittaker function) ([7]):

$$W_{\lambda}(\phi_{i}) = e^{-\sum_{i} \lambda_{i} \phi_{i}} \int_{X=B \setminus G} \prod_{i < j} dx_{ij} \prod_{i=1}^{N-1} \Delta_{i}^{-(\sum_{k} A_{k,i} \lambda_{k} + 1)} (xS^{-1}) \times e^{\mu_{i}^{R} x_{i,i+1} e^{(\sum_{k} A_{k,i} \lambda_{k})} - \mu_{N-i}^{L} \frac{\Delta_{i,i+1}(xS^{-1})}{\Delta_{i}(xS^{-1})}}.$$
 (82)

where  $\Delta_i$  - is i-th principal minor of matrix  $x_{i,j}$ ,  $\Delta_{i,i+1}$  - determinant (n-1)\*(n-1) submatrix, which obtained by interchanging n-1 and n column. Matrix S is given by  $S_{i,j}=\delta_{i+j,n}$ .

The theorem is corollary of the facts that the invariant pairing  $\langle v|u\rangle$  is realized as integral with the flat measure, and the realizations of the Whittaker and the dual Whittaker vectors. The most nontrivial part is to find dual Whittaker vector. We refer to [7] for detailed exposition.

4.3. **Intertwining operators.** Denote by  $V_{(1,0,\ldots,0)}$  the first fundamental representation of the  $SL(n,\mathbb{R})$ , which is just the standard action of the n\*n matrices on  $\mathbb{C}^n$ , denote by |0> the highest weight vector, by |1> the vector  $f_1|0>$ , by |2> the vector  $f_2f_1|0>=f_3|0>$ , etc.

To construct relations for wave functions, one needs to express the action of the intertwiner on Whittaker vector, in terms of the generators  $f_i, h_i$ . For example the crucial formula for us in the case of SL(3) was:

$$|w>_{\lambda+1}^{\mu_1,mu_2} \to \frac{1}{(\lambda_1+1)(\lambda_1+\lambda_2+2)} (\mu_1\mu_2|w>_{\lambda}^{\mu_1,mu_2} \otimes |2> +$$
 (83)

$$\frac{1}{\mu_2}(\tilde{\Lambda}_2 + 2)|w>_{\lambda}^{\mu_1, mu_2} \otimes |1> + \frac{1}{\mu_2}(f_1 + \frac{1}{\mu_2}(\tilde{\Lambda}_1^2 - (\lambda_2 - 3)\tilde{\Lambda}_1))|w>_{\lambda}^{\mu_1, mu_2} \otimes |0>). \tag{84}$$

Where  $\tilde{\Lambda}_i$  were fundamental coweights minus some constants.

We need analogous formulas for the case of SL(n). We are unable to find them. We will present only the equations solutions to which will give necessary formulas and first steps towards their solution.

Let us consider the most simple intertwiner  $\Phi: V_{\lambda+(1,0,\ldots,0)} \to V_{\lambda} \otimes V_{(1,0,\ldots,0)}$ . We need to find such polynomials  $P_j(f_i,h_i), 0 \leq j < n$  such that:

$$\Phi|w>_{\lambda+(1,0,\dots,0)} = \sum_{j=0}^{n-1} P_j(f_i, h_i)|w>_{\lambda} \otimes |j>$$
(85)

Proposition 4.6 The following relations holds:

$$e_k P_i(f_i, h_i)|w\rangle_{\lambda} + \delta_{i,k-1} P_k(f_i, h_i)|w\rangle_{\lambda} = \mu_k P_i(f_i, h_i)|w\rangle_{\lambda}$$
(86)

The proof is obvious.

Corollary 4.7: One can look for  $P_i(f_i, h_i)$  as an elements of the U(sl(n)), satisfying the following relations:

$$[e_k, P_j(f_i, h_i)] = 0, \text{ for } k \neq j+2$$
 (87)

$$[e_k, P_{k-2}(f_i, h_i)] = \alpha_k P_{k-1}(f_i, h_i) e_k + \beta_k P_{k-1}(f_i, h_i), \text{ where } \alpha_k + \beta_k = 1.$$
(88)

This allows one to organize itterative search of polynomials  $P_j$ : if  $P_j$  already found then  $P_{j-1}$  is defined by conditions:  $[e_k, P_{j-1}(f_i, h_i)] = 0$  for  $k \neq j+1$  and  $[e_{j+1}, P_{j-1}(f_i, h_i)] = \alpha_k P_{k-1}(f_i, h_i)e_k + \beta_k P_{k-1}(f_i, h_i)$ .

We can conclude from the considered examples that the highest degree part of the  $P_{j-1}$  can be defined uniquely from this relations. The problem is that we do not know the way of solving such reccurence relations. Let us present the first 3 polynomials  $P_i$  found by hands.

Proposition 4.8: Polynomials  $P_{n-1}, P_{n-2}, P_{n-3}$  are given by the formulas:

$$P_{n-1} = \mu_1 \mu_2 \dots \mu_{n-1}, \quad P_{n-2} = (\Lambda_{n-1} + Constant) \mu_1 \mu_2 \dots \mu_{n-2}, \tag{89}$$

$$P_{n-3} = (\mu_{n-2}f_{n-2} + \mu_{n-3}f_{n-3} + \dots + \mu_1f_1 + \dots + \mu_nf_n + \dots + \mu_nf_n$$

$$+(\Lambda_{n-2}^2 - \Lambda_{n-2}\Lambda_{n-3}) + (\Lambda_{n-3}^2 - \Lambda_{n-3}\Lambda_{n-4}) + \dots + (\Lambda_1)^2)\mu_1\mu_2\dots\mu_{n-3} + F_1(\Lambda_i). \tag{90}$$

where  $\Lambda_i$  - fundamental coweights,  $F_1$  - some polynomial of degree 1.

The proof is straightforward check that given polynomials satisfy relations (87) and one can see that they are the only with this property.

So we see polynomials  $P_{n-1-j}$  are of degree j. It's easy to guess that it also holds true for j > 3. It's easy to understand that to write the simplest raising relation one should find  $P_0$ , which should be some polynomial of degree n-1. Our first hope to do that was to find some recurrence formula:  $P_{j-1} = A_j P_j$ , but we did not succeed, looking at 90 it seems that such formula may not exist. Another idea is that may be  $P_j$  are somehow related to the Casimirs of subalgebras SL(n-j), which one guess looking at 90 and to commutation relation:  $[e_k, P_j] = 0$  for  $k \neq j+2$ .

Let us also note that it is possible to write intertwiner  $V_{\lambda+(1,0,...,0)} \to V_{\lambda} \otimes V_{(1,0,...,0)}$  in Borel-Weil realization so it is possible to find the expression for  $P_j$  as some polynomial of operators of multiplication on monomials  $x_{i,j}$  but the problem is then express such operators via the polynomial of the  $f_i, h_i$ .

### 5. Concluding remarks.

Let us formulate the main points of our work. First, we recalled the representation theory approach to integrable systems, i.e. we recalled that appropriate matrix elements in irreducible representations are the wave functions of integrable systems, Casimirs of the group are the Hamiltonians, the integral representation can be obtained by realizing the representation of the group in space of functions, where the pairing is given by the integral. Second, we demonstrated in our paper that different relations between the wave functions, can be obtained from considerations of intertwining operators between above representations. We saw that raising, bilinear etc. relations can be obtained this way.

In this paper we did not succeed in obtaining the general formula for the case of SL(n), it seems such formula can be quite complicated. We only found some recurrence relations, whose solution leads to desired formula. It seems that one should guess some ansatz for the desired polynomial, something like determinant or trace of some matrix of the generators of the algebra, like the formula for Casimirs.

Group theory approach to integrable systems is very powerful, it allows to give transparent explanations for different properties of the system. So let us mention some questions about integrable systems which should be interpreted by the group theory methods. It will be interesting to understand the relation between our approach to raising relations and approach based on Hecke algebra (see [18]). It is well-known that Bessel functions, which are very close to the SL(2) Toda wave functions, satisfy the following properties:

$$\sum_{n=-\infty}^{\infty} J_n(x)t^n = exp(\frac{1}{2}x(t-\frac{1}{t})) \text{ and hence } J_{-n}(x) = (-1)^n J_n(x).$$

It will be interesting to find analogues formulas for the wave functions of SL(n) Toda chain and find their group theory interpretation. In the studding integrable systems recently appeared such new important ideas as shift operator and Dunkl operator [25], another new important concept is Cherednik-Matsuo correspondence (see Cherednik in [2])

between solutions of integrable systems and solutions of Knizhnik-Zamolodchikov equation it seems to be interesting to understand their group theory meaning.

Let us note the most straight-forward generalization of relations obtained here is their transfer to the other groups: instead of SL(2, R) we plan to consider the affine version  $\hat{SL}(2)$  related to periodic Toda chain, quantum analogue  $U_q(SL(2))$  related to Mcdonald polynomials and q-special functions, and may be finite field analog  $SL(2, F_q)$ , which may be related to the questions considered in [26].

### 6. Acknowledgments.

The author is especially grateful to S. Kharchev, for excellent explanations and discussions, and sharing with the author his ideas, which led to writing this paper. We are also grateful to A. Mironov for the significant help during working on text, to A. Morozov for initiating our work in this direction, and to our scientific advisor S. Khoroshkin for constant support. The work was supported by INTAS -93-10183 Ext., Russian President's grant 96-15-96939 and RFBR-98-0100344.

Appendix.

In this Appendix we collect some more formulas on intertwining operators for SL(3). We use notations of section 3.

Proposition: The representation  $V_{\lambda} \otimes V_1$  is isomorphic to the:  $V_{\lambda+(1,0)} \oplus V_{\lambda+(-1,1)} \oplus V_{\lambda+(0,-1)}$ . The highest weight vectors in  $V_{\lambda} \otimes V_1$  are given by the formulas:

$$|0\rangle_{\lambda} \otimes |0\rangle$$
 - weight  $\lambda + (1,0)$  (91)

$$f_1|0>_{\lambda}\otimes|0>-\lambda_1|0>_{\lambda}\otimes|1> - \text{ weight }\lambda+(-1,1)$$
 (92)

$$\lambda_2(\lambda_1 + \lambda_2 + 1)|0>_{\lambda} \otimes |2> -(\lambda_1 + \lambda_2 + 1)f_2|0>_{\lambda} \otimes |1> +(f_1f_2 + \lambda_2f_3)|0>_{\lambda} \otimes |0> - \text{ weight } \lambda + (0, -1)$$
 (93)

The intertwiner  $\Phi_{\lambda} = \Phi_{\lambda,(+1,0)} \oplus \Phi_{\lambda,(-1,1)} \oplus \Phi_{\lambda,(0,-1)} : V_{\lambda+(1,0)} \oplus V_{\lambda+(-1,1)} \oplus V_{\lambda+(0,-1)} \to V_{\lambda} \otimes V_1$  in "Verma" basis is given by the formulas:

$$\begin{split} &\Phi_{\lambda,(+1,0)}: f_{1}^{n_{1}}f_{2}^{n_{2}}f_{3}^{n_{3}}|0>_{\lambda+(1,0)}\rightarrow f_{1}^{n_{1}}f_{2}^{n_{2}}f_{3}^{n_{3}}|0>_{\lambda}\otimes|0> +n_{1}f_{1}^{n_{1}-1}f_{2}^{n_{2}}f_{3}^{n_{3}}|0>_{\lambda}\otimes|1> +n_{3}f_{1}^{n_{1}}f_{2}^{n_{2}}f_{3}^{n_{3}-1}|0>_{\lambda}\otimes|2>, \quad (94) \\ &\Phi_{\lambda,(-1,1)}: f_{1}^{n_{1}}f_{2}^{n_{2}}f_{3}^{n_{3}}|0>_{\lambda+(-1,1)}\rightarrow (f_{1}^{n_{1}+1}f_{2}^{n_{2}}f_{3}^{n_{3}}-n_{2}f_{1}^{n_{1}}f_{2}^{n_{2}-1}f_{3}^{n_{3}+1})|0>_{\lambda}\otimes|0> +\\ &((-\lambda_{1}+n_{1})f_{1}^{n_{1}}f_{2}^{n_{2}}f_{3}^{n_{3}}-n_{1}n_{2}f_{1}^{n_{1}-1}f_{2}^{n_{2}-1}f_{3}^{n_{3}+1})|0>_{\lambda}\otimes|1> +(n_{2}(\lambda_{2}-n_{3})f_{1}^{n_{1}}f_{2}^{n_{2}-1}f_{3}^{n_{3}}+n_{3}f_{1}^{n_{1}+1}f_{2}^{n_{2}}f_{3}^{n_{3}-1})|0>_{\lambda}\otimes|2\%\rangle \\ &\Phi_{\lambda,(0,-1)}: f_{1}^{n_{1}}f_{2}^{n_{2}}f_{3}^{n_{3}}|0>_{\lambda+(0,-1)}\rightarrow (f_{1}^{n_{1}+1}f_{2}^{n_{2}+1}f_{3}^{n_{3}}+(\lambda_{2}-n_{2})f_{1}^{n_{1}}f_{2}^{n_{2}}f_{3}^{n_{3}+1})|0>_{\lambda}\otimes|0> +\\ &(n_{1}(\lambda_{2}-n_{2})f_{1}^{n_{1}-1}f_{2}^{n_{2}}f_{3}^{n_{3}+1}+(n_{1}-(\lambda_{1}+\lambda_{2}+1))f_{1}^{n_{1}}f_{2}^{n_{2}+1}f_{3}^{n_{3}})|0>_{\lambda}\otimes|1> +\\ &((\lambda_{2}-n_{2})(n_{3}-(\lambda_{1}+\lambda_{2}+1))f_{1}^{n_{1}}f_{2}^{n_{2}}f_{3}^{n_{3}}+n_{3}f_{1}^{n_{1}+1}f_{2}^{n_{2}+1}f_{3}^{n_{3}-1})|0>_{\lambda}\otimes|2>. \quad (96) \end{split}$$

The formula 66 for the intertwiner  $(\Phi_{\lambda,(+1,0)})$  in Borel-Weil realization can be rewritten in more invariant terms as follows:

$$(\Phi_{\lambda,(+1,0)})^{-1}f(x_{1},x_{2},x_{12})|0>_{\lambda+(1,0)} \to \frac{1}{(\lambda_{1}+1)(\lambda_{1}+\lambda_{2}+2)}((\partial_{x_{1}}\partial_{x_{2}}+((\lambda_{1}+2)+x_{2}\partial_{x_{2}})\partial_{x_{12}})f(x_{1},x_{2},x_{12})|0>_{\lambda}\otimes|2> + ((\lambda_{1}+\lambda_{2}+2-x_{2}\partial_{x_{2}})\partial_{x_{1}}+(\lambda_{2}-x_{2}^{2}\partial_{x_{2}})\partial_{x_{12}})f(x_{1},x_{2},x_{12})|0>_{\lambda}\otimes|1> + (-x_{12}\partial_{x_{1}})\partial_{x_{2}}+x_{1}x_{2}(-\lambda_{2}+x_{2}\partial_{x_{2}})\partial_{x_{12}}+((\lambda_{1}+1)(\lambda_{1}+\lambda_{2}+2)-(\lambda_{1}+\lambda_{2}+2)x_{1}\partial_{x_{1}}+x_{1}x_{2}\partial_{x_{1}}\partial_{x_{2}}-(\lambda_{1}+2)x_{1}\partial_{x_{1}}-x_{2}x_{12}\partial_{x_{2}}\partial_{x_{12}})f(x_{1},x_{2},x_{12})|0>_{\lambda}\otimes|0>).$$

$$(97)$$

Proposition: In Borel-Weil realization the action of the intertwiner on the Whittaker vector can be written as follows:

$$(\Phi_{\lambda,(+1,0)})^{-1} : exp(\mu_{1}x_{1} + \mu_{2}x_{2})|0>_{\lambda+1} \rightarrow \frac{1}{(\lambda_{1}+1)(\lambda_{1}+\lambda_{2}+2)}(\mu_{1}\mu_{2}exp(\mu_{1}x_{1}+\mu_{2}x_{2})|0>_{\lambda}\otimes|2> + (\mu_{1}(\lambda_{1}+\lambda_{2}+2) - \mu_{1}\mu_{2}x_{2})exp(\mu_{1}x_{1} + \mu_{2}x_{2})|0>_{\lambda}\otimes|1> + (\mu_{1}\mu_{2}(x_{1}x_{2} - x_{12}) - \mu_{1}(\lambda_{1}+\lambda_{2}+2)x_{1} + ((\lambda_{1}+1)(\lambda_{1}+\lambda_{2}+2))exp(\mu_{1}x_{1} + \mu_{2}x_{2})|0>_{\lambda}\otimes|0>).$$
(98)

### References

[1] E.Date, M.Jimbo, M.Kashiwara, T.Miwa, *Transformation groups for soliton equations*, in: Proc.RIMS symp. Nonlinear integrable systems — classical theory and quantum theory, M.Jimbo, T.Miwa, eds., World Scientific, Singapore 1983, p.39

- [2] M.A.Olshanetsky, A.M.Perelomov, Classical integrable finite-dimensional systems related to Lie algebras, Phys.Rept., 71 (1981) 313-400
  - M.Olshanetsky, A.Perelomov, Quantum integrable systems related to Lie algebras, Phys.Rept., 94 (1983) 313-404
  - J.-L. Verdier, Algebras de Lie, systems Hamiltonienes, courbes algebriques, Seminaire Bourbaki, 34-e annee, 1980/81, N 566, p. 1-10. Asterisque 1980. (Russian translation: Algebra i teoria chisel,pp. 8-29, sbornik "Matemeika.Novoe v zarubeznoi nauke." 43, Mir 1987)
  - J.-L. Verdier, Les representations des algebras de Lie affines: applications a quelques problemes de physique, Seminaire Bourbaki, 34-e annee, 1981/82, N 596, p. 1-13. Asterisque 1982. (Russian translation: Algebra i teoria chisel, pp. 84-99, sbornik "Matemeika. Novoe v zarubeznoi nauke." 43, Mir 1987)
  - A. Gorsky, Integrable many body systems in the field theories, in Theor. Mat. Phys. vol. 103, N 3, 1995; hep-th/9410228.
  - Jimbo, M., and Miwa, T. Algebraic Analysis of Solvable Lattice Models. Conference Board of the Math. Sci., Regional Conference Series in Mathematics, 85 (1995).
  - E. Frenkel, Five lectures on soliton equations, q-alg/9712005.
  - I. Cherednik, Lectures on affine Knizhnik-Zamoldchikov equations, quantum many-body problems, Hecke algebras, and Macdonald theory RIMS 1998
  - S. Kharchev, Kadomtsev-Petviashvili Hierachy and generalized Kontsevich model hep-th/9810091
- [3] V.Drinfeld and Sokolov, Lie algebras and evolutions of KdV type, J.Sov.Math., 30 (1985) 1975-2036
- [4] A. Pressley, G. Segal Loop groups, Oxford: Claredon Press 1986.
  - V. Kac, Infinite-dimensional Lie algebras, third edition, Cambridge University Press.
  - A.N.Leznov, M.V.Saveliev, Theory group methods for integrating non-linear dynamical systems, Moscow, "Nauka", 1985.
- [5] P. Etingof, I. Frenkel, A. Kirillov Representation theory and Kniznik-Zamolodchikov equation. AMS-1998
- [6] A. Mironov, Tau-function within group theory approach and its quantization, hep-th/9711006.
- [7] A.Gerasimov, S.Kharchev, A.Marshakov, A.Mironov, A.Morozov, M.Olshanetsky, Liouville type models in group theory framework.
   I. Finite-dimensional algebras, preprint FIAN/TD-18/95, ITEP M4/TH-7/95; hep-th/9601161
- [8] A.Gerasimov, S.Khoroshkin, D.Lebedev, A.Mironov, A.Morozov, Generalized Hirota equations and representation theory. I. The case of SL(2) and SL<sub>q</sub>(2), Int.J.Mod.Phys., A10 (1995) 2589-2614
- [9] M. Golenisheva-Kutozova, D. Lebedev, Itertwining operators and soliton equations, hep-th/9805186.
- [10] A. Antonov, B. Feigin, Quantum Group Representations and Baxter Equation, Phys.Lett. B392 (1997) 115-122, hep-th/9603105
- [11] A. Tsuchiya, Y. Kanie Vertex operators in conformal field theory on P<sup>1</sup> and monodromy representations of braid group, Adv. Stud. Pure Math. 16 297-372 (1988).
- [12] I.Frenkel and N.Reshetikhin, Comm.Math.Phys., 146 (1992) 1.
- [13] N. Vilenkin, A. Klimyk Representations of Lie groups and special functions, Kluwer Academic Publishers 1995.
- [14] J. Van der Jeugt, R. Jagannathan Realizations of su(1,1) and  $U_q(su(1,1))$  and generating functions for orthogonal polynomials, math-ph/9807019.
- [15] O.I.Bogoyavlensky, On perturbations of the periodic Toda lattice, Comm. Math. Phys., 51 (1976) 201-209.
- [16] M.Semenov-Tian-Shansky, Quantization of the open Toda chains, in "Sovremenie problemi matematiki" VINITI, 16 (1987) 194-226.
- [17] B.Kostant, Whittaker vectors and representation theory, Invent. Math., 48 (1978) 101-184.
- [18] Anatol N. Kirillov, Masatoshi Noumi, Affine Hecke algebras and raising operators for Macdonald polynomials, q-alg/9605004.
  Anatol N. Kirillov, Masatoshi Noumi, q-Difference raising operators for Macdonald polynomials and the integrality of transition coefficients, q-alg/9605005.
  - Luc Lapointe, Luc Vinet Creation operators for the Macdonald and Jack polynomials, q-alg/9607024.
  - Luc Lapointe, Luc Vinet, Rodrigues formulas for the Macdonald polynomials q-alg/9607025.
  - Yasushi Kajihara, Masatoshi Noumi, Raising operators of row type for Macdonald polynomials, math/9803151.
- [19] I.Gradshtein and I.Ryzhik, Tables of integrals, sums, series and products, Fizmatgiz, Moscow 1963.
- [20] P. Etingof, Whittaker functions on quantum groups and q-deformed Toda operators, math.QA/9901053.
- [21] A. Sevostyanov, The Whittaker model of the center of the quantum group and Hecke algebras, math.QA/9904075.

- [22] V. Bazhanov, S. Lukyanov, A. Zamolodchikov, Itegrable structure of Conformal field theory I,II,III, hep-th/9412229, hep-th/9604044, hep-th/9805008.
- [23] P. Etingof, Integral formulas for wave functions of quantum many-body problems and representations of  $gl_n$ , hep-th/9405038.
- [24] B. Feigin, E. Frenkel, in Knizhnik Memorial volume World Scientific, Singapore, 1990, pp. 271-316.
- [25] G. Heckmann, Invent. Math. 98, (1991) 341; C. Dunkl, Trans. Amer. Math. Soc., 311, (1989); E. Opdam, Invent. Math. 98 (1989) 1;
- [26] J. Denef, F. Loeser, Character sums associated to finite Coxeter groups, math.AG/9803033.