

## A construction of finite-dimensional faithful representation of Lie algebra

YURI A. NERETIN

The Ado theorem is a fundamental fact, which has a reputation to be a 'strange theorem'. We give its natural proof.

**1. Construction of faithful representation.** Consider a finite-dimensional Lie algebra  $\mathfrak{g}$ . Assume that  $\mathfrak{g}$  is a semidirect product  $\mathfrak{p} \ltimes \mathfrak{n}$  of a subalgebra  $\mathfrak{p}$  and a nilpotent ideal  $\mathfrak{n}$ . Assume that the adjoint action of  $\mathfrak{p}$  on  $\mathfrak{n}$  is faithful, i.e., for any  $z \in \mathfrak{p}$ , there exists  $x \in \mathfrak{n}$  such that  $[z, x] \neq 0$ .

Consider the minimal  $k$  such that all the commutators

$$[\dots [x_1, x_2], x_3], \dots, x_k], \quad x_j \in \mathfrak{n}$$

are 0.

Denote by  $U(\mathfrak{n})$  the enveloping algebra of  $\mathfrak{n}$ . The algebra  $\mathfrak{n}$  acts on  $U(\mathfrak{n})$  by the left multiplications. The algebra  $\mathfrak{p}$  acts on  $U(\mathfrak{n})$  by the derivations

$$d_z x_1 x_2 x_3 \dots x_l = [z, x_1] x_2 x_3 \dots x_l + x_1 [z, x_2] x_3 \dots x_l + \dots, \quad \text{where } z \in \mathfrak{p}.$$

This defines the action of the semidirect product  $\mathfrak{p} \ltimes \mathfrak{n} = \mathfrak{g}$  on  $U(\mathfrak{n})$ .

Denote by  $I$  the subspace in  $U(\mathfrak{n})$  spanned by all the products  $x_1 x_2 \dots x_N$ , where  $N > k + 2$ . Obviously,

1.  $I$  is the two-side ideal in  $U(\mathfrak{n})$ .
2. Consider the linear span  $\mathcal{A}$  of all the elements having the form  $1, x, x_1 x_2 \in U(\mathfrak{n})$ . Obviously,  $I \cap \mathcal{A} = 0$ .
3.  $I$  is invariant with respect to the derivations  $d_z$ .

Obviously, the module  $U(\mathfrak{n})/I$  is a finite-dimensional faithful module over  $\mathfrak{g}$ .

### 2. The Ado theorem.

LEMMA 1. *Any finite-dimensional Lie algebra  $\mathfrak{q}$  admits an embedding to an algebra  $\mathfrak{g}$  such that*  
*a)  $\mathfrak{g}$  is a semidirect product of a reductive subalgebra  $\mathfrak{p}$  and a nilpotent ideal  $\mathfrak{n}$ ;*  
*b) the action of  $\mathfrak{p}$  on  $\mathfrak{n}$  is completely reducible.*

Obviously, Lemma 1 implies the Ado theorem. Indeed,  $\mathfrak{g}$  admits a decomposition

$$\mathfrak{g} = \mathfrak{p}' \oplus (\mathfrak{p}'' \ltimes \mathfrak{n})$$

where  $\mathfrak{p}'$ ,  $\mathfrak{p}''$  are reductive subalgebras and the action of  $\mathfrak{p}''$  on  $\mathfrak{n}$  is faithful. After this, it is sufficient to apply the construction of p.1.

REMARK. The Ado theorem implies Lemma 1 modulo the Chevalley construction of algebraic envelope of a Lie algebra. But Lemma 1 itself can be easily proved directly.

**3. Killing lemma.** Let  $\mathfrak{g}$  be a Lie algebra, let  $d$  be its derivation. For an eigenvalue  $\lambda$ , denote by  $\mathfrak{g}_\lambda$  its root subspace  $\mathfrak{g}_\lambda = \cup_k \ker(d - \lambda)^k$ ; we have  $\mathfrak{g} = \oplus \mathfrak{g}_\lambda$ . As it was observed by Killing,  $x \in \mathfrak{g}_\lambda$ ,  $y \in \mathfrak{g}_\mu$  implies  $[x, y] \in \mathfrak{g}_{\lambda+\mu}$ .

Thus the Lie algebra  $\mathfrak{g}$  admits the gradation by the eigenvalues of  $d$ . Consider the gradation operator  $d_s : \mathfrak{g} \rightarrow \mathfrak{g}$  defined by  $d_s v = \lambda v$  if  $v \in \mathfrak{g}_\lambda$ . Obviously,  $d_s$  is a derivation, and  $dd_s = d_s d$ . We also consider the derivation  $d_n := d - d_s$ , this operator is nilpotent (the equality  $d = d_n + d_s$  is called the Jordan–Chevalley decomposition). Clearly,

$$\ker d_s \supset \ker d; \quad \ker d_n \supset \ker d; \tag{1}$$

$$\operatorname{im} d_s \subset \operatorname{im} d; \quad \operatorname{im} d_n \subset \operatorname{im} d_s \tag{2}$$

**4. Elementary expansions.** Let  $\mathfrak{q}$  be a Lie algebra, let  $I$  be an ideal of codimension 1. Let  $x \notin I$ . Denote by  $d$  the operator  $\operatorname{Ad}_x : I \rightarrow I$ . Consider the corresponding pair of derivations  $d_s$ ,  $d_n$ . Consider the space

$$\mathfrak{q}' = \mathbb{C}y + \mathbb{C}z + I$$

where  $y, z$  are formal vectors. We equip this space with a structure of a Lie algebra by the rule

$$[y, z] = 0, \quad [y, u] = d_s u, \quad [z, u] = d_n u, \quad \text{for all } u \in I$$

and the commutator of  $u, v \in I$  is the same as it was in  $I$ .

The subalgebra  $\mathbb{C}(y + z) \oplus I \subset \mathfrak{q}'$  is isomorphic  $\mathfrak{q}$ . We say that  $\mathfrak{q}'$  is an *elementary expansion* of  $\mathfrak{q}$ .

Obviously,  $[\mathfrak{q}', \mathfrak{q}'] = [\mathfrak{q}, \mathfrak{q}]$ .

For a general Lie algebra, the required embedding to a semidirect product can be obtained by a sequence of elementary expansions.

**5. Proof of Lemma 1.** Let  $\mathfrak{q}$  be a Lie algebra. Let  $\mathfrak{h}$  be its Levi part, and  $\mathfrak{r}$  be the radical. Denote by  $\mathfrak{m}$  the nilradical of  $\mathfrak{q}$ , i.e.,  $\mathfrak{m} = [\mathfrak{q}, \mathfrak{r}]$ ; recall that  $\mathfrak{m}$  is a nilpotent ideal, and  $[\mathfrak{q}, \mathfrak{q}] = \mathfrak{h} \ltimes \mathfrak{m}$  (see [1], 1.4.9).

Consider a nilpotent ideal  $\mathfrak{n}$  of  $\mathfrak{q}$  containing the nilradical  $\mathfrak{m}$ . Consider a subalgebra  $\mathfrak{p} \supset \mathfrak{h}$  such that the adjoint action of  $\mathfrak{p}$  on  $\mathfrak{q}$  is completely reducible and  $\mathfrak{p} \cap \mathfrak{n} = 0$ ; for instance, the can choice  $\mathfrak{n} = \mathfrak{m}$ ,  $\mathfrak{p} = \mathfrak{h}$ .

Obviously, the  $\mathfrak{q}$ -module  $\mathfrak{q}/(\mathfrak{p} \ltimes \mathfrak{n})$  is trivial. Consider any subspace  $I$  of codimension 1 containing  $\mathfrak{p} \ltimes \mathfrak{n}$ , obviously  $I$  is an ideal in  $\mathfrak{q}$ . Since the action of  $\mathfrak{p}$  on  $\mathfrak{q}$  is completely reducible, there exists a  $\mathfrak{p}$ -invariant complementary subspace for  $I$ . Let  $x$  be an element of this subspace. Since the  $\mathfrak{p}$ -module  $\mathfrak{q}/I$  is trivial,  $x$  commutes with  $\mathfrak{p}$ . We apply the elementary expansion to these data.

We obtain the new algebra  $\mathfrak{q}' = \mathbb{C}y + \mathbb{C}z + I$  with the nilpotent ideal  $\mathfrak{n}' = \mathbb{C}z + \mathfrak{n}$  and with the reductive subalgebra  $\mathfrak{p}' = \mathbb{C}y \oplus \mathfrak{p}$  (by (1),  $y$  commutes with  $\mathfrak{p}$ ).

It remains to notice that

$$\dim \mathfrak{q}' - \dim \mathfrak{p}' - \dim \mathfrak{n}' = \dim \mathfrak{q} - \dim \mathfrak{p} - \dim \mathfrak{n} - 1$$

and we can repeat the same construction.

## References

- [1] J.Dixmier, *Enveloping algebras*, North.Holland Publ.Co, 1977

Addres:

Math.Physics Group, ITEP, B.Chermushkinskaya, 25, Moscow, Russia

&

Independent University of Moscow

&

ESI, Vienna, Austria (January, 2002).

neretin@main.mccme.rssi.ru