

# q-rook monoid algebras, Hecke algebras, and Schur-Weyl duality

Tom Halverson\*

Department of Mathematics and Computer Science

Macalester College

St. Paul, MN 55105

halverson@macalester.edu

and

Arun Ram\*\*

Department of Mathematics

University of Wisconsin, Madison

Madison, WI 53706

ram@math.wisc.edu

*In memory of Sergei Kerov 1946-2000*

*When we were at the beginnings of our careers Sergei's support helped us to believe in our work. He generously encouraged us to publish our results on Brauer and Birman-Murakami-Wenzl algebras, results which had in part, or possibly in total, been obtained earlier by Sergei himself. He remains a great inspiration for us, both mathematically and in our memory of his kindness, modesty, generosity, and encouragement to the younger generation.*

## 0. Introduction

The rook monoid  $R_k$  is the monoid of  $k \times k$  matrices with entries from  $\{0, 1\}$  and at most one nonzero entry in each row and column. Recently, the representation theory of its “Iwahori-Hecke” algebra  $R_k(q)$ , called the  $q$ -rook monoid algebra, has been analyzed. In particular, a Schur-Weyl type duality on tensor space was found for the  $q$ -rook monoid algebra and its irreducible representations were given explicit combinatorial constructions. In this paper we show that, in fact, the  $q$ -rook monoid algebra is a quotient of the affine Hecke algebra of type A. With this knowledge in hand, we show that the recent results on the  $q$ -rook monoid algebras actually come from known results about the affine Hecke algebra. In particular

- (a) The recent combinatorial construction of the irreducible representations of  $R_k(q)$  by Halverson [Ha] turns out to be a special case of the construction of irreducible calibrated representations of affine Hecke algebras of Cherednik [Ch] (see also Ram [Ra]), the construction of irreducible representations of cyclotomic Hecke algebras by Ariki and Koike

---

\* Research supported in part by National Science Foundation grant DMS-0100975.

\*\* Research supported in part by National Science Foundation grant DMS-9971099, the National Security Agency and EPSRC grant GR K99015.

[AK], and the construction of the irreducible representations of Iwahori-Hecke algebras of type B by Hoefsmit [Ho].

- (b) The Schur-Weyl duality for the  $q$ -rook monoid algebra discovered by Solomon [So2-4] and studied by Halverson [Ha] turns out to be a special case of the Schur-Weyl duality for cyclotomic Hecke algebras given by Sakamoto and Shoji [SS].

Though these results show that the representation theory of the  $q$ -rook monoid algebra is “just” a piece of the representation theory of the affine Hecke algebra, this was not at all obvious at the outset. It was only on the analysis of the recent results in [So4] and [Ha] that the similarity to affine Hecke algebra theory was noticed. This observation then led us to search for and establish a concrete connection between these algebras.

The  $q$ -rook monoid algebra was first studied in its  $q = 1$  version in the 1950’s by Munn [Mu1-2]. Solomon [Sol] discovered the general  $q$ -version of the algebra as a Hecke algebra (double coset algebra) for the finite algebraic monoid  $M_n(\mathbb{F}_q)$  of  $n \times n$  matrices over a finite field with  $q$  elements, with respect to the “Borel subgroup”  $B$  of invertible upper triangular matrices. Later Solomon [So2] found a Schur-Weyl duality for  $R_k(1)$  in which  $R_k(1)$  acts as the centralizer algebra for the action of the general linear group  $GL_n(\mathbb{C})$  on  $V^{\otimes k}$  where  $V = L(\varepsilon_1) \oplus L(0)$  is the direct sum of the “fundamental”  $n$ -dimensional representation and the trivial module  $L(0)$  for  $GL_n(\mathbb{C})$ . Then Solomon [So3,4] gave a presentation of  $R_k(q)$  by generators and relations and defined an action of  $R_k(q)$  on tensor space.

Halverson [Ha] found a new presentation of  $R_k(q)$  and used it to show that Solomon’s action of  $R_k(q)$  on tensor space extends the Schur-Weyl duality so that  $R_k(q)$  is the centralizer of the quantum general linear group  $U_q \mathfrak{gl}(n)$  on  $V^{\otimes k}$  where now  $V = L(\varepsilon_1) \oplus L(0)$  is the direct sum of the “fundamental” and the trivial module for  $U_q \mathfrak{gl}(n)$ . Halverson also exploited his new presentation to construct, combinatorially, all the irreducible representations of  $R_k(q)$  when  $R_k(q)$  is semisimple.

The main results of this paper are the following:

- (a) We find yet another presentation (1.6) of  $R_k(q)$  by generators and relations.
- (b) Our new presentation shows that

$$R_k(q) = H_k(0, 1; q)/I,$$

where  $H_k(0, 1; q)$  is the Iwahori-Hecke algebra of type  $B_k$  with parameters specialized to 0 and 1, and  $I$  is the ideal generated by the minimal ideal of  $H_2(0, 1; q)$  corresponding to the pair of partitions  $\lambda = ((1^2), \emptyset)$ .

- (c) We show that the irreducible representations of  $R_k(q)$  found in [Ha] come from the constructions of irreducible representations of  $H_k(0, 1; q)$ .
- (d) We use the fact that  $R_k(q)$  is a quotient of  $H_k(0, 1; q)$  and the fact that the Iwahori-Hecke algebra  $H_k(q)$  of type  $A_{k-1}$  is a quotient of  $R_k(q)$  to easily determine, in Corollary 2.21, the values of  $q$  for which  $R_k(q)$  is semisimple. These values were first found in [So4] using other methods.
- (e) We show that the Schur-Weyl duality between  $R_k(q)$  and  $U_q \mathfrak{gl}(n)$  comes from the Schur-Weyl duality of Sakamoto and Shoji [SS] for the cyclotomic Hecke algebras (Theorem 3.5).
- (f) We give a *different* Schur-Weyl duality for algebras  $A_k(u_1, u_2; q) = H_k(u_1, u_2; q)/I$ , where  $u_1, u_2 \neq 0$ ,  $H_k(u_1, u_2; q)$  is the Iwahori-Hecke algebra of type  $B_k$  and  $I$  is the ideal generated by the minimal ideal of  $H_2(u_1, u_2; q)$  corresponding to the pair of partitions  $\lambda = ((1^2), \emptyset)$ . This Schur-Weyl duality comes from the Schur-Weyl duality of Orellana and Ram for the affine Hecke algebra (Theorem 3.3).

**Acknowledgements.** A. Ram thanks the Isaac Newton Institute for the Mathematical Sciences at Cambridge University for support for a very pleasant residency during which the research in this paper was completed.

## 1. Presentations of the $q$ -rook monoid algebras

Fix  $q \in \mathbb{C}^*$ . The  $q$ -rook monoid algebra is the algebra  $R_k(q)$  given by generators

$$P_1, P_2, \dots, P_k \quad \text{and} \quad T_1, T_2, \dots, T_{k-1}$$

with relations

$$\begin{aligned} \text{(A1)} \quad & T_i^2 = (q - q^{-1})T_i + 1, & 1 \leq i \leq k-1, \\ \text{(A2)} \quad & T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, & 1 \leq i \leq k-2, \\ \text{(A3)} \quad & T_i T_j = T_j T_i, & |i - j| > 1, \\ \text{(R1)} \quad & P_i^2 = P_i, & 1 \leq i \leq k, \\ \text{(R2)} \quad & P_i P_j = P_j P_i, & 1 \leq i, j \leq k, \\ \text{(R3)} \quad & P_i T_j = T_j P_i, & 1 \leq i < j \leq k, \\ \text{(R4)} \quad & P_i T_j = T_j P_i = q P_i, & 1 \leq j < i \leq k, \\ \text{(R5)} \quad & P_{i+1} = q P_i T_i^{-1} P_i = q(P_i T_i P_i - (q - q^{-1})P_i), & 1 \leq i \leq k-1. \end{aligned} \tag{1.1}$$

The algebra  $R_k(q)$  was introduced by Solomon [Sol] as an analogue of the Iwahori-Hecke algebra for the finite algebraic monoid  $M_k(\mathbb{F}_q)$  of  $k \times k$  matrices over a finite field with  $q$  elements with respect to its “Borel subgroup” of invertible upper triangular matrices. The presentation of  $R_k(q)$  given above is due to Halverson [Ha].

When  $q = 1$ ,  $R_k(q)$  specializes to the algebra of the rook monoid  $R_k$  that consists of  $k \times k$  matrices with entries from  $\{0, 1\}$  and *at most* one nonzero entry in each row and column. These correspond with the possible placements of nonattacking rooks on an  $k \times k$  chessboard. In this specialization,  $T_i$  becomes the matrix obtained by switching rows  $i$  and  $i+1$  in the identity matrix  $I$  and, for  $1 \leq i \leq k-1$ ,  $P_i$  becomes the matrix  $E_{i+1,i+1} + E_{i+1,i+2} + \dots + E_{k,k}$ , where  $E_{i,j}$  is the matrix with a 1 in position  $(i, j)$  and zeros elsewhere. The generator  $P_k$  specializes to the 0 matrix (which is not the 0 element in the monoid algebra).

**Remark 1.2.** The definition of  $R_k(q)$  in [Sol, 3, 4] and [Ha] uses generators  $\tilde{T}_i$  in place of  $T_i$ . These generators satisfy  $\tilde{T}_i^2 = (q-1)\tilde{T}_i + q$  in place of (A1). In our presentation, if we let  $\tilde{T}_i = qT_i$ , then  $\tilde{T}_i^2 = q^2((q - q^{-1})T_i + 1) = (q^2 - 1)qT_i + q^2 = (q^2 - 1)\tilde{T}_i + q^2$ , which shows that our algebra is the same except with parameter  $q^2$  instead of  $q$ .

Define

$$X_i = T_{i-1} T_{i-2} \dots T_1 (1 - P_1) T_1 T_2 \dots T_{i-1}, \quad 1 \leq i \leq k, \tag{1.3}$$

so that  $X_{i+1} = T_i X_i T_i$ .

**Lemma 1.4.** In  $R_k(q)$  we have the following relations

- (a)  $P_i P_j = P_j P_i = P_j$ , for  $i \leq j$ .
- (b)  $P_1 X_2 = P_1 - P_2$ .

*Proof.* (a) If  $i = j$  this is (R1). If  $i < j$  then, by (R5) and induction,

$$P_i P_j = P_i (P_{j-1} T_{j-1} P_{j-1} - (q - q^{-1})P_{j-1}) = P_{j-1} T_{j-1} P_{j-1} - (q - q^{-1})P_{j-1} = P_j.$$

(b) We use relations (A1), (R4), and (R5) to get

$$\begin{aligned}
P_1 X_2 &= P_1 T_1 (1 - P_1) T_1 = P_1 T_1^2 - P_1 T_1 P_1 T_1 \\
&= (q - q^{-1}) P_1 T_1 + P_1 - P_1 T_1 P_1 T_1 \\
&= (q - q^{-1}) P_1 T_1 + P_1 - q^{-1} P_2 T_1 - (q - q^{-1}) P_1 T_1 \\
&= P_1 - q^{-1} P_2 T_1 = P_1 - P_2. \quad \blacksquare
\end{aligned}$$

**Proposition 1.5.** *The  $q$ -rook monoid algebra  $R_k(q)$  is generated by the elements  $X_1, T_1, \dots, T_{k-1}$ , and these elements satisfy the relations (A1), (A2), (A3), and*

- (B1)  $X_1 T_j = T_j X_1$ , for  $2 \leq j \leq k$ ,
- (B2)  $X_1^2 = X_1$ ,
- (B3)  $X_1 T_1 X_1 T_1 = T_1 X_1 T_1 X_1$ ,
- (B4)  $(1 - X_1)(T_1 - q)(1 - X_1)(1 - X_2) = 0$ , where  $X_2 = T_1 X_1 T_1$ .

*Proof.* By (R5),  $R_n(q)$  is generated by  $X_1 = 1 - P_1, T_1, \dots, T_{k-1}$ .

(B1) By (R3),  $X_1 T_j = (1 - P_1) T_j = T_j (1 - P_1) = T_j X_1$ .

(B2) By (R1),  $X_1^2 = (1 - P_1)^2 = 1 - 2P_1 + P_1^2 = 1 - P_1 = X_1$ .

(B3) Using (R4) and (R5),

$$q(T_1 P_1 T_1 P_1 - (q - q^{-1}) T_1 P_1) = T_1 P_2 = P_2 T_1 = q(P_1 T_1 P_1 T_1 - (q - q^{-1}) P_1 T_1),$$

and so

$$P_1 T_1 P_1 T_1 = T_1 P_1 T_1 P_1 + (q - q^{-1})(P_1 T_1 - T_1 P_1). \quad (*)$$

Now, using (\*) and (A1),

$$\begin{aligned}
X_1 T_1 X_1 T_1 &= (1 - P_1) T_1 (1 - P_1) T_1 \\
&= T_1^2 - P_1 T_1^2 - T_1 P_1 T_1 + P_1 T_1 P_1 T_1 \\
&= T_1^2 - (q - q^{-1}) P_1 T_1 - P_1 - T_1 P_1 T_1 + T_1 P_1 T_1 P_1 + (q - q^{-1})(P_1 T_1 - T_1 P_1) \\
&= T_1^2 - P_1 - T_1 P_1 T_1 + T_1 P_1 T_1 P_1 - (q - q^{-1}) T_1 P_1 \\
&= T_1^2 - T_1^2 P_1 - T_1 P_1 T_1 + T_1 P_1 T_1 P_1 \\
&= T_1 (1 - P_1) T_1 (1 - P_1) \\
&= T_1 X_1 T_1 X_1.
\end{aligned}$$

Finally, to show (B4), we use Lemma 1.4(b),

$$\begin{aligned}
(1 - X_1)(T_1 - q)(1 - X_1)(1 - X_2) &= P_1 (T_1 - q) P_1 (1 - X_2) \\
&= P_1 T_1 P_1 - q P_1 - P_1 T_1 P_1 X_2 + q P_1 X_2 \\
&= P_1 T_1 P_1 - q P_1 - P_1 T_1 (P_1 - P_2) + q (P_1 - P_2) \\
&= -q P_2 + P_1 T_1 P_2 \\
&= -q P_2 + q P_1 P_2 \quad \text{by (R4)} \\
&= -q P_2 + q P_2 \quad \text{by Lemma 1.4(a)} \\
&= 0. \quad \blacksquare
\end{aligned}$$

Define a new algebra  $A_k(q)$  by generators

$$X_1 \quad \text{and} \quad T_1, T_2, \dots, T_{k-1}$$

and relations

$$\begin{aligned} \text{(A1)} \quad & T_i^2 = (q - q^{-1})T_i + 1, & 1 \leq i \leq k-1, \\ \text{(A2)} \quad & T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, & 1 \leq i \leq k-2, \\ \text{(A3)} \quad & T_i T_j = T_j T_i, & |i - j| > 1, \\ \text{(B1)} \quad & X_1 T_j = T_j X_1, & 2 \leq j \leq k, \\ \text{(B2)} \quad & X_1^2 = X_1, \\ \text{(B3)} \quad & X_1 T_1 X_1 T_1 = T_1 X_1 T_1 X_1, \\ \text{(B4)} \quad & (1 - X_1)(T_1 - q)(1 - X_1)(1 - X_2) = 0, \quad \text{where } X_2 = T_1 X_1 T_1. \end{aligned} \tag{1.6}$$

We will show that  $R_k(q) \cong A_k(q)$ . Define

$$P_1 = (1 - X_1) \quad \text{and} \quad P_{i+1} = q(P_i T_i P_i - (q - q^{-1})P_i), \quad 1 \leq i \leq k-1. \tag{1.7}$$

**Lemma 1.8.** *Relations (B1)-(B4) are equivalent, respectively, to*

$$\begin{aligned} \text{(B1')} \quad & P_1 T_j = T_j P_1, \text{ for } 2 \leq j \leq n, \\ \text{(B2')} \quad & P_1^2 = P_1, \\ \text{(B3')} \quad & P_2 T_1 = T_1 P_2, \\ \text{(B4')} \quad & P_2^2 = P_2. \end{aligned}$$

*Proof.* Subtracting  $T_j$  from each side of

$$T_j - P_1 T_j = (1 - P_1)T_j = X_1 T_j = T_j X_1 = T_j(1 - P_1) = T_j - T_j P_1,$$

shows that (B1) is equivalent to (B1'). Relations (B2) and (B2') are equivalent since

$$1 - P_1 = X_1 = X_1^2 = (1 - P_1)^2 = 1 - 2P_1 + P_1^2.$$

Since

$$\begin{aligned} X_1 T_1 X_1 T_1 &= (1 - P_1)T_1(1 - P_1)T_1 = (1 - P_1)T_1^2 - T_1 P_1 T_1 + P_1 T_1 P_1 T_1 \\ &= (1 - P_1)((q - q^{-1})T_1 + 1) - T_1 P_1 T_1 + (q^{-1}P_2 + (q - q^{-1})P_1)T_1 \quad (\text{by (A1) and (1.7)}) \\ &= (q - q^{-1})T_1 + 1 - (q - q^{-1})P_1 T_1 - P_1 - T_1 P_1 T_1 + q^{-1}P_2 T_1 + (q - q^{-1})P_1 T_1 \\ &= (q - q^{-1})T_1 + 1 - P_1 - T_1 P_1 T_1 + q^{-1}P_2 T_1 \end{aligned}$$

is equal to

$$\begin{aligned} T_1 X_1 T_1 X_1 &= T_1(1 - P_1)T_1(1 - P_1) = T_1^2(1 - P_1) - T_1 P_1 T_1 + T_1 P_1 T_1 P_1 \\ &= ((q - q^{-1})T_1 + 1)(1 - P_1) - T_1 P_1 T_1 + T_1(q^{-1}P_2 + (q - q^{-1})P_1) \quad (\text{by (A1) and (1.7)}) \\ &= (q - q^{-1})T_1 + 1 - (q - q^{-1})T_1 P_1 - P_1 - T_1 P_1 T_1 + q^{-1}T_1 P_2 + (q - q^{-1})T_1 P_1 \\ &= (q - q^{-1})T_1 + 1 - P_1 - T_1 P_1 T_1 + q^{-1}T_1 P_2, \end{aligned}$$

(B3) is equivalent to (B3').

Expanding

$$\begin{aligned}
(T_1 - q)(1 - X_2) &= (T_1 - q)(1 - T_1 X_1 T_1) = (T_1 - q)(1 - T_1(1 - P_1)T_1) \\
&= (T_1 - q)(1 - (q - q^{-1})T_1 - 1 + T_1 P_1 T_1) \quad (\text{by (A1)}) \\
&= (T_1 - q)T_1(-(q - q^{-1}) + P_1 T_1) \\
&= ((q - q^{-1})T_1 + 1 - qT_1)(q^{-1} - q + P_1 T_1) \quad (\text{by (A1)}) \\
&= (1 - q^{-1}T_1)(q^{-1} - q + P_1 T_1) \\
&= q^{-1} - q + P_1 T_1 - q^{-2}T_1 + T_1 - q^{-1}T_1 P_1 T_1 \\
&= -(q - q^{-1}) + q^{-1}(q - q^{-1})T_1 + P_1 T_1 - q^{-1}T_1 P_1 T_1,
\end{aligned}$$

gives

$$\begin{aligned}
(1 - X_1)(T_1 - q)(1 - X_2)(1 - X_1) &= P_1(T_1 - q)(1 - X_2)P_1 \\
&= -(q - q^{-1})P_1 + q^{-1}(q - q^{-1})P_1 T_1 P_1 + P_1 T_1 P_1 - q^{-1}P_1 T_1 P_1 T_1 P_1 \quad (\text{by (B2')}) \\
&= -(q - q^{-1})P_1 + (2 - q^{-2})P_1 T_1 P_1 - q^{-1}(P_1 T_1 P_1)^2 \quad (\text{by (B2')}) \\
&= -(q - q^{-1})P_1 + (2 - q^{-2})(q^{-1}P_2 + (q - q^{-1})P_1) - q^{-1}(q^{-1}P_2 + (q - q^{-1})P_1)^2 \quad (\text{by (1.7)}) \\
&= (q - q^{-1})(-1 + 2 - q^{-2})P_1 + q^{-1}(2 - q^{-2})P_2 \\
&\quad - q^{-1}(q^{-2}P_2^2 + 2q^{-1}(q - q^{-1})P_2 + (q - q^{-1})^2 P_1) \\
&= (q - q^{-1})^2(q^{-1} - q^{-1})P_1 + q^{-1}(q^{-2}P_2 - q^{-2}P_2^2) \\
&= q^{-3}(P_2 - P_2^2).
\end{aligned}$$

and so  $(1 - X_1)(T_1 - q)(1 - X_2)(1 - X_1) = 0$  if and only if  $P_2^2 = P_2$ . Thus (B4) is equivalent to (B4'). ■

**Proposition 1.9.** *The algebra  $A_k(q)$  is generated by  $T_1, \dots, T_{k-1}, P_1, \dots, P_k$ , and these elements satisfy the relations (A1), (A2), (A3), and*

- (E1)  $P_i T_j = T_j P_i$ , for  $1 \leq i < j \leq k$ ,
- (E2)  $P_i P_j = P_j P_i = P_i$ , for all  $1 \leq j < i \leq k$ ,
- (E3)  $P_i^2 = P_i$ , for  $1 \leq i \leq k$ ,
- (E4)  $P_i T_j = T_j P_i = q P_i$ , for  $1 \leq j < i \leq k$ ,
- (E5)  $P_{i+1} = q P_i T_i^{-1} P_i = q(P_i T_i P_i - (q - q^{-1})P_i)$ , for  $1 \leq i \leq k - 1$ .

*Proof.* Since  $X_1 = 1 - P_1$ , the elements  $T_1, \dots, T_{n-1}, P_1$  generate  $A_k(q)$ . Relation (E5) is the definition of  $P_{i+1}$ .

We prove (E1) by induction on  $i$ . The case  $i = 1$  is (B1'). Assume that  $j > i + 1 > 1$ , then  $T_j$  commutes with  $P_i$  by induction and  $T_j$  commutes with  $T_i$  by (A3), so

$$T_j P_{i+1} = q^{-1} T_j P_i T_j P_i - (q - q^{-1}) T_j P_i = q^{-1} P_i T_j P_i T_j - (q - q^{-1}) P_i T_j = P_{i+1} T_j,$$

proving (E1).

We now prove (E2)-(E4) collectively by induction on  $i$ . The case  $i = 1$  for (E2) follows from (B2') since

$$P_2 P_1 = q(P_1 T_1 P_1 - (q - q^{-1})P_1)P_1 = q(P_1 T_1 P_1 - (q - q^{-1})P_1) = P_2. \quad (*)$$

The relation  $P_1P_2 = P_2$  is similar. The first two cases of (E3) are (B2') and (B4'). The  $i = 1$  case of (E4) follows from (B4') since

$$\begin{aligned} P_2^2 &= P_2q(P_1T_1P_1 - (q - q^{-1})P_1) \\ &= q(P_2T_1P_1 - (q - q^{-1})P_2) \quad (\text{by } (*)) \\ &= q(T_1P_2P_1 - (q - q^{-1})P_2) \quad (\text{by (B3')}) \\ &= q(T_1P_2 - (q - q^{-1})P_2) \quad (\text{by } (*)). \end{aligned}$$

Since  $P_2^2 = P_2$ , we have  $T_1P_2 = qP_2$ . The case  $P_2T_1 = qP_2$  is similar.

Now fix  $i > 1$  and assume the following relations,

$$(E2^*) \quad P_iP_j = P_jP_i = P_i, \text{ for all } 1 \leq j < i,$$

$$(E3^*) \quad P_i^2 = P_i,$$

$$(E4^*) \quad P_iT_j = T_jP_i = qP_i, \text{ for } 1 \leq j < i,$$

We show each of these relations for  $i + 1$ .

For (E2) we use (E3\*) to get

$$P_{i+1}P_i = q(P_iT_iP_i^2 - (q - q^{-1})P_i^2) = q(P_iT_iP_i - (q - q^{-1})P_i) = P_{i+1},$$

and when  $j < i$ , we use (E2\*) to get

$$P_{i+1}P_j = q(P_iT_iP_iP_j - (q - q^{-1})P_iP_j) = q(P_iT_iP_i - (q - q^{-1})P_i) = P_{i+1}.$$

The relations  $P_iP_{i+1} = P_{i+1}$  and  $P_jP_{i+1} = P_{i+1}$  are similar, and so (E2) is established.

To establish (E3), let  $i > 2$  (note that we have established  $i = 1, 2$ ),

$$\begin{aligned} P_{i+1}^2 &= q^2(P_iT_iP_i - (q - q^{-1})P_i)^2 \\ &= q^2(P_iT_iP_iT_iP_i - 2(q - q^{-1})P_iT_iP_i + (q - q^{-1})^2P_i) \quad (\text{by (E3*)}) \\ &= q^2(P_iT_iqP_{i-1}T_{i-1}P_{i-1}T_iP_i - q(q - q^{-1})P_iT_iP_{i-1}T_iP_i \\ &\quad - 2(q - q^{-1})P_iT_iP_i + (q - q^{-1})^2P_i) \quad (\text{by (E5)}) \\ &= q^2(qP_iP_{i-1}T_iT_{i-1}T_iP_{i-1}P_i - q(q - q^{-1})P_iP_{i-1}T_i^2P_i \\ &\quad - 2(q - q^{-1})P_iT_iP_i + (q - q^{-1})^2P_i) \quad (\text{by (E1)}) \\ &= q^2(qP_iT_iT_{i-1}T_iP_i - q(q - q^{-1})P_iT_i^2P_i - 2(q - q^{-1})P_iT_iP_i + (q - q^{-1})^2P_i) \quad (\text{by (E2)}) \\ &= q^2(qP_iT_{i-1}T_iT_{i-1}P_i - q(q - q^{-1})P_iT_i^2P_i - 2(q - q^{-1})P_iT_iP_i + (q - q^{-1})^2P_i) \quad (\text{by (A2)}) \\ &= q^2(qP_iT_{i-1}T_iT_{i-1}P_i - q(q - q^{-1})^2P_iT_iP_i - q(q - q^{-1})P_i \\ &\quad - 2(q - q^{-1})P_iT_iP_i + (q - q^{-1})^2P_i) \quad (\text{by (A1)}) \\ &= q^2(q^3P_iT_iP_i - q(q - q^{-1})^2P_iT_iP_i - q(q - q^{-1})P_i \\ &\quad - 2(q - q^{-1})P_iT_iP_i + (q - q^{-1})^2P_i) \quad (\text{by (E4*)}) \\ &= q^2((q^3 - q(q - q^{-1})^2 - 2(q - q^{-1}))P_iT_iP_i + (q - q^{-1})(-q + q - q^{-1})P_i) \\ &= q(P_iT_iP_i - (q - q^{-1})P_i) \\ &= P_{i+1}. \end{aligned}$$

Finally, we prove (E4). First let  $j < i$ . Then, by (E4\*),

$$P_{i+1}T_j = q(P_iT_iP_iT_j - (q - q^{-1})P_iT_j) = q(qP_iT_iP_i - (q - q^{-1})qP_i) = qP_{i+1},$$

and  $T_j P_{i+1} = q P_{i+1}$  is similar. Now we consider the case where  $j = i$ ,

$$\begin{aligned}
P_{i+1} T_i &= q(P_i T_i P_i T_i - (q - q^{-1}) P_i T_i) \quad (\text{by (E5)}) \\
&= q(P_i T_i q P_{i-1} T_{i-1} P_{i-1} T_i - q(q - q^{-1}) P_i T_i P_{i-1} T_i - (q - q^{-1}) P_i T_i) \quad (\text{by (E5)}) \\
&= q(q P_i P_{i-1} T_i T_{i-1} T_i P_{i-1} - q(q - q^{-1}) P_i P_{i-1} T_i^2 - (q - q^{-1}) P_i T_i) \quad (\text{by (E1)}) \\
&= q(q P_i T_i T_{i-1} T_i P_{i-1} - q(q - q^{-1}) P_i T_i^2 - (q - q^{-1}) P_i T_i) \quad (\text{by (E2)}) \\
&= q(q P_i T_{i-1} T_i T_{i-1} P_{i-1} - q(q - q^{-1}) P_i T_i^2 - (q - q^{-1}) P_i T_i) \quad (\text{by (A2)}) \\
&= q(q^2 P_i T_i T_{i-1} P_{i-1} - q(q - q^{-1}) P_i T_i^2 - (q - q^{-1}) P_i T_i) \quad (\text{by (E4*)}) \\
&= q(q^2 P_i T_i T_{i-1} P_{i-1} - q(q - q^{-1})^2 P_i T_i - q(q - q^{-1}) P_i - (q - q^{-1}) P_i T_i) \quad (\text{by (A1)}) \\
&= q(q^2 P_i T_i T_{i-1} P_{i-1} - q^2(q - q^{-1}) P_i T_i - q(q - q^{-1}) P_i) \\
&= q(q^2 P_i P_{i-1} T_i T_{i-1} P_{i-1} - q^2(q - q^{-1}) P_i T_i - q(q - q^{-1}) P_i) \quad (\text{by (E2)}) \\
&= q(q^2 P_i T_i P_{i-1} T_{i-1} P_{i-1} - q^2(q - q^{-1}) P_i T_i - q(q - q^{-1}) P_i) \quad (\text{by (E1)}) \\
&= q(q P_i T_i P_i + q^2(q - q^{-1}) P_i T_i P_{i-1} - q^2(q - q^{-1}) P_i T_i - q(q - q^{-1}) P_i) \quad (\text{by (E5)}) \\
&= q(P_{i+1} + q^2(q - q^{-1}) P_i T_i P_{i-1} - q^2(q - q^{-1}) P_i T_i) \quad (\text{by (E5)}) \\
&= q(P_{i+1} + q^2(q - q^{-1}) P_i P_{i-1} T_i - q^2(q - q^{-1}) P_i T_i) \quad (\text{by (E1)}) \\
&= q(P_{i+1} + q^2(q - q^{-1}) P_i T_i - q^2(q - q^{-1}) P_i T_i) \quad (\text{by (E2)}) \\
&= q P_{i+1}.
\end{aligned}$$

The case  $T_i P_{i+1} = q P_{i+1}$  is similar. ■

Propositions 1.5 and 1.9 give the following theorem.

**Theorem 1.10.**  $R_k(q) \cong A_k(q)$  and thus (1.6) is a new presentation of  $R_k(q)$ .

## 2. Hecke algebras

The affine Hecke algebra  $\tilde{H}_k$

Fix  $q \in \mathbb{C}^*$ . The affine Hecke algebra  $\tilde{H}_k$  is the algebra given by generators

$$X_1, \dots, X_k \quad \text{and} \quad T_1, \dots, T_{k-1}$$

with relations

- (1)  $T_i T_j = T_j T_i, \quad |i - j| > 1,$
- (2)  $T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, \quad 1 \leq i \leq k - 2,$
- (3)  $T_i^2 = (q - q^{-1}) T_i + 1, \quad 1 \leq i \leq k,$
- (4)  $X_i X_j = X_j X_i, \quad 1 \leq i, j \leq k,$
- (5)  $X_i T_i = T_i X_{i+1} + (q - q^{-1}) X_i, \quad 1 \leq i \leq k - 1.$

It follows from relations (3) and (5) that

$$X_i = T_{i-1} \cdots T_2 T_1 X_1 T_1 T_2 \cdots T_{i-1}, \quad \text{for } 1 \leq i \leq k,$$



and from (4) that

$$(6) \quad X_1 T_1 X_1 T_1 = T_1 X_1 T_1 X_1.$$

In fact,  $\tilde{H}_k$  can be presented as the algebra generated by  $X_1$  and  $T_1, \dots, T_{k-1}$  with relations (1-3) and (6).

Let

$$[k]! = [1][2] \cdots [k], \quad \text{where} \quad [i] = 1 + q^2 + \cdots + q^{2(i-1)}.$$

When  $[k]! \neq 0$  a large class of irreducible representations of the affine Hecke algebra  $\tilde{H}_k(q)$ , the integrally calibrated irreducible representations, have a simple combinatorial construction. An  $\tilde{H}_k$ -module  $M$  is *integrally calibrated* if  $M$  has a basis of simultaneous eigenvectors for  $X_1, X_2, \dots, X_k$  for which the eigenvalues are all of the form  $q^j$  with  $j \in \mathbb{Z}$ . The construction of these  $\tilde{H}_k$ -modules is originally due to Cherednik [Ch] (see [Ra] for greater detail) and is a generalization of the classical seminormal construction of the irreducible representations of the symmetric group by A. Young. Young's construction had been generalized to Iwahori-Hecke algebras of classical type (see Theorem 2.8 below) by Hoefsmit [Ho] in 1974.

To describe the construction we shall use the notations of [Mac] for partitions so that a partition is identified with a collection of boxes in a corner,  $\ell(\lambda)$  is the number of rows of  $\lambda$ , and  $|\lambda|$  is the number of boxes in  $\lambda$ . For example, the partition

$$\lambda = (5, 5, 3, 1, 1) = \begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \square & \square & \square & \square & \\ \hline \square & \square & \square & & \\ \hline \square & & & & \\ \hline \square & & & & \\ \hline \end{array} \quad \text{has } \ell(\lambda) = 5 \text{ and } |\lambda| = 15.$$

If  $\lambda$  is a partition that is obtained from  $\mu$  by adding  $k$  boxes let  $\lambda/\mu$  be the *skew shape* consisting of those boxes of  $\lambda$  that are not in  $\mu$ . A *standard tableau* of shape  $\lambda/\mu$  is a filling of the boxes of  $\lambda/\mu$  with  $1, 2, \dots, k$  such that

- (a) the entries in the rows increase left to right, and
- (b) the entries in the columns increase top to bottom.

If  $b$  is a box in  $\lambda/\mu$  define

$$CT(b) = q^{2(c-r)}, \quad \text{if } b \text{ is in position } (r, c) \text{ of } \lambda. \quad (2.1)$$

**Theorem 2.2.** ([Ch], see also [Ra, Theorem 4.1], and [OR, Theorem 6.20a]) Assume that  $[k]! \neq 0$ . Then the calibrated irreducible representations  $H^{\lambda/\mu}$  of the affine Hecke algebra  $\tilde{H}_k$  are indexed by skew shapes and can be given explicitly as the vector space

$$H^{\lambda/\mu} = \mathbb{C}\text{-span}\{v_L \mid L \text{ is a standard tableau of shape } \lambda/\mu\}$$

(so that the symbols  $v_L$  form a basis of  $H^{\lambda/\mu}$ ) with  $\tilde{H}_k$ -action given by

$$\begin{aligned} X_i v_L &= CT(L(i)) v_L, \\ T_i v_L &= \left( \frac{CT(L(i+1))(q - q^{-1})}{CT(L(i+1)) - CT(L(i))} \right) v_L + \left( q^{-1} + \frac{CT(L(i+1))(q - q^{-1})}{CT(L(i+1)) - CT(L(i))} \right) v_{s_i L}, \end{aligned}$$

where

$s_i L$  is the same as  $L$  except  $i$  and  $i+1$  are switched, and

$v_{s_i L} = 0$ , if  $s_i L$  is not a standard tableau.

**Remark 2.3.** In [OR] it is explained how the basis  $v_L$  of  $H^{\lambda/\mu}$  and the action of  $\tilde{H}_k$  in Theorem 2.2 can be derived in a natural way from the general mechanism of quantum groups ( $\mathcal{R}$ -matrices, quantum Casimirs, the tensor product rule in (3.1)) and a Schur-Weyl duality theorem (Theorem 3.3 below) for the affine Hecke algebra.

The cyclotomic Hecke algebra  $H_k(u_1, \dots, u_r; q)$

Let  $u_1, \dots, u_r \in \mathbb{C}$  and  $q \in \mathbb{C}^*$ . The cyclotomic Hecke algebra  $H_k(u_1, \dots, u_r; q)$  is the quotient of the affine Hecke algebra  $H_k$  by the ideal generated by the relation

$$(X_1 - u_1)(X_1 - u_2) \cdots (X_1 - u_r) = 0. \quad (2.4)$$

The algebra  $H_k(u_1, \dots, u_r; q)$  is a deformation of the group algebra of the complex reflection group  $G(r, 1, k) = (\mathbb{Z}/r\mathbb{Z}) \wr S_k$  and is of dimension  $\dim(H_k(u_1, \dots, u_r; q)) = r^k k!$ . These algebras were introduced by Ariki and Koike [AK]. Ariki has generalized the classical result of Gyoja and Uno [GU] and given precise conditions for the semisimplicity of the cyclotomic Hecke algebras.

**Theorem 2.5.** ([Ar]) The algebra  $H_k(u_1, \dots, u_r; q)$  is semisimple if and only if

$$q^{2d} u_i \neq u_j \text{ for all } -k < d < k, 1 \leq i < j \leq r, \quad \text{and} \quad [k]! \neq 0,$$

where  $[k]! = [1][2] \cdots [k]$  and  $[i] = 1 + q^2 + \cdots + q^{2(i-1)}$ .

*Proof.* Let us only explain the conversion between the statement in [Ar] and the statement here. This conversion is the same as in Remark 1.2. If  $\tilde{T}_i = qT_i$  then  $\tilde{T}_i^2 = q^2((q - q^{-1})T_i + 1) = (q^2 - 1)qT_i + q^2 = (q^2 - 1)\tilde{T}_i + q^2$ , which shows that our algebra is the same as Ariki's except with parameter  $q^2$ . ■

Ariki and Koike give a combinatorial construction of the irreducible representations of the cyclotomic Hecke algebra  $H_k(u_1, \dots, u_r; q)$  when it is semisimple. Define

$$\hat{H}_k^{(r)} = \{r\text{-tuples } \lambda = (\lambda^{(1)}, \dots, \lambda^{(r)}) \text{ of partitions with } k \text{ boxes total}\}. \quad (2.6)$$

Let  $\lambda \in \hat{H}_k^{(r)}$ . A *standard tableau of shape*  $\lambda$  is a filling of the boxes of  $\lambda$  with  $1, 2, \dots, k$  such that for each  $\lambda^{(i)}$ ,  $1 \leq i \leq r$ ,

- (a) the entries in the rows are increasing left to right, and
- (b) the entries in the columns are increasing top to bottom.

Let  $L(i)$  denote the box of  $L$  containing  $i$ , and define

$$CT(b) = u_i q^{2(c-r)}, \quad \text{if the box } b \text{ is in position } (r, c) \text{ of } \lambda^{(i)}. \quad (2.7)$$

**Theorem 2.8.** [AK, Theorem 3.7] If  $H_k(u_1, u_2, \dots, u_r; q)$  is semisimple its irreducible representations  $H^\lambda$ ,  $\lambda \in \hat{H}_k^{(r)}$ , are given by

$$H^\lambda = H^{(\lambda^{(1)}, \dots, \lambda^{(r)})} = \mathbb{C}\text{-span}\{v_L \mid L \text{ is a standard tableau of shape } \lambda\}$$

(so that the symbols  $v_L$  form a basis of the vector space  $H^\lambda$ ) with  $H_k(u_1, \dots, u_r; q)$ -action given by

$$X_i v_L = CT(L(i)) v_L, \quad \text{and} \quad T_i v_L = (T_i)_{LL} v_L + (q^{-1} + (T_i)_{LL}) v_{s_i L},$$

where

$$(T_i)_{LL} = \begin{cases} q, & \text{if } L(i) \text{ and } L(i+1) \text{ are in the same row,} \\ & \text{of } \lambda^{(j)} \text{ for some fixed } j, \\ -q^{-1}, & \text{if } L(i) \text{ and } L(i+1) \text{ are in the same column,} \\ & \text{of } \lambda^{(j)} \text{ for some fixed } j, \\ \frac{CT(L(i+1))(q - q^{-1})}{CT(L(i+1)) - CT(L(i))}, & \text{otherwise,} \end{cases}$$

$s_i L$  is the same as  $L$  except  $i$  and  $i+1$  are switched, and

$v_{s_i L} = 0$ , if  $s_i L$  is not a standard tableau.

It is interesting to note that Theorem 2.8 is almost an immediate consequence of Theorem 2.2.

The Iwahori-Hecke algebra  $H_k(u_1, u_2; q)$  of type  $B_k$

The Iwahori-Hecke algebra of type  $B_k$  is the cyclotomic Hecke algebra  $H_k(u_1, u_2; q)$ . Thus, for  $u_1, u_2 \in \mathbb{C}$  and  $q \in \mathbb{C}^*$ ,  $H_k(u_1, u_2; q)$  is the quotient of the affine Hecke algebra by the ideal generated by the relation

$$(X_1 - u_1)(X_1 - u_2) = 0. \quad (2.9)$$

The algebra  $H_k(1, -1; q)$  is the group algebra of the Weyl group of type  $B_k$  (the hyperoctahedral group of signed permutations).

In the case of the Iwahori-Hecke algebra  $H_k(u_1, u_2; q)$  of type  $B_k$  Theorem 2.8 is due to Hoefsmit [Ho]. When  $H_k(u_1, u_2; q)$  is semisimple Hoefsmit's construction of the irreducible representations of  $H_k(u_1, u_2; q)$  implies that, as  $H_{k-1}(u_1, u_2; q)$ -modules

$$\text{Res}_{H_{k-1}}^{H_k} H^\lambda = \bigoplus_{\lambda^-} H^{\lambda^-}, \quad (2.10)$$

where the sum runs over all pairs of partitions  $\lambda^-$  which are obtained from  $\lambda$  by removing a single box, and

$$H^{\lambda^-} = \mathbb{C}\text{-span} \left\{ v_L \mid \begin{array}{l} L \text{ is a standard tableau of shape } \lambda \\ \text{and } L^- \text{ has shape } \lambda^- \end{array} \right\}, \quad (2.11)$$


where  $L^-$  is the standard tableau with  $k-1$  boxes which is obtained by removing the entry  $k$  from  $L$ . The restriction rules (2.10) can be encoded in the *Bratteli diagram* for the sequence of algebras

$$H_1(u_1, u_2; q) \subseteq H_2(u_1, u_2; q) \subseteq H_3(u_1, u_2; q) \subseteq \cdots \quad (2.12)$$

i.e., the graph which has

$$\text{vertices on level } k \text{ indexed by } \lambda \in \hat{H}_k^{(2)} \quad \text{and} \quad \text{edges } \lambda \longleftrightarrow \lambda^-$$

if  $\lambda^-$  is obtained from  $\lambda$  by removing a single box. The first few rows of the Bratteli diagram for  $H_k(u_1, u_2; q)$  are displayed in Figure 1.

Figure 1. Bratteli Diagram for  $H_k(u_1, u_2; q)$ 

If  $H_k(u_1, u_2; q)$  is semisimple then

$$H_k(u_1, u_2; q) \cong \bigoplus_{\lambda \in \hat{B}_k} M_{d_\lambda}(\mathbb{C}), \quad (2.13)$$

where  $d_\lambda$  is the number of standard tableaux  $L$  of shape  $\lambda$ , and  $M_d(\mathbb{C})$  is the algebra of  $d \times d$  matrices with entries from  $\mathbb{C}$ .

The minimal ideals  $I^\lambda$  of  $H_k(u_1, u_2; q)$  are in one-to-one correspondence with the summands in (2.13). Let  $m < k$ , let  $I^\mu$  be a fixed minimal ideal of  $H_m(u_1, u_2; q)$  and define

$$\langle I^\mu \rangle_k \text{ is the ideal of } H_k(u_1, u_2; q) \text{ generated by } I^\mu$$

( $I^\mu \subseteq H_m(u_1, u_2; q) \subseteq H_k(u_1, u_2; q)$ ). The restriction rules (2.10) imply that

$$\langle I^\mu \rangle_k = \bigoplus_{\lambda \supseteq \mu} I^\lambda, \quad (2.14)$$

where the sum is over all pairs of partitions  $\lambda = (\lambda^{(1)}, \lambda^{(2)}) \in H_k^{(2)}$  which are obtained from  $\mu = (\mu^{(1)}, \mu^{(2)}) \in \hat{H}_m^{(2)}$  by adding  $(k - m)$  boxes.

The ideal  $I^{((1^2), \emptyset)}$

**Lemma 2.15.** Assume that  $H_2(u_1, u_2; q)$  is semisimple. The minimal ideal  $I^{((1^2), \emptyset)}$  of  $H_2(u_1, u_2; q)$  is generated by the element

$$p = \begin{cases} (X_1 - u_2)(X_2 - u_2)(X_2 - q^2 u_1), & \text{if } u_1 \neq 0, \\ (X_1 - u_2)(T_1 - q)(X_1 - u_2)(X_2 - u_2), & \text{if } u_1 = 0, \end{cases} \quad \text{where } X_2 = T_1 X_1 T_1.$$

*Proof.* Using the construction of the simple  $H_2(u_1, u_2; q)$ -modules in Theorem 2.8 it is not tedious to check that, when  $u_1 \neq 0$ ,

$$pv_L = \begin{cases} (u_1 - u_2)(q^{-2}u_1 - u_2)(q^{-2}u_1 - q^2u_1)v_L, & \text{if } L \text{ has shape } ((1^2), \emptyset), \\ 0, & \text{otherwise,} \end{cases}$$

and, when  $u_1 = 0$ ,

$$pv_L = \begin{cases} (0 - u_2)(-q^{-1} - q)(0 - u_2)(0 - u_2)v_L, & \text{if } L \text{ has shape } ((1^2), \emptyset), \\ 0, & \text{otherwise.} \end{cases}$$

Thus  $p$  is an element of the ideal  $I^{((1^2), \emptyset)}$ . Since  $I^{((1^2), \emptyset)}$  is a minimal ideal it is generated by any one of its (nonzero) elements. ■

The algebra  $A_k(u_1, u_2; q)$

Let  $u_1 \in \mathbb{C}$  and  $u_2, q \in \mathbb{C}^*$ . Let  $A_k(u_1, u_2; q)$  be the algebra given by generators

$$X_1 \quad \text{and} \quad T_1, T_2, \dots, T_{k-1}$$

and relations

$$\begin{aligned} (1) \quad & T_i T_j = T_j T_i, & |i - j| > 1, \\ (2) \quad & T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, & 1 \leq i \leq n - 2, \\ (3) \quad & T_i^2 = (q - q^{-1})T_i + q, & 1 \leq i \leq k - 1, \\ (4) \quad & X_1 T_1 X_1 T_1 = T_1 X_1 T_1 X_1, \\ (5) \quad & (X_1 - u_1)(X_1 - u_2) = 0, \\ (6) \quad & (X_1 - u_2)(X_2 - u_2)(X_2 - q^2 u_1) = 0, \quad \text{if } u_1 \neq 0, \\ & (X_1 - u_2)(T_1 - q)(X_1 - u_2)(X_2 - u_2), \quad \text{if } u_1 = 0, \quad \text{where } X_2 = T_1 X_1 T_1. \end{aligned}$$

Let

$$\hat{A}_k = \{(\lambda^{(1)}, \lambda^{(2)}) \in \hat{H}_k^{(2)} \mid \lambda^{(1)} \text{ has at most one row}\}. \quad (2.16)$$

**Theorem 2.17.** Assume  $H_k(u_1, u_2; q)$  is semisimple.

- (a)  $A_k(u_1, u_2; q)$  is semisimple.
- (b) As in (2.14) let  $\langle I^{((1^2), \emptyset)} \rangle_k$  be the ideal of  $H_k(u_1, u_2; q)$  generated by the minimal ideal  $I^{((1^2), \emptyset)}$  of  $H_2(u_1, u_2; q)$ . Then

$$A_k(u_1, u_2; q) \cong \frac{H_k(u_1, u_2; q)}{\langle I^{((1^2), \emptyset)} \rangle_k}.$$

- (c) As in (2.13) let  $d_\lambda$  denote the number of standard tableaux of shape  $\lambda = (\lambda^{(1)}, \lambda^{(2)})$  and  $M_d(\mathbb{C})$  the algebra of  $d \times d$  matrices with entries from  $\mathbb{C}$ . Then

$$A_k(u_1, u_2; q) \cong \bigoplus_{\lambda \in \hat{A}_k} M_{d_\lambda}(\mathbb{C}).$$

- (d) The irreducible  $A_k(u_1, u_2; q)$ -modules  $H^\lambda$ ,  $\lambda \in \hat{A}_k$ , are given by Hoefsmit's construction (Theorem 2.8).
- (e) The Bratteli diagram for the sequence of algebras  $A_1(u_1, u_2; q) \subseteq A_2(u_1, u_2; q) \subseteq \dots \subseteq A_k(u_1, u_2; q)$  has vertices on level  $m$  indexed by  $\lambda \in \hat{A}_m$  and edges  $\lambda \longleftrightarrow \lambda^-$  if  $\lambda^-$  is obtained from  $\lambda$  by removing a box. See Figure 2.

*Proof.* (a) follows from the fact that  $A_k(u_1, u_2; q)$  is a quotient of  $H_k(u_1, u_2; q)$ .

(b) By Lemma 2.15, the element  $p$  generates the ideal  $\langle I^{((1^2), \emptyset)} \rangle$  in  $H_k(u_1, u_2; q)$  and so this is a consequence of the definition of  $A_k(u_1, u_2; q)$ .

(c) By (2.14)

$$\langle I^{((1^2), \emptyset)} \rangle_k = \bigoplus_{\lambda \supseteq ((1^2), \emptyset)} I^\lambda, \quad (2.18)$$

and so, by (b) and (2.13), the simple components of  $A_k(u_1, u_2; q)$  are indexed by those elements of  $\lambda \in \hat{H}_k^{(2)}$  which do not contain  $((1^2), \emptyset)$ . These are exactly the elements of  $\hat{A}_k$ .

(d) and (e) are consequences of (b), (c) and (2.18). ■




Figure 2. Bratteli Diagram of  $A_k(u_1, u_2; q)$  and  $R_k(q)$

By construction it is clear that the Bratteli diagram for  $A_k(u_1, u_2; q)$  is a subgraph of the Bratteli diagram for  $H_k(u_1, u_2; q)$ . It is the subgraph which is obtained by removing all pairs of partitions  $\lambda = (\lambda^{(1)}, \lambda^{(2)})$  which appear in (2.18) (for all  $k$ ). Thus, it is the subgraph which is obtained by removing the vertex  $((1^2), \emptyset)$  and all its *descendants*, i.e., all pairs of partitions  $\lambda = (\lambda^{(1)}, \lambda^{(2)})$  which are obtained by adding boxes to  $((1^2), \emptyset)$ .

**Theorem 2.19.** *The algebra  $A_k(u_1, u_2; q)$  is semisimple if and only if*

$$q^{2d}u_1 \neq u_2 \text{ for all } -k < d < k, \quad \text{and} \quad [k]! \neq 0,$$

where  $[k]! = [1][2] \cdots [k]$  and  $[i] = 1 + q^2 + \cdots + q^{2(i-1)}$ .

*Proof.* Since  $A_k(u_1, u_2; q)$  is a quotient of  $H_k(u_1, u_2; q)$ , we know that  $A_k(u_1, u_2; q)$  is semisimple when  $H_k(u_1, u_2; q)$  is. Thus  $A_k(u_1, u_2; q)$  is semisimple when (a) and (b) hold.

The Iwahori-Hecke algebra  $H_k(q)$  of type  $A_{k-1}$  is the cyclotomic Hecke algebra  $H_k(1, 1; q)$ . Thus,  $H_k(q)$  is the quotient  $A_k(u_1, u_2; q)$  by the relation  $X_1 = u_1$ . By Theorem 2.5 (in this case originally due to Gyoja and Uno)  $H_k(q)$  is semisimple if and only if  $[k]! \neq 0$ . Since  $H_k(q)$  is a quotient of  $A_k(u_1, u_2; q)$ , the algebra  $A_k(u_1, u_2; q)$  is not semisimple when  $H_k(q)$  is not semisimple. Thus,  $A_k(u_1, u_2; q)$  is not semisimple when  $[k]! = 0$ .

If  $u_1 = u_2$  then the representation  $\rho: A_k(u_1, u_2; q) \rightarrow M_2(\mathbb{C})$  given by setting

$$\rho(X_1) = \begin{pmatrix} u_1 & 1 \\ 0 & u_1 \end{pmatrix} \quad \text{and} \quad \rho(T_i) = \begin{pmatrix} q & 0 \\ 0 & q \end{pmatrix}$$

is an indecomposable representation which is not irreducible. Thus  $A_k(u_1, u_1; q)$  is not semisimple.

The Tits deformation theorem (see [CR, (68.17)]) says that the algebra  $A_k(u_1, u_2; q)$  has the same structure for any choice of the parameters  $u_1, u_2$  for which it is semisimple. Assume that  $[k]! \neq 0$  and  $u_2 = q^{2d}u_1$ ,  $u_1 \neq 0$ . Let  $\lambda/\mu$  be the skew shape given by  $\lambda = (k-1, d)$  and  $\mu = (d-1)$  and define

$$CT(b) = u_1 q^{2(c-r)+2}, \quad \text{if } b \text{ is a box in position } (r, c) \text{ of } \lambda.$$

With these definitions the formulas in Theorem 2.2 define an  $H_k(u_1, q^{2d}u_1; q)$ -module  $H^{\lambda/\mu}$ . A check that

$$(X_1 - q^{2d}u_1)(X_2 - q^{2d}u_1)(X_2 - q^2u_1)v_L = 0,$$

for all standard tableaux  $L$  of shape  $\lambda/\mu$  shows that that  $H^{\lambda/\mu}$  is an  $A_k(u_1, q^{2d}u_1; q)$ -module. The standard proof (see [Ra, Theorem 4.1]) of Theorem 2.5 applies in this case to show that  $H^{\lambda/\mu}$  is an irreducible  $A_k(u_1, q^{2d}u_1; q)$ -module. It has dimension

$$\dim(H^{\lambda/\mu}) = (\text{the number of standard tableaux of shape } \lambda/\mu) = \binom{k}{d} - 1.$$

When we restrict this  $A_k(u_1, q^{2d}u_1; q)$ -module to  $H_k(q)$ , it is a direct sum of irreducible  $H_k(q)$ -modules indexed by partitions  $\nu \vdash k$  with multiplicity given by the classical Littlewood-Richardson coefficient  $c_{\mu\nu}^\lambda$  (see [Ra, Theorem 6.1]). Since  $\lambda = (k-1, d)$  and  $\mu = (d-1)$  we have  $c_{\mu\nu}^\lambda \neq 0$  only if  $\nu$  has  $\leq 2$  rows and  $c_{\mu\nu}^\lambda = 0$  when  $\nu = (k)$ . So  $H^{\lambda/\mu}$  is an irreducible  $A_k(u_1, q^{2d}u_1; q)$ -module such that upon restriction to  $H_k(q)$  is a direct sum of irreducible representations indexed by partitions with length  $\leq 2$ , and which does not contain the “trivial” representation of  $H_k(q)$ .

When  $A_k(u_1, u_2; q)$  is semisimple its irreducible representations  $H^\lambda$  are indexed by pairs of partitions  $\lambda = (\lambda^{(1)}, \lambda^{(2)})$  such that  $\lambda^{(1)}$  has at most one row. If  $\lambda^{(1)}$  has length  $r$ , then, on restriction to  $H_k(q)$ ,  $H^\lambda$  is a direct sum of irreducibles indexed by partitions  $\nu \vdash k$  with multiplicities  $c_{\lambda^{(1)}\lambda^{(2)}}^\nu$ . By the Pieri rule [Mac, (5.16)] the resulting  $\nu$  are those obtained by adding a horizontal strip of length  $r$  to  $\lambda^{(2)}$ . Only when  $\lambda^{(2)}$  has a single row will all the  $\nu$  have  $\leq 2$  rows and, in this case,  $c_{\lambda^{(1)}\lambda^{(2)}}^\nu = c_{(r), (k-r)}^\nu = 1$  for  $\nu = (k)$ . Thus, when  $A_k(u_1, u_2; q)$  is semisimple every irreducible representation which, on restriction to  $H_k(q)$ , decomposes as a direct sum of components indexed by partitions with  $\leq 2$  rows does contain the “trivial” representation of  $H_k(q)$ .

Thus, the Tits deformation theorem implies that  $A_k(u_1, q^{2d}u_1; q)$  is not semisimple. ■

**Remark 2.20.** It is interesting to note that the blob algebras (see [MW]) are also quotients of  $A_k(u_1, u_2; q)$ .

*The  $q$ -rook monoid algebras  $R_k(q)$*

The new presentation of the  $q$ -rook monoid given in Section 1 shows that

$$R_k(q) = A_k(0, 1; q),$$

and thus  $R_k(q)$  is a quotient of Iwahori-Hecke algebra  $H_n(0, 1; q)$  of type  $B_k$ .

**Corollary 2.21.**

- (a) *The  $q$ -rook monoid algebra  $R_k(q)$  is semisimple if and only if  $[k]_q! \neq 0$ .*
- (b) *If  $R_k(q)$  is semisimple then the irreducible representations of  $R_k(q)$  are indexed by  $\lambda \in \hat{A}_k$  (see (2.16)) and are given explicitly by the construction in Theorem 2.8.*

Part (a) of Corollary 2.21 is Theorem 2.19 applied to  $R_k(q)$  and (b) is Theorem 2.17(e) for  $R_k(q)$ . Part (a) was proved in a different way by Solomon [So4] and part (b) is the result of Halverson [Ha, Theorem 3.2] which was the catalyst for the results of this paper.

As in (2.10) it follows that, as  $R_{k-1}(q)$ -modules

$$\text{Res}_{R_{k-1}}^{R_k} R^\lambda = \bigoplus_{\lambda^-} R^{\lambda^-},$$

where the sum runs over all pairs of partitions  $\lambda^-$  which are obtained from  $\lambda$  by removing a single box, and

$$R^{\lambda^-} = \mathbb{C}\text{-span}\{v_L \mid L^- \text{ has shape } \lambda^-\},$$

where  $L^-$  is the standard tableau with  $k-1$  boxes which is obtained by removing the  $k$  from  $L$ . The first few rows of the Bratteli diagram for the sequence of algebras

$$R_1(q) \subseteq R_2(q) \subseteq R_3(q) \subseteq \cdots$$

are as displayed in Figure 2.

### 3. Schur-Weyl dualities

Let  $U_q \mathfrak{gl}(n)$  be the quantum group corresponding to  $GL_n(\mathbb{C})$ . This is the algebra given by generators

$$E_i, \quad F_i, \quad (1 \leq i < n), \quad \text{and} \quad q^{\pm \varepsilon_i}, \quad (1 \leq i \leq n),$$

with relations

$$\begin{aligned} q^{\varepsilon_i} q^{\varepsilon_j} &= q^{\varepsilon_j} q^{\varepsilon_i}, \quad q^{\varepsilon_i} q^{-\varepsilon_i} = q^{-\varepsilon_i} q^{\varepsilon_i} = 1, \\ q^{\varepsilon_i} e_j q^{-\varepsilon_i} &= \begin{cases} q^{-1} e_j, & \text{if } j = i-1, \\ q e_j, & \text{if } j = i, \\ e_j, & \text{otherwise,} \end{cases} \quad q^{\varepsilon_i} f_j q^{-\varepsilon_i} = \begin{cases} q f_j, & \text{if } j = i-1, \\ q^{-1} f_j, & \text{if } j = i, \\ f_j, & \text{otherwise,} \end{cases} \\ e_i f_j - f_j e_i &= \delta_{ij} \frac{q^{\varepsilon_i - \varepsilon_{i+1}} - q^{-(\varepsilon_i - \varepsilon_{i+1})}}{q - q^{-1}}, \end{aligned}$$

$$e_{i\pm 1} e_i^2 - (q + q^{-1}) e_i e_{i\pm 1} e_i + e_i^2 e_{i\pm 1} = 0, \quad f_{i\pm 1} f_i^2 - (q + q^{-1}) f_i f_{i\pm 1} f_i + f_i^2 f_{i\pm 1} = 0,$$

$$e_i e_j = e_j e_i, \quad f_i f_j = f_j f_i, \quad \text{if } |i - j| > 1.$$

Part of the data of a quantum group is an  $\mathcal{R}$ -matrix, which provides a canonical  $U_q \mathfrak{gl}(n)$ -module isomorphism

$$\check{R}_{MN}: M \otimes N \longrightarrow N \otimes M$$

for any two  $U_q \mathfrak{gl}(n)$ -modules  $M$  and  $N$ .

The irreducible polynomial representations  $L(\lambda)$  of  $U_q \mathfrak{gl}(n)$  are indexed by dominant integral weights  $\lambda \in L^+$  where

$$L = \sum_{i=1}^n \mathbb{Z} \varepsilon_i = \{\lambda = \lambda_1 \varepsilon_1 + \cdots + \lambda_n \varepsilon_n \mid \lambda_i \in \mathbb{Z}\},$$

and  $L^+ = \{\lambda \in L \mid \lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n\}$ . The elements of  $L^+$  can be identified with partitions  $\lambda$  with  $\leq n$  rows.

The irreducible representation

$$V = L(\varepsilon_1) \quad \text{has} \quad \dim(V) = n, \quad \text{and} \quad L(\mu) \otimes V \cong \bigoplus_{\mu^+} L(\mu^+) \quad (3.1)$$



as  $U_q \mathfrak{gl}(n)$ -modules, where the direct sum is over all partitions  $\mu^+$  which are obtained from  $\mu$  by adding a box. The  $U_q \mathfrak{gl}(n)$ -module  $V$  can be given explicitly as the vector space

$$V = \mathbb{C}\text{-span}\{v_1, \dots, v_n\}$$

(so that the symbols  $v_i$  form a basis of  $V$ ) with  $U_q \mathfrak{gl}(n)$ -action given by

$$\begin{aligned} e_i v_j &= \begin{cases} v_{j-1}, & \text{if } j = i + 1, \\ 0, & \text{if } j \neq i + 1, \end{cases} & f_i v_j &= \begin{cases} v_{j+1}, & \text{if } j = i, \\ 0, & \text{if } j \neq i, \end{cases} & \text{and} \\ q^{\pm \varepsilon_i} v_j &= \begin{cases} q^{\pm 1} v_j, & \text{if } j = i, \\ v_j, & \text{if } j \neq i. \end{cases} \end{aligned}$$

With this notation the  $R$ -matrix for  $V \otimes V$  is given explicitly by

$$\begin{aligned} \check{R}_{VV}: V \otimes V &\rightarrow V \otimes V, & \text{where} \\ \check{R}_{VV}(v_i \otimes v_j) &= \begin{cases} q v_j \otimes v_i, & \text{if } i = j, \\ v_j \otimes v_i, & \text{if } i > j, \\ v_j \otimes v_i + (q - q^{-1})(v_i \otimes v_j), & \text{if } i < j. \end{cases} \end{aligned} \quad (3.2)$$

*A Schur-Weyl duality for affine and cyclotomic Hecke algebras*

**Theorem 3.3.** (see [OR, Theorem 6.17ab and Theorem 6.18])

(a) For any  $\mu \in L^+$  there is an action of the affine Hecke algebra  $\tilde{H}_k$  on  $L(\mu) \otimes V^{\otimes k}$  given by  $\Phi: \tilde{H}_k \longrightarrow \text{End}(L(\mu) \otimes V^{\otimes k})$  where

$$\Phi(X_1) = \check{R}_{V, L(\mu)} \check{R}_{L(\mu), V} \otimes \text{id}_V^{\otimes(k-1)} \quad \text{and} \quad \Phi(T_i) = \text{id}_{L(\mu)} \otimes \text{id}_V^{\otimes(i-1)} \otimes \check{R}_{VV} \otimes \text{id}_V^{\otimes(k-i-1)}.$$

(b) The  $\tilde{H}_k$  action on  $L(\mu) \otimes V^{\otimes k}$  commutes with the  $U_q \mathfrak{gl}(n)$ -action and the map

$$\Phi: \tilde{H}_k \longrightarrow \text{End}_{U_q \mathfrak{gl}(n)}(L(\mu) \otimes V^{\otimes k}) \quad \text{is surjective.}$$

(c) As a  $(U_q \mathfrak{gl}(n), \tilde{H}_k)$  bimodule

$$L(\mu) \otimes V^{\otimes k} \cong \bigoplus_{\lambda} L(\lambda) \otimes H^{\lambda/\mu},$$

where the sum is over all partitions  $\lambda$  which are obtained from  $\mu$  by adding  $k$  boxes and  $H^{\lambda/\mu}$  is a simple  $\tilde{H}_k$ -module.

(d) The representation  $\Phi$  given in part (a) is a representation of the cyclotomic Hecke algebra  $H_k(u_1, \dots, u_r; q)$ , i.e.

$$\Phi: H_k(u_1, \dots, u_r; q) \longrightarrow \text{End}_{U_q \mathfrak{gl}(n)}(L(\mu) \otimes V^{\otimes k}),$$

for any (multi)set of parameters  $u_1, \dots, u_r$  containing the (multi)set of values  $CT(b)$  (defined in (2.1)) as  $b$  runs over the boxes which can be added to  $\mu$  (to get a partition).

**Remark 3.4.** The affine Hecke algebra module  $H^{\lambda/\mu}$  which appears in Theorem 3.3(c) is the same as the module  $H^{\lambda/\mu}$  constructed in Theorem 2.2.

*A Schur-Weyl duality for Iwahori-Hecke algebras of type B*

Suppose that  $\mu$  is a partition with two addable boxes, i.e.

$$\mu = \underbrace{\quad}_{\ell} = \ell^d, \quad \text{for some } 0 < d \leq n, \ell \in \mathbb{Z}_{\geq 0}.$$

**Proposition 3.5.** *Let  $\mu = \ell^d$ ,  $0 < d \leq n$ ,  $\ell \in \mathbb{Z}_{\geq 0}$ . Then Theorem 3.3(d) provides a Schur-Weyl duality for  $H_k(u_1, u_2; q)$  with  $u_1 = q^{2\ell}$  and  $u_2 = q^{2d}$ .*

*Proof.* One only needs to note that if  $b_1, b_2$  are the addable boxes of  $\mu$  and  $CT(b)$  is as defined in (2.1) then

$$u_1 = CT(b_1) = q^{2\ell}, \quad \text{and} \quad u_2 = CT(b_2) = q^{2d}. \quad \blacksquare$$

*A Schur-Weyl duality for  $A_n(u_1, u_2; q)$*

Keeping the notation of Proposition 3.5, consider the special case  $\ell = n - 1$  and  $d \geq k$ , so that

$$\mu = \underbrace{\quad}_{\ell} = \ell^{(n-1)}, \quad \text{for some } \ell \in \mathbb{Z}_{\geq 0}.$$

Then, as a  $(U_q \mathfrak{gl}(n), H_k(u_1, u_2; q))$  bimodule

$$L(\mu) \otimes V^{\otimes k} \cong \bigoplus_{\lambda} L(\lambda) \otimes H^{\lambda}, \quad \text{where} \quad \lambda = \begin{array}{|c|c|} \hline \mu & \lambda^{(2)} \\ \hline \lambda^{(1)} & \\ \hline \end{array} \quad (3.6)$$

and  $H^{\lambda}$  is a simple  $H_k(u_1, u_2; q)$  module indexed by a pair of partitions  $\lambda = (\lambda^{(1)}, \lambda^{(2)})$  with  $k$  boxes total and such that  $\lambda^{(1)}$  has at most one row. The following result shows that, in this case, the Schur-Weyl duality in Theorem 3.3 becomes a Schur-Weyl duality for the algebra

$$A_k(u_1, u_2; q), \quad \text{where} \quad u_1 = q^{2(n-1)} \quad \text{and} \quad u_2 = q^{2\ell}.$$

**Proposition 3.7.** *Let  $\mu = (n-1)^d$ . Then the  $\tilde{H}_k$  action on  $L(\mu) \otimes V^{\otimes k}$  which is given by Theorem 3.3 factors through the algebra  $A_k(u_1, u_2; q)$ , where  $u_1 = q^{2(n-1)}$  and  $u_2 = q^{2\ell}$ .*

*Proof.* By Theorem 3.3(d), the  $\tilde{H}_k$  action on  $L(\mu) \otimes V^{\otimes k}$  factors through the algebra  $H_k(u_1, u_2; q)$  where  $u_1 = q^{2(n-1)}$  and  $u_2 = q^{2\ell}$ . It remains to check that  $(X_1 - u_2)(X_2 - u_2)X_2 - q^2 u_1 = 0$  as operators on  $L(\mu) \otimes V^{\otimes k}$ . To do this it is sufficient to show that  $(X_1 - u_2)(X_2 - u_2)X_2 - q^2 u_1 = 0$  as operators on  $H^{\lambda}$  for each  $H^{\lambda}$  which appears in the decomposition (3.6). The  $H_k(u_1, u_2; q)$ -module  $H^{\lambda}$  has basis indexed by the standard tableaux  $L$  of shape  $\lambda$  and

$$(X_1 - u_2)(X_2 - u_2)(X_2 - q^2 u_1)v_L = (CT(L(1)) - u_2)(CT(L(2)) - u_2)(CT(L(2)) - q^2 u_1)v_L.$$

For each of the possible positions of the first two boxes of  $L$  at least one of the factors in the last product is 0. Thus  $(X_1 - u_2)(X_2 - u_2)X_2 - q^2 u_1 = 0$  as operators on  $H^{\lambda}$ .  $\blacksquare$

*Another Schur-Weyl duality for cyclotomic Hecke algebras*

Let  $m_1, \dots, m_r$  be positive integers such that  $m_1 + \dots + m_r = n$ . Then  $\mathfrak{g}_P = \mathfrak{gl}(m_1) \oplus \dots \oplus \mathfrak{gl}(m_r)$  is a Lie subalgebra of  $\mathfrak{gl}(n)$ , and correspondingly

$$U_P = U_q \mathfrak{gl}(m_1) \otimes \dots \otimes U_q \mathfrak{gl}(m_r) \quad \text{is a subalgebra of} \quad U_q \mathfrak{gl}(n).$$

There is a corresponding decomposition of the fundamental representation  $V$  of  $U_q \mathfrak{gl}(n)$  as a  $U_P$ -module:

$$V = V_1 \oplus \dots \oplus V_r, \quad \text{where} \quad \dim(V_j) = m_j,$$

and  $V_j$  is the fundamental representation for  $U_q \mathfrak{gl}(m_j)$ .

$$\text{If } v \in V_j, \quad \text{we write} \quad \deg(v) = j$$

and say that  $v$  is *homogeneous of degree*  $j$ .

Let  $u_1, \dots, u_r \in \mathbb{C}$  and define

$$d: V \rightarrow V \quad \text{by} \quad d(v) = u_j v, \quad \text{if } \deg(v) = j. \quad (3.8)$$

Recall the action of  $\check{R}_{VV}: V \otimes V \rightarrow V \otimes V$  as given in (3.2), define

$$\check{S}_{VV}: V \otimes V \rightarrow V \otimes V \quad \text{by} \quad \check{S}_{VV}(v \otimes w) = \begin{cases} \check{R}_{VV}(v \otimes w), & \text{if } \deg(v) = \deg(w), \\ w \otimes v, & \text{if } \deg(v) \neq \deg(w), \end{cases}$$

(for homogeneous  $v, w \in V$ ), and define  $d_i, R_i, S_i \in \text{End}(V^{\otimes k})$  by

$$\begin{aligned} d_i &= \text{id}_V^{\otimes(i-1)} \otimes d \otimes \text{id}_V^{(k-i)}, & 1 \leq i \leq k, \\ \check{R}_i &= \text{id}_V^{\otimes(i-1)} \otimes \check{R}_{VV} \otimes \text{id}_V^{(k-i-1)}, & 1 \leq i \leq k-1, \\ \check{S}_i &= \text{id}_V^{\otimes(i-1)} \otimes \check{S}_{VV} \otimes \text{id}_V^{(k-i-1)}, & 1 \leq i \leq k-1. \end{aligned} \quad (3.9)$$

**Theorem 3.10.** (Sakamoto-Shoji [SS])

(a) There is an action of  $H_k(u_1, u_2, \dots, u_r; q)$  on  $V^{\otimes k}$  given by  $\Phi_P: H_k(u_1, \dots, u_r; q) \rightarrow \text{End}(V^{\otimes k})$  where

$$\Phi_P(T_i) = \check{R}_i \quad \text{and} \quad \Phi_P(X_1) = \check{R}_1^{-1} \dots \check{R}_k^{-1} \check{S}_k \dots \check{S}_1 d_1.$$

(b) The action of  $H_k(u_1, \dots, u_r; q)$  commutes with the action of  $U_P$  on  $V^{\otimes k}$ ,

$$\text{i.e., } \Phi_P: H_k(u_1, \dots, u_r; q) \rightarrow \text{End}_{U_P}(V^{\otimes k}).$$

(c) As a  $(U_P, H_k(u_1, \dots, u_r; q))$ -bimodule

$$V^{\otimes k} \cong \bigoplus_{\lambda=(\lambda^{(1)}, \dots, \lambda^{(r)})} L_P(\lambda) \otimes H^\lambda,$$

where the sum is over all  $r$ -tuples  $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$  of partitions such that  $\ell(\lambda^{(j)}) \leq m_j$ ,  $L_P(\lambda)$  is the simple  $U_P$ -module given by

$$L_P(\lambda) = L^{(1)}(\lambda^{(1)}) \otimes \dots \otimes L^{(r)}(\lambda^{(r)}),$$

where  $L^{(j)}(\lambda^{(j)})$  is the simple  $U_q(\mathfrak{gl}(m_j))$ -module corresponding to the partition  $\lambda^{(j)}$ , and  $H^\lambda$  is a (not necessarily simple)  $H_k(u_1, \dots, u_r; q)$ -module.

**Remark 3.11.** The  $H_k(u_1, \dots, u_r; q)$ -module  $H^\lambda$  appearing in Theorem 3.10(c) is simple whenever  $H_k(u_1, \dots, u_r; q)$  is semisimple. In that case  $H^\lambda$  coincides with the  $H_k(u_1, \dots, u_r; q)$ -module constructed in Theorem 2.8.

### A Schur-Weyl duality for $R_k(q)$

Consider the case of Theorem 3.10 when

$$r = 2, \quad m_1 = 1, \quad m_2 = n, \quad u_1 = 0, \quad u_2 = 1.$$

Then

$$V = V_1 \oplus V_2 \quad \text{where} \quad V_1 = \mathbb{C}v_0, \quad \text{and} \quad V_2 = \mathbb{C}\text{-span}\{v_1, \dots, v_n\}.$$

Let us analyze the action of  $H_k(0, 1; q)$  on  $V^{\otimes k}$  as given by Sakamoto and Shoji. Since  $dv_0 = u_1v_0 = 0$ ,

$$\Phi_P(X_1)(v_0 \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) = 0,$$

and, for  $\ell > 0$ ,

$$\begin{aligned} \Phi_P(X_1)(v_\ell \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) &= \check{R}_1^{-1} \cdots \check{R}_{k-1}^{-1} \check{S}_{k-1} \cdots \check{S}_1 d_1(v_\ell \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) \\ &= \check{R}_1^{-1} \cdots \check{R}_{k-1}^{-1} \check{S}_{k-1} \cdots \check{S}_1(v_\ell \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) \\ &= \check{R}_1^{-1} \cdots \check{R}_{k-1}^{-1} \check{R}_{k-1} \cdots \check{R}_1(v_\ell \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) \\ &= v_\ell \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}, \end{aligned}$$

and thus  $\Phi_P(X_1) = d_1$ . This calculation shows that the Sakamoto-Shoji action of  $H_k(0, 1; q)$  coincides exactly with action for the Schur-Weyl duality for the  $q$ -rook monoid algebra  $R_k(q)$  in the form given by Halverson [Ha, Corollary 6.3].

## 4. References

- [AK] S. ARIKI AND K. KOIKE, *A Hecke algebra of  $(\mathbb{Z}/r\mathbb{Z}) \wr S_n$  and construction of its irreducible representations*, Adv. Math. **106** (1994), 216–243.
- [Ar] S. ARIKI, *On the semisimplicity of the Hecke algebra of  $(\mathbb{Z}/r\mathbb{Z}) \wr S_n$* , J. Algebra **169** (1994), 216–225.
- [Ch] I. CHEREDNIK, *A new interpretation of Gelfand-Tsetlin bases*, Duke Math. J. **54** (1987), 563–577.
- [CR] C. CURTIS AND I. REINER, *Methods of Representation Theory: With Applications to Finite Groups and Orders, Volume II*, Wiley, New York, 1987.
- [GU] A. GYOJA AND K. UNO, *On the semisimplicity of Hecke algebras*, J. Math. Soc. Japan **41** (1989), 75–79.
- [Ha] T. HALVERSON, *Representations of the  $q$ -rook monoid*, preprint (2001).
- [Ho] P. N. HOEFSMIT, *Representations of Hecke algebras of finite groups with BN-pairs of classical type*, Thesis, Univ. of British Columbia, 1974.
- [Mac] I. G. MACDONALD, *Symmetric Functions and Hall polynomials*, Second edition, Oxford University Press, New York, 1995.
- [MW] P. P. MARTIN AND D. WOODCOCK, *On the structure of the blob algebra*, J. Algebra **225** (2000), 957–988.
- [Mu1] W. D. MUNN, *Matrix representations of semigroups*, Proc. Camb. Phil. Soc., **53** (1957), 5–12.

- [Mu2] W. D. MUNN, *The characters of the symmetric inverse semigroup*, Proc. Camb. Phil. Soc., **53** (1957), 13-18.
- [OR] R. ORELLANA AND A. RAM, *Affine braids, Markov traces and the category  $\mathcal{O}$* , preprint (2001).
- [Ra] A. RAM, *Skew shape representations are irreducible*, preprint (1998).
- [So1] L. SOLOMON, *The Bruhat decomposition, Tits system and Iwahori ring for the monoid of matrices over a finite field*, Geom. Dedicata **36** (1990), 15–49.
- [So2] L. SOLOMON, *Representations of the rook monoid*, J. Algebra, to appear.
- [So3] L. SOLOMON, Abstract No. 900-16-169, Abstracts Presented to the American Math. Soc., Vol. 16, No. 2, Spring 1995.
- [So4] L. SOLOMON, *The Iwahori algebra of  $\mathbf{M}_n(\mathbf{F}_q)$ , a presentation and a representation on tensor space*, preprint (2001).
- [SS] S. SAKAMOTO AND T. SHOJI, *Schur-Weyl reciprocity for Ariki-Koike algebras*, J. Algebra **221** (1999), 293-314.
- [Yg] A. YOUNG, *On quantitative substitutional analysis VI*, Proc. London Math. Soc. **31** (1931), 253-289.