

# HOROSPHERICAL CAUCHY-RADON TRANSFORM ON COMPACT SYMMETRIC SPACES

SIMON GINDIKIN

**ABSTRACT.** Harmonic analysis on noncompact Riemannian symmetric spaces is in a sense equivalent to the theory of the horospherical transform. There are no horospheres on compact symmetric spaces, but we define a complex version of horospherical transform which plays a similar role for the harmonic analysis on them.

In the horospherical transform we integrate test functions along horospheres or, in other words, we integrate these functions with  $\delta$ -functions with supports on horospheres. The idea of the complex horospherical transform [Gi00] is to replace the  $\delta$ -function by the Cauchy kernel: more precisely, we take Cauchy kernels with singularities along complex horospheres without real points. Such a transform can work when the real transform either does not exist or has too big a kernel [Gi00, Gi01, Gi04]. We define such a transform here on compact symmetric spaces. Earlier we saw [Gi04] that it already gives a new interesting integral transform for the sphere  $S^n$ .

Let  $X = G/K$  be a compact Riemannian symmetric space;  $G$  here is a simply connected compact semisimple Lie group;  $K$  is its connected involutive subgroup. Let  $\mathfrak{g} = \mathfrak{k} + \mathfrak{n}$  be the corresponding involutive decomposition in the Lie algebras. Let  $X_n = G_n/K$  be the dual noncompact Riemannian symmetric space and  $\mathfrak{g}_n = \mathfrak{k} + i\mathfrak{n}$  be the corresponding decomposition in Lie algebras. Let  $\mathfrak{a}$  be a Cartan subspace in  $i\mathfrak{n}$ ,  $\Sigma_+$  be a system of positive (restricted) roots;  $(\beta_1, \dots, \beta_l)$  be a basis of unmultipliable roots and  $(\mu_1, \dots, \mu_l)$  be the dual system of fundamental weights  $(\frac{\langle \mu_i, \beta_j \rangle}{\langle \beta_j, \beta_j \rangle} = \delta_{ij})$ . Then

$$\Lambda = \{\mu = \lambda_1 \mu_1 + \dots + \lambda_l \mu_l; \lambda_j \in \mathbb{Z}^+\}$$

is the set of highest (restricted) weights of spherical representations corresponding to the symmetric space  $X$ .

Let  $A = \exp(\mathfrak{a})$ ,  $A_I = \exp(i\mathfrak{a})$  be the corresponding commutative subgroups in  $G_n, G$  correspondingly. Let  $G_n = KAN$  be the Iwasawa decomposition, where  $N$  is the maximal unipotent subgroup in  $G$ . Let  $G_{\mathbb{C}}^0 = K_{\mathbb{C}} A_{\mathbb{C}} N_{\mathbb{C}}$  be the holomorphic complexification of this decomposition in  $G_{\mathbb{C}}$ . For a group  $H$  we denote through  $H_{\mathbb{C}}$  its complexification. The set  $G_{\mathbb{C}}^0$  is a Zariski open set in  $G_{\mathbb{C}}$ . Let us consider the projection  $a(g)$  of  $G_{\mathbb{C}}^0$  on the factor  $A_{\mathbb{C}}$ . We can consider this function also as a function on the symmetric Stein space  $X_{\mathbb{C}} = G_{\mathbb{C}}/K_{\mathbb{C}}$ .

For  $a \in A_{\mathbb{C}}$  we define

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

$$a_j = \exp(\mu_j(\log a)), \quad a^\mu = \prod_{1 \leq j \leq l} (a_j)^{\lambda_j}, \mu \in \Lambda.$$

For noncompact symmetric spaces  $X_n$  zonal spherical functions can be obtained by the averaging of  $a(x)^\mu$  along  $K$ -orbits (Harish-Chandra's integral representation). Clerc [Cl88] suggested in the case of compact symmetric spaces  $X$ , using the possibility of the holomorphic extension of zonal spherical functions on  $X_{\mathbb{C}}$ , to write the analogue of Harish-Chandra representation on  $X_{\mathbb{C}}$ . This representation can be restricted on the compact group  $G$  and the compact space  $X$  but there is involved the complex function  $a^\mu(x)$ . For the zonal spherical function  $\phi_\mu(x)$  with the center at the point  $x_0 \in X$  (corresponding to a fixed isotropy subgroup  $K$ ) we have

$$\phi_\mu(x; x_0) = \int_K a^\mu(xk) \nu(dk), x \in X_{\mathbb{C}}$$

where we integrate on the invariant measure on  $K$ . Here and everywhere below we integrate on a compact homogeneous manifold  $Y$  on the invariant measure  $\nu(dy)$  normalized by the condition  $\int_Y \nu(dy) = 1$ . The integrand is well-defined for almost all  $k \in K$ . Clerc used this representations for the computation of the asymptotic of zonal spherical functions on compact symmetric spaces similar to the famous Harish-Chandra's computation for noncompact spaces. The construction of the complex horospherical transform on  $X$  in this note continues this application of complex geometry to the harmonic analysis on compact symmetric spaces.

The functions  $a^\mu(x), x \in X_{\mathbb{C}}$ , are elements of eigen subspaces of the invariant differential operators which are  $N_{\mathbb{C}}$ -invariant. They sometimes are called zonal horospherical functions. Let us extend them for different choices of  $A, N$ . Let

$$h_\mu(x, g) = a^\mu(xg), x \in X_{\mathbb{C}}, g \in G_{\mathbb{C}}.$$

This function will not change under right multiplications on elements of  $N_{\mathbb{C}}$  as well as on elements of  $M_{\mathbb{C}}$  (as usual,  $M$  is the centralizer of  $A$  in  $K$ ). Let  $\Xi = G_{\mathbb{C}}/M_{\mathbb{C}}N_{\mathbb{C}}$ . We will realize its points as classes  $\zeta = (gM_{\mathbb{C}}N_{\mathbb{C}})$ . Then the function  $h_\mu$  will depend only on the class  $\zeta(g) \in \Xi$ . Correspondingly, we will denote it as  $h_\mu(x, \zeta), x \in X_{\mathbb{C}}, \zeta \in \Xi$ .

We will call  $\Xi$  the horospherical manifold. This name is connected with the possibility to realize points of  $\Xi$  as the horospheres on  $X_{\mathbb{C}}$ : nondegenerate orbits of the subgroup  $N_{\mathbb{C}}$  and its  $G_{\mathbb{C}}$ -translations.

Let  $F = G_{\mathbb{C}}/P_{\mathbb{C}} = G_{\mathbb{C}}/M_{\mathbb{C}}A_{\mathbb{C}}N_{\mathbb{C}} \cong G/MA$  be the corresponding flag manifold. There is a natural fibering  $\Xi \rightarrow F$  with fibers  $A_{\mathbb{C}}$ . The group  $G_{\mathbb{C}}$  acts on  $\Xi$  by the "left multiplications" ( $\zeta \mapsto g \cdot \zeta$ ) and  $A_{\mathbb{C}}$  by the "right multiplications" ( $\zeta \mapsto \zeta \cdot a$ ) and these actions commute. The existence of the right action of  $A_{\mathbb{C}}$  is a crucial circumstance for the horospherical transform. We have

$$h_\mu(x, \zeta \cdot a) = a^\mu h_\mu(x, \zeta), x \in X_{\mathbb{C}}, \zeta \in \Xi, a \in A_{\mathbb{C}}.$$

Choosing an initial point  $\zeta_0$  we can parameterize points  $\zeta \in \Xi$  as pairs  $\zeta = (a, u), a \in A_{\mathbb{C}}, u \in F$  such that the right multiplications on elements of  $A_{\mathbb{C}}$  act on the  $a$ -components of  $\zeta = (a, u)$ . We can define the horospheres  $\Omega(\zeta), \zeta \in \Xi$ , on  $X_{\mathbb{C}}$  by the equations

$$h_j(\cdot, \zeta) = 1, \quad 1 \leq j \leq l,$$

where  $h_j = h_{\mu_j}$ . Of course, then  $h_\mu(x, \zeta) = 1$  for all  $\mu$ .

We have on  $X$  [Cl88, Corollary, p.427]:

$$|a_j(x)| \geq 1, 1 \leq j \leq l$$

and the conditions  $|a_j(x)| = 1$  distinguish the image of the subgroup  $A_I$  in  $X$  ( $a_j(x) = 1$  for the initial point  $x_0$ ). Let  $A_+$  be the semigroup  $\{|a_j| < 1, 1 \leq j \leq l\}$  in  $A_{\mathbb{C}}$  and we define the domain  $\Xi_+$  in  $\Xi$  as  $\{\zeta = (a, u); a \in A_+\}$  (the  $(G \cdot \zeta_0 \cdot A_+)$ -orbit of the initial point  $\zeta_0$ ). So the domain  $\Xi_+$  is fibering over the flag domain  $F$  with the fibers  $A_+$ . We can reformulate the definition of  $\Xi_+$ :

$$\Xi_+ = \{\zeta \in \Xi; h_j(x, \zeta) < 1 \text{ for all } x \in X \text{ and } j\}.$$

In other words, the domain  $\Xi_+$  parameterizes horospheres  $\Omega(\zeta)$  on  $X_{\mathbb{C}}$  without real points (they do not intersect  $X$ ). The points of the boundary  $\zeta \in \partial\Xi_+$  correspond to the horospheres which intersect  $X$  in one point. It means that the boundary admits the fibering over the symmetric space  $X$  with the fiber  $\Xi(x)$  over  $x \in X$  defined by the equations

$$h_j(x, \zeta) = 1, 1 \leq j \leq l,$$

(or  $a(x \cdot \zeta) = e$ ). The fiber  $\Xi_x$  parameterizes the horospheres which intersect  $X$  in the point  $x$ . The fibers are isomorphic to  $K/M$ .

Now we are ready to give the basic definition. For  $f \in C^\infty(X)$  we define its horospherical Cauchy-Radon transform as

$$\hat{f}(\zeta) = \int_X \frac{f(x)}{\prod_{1 \leq j \leq l} (1 - h_j(x, \zeta))} \nu(dx), \zeta \in \Xi_+.$$

For  $\zeta \in \Xi_+$  the integrand has no singularities on  $X$  and is holomorphic on  $\zeta$ . Hence the function  $\hat{f}$  is holomorphic in the domain  $\Xi_+$ . The singular set of the integrand on  $X_{\mathbb{C}}$  has a very simple structure and has as the edge exactly the horosphere  $\Omega(\zeta)$ . For  $\zeta \in \partial\Xi_+$  the integral exists in the distribution sense and gives the regular boundary values of  $\hat{f}$ .

As usual in integral geometry, the principal problem is the problem of the inversion of this horospherical transform. Let  $D_j$  be the partial logarithmic derivative in the direction of the basic root  $\beta_j$  on  $A_{\mathbb{C}}$  and its transfer on  $\Xi$  relative to the right multiplications on  $A_{\mathbb{C}}$ . Then  $D_j h^\mu(x, \zeta) = \mu_j h^\mu(x, \zeta)$ . Let for any root  $\alpha \in \Sigma_+$ :

$$D(\alpha) = 1 + \frac{\langle \alpha, D \rangle}{\langle \alpha, \rho \rangle}$$

where  $\rho$  is the half-sum of the positive roots and

$$\mathcal{L} = \prod_{\alpha \in \Sigma_+} D(\alpha).$$

**Theorem.** *There is an inversion formula for the horospherical Cauchy-Radon transform*

$$f(x) = \int_{\Xi(x)} \mathcal{L}\hat{f}(\zeta)\nu(d\zeta).$$

Here we integrate the boundary values of the holomorphic function  $\hat{f}$  over the normalized invariant measure on  $K/M$ .

This result is a simple corollary of the Plancherel formula on compact symmetric spaces [Sh77,H94]. We define the spherical Fourier transform in "complex" form:

$$\tilde{f}(\zeta; \mu) = \int_X f(x) h^\mu(x, \zeta) \nu(dx), \quad \mu \in \Lambda, \zeta \in \Xi.$$

The function  $\tilde{f}(\zeta; \mu)$  is holomorphic on  $\zeta \in \Xi$  and satisfies the condition of homogeneity:

$$\hat{f}(\zeta \cdot a; \mu) = a^\mu(\hat{f}(\zeta; \mu)).$$

In other words,  $\tilde{f}(\zeta; \mu)$  is the section of the line bundle on  $F$  corresponding to  $\mu$ . So we have maps in the spaces of irreducible spherical representations in the flag realization.

Similarly, to noncompact symmetric spaces, the spherical Fourier transform and horospherical Cauchy-Radon transform are connected by the commutative Fourier transform, this time, by the discrete one. Since the right multiplications on elements  $a \in A_{\mathbb{C}}(\zeta \mapsto \zeta \cdot a)$  commute with the action of  $G$  we can decompose  $\hat{f}(\zeta)$  in the Fourier series relative to this action of Abelian compact group  $A_I$ . The components (which are invariant relative to the action of  $G$ ) are just  $\tilde{f}(\zeta; \mu)$ :

$$\hat{f}(\zeta) = \sum_{\mu \in \Lambda} \tilde{f}(\zeta; \mu).$$

It is the result of the direct decomposition the kernel in the definition of the horospherical transform. The inversion formula for the spherical Fourier transforms equivalent to the Plancherel formula on  $X$  is

$$f(x) = \sum_{\mu \in \Lambda} d(\mu) \int_{\Xi(x)} \tilde{f}(\zeta; \mu) \nu(d\zeta),$$

where for the dimension we have Weyl's formula

$$d(\mu) = \prod_{\alpha \in \Sigma_+} \frac{\langle \mu + \rho, \alpha \rangle}{\langle \rho, \alpha \rangle}.$$

It follows from the inversion formula on  $X$  [Sh77,H94] since

$$\int_{\Xi(x)} \tilde{f}(\zeta, \mu) \nu(d\zeta) = \int_X f(y) \phi(y; x) \nu(dy).$$

Here  $\phi(\cdot, x)$  is the zonal spherical function with the center  $x$  and it follows from Clerk's representation for zonal spherical functions. Now the inversion formula is

the consequence of the fact that the multiplication of the Fourier coefficients on the polynomial  $d(\mu)$  corresponds to the action of the differential operator  $\mathcal{L}$ .

In this note we obtain the results on horospherical transform as a reformulation of known facts of the harmonic analysis on compact symmetric spaces. Our aim here is to emphasize that formulas which we receive for the complex horospherical transform on compact symmetric spaces are completely similar to the formulas for the horospherical transform on noncompact Riemannian symmetric spaces. In our next publication we will obtain them by direct methods of integral geometry and it gives a new approach to harmonic analysis on compact homogeneous manifolds.

Let us consider a simplest example when  $X$  is the sphere  $S^n \subset \mathbb{R}^{n+1}$  [Gi04]:

$$\Delta(x) = x \cdot x = (x_1)^2 + \cdots + (x_{n+1})^2 = 1.$$

Let  $\mathbb{C}S^n$  be its complexification defined in  $\mathbb{C}^{n+1}$  by the equation  $\Delta(z) = 1$ . Then  $\Xi$  can be realized as the cone

$$\Delta(\zeta) = 0, \zeta \in \mathbb{C}^{n+1}, \zeta \neq 0,$$

and the flag manifold  $F$  as its projectivization in  $PC^n$ . The horospheres  $\Omega(\zeta), \zeta \in \Xi$ , can be defined as the intersection of  $S^n$  by the isotropic hyperplanes

$$\zeta \cdot z = \zeta_1 z_1 + \cdots + \zeta_{n+1} z_{n+1} = 1.$$

The parameters  $\zeta$  of horospheres without real points give the domain  $\Xi_+$ :

$$\Delta(\xi) = \Delta(\eta) < 1, \zeta = \xi + i\eta.$$

The fibers  $\Xi(x)$  in the fibering of  $\partial\Xi_+$  over  $S^n$  are  $\{\zeta = x + i\eta, x \cdot \eta = 0, \eta \cdot \eta = 1\}$ . Finally the differential operator in the inversion formula is  $\mathcal{L} = (1 + cD)^{n-1}$ , where  $D$  is the logarithmic differentiation along generators in the cone  $\Xi$  and  $c$  is a normalized constant.

*Remarks.* 1. The domain  $\Xi$  covers the compact complex manifold  $F$ . On  $F$  all holomorphic functions are constant, but on  $\Xi_+$  there is a lot of holomorphic functions (they separate points of  $\Xi_+$ ). The domain  $\Xi_+$  is not holomorphically complete since  $\Xi$  does not have this property. The extension reduces to the extension of the fibers  $A_{\mathbb{C}}$ . If  $F = \mathbb{C}P^n$  and  $z_0, \dots, z_n$  are homogeneous coordinates, then  $\Xi_+$  is the complex ball without the point:  $1 > |z_0|^2 + \cdots + |z_n|^2 > 0$  and all holomorphic functions in  $\Xi_+$  extend on the ball. In the example of the sphere, which we considered earlier, the natural extension of  $\Xi_+$  has the singularity.

It is simple to construct a Hardy type norm in  $H(\Xi_+)$  corresponding to the action of  $A_+$  and then we will have the model of spherical series of representations for  $X$ .

2. The horospherical transform and its inversion connect functions on real manifold  $X$  and holomorphic functions in  $\Xi_+$ . The natural understanding of this situation lies in the consideration of hyperfunctions on  $X$  as  $\bar{\partial}$ -cohomology classes of  $H^{(m)}(X_{\mathbb{C}} \setminus X)$ , where  $m = n - l, n = \dim X, l = \text{rank } X$  (the dimension of horospheres). Then we can interpret the horospherical transform as a version of the Penrose transform: we integrate cohomology along complex horospheres in  $X_{\mathbb{C}} \setminus X$ . A complication is that horospheres are not cycles and we need to be careful when

integrating cohomology. It would be interesting to define natural subspaces of cohomology in  $X_{\mathbb{C}} \setminus X$  and holomorphic functions in  $\Xi_+$  which can be intertwined. Intermediate Paley-Wiener theorems can be also interesting.

To receive the cohomological interpretation of the inverse horospherical transform we need to appeal to holomorphic language for  $\bar{\partial}$ -cohomology from [EGW95]. For each  $u \in F$  we take the union  $D(u)$  of "parallel" horospheres parameterizing by points in  $\Xi_+$  lying over  $u$ . We have the covering of  $X_{\mathbb{C}} \setminus X$  by the Stein domains  $D(u)$ ,  $u \in F$ . This covering satisfies the conditions from [EGW95] and we can construct Dolbeault cohomology using the complex of holomorphic forms  $\varphi(z|u, du)$  on  $F$  depending holomorphically on parameters  $z \in D(u)$ . We can extend the integrand in the inversion formula as such kind of a form.

There is another possibility interesting from point of view of complex analysis: to consider the horospherical Cauchy-Radon transform as the intertwining operator between holomorphic functions on whole manifolds  $X_{\mathbb{C}}$  and  $\Xi$ . We will discuss this operator in another paper.

3. When we want to construct the inverse horospherical transform we seek an integral operator on the space of all in a sense functions on the manifold of horospheres. In our case they are holomorphic functions. The restrictions of this operator on subspaces of sections of line bundles on  $F$ , corresponding to irreducible representations, give eigen functions of invariant differential functions (spherical polynomials). It is a Poisson's type integral. For the sphere such formulas go back to Maxwell. It would be interesting to deliberate the analogy with noncompact case. In a sense the flag manifold  $F$  plays the role of the "complex" boundary of  $X$ .

#### REFERENCES

- [Gi00] S.Gindikin, *Integral Geometry on  $SL(2; \mathbb{R})$* , Math.Res.Letters (2000), 417-432.
- [Gi02] S. Gindikin, *An analytic separation of series of representations for  $SL(2; \mathbb{R})$* , Moscow Math. J. **2** (2002), no. 4, 1-11.
- [Gi04] S.Gindikin, *Complex horospherical transform on real sphere* (2004).
- [Cl88] J.-L.Clerc, *Fonctions sphériques des espaces symétriques compacts*, Trans.AMS **306** (1988), 421-431.
- [EGW95] M.G.Eastwood, S.G.Gindikin, and H.-W.Wong, *Holomorphic realization of  $\bar{\partial}$ -cohomology and constructions of representations*, J.Geom.&Phys. **17** (1995), 231-244..
- [H94] S.Helgasson, *Geometric analysis on symmetric spaces*, AMS, 1994.
- [Sh77] T.Sherman, *Fourier analysis on compact symmetric spaces*, Bulletin AMS (1977), 73-144.

DEPARTM. OF MATH., HILL CENTER, RUTGERS UNIVERSITY, 110 FRELINGHYSEN ROAD,  
PISCATAWAY, NJ 08854-8019

E-mail address: gindikin@math.rutgers.edu