

In order to deliver a personalised, responsive service and to improve the site, we remember and store information about how you use it. This is done using simple text files called cookies which sit on your computer. By continuing to use this site and access its features, you are consenting to our use of cookies. To find out more about the way BioTechniques uses cookies please go to our [‘Cookie Policy’](#) page.



In order to deliver a personalised, responsive service and to improve the site, we remember and store information about how you use it. This is done using simple text files called cookies which sit on your computer. By continuing to use this site and access its features, you are consenting to our use of cookies. To find out more about the way BioTechniques uses cookies please go to our [‘Cookie Policy’](#) page.



 **EXCLUSIVE OFFER**

Purchase 2 STR Profiling Services and receive a 3rd service free!

START TODAY ▶



Restrictions may apply

eCLIPsing CLIP: RNA-Protein Interactions Revealed

04/15/2016
Josh P. Roberts
Hundreds of proteins can bind RNA, but finding out which ones bind where and what they do once they bind has been a challenge. That job just got a lot more manageable. Learn more...

 Like 8

 Tweet

 G+

 21 Share

 21 Share

 Tweet

 G+


There is [more to genetic regulation](#) than just [transcriptional control](#). As researchers become increasingly aware of the diverse roles played by RNA, the search for interactions between these nucleic acids and [RNA-binding proteins](#) (RBPs) takes on even greater importance. Hundreds of RBPs are known—many of which have been implicated in a variety of diseases such as [neurodegeneration](#), [autoimmune defects](#), and [cancer](#)—but most of their targets remain to be discovered. Finding sequences where RBPs bind can give some insight into the mechanisms by which they manage the trafficking, processing, and sometimes translation of RNAs both big and small.

State-of-the-art methods to discover where RBPs bind rely on crosslinking and immunoprecipitation (CLIP) of the ribonucleoprotein complex, followed by sequencing. Such protocols are technically challenging, require radioactivity, have a high experimental failure rate, and generate a high percentage of duplicate reads. “When performing large-scale CLIP experiments as part of the [ENCODE](#) consortium, we determined that the current methods were not robust enough to profile hundreds of factors,” related [Eric Van Nostrand](#) and [Gene Yeo](#) from the [University of California, San Diego](#) by email. They and their colleagues created enhanced CLIP (eCLIP) to solve these problems.

eCLIP borrows its 2-step multiplex library preparation technique from [ChIP-seq](#). First, the researchers ligate an indexed 3’ adapter (which allows for subsequent pooling) to the RNA fragments attached to immunoprecipitation beads. They then separate the complexes by electrophoresis and digest away the protein. After reverse transcribing the RNA, they ligate a second 3’ adapter containing an inline random 5- or 10-mer to the now single-stranded DNA before PCR amplification.

The fact that reverse transcription often terminates at the site of cross-linking allows the researchers to obtain nucleotide resolution. In the case of identical sequencing reads, the random single-stranded DNA adapter sequence will distinguish unique fragments from PCR duplicates, allowing the latter to be discarded, thus making the number of mapped reads more quantitative and minimizing wasted sequencing. In addition, eCLIP adds a size-matched pre-immunoprecipitation control for non-specific background and inherent biases.

These modifications mean that eCLIP requires about 1000-fold less amplification than its predecessors to achieve a higher percentage of useable reads and generate libraries with enough



- [A Diagnostic and Scientific Odyssey](#)
- [The Rules of Attraction for Double-Stranded DNA](#)
- [Gene Regulation Analysis Gets Loopy](#)

Reference

Keywords: ecLIP CLIP RNA-protein interactions RNA binding transcription RBP RNA-binding proteins neurodegeneration autoimmune defects cancer crosslinking immunoprecipitation ENCODE consortium ENCODE Eric Van Nostrand Gene Yeo University of California San Diego cross-linking Nature Methods



feedback