




ACUTE, REGIONAL ANESTHESIOLOGY & PERIOPERATIVE PAIN SECTION

Predictors of Persistent Post-Surgical Pain Following Total Knee Arthroplasty: A Systematic Review and Meta-Analysis of Observational Studies

Vahid Ashoorion , MD PhD,* Behnam Sadeghirad, PharmD, MPH, PhD,[†] Li Wang , PhD,^{‡,§} Atefeh Noori, MSc, PhD,[¶] Meisam Abdar, MD, MSc (cand),[¶] Yechan Kim, MSc, MD (cand),[¶] Yaping Chang, PhD,^{||} Nadia Rehman, BDS, MS (cand),** Luciane C. Lopes, ScD, MS,^{††} Rachel J. Couban, MA, MIST,^{‡‡} Mahmood Aminilari , MA, PhD,[¶] Alireza Malektojari, MD (cand),^{§§} Sara Ghazizadeh, MD (cand),^{§§} Yasir Rehman, MD, PhD (cand),**^{¶¶} Mehdi Ghasemi, MD,^{||} Anthony Adili, MD,^{||||} Gordon H. Guyatt, MD, MSc[¶] and Jason W. Busse , DC, PhD***.†††

*The Michael G. DeGroote Centre for Medicinal Cannabis Research, The Michael G. DeGroote National Pain Centre, McMaster University, Hamilton, Ontario, Canada; [†]The Michael G. DeGroote National Pain Centre, Department of Anesthesia, Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada; [‡]The Michael G. DeGroote National Pain Centre, Department of Anesthesia, McMaster University, Hamilton, Ontario, Canada; [§]Chinese Cochrane Centre, West China Hospital, Sichuan University, Chengdu, China; [¶]Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada; ^{||}Faculty of Medicine, University of Western Ontario, London, Ontario, Canada; ^{||}OrthoEvidence, Burlington, Ontario, Canada; ^{**}Canadian Academy of Osteopathy, Hamilton, Ontario, Canada; ^{††}Graduate Program in Pharmaceutical Sciences, University of Sorocaba, Sorocaba, State of São Paulo, Brazil; ^{‡‡}The Michael G. DeGroote National Pain Centre, McMaster University, Hamilton, Ontario, Canada; ^{§§}Hormozgan University of Medical Sciences, Bandarabbas, Iran; ^{¶¶}The Michael G. DeGroote National Pain Centre, Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada; ^{||||}Isfahan University of Medical Sciences, Isfahan, Iran; ^{***}The Michael G. DeGroote Centre for Medicinal Cannabis Research, The Michael G. DeGroote National Pain Centre, Department of Anesthesia, Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada; ^{†††}The Chronic Pain Centre of Excellence for Canadian Veterans, Hamilton, Ontario, Canada

Correspondence to: Jason W. Busse, DC, PhD, Department of Anesthesia, Michael G. DeGroote School of Medicine, McMaster University, HSC-2V9, 1280 Main St. West, Hamilton L8S 4K1, Canada. Tel: 905-525-9140 (x21731); Fax: 905-523-1224; E-mail: bussejw@mcmaster.ca.

Funding sources: This was an unfunded study. Dr. Ashoorion was supported by a Michael G. DeGroote Institute for Pain Research and Care post-doctoral fellowship.

Conflicts of interest: The corresponding author confirms on behalf of all authors that there has been no conflict of interest that might raise the question of bias in the work reported or in the conclusions, implications, or opinions stated.

Disclosures: Reproducible Research Statement: Study protocol: Available online (PROSPERO: CRD42018104968). Statistical code: Available on request from Dr. Ashoorion (e-mail, ashooriv@mcmaster.ca) Data set: Not applicable.

Systematic review registration: PROSPERO: CRD42018065943.

Received on 14 February 2022; revised on 8 September 2022; Accepted on 15 September 2022

Abstract

Objective. Approximately one in four total knee replacement patients develop persistent pain. Identification of those at higher risk could help inform optimal management. **Methods.** We searched MEDLINE, EMBASE, CINAHL, AMED, SPORTDiscus, and PsycINFO for observational studies that explored the association between risk factors and persistent pain (≥ 3 months) after total knee replacement. We pooled estimates of association for all independent variables reported by > 1 study. **Results.** Thirty studies (26,517 patients) reported the association of 151 independent variables with persistent pain after knee replacement. High certainty evidence demonstrated an increased risk of persistent

pain with pain catastrophizing (absolute risk increase [ARI] 23%, 95% confidence interval [CI] 12 to 35), younger age (ARI for every 10-year decrement from age 80, 4%, 95% CI 2 to 6), and moderate-to-severe acute post-operative pain (ARI 30%, 95% CI 20 to 39). Moderate certainty evidence suggested an association with female sex (ARI 7%, 95% CI 3 to 11) and higher pre-operative pain (ARI 35%, 95% CI 7 to 58). Studies did not adjust for both peri-operative pain severity and pain catastrophizing, which are unlikely to be independent. High to moderate certainty evidence demonstrated no association with pre-operative range of motion, body mass index, bilateral or unilateral knee replacement, and American Society of Anesthesiologists score. **Conclusions.** Rigorously conducted observational studies are required to establish the relative importance of higher levels of peri-operative pain and pain catastrophizing with persistent pain after knee replacement surgery.

Key Words: Chronic Pain; Meta-Analysis; Prognosis; Risk Factors; Systematic Review; Total Knee Replacement

Introduction

Total knee arthroplasty (TKA) is commonly performed for patients with advanced osteoarthritis who find insufficient relief with non-operative treatment [1]. The UK National Joint Registry recorded 312,167 knee replacement surgeries between 2017 and 2019 [2]; however, the prevalence of osteoarthritis is rising due to high rates of obesity and increasing life expectancy in western societies [3]. A modelling study using data from the UK National Joint Registry estimated that 186,302 patients would require primary knee replacement in 2030 [4].

The US Nationwide Inpatient Sample estimates that 1.3M Americans will undergo knee replacement in 2025, and that this figure will increase to 1.9M by 2030 [5].

Pain and consequent functional limitations are the primary indication for knee replacement, and patients anticipate that surgery will provide relief [6]; however, approximately 25% of patients report persistent post-surgical pain after TKA with higher rates after revision surgery [7, 8]. In 2017, the International Association for the study of Pain highlighted the importance of identifying surgical patients at high risk for the development of persistent pain to optimize care [9]. The 2021 Canadian Pain Task Force report has similarly called for guidance in this area [10]. We conducted a systematic review and meta-analysis of observational studies to identify predictors of persistent pain following TKA.

Methods

We followed the Meta-Analysis of Observational Studies in Epidemiology (MOOSE) statement [11], registered our review (PROSPERO Identifier: CRD42018065943), and published our protocol [12].

Data Sources and Searches

An academic librarian developed our search strategies (Supplementary Data eTable 1) that we implemented in MEDLINE, EMBASE, CINAHL, AMED, SPORTDiscus, and PsycINFO, from inception to April 30, 2021. We reviewed reference lists of eligible studies and related reviews for additional potentially eligible articles.

Eligibility Criteria and Study Selection

We included cohort and case-control studies in any language that: (1) enrolled adults (≥ 18 years) who received knee replacement surgery and (2) investigated, in an adjusted analysis, risk factors for persistent post-surgical pain after total knee replacement. Specifically, we required that eligible primary studies report an adjusted logistic regression model exploring predictors of persistent pain following TKA. We excluded randomized trials as strict eligibility criteria may systematically exclude patients with potentially important prognostic factors. We compared the definition of persistent post-surgical pain used in eligible studies with the definition introduced by the World Health Organization in 2015: (1) pain that began after surgery or a tissue trauma is experienced, (2) pain is in an area of preceding surgery, (3) pain has persisted for ≥ 3 months after surgery, and (4) the pain is not better explained by an infection, a malignancy, a pre-existing condition or other alternative cause [13]. We excluded conference abstracts.

Studies were ineligible if their predictive model(s) included significant associations with variables collected at the same time the outcome was measured. In such instances, the status of the predictor may be a result, rather than a cause, of persistent pain. If study populations overlapped by $>50\%$ among eligible articles, we included only the study with the largest sample size, longest follow-up, or larger number of independent factors explored in their regression model. Pairs of reviewers screened titles, abstracts, and full-text articles of potentially eligible studies, independently and in duplicate. We used online systematic review software (DistillerSR, Evidence Partners, Ottawa, Canada; <http://systematic-review.net>) to facilitate literature screening.

Data Extraction

Pairs of reviewers extracted, independently and in duplicate, the following information from all eligible studies: (1) study characteristics; (2) study population features; (3) surgical details; and (4) adjusted measures of association with persistent post-surgical pain for all reported risk factors. Reviewers resolved disagreements by discussion or by consultation with an adjudicator when

required. When a study reported more than one regression model, we used the model with the largest number of risk factors. We contacted authors for clarification of eligibility, to request missing data, or verify information.

Risk of Bias Assessment

We used the following criteria to assess risk of bias among eligible studies: (1) representativeness of the study population; (2) validity of outcome assessment; (3) proportion of missing data ($\geq 20\%$ was considered high risk of bias); and (4) whether predictive models included, at minimum, age, sex, and a measure of disease severity. ([Supplementary Data eTable 2](#)) [14].

Data Synthesis

We converted all pain scales used to measure intensity of persistent post-surgical pain to a 10 cm visual analogue scale (VAS) [15, 16]. When age or body mass index (BMI) were reported as categorical variables in regression models, we assumed linearity and that the association across categories were independent of each other. We then calculated the odds ratio (OR) and 95% confidence interval (CI) for each category using Bucher's approach and combined ORs using the inverse variance method to produce a single value [17, 18].

We pooled all independent factors that were reported by more than 1 study as an OR and associated 95% CI. We complemented relative measures of association (OR) with the absolute risk increase for each predictor amenable to meta-analysis. We used a baseline risk for persistent post-surgical pain of 25%, which we acquired from the low risk group in the study with the largest sample size among studies eligible for our review at low risk of bias [19]. We used the following formula to calculate absolute effects of all pooled predictors: Absolute effect = $((OR_{\text{predictor}} * \text{Baseline risk}) / ((1 - \text{Baseline risk}) + (OR_{\text{predictor}} * \text{Baseline risk}))) - \text{Baseline risk}$. See [Supplementary file](#) for a practical example ([eEquation 1](#)).

We used DerSimonian–Laird random-effects models for all meta-analyses [20]. It was decided a priori in the research protocol that random effects models would be used for all analyses to be conservative. Thus, random effects analyses was presented even for predictors for which the results were quite homogeneous and I^2 was low. We performed all analyses using Stata statistical software version 15.1 (StataCorp). All comparisons were two-tailed with a threshold P of .05.

When individual studies did not provide data that allowed their inclusion in pooled estimates, we explored the consistency of association between pooled results and those. When studies explored predictors in which no pooling was possible, we labelled them as promising for future study if they met the following criteria: (1) a statistically significant association with persistent pain of $P \leq .01$, (2) a large magnitude of association ($OR \geq 2.0$ or ≤ 0.5), and (3) a sample size of ≥ 500 patients. We

were unable to pool predictors reported by a single study, or when two studies reported the same predictor but one as a binary variable and the other as a continuous variable.

Subgroup Analyses, Meta-Regression, and Sensitivity Analyses

We evaluated heterogeneity for all pooled estimates through visual inspection of forest plots and I^2 statistic was also provided for all figures, for use by readers. We explored three a priori hypotheses to explain variability between studies assuming larger associations with persistent pain and: (1) shorter duration of follow up, (2) lower pain threshold (i.e., pain vs no pain) rather than a higher threshold (e.g., no pain/mild pain vs moderate to severe pain), and (3) greater risk of bias on a criterion-by-criterion basis. We conducted subgroup analyses if there were two or more studies in a given subgroup and assessed credibility of significant subgroup effects using modified ICEMAN criteria [21].

We performed sensitivity analyses to explore the impact of imputing data for non-significant predictors and converting categorical to continuous data. Specifically, in studies that excluded predictors from their final adjusted analysis due to non-significant association in univariate analysis, or the magnitude of association was not reported due to non-significant association in the final adjusted model, we used an OR of 1 and imputed the associated variance using the hot deck approach [18, 22]. However, imputation requires assumptions that may or may not be met, and we therefore imputed data only when the proportion of trials not reporting measures of association was $< 25\%$ of the total number of studies contributing to a meta-analysis.

Certainty of Evidence

We used the GRADE (Grading of Recommendations, Assessment, Development and Evaluations) approach to assess the certainty of evidence for all pooled measures of association as high, moderate, low, or very low [23]. With GRADE, evidence from prognostic studies begins as high certainty but can be rated down for risk of bias, indirectness, imprecision, or publication bias [24]. If, in subgroup analysis for risk of bias, we did not find a significant association, we included all studies and did not rate down for risk of bias. If we found a credible subgroup effect for risk of bias, we pooled only low risk studies. When we could not conduct subgroup analysis due to small number of studies, we followed GRADE guidance for assessing certainty of evidence [25].

Given the high baseline risk of 25% for persistent pain after TKA, our clinical experts (A.A., M.R.) estimated that a 5% increase in absolute risk would be sufficient for clinicians to address modifiable risk factors, which can be directly targeted in an effort to prevent persistent pain. The same experts estimated that an absolute

difference in risk of 10% between groups at low and high risk for persistent pain would be sufficient for clinicians to selectively target nonmodifiable risk factors to identify high-risk candidates for additional intervention. Therefore, we rated down for imprecision if the 95% CI associated with the risk difference included 5% for modifiable risk factors, or 10% for nonmodifiable risk factors. When at least 10 studies were included in a meta-analysis, we assessed publication bias by visual assessment of funnel plots and Egger's test [23, 26]. When we were able to pool the same predictor as both a continuous and binary variable, we reported the result supported by higher certainty of evidence.

Results

We identified 9,652 unique records, of which 30 English reports met our eligibility criteria. (Supplementary Data eFigure 1, Supplementary Data eTable 3) We attempted contact with four authors to verify data; one responded [27], providing additional details on their regression model and dependent variable. Ten studies enrolled patients from North American [28–37], 12 from Europe [38–49], and the remainder from Australia [50, 51], New Zealand [52], and Asia [53–57]. (Supplementary Data eFigure 2). Definitions of persistent pain varied across studies (Supplementary Data eTable 4). All studies reported persistent pain at least 3 months after knee arthroplasty; however, no study explicitly reported if the three remaining World Health Organization's criteria for persistent post-surgical pain were met. The median sample size among eligible studies was 350 (interquartile range [IQR]: 245 to 617), and the median duration of follow-up was 13 months (IQR: 6 to 24 months). The median proportion of female patients among eligible studies was 67% (IQR: 59 to 70) and median age was 68 years (IQR: 67 to 69) (Supplementary Data eTable 4).

Risk of Bias

Among eligible studies, 73% (22 of 30) were at high risk of bias for at least one criterion. Two studies did not enroll a representative study population, four did not use a valid tool to assess persistent pain, 18 reported high loss to follow up, and the regression models reported in eight studies did not adjust for one of age, sex, or disease severity (Supplementary Data eTable 5). We detected no evidence of publication bias for sex (Supplementary Data eFigures 3), and Egger's test was non-significant ($P = .43$).

Prevalence and Intensity of Persistent Pain after TKA

The median prevalence of persistent pain after TKA among the 29 studies ($n = 13,863$) reporting this information was 25% (IQR: 14% to 33%); there was no credible subgroup effect based on the threshold used for

defining persistent pain (Supplementary Data eTable 11). The overall pooled pain intensity among TKA patients reporting persistent pain, among 6 studies ($n = 2,388$) that reported pain severity, was 6.1 cm on a 10 cm VAS (95% CI 5.4 to 6.8) (Supplementary Data eFigure 4).

Predictors of Persistent Pain after TKA

Thirty studies involving 26,517 patients reported the association of 151 independent variables with persistent pain following TKA. Twenty (67%) included only variables significant in univariate analysis in their final regression model, and 13 studies (43%) failed to present data for non-significant predictors in their adjusted analysis.

High certainty evidence demonstrated a significant association between three factors and persistent post-surgical pain after TKA: (1) younger age (4% absolute risk increase [ARI] for every 10-year decrement from age 80, 95% CI 1.7% to 6.4%, $P = .015$; six studies), (2) moderate-to-severe acute post-operative pain (29.5% ARI, 95% CI 20.2% to 38.5%, $P < .001$; three studies), and (3) pain catastrophizing (23% ARI, 95% CI 11.7% to 34.5%, $P < .001$; three studies). (Figures 1–3) Moderate-to-severe acute post-operative pain was defined as $\geq 4/10$ on a 10-point scale. All 3 studies contributing to the pooled measure of association for pain catastrophizing used the Pain Catastrophizing Scale (PCS), which generates scores from 0 (no catastrophizing) to 52 (severe catastrophizing). A single cutoff on the PCS has not been agreed upon [58, 59], and studies used either 16, 20, or 21 as their threshold for designating meaningful pain catastrophizing.

Moderate certainty evidence suggested an association with pre-operative pain (35% ARI, 95% CI 7.3% to 57.7%, $P = .010$; 5 studies), female sex (6.7% ARI, 95% CI 2.5% to 11.2%, $P = 0.001$; 10 studies), non-white race (9.5% ARI, 95% CI 0.4% to 20%, $P = .039$; two studies) and comorbid diabetes mellitus (11% ARI, 95% CI 1.1% to 22.3%, $P = 0.026$; 2 studies). (Figure 4, Supplementary Data eFigures 5–7, Table 1) Four studies [40, 50, 55, 60] reporting the association with pre-operative pain compared lower vs higher pain severity on a 10-point scale, and one study [42] explored no pain (0/10) vs any pain ($\geq 1/10$) at rest; only 22 patients did not report pain at rest before surgery in this study which resulted in very little variability to detect an association with persistent pain. The association between pre-operative pain and persistent pain was rated down to moderate certainty evidence based on inconsistency, due to the one study that used a low threshold for pain severity.

High certainty evidence showed no significant association between persistent pain and preoperative range of motion. (Supplementary Data eFigure 8, Table 1) Moderate certainty evidence suggested no association with pre-surgical BMI, bilateral vs unilateral TKA,




Figure 1. Meta-analysis of the association between every 10 years decrease from age 80 years old and persistent pain.




Figure 2. Meta-analysis of the association between acute post-operative pain and persistent post-surgical pain (binary predictor: acute moderate to severe post operative pain vs no to mild acute post operative pain).




Figure 3. Meta-analysis of the association between pain catastrophizing and persistent post-surgical pain (binary predictor: pain catastrophizing vs no pain catastrophizing).




Figure 4. Meta-analysis of the association between pre-operative pain and persistent pain (binary predictor: pre-operative pain vs no pre-operative pain).

Table 1. GRADE evidence profile predictors of persistent pain after total knee arthroplasty

GRADE Evidence Profile: Predictors of Persistent Pain after Total Knee Arthroplasty									
Predictor	Quality Assessment						Relative Effect OR (95% CI)	Anticipated Absolute Effect	
	Risk of Bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Overall		Baseline Risk	Risk Difference (95% CI)*
Age: Every 10 yr. decrease									
6,717 patients, (6 studies) Median follow-up 18 mo	No serious risk of bias ^a	No serious inconsistency	No serious Indirectness	No serious Imprecision	Undetected; only 6 studies	High	1.22 (1.09– 1.37)	25%	3.9% more (95% CI: 1.7% more to 6.4% more) patients experience persistent pain per 10 years decrease from age 80
Sex: Female vs Male									
7,632 patients, (10 studies) Median follow-up 14 mo	No Serious risk of bias ^a	No serious inconsistency	No serious Indirectness	Serious Imprecision ^b	Symmetric funnel plot; Egger test <i>P</i> = .430	Moderate	1.39 (1.14–1.7)	25%	Females experience 6.7% more (2.5% more to 11.2% more) persistent pain
BMI: Every 5-score Increase									
5,385 patients, (3 studies) Median follow-up 24 mo	No serious risk of bias ^a	No serious inconsistency	No serious Indirectness	Serious Imprecision ^d	Undetected; only 3 studies	Moderate	1.16 (1–1.35)	25%	2.9% more (95% CI: 0% to 6 % more) patients experience persistent pain for every 5-point increase in BMI
Pre-operative Pain in Knee (Threshold used was no vs some pain in 1 study, and lower vs higher pain in 4 studies ^c)									
601 patients, (5 studies) median follow-up 14 mo	No serious risk of bias ^a	Serious inconsistency	No serious Indirectness	No Serious Imprecision	Undetected; only 5 studies	Moderate	4.52 (1.43–14.29)	25%	35.1% more (7.3% more to 57.7% more) patients experience persistent pain

(continued)

GRADE Evidence Profile: Predictors of Persistent Pain after Total Knee Arthroplasty

Predictor	Quality Assessment						Relative Effect OR (95% CI)	Anticipated Absolute Effect	
	Risk of Bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Overall		Baseline Risk	Risk Difference (95% CI)*
Acute Post-operative Pain (Binary: Moderate/Severe pain vs No/Mild pain)									
935 patients, (3 studies) median follow up 6 mo	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	Undetected; only 3 studies	High	3.59 (2.47–5.21)	25%	29.5% more (95% CI: 20.2% more to 38.5% more) patients experience persistent pain
Pain Catastrophizing (threshold varies between 16 and 21 on pain catastrophizing scale 0–52, 0 [no catastrophizing] to 52 [severe catastrophizing])									
796 patients, (3 studies) median follow-up 12 mo	No serious risk of bias ^a	No serious inconsistency	No serious indirectness	No serious imprecision	Undetected; only 3 studies	High	2.77 (1.74–4.41)	25%	23% more (11.7% more to 34.5% more) patients experience persistent pain
Race (Non-White vs White)									
1,484 patients, (2 studies) median follow-up 9 mo	No serious risk of bias ^a	No serious inconsistency	No serious indirectness	Serious imprecision ^b	Undetected; only 4 studies	Moderate	1.58 (1.02–2.45)	25%	9.5% more (95% CI: 0.4% more to 20% more) patients experience persistent pain
Diabetes comorbidity									
824 patients, (2 studies) median follow-up 15 mo	No serious risk of bias ^a	No serious inconsistency	No serious indirectness	Serious imprecision ^b	Undetected; only 4 studies	Moderate	1.69 (1.06–2.69)	25%	11% more (95% CI: 1.11% more to 22.3% more) patients experience persistent pain
Bilateral vs Unilateral (Simultaneous or Staged Bilateral vs Unilateral)									
1,140 patients, (2 studies) median follow-up 8 mo	No serious risk of bias ^a	No serious inconsistency	No serious Indirectness	Serious imprecision	Undetected; only 2 studies	Moderate	0.74 (0.50–1.09)	25%	5.2% fewer (10.7% less to 1.7% more) patients experience persistent pain

(continued)

GRADE Evidence Profile: Predictors of Persistent Pain after Total Knee Arthroplasty

Predictor	Quality Assessment						Relative Effect OR (95% CI)	Anticipated Absolute Effect	
	Risk of Bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Overall		Baseline Risk	Risk Difference (95% CI)*
Preoperative Range of Motion (Change in 1 degree in range of motion, measured 2–4 week before surgery considered as non-modifiable factor)									
491 patients, (2 studies) range of follow-up 9 mo	No serious risk of bias	No serious Inconsistency	No serious Indirectness	No serious imprecision ^e	Undetected; only 3 studies	High	1.04 (0.98–1.10)	25%	0.7% more (0.4% less to 1.8% more) patients experience persistent pain
ASA III vs II									
823 patients, (2 studies) median follow-up 4.5 mo	No serious risk of bias ^a	No serious inconsistency	No serious indirectness	Serious imprecision	Undetected; only 2 studies	Moderate	0.60 (0.32–1.14)	25%	8.3% less (15.4% less to 2.5% more) patients experience persistent pain

^aCertainty of evidence was not rated down on the basis of risk of bias, because the subgroup analyses and meta-regression did not show a significant difference between each risk-of-bias component and the estimates of association.

^bCertainty of evidence was rated down on the basis of imprecision. Although the 95% CI for the pooled effect did not include a risk difference of 1 (no effect), clinical actions based on the estimate of the lower or upper boundary would change, according to the predefined threshold of $\geq 10\%$ for non-modifiable factors.

^cFour studies compared less vs more pre-operative pain (using thresholds from 2 to 6 on a 0–10 VAS, 10 = worst pain); one study compared no pre-operative pain vs any pain at rest.

^dThree studies reported different thresholds for pain catastrophizing: 16, 20, and 21, on the 0- to 52-point pain catastrophizing scale.

^eCertainty of evidence was not rated down on the basis of imprecision. Although the 95% CI for the pooled effect overlap a risk difference of 1 (no effect), clinical actions based on the estimate of the lower or upper boundary would not change, according to the predefined threshold of $\geq 10\%$ for non-modifiable factors.

*We used the following formula to calculate absolute effects of all pooled predictors: Absolute effect = $((OR_{\text{predictor}} * \text{Baseline risk}) / ((1 - \text{Baseline risk}) + (OR_{\text{predictor}} * \text{Baseline risk}))) - \text{Baseline risk}$. A practical example is provided in the [Supplementary File \(eEquation 1\)](#).

or American Society of Anesthesiology (ASA) score (Supplementary Data eFigures 9–11, and Table 1).

The results from seven studies that reported 1 or more of the 11 predictors that we subjected to meta-analysis but whose data could not be pooled were reasonably consistent with our pooled analyses (Supplementary Data eTable 6). Of the 134 factors that were assessed for an association with persistent pain but not amenable to meta-analysis; none met our criteria as promising for future study (Supplementary Data eTables 7 and 8).

Subgroup Analysis, Meta-Regression and Sensitivity Analysis

Our sensitivity analysis imputing unreported non-significant associations rendered non-white race and comorbid diabetes as no longer significantly associated with persistent post operation pain. (Supplementary Data eTable 9) No additional subgroup analysis or meta-regression was credible (Supplementary Data eTables 10–12, Supplementary Data eFigure 12).

Discussion

We found high certainty evidence that pain catastrophizing, younger age and moderate to severe acute postoperative pain are associated with persistent pain after TKA, and moderate certainty evidence for an association with female sex and moderate to severe pre-operative pain. We also found moderate certainty evidence for an association with non-white race and comorbid diabetes; however, these associations became non-significant when imputing results for studies that found no association but reported no data. The strongest of these associations was for pain catastrophizing and greater peri-operative pain, with an absolute increase in the risk for persistent pain ranging from 23% to 35%. High to moderate certainty evidence suggests no association with BMI, preoperative range of motion, bilateral vs unilateral TKA, or ASA score. Investigators have tested 134 additional predictors that could not be statistically pooled, of which none clearly warrant additional study.

The most recent meta-analysis that explored risk factors for persistent pain after knee arthroplasty identified 32 studies, of which 16 were included in our review [61] (Supplementary Data eTable 13). Of the 16 not included in our review, seven reported only unadjusted associations with persistent pain [62–68], populations enrolled in seven overlapped with patients in other studies [7, 8, 69–73], and two studies were randomized trials [74, 75]. This prior review pooled measures of association as Fisher's Z and found that pain catastrophizing, preoperative pain, and presence of comorbidities were associated with persistent pain after TKA; better mental health before surgery had a protective effect [61]. They did not assess the certainty of evidence supporting these associations. We have confirmed and quantified the associations

of pre-operative pain and pain catastrophizing with persistent pain, and established three additional risk factors: younger age, female sex, and acute postoperative pain. We also identified moderate to high certainty evidence that BMI, bilateral vs unilateral TKA, ASA score, preoperative range of motion are not important predictors.

We found three associations in which the absolute increase in risk would be sufficient to suggest intervening: pre-operative (ARI 35%) and acute postoperative pain (ARI 30%), and pain catastrophizing (ARI 23%). However, these factors are unlikely to be independent as pain catastrophizing is associated with reporting higher levels of pain [76]. None of the studies that contributed to our pooled estimates of association for pre-operative or acute post-operative pain were adjusted for pain catastrophizing. Clarifying this issue may be important to informing whether promising interventions to reduce the risk of persistent pain after TKA should focus on behavioural modification, further optimization of peri-operative pain relief, or both.

Strengths and Limitations

Strengths of our review include explicit eligibility criteria and a comprehensive search that identified 14 studies that were not included in the most recent published systematic review exploring factors associated with persistent pain after TKA [61, 77–80]. We pooled measures of association for predictive factors, conducted sensitivity analyses imputing data for missing nonsignificant predictors, used the GRADE approach to appraise the certainty of evidence and, presented associations as both relative and absolute measures to optimize interpretation.

Our study has limitations. We pooled measures of association reported at the longest follow-up time, which ranged from 3 months to 7 years; however, we found no credible subgroup effects based on length of follow-up. We assessed heterogeneity of pooled measures of association with I^2 , and statistical tests of heterogeneity can be misleading when sample sizes are large and CIs are therefore narrow [81]; however, the number of patients enrolled in studies eligible for our review were typically modest with a median sample size of 350 (IQR: 245 to 617). We used the World Health Organization's criteria for the definition of persistent pain in this review; however, none of the included studies met all criteria and, as such, they may have overestimated the prevalence of persistent pain after TKA. Finally, the estimate of precision for the association with age and persistent pain from Singh et al.'s cohort of patients undergoing primary TKA is narrower than expected given their large sample size ($n = 4094$) [31]. This study reported a much lower event rate (8%) than other studies (e.g., 30% for Gungor et al.) [33], and reported ORs by age category which we then converted to a single estimate vs a reference of age ≥ 80 . Our imputation approach assumed the correlation between age categories was zero, which is a conservative

assumption that can exaggerate the variances resulting in a wider CI. This is a limitation of estimating the variance when a correlation coefficient is not available; however, our sensitivity analyses showed consistent results for an association with younger age and persistent pain when excluding converted estimates (see [Supplementary Data eTable 9](#)).

Conclusion

Higher pre- and acute post-operative pain, and pain catastrophizing, were associated with large increases in the risk of persistent post-surgical pain after TKA; however, these factors are unlikely to be independent. Rigorously conducted observational studies are required to establish the relative importance of higher levels of peri-operative pain and pain catastrophizing in the development of persistent pain after knee replacement surgery. These factors may represent high-yield targets to prevent the development of persistent post-surgical pain.

Acknowledgments

The authors thank Dr. Mehran Radi, MD, for guidance regarding selection of absolute risk increase thresholds for modifiable and non-modifiable factors, Dr. Stefan Schandelmaier, MD, PhD, for adjudicating one study published in Germany, and Mr. Pirooz Emami for screening citations and full text studies.

Dr. Mehran Radi, MD, Isfahan University of Medical Sciences, Isfahan, Iran.

Dr. Stefan Schandelmaier, MD, PhD Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada

Mr. Pirooz Emami MSc (Cand.), Department of Radiation Sciences, McMaster University, Hamilton, ON, Canada

Supplementary Data

[Supplementary Data](#) may be found online at *Pain Medicine* online.

References

- Singh JA, Lewallen DG. Cerebrovascular disease is associated with outcomes after total knee arthroplasty: A us total joint registry study. *J Arthroplasty* 2014;29(1):40–3.
- National Joint Registry, 17th Annual Report 2020. 2020. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK566660/> (accessed August 8, 2022).
- Dowsey MM, Nikpour M, Dieppe P, et al. Associations between pre-operative radiographic changes and outcomes after total knee joint replacement for osteoarthritis. *Clin Orthopaed Related Res* 2010;15:804–12.
- Patel A, Pavlou G, Mújica-Mota R, Toms A. The epidemiology of revision total knee and hip arthroplasty in England and Wales: A comparative analysis with projections for the United States. A study using the National Joint Registry dataset. *Bone Joint J* 2015;97-B (8):1076–81.
- Singh JA, Yu S, Chen L, Cleveland JD. Rates of total joint replacement in the United States: Future projections to 2020–2040 using the national inpatient sample. *J Rheumatol* 2019;46 (9):1134–40.
- Beswick AD, Wylde V, Gooberman-Hill R, Blom A, Dieppe P. What proportion of patients report long-term pain after total hip or knee replacement for osteoarthritis? A systematic review of Prospective studies in unselected patients. *BMJ Open* 2012;2 (1):e000435.
- Brander VA, Stulberg SD, Adams AD, et al. Ranawat award paper: Predicting total knee replacement pain. *Clin Orthopaed Relat Res* 2003;416:27–36.
- Wylde V, Hewlett S, Learmonth ID, Dieppe P. Persistent pain after joint replacement: Prevalence, sensory qualities, and post-operative determinants. *Pain* 2011;152(3):566–72.
- Carr DB, Morlion B. *Pain After Surgery: What Health-Care Professionals Should Know*. Washington, DC: IASP; 2017.
- Canadian Pain Task Force. An action plan for pain in Canada. 2021. Available at: https://www.canada.ca/en/health-canada/corporate/about-health-canada/public-engagement/external-advisory-bodies/canadian-pain-task-force/report-2021.html?utm_source=action-plan-report&utm_medium=partner&utm_content=en&utm_campaign=hc-sc-pain-task-force-21-22 (accessed August 8, 2022).
- Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: A proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283(15):2008–12.
- Ashoorion V, Sadeghirad B, Wang L, et al. Predictors of persistent postsurgical pain following total knee arthroplasty: A protocol for systematic review and meta-analysis. *Can J Pain* 2019;3 (2):10–5.
- Treede R-D, Rief W, Barke A, et al. A classification of chronic pain for ICD-11. *Pain* 2015;156(6):1003–7.
- Randolph A, Cook DJ, Guyatt GH. Prognosis. In: Guyatt GH, Rennie D, Meade MO, Cook DJ, eds. *Users' Guides to the Medical Literature*. New York, Chicago, San Francisco, Athens, London, Toronto: McGraw Hill; 2015:421–30.
- Thorlund K, Walter SD, Johnston BC, Furukawa TA, Guyatt GH. Pooling health-related quality of life outcomes in meta-analysis—a tutorial and review of methods for enhancing interpretability. *Res Synth Methods* 2011;2(3):188–203.
- Johnston BC, Patrick DL, Thorlund K, et al. Patient-reported outcomes in meta-analyses –Part 2: Methods for improving interpretability for decision-makers. *Health QualLife Outcomes* 2013;11:211.
- Bucher HC, Guyatt GH, Griffith LE, Walter SD. The results of direct and indirect treatment comparisons in meta-analysis of randomized controlled trials. *J Clin Epidemiol* 1997;50(6): 683–91.
- Wang L, Guyatt GH, Kennedy SA, et al. Predictors of persistent pain after breast cancer surgery: A systematic review and meta-analysis of observational studies. *CMAJ* 2016;11:1–10.
- Valdes AM, Doherty SA, Zhang W, Muir KR, Maciewicz RA, Doherty M. Inverse relationship between preoperative radiographic severity and postoperative pain in patients with osteoarthritis who have undergone total joint arthroplasty. *Semin Arthritis Rheum* 2012;41(4):568–75.
- Spencer FA, Iorio A, You J, et al. Uncertainties in baseline risk estimates and confidence in treatment effects. *BMJ* 2012;345: e7401.
- Schandelmaier S, Briel M, Varadhan R, et al. Development of the Instrument to assess the Credibility of Effect Modification

- Analyses (ICEMAN) in randomized controlled trials and meta-analyses. *CMAJ* 2020;192(32):E901–E906.
22. Andridge RR, Little RJA. A review of hot deck imputation for survey non-response. *Int Stat Rev = Revue Internationale de Statistique* 2010;78(1):40–64.
 23. Iorio A, Spencer FA, Falavigna M, et al. Use of GRADE for assessment of evidence about prognosis: Rating confidence in estimates of event rates in broad categories of patients. *BMJ* 2015;350:h870.
 24. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction—GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64(4):383–94.
 25. Foroutan F, Iorio A, Thabane L, Guyatt G. Calculation of absolute risk for important outcomes in patients with and without a prognostic factor of interest. *J Clin Epidemiol* 2020;117:46–51.
 26. Higgins JPT, Thomas J, Chandler J, et al., eds. *Cochrane Handbook for Systematic Reviews of Interventions*, 2nd edition. Chichester, UK: John Wiley & Sons; 2019.
 27. Skinner J, Zhou W, Weinstein J. The influence of income and race on total knee arthroplasty in the United States. *J Bone Joint Surg Ser A* 2006;88(10):2159–66.
 28. Riddle DL, Wade JB, Jiranek WA, Kong X. Preoperative pain catastrophizing predicts pain outcome after knee arthroplasty. *Clin Orthopaed Relat Res* 2010;468(3):798–806.
 29. Sakellariou VI, Poultides LA, Ma Y, Bae J, Liu S, Sculco TP. Risk assessment for chronic pain and patient satisfaction after total knee arthroplasty. *Orthopedics* 2016;39(1):55–62.
 30. Dave AJ, Selzer F, Losina E, et al. The association of pre-operative body pain diagram scores with pain outcomes following total knee arthroplasty. *Osteoarthritis Cartil* 2017;25(5):667–75.
 31. Singh JA, Gabriel SE, Lewallen DG. Higher body mass index is not associated with worse pain outcomes after primary or revision total knee arthroplasty. *J Arthroplasty* 2011;26(3):366–74.e1.
 32. Katz JN, Mahomed NN, Baron JA, et al. Association of hospital and surgeon procedure volume with patient-centered outcomes of total knee replacement in a population-based cohort of patients age 65 years and older. *Arthritis Rheum* 2007;56(2):568–74.
 33. Gungor S, Fields K, Aiyer R, Della Valle AG, Su EP. Incidence and risk factors for development of persistent postsurgical pain following total knee arthroplasty: A retrospective cohort study. *Medicine* 2019;98(28):e16450.
 34. Buvanendran A, Della Valle CJ, Kroin JS, et al. Acute postoperative pain is an independent predictor of chronic postsurgical pain following total knee arthroplasty at 6 months: A prospective cohort study. *Regional Anesth Pain MED* 2019;44(3):e100036.
 35. Vina ER, Ran D, Ashbeck EL, Kwoh CK. Widespread pain is associated with increased risk of no clinical improvement after TKA in women. *Clin Orthopaed Relat Res* 2020;478(7):1453.
 36. Vila MR, Todorovic MS, Tang C, et al. Cognitive flexibility and persistent post-surgical pain: The FLEXCAPP prospective observational study. *Br J Anaesth* 2020;124(5):614–22.
 37. Barrack RL, Ruh EL, Chen J, et al. Impact of socioeconomic factors on outcome of total knee arthroplasty. *Clin Orthopaed Relat Res* 2014;472(1):86–97.
 38. König A, Scheidler M, Rader C, Eulert J. The need for a dual rating system in total knee arthroplasty. *Clin Orthopaed Relat Res* 1997;345:161–7.
 39. Elson DW, Brenkel IJ. Predicting pain after total knee arthroplasty. *Journal of Arthroplasty* 2006;21(7):1047–53.
 40. Lundblad H, Kreicbergs A, Jansson KA. Prediction of persistent pain after total knee replacement for osteoarthritis. *J Bone Joint Surg (BR)* 2008;90(2):166–71.
 41. Puolakka PAE, Rorarius MGF, Roviola M, Puolakka TJS, Nordhausen K, Lindgren L. Persistent pain following knee arthroplasty. *Eur J Anaesthesiol* 2010;27(5):455–60.
 42. Thomazeau J, Rouquette A, Martinez V, et al. Predictive factors of chronic post-surgical pain at 6 months following knee replacement: Influence of postoperative pain trajectory and genetics. *Pain Physician* 2016;19(5):E729.
 43. Birch S, Stilling M, Mechlenburg I, Hansen TB. The association between pain catastrophizing, physical function and pain in a cohort of patients undergoing knee arthroplasty. *BMC Musculoskelet Disord* 2019;20(1):421.
 44. Bugada D, Allegri M, Gemma M, et al. Effects of anaesthesia and analgesia on long-term outcome after total knee replacement: A prospective, observational, multicentre study. *Eur J Anaesthesiol* 2017;34(10):665–72.
 45. Nielsen CS, Nebergall A, Huddleston J, Kallemose T, Malchau H, Troelsen A. Medial overhang of the tibial component is associated with higher risk of inferior knee injury and osteoarthritis outcome score pain after knee replacement. *J Arthroplasty* 2018;33(5):1394–8.
 46. Scott CEH, Clement ND, Yapp LZ, MacDonald DJ, Patton JT, Burnett R. Association between femoral component sagittal positioning and anterior knee pain in total knee arthroplasty: A 10-year case-control follow-up study of a cruciate-retaining single-radius design. *J Bone Joint Surg Am* 2019;101(17):1575–85.
 47. Skrejborg P, Petersen KK, Kold S, Kappel A, et al. Presurgical comorbidities as risk factors for chronic postsurgical pain following total knee replacement. *Clin J Pain* 2019;35(7):577–82.
 48. Karsdal MA, Bihlet A, Byrjalsen I, et al. OA phenotypes, rather than disease stage, drive structural progression: Identification of structural progressors from 2 phase III randomized clinical studies with symptomatic knee OA. *Osteoarthritis Cartilage* 2015;23(4):550–8.
 49. Getachew M, Lerdal A, Småstuen MC, et al. High levels of pre-operative pain and fatigue are red flags for moderate-severe pain 12 months after total knee arthroplasty—A longitudinal cohort study. *Musculoskelet Care* 2021;19(2):186–92.
 50. Smith AJ, Lloyd D, Wood D. Pre-surgery knee joint loading patterns during walking predict the presence and severity of anterior knee pain after total knee arthroplasty. *J Orthopaed Res* 2004;22(2):260–6.
 51. Dowsey MM, Dieppe P, Lohm e. S, Castle D, Liew D, et al. The association between radiographic severity and pre-operative function in patients undergoing primary knee replacement for osteoarthritis. *Knee* 2012;19(6):860–5.
 52. Rice D, Kluger M, McNair P, et al. Persistent postoperative pain after total knee arthroplasty: A prospective cohort study of potential risk factors. *Br J Anaesth* 2018;121(4):804–12.
 53. Kim S, Yoon K, Yoon D, Yoo J, Ahn K. Influence of centrally mediated symptoms on postoperative pain in osteoarthritis patients undergoing total knee arthroplasty: A prospective observational evaluation. *Pain Pract* 2015;15(6):E46–e53.
 54. Nashi N, Hong CC, Krishna L. Residual knee pain and functional outcome following total knee arthroplasty in osteoarthritic patients. *Knee Surg Sports Traumatol Arthrosc* 2015;23(6):1841–7.
 55. Albayrak I, Apiliogullari S, Erkocak OF, Kavalci H, Ozerbil OM, Levendoglu F. Total knee arthroplasty due to knee osteoarthritis: Risk factors for persistent postsurgical pain. *J Natl Med Assoc* 2016;108(4):236–43.
 56. Sugiyama Y, Iida H, Amaya F, et al. Prevalence of chronic post-surgical pain after thoracotomy and total knee arthroplasty: A retrospective multicenter study in Japan (Japanese Study Group of Subacute Postoperative Pain). *J Anesth* 2018;32(3):434–8.

57. Aso K, Ikeuchi M, Takaya S, et al. Chronic postsurgical pain after total knee arthroplasty: A prospective cohort study in Japanese population. *Modern Rheumatol* 2021;31(5):1038–44.
58. Darnall BD, Sturgeon JA, Cook KF, et al. Development and validation of a daily pain catastrophizing scale. *J Pain* 2017;18(9):1139–49.
59. Petrini L, Arendt-Nielsen L. Understanding pain catastrophizing: Putting pieces together. *Front Psychol* 2020;11:603420.
60. Riis A, Rathleff MS, Jensen MB, Simonsen O. Low grading of the severity of knee osteoarthritis pre-operatively is associated with a lower functional level after total knee replacement: A prospective cohort study with 12 months' follow-up. *Bone Joint J* 2014;96-B(11):1498–502.
61. Lewis GN, Rice DA, McNair PJ, Kluger M. Predictors of persistent pain after total knee arthroplasty: A systematic review and meta-analysis. *Br J Anaesth* 2015;114(4):551–61.
62. Bhan S, Malhotra R, Eachempati KK. Total knee arthroplasty without patellar resurfacing in patients with rheumatoid arthritis. *Clin Orthop Related Res* 2006;450(1):157–63.
63. Murray DW, Frost SJ. Pain in the assessment of total knee replacement. *J Bone Joint Surg Br* 1998;80(3):426–31.
64. Forsythe ME, Dunbar MJ, Hennigar AW, Sullivan MJL, Gross M. Prospective relation between catastrophizing and residual pain following knee arthroplasty: Two-year follow-up. *Pain Res Manage* 2008;13(4):335–41.
65. Nicoll D, Rowley DI. Internal rotational error of the tibial component is a major cause of pain after total knee replacement. *J Bone Joint Surg Br* Vol 2010;92(9):1238–44.
66. Ritter MA, Wing JT, Berend ME, Davis KE, Meding JB. The clinical effect of gender on outcome of total knee arthroplasty. *J Arthroplasty* 2008;23(3):331–6.
67. Whiteside LA. The effect of patient age, gender, and tibial component fixation on pain relief after cementless total knee arthroplasty. *Clin Orthop Relat Res* 1991;271:21–7.
68. Cross M, Lapsley H, Barcenilla A, Parker D, Coolican M, March L. Patient expectations of hip and knee joint replacement surgery and postoperative health status. *Patient* 2009;2(1):51–60.
69. Creameans-Smith JK, Boarts JM, Greene K, Delahanty DL. Patients' reasons for electing to undergo total knee arthroplasty impact post-operative pain severity and range of motion. *J Behav Med* 2009;32(3):223–33.
70. Lingard EA, Riddle DL. Impact of psychological distress on pain and function following knee arthroplasty. *J Bone Joint Surg Am* 2007;89(6):1161–9.
71. Quintana JM, Escobar A, Arostegui I, et al. Health-related quality of life and appropriateness of knee or hip joint replacement. *Arch Intern Med* 2006;166(2):220–6.
72. Singh JA, Gabriel S, Lewallen D. The impact of gender, age, and preoperative pain severity on pain after TKA. *Clin Orthopaed Relat Res* 2008;466(11):2717–23.
73. Singh JA, O'Byrne MM, Colligan RC, Lewallen DG. Pessimistic explanatory style: A psychological risk factor for poor pain and functional outcomes two years after knee replacement. *J Bone Joint Surg (BR)* 2010;92(6):799–806.
74. Barrack RL, Bertot AJ, Wolfe MW, Waldman DA, Milicic M, Myers L. Patellar resurfacing in total knee arthroplasty: A prospective, randomized, double-blind study with five to seven years of follow-up. *J Bone Joint Surg (Am)* 2001;83(9):1376–81.
75. Barrack RL, Wolfe MW, Waldman DA, Milicic M, Bertot AJ, Myers L. Resurfacing of the patella in total knee arthroplasty: A prospective, randomized, double-blind study. *J Bone Joint Surg Am* 1997;79(8):1121–31.
76. Wertli MM, Eugster R, Held U, Steurer J, Kofmehl R, Weiser S. Catastrophizing—a prognostic factor for outcome in patients with low back pain: A systematic review. *Spine J* 2014;14(11):2639–57.
77. Duan G, Liu C, Lin W, et al. Different factors conduct anterior knee pain following primary total knee arthroplasty: A systematic review and meta-analysis. *J Arthroplasty* 2018;33(6):1962–71.e3.
78. Wylde V, Beswick AD, Dennis J, Gooberman-Hill R. Post-operative patient-related risk factors for chronic pain after total knee replacement: A systematic review. *BMJ Open* 2017;7(11):e018105.
79. Harmelink KEM, Zeegers AVCM, Hullegie W, Hoogeboom TJ, Nijhuis-van der Sanden MWG, Staal JB. Are there prognostic factors for one-year outcome after total knee arthroplasty? A Systematic Review. *J Arthroplasty* 2017;32(12):3840–53.e1.
80. Lindberg MF. Factors that predict the trajectories of acute pain, pain with walking, and self-rated health after total knee arthroplasty. 2017.
81. Rücker G, Schwarzer G, Carpenter JR, Schumacher M. Undue reliance on I 2 in assessing heterogeneity may mislead. *BMC Med Res Methodol* 2008;8:79.