




C01 | How much recovery improves over the stored pool



C02 | Additional recoveries and lost recoveries




C03 | Which ligands are recovered by which methods?





C04 | Pairwise complementarity between generators





Method recovering the ligand




C05 | Best available conformers versus typical conformers




C06 | Per-ligand geometric agreement in selected comparisons




C07 | Where the core cohort has statistical support





C08 | Different summaries of geometric diversity



C09 | Physical validity and bound-conformer recovery



C10 | Classical raw and minimized pool comparisons



C11 | Coverage and precision for every drug molecule


Reference coverage / recall (COV-R, %)

4-Hydroxytamoxifen	100	100	100	100	100	100
Acetazolamide	98	100	96	96	100	100
Actinonin	68	22	1	0	0	4
Amprenavir	93	0	0	0	0	77
Bortezomib	68	50	25	29	0	37
Bromosporine	96	100	100	100	96	100
Chloramphenicol	96	98	99	98	98	99
Colchicine	100	0	100	100	100	100
Cyclothiazide	100	100	100	99	100	100
Dexamethasone	97	100	97	97	100	97
Diclofenac	89	100	96	96	100	100
Fluconazole	97	100	94	97	100	100
Fosmidomycin	96	100	100	96	100	100
IBMX	100	100	100	100	100	100
Imatinib	5	0	0	2	0	79
Indomethacin	94	98	94	94	100	100
Nicotine	100	100	100	100	100	100
Pleconaril	100	43	100	100	80	100
S-thalidomide	100	100	100	100	100	100
Staurosporine	100	100	100	100	100	100
Trichostatin A	100	97	100	97	97	100
Triclosan	100	100	100	100	100	100
Trimethoprim	100	100	100	100	100	100
	LoQI	Torsional Diffusion	MCF drugs-L	NEXT-Mol DMT-L	FlowR	Qwen 1.7B FSQ


Observed-reference precision (COV-P, %)

4-Hydroxytamoxifen	46	8	17	21	7	21
Acetazolamide	100	49	50	51	65	85
Actinonin	6	0	0	0	0	0
Amprenavir	0	0	0	0	0	0
Bortezomib	26	2	2	2	0	6
Bromosporine	39	19	34	37	4	8
Chloramphenicol	90	50	32	25	24	42
Colchicine	52	0	6	16	4	7
Cyclothiazide	48	52	14	7	40	62
Dexamethasone	83	86	85	36	50	99
Diclofenac	99	52	48	58	62	66
Fluconazole	3	7	13	9	8	6
Fosmidomycin	41	9	13	5	27	52
IBMX	100	100	100	100	98	100
Imatinib	1	0	0	1	0	1
Indomethacin	100	97	99	98	78	73
Nicotine	100	100	80	86	94	97
Pleconaril	4	0	4	8	1	3
S-thalidomide	100	98	65	82	89	78
Staurosporine	87	100	76	45	29	89
Trichostatin A	50	10	8	15	4	8
Triclosan	95	59	65	62	80	84
Trimethoprim	88	82	87	86	76	90
	LoQI	Torsional Diffusion	MCF drugs-L	NEXT-Mol DMT-L	FlowR	Qwen 1.7B FSQ

C12 | Coverage and concentration define different preferences




C13 | Similar mean recall can hide opposing molecule-level outcomes




C14 | Chemical validity varies strongly across molecules and methods

4-Hydroxytamoxifen	100	53	100	94	0	99
Acetazolamide	100	49	25	21	44	96
Actinonin	98	68	25	18	96	77
Amprenavir	92	48	9	14	82	73
Bortezomib	96	81	27	25	0	4
Bromosporine	100	77	87	94	62	75
Chloramphenicol	100	95	24	25	97	97
Colchicine	99	51	60	50	70	17
Cyclothiazide	91	97	8	6	94	16
Dexamethasone	6	100	80	24	72	99
Diclofenac	100	95	100	100	96	96
Fluconazole	100	98	100	100	96	92
Fosmidomycin	100	86	99	100	100	99
IBMX	100	100	100	100	98	100
Imatinib	100	56	66	87	76	99
Indomethacin	100	100	100	100	93	100
Nicotine	100	100	46	48	98	96
Pleconaril	100	93	100	100	95	98
S-thalidomide	100	97	42	54	95	99
Staurosporine	9	95	20	20	46	95
Trichostatin A	99	79	48	51	95	97
Triclosan	100	100	100	100	98	100
Trimethoprim	100	100	100	100	94	100
	LoQI	Torsional Diffusion	MCF drugs-L	NExT-Mol DMT-L	FlowR	Qwen 1.7B FSQ


C15 | Continuous matching distances complement coverage



C16 | The experimental reference collection is unevenly sampled




C17 | How uncertain are differences from LoQI?




C18 | Illustrative failures differ across constraints


Imatinib · 57 references



Actinonin · 99 references



Bortezomib · 189 references



Dexamethasone · 32 references

